

Test Cases, Resolvability, and Group Selection: A Critical Examination of the Myxoma Case*

Robert A. Wilson^{†‡}

The evolution of the myxoma virus in Australia has been presented for many years as a test case for the hypothesis that group selection can function effectively ‘in the wild.’ This paper critically examines the myxoma case, and argues that its failure as a test case for this hypothesis has broader implications for debates over the levels of selection.

1. Introduction. In virtue of their controversial or novel nature, some views and theories carry a special burden of proof. A dialectically powerful way to discharge that burden is to present a clear and decisive example that can only or can best be explained by the view or theory. In science, such examples take the form of phenomena that serve as test cases for the view or theory under consideration.

One such view in evolutionary biology is the theory of group selection, particularly as articulated and defended in a series of publications by David Sloan Wilson and Elliott Sober (Wilson 1975, 1980, 1983, 1989, 1997, 2002; Sober 1984, 1993; Wilson and Sober 1994; Sober and Wilson 1994, 1998). Amongst the theoretical work that Wilson and Sober have done in articulating group selection as a coherent and potentially powerful force directing evolution, and the empirical evidence they have adduced in

*Received April 2003; revised October 2003.

† To contact the author write to Department of Philosophy, 4-115 Humanities, University of Alberta, Edmonton, Alberta, T6G 2E5, Canada; e-mail: rob.wilson@ualberta.ca.

‡ I would like to thank audiences at the University of Alberta and at the 2002 meetings of Western Canadian Philosophical Association held in Calgary, as well as two referees for the journal, for their constructive feedback on earlier versions of this paper. The work presented here was written with the financial support of the Social Sciences and Humanities Research Council of Canada.

Philosophy of Science, 71 (July 2004) pp. 380–401. 0031-8248/2004/7103-0006\$10.00
Copyright 2004 by the Philosophy of Science Association. All rights reserved.

support of the efficacy of group selection in actual populations, are several examples they represent as test cases of their theory. In their recent *Unto Others*, they present two such examples: the evolution of female-biased sex ratios, and the evolution of virulence (1998, 35–50). In this paper I shall focus exclusively on the latter case, particularly on the most detailed example they discuss, that of the evolution of avirulence in the myxoma virus, an example originally introduced into the debate over the levels of selection by Richard Lewontin 1970.

There are two immediate conclusions that I shall argue for from an examination of this case, and a third that warrants further discussion. These conclusions carry with them a number of broader implications for how we do and might think about biological reality that I discuss elsewhere (Wilson 2003). The immediate conclusions themselves are of increasing generality, moving from a weaker to a stronger claim about the myxoma case, and then in turn to the debate over the levels of selection more generally.

First, as described in the philosophical literature on the units of selection, the myxoma case does not, *contra* Wilson and Sober, “provide compelling evidence for group selection” (1998, 46): it is not the decisive test case of group selection theory that they claim it is. This is not simply because there is a possible, competing individual-level explanation of the case, but because there is no independent basis for preferring the group selection account of the phenomenon to alternative explanations (Sections 2–3).

Second, the myxoma case is likely to remain subject to multiple, alternative explanations, only some of which appeal to group selection. This second conclusion goes beyond the first not only in strength, but also by taking us beyond the abbreviated characterizations of the myxoma case that have featured in the debate over the units of selection into the epidemiological and virological literatures in which myxomatosis is discussed more fully. One could reasonably expect the addition of further empirical details to the case to reveal the true level or levels at which the evolution of virulence is governed by natural selection. Yet far from achieving this goal, restoring some of the complexities to the myxoma case usually omitted or glossed over by philosophers and biologists alike reinforces and strengthens the first conclusion. The myxoma case is likely to be *irresolvable* vis-à-vis the debate over the levels of selection (Sections 4–6).

Third, what is true of the myxoma case is true more generally, and so the general debate between proponents of group selection and other units of selection is also likely to be irresolvable. Invoking the metaphor of viewpoints or standpoints, many evolutionary phenomena that can be adequately explained by invoking one viewpoint—whether it be that of

the gene, the individual, or the group—can be explained as adequately by invoking at least one other viewpoint. A defense of this final claim would involve at least a detailed examination of the other test case for group selection theory that Wilson and Sober provide, that of the evolution of female-biased sex ratios. And a version of this generalization that treated various levels of selection symmetrically would require exploring other phenomena, such as meiotic drive, that have been claimed by many to constitute decisive cases vis-à-vis the theory of *genic* selection.

A strong form of this third conclusion is entailed by a widespread form of *pluralism* about the levels of selection that maintains that these viewpoints are, in an important sense, equivalent (Dugatkin and Reeve 1994; Sterelny 1996; Kerr and Godfrey-Smith 2002). Since I am skeptical of stronger versions of this conclusion and of these forms of pluralism, my argument here does not rest on pluralism about the levels of selection (Section 7).

I begin with a familiar sketch of the myxoma case and the debate over it in Section 2, going on in Section 3 to argue against the most commonly cited reasons for resolving this debate one way or the other. Included here is the appeal that Sober and Wilson make to “the averaging fallacy” in criticizing individualistic explanations of the myxoma case. These authors have invoked this fallacy, separately and together, for many years in their defense of group selection. Yet attributions of the fallacy do little to support the idea that the myxoma case is a paradigm case of group selection in the wild.

2. The Case of the Myxoma Virus: A Sketch and Two Interpretations.

The myxoma virus was introduced into Australia as a way to control an exploding rabbit population in 1950, a solution that met with unexpected limits to its success. After myxomatosis initially killed 99% of the hosts infected, and seemingly quickly, the rabbit population began to rise steadily again. As expected from individual selection theory, rabbits from the wild tested against laboratory strains of the virus had an increased resistance to the virus. But it was also found that the virus in the wild had *decreased* its virulence, compared to those very laboratory strains, a finding anomalous within the theory of individual selection.

In Australia in 1950, the virus was primarily transmitted between host rabbits by an arthropod vector, the mosquito, and, as Lewontin notes, “mosquitoes do not bite dead rabbits” (1970, 15). Thus, the transmission of the virus from dead hosts is minimal. More virulent forms of the virus kill the host organism more quickly, and thus are less likely to be transmitted to other host organisms. Less virulent forms of the virus tend to fixation in the overall host population of rabbits, with more virulent forms disappearing. The chief phenomenon to be explained is the apparent

evolution of what is often termed “avirulence” in the myxoma virus. Why did avirulent strains of the virus predominate? What was the mechanism that brought this observed result about?

The standard group selectionist interpretation and explanation of this phenomenon is as follows. Within single host organisms, more virulent forms of the virus are favored, since these have a higher reproductive rate, and come to replace less virulent strains within any given rabbit. Thus, higher virulence is fitter *within* an organismic host than is lower virulence and evolves by individual selection. But forms of the virus with lower virulence are likely to infect more hosts by allowing their hosts to live longer, and so are fitter *across* organismic hosts. In effect, viruses on a given rabbit are a *trait group*: they are “populations enclosed in areas smaller than the boundary of the deme” (Wilson 1975, 143), or sets “of individuals that influence each other’s fitness with respect to a certain trait but not the fitness of those outside the group” (Wilson and Sober 1998, 92). If we assume that an individual rabbit delineates a group of viruses, then this implies that low virulence is altruistic, and that it evolves because of the higher fitness of less virulent groups of viruses. This is a process of selection between groups, and thus an example of *group selection* (Wilson 1989, 148; Sober 1993, 109).

There is, however, an obvious, contrasting individualist explanation for the phenomenon, one provided by Alexander and Borgia (1978, 451–453) and by Futuyma (1979, 455). A viral strain that kills its host is less fit than one that does not kill its host. Low virulence in a viral strain reduces the chance that it will kill its host. Thus, low virulence is an adaptation of individual viral units, and it evolves by a process of selection that acts on individuals. As a result, lower levels of virulence spread through the population of viruses as a whole via *individual selection*.

On this view, the only relevant groups are the overall population of rabbits and the overall population of viruses. Individual selection changes the composition of the latter of these from one with predominantly highly virulent individuals to one with predominantly less virulent individuals over evolutionary time. In positing low virulence as an adaptation of individual viruses, this view also denies the existence of low virulence as an altruistic trait, on a range of views of altruism, since low virulence is depicted as being to the evolutionary advantage of the individual virus.

Sober and Wilson (1998, 31–34, 46–50) have made two related criticisms of this sort of individualistic interpretation, criticisms first made twenty years ago (Sober 1984, 327–335). First, it involves what they call the *averaging approach* to calculating fitness: it calculates the fitness of traits across the entire metapopulation, rather than within the constituent populations, i.e., groups. Sober and Wilson regard the averaging approach to fitness in this context as involving a fallacy, the

averaging fallacy. As a fallacy, this is the mistake of using the averaging approach to fitness to draw a conclusion about the level of selection that governs its distribution. This inference is a fallacy, according to Sober and Wilson, because in adopting the averaging approach one collapses the distinction between within-and between-group selection, itself crucial to characterizing, respectively, the process of individual selection and that of group selection. Thus, once one adopts the averaging approach, no inferences can be validly drawn about the level at which selection operates (cf. Wilson 1989). Second (and subsequently), by focusing exclusively on the products of natural selection, overall fitness values, this sort of interpretation ignores the processes that generate those products. The debate over the units of selection, however, is precisely one over *how* natural selection operates. Thus, the individualistic explanation begs the question at issue.

Individualistic explanations of the myxoma case have been motivated by a general concern over the coherence or prevalence of group selection, one that turns in large part on doubts about the *ontological* status of groups. One suspicion is that there is no way to demarcate groups of organisms from mere sets of organisms, thus implying that any n-tuple of individuals could be said to constitute a group. Call this the *arbitrariness problem*. And there is the objection that groups, unlike individuals or genes, are not sufficiently permanent arrangements in the biological world to be the units on which natural selection operates over evolutionary time. Call this the *ephemerality problem*. Both problems reflect a concern over the ontological status of groups insofar as they express the view that groups are “not real” in the way that genes and individuals are. Such existence as they have makes them unsuitable to play the role of a unit of selection in general, and in the myxoma case in particular.

3. The Dialectics of the Debate: Averaging and the Status of Groups. If the individual selectionist account of the myxoma case sketched above were to commit a fallacy, as Sober and Wilson claim, then clearly that would provide a reason to favor the group selectionist account of it. And, conversely, if the group selectionist account of the myxoma case were to suffer from either the arbitrariness or the ephemerality problem, then the *prima facie* gridlock between the two explanations would be broken in favor of the individual selection account. It is important to recognize that neither resolution of the debate over the myxoma case is plausible.

Consider first the claim that the group selectionist account of the myxoma case suffers from the arbitrariness and ephemerality problems, and that this provides a conclusive reason for preferring the individual selectionist account. Neither charge holds of the groups of viruses bounded by individual host organisms. A rabbit-bound population

of viruses constitutes an entity that faces shared ecological and evolutionary circumstances: individual viruses in such a group draw on the same resources, and in virtue of physical proximity to one another, face the possibility of host resistance or vector transmission together. Importantly, populations of viruses on different rabbits are likely to differ from one another in these respects, and so differ in fitness. Thus rabbit-bound groups of viruses are not arbitrary from an evolutionary point of view (cf. Section 6 below, however). Furthermore, given that typically there will be hundreds of generations of viruses on any given rabbit, these groups are not ephemeral either. In short, even if there *are* putative instances of group selection that are faced with the arbitrariness and ephemerality problems, group selection in the myxoma case cannot be rejected because the case invokes arbitrary or ephemeral groups.

Let us turn now to the averaging fallacy. There are two responses that can be made to the charge that such a fallacy is committed in the individual level explanation of the myxoma case. The first of these concedes that the inference from adopting the averaging approach to fitness to a conclusion about the level at which selection operates *is* a fallacy, but denies that this inference is made in explaining the myxoma case. The second response argues that the attribution of the averaging fallacy in this case, and perhaps more generally, itself begs the question against the proponent of individual selection. Let us take each in turn.

If the averaging fallacy is committed in the individual level explanation of the myxoma case, then that explanation must somewhere invoke the averaging approach to calculating fitness. Thus, whether the averaging fallacy is committed turns on what basis there is for the claims that:

- (1) A viral strain that kills its host is less fit than one that does not.
- (2) Low virulence reduces the chances of killing one's host.

Thus,

- (3) Low virulence is an adaptation of individual viruses.

So,

- (4) Low virulence evolves by individual selection.

For the averaging fallacy to be committed here, the fitness explicit in (1) and implicit in (3) must be conceptualized or calculated as an average over the whole population of viruses. Furthermore, this averaging must be the reason for holding at least one of (1)–(4) to be true.

There are, however, at least two related reasons for thinking that (1)–(4) are true that are independent of whether fitness is calculated over the whole population. Both reasons pack some dialectical punch in part as a result of the distance that proponents of the “new” group selection have

self-consciously promoted between their views and the “old” group selection tradition.

First, it has been well known at least since the work of David Lack (1954) on optimal clutch size in birds that individuals can increase their reproductive success by limiting the number of offspring they produce. One might think that reduced virulence in individual viruses is simply an instance of such a reproductive strategy, and so an adaptation of individual viruses that reflects the increased fitness of those viruses. Explanations such as Lack’s have not been seriously disputed by proponents of group selection as appropriate in at least some cases. This suggests that it does not commit the averaging fallacy. Given that, there would have to be some relevant difference in the myxoma case that vitiates the same reasoning that Lack used, were any “averaging fallacy” charge to be plausible in that case.

Second, since the fitness of any trait needs to be relativized to an environment, low virulence may be viewed as having higher fitness once we consider an environment in which host death constitutes a major limitation on long-term reproductive success. One comes to view lower virulence as an adaptation by considering the nature of this environment more fully, rather than simply by defining fitness and adaptation as “what evolves.” Again, there is a basis for (1)–(4) that appeals to points that group selectionists themselves appear to be committed to.

Such reasons are independent of whether we calculate fitnesses within or across local groups of viruses (= rabbits) in that they are factors “in play,” so to speak, prior to *any* calculations of fitness at all. Of course, if the individual selectionist viewpoint is adopted, then fitnesses *will* be calculated across the whole population of viruses. But provided that this is not the basis for adopting (1)–(4), the averaging fallacy has not been committed.

So much for the first response to the averaging fallacy charge. The second response is more dialectically powerful in the context of the overall argument of the paper. The idea behind the averaging fallacy charge is that individualists illegitimately collapse the distinction between within and between group fitnesses by averaging across groups to begin with. But one can collapse this distinction (illegitimately or not) only if the distinction is already there to collapse. This in turn presupposes not simply that there are groups of viruses (= rabbits) in the overall population, but that these groups constitute a causal structure to which natural selection itself is sensitive. Yet this is precisely what proponents of the individual level explanation will deny—not necessarily in general but in this particular case. For although there is a sense in which rabbits are groups of viruses, it is not this group structure on which natural selection operates. Rather, it operates on individual viruses, viruses whose fitness is sensitive to features

of their environment, including the nature of their hosts (= groups of viruses) and limitations on transmission imposed by the nature of the vectors that mediate the spread of the virus. In short, an averaging fallacy can be committed only if we already concede that groups themselves are acted on by natural selection, and this is precisely what is at issue between proponents of the two explanations. Ironically, it is Sober and Wilson's attribution of an averaging fallacy itself that begs the question against the individualist.

To summarize: Sober and Wilson argue that at least some individualistic views of the myxoma case commit the averaging fallacy, and that interpretations that rely crucially on a misplaced skepticism about the reality of groups (of viruses, in this case) should be rejected. I concur. But in generalizing this claim and presenting the myxoma case as a "test case" for the theory of group selection, a test that the theory passes, Sober and Wilson are mistaken. The test can also be passed by the theory of individual selection. Moreover, their general invocation of the averaging fallacy itself begs the very question at issue. At this point, I draw my first conclusion: that the myxoma case, as it is usually presented, is not a decisive example of group selection.

I have already remarked that the dispute over the myxoma case has involved an abbreviated characterization of the case. One healthy suspicion fostered by interplay between the philosophy, history, and social studies of science over the last 40 years is directed at just this sort of impoverished treatment of actual examples: that our anemic characterization of "the evolution of avirulence" in myxoma actually creates or maintains the standoff in the debate over it. Shouldn't a more complete rendering of the facts allow a resolution of the apparent deadlock between proponents of our two perspectives?

In the next section I fill out the picture of the pathology of the disease and the nature of its transmission, drawing largely on the book-length treatments of Fenner and Ratcliffe (1965) and Fenner and Fantinni (1999). I shall argue that not only does a more complete rendering of the case fail to challenge my conclusion about the irresolvability of the case, but that it actually *reinforces* that conclusion in a way that suggests the pair of stronger conclusions about irresolvability stated in the introduction.

4. A Little More on the Virology and Epidemiology of Myxomatosis.

The myxoma virus is a member of the family *Poxviridae*, and the disease it causes in rabbits was first described by Sanarelli in 1898 when European rabbits he received for his laboratory work in Uruguay (sent from Brazil) developed the disease. The virus is relatively host specific, being largely restricted to a variety of "rabbit species," including those from

the *Sylvilagus* and *Oryctolagus* genera, commonly called, respectively, the Californian and European rabbit. The strain of the virus introduced into Australia was isolated from an *Oryctolagus* rabbit in Brazil in 1911, and was maintained through serial passage from host to host until it was introduced to Australia in 1950.

At the initial site of myxoma infection in rabbits a skin lesion appears, and the virus begins replicating in the body of the host, working its way through lymph nodes and the circulatory system. Secondary lesion sites also appear, and there is often a discharge from the eyes and a swelling of the genitalia. The concept of virulence used by epidemiologists is host-fatality, the percentage of infected rabbits that die, though this is commonly estimated from a more readily operationalized measure, the average number of days to death. An estimate was necessitated in Australia by the impracticality of keeping large numbers of rabbits, and was made possible by the correlation between case-mortality rate and the more readily measured-in-the-lab variable of average number of days to death. (For questions about the adequacy of this basic measure of virulence, however, see Parer 1995; Fenner and Fantini 1999, 193, 297–298).

With the original strain of the virus introduced into Australia, infected laboratory rabbits survived about five days, making the virus extremely lethal. In one of the first systematic field counts, at Lake Urana in 1951, the lethality of the virus was 99.8%, this being reduced to 90% in 1952 (Fenner and Ratcliffe 1965, 284–286).

Since the concept of virulence used here is relational, this reduction in virulence alone could in principle be accounted for by a change in host resistance in the population. But suspecting that the virus had changed, researchers checked virus serum taken from infected rabbits against captive rabbits that had been isolated from the virus until that point, and found that the virus itself was reduced in its lethality. In their attempts to measure the changes in the virulence of the virus itself, from 1951 researchers began gathering multiple samples of the virus from wild populations of infected rabbits. By 1959 they had 672 such strains (no doubt, not all distinct), and they introduced five grades of virulence for these, rated by their estimated case-mortality rates, starting with >99% for Grade I, the original strain (rarely recovered from the wild), through to <50% for Grade V. The virus stabilizes at Grade III, with this constituting roughly 50% of the strains throughout the 1950s, with an estimated lethality of 70–95%. The incidence of Grade II viruses (lethality: 95–99%) remains over 10% of the sampled population for much of the period surveyed, and increases significantly from 1958 to 1959; the incidence of Grade V viruses remains extremely low throughout, and Fenner and Ratcliffe note that “it is unlikely that a virus of this type would survive in

nature” (1965, 221). Simple talk of “the evolution of avirulence” neither conveys nor does justice to the complexities here.¹

Transmission is also more complicated than the standard picture from Section 2 suggests. While the most prevalent and far-reaching form of transmission is via an arthropod vector, the virus can also be transmitted through direct contact or contagion between hosts, or through non-animate vectors (such as thornbushes or warrens). Given that rabbits live in close proximity to one another in hutches, these are actual (not merely possible) mechanisms for transmission. Arthropod vectors pick up the virus on their mouthparts when they feed on an infected host animal, particularly at a lesion site, and then transmit the virus when they move to another host, or to another part of the same host. Vectors preferentially feed on the head of the rabbit, perhaps because it is the primary site of lesions. Thus, as well as between-host transmission of the virus, there is within-host transmission facilitated by vectors themselves and by the host’s circulatory system, and differential vector transmission within a host.

Transmissibility of the virus by arthropod vectors, such as the mosquito, depends on facts about the ecology and life cycle of those vectors. The concentration of mosquitoes varies in accord with heat and moisture, and thus their abundance follows a seasonal cycle. This represented a problem for the use of myxoma in controlling the wild population of rabbits in Australia, not only in areas in which the concentration of mosquitoes was insufficient to transmit the disease effectively at any time of the year (e.g., in the drier plains of Western Australia), but in all areas during the winter months. In the late 1960s, in response to this problem the Australian government explored possibilities for introducing another vector for transmission, the European rabbit flea (*Spilopsyllus cuniculi*) that infests rabbits and does so year-round. *S. cuniculi* was, in general, an effective vector for transmitting the myxoma virus throughout the year, and was particularly effective in infecting young rabbits, which significantly reduced the population growth for the following year. However, *S. cuniculi* survived poorly in arid conditions, and in the early 1990s another flea species, *Xenopsylla cunicularis*, was introduced from Spain.

The introduction of vectors alternative to the naturally occurring mosquito was thus aimed primarily at solving what has been called *the problem of overwintering*. Due to vector reduction during the winter months, more virulent forms of the myxoma virus could not survive the winter: they killed their hosts before being spread to other hosts. Fenner and Ratcliffe

1. See Fenner and Ratcliffe (1965, chap.13, esp. 220–225). The core data is summarized in their Figure 16 (223), which is extended to include data up to 1967 in Table 7.2 in Fenner and Fantini (1999, 173). On Grade V myxoma (neuromyxoma), see also Fenner and Ratcliffe (1965, 163, Table 30).

(1965, 222–223) identify the occurrence of attenuated strains in geographically distinct areas as a reason to think that the virus was mutating within local populations rather than being vector-transmitted there, since it was unlikely that vectors could reach these areas from regions in which attenuation had already taken place (cf. Fenner and Fantinni 1999, 200–201, 300–301 on latent infections and overwintering). The switch in vectors met, however, with mixed and limited success as a way to control the rabbit population, a point I shall return to in the next section.

Finally, given that the attention commanded by the myxoma case amongst those interested in the mechanisms of *natural* selection derives in part from its “in the wild” status, it is relevant that the bulk of the long-term data reported is the result of the natural overwintering of the virus in rabbit populations *together with* supplemental “inoculation campaigns.” This program of human-induced infection was systematic and sustained, involving direct transmission of the virus by human beings (researchers and farmers) to over 20,000 rabbits on average in each of the twelve years following its original introduction (Fenner and Ratcliffe 1965, 292, Table 50). In addition, farmers intervened in another way in engaging their fondness for shooting and poisoning rabbits, which had become a major threat to their livelihood. Thus, human agents play a key role in the spread and maintenance of myxomatosis, not just through the selection of additional vectors, but also by serving as initiating vectors for transmission of the virus and in culling the (infected) rabbit population. This is what one would expect in a case that is, after all, a form of pest control. Whether these facts impugn the status of the myxoma case as one of group selection *in the wild*, I leave as an exercise for the reader.

5. Why Further Details Reinforce the Irresolvability Claim. A surprisingly large gap exists between the epidemiological literature on myxomatosis and the evolutionary debate that the phenomena it studies has inspired. The tendency in virology and epidemiology is to focus on the documented findings in natural populations of rabbits via sampling techniques, approaching the question of the mechanisms mediating changes here through laboratory studies of both rabbit and virus. There has been little explicit attention to the question of *just how* natural selection operates in this case. For example, Fenner and Ratcliffe attribute the reduction in virulence they document as “being due to mutations of the virus which were subsequently selected for” (1965, 222), but do not specify the selective mechanism. Later, in discussing the evolution of attenuated strains of the virus, they appeal to both the greater survival over the winter months of rabbits with attenuated strains of the virus and to lower temperatures during the winter as facilitating this survival (of both rabbits and virus). Again, the nature of the selective mechanism itself is not further

specified (1965, 341). In the extensive overview that Fenner and Fantini (1999) provide of the history of myxomatosis, this question of the level at which selection operates remains not simply unanswered, but (to a close approximation) unasked.

General textbook treatments of the example (e.g., Schmid-Hempel 1998, 251; cf. Ewald 1993, chap. 3) have sometimes taken the “classic view” of the evolution of lower levels of virulence to be group selection. I have found it difficult, however, to find such a view—indeed, any explicit view on the units of selection issue—in Fenner and Ratcliffe (1965) and Fenner and Fantini (1999). Those involved most directly in the field and laboratory work on the myxoma virus seem to have *individual* selection, if anything, in mind as the form that natural selection takes (cf. also Levine 1992, 209–210). This is largely because of the understandable emphasis on a range of factors that affect the spread of lower levels of virulence—host resistance, temperature, means of transmission—all discussed in terms of their effects on individual fitness.

Consider, by contrast, the literature on the levels of selection that appeals to the myxoma example. Typically describing the example in a short paragraph or so, the details sketched above are omitted. In particular, transmission by contagion is ignored, as are secondary lesion sites and preferential vector feeding, the effect of which is to highlight the significance of vector-mediated rabbit-to-rabbit transmission, i.e., transmission between groups of viruses. For group selectionists, population structure “pops out” as a salient cause of the spread of what is observed to evolve, i.e., lower levels of virulence. By contrast, although individualists ascribe a role to population structure, as we will see later in this section, they oppose this interpretation on quite general grounds, and see little need to appeal more than minimally to the details of the case, being concerned primarily to debunk the claims of group selectionists.

A range of putative examples of decisive empirical findings, actual or possible, are suggested by the details sketched in the previous section. These include those from studies that artificially increase (or decrease) the respective strengths of individual and group selection in an experimental group, and the examination of evolutionary patterns of viral replication both within groups (rabbits) and between groups. The basic problem with such scenarios being viewed as potentially decisive resolutions to the debate, however, is that it is fairly easy to construct explanations from either of these perspectives for whatever data is found or even merely envisaged. Lest this sound a little too *a priori* as a pronouncement of what we must find, let us consider a few good candidates for just the sort of decisive result we seek. Since the primary interest in the case is whether it constitutes a decisive test for group selection, I focus on results that might be taken to offer such decisive support.

For example, consider the basic finding that virulence increases within hosts but decreases over the whole population over time. Surely, one might think, this at least *prima facie* supports the group selectionist view, since that view posits opposed forces of selection at different levels that directly explain the within-and between-group trends. Yet an individualist should respond that while individual selection should indeed increase virulence, faced with the environmental circumstances of shorter-lived hosts, it will also drive lower levels of virulence. That is, rather than seeing individual and group selection as competing forces that pull in different directions, as does the proponent of group selection, the individualist views both forces as being at the individual level, each reflecting a different adaptive pressure that individuals face, with the one driving lower virulence winning out.

Likewise, consider experimental or naturalistic conditions that putatively reduce the effects of group selection and that thus lead to increases in levels of virulence. Lewontin 1970 had predicted that the introduction of the European rabbit flea, *S. cuniculi*, would increase virulence, presumably because doing so would effectively reduce the “groupishness” of viruses located on particular rabbits, and so limit the effects of group selection in promoting reduced levels of virulence by increasing its transmissibility between rabbits. *Prima facie*, such manipulations would support the group selectionist explanation, in much the way that the findings of Michael Wade’s (1977, 1978) celebrated *Tribolium* experiments that used an artificial group selection paradigm were taken to support group selection. In both cases, we manipulate factors that would adjust the strength of the force of group selection, were it to exist, and observe a phenotypic change that confirms the hypothesis that group selection is acting in the natural environment.

There are two problems with this idea, one pertaining to the inconclusiveness of the empirical data relevant to Lewontin’s particular suggestion, the other more general.

The first is that the effects of introducing an alternate vector on the virulence of the myxoma virus remain unclear. In part this is due to regional variation and the environmental sensitivities of this new vector. Studies at government sites around Canberra in 1977 and at Lake Urana in New South Wales in 1981 suggested that the introduction of rabbit fleas had little effect on the transmission of the myxoma virus, while results from South Australia in 1983 supported the effectiveness of the new vector in transmitting the virus (see Fenner and Fantini 1999, 183–185). Temperature and moisture are crucial variables here, and the myxoma virus continued to be transmitted by the pre-existing vector, the mosquito, in all three locations over the summer months.

The second and deeper problem is that even had the empirical results been more clear-cut, they could also have been explained from the

individualist's point of view. This is because the relevant manipulation changes the nature of the population-structured environment in which individual organisms exist, and with it the relationship between individuals and this environment. The explanation encapsulated in (1)–(3) would no longer apply, true enough, but that is because, in the new environmental circumstances, killing one's host is no longer detrimental to the fitness of the individual virus, and so low virulence loses the fitness advantage it had in the pre-flea environment. In the new environment, increased virulence has a higher level of fitness, and so we would expect it to be the trait that evolved.

Consider a third example. It is possible for a micro-organism to have a high reproductive rate without killing its host. *Escherichia coli* bacteria, for example, are extremely rapid reproducers in the human gut, but do not usually pose a threat to the life of the host organism. In the case of the myxoma virus, there could be a strain of the virus that approximated *E. coli* in this respect, somewhat like how myxoma actually operates in *Sylvilagus* hosts (cf. Fenner and Ratcliffe 1965, 346–47). For example, a viral strain that increased the number of lesions on a rabbit, thus increasing transmissibility, with reduced infection of vital organs in the host (say, due to host adaptation), would have this effect. Likewise, a strain that concentrated lesions on the head of the rabbit (where *S. cuniculi* concentrates, especially the ears) would have a similar transmission advantage. With an increase in transmissibility, we could expect that, *ceteris paribus*, levels of virulence would increase. I should like to make two points about the gap between this prediction and any conclusion about the level or levels at which such a selective force acts.

First, the intricacy of the relationships between host, vector, and virus, and how each of these are affected by environmental parameters that are subject to change, makes it unlikely that all other things *will* be equal. Second, plausible explanations could be provided for such an outcome from each of these perspectives. Whether such an increase in virulence brought about by increased transmissibility does so in virtue of selection operating directly on real population structures, groups of viruses, or does so in virtue of its operation on an individual's sensitivity to changes in its environment, would remain open.

These examples indicate the sorts of problems with the appeal to decisiveness. There is a principled reason why the problem here is general. We have two paradigms of evolutionary explanation, that of group selection and that of individual selection. Each has a basic repertoire of tools that can be adapted to much the same data sets in the myxoma case. Both can acknowledge the reality of a group structure (or a variety of group structures) within the overall virus population, and each accommodates this fact in its own way. The group selectionist sees natural selection as

operating on the groups themselves, eliminating those groups that are less fit, which are groups whose members have high levels of virulence. The individual selectionist, by contrast, sees natural selection as operating on individuals sensitive to features of their environment, including the group structures imposed by the nature of the hosts and vectors present for transmission. If the resources of each view are rich enough to explain *any* putatively decisive result, as I have been suggesting, then this provides support for my second conclusion, that the myxoma case is in fact *irresolvable* vis-à-vis the question of how selection is really operating in it.²

If this is so, then it might seem a short step down the path to generalizations of this conclusion, perhaps even to the debate over group selection itself. But as I implied in the introduction, even this “short step” involves tasks that lie beyond the current paper. As a way of reinforcing my conclusion about the irresolvability of the myxoma case, and to move to some broader issues that this conclusion itself raises, I focus next on a metaphor occasionally made explicit in the myxoma debate and otherwise often not far beneath the surface of the debate. This is the metaphor of particular units of selection as having *standpoints* or *viewpoints*.

6. Decisiveness and Viewpoints. Much of the myxoma debate can be cast in terms of talk of the *viewpoints* of particular organisms, and indeed has been (e.g., Lewontin 1970, 15; Sober and Wilson 1998, 45). The conception of evolution in terms of the *selfish gene* has perhaps made most effective use of such viewpoint talk in evolutionary biology more generally (see Okasha 2002 for a recent example).

The primary viewpoint adopted in the debate over myxomatosis is that of the virus. Group selectionists hold that the viewpoint of the group of viruses that live on a rabbit is necessary for understanding, or sheds light on, just how natural selection operates in the myxoma case. We could encapsulate the concern that individualists have about the group selectionist account in terms of whether the viewpoint of virus groups adds anything to our understanding of the phenomena to that provided by the viewpoint of individual viruses. I propose to use this metaphor in conjunction with the details sketched in sections 4 and 5 to raise a series of further complications.

The first is that the viewpoints of viruses (individuals or groups) are not independent of those of rabbits, especially given that the central concept in play, virulence, is often characterized in terms of effects of the

2. The reasoning here bears a similarity, and perhaps a debt, to that in Sterelny and Kitcher's 1988 response to the argument of Sober and Lewontin 1982 that claimed that heterozygote superiority in the case of sickle cell anemia could not be explained by genic selection.

virus on the well being of rabbits. The second is that even these four viewpoints are not exhaustive, for there is that also of the *vectors*, as well as that of a nested hierarchy of groups, especially of viruses. As I implied in Section 4, if rabbits constitute a group of viruses, then surely (given contagious infection) a *hutch* of rabbits does as well, as does the *local deme* of rabbit hutches. And given the differential transmissibility of viruses located on different parts of the rabbit, all those viruses on the rabbit's head, or its ear, or around its eyes, also constitute groups of viruses. Note that here a variation on the arbitrariness problem from Section 3 arises, for although rabbit-bound viruses are not an evolutionary arbitrary group, they are far from a unique such group, and the focus just on rabbit-bound viruses as the object of group selection seems either arbitrary or in need of further justification.

On the group selectionist view, within each of these groups individual selection should promote increased virulence, while group selection should promote decreased virulence. Individual selectionists, of course, have their own account of how to understand this added complexity. The third complication, in light of the first two, is that there seems to be no fact of the matter as to which viewpoint is (or viewpoints are) *the* correct or preferred viewpoint(s) for thinking about myxomatosis.

Consider now the viewpoint not of the virus but of the *rabbit*. When researchers found that rabbits had increased their resistance to the myxoma virus in the years following its introduction, this result was readily intelligible within the parameters of the theory of individual selection. What would have been paradoxical, in much the way as was the discovery of lower levels of virulence, would have been the finding that rabbits had *decreased* their levels of resistance. Although this was *not* found, I want to explore how this possible finding would be explained from each side of the debate over the myxoma case.

One initial thought is that if high resistance evolves by individual selection, then low resistance should evolve by *group* selection. (After all, this would parallel what has been said when considering the viewpoint of the virus.) To make the explanation of this explicit, consider the following argument sketch:

- (A) High resistance has a higher level of fitness for an individual rabbit.
- (B) But highly resistant rabbits spread the disease throughout the population.

Thus,

- (C) Groups of low resistance rabbits are fitter than groups of high resistance rabbits.

So,

(D) Low resistance evolves by group selection.

Note that (A)–(D) provide an account of the evolution of low resistance in rabbits that closely parallels the group selectionist account of the evolution of low levels of virulence in the virus. In both cases, a trait that reduces within-group fitness nevertheless evolves because of group selection.

Low resistance would not, however, provide decisive support for the theory of group selection, since there is another explanation available that appeals only to *individual* selection. Again, to parallel the explanation of the evolution of low levels of virulence encapsulated in (1)–(3), consider the following individual-level explanation:

(1*) A rabbit killed by a viral strain is less fit than one not so killed.

(2*) Low resistance reduces the chances of being killed by a viral strain.

Thus,

(3*) Low resistance is an adaptation of individual rabbits.

So,

(4*) Low resistance evolves by individual selection.

Like the explanation of the evolution of avirulence by an appeal to individual selection, this explanation need not commit the averaging fallacy. Unlike that explanation, at least one of its premises seems strangely counter-intuitive and stands in need of special justification.

This is (2*): how could *low* resistance increase one's fitness by reducing one's chance of parasite-induced death? Here an appeal to the sensitivity of individual rabbits to the nature of their environment needs to be made explicit. Given that the virus itself varies in its virulence, if the contraction of less virulent strains were to provide some immunization against the contraction of more virulent strains, then given other environmental conditions (e.g., the prevalence of less virulent strains) lower resistance could promote longevity in rabbits that have it. This is how other *pox* viruses operate on humans, and why having chickenpox as a child is not altogether a bad thing. Fenner and Ratcliffe (1965, 233–235) in fact present some evidence for just this sort of effect in myxomatosis, though they remain neutral as to whether this is caused by an interference effect between competing strains or a heightened immune response from the host.

Alternatively, (2*) might be true not in virtue of the structure of the viral environment and its interaction with individual rabbits, but because

of facts about *vectors* and how they interact with rabbits with various levels of resistance. Infected rabbits may have less appeal for arthropod vectors than do uninfected rabbits, and so lower resistance may be a strategy that individual rabbits adopt in order to limit their exposure to highly virulent strains of the virus.

The counterfactual nature of the “finding” of the evolution of low resistance in rabbits should make it clear that this is intended as part of a “how possibly” rather than a “how actually” story. The interest of such a Just So Story is twofold in the present context.

First, it reminds us that we need to consider the full environment of an individual organism, including the nature of both its parasites and whatever group structure they have, as well as that of the vectors that mediate between host and parasite. In so doing, it draws attention to the toolkit of an individualist who appeals to the sensitivity of an individual to its (complicated) environmental circumstances. This reinforces the conclusion that such an individualist explanation will likely be available, once the empirical details of any such case are filled in, and so adds to the conclusions for which I have already argued.

Second, this just-so story leads us fairly naturally into another, one that reminds us that there remains a further viewpoint, that of the gene. For we can now consider the following explanation of the “evolution of low resistance”:

- (1+) Genes whose vehicles are killed by viral strains in E are less fit than those in vehicles not so killed.
- (2+) Genes for low resistance in E reduce the chances of one’s vehicle being killed by viral strains.

Thus,

- (3+) Low resistance is a genetic adaptation.

So,

- (4+) Low resistance evolves by genic selection.

(1+)–(4+) make explicit the thus-far implicit appeal to an environment. I suggest that the genic selectionist defending (2+) has at least as rich a repertoire of tools as has the individualist defending (2*).

If this is correct, then the basic problem for resolving the debate over the myxoma case is compounded, my conclusion about its likely irresolvability is further reinforced, and we have some additional reason to think that the extension of this conclusion beyond the myxoma case more generally is defensible. But this also raises two broader issues about the nature of selection and how we conceptualize it. To raise these issues as provocative questions: Doesn’t the irresolvability of the debate over the

myxoma virus, and perhaps the levels of selection more generally, mean that the viewpoint of the individual and that of the group are simply two equivalent ways of viewing natural selection? And does the irresolvability claim here call into question the very conception of selection as operating at distinct levels? I shall say something briefly about these issues in closing.

7. Conclusion. In the introduction I noted that pluralism has some vogue as a position about the levels of selection. A prominent form of pluralism that has proven especially popular amongst biologists and philosophers of biology holds that there is an important sense in which different models of selection are equivalent, such that the choice between them is to be made chiefly on heuristic or pragmatic grounds (see Dugatkin and Reeve 1994; Reeve 2000; Bourke and Franks 1995; Crozier and Pamilo 1996; Frank 1998; Michod 1999; Kerr and Godfrey-Smith 2002; Dugatkin 2002; Maynard 2002; Sober and Wilson 2002). This view has been called *model pluralism*, since it adopts a pluralistic stance toward various models of selection, including individual and group selection models (Wilson 2003).

Model pluralism provides a seemingly natural view to adopt if we accept either the specific irresolvability claim I have argued for about the myxoma case, or the more general irresolvability claim that I have groped toward. But despite appearances, the position I have defended in this paper does not have a deep affinity with model pluralism, and does not rest on it.

What I share with model pluralists is the idea that there is some kind of interdependence between the various putative “levels of selection.” Model pluralists locate this interdependence in equivalences between the different models of natural selection, equivalences that provide a means for translating between different explanatory frameworks. This constitutes the basis for their deflationary view of disputes between proponents of one or another view of “the” level of selection. By contrast, on my view a deep, ontological difference remains between (say) proponents of individual and group selection that concerns what sort of biological agents drive natural selection in particular cases. It is just that the structure of the biological world itself thwarts our attempts to resolve the resulting disagreement.

Another way to understand the difference here is to consider the perspective that each view adopts towards the very idea of biological reality being structured into distinct levels. Model pluralists typically take the level metaphor for granted and call into question the significance of the distinctness of various levels. But I view the metaphor of levels itself is suspect. Biological reality is more aptly conceived of as fused, enmeshed, or *entwined*, rather than hierarchically structured into neat levels, in that

the properties on which selection acts, and indeed the mechanism of selection itself, do not come prepackaged at distinct levels. Conceiving of natural selection as operating at various “levels” is a simplification of the entwined, messy reality, a conception that imposes structure on, rather than simply reflects, biological reality. Reduced virulence in the myxoma virus can be viewed either as an adaptation of individual viruses or as a product of inter-group competition not because we can translate between different models that correspond to different levels of selection, but because biological reality is more complicated than our levels metaphor recognizes (see also Wilson 2004).

The evolution of the myxoma virus in natural populations of rabbits in Australia is less conclusive as an example of group selection than proponents of group selection think, but not because their individualistic critics are correct in their analysis of the example. Rather, the debate here stands as an example of a scientific dispute that cannot be resolved by the current evidence, and I have argued that it seems unlikely to yield to further evidence (cf. Okasha 2001). Reflection on some of the complexities to the phenomenon of myxomatosis reinforce rather than ameliorate the irresolvability of the myxoma case. Thus, there are reasons to think that the scientific dispute here cannot be rationally resolved in a conclusive manner.

This conclusion almost certainly generalizes to other cases invoked in the debate over the levels of selection. While one might think that it generalizes to the whole debate between proponents of individual and group selection, that would require significantly more and I think a somewhat different argument than I have provided here. Those who endorse such a view of the general debate over the levels of selection may be tempted by model pluralism, but this is not my temptation. Rather, I think that the argument here raises questions about the adequacy of the very conception of natural selection operating at distinct levels, a conception ubiquitous in debates over how natural selection operates. If this is correct, then perhaps we need to rethink some large questions in the field.

REFERENCES

- Alexander, Richard, and Garcia Borgia (1978), “Group Selection, Altruism, and the Levels of Organization of Life”, *Annual Review of Ecological Systematics* 9: 449–474.
- Bourke, Andrew F.G., and Nigel R. Franks (1995), *Social Evolution in Ants*. Princeton, NJ: Princeton University Press.
- Crozier, Ross H., and Pekka Pamilo (1996), *Evolution of Social Insect Colonies*. New York: Oxford University Press.
- Dugatkin, Lee (2002), “Will Peace Follow?”, *Biology and Philosophy* 17: 519–522.
- Dugatkin, Lee, and Hudson K. Reeve (1994), “Behavioral Ecology and Levels of Selection: Dissolving the Group Selection Controversy”, in Peter J.B. Slater et al. (eds.), *Advances in the Study of Behavior*, vol. 23. New York: Academic Press, 101–133.
- Ewald, Paul W. (1993), *The Evolution of Infectious Disease*. New York: Oxford University Press.

- Fenner, Frank, and Bernadino Fantini (1999), *Biological Control of Vertebrate Pests: The History of Myxomatosis—an Experiment in Evolution*. Oxford: CABI Publishing.
- Fenner, Frank, and F.N. Ratcliffe (1965), *Myxomatosis*. Cambridge: Cambridge University Press.
- Frank, Steve A. (1998), *Foundations of Social Evolution*. Princeton, NJ: Princeton University Press.
- Futuyma, Douglas J. (1979), *Evolutionary Biology*. Sunderland, MA: Sinauer.
- Kerr, Benjamin, and Peter Godfrey-Smith (2002), “Individualist and Multi-Level Perspectives on Selection in Structured Populations”, *Biology and Philosophy* 17: 477–517.
- Lack, David (1954), *The Natural Regulation of Animal Numbers*. New York: Oxford University Press.
- Levine, Andrew J. (1992), *Viruses*. New York: Scientific American Library.
- Lewontin, Richard (1970), “The Units of Selection”, *Annual Review of Ecology and Systematics* 1: 1–18.
- Maynard Smith, John (2002), “Commentary on Kerr and Godfrey-Smith” *Biology and Philosophy* 17: 523–527.
- Michod, Richard (1999), *Darwinian Dynamics*. Princeton, NJ: Princeton University Press.
- Okasha, Samir (2001), “Why Won’t the Group Selection Controversy Go Away?” *British Journal for the Philosophy of Science* 52: 25–50.
- (2002), “Genetic Relatedness and the Evolution of Altruism”, *Philosophy of Science* 69: 138–149.
- Parer, Ian P. (1995), “Relationship Between Survival Rate and Survival Time of Rabbits, *Oryctolagus cuniculus* (L.) Challenged with Myxoma Virus”, *Australian Journal of Zoology* 43: 303–311.
- Reeve, Hudson K. (2000), “Multi-Level Selection and Human Cooperation”, *Evolution and Human Behavior* 21: 65–72.
- Schmid-Hempel, Paul (1998), *Parasitism in Social Insects*. Princeton, NJ: Princeton University Press.
- Sober, Elliott (1984), *The Nature of Selection*. Cambridge, MA: MIT Press.
- (1993), *Philosophy of Biology*. Boulder, CO: Westview Press.
- Sober, Elliott, and Richard Lewontin (1982), “Artifact, Cause, and Genic Selection”, *Philosophy of Science* 49: 157–198.
- Sober, Elliott, and David Sloan Wilson (1994), “A Critical Review of Philosophical Work on the Units of Selection Problem”, *Philosophy of Science* 61: 534–555.
- (1998), *Unto Others: The Evolution and Psychology of Unselfish Behavior*. Cambridge, MA: Harvard University Press.
- (2002), “Perspectives and Parameterizations: Commentary on Benjamin Kerr and Peter Godfrey-Smith’s ‘Individualist and Multi-Level Perspectives on Selection in Structured Populations’”, *Biology and Philosophy* 17: 529–537.
- Sterelny, Kim (1996), “The Return of the Group”, *Philosophy of Science* 63: 562–84.
- Sterelny, Kim, and Philip Kitcher (1988), “The Return of the Gene”, *Journal of Philosophy* 85: 339–361.
- Wade, Michael J. (1977), “An Experimental Study of Group Selection”, *Evolution* 31: 134–153.
- (1978), “A Critical Review of the Models of Group Selection”, *Quarterly Review of Biology* 53: 101–114.
- Wilson, David Sloan (1975), “A Theory of Group Selection”, *Proceedings of the National Academy of Sciences USA* 72: 143–146.
- (1980), *The Natural Selection of Populations and Communities*. Menlo Park, CA: Benjamin Cummins.
- (1983), “The Group Selection Controversy: History and Current Status”, *Annual Review of Ecology and Systematics* 14: 159–187.
- (1989), “Levels of Selection: An Alternative to Individualism in the Human Sciences”, *Social Networks* 11: 257–272. Reprinted in Elliott Sober (ed.), *Conceptual Issues in Evolutionary Biology*. 2d ed.. Cambridge, MA: MIT Press, 143–154.

- (1997), “Altruism and Organism: Disentangling the Themes of Multilevel Selection Theory”, *American Naturalist* 150 (supp.): S122–S134.
- (2002), *Darwin’s Cathedral: Evolution, Religion, and the Nature of Society*. Chicago: University of Chicago Press.
- Wilson, David Sloan, and Elliott Sober (1994), “Reintroducing Group Selection to the Human Behavioral Sciences”, *Behavioral and Brain Sciences* 17: 585–654.
- Wilson, Robert A. (2003), “Pluralism, Entwinement, and the Levels of Selection”, *Philosophy of Science* 70: 531–552.
- (2004), *Genes and the Agents of Life: The Individual in the Fragile Sciences: Biology*. New York: Cambridge University Press.