



National Library  
of Canada

Bibliothèque nationale  
du Canada

Canadian Theses Service

Services des thèses canadiennes

Ottawa, Canada  
K1A 0N4

## CANADIAN THESES

## THÈSES CANADIENNES

### NOTICE

The quality of this microfiche is heavily dependent upon the quality of the original thesis submitted for microfilming. Every effort has been made to ensure the highest quality of reproduction possible.

If pages are missing, contact the university which granted the degree.

Some pages may have indistinct print especially if the original pages were typed with a poor typewriter ribbon or if the university sent us an inferior photocopy.

Previously copyrighted materials (journal articles, published tests, etc.) are not filmed.

Reproduction in full or in part of this film is governed by the Canadian Copyright Act, R.S.C. 1970, c. C-30. Please read the authorization forms which accompany this thesis.

**THIS DISSERTATION  
HAS BEEN MICROFILMED  
EXACTLY AS RECEIVED**

### AVIS

La qualité de cette microfiche dépend grandement de la qualité de la thèse soumise au microfilmage. Nous avons tout fait pour assurer une qualité supérieure de reproduction.

S'il manque des pages, veuillez communiquer avec l'université qui a conféré le grade.

La qualité d'impression de certaines pages peut laisser à désirer, surtout si les pages originales ont été dactylographiées à l'aide d'un ruban usé ou si l'université nous a fait parvenir une photocopie de qualité inférieure.

Les documents qui font déjà l'objet d'un droit d'auteur (articles de revue, examens publiés, etc.) ne sont pas microfilmés.

La reproduction, même partielle, de ce microfilm est soumise à la Loi canadienne sur le droit d'auteur, SRC 1970, c. C-30. Veuillez prendre connaissance des formules d'autorisation qui accompagnent cette thèse.

**LA THÈSE A ÉTÉ  
MICROFILMÉE TELLE QUE  
NOUS L'AVONS REÇUE**

**Canada**



National Library of Canada

Bibliothèque nationale du Canada

Ottawa, Canada  
K1A 0N4

TC -

0-315-22948-9

CANADIAN THESES ON MICROFICHE SERVICE - SERVICE DES THÈSES CANADIENNES SUR MICROFICHE

PERMISSION TO MICROFILM - AUTORISATION DE MICROFILMER

Please print or type - Écrire en lettres moulées ou dactylographier

AUTHOR - AUTEUR

Full Name of Author - Nom complet de l'auteur

Glen Douglas Wheeler

Date of Birth - Date de naissance

March 8, 1958

Canadian-Citizen - Citoyen canadien

Yes / Oui

No / Non

Country of Birth - Lieu de naissance

Canada

Permanent Address - Résidence fixe

10552 - 66 Ave  
Edmonton, Alberta  
T6H5R3 1X4

THESIS - THÈSE

Title of Thesis - Titre de la thèse

Immunolocalization of kallikrein  
in various tissues

Degree for which thesis was presented  
Grade pour lequel cette thèse fut présentée

Ph. D.

Year this degree conferred  
Année d'obtention de ce grade

1985

University - Université

Univ. of Alberta

Name of Supervisor - Nom du directeur de thèse

Dr. M. Schachter

AUTHORIZATION - AUTORISATION

Permission is hereby granted to the NATIONAL LIBRARY OF CANADA to microfilm this thesis and to lend or sell copies of the film.

L'autorisation est, par la présente, accordée à la BIBLIOTHÈQUE NATIONALE DU CANADA de microfilmer cette thèse et de prêter ou de vendre des exemplaires du film.

The author reserves other publication rights, and neither the thesis nor extensive extracts from it may be printed or otherwise reproduced without the author's written permission.

L'auteur se réserve les autres droits de publication; ni la thèse ni de longs extraits de celle-ci ne doivent être imprimés ou autrement reproduits sans l'autorisation écrite de l'auteur.

ATTACH FORM TO THESIS - VEUILLEZ JOINDRE CE FORMULAIRE À LA THÈSE

Signature

*Glen Wheeler*

Date

2 Oct, 1985

THE UNIVERSITY OF ALBERTA

IMMUNOLOCALIZATION OF KALLIKREIN IN VARIOUS TISSUES

by

GLEN D. WHEELER

A THESIS

SUBMITTED TO THE FACULTY OF GRADUATE STUDIES AND RESEARCH  
IN PARTIAL FULFILMENT OF THE REQUIREMENTS FOR THE DEGREE  
OF DOCTOR OF PHILOSOPHY

DEPARTMENT OF PHYSIOLOGY

EDMONTON, ALBERTA

FALL 1985

THE UNIVERSITY OF ALBERTA

RELEASE FORM

NAME OF AUTHOR GLEN D. WHEELER

TITLE OF THESIS IMMUNOLOCALIZATION OF KALLIKREIN IN VARIOUS TISSUES


DEGREE FOR WHICH THESIS WAS PRESENTED DOCTOR OF PHILOSOPHY

YEAR THIS DEGREE GRANTED FALL 1985

Permission is hereby granted to THE UNIVERSITY OF ALBERTA LIBRARY to reproduce single copies of this thesis and to lend or sell such copies for private, scholarly or scientific research purposes only.

The author reserves other publication rights, and neither the thesis nor extensive extracts from it may be printed or otherwise reproduced without the author's written permission.

(SIGNED)



PERMANENT ADDRESS:

10552 - 66 Avenue

Edmonton, Alberta, Canada

T6H 1X4

DATED Sept 27 19 85

THE UNIVERSITY OF ALBERTA  
FACULTY OF GRADUATE STUDIES AND RESEARCH

The undersigned certify that they have read, and recommend to the Faculty of Graduate Studies and Research, for acceptance, a thesis entitled IMMUNOLOCALIZATION OF KALLIKREIN IN VARIOUS TISSUES submitted by GLEN D. WHEELER in partial fulfilment of the requirements for the degree of DOCTOR OF PHILOSOPHY.

*M. Schacter*  
.....

Supervisor

*R. Kaye*  
.....

*Suzanne Barton*  
.....

*E. J. Sanders*  
.....

*Ronald Winton*  
.....

*A. Murray Perle*  
.....

External Examiner

Date *Sept. 27, 1985*

**DEDICATION**

To My Parents

## ABSTRACT

Many techniques were employed for the immunocytochemical localization of tissue kallikreins (EC 3.4.21.35) and nerve growth factor (NGF). Freeze-drying and chemical fixatives were successful but to varying degrees. Resin-embedded tissues were better preserved and yielded better results than paraffin embedding. Two methods of immunostain were most successful, namely peroxidase-antiperoxidase (PAP) and avidin-biotin complex (ABC).

Tissue kallikrein was immunocytochemically localized in various mammalian tissues, including the submandibular gland of cat and guinea-pig where it was localized in apical regions of striated duct cells. It was also localized in the goblet or mucous cells of the colon and in the mucous cells of the stomach of the cat and rat. At the ultrastructural level, it was localized in the small apical secretory granules of striated duct cells of the submandibular gland in cat and guinea-pig. Kallikrein was also immunolocalized in surface epithelial secretory cells of the guinea-pig coagulating and prostate glands; also in secretory cells of the submucosa of cat trachea. Nerve growth factor (NGF) was immunolocalized in the surface epithelial cells in the guinea-pig prostate gland.

The biological and evolutionary significance of the broad group of serine proteases is discussed. It is suggested that these enzymes play multiple and diverse roles in bioregulatory processes, and these various possibilities are considered and discussed.

## ACKNOWLEDGEMENTS

To Dr. M. Schachter for being a great spirit.

I also thank Dr. R.W. Matthews, Dr. M. Peret, Dr. S. Barton, Mr. D.J. Longridge, Mr. K. Burt, Mrs. T. Labeledz and Mrs. O.M. Findlay.

I am also thankful to the Alberta Heritage Foundation for Medical Research and the Medical Research Council of Canada for support of this project.



Chapter	Table of Contents	Page
I. INTRODUCTION .....		1
A. Kallikrein-Kinin System .....		1
1. Salivary gland .....		2
2. Pancreas .....		6
3. Colon .....		7
4. Intestine .....		8
3. Stomach .....		9
6. Kidney (Urinary) .....		9
7. Respiratory Tract .....		12
8. Male Sex Glands .....		12
9. Brain .....		13
10. Other Tissues .....		13
B. Nerve Growth Factor .....		14
II. MATERIALS AND METHODS .....		17
A. Animals, Organ Dissection, Tissue Preparation and Extracts .....		17
B. Fixation of Tissues .....		18
1. Chemical Fixation .....		18
2. Freeze-Drying .....		22
C. Embedding .....		23
D. Immunocytochemistry .....		26
1. Antisera .....		26
2. Immunostaining systems .....		29
a. Peroxidase-antiperoxidase .....		29
i. Paraffin sections .....		29
ii. Semi-thin resin sections .....		32
iii. Ultrathin resin sections .....		33

b. Avidin-biotin complex .....	35
i. Semi-thin resin sections .....	35
ii. Ultrathin resin sections .....	38
iii. Pre-embedding .....	39
c. Protein A-colloidal gold .....	40
i. Ultrathin resin sections .....	40
3. Specificity and controls .....	40
E. Microscopy and Photography .....	41
III. RESULTS .....	42
A. Summarized Results of Immunostaining .....	42
B. Localization of Kallikrein .....	42
1. Submandibular Gland - Light and Electron Microscopy .....	42
a. Cat .....	42
b. Guinea-pig .....	57
2. Colon .....	57
a. Cat .....	57
b. Rat .....	67
3. Stomach .....	72
a. Cat .....	72
b. Rat .....	81
4. Respiratory Tract .....	84
a. Cat .....	84
C. Localization of Kallikrein and Nerve Growth Factor .....	91
1. Male Sex Glands - Light Microscopy .....	91
a. Guinea-pig .....	91
IV. DISCUSSION .....	103
A. Kallikrein and Related Proteases in Salivary Glands .....	103

B. Kallikreins, Nerve Growth Factor and Related Proteases in Male Sex Glands and Their Possible Roles .....	106
C. Gastrointestinal Kallikreins and Their Possible Roles .....	107
D. Interpretation of Immunocytochemical Findings .....	108
E. Possible Sources of Circulating Tissue Kallikrein .....	111
V. CONCLUSION .....	112
VI. REFERENCES .....	113

## LIST OF TABLES

Table	Description	Page
1	Summarized results of light microscopic immunostained tissues fixed in various ways	43

## LIST OF FIGURES

Figure	Description	Page
1	Low temperature freeze drier and copper block freezing device	24-25
2	Double diffusion analysis of $\alpha$ CSK with cat colon and cat submandibular gland	27-28
3	Summarized PAP method	30-31
4	Summarized ABC method	36-37
5	Kallikrein localization in striated duct cells of the cat submandibular gland (paraffin sections)	44-45
6	Control section for Figure 5	44-45
7	Kallikrein localization in apical regions of striated duct cells of the cat submandibular gland (semi-thin sections)	46-47
8	Control section for Figure 7	46-47
9	Electron microscopic localization of kallikrein in apical secretory granules of the cat submandibular gland, striated duct cells	49-50
10	Control section for Figure 9	49-50

11	Ultrastructure of apical region of striated duct cells fixed by strong aldehyde fixative	51-52
12	Ultrastructure of apical region of striated duct cells fixed by weak immunocytochemical fixative	51-52
13	Localization of kallikrein in apical granules, stained as in Figure 9, fixed as in Figure 12	51-52
14	Kallikrein localization in apical regions of striated duct cells of the cat submandibular gland, post-fixation with osmium (semi-thin sections)	53-54
15	Control section for Figure 14	53-54
16	Electron microscopic localization of kallikrein in apical secretory granules of the cat submandibular gland, striated duct cells, post-fixed with osmium	55-56
17	Control section for Figure 16	55-56
18	Electron microscopic localization of kallikrein in apical secretory granules of the guinea-pig submandibular gland, striated duct cells	58-59
19	Control section for Figure 18	58-59
20	Kallikrein localization in mucous cells and secretions of the cat colon (paraffin sections)	61-62
21	Control section for Figure 20	61-62

22	Kallikrein localization in mucous cells and secretions of the cat colon (semi-thin sections)	63-64
23	Control section for Figure 22	63-64
24	Kallikrein localization in mucous cells and secretions of the cat colon (freeze-dried tissue, semi-thin sections)	65-66
25	Control section for Figure 24	65-66
26	Electron microscopic localization of kallikrein in the mucous granule region of cat colon mucous cells	68-69
27	Control section for Figure 26	68-69
28	Kallikrein localization in mucous cells of the rat colon	70-71
29	Control section for Figure 28	70-71
30	Kallikrein localization in apical regions of mucous cells of the cat stomach antrum (osmium post-stain)	73-74
31	Control section for Figure 30	73-74
32	Kallikrein localization in apical regions of mucous cells of the cat stomach antrum (without osmium post-stain)	75-76
33	Control section for Figure 32	75-76

34	PAS stain with immunostain for kallikrein in cat stomach antrum (high magnification)	77-78
35	Kallikrein localization in periphery of mucous cells of cat stomach antrum	77-78
36	Control section for Figures 34, 35 and 37	77-78
37	PAS stain with immunostain for kallikrein in cat stomach antrum (low magnification)	77-78
38	Kallikrein localization in mucous cells of cat stomach antrum (PAP method)	79-80
39	Control section for Figure 38	79-80
40	Kallikrein localization in mucous cells and secretions of cat stomach antrum (ABC method)	79-80
41	Control section for Figure 40	79-80
42	Kallikrein localization in mucous neck cells of the rat stomach antrum (paraffin sections)	82-83
43	Control section for Figure 42	82-83
44	Kallikrein localization in mucous cells and secretions of the rat stomach antrum (semi-thin sections)	85-86
45	Control section for Figure 44	85-86



46	Kallikrein localization in apical regions of mucous cells of the rat stomach antrum (without treatment)	87-88
47	Kallikrein localization in apical regions of mucous cells of the rat stomach antrum (EAI treatment)	87-88
48	Kallikrein localization in apical regions of mucous cells of the rat stomach antrum (PVPP treatment)	87-88
49	Kallikrein localization in submucosal cells of cat respiratory tissue	89-90
50	Control section for Figure 49	89-90
51	Kallikrein localization in secretory epithelial cells of the guinea-pig coagulating gland	92-93
52	Control section for Figure 51	92-93
53	Kallikrein localization in secretory epithelial cells of the guinea-pig coagulating gland (freeze-dried)	94-95
54	Control section for Figure 53	94-95
55	Kallikrein localization in secretory epithelial cells of the guinea-pig coagulating gland (PVPP treatment)	96-97
56	Control section for Figure 55	96-97

57	Kallikrein localization in apical cytoplasm and secretions of the guinea-pig prostate gland	99-100
58	Control section for Figure 57	99-100
59	NGF localization in apical cytoplasm and secretions of the guinea-pig prostate gland	101-102 ○
60	Control section for Figure.59	101-102

## LIST OF ABBREVIATIONS AND SYMBOLS

ABC	Avidin-biotin complex
$\alpha$	Denotes antiserum
$\alpha$ -	Denotes subunit of NGF
$\beta$ -	Denotes subunit of NGF
$\gamma$ -	Denotes subunit of NGF
BG	Biotinylated goat anti-rabbit immunoglobulin
BSA	Bovine serum albumin
CDI	1 ethyl-3-(3-dimethylaminopropyl) carbodiimide HCl
CGK	Guinea-pig coagulating gland kallikrein
CSK	Cat salivary kallikrein
DAB	3,3' diaminobenzidine tetrahydrochloride
DMS	Dimethylsuberimidate
EAI	Ethyl acidimidate
EDTA	Ethylene diaminetetraacetate
EGF	Epidermal growth factor
EPTD	Edwards-Pearse tissue drier

FAB	Goat serum immunized against rabbit immunoglobulin G Fab fragments
GMA	Glycol methacrylate
GPCG	Guinea-pig coagulating gland
GPPG	Guinea-pig prostate gland
GPSK	Guinea-pig salivary kallikrein
GTA	Glutaraldehyde
HE	Hematoxylin and eosin
HSK	Human salivary kallikrein
ml	Milliliter, $10^{-3}$ liter
mm	Millimeter, $10^{-3}$ meter
mOsm	Milliosmol
$\mu$ l	Microliter, $10^{-6}$ liter
$\mu$ m	Micrometer, $10^{-6}$ meter
NGF	Nerve growth factor
nm	Nanometer, $10^{-9}$ meter
Osmium	Osmium tetroxide
PAG	Protein A-colloidal gold complex

PAP	Rabbit peroxidase-antiperoxidase
PAS	Periodic acid-Schiff reagent
PBS	Phosphate-buffered saline
PEG	Polyethylene glycol
PFA	Paraformaldehyde
PLP	Periodate-lysine-paraformaldehyde
PVPP	Polyvinylpyrrolidone
RIK	Rat intestinal kallikrein
RPK	Rat pancreatic kallikrein
SDS	Sodium dodecylsulfate
Tris	Trizma base
Triton	Triton X-100 detergent

## I. INTRODUCTION

### A. Kallikrein-Kinin System

Tissue kallikreins are a group of serine proteases with common chemical properties and limited proteolytic activity. They release an active peptide from a precursor, or substrate, the peptide being bradykinin, or lysyl bradykinin, the latter also being known as kallidin (see Schachter, 1969, 1980). They release lysyl bradykinin from kininogen, a substrate that is present in lymph, plasma, and interstitial fluid and are, therefore, also called kininogenases. They have effects on blood vessels and smooth muscles *in vivo* that are indirect, i.e. due to the release of the pharmacologically active kinin (lysyl bradykinin). Physiologically significant proteolytic actions of these enzymes may, however, sometimes be unrelated to kinin release (Beraldo *et al.*, 1956; Chao *et al.*, 1981).

The kallikrein story begins in 1909 when Abelous and Bardier described a substance from human urine which lowered blood pressure. The presence of this hypotensive substance in urine was confirmed and it was found to be nondialyzable (Frey and Kraut, 1926). Thought to originate from the pancreas, it was named kallikrein (Greek, "kallikreas" = pancreas) by Kraut *et al.* (1930). The pancreas was later discounted as a major source of the urinary agent since kallikrein levels in dog urine remained high following pancreatectomy or pancreatic vessel occlusion (Beraldo *et al.*, 1956).

Plasma kallikrein differs from the tissue varieties on many grounds. It has a higher molecular weight, resembles trypsin in its substrate requirements, the latter being less specific, and releases bradykinin instead of lys-bradykinin (Colman and Bagdasarian, 1976). The plasma kallikrein-kinin system has been implicated in various pathological states such as inflammation and shock (Colman and Wong, 1979).

The splitting of the arginine-arginine bond by these enzymes indicates that they may be involved in processing prohormones, proenkephalin being a good example (Prado *et al.*, 1983). Kallikrein has also been reported to activate proinsulin (ole-Moi Yoi *et al.*, 1979b) and prorenin

(Sealey *et al.*, 1978).

Measurement of kallikreins is commonly performed chemically by the hydrolysis of various synthetic esters. Bioassay is also performed by recording hypotensive activity of the substance on the dog blood pressure or by measuring the substance's ability to release kinin from a substrate. Aprotinin (or Kunitz's trypsin inhibitor) inhibits tissue kallikrein, which results in prevention of kinin production.

Localization techniques are widely used and allow proteins to be identified and characterized at their sites of synthesis, storage and secretion. Several immunocytochemical techniques have been used to localize kallikrein in various tissues. Frequently, however, immunolocalizations by different authors produced different results. The purpose of this thesis is to immunolocalize tissue kallikrein in various tissues. The diverse cellular and subcellular locations of these enzymes indicate that they have different functions in different organs despite having similar chemical and pharmacological properties (Schachter, 1980).

In this thesis, several related kallikreins are localized immunocytochemically in various gastrointestinal and male reproductive tissues. Freeze drying and chemical fixation techniques are used successfully to immunocytochemically localize tissue kallikrein in the submandibular gland of the cat and guinea-pig where it is in apical regions of striated duct cells. It is also localized in goblet (mucous) cells of the colon and mucous cells of the stomach of the cat and rat. Ultrastructurally, it is localized in the apical secretory granules of striated duct cells in the cat and guinea-pig. Furthermore, kallikrein is immunolocalized in surface epithelial cells of the guinea-pig coagulating and prostate glands; also in submucosa glands of the cat trachea. Nerve growth factor is also immunolocalized in surface epithelial cells in the guinea-pig prostate gland.

### 1. Salivary gland

A vasodilator substance was found in saliva some fifty years ago (Secker, 1934 a,b) with vascular effects unlike acetylcholine or adrenaline (Gibbs, 1935; Larson, 1935). Feldberg

and Guimaraes (1935) also concluded that this vasodepressor substance was not acetylcholine, choline, histamine or any other known depressor substance. They also observed that sympathetic saliva had more depressor activity than parasympathetic saliva and that this activity decreased in successive saliva samples. A vasodepressor substance resembling kallikrein of pancreas and urine was found in salivary glands of several species (Werle and von Roden, 1936). It was also in saliva, presumably in the same form as in the gland. It would seem that all of the above vasodilator substances were, in fact, "salivary kallikrein".

Highest concentrations of kallikrein are in the submandibular salivary glands (Werle and von Roden, 1936) with only low concentrations in parotid and sublingual glands and the rat submandibular gland has the highest concentration of any tissue. Salivary kallikrein was isolated from saliva (Fujimoto *et al.*, 1973) and from submandibular glands of man (Fujimoto *et al.*, 1973; Hare and Verpoorte, 1982), cat (Moriwaki *et al.*, 1976; Fukuoka *et al.*, 1979; Fujimori *et al.*, 1985a), mouse (Porcelli *et al.*, 1976), rat (Brandtzaeg *et al.*, 1976) and guinea-pig (Fiedler *et al.*, 1982). In the rat, submandibular gland kallikrein occurs as multiple isoenzymes (Brandtzaeg *et al.*, 1976) as is also the case in the cat (Fukuoka *et al.*, 1979).

Kallikrein is not present in submandibular glands at birth (Werle *et al.*, 1960), and it is synthesized in the adult rat submandibular gland at a slow rate (Nustad and Vaaje, 1975). Kallikrein content of saliva of older people was found to be five times higher than that of teenagers (Sallay and Nador, 1950).

Kallikrein secretion into saliva is mediated primarily by sympathetic nerve stimulation and probably involves alpha adrenergic receptors (Beilenson *et al.*, 1968; Barton *et al.*, 1975; Ørstavik and Gautvik, 1977; Schachter *et al.*, 1977; Ørstavik, 1978b; Simson *et al.*, 1979; Izumi and Aoki, 1981; Izumi, 1984). Parasympathetic denervation caused the kallikrein content of submandibular glands to fall (Emmelin and Henriksson, 1953), and this reduction in content correlated with a reduction in the size of demilune cells. Kallikrein was therefore originally thought to be present in these cells. Chronic duct ligation also caused kallikrein concentrations in submandibular glands to fall in rabbits (Mattioli and Mattioli, 1947a,b), rats, mice, dogs



(Werle, 1960; Werle and Trautschold, 1963) and cats (Beilenson *et al.*, 1968). Kallikrein was thought to be stored in acinar granules of the guinea-pig submandibular gland (Bhoola and Ogle, 1966). Based on fractionation of gland extracts, the enzyme was absent from mitochondria and lysosomes (Bhoola and Dorey, 1969). Bhoola and Heap (1970) used electron microscopy to compare isolated granules to fractions with kallikrein activity. Based on size and shape, they concluded that granules with the highest activity resembled those from acinar cells and the suggestion was therefore made again that kallikrein was stored there. This conclusion was extended to the cat, dog and rabbit (Bhoola, 1969). Other morphological data also seemed to support a connection between kallikrein release and acinar cell secretion in gland slices (Albano *et al.*, 1976).

A ductal localization of kallikrein was, however, later suggested by morphological studies after sympathetic nerve stimulation and correlations of the concentration of ductal granules with kallikrein content of the gland. In cats, sympathetic stimulation depleted striated duct cells of secretory granules, and depleted the kallikrein content of the gland, but acinar cells were left intact (Barton *et al.*, 1975; Garrett and Kidd, 1975; Schächter *et al.*, 1977). Similar results were reported in guinea-pigs and dogs (Uddin and Tyler, 1978, 1981; Schachter *et al.*, 1980). Parasympathetic nerve stimulation in the cat resulted in loss of acinar secretory granules but no reduction in kallikrein content (Barton *et al.*, 1975; Garrett and Kidd, 1975). Secretory granules from demilune cells were depleted after either sympathetic or parasympathetic stimulation, suggesting that these cells were unlikely sources of kallikrein (Uddin and Tyler, 1978).

Immunocytochemistry clearly established that kallikrein was localized in granular convoluted tubule and striated duct cells in the submandibular glands of rat (Ørstavik *et al.*, 1975; Brandtzaeg *et al.*, 1976; Ørstavik, 1978a; Ørstavik and Glenner, 1978; Simson *et al.*, 1979; Ørstavik *et al.*, 1982a), mouse (Ekfors and Hopsu-Havu, 1971; Simson *et al.*, 1978, 1979), cat (Hojjima *et al.*, 1977a; Maranda *et al.*, 1978; Schachter *et al.*, 1980), pig (Dietl *et al.*, 1978), guinea-pig (Schachter *et al.*, 1978, 1980), dog and man (Ørstavik *et al.*, 1980a; Schachter *et al.*,

1980). In sublingual glands of the rat, kallikrein was also localized in the striated duct cells (Ørstavik *et al.*, 1975; Simson *et al.*, 1983) and in the luminal rim of striated and larger duct cells (Ørstavik *et al.*, 1975). In parotid and sublingual glands of the rat, kallikrein was again localized in ductal segments (Ørstavik 1978a; Simson *et al.*, 1979; Ørstavik *et al.*, 1982a). In human parotid glands, kallikrein was localized luminally in striated duct cells and to a lesser degree in the larger ducts (Kimura and Moriya, 1984a), and some immunostaining was also reported extracellularly in duct and acinar basement membrane regions.

Enzyme histochemical techniques, using the substrate, val-leu-arg-methoxy-2-naphthylamide, supported the ductal localization of kallikrein. Active enzyme was localized periluminally in submandibular gland granular tubules and striated duct cells in the rat, mouse, hamster, guinea-pig, cat, dog (Garrett *et al.*, 1982) and man (Garrett *et al.*, 1984). An active enzyme, resembling kallikrein, was also demonstrated in salivary gland mast cells by this enzyme histochemical method. Submandibular, parotid and sublingual glands of monkeys (Arnold, 1984) and other mammals (Arnold, 1983) had kallikrein-like activity by this method also in apical regions of striated duct cells. Active enzyme was also demonstrated histochemically in luminal regions of striated duct cells of the human parotid gland (Kimura and Moriya, 1984a).

Kallikrein was finally localized immunocytochemically by electron microscopy in secretory granules of striated duct cells in cat submandibular glands (Schächter *et al.*, 1983c) (see Results). In rat submandibular glands, the enzyme was similarly localized in secretory granules of granular convoluted tubule and striated duct cells (Simson *et al.*, 1983). Apical membrane and basal lysosomal staining were also reported. Ultrastructural enzyme histochemistry also localized kallikrein-like activity in apical secretory granules in striated duct cells of monkey submandibular glands (Arnold, 1985).

Salivary gland kallikrein may be under endocrine control in some cases. Although submandibular glands of rats and mice failed to show a sexual difference and testosterone treatment did not affect kallikrein levels (Bhoola *et al.*, 1973) Van Leeuwen *et al.* (1984) found

that testosterone increased kallikrein levels in rat submandibular glands. In the hamster, which shows an estrogen-linked biochemical dimorphism of the submandibular glands, female glands contain more kallikrein than male glands (Bhoola *et al.*, 1973), but human saliva kallikrein concentration is probably not estrogen dependent since it did not vary during the menstrual cycle (Bhoola *et al.*, 1978).

## 2. Pancreas

The pancreas was found to contain large amounts of kallikrein (Kraut *et al.*, 1930). The enzyme exists in the pancreas as a precursor (prokallikrein), and also in the pancreatic secretion (Amouric and Figarella, 1979) and in fresh organ extracts (Hojima *et al.*, 1977b). It is secreted into the duodenum and activated by trypsin and enterokinase (Werle and Semm, 1955). Purification of pancreas kallikrein was achieved in the dog (Hojima *et al.*, 1977b), hog (Fiedler and Gebhard, 1980), rat (Fiedler *et al.*, 1977) cat (Fujimori *et al.*, 1985b) and man (Moriya *et al.*, 1963; ole-Moi Yoi *et al.*, 1979a). Purified enzymes usually existed in multiple forms.

Kallikrein was found to be synthesized as well as secreted by the pancreas (Frey *et al.*, 1968), where the enzyme was suggested to have an exocrine function. Support for this idea came from observations that it was secreted in parallel with trypsin and lipase and because its concentration fell like other digestive enzymes after pancreatic duct ligation (Forell, 1960).

It was hypothesized that kallikrein originated from acinar cells because of the above observations and because its concentration was not affected by treatment which destroyed  $\beta$ -cells of the islets of Langerhans (Forell, 1960). Based on subcellular fractionation, kallikrein activity was detected within secretory granules (Bhoola and Dorey, 1971), and by immunocytochemistry in apical regions of acinar cells in the pancreas of pig (Dietl *et al.*, 1978), rat (Proud *et al.*, 1977; Ørstavik and Glenner, 1978; Ørstavik *et al.*, 1980a, 1981a) and man (Ørstavik *et al.*, 1980a, 1981a). Ultrastructural immunolocalization demonstrated kallikrein in different cellular compartments of rat pancreatic acinar cells (Bendayan and Ørstavik, 1982). It was thus

localized in rough endoplasmic reticulum, Golgi body, condensing vacuoles and secretion granules which supports the view that it is synthesized and secreted by acinar cells. In accord with the above, kallikrein was localized ultrastructurally in porcine pancreatic acinar cells, secretory granules and the Golgi body (Lechene de la Porte *et al.*, 1981).

By radioimmunoassay and bioassay, kallikrein was present in isolated acinar cells, but not in isolated islet cells (Chao *et al.*, 1980). To further resolve localization in different cell types, enzyme concentration was measured following either  $\beta$ -cell destruction by streptozotocin or acinar atrophy by duct occlusion (Ørstavik *et al.*, 1981b). Kallikrein content was decreased with acinar atrophy only, which again indicates that these cells are the site of storage in this organ.

The situation has recently been rendered more complex by ole-Moy Yoi *et al.* (1979a), who, using human antisera to urinary kallikrein, immunolocalized kallikrein in  $\beta$ -cells of human pancreatic islets. Furthermore, by varying experimental conditions, kallikrein-like antigenicity was localized in either islet  $\beta$ -cells or acinar cells (Pinkus *et al.*, 1983). Untreated tissue sections immunostained exclusively in acinar cells but when enzyme-digestion methods were used kallikrein was detected in  $\beta$ -cells. The question of the presence of kallikrein in islet  $\beta$ -cells requires further study.

### 3. Colon

Hypotensive activity due to kallikrein was first reported in extracts of the colon of several mammals (Werle, 1960). Trypsin-activated kallikrein was found in the colon of man, dog, cat, rat, pig and cow (Frankish and Zeitlin, 1980) and also in colonic mucosa of rat and rabbit (Amundsen and Nustad, 1965). Rat colon extracts also contained kallikrein (Zeitlin, 1970). In the human (Zeitlin and Smith, 1973) and cat colon (Fath *et al.*, 1978), kallikrein was found mainly in the mucosal layer.

Prokallikrein, activatable by trypsin was reported to be in the colon wall of human, monkey and dog tissues (Seki *et al.*, 1972). On enzymatic and chemical grounds this enzyme

was earlier thought to resemble the plasma kallikreins, but purified human (Zimmermann *et al.*, 1979), cat and rat colon kallikrein (Frankish and Zeitlin, 1980; Fujimori *et al.*, 1985a) have since been consistently found to resemble tissue kallikreins on enzymatic, immunological and chemical grounds.

Kallikrein was localized immunocytochemically in mucous or goblet cells of human, rat and cat colon (Schachter *et al.*, 1983a) (see Results). By enzyme histochemistry, kallikrein-like activity was also detected in mucous cells of the cat and rat colon (Schachter *et al.*, unpublished observations) and also in gastrointestinal mast cells.

The above studies strongly indicate that colon kallikrein is of the tissue variety and is present in goblet cells.

#### 4. Intestine

Kallikrein, activated by trypsin, was detected in the small intestine of many mammals, viz., man, dog, cat, rat, pig and cow tissues (Werle, 1960; Werle and Vogel, 1960). Intestinal mucosal cells of rats and rabbits (Amundsen and Nustad, 1965), extracts of rabbit small intestine (Burger *et al.*, 1968) and extracts of rat intestinal tissues also displayed enzyme activity (Zeitlin, 1970; Frankish & Zeitlin, 1980).

Kallikrein in rat intestinal mucosa was present as an inactive precursor (Seki *et al.*, 1972; Zeitlin *et al.*, 1976), activatable by autolytic processes or by incubation with trypsin (Werle, 1960; Zeitlin, 1971). It behaved enzymatically like a tissue kallikrein, rather than like a plasma kallikrein (Zeitlin 1971) and its molecular weight also indicated that it was similar to other tissue kallikreins (Zeitlin *et al.*, 1976). Kallikrein was localized immuno- and enzyme-histochemically in mucous cells of rat and cat small intestine (Schachter *et al.*, unpublished observations).

### 3. Stomach

Kallikrein-like activity was found in extracts of rat and rabbit stomach (Amundsen and Nustad, 1965; Zeitlin, 1970; Frankish and Zeitlin, 1980). Activity was detected in rat stomach tissue with an unusually low pH optimum of 5 (Kobayashi *et al.*, 1979); based on enzyme inhibition profiles, this enzyme resembled cathepsin D (Kobayashi and Ohata, 1981). A kallikrein was also purified from rat stomach similar to other tissue kallikreins (Uchida *et al.*, 1980) with a pH optimum of 11. Similarly, a human stomach kallikrein was purified and found to be similar to tissue kallikreins (Uetsuji *et al.*, 1982). A kallikrein-like enzyme was also localized in the mucous cells of cat stomach (Schachter *et al.*, unpublished observations) (see Results).

### 6. Kidney (Urinary)

A hypotensive substance detected in human urine by Frey and his colleagues (Frey and Kraut, 1926) was first described as a "Kreislaufhormon" (circulating hormone). This hormone, of uncertain origin, was later named kallikrein (Greek, *kallikreas* = pancreas) since it was thought to originate in the pancreas (Frey *et al.*, 1968).

Purified human urinary kallikrein resembled other tissue kallikreins (Moriya *et al.*, 1963; ole-Moi Yoi *et al.*, 1977, 1978, 1979b). Various other kidney and urinary kallikreins have been purified and found to resemble tissue kallikreins from the same species (Nustad, 1970a; Nustad and Pierce, 1974; Moriwaki *et al.*, 1976; Porcelli *et al.*, 1976; Fujimori *et al.*, 1985b).

Kallikrein activity was first demonstrated in rat kidney homogenates and its activity was shown to be enhanced by trypsin treatment (Werle and Vogel, 1961). The suggestion was therefore made that the enzyme first isolated by Kraut *et al.* (1930) was a prohormone. The fact that the molecular weight of renal enzyme was larger than the urinary enzyme suggested to Moriwaki *et al.* (1976) that the urinary kallikrein might be a degradation product of renal kallikrein.

Subcellular localization of kallikrein within the kidney was determined on the basis of biochemical fractionation. Kallikrein activity was detected in the microsomal fraction (Nustad

and Rubin, 1970), in the endoplasmic reticulum (Nustad, 1970b; Ward *et al.*, 1976a), in the plasma membrane (Nustad, 1970b; Ward *et al.*, 1975, 1976a, b), and in lysosomes of kidney cells (de Carvalho and Diniz, 1966; Baggio *et al.*, 1975; Heidrich and Geiger, 1980).

Kallikrein was found to be synthesized by the renal cortex (Nustad, 1970b) and suspensions of renal cortical cells (Kaizu and Margolius, 1975), and dog renal homogenates had the highest levels of such activity in extracts of cortex (Scicli *et al.*, 1976b). Release of kallikrein from rat kidney occurred into the venous effluent and urine (Roblero *et al.*, 1976).

Since the time that Werle and Vogel (1960) showed a reduced urinary excretion of kallikrein following pharmacological destruction of tubule cells, most evidence suggests the presence of kallikrein in the distal nephron tubules. Extracellular secretion of urinary kallikrein from suspended renal cortical cells was increased by aldosterone and decreased by spironolactone, an aldosterone antagonist in distal tubules (Kaizu and Margolius, 1975). Secretion of kallikrein occurred in the distal nephron based on stop-flow analysis of dog kidney (Scicli *et al.*, 1976a).

The proximal convoluted tubule was also reported to contain kallikrein activity (Nustad, 1970b). Reabsorption droplets in proximal tubule cells were immunocytochemically stained (Simson *et al.*, 1979; Pinkus *et al.*, 1981), which suggested that tissue kallikrein was reabsorbed from tubule fluid (Kimura and Moriya, 1984b). By enzyme histochemistry, active rat kidney kallikrein was localized in tubule cytoplasm of the outer medulla and deep cortex, where proximal tubules, ascending limb of Henlé and collecting ducts reside (Kimura *et al.*, 1982). In the human kidney, Kimura and Moriya (1984b) unexpectedly observed kallikrein-like enzyme activity exclusively in proximal tubule cells. Since kallikrein immune staining did not occur in these or other tubule cells, it was suggested that the enzyme activity was due to non-kallikrein esterases. These authors who found kallikrein immune staining in interstitium and basement membrane regions, viewed renal and urinary kallikreins as derived from circulating tissue kallikrein.

In microdissected rabbit kidney, active and inactive kallikreins were detected in connecting tubule cells (Tomita *et al.*, 1981; Omata *et al.*, 1982; Proud *et al.*, 1983; Marchetti *et al.*, 1984). Using immunocytochemistry, kallikrein secretion was indicated by its localization in apical regions of distal tubule cells and in the tubule lumen of rat (Ørstavik *et al.*, 1976; Simson *et al.*, 1979; Ørstavik and Inagami, 1982), mouse (Simson *et al.*, 1979), and human kidney (Pinkus *et al.*, 1981). Kimura and Moriya (1984b) postulated that kallikrein secretion in the distal tubule may not necessarily indicate *de novo* synthesis in tubule cells, where it is secreted into the urine. Kallikrein antigenicity in distal tubule cells may therefore represent tissue enzyme from other sources. Figueroa *et al.* (1984) localized kallikrein immunocytochemically in the cytosol of apical and perinuclear areas of distal connecting tubule cells. The antigenicity pattern within organelles suggested that kallikrein synthesis occurred in this cell type. Basal infoldings of connecting tubule cells also stained, which may explain the appearance of kallikreins in isolated kidney venous effluent (Roblero *et al.*, 1976; Vio *et al.*, 1981, 1982, 1983).

Although the physiological significance is not completely understood, Margolius *et al.* (1972, 1974) observed that urinary kallikrein excretion is abnormal in various hypertensive animal models and in human hypertensive disease. Low sodium diet or high levels of sodium-retaining steroid hormones also increased urinary kallikrein excretion in rat and man (Geller *et al.*, 1972; Margolius *et al.*, 1974). Further observations have been made indicating that kallikrein is under hormonal control in the kidney but the physiological significance of this fact remains obscure (Geller *et al.*, 1972; Margolius *et al.*, 1974).

Kallikrein activity in the nephron tubule was shown to be under corticosteroid hormone control. Urinary kallikrein secretion was stimulated by aldosterone or deoxycorticosterone (Geller *et al.*, 1972; Margolius *et al.*, 1974) and kallikrein activity in the rabbit connecting tubule decreased after adrenalectomy (Marchetti *et al.*, 1984). Chronic deoxycorticosterone treatment also increased activity of kallikrein in the connecting tubule, though not its synthesis (Marchetti *et al.*, 1984).



Possible physiological roles of renal kallikreins were also suggested to be related to the activation of prorenin by kallikrein (Sealey *et al.*, 1978). Though prorenin cleavage resulted in activation, some structural alteration of prorenin was required before activation could occur (Hsueh *et al.*, 1980).

## 7. Respiratory Tract

Human bronchial secretion has been reported to contain kallikrein (Havez *et al.*, 1966a,b). It was detected in normal "respiratory sputum" and found to resemble tissue kallikreins on enzymatic and immunological grounds. The problem is whether the sputum contained salivary kallikrein, but the authors do not discuss this obvious problem.

## 8. Male Sex Glands

An arginine esterase was described in the different lobes of rodent prostate glands (Gotterer *et al.*, 1956). The guinea-pig coagulating gland was shown to secrete "toxic" substances (Freund and Thompson, 1957) and in further studies this factor was characterized as a glycoprotein with arginine esterase activity that increased vascular permeability (Freund *et al.*, 1958). Bhoola *et al.* (1962) and Moriwaki and Schachter (1971) found a kinin-releasing "kallikrein" (coagulating gland kininogenase, CGK) responsible for hypotensive, permeability-enhancing and esterolytic actions of coagulating gland extracts. They emphasized the kallikrein-like nature of this kininogenase (CGK). Purified CGK resembled other tissue kallikreins on enzymatic and chemical grounds (Moriwaki *et al.*, 1974). Kinin-producing activity was not detected in coagulating or prostate gland extracts of rat, rabbit, dog or man (Bhoola *et al.*, 1962). CGK is now known to resemble the kallikreins chemically (Fiedler, 1979).

Synthesis of a major protein of canine seminal plasma was found in prostate slices and its concentration was increased by androgens and decreased by castration (Dubé *et al.*, 1983). Isaacs and Shaper (1983) and Chapdelaine *et al.* (1984) determined that this major androgen-dependent protein was a serine arginine esterase similar to tissue kallikreins. This

enzyme was identified as a serine protease of the tissue kallikrein type but its ability to release kinin was apparently not tested (Lazure *et al.*, 1984).

A cytosolic localization of CGK from guinea-pig was found by Barton *et al.* (1973), who demonstrated approximately 90% of the activity was in the soluble fraction of guinea-pig coagulating gland extracts. Immunocytochemically, kallikrein was localized apically in the secretory epithelial cells of the guinea-pig coagulating (GPCG) (Schachter *et al.*, 1978) and prostate glands (GPPG) (see Results).

## 9. Brain

Kallikrein-like activity was first described in nervous tissue by Werle and Vogel (1961). Rabbit and rat brain homogenates (Shikimi *et al.*, 1972) also had kallikrein-like activity. Highest concentrations were in the cerebral cortex (Shikimi *et al.*, 1972) where kallikrein activity was also reported to be inhibited by aprotinin. Scicli *et al.* (1984) found kallikrein in many regions of the brain, including the hypothalamus.

Simson *et al.* (1984) immunolocalized tissue kallikrein in rat brain. Although staining did not occur in specific cell types, it was found in hypothalamus, ventricular ependymal cells, and cell bodies of certain nuclei. The kallikrein-kinin system was therefore postulated to participate in nervous and neuroendocrine functions (Scicli *et al.*, 1984).

## 10. Other Tissues

A kallikrein-like enzyme was isolated from porcine pituitary glands and found to resemble other tissue kallikreins (Polivka *et al.*, 1982). Powers and Nasjletti (1983) also reported kallikrein-like activity in the rat neuro-intermediate lobe where it may be involved in pro-opiomelanocortin processing. Synthesis of kallikrein in the anterior lobe was suggested by the finding of kallikrein mRNA in normal and hypersecretory rat pituitaries (Fuller *et al.*, 1985). Powers *et al.* (1984) suggested an involvement with hypertension since lower levels of tissue kallikrein activity were found in genetically hypertensive rats. A possible role in hormone

processing was implied by the observation that kallikrein-like activity of the female rat anterior pituitary was fifteen times higher than that of the male (Powers and Nasjletti, 1984).

Kinin-releasing proteases were also isolated from blood vessels (Nolly and Lama, 1982; Nolly *et al.*, 1983), rat cardiac tissue (Britos and Nolly, 1981) and rat spleen (Chao *et al.*, 1984b). With allowances for minor variations, these enzymes can be included in the tissue kallikrein family. Tissue kallikrein, activatable by trypsin (Chao *et al.*, 1984a), was found in intact human erythrocytes and ghosts. Its possible relation to erythrocyte maturation or to membrane transport remains to be established.

Rat thyroid gland also contains a kinin-releasing esterase resembling other tissue kallikreins (Uchida *et al.*, 1982). Chemical and physical properties of this enzyme were established but its physiological role is unknown.

## B. Nerve Growth Factor

This substance is included because, as described below, it often co-exists in tissues with kallikrein, and is also associated with a kallikrein-like molecule in a macromolecular complex or aggregate.

Nerve growth factor (NGF) is a molecular complex possessing a subunit with some chemical and physical properties which are similar to those of the kallikreins (Thoenen and Barde, 1980; Darling *et al.*, 1983; Calissano *et al.*, 1984). NGF was discovered when a substance was found in sarcomas, snake venoms, and elsewhere, which stimulated growth of chick sympathetic neurons (Bueker, 1948; Levi-Montalcini and Hamburger, 1951; Levi-Montalcini and Booker, 1960).

The richest known source of NGF is the submandibular gland of the male mouse (Levi-Montalcini and Angelletti, 1961). Submandibular glands of the rat, guinea-pig, cow, pig, rabbit and man contain little or no NGF (Harper and Thoenen, 1980). NGF was recently found in prostate glands, though not all prostate glands contain it. Of several animals surveyed, only guinea-pig, rabbit and bull showed NGF activity in prostatic tissue (Harper *et al.*, 1979;

Harper and Thoenen, 1984). Purification of guinea-pig prostatic NGF was achieved by Chapman *et al.* (1981). Bovine seminal plasma NGF was purified and found to resemble mouse submandibular gland NGF in biological and immunological properties (Harper and Thoenen, 1984). Other rich sources of NGF include snake venom (Cohen, 1959), human placenta at term (Goldstein *et al.*, 1983) and submandibular and prostate glands of the house musk shrew (*Suncus murinus*) (Ueyama *et al.*, 1981).

Male mouse submandibular gland NGF was purified and characterized (Varon *et al.*, 1968; Server and Shooter, 1977) and found to be composed of three subunits ( $\alpha$ ,  $\beta$ ,  $\gamma$ ), each of which is dimeric (Varon *et al.*, 1968). One or two bound zinc ions further stabilize the complex (Pattison and Dunn, 1975). The  $\alpha$ -subunit has not been shown to have enzymatic activity; the complex's biological activity on nerve growth residing with the  $\beta$ -subunit (Varon *et al.*, 1968). The  $\beta$ -subunit promotes survival, differentiation and maintained function of sympathetic nerve cells (Mobley *et al.*, 1977a, b, c) and may also be the chemotactic agent for neurite outgrowth (Campanot, 1977). The  $\gamma$ -subunit is a potent kallikrein-like arginyl esterpeptidase (Green *et al.*, 1969; Moore *et al.*, 1974; Thomas and Bradshaw, 1981). Bothwell (1979) and Fiedler and Fritz (1981) described large sequence homologies between  $\gamma$ -NGF and tissue kallikreins. The activity of  $\gamma$ -NGF is to cleave the  $\beta$ -NGF precursor (pro- $\beta$ -NGF) to yield  $\beta$ -NGF (Berger and Shooter, 1977). Morris *et al.* (1981) found that the  $\gamma$ -NGF subunit also activates inactive renin.

Synthesis of NGF and differentiation of secretory tubule cells of rodent submandibular glands were shown to be under control of testosterone (Levi-Montalcini and Angeletti, 1961; Goldstein and Burdman, 1965). NGF was reported to be located in the granular convoluted tubules of the rodent submandibular gland (Levi-Montalcini and Angeletti, 1964).

Light microscopy and subcellular immunocytochemical localization indicated an association of NGF with cytoplasm, secretory granules, the basal part of the cell and the nucleus (Levi-Montalcini and Angeletti, 1961; Goldstein and Burdman, 1965; Kumar *et al.*, 1972). These early conclusions were based on inadequate methods, and later light microscopic

and ultrastructural immunocytochemical studies showed NGF localization in apical secretory granules of secretory tubule (or duct) cells and in saliva (Schwab *et al.*, 1976). The African rodent, *Praomys (mastomys) natalensis*, has high levels of NGF in submandibular glands, where it was also immunolocalized in the granules of the convoluted tubules (Aloe *et al.*, 1981). NGF was also immunolocalized in Suzmann cells or other associated dense fibres in the rat iris (Rush, 1984), and in GPPG within epithelial cells and luminal secretions (Shikata, 1984).

## II. MATERIALS AND METHODS

### A. Animals, Organ Dissection, Tissue Preparation and Extracts

Animals were kept in a stable environment prior to surgery. Animals were then starved for 24 hours and water allowed *ad libitum* for several hours prior to surgery.

Cats of either sex, weighing 2-4 kg, were anaesthetized with intraperitoneal sodium pentobarbital (M.T.C. Pharmaceuticals, 30-35 mg/kg). A midline incision was followed by blunt dissection to reveal submandibular glands on either side of the trachea near the junction of the transverse vein with the external jugular vein. A small area of gland was separated with clamps and removed while blood flow was intact. Tissue was rinsed briefly in saline and minced on a paraffin-covered dish. Mincing usually took place in fixative unless tissues were to be freeze-dried, where mincing took place in saline. Final tissue sizes were typically less than 1 mm<sup>3</sup>. Cat trachea was obtained similarly. Pieces of trachea near the bifurcation were removed, rinsed in saline and minced. The stomach was delivered through an incision and small pieces of tissue were obtained from fundus, corpus or antrum regions. Ascending, transverse and descending portions of the colon were also removed. Upon removal, tissues were rinsed in saline and minced. In some cases, only the mucosal layer, which was stripped of muscular layers, was used.

Rats of either sex weighing 300-400 grams were anaesthetized with intraperitoneal sodium pentobarbital (40-50 mg/kg). Gastric and colonic tissues were removed as before. Usually, only stomach antrum mucosa or descending colon mucosa were removed, rinsed in saline and minced.

Male guinea-pigs, weighing 600-1200 grams, were anaesthetized with intraperitoneal sodium pentobarbital (40 mg/kg). Animals were prepared as above and submandibular glands and sex glands identified. Pieces were removed, rinsed with saline and minced. Seminal vesicles, coagulating gland, and dorsal and ventral prostate glands were identified. Pieces of paired coagulating glands, consisting of larger, more transparent lobes, and paired dorsal or lateral

prostate glands were removed, rinsed in saline and minced. Pieces were also transferred into fresh fixative or onto prepared surfaces for rapid freezing.

Extracts were prepared for immunodiffusion experiments from various tissues. Tissues were cut into small pieces, mixed in 5 volumes of distilled water and stirred for 4 hours at room temperature. Mixtures were centrifuged for 10 minutes at 10,000 g and pellets were homogenized and resuspended with distilled water. Two more centrifugations followed, with pellets being homogenized and resuspended. Supernatants were collected, pooled and freeze-dried. Powders were resuspended in distilled water, centrifuged and freeze-dried.

## **B. Fixation of Tissues**

### **1. Chemical Fixation**

Fixative solutions were developed for the dual purpose of preserving ultrastructure while retaining antigenicity. Hundreds of combinations were tested for various tissues, with most intense investigation of the cat submandibular gland. Factors which determined a fixative's suitability were concentration of cross-linking reagent, fixative osmolarity and time of fixation. Factors which were less important were type of buffer, fixation temperature, pH (range 7.0-7.5) and addition of  $\text{CaCl}_2$ . A summary of major fixatives follows.

For the cat submandibular gland, most commonly employed cross-linking reagents were aldehydes. Fixatives which used only one type of aldehyde were 1% glutaraldehyde (GTA) (Analychem Corp. Ltd. or J.B. EM Services, Inc.) and 2% or 4% paraformaldehyde (PFA) (Fisher Scientific Ltd.) made from depolymerized powder. Fixatives were made in 0.1 M sodium cacodylate (Analychem Corp. Ltd.) and adjusted to a pH between 7.0 and 7.5 with either NaOH or HCl. Solutions were prepared 1-2 hours before use. Tissues were kept in fixative for 1-2 hours at room temperature. PFA and GTA combinations were often employed. Concentrations of GTA ranged from 0.7%-1.5%, most often held at 1%, while PFA concentrations ranged from 2%-4.5%, often held at 3%. Acrolein (Analychem Corp. Ltd.) was

sometimes added at a concentration of either 0.1% or 1%, but this caused masking of antigenicity. Buffering of fixatives was achieved with either 0.05 M or 0.1 M cacodylate adjusted to a final pH between 7.0-7.5. Sodium phosphate buffers (pH 7.2) were also used. Molarity varied from 0.075 (180 mOsm, measured by freezing point depression with an Osmette precision osmometer) (Precision Systems) - 0.135 M (300 mOsm). Time of fixation for these aldehyde combinations varied from 1-2 hours except for those using acrolein, which was 15 minutes. Often used to determine ultrastructure for a gland was a relatively concentrated combination of aldehydes arrived at by Barton *et al.* (1975). It consisted of a mixture of 2.5% GTA, 4% PFA and 1% acrolein in 0.05 M cacodylate, pH 7.3. Primary fixation was for 2 hours followed by 1 hour post-fixation with cacodylate-buffered 1% osmium tetroxide (osmium) (Analychem Corp. Ltd), pH 7.3.

Cat submandibular gland fixation was examined using the bifunctional reagent, 1-ethyl-3-(3-dimethylaminopropyl) carbodiimide HCl (CDI) (Sigma Chemical Co.) in concentrations of 1% and 10%. CDI was also combined with aldehyde reagents. GTA ranged from 0.5%-1%, while CDI levels were either 2%, 4% or 6%. Also, combinations were used where CDI was 1% or 2%, GTA was 0.5% and PFA was 1%, 2% or 3%. These primary fixatives were used at 4°C for 0.5-4 hours. In some cases, tissues were post-fixed with 1% osmium buffered with 0.05 M cacodylate, pH 7.3, for 15 or 30 minutes at room temperature. Although the buffering system for CDI combinations was occasionally 0.05 M Trizma base (Tris) (Sigma Chemical Co.), 0.1 M sodium phosphate or phosphate-buffered physiological saline, 265 mOsm (Lewis and Knight, 1980), all at pH 7.2, usually a phosphate-buffered saline (PBS)/Tris buffer (Willingham and Yamada, 1979) was used which had a final pH of 7.0 and an osmolarity of 360 mOsm. Osmolarities for solutions containing combinations of CDI, GTA and PFA ranged from 260-1200 mOsm with most suitable osmolarities below 500 mOsm.

Other fixatives were used to compare with the reagents described above. The periodate-lysine-paraformaldehyde (PLP) solution (McLean and Nakane, 1974) was used. It was modified according to Reissig *et al.* (1978) by varying PFA concentrations from 1%-2%.



periodate kept at 0.01 M, lysine at 0.075 M and buffered with either sodium phosphate or cacodylate. The pH was adjusted before use to 7.2 and fixation was for 4 hours at room temperature. Also, 1% acrolein was added to the PLP fixative and used for 15 minutes. Other schemes included Bouin solution (Humason, 1979), 80% ethanol or 1% osmium in Millonig's buffer (Millonig, 1961). Most of these fixatives distorted ultrastructure or masked antigenicity. cat submandibular gland tissue was also perfusion-fixed by injecting 100 ml of fixative into the cannulated carotid artery of an anaesthetized animal. PLP, with or without 3% dextran (Sigma Chemical Co.) or 0.05 M cacodylate-buffered 1% GTA, pH 7.2, was used as perfusant. After 5 minutes, glands were removed and processed in fixative for an additional 4 hours. Although perfusion techniques were successful, they were also cumbersome and results were comparable to immersion fixation.

In preserving guinea-pig submandibular gland ultrastructure, similar fixatives to those used for the cat submandibular gland were developed. Combinations were used where GTA ranged from 0.5%-1.3% and PFA was kept at either 2% or 3%. Acrolein was sometimes added to a level of 0.1%, 1%, 2% or 5% CDI was used in conjunction with either 0.5% GTA or 0.5% GTA and 1% PFA. Post-fixation with 0.1 M cacodylate-buffered 1% osmium, pH 7.1, for 15 minutes was sometimes tried. 10% CDI was also used for 24 hours at 4°C. When CDI was included in the fixative, PBS/Tris buffer was used and the final pH adjusted to between 7.0 and 7.5. 0.05 M or 0.1 M cacodylate was used to buffer these mixtures in the same pH range. The concentrated aldehyde fixative used for the cat submandibular gland was also used for the guinea-pig submandibular gland to examine regular ultrastructure. Other unsuccessful fixatives included cold 95% ethanol or 2% dimethylsuberimidate (DMS) (Sigma Chemical Co.) in 0.15 M Tris, pH 9.5, to which 0.02 M CaCl<sub>2</sub> was added (Hassel and Hand, 1974). These last two fixatives were used for 4 and 2 hours, respectively, but caused cell distortion.

Solutions used to fix pieces of cat colon included aldehyde-based mixtures such as 1% GTA-3% PFA for 2 hours, 1% CDI-0.5% GTA for 1 or 2 hours, all in PBS/Tris buffer. 1% CDI was also used alone or in combination with 2% PFA with or without the presence of 0.1%

saponin (Fisher Scientific Co.) made in either PBS/Tris buffer or 0.1 M sodium phosphate, pH 7.4. 4% PFA was used in sodium phosphate buffer, pH 7.4, for 24 hours at 4°C. PLP, cold 95% ethanol or 0.5% osmium in 0.075 M cacodylate, pH 7.0, were also used to fix tissue.

Rat colon pieces were fixed in solutions of 1% CDI in PBS/Tris buffer, pH 7.0, or PLP for 3 hours. Osmium was used at three concentrations: 0.5%, 1% and 2% in 0.1 M cacodylate buffer, pH 7.0.

Fixatives for the cat stomach were 1% CDI, 1% CDI-0.5% GTA or 1% GTA-3% PFA in PBS/Tris buffer, pH 7.3, for 1-2 hours. 1% CDI-2% PFA in 0.1 M phosphate buffer, pH 7.4, with or without 0.1% saponin, was also used for 2 hours. 4% PFA was used in either 0.1 M phosphate buffer for 24 hours at 4°C or 0.1 M cacodylate buffer for 30 minutes followed by post-fixation with 0.7% osmium in 0.1 M cacodylate buffer, pH 7.0, for 2 hours. Three concentrations of osmium, 0.5%, 0.7% and 1%, were used in 0.05 M, 0.075 M or 0.1 M cacodylate buffer, pH 7.0-7.4, for 1-2.5 hours. PLP and cold 95% ethanol were also used.

Rat stomach was fixed in either 1%, 3% or 5% CDI or 1% CDI-2% PFA in PBS/Tris buffer. PLP was used for 3 hours, or 0.5%, 1% and 2% osmium in 0.1 M cacodylate buffer, pH 7.2, were used for 1.5 hours.

Fixatives for cat trachea were PLP, 0.5% osmium in 0.05 M cacodylate buffer, pH 7.2, or a combination of 1% CDI and either 2% or 3% PFA. The effect of adding 0.1% saponin was also tested in 1% CDI-2% PFA mixtures.

For ultrastructure, guinea-pig male sex glands were fixed in a modified aldehyde solution after Barton *et al.* (1975). GTA was either 2.5% or 3% with 3% PFA in PBS/Tris buffer. Post-fixation was with 1% osmium for 1 hour. Other combinations included 0.5% or 1% GTA with 2% or 3% PFA in 0.05 M cacodylate buffer. CDI concentrations of 1% or 2% were used with or without 0.5% GTA, and PFA levels ranged from 1%-3%. In some cases, a 15 minute post-fixation step with 1% osmium in 0.05 M cacodylate buffer was added. Osmium as a primary fixative was also used in concentrations of 0.5%, 1% or 2%. Other fixatives included cold 95% ethanol or 2% DMS as described for guinea-pig submandibular gland.

Fixatives that were specifically tested on GPCG pieces were 0.2% or 0.5% GTA with either 1% or 2% PFA. 2%, 3% or 6% PFA was also used. Fixatives used for male sex glands which were used for pre-embedding immunostaining systems are described in that section.

Following fixation, all tissues were washed for at least 1 hour in several changes of fixative buffer. If post-fixation was involved, it followed a buffer wash and after post-fixation, a buffer wash corresponding to this step was used. Washes were followed by dehydration in ascending ethanol concentrations (30%, 50%, 70%, 90%, 95% and 100% twice). If acrolein was part of the fixative, methanol instead of ethanol was used. Xylene was used as intermediate solvent for tissues to be embedded in paraffin (Paraplast Plus) (Fisher Scientific Co.). Propylene oxide was used as an intermediate solvent for tissues to be embedded in epoxy resin. Tissues processed into glycol methacrylate (GMA) (Analychem Corp. Ltd.) were dehydrated without alcohol.

To enhance antigenicity by protecting tissue with chemical treatments, a modified method of Kuhlmann and Krischan (1981) was used. Cat and rat stomach, GPCG and cat trachea were treated with ethyl acidimidate HCl (EAI) (Sigma Chemical Co.) as follows. After fixation, tissues were washed in several buffer changes for at least 2 hours and then treated with 0.2 M EAI dissolved in 0.2 M  $K_2HPO_4$ , pH 7.3, for 2 hours at 4°C. A 1 hour buffer wash following treatment was followed by routine dehydration. Rat stomach and GPCG were additionally used to examine polyvinylpyrrolidone (PVPP) (Sigma Chemical Co.) treat-

ment to enhance antigenicity. After fixation and buffer wash, tissues were treated with 10% PVPP in buffer for 2 hours. A routine dehydration schedule followed, with all concentrations of ethanol containing 10% PVPP. Tissues were then transferred through two changes of propylene oxide before embedding.

## 2. Freeze-Drying

Tissue pieces were rapidly frozen either by copper block or cryogenic fluid methods. Cryogenic fluids such as 2-methylbutane (Eastman Kodak Co.) or freon (Analychem Corp.


Ltd.) were cooled to their melting point with liquid nitrogen. Tissue pieces were frozen by immersion in cryogenic fluid and then freeze-dried in an Edwards-Pearse tissue drier (EPTD) (Edwards High Vacuum) at  $-60^{\circ}\text{C}$  for 24-48 hours. No vapor fixation step was used before embedding in vacuum-treated epoxy resin. This method was used to preserve cat submandibular gland, guinea-pig submandibular gland, GPCG and GPPG tissues.

The copper block method of freezing (Figure 1) (Coulter and Terracio, 1977) was used for cat submandibular gland, guinea-pig submandibular gland, cat colon, rat colon, cat stomach, rat stomach, cat trachea, GPCG and GPPG tissues. Pieces of tissue were placed on thin-gauge aluminum foil on cork stoppers and blotted of excess moisture. Stopper bases were attached to the brass plunger which rapidly delivered tissues onto the polished surface of copper, maintained at liquid nitrogen temperature. Pieces were kept in contact with the block for at least 20 seconds and then delivered into the freeze-drier. Some tissues were processed in the EPTD as above, and others were freeze-dried in an all-glass freeze-drier (Coulter and Terracio, 1978). This device was cooled with liquid nitrogen to keep tissues below  $-137^{\circ}\text{C}$  while being dried and an E2M2 doublestage vacuum pump (Edwards High Vacuum) kept pressure below  $10^{-3}$  torr. In some cases, osmium was used for 1-4 hours to vapor-fix tissues. Tissues were embedded into vacuum-treated epoxy resins. Both cryogenic fluids or the copper block method to freeze tissues produced reasonable structure for light microscopy. Ultrastructure, however, was poor and ice crystal artifact was evident.

### C. Embedding

Tissues were embedded in paraffin using at least two changes of melted paraffin (melting point  $56^{\circ}\text{C}$ ). Araldite and Epon epoxy resins (Analychem Corp. Ltd.) were used according to Hayat (1970). Following the propylene oxide stage of dehydration, a 1:1 ratio of resin:propylene oxide mixture was used on tissues for 24 hours. Tissues were then embedded in capsules filled with resin and polymerized at  $60^{\circ}\text{C}$  for 48 hours.

FIGURE 1. All-glass, low-temperature freeze-drier and copper block freezing device (inset). O, osmium port; P, plunger; R, resin port; T, tissue trough, holds tissue pieces which are cooled by liquid nitrogen (in dewar flask below trough). (After Coulter and Terracio, 1977, 1978).



Tissues that were freeze-dried were usually embedded in Spurr low-viscosity embedding medium (Analychem Corp. Ltd.) according to Spurr (1969). Polymerization occurred at 60°C for 24 hours. Other tissues were embedded in water-miscible GMA. According to Leduc and Bernhard (1967), dehydration with water-GMA monomer mixture was followed by overnight GMA polymer impregnation. Polymerization of embedded tissues was with long wavelength UV light for 8-48 hours.

#### D. Immunocytochemistry

##### 1. Antisera

Antigens, previously purified, had high esterolytic activities and yielded single bands on disc electrophoresis. Cat salivary kallikrein (CSK), guinea-pig salivary kallikrein (GPSK) and human salivary kallikrein (HSK), purified according to Fujimoto *et al.* (1973) and Fukuoka *et al.* (1979), rat pancreatic kallikrein (RPK) (Hojima *et al.*, 1977b), rat intestinal kallikrein (RIK) (Moriwaki *et al.*, 1974, 1980) and guinea-pig coagulating gland kallikrein (CGK) (Moriwaki *et al.*, 1974) were all generous gifts of the late Dr. C. Moriwaki.

Kallikrein antisera were raised in New Zealand rabbits (Schachter *et al.*, 1980) and blood for immune sera was collected from ear veins. Blood from hyperimmunized rabbits was also collected. NGF rabbit antisera were purchased from Collaborative Research Inc. They were raised against 2.5s NGF.

Antisera produced single precipitin lines when reacted with purified antigen or tissue extracts. Although stomach and trachea extracts did not show immune precipitation when reacted with  $\alpha$ CSK, double diffusion analysis may not have been sensitive enough to detect small amounts of antigen. Specificity of  $\alpha$ CSK was shown (Figure 2) by its ability to form single precipitin lines with cat submandibular gland extracts (peripheral wells 2, 4 and 6). Single precipitin lines were also shown for diffusion of  $\alpha$ CSK (center well) and cat colon extracts (peripheral wells 1, 3 and 5). Partial antigenic non-identity between cat submandibular gland

**FIGURE 2.** Double diffusion analysis of  $\alpha$ CSK (center well) with cat colon (well 1, 10  $\mu$ g; well 3, 20  $\mu$ g; well 5, 30  $\mu$ g) and cat submandibular gland (well 2, 10  $\mu$ g; well 4, 20  $\mu$ g; well 6, 30  $\mu$ g) extracts. Note single precipitin line and spur formation between lines from peripheral wells 2 and 3.





and cat colon antigens was disclosed by spur formation between lines from peripheral wells 2 and 3.


## 2. Immunostaining systems

### a. Peroxidase-antiperoxidase

#### i. Paraffin sections

Peroxidase-antiperoxidase staining procedures (Sternberger, 1979) (Figure 3) were modified for paraffin-embedded tissues. Seven  $\mu\text{m}$  thick sections were cut and mounted on gelatin-coated glass slides. Removal of paraffin with several changes of xylene was followed by rehydration through a descending ethanol concentration gradient (100% twice, 95% twice, 70%) and washing in PBS or 0.05 M Tris buffer, pH 7.6. Excess moisture was removed and slides were put into a humidity chamber. Sections were incubated for 30 minutes at room temperature in normal goat serum (diluted 1:30) (Gibco Laboratories). Unless otherwise stated, all dilutions are in 0.05 M Tris buffer, pH 7.6, and all normal goat sera dilutions are 1:30. They were blotted and covered with either immune or normal rabbit serum (diluted 1:1000-1:5000 with buffer containing 1% normal goat serum, optimally diluted 1:2000). Unless otherwise stated, all primary sera were diluted in buffer with 1% normal goat serum. Humidity chambers were covered and sections were incubated for 24 hours at 4°C. Sections were rinsed with buffer, incubated for 30 minutes with goat serum immunized against rabbit immunoglobulin G Fab fragments (FAB) (Miles Laboratories Inc.) (diluted 1:20-1:80, optimally diluted 1:40), rinsed in buffer, incubated for 30 minutes with rabbit peroxidase-antiperoxidase (PAP) complex (Miles Laboratories Inc., diluted 1:30-1:100, or Cappel Laboratories, diluted 1:50-1:120, optimally diluted 1:100), rinsed with buffer, and finally incubated with a freshly prepared, filtered solution of 0.05% 3,3'-diaminobenzidine tetrahydrochloride (DAB) (Sigma Chemical Co.) and 0.01%  $\text{H}_2\text{O}_2$  (J.T. Baker Chemical Co.) in buffer. Reaction was stopped after

**FIGURE 3.** Summarized PAP method of immunostaining. LM, light microscopically; EM, electron microscopically. (After Sternberger, 1979.)



10-15 minutes by rinsing slides in distilled water. Slides were dehydrated in xylene and mounted with Permount (Fisher Scientific Co.). Paraffin-embedded tissues and antisera tested for immunostaining were as follows. The cat submandibular gland was tested with  $\alpha$ CSK and  $\alpha$ CGK. The cat colon was tested for reactivity with  $\alpha$ CSK and  $\alpha$ CGK, and the rat stomach was tested with  $\alpha$ CSK,  $\alpha$ RPK and  $\alpha$ RIK.

For routine light microscopic examination of structure, paraffin sections were stained with hematoxylin and eosin (HE) (Humason, 1979) or by the periodic acid-Schiff reagent (PAS) (Fisher Scientific Co.) of Humason (1979).

## ii. Semi-thin resin sections

Procedures for semi-thin resin section immunostaining were similar to those described by Schachter *et al.* (1983a). Sections measuring 0.5  $\mu$ m to 1.5  $\mu$ m thickness were cut on a Reichert-Jung ultracut (Sargent-Welch Scientific Co.) and dried onto gelatin-coated glass slides on a hot plate at 200°C. Resin was removed by treating slides for 30-90 minutes with 30%-100% saturated ethanolic or methanolic sodium ethoxide (Aldrich Chemical Co.). In some cases, slides were treated with a similar solution made at least 1 week before use by dissolving NaOH pellets in absolute ethanol (Lane and Europa, 1965). Following etching, slides were rinsed twice in absolute ethanol (or methanol), twice in distilled water and immersed in buffer for at least 30 minutes. In some cases to reduce background staining for tissues fixed with osmium, a distilled water rinse-H<sub>2</sub>O<sub>2</sub> treatment (5% or 10% for 5-15 minutes) -distilled water rinse sequence was introduced between the earlier distilled water rinses. For the cat colon, detergent treatment was sometimes attempted to enhance penetration of staining reagents into fixed tissue (DeArmond *et al.*, 1981). 2% sodium dodecyl sulfate (SDS) (Terochem Laboratories Ltd.) in PBS with 10<sup>-3</sup> M ethylene diaminetetraacetate (EDTA) (Sigma Chemical Co.) was used for 10 minutes. Slides were rinsed twice with PBS-EDTA and then Tris buffer. Following etching, sections were incubated in normal goat serum for 15-30 minutes at room temperature. In some experiments, this step was omitted to test normal goat serum's

effectiveness in blocking sites of non-immune or unspecific binding of reagents. They were blotted with Whatman No. 4 filter paper (Fisher Scientific Co.) and covered with either immune or normal rabbit serum (diluted 1:500-1:10,000). Incubations were carried out in humidity chambers for 48-72 hours at 4°C. Sections, at room temperature, were rinsed with buffer, blotted, incubated for 30 minutes in FAB (diluted 1:10-1:80), rinsed with buffer, blotted, incubated for 30 minutes in PAP (diluted 1:30-1:120), rinsed in buffer, blotted and incubated in a freshly prepared, filtered solution of 0.025% DAB and 0.005% H<sub>2</sub>O<sub>2</sub> in buffer, until a brown reaction product appeared (usually 5-10 minutes). Slides were rinsed in distilled water, dried at 60°C for at least 1 hour and mounted with Permount.

Semi-thin resin-embedded tissues and antisera tested with the PAP method were prepared as follows. The cat submandibular gland was tested with  $\alpha$ CSK and the guinea-pig submandibular gland was tested with  $\alpha$ GPSK. The cat colon was tested with  $\alpha$ CSK,  $\alpha$ RPK,  $\alpha$ RIK,  $\alpha$ GPSK,  $\alpha$ HSK and  $\alpha$ CGK and the rat colon was tested with  $\alpha$ CSK. Cat stomach tissue was tested with  $\alpha$ CSK and  $\alpha$ RPK and the rat stomach was examined using  $\alpha$ CSK. Cat trachea was tested with  $\alpha$ CSK, GPCG was tested with  $\alpha$ CGK and GPPG was tested with  $\alpha$ CGK,  $\alpha$ CSK and  $\alpha$ NGF. Cat submandibular gland, cat colon, cat and rat stomach semi-thin sections were also stained by the PAS method, either alone or in conjunction with immunostaining. When both staining methods were used together, PAS staining for carbohydrates followed PAP immunostaining. Sometimes, semi-thin resin sections were stained for regular light microscopic histology, after being etched, with HE and PAS. Usually, they were stained before etching with methylene-blue-azur B (Richardson *et al.*, 1960). Slides were dried and coverslipped as before.

### iii. Ultrathin resin sections

Ultrathin immunostaining procedures were modified after Schachter *et al.* (1983a, 1983c). 60 to 80 nm sections were cut on a Reichert-Jung ultracut, expanded with xylene vapor and mounted on 150 or 200 mesh nickel grids, previously coated with Formvar (Ladd Research Industries Ltd.). In some instances, sections were etched with either 1% or 2% saturated

ethanolic sodium ethoxide for 3-10 minutes or 5% or 10% aqueous  $H_2O_2$  for 5-10 minutes. An attempt to reduce background staining was also made by incubating grids in methanol for 5 minutes followed by 0.01%  $H_2O_2$  in methanol for 5 minutes to block endogenous peroxidase activity (Streefkerk, 1972). Grids were rinsed in distilled water, buffer, and incubated with normal goat serum for 5 minutes at room temperature. They were blotted with Whatman filter paper and submerged in drops of immune or normal rabbit serum (diluted 1:500-1:15,000 as before, optimally diluted 1:5000) or in the presence of 0.001%, 0.01% or 0.05% Triton X-100 detergent (triton) (Sigma Chemical Co.) in a humidified chamber after Pelliniemi and Van Noorden (1977). Incubations were carried out for 24-96 hours at 4°C. Staining took place at room temperature as follows. Grids were rinsed with buffer, blotted, floated for 5 minutes on normal goat serum, blotted, incubated for 5 minutes with FAB (diluted 1:10-1:90, optimally diluted 1:30), rinsed with buffer, blotted, floated on normal goat serum for 5 minutes, blotted, incubated for 5 minutes in PAP (diluted 1:40-1:100, optimally diluted 1:100), rinsed with buffer and loaded into a modified grid storage box with holes of 2 mm diameter drilled into each chamber to allow good flow of reagents. In some cases, an e.m. multiple grid staining kit was used (Analychem Corp. Ltd.). Without allowing them to dry completely, up to 20 grids were reacted together with a freshly prepared, filtered solution of 0.0125% DAB and 0.0025%  $H_2O_2$  in buffer. The reagents were magnetically stirred while grids were incubated for 3-5 minutes. Grids were rinsed twice in distilled water, dried and floated on 4% aqueous osmium for 15-30 minutes. They were again rinsed twice in distilled water and allowed to dry. In some cases, this osmium step was omitted.

Ultrathin resin-embedded tissues and antisera tested with the PAP method were as follows. The cat submandibular gland was tested with  $\alpha$ CSK and guinea-pig submandibular gland with  $\alpha$ GPSK and  $\alpha$ CSK. The GPCG and GPPG were examined with  $\alpha$ CGK and cat colon tissue was examined with  $\alpha$ CSK. Ultrastructure was examined by staining ultrathin sections with a double staining procedure (Pease, 1964). Grids were first processed in 5% uranyl acetate in methanol for 10-60 minutes, rinsed several times in methanol and dried.

Secondly, grids were floated on drops of freshly prepared lead citrate solution (Reynolds, 1963) for 1-3 minutes, rinsed in distilled water and dried before being observed in the electron microscope.

#### b. Avidin-biotin complex


##### i. Semi-thin resin sections

Avidin-biotin-complex staining procedures (Hsu *et al.*, 1981) (Figure 4) were used for semi-thin resin sections. 0.5-1.5  $\mu\text{m}$  sections were cut and dried onto gelatin-coated glass slides. Sections were treated with sodium ethoxide to remove resin and, where appropriate, treated with  $\text{H}_2\text{O}_2$  as described before. Enhancement of immunostaining by protease digestion was tried on GPCG sections by using 0.1% trypsin (from bovine pancreas) (Sigma Chemical Co.) in 0.05 M Tris buffer containing 0.1%  $\text{CaCl}_2$ , pH 7.6. Incubation took place at 37°C for 10-20 minutes and was followed by rinsing in buffer before proceeding. Following etching, sections were incubated in normal goat serum for 15-30 minutes. They were blotted and covered with either immune or normal rabbit serum (diluted 1:500-1:10,000, optimally diluted 1:5000). Incubations were carried out in humidity chambers for 24-72 hours at 4°C. At room temperature, sections were rinsed with buffer, blotted, incubated for 30-60 minutes in biotinylated goat anti-rabbit immunoglobulin (BG) (Cedarlane Laboratories Ltd.), diluted 1:200-1:500 (optimally diluted 1:200) with 0.05 M Tris buffer, pH 7.6, rinsed with buffer, blotted, incubated for 30-60 minutes with a complex of avidin DH (or avidin D) and biotinylated horseradish peroxidase (ABC) (Cedarlane Laboratories Ltd.), each diluted 1:100-1:1000 (optimally diluted 1:100) and rinsed with buffer. Peroxidase visualization was achieved by incubating sections for 10-15 minutes in a freshly prepared, filtered solution of 0.025% DAB and 0.005%  $\text{H}_2\text{O}_2$  in buffer. In some experiments, slides were further treated, following a distilled water rinse, by incubating them in a 1% aqueous solution of osmium for 10-15 minutes. Slides were rinsed in distilled water, dried and mounted with Permount.



FIGURE 4. Summarized ABC method of immunostaining. LM, light microscopically; EM, electron microscopically.

ABC



Semi-thin resin-embedded tissues and antisera tested with the ABC method were processed as follows. The cat submandibular gland was tested with  $\alpha$ CSK and guinea-pig submandibular gland with  $\alpha$ GPSK and  $\alpha$ CSK. The cat colon and rat colon were tested with  $\alpha$ CSK, the cat stomach was tested with  $\alpha$ CSK or  $\alpha$ CGK and the rat stomach was tested with  $\alpha$ CSK. Cat trachea was examined with  $\alpha$ CSK and GPCG and GPPG were tested with either  $\alpha$ CGK or  $\alpha$ NGF. Cat stomach sections were also stained for carbohydrates by the PAS method following ABC immunostaining. Light microscopic structure was examined by staining methods outlined before.

#### ii. Ultrathin resin sections

Ultrathin sections were cut, xylene-expanded, mounted on 150 or 200 mesh, Formvar-coated, nickel grids and processed by the ABC method. In some cases, digestion with 0.5% trypsin was performed in 0.05 M Tris buffer, pH 7.5, with 0.1%  $\text{CaCl}_2$  for 10-20 minutes at 37°C. Grids were rinsed in buffer, distilled water and incubated at room temperature with normal goat serum for 5 minutes. They were blotted and submerged in drops of immune or normal rabbit sera (diluted 1:1000-1:5000) in a humidity chamber. GPCG and GPPG were also incubated in antisera containing 0.1% saponin, used to aid antibody penetration and immunostaining. Incubations at 4°C were performed for 48-72 hours. Grids were rinsed in Tris buffer, blotted, floated for 5 minutes on normal goat serum, blotted, incubated for 5 minutes on BG (diluted 1:200), rinsed with buffer, blotted, floated on normal goat serum for 5 minutes, blotted, incubated for 5 minutes in ABC (components diluted 1:100), rinsed with buffer and loaded into a multiple grid staining unit, as before. Grids were reacted with a freshly prepared, filtered solution of 0.0125% DAB and 0.0025%  $\text{H}_2\text{O}_2$  in buffer. The reagents were stirred while grids were incubated for 3-5 minutes. Grids were rinsed twice in distilled water, dried and floated on 4% aqueous osmium for 15-25 minutes. After being rinsed twice with distilled water, they were allowed to dry.

Using this method, cat submandibular gland tissue was tested with  $\alpha$ CSK and GPCG and GPPG were tested with  $\alpha$ CGK. Ultrastructure was also examined by double staining methods outlined before.

### iii. Pre-embedding

GPCG or GPPG tissues were processed for pre-embedding ultrastructural immunostaining after the method of Pignat *et al.* (1982). Anaesthetized animals were perfused via the left ventricle with 50 ml of 4% PFA in 0.1 M phosphate buffer, pH 7.3, for 10 minutes. Although some animals were not perfused, all glands were removed, cut into small pieces and immersed in fixative for 2 hours at room temperature. A wash followed with three changes of buffer at 4°C. Usually the following steps were done at 4°C with solutions containing 0.1% saponin. Tissue pieces were incubated in primary antisera ( $\alpha$ CGK or normal rabbit serum, diluted 1:100-1:5000) for 48-72 hours, washed twice in buffer, incubated in normal goat serum for 10 minutes, incubated in BG (diluted 1:200) for 1 hour, washed twice in buffer, incubated in ABC (components diluted 1:100) and washed twice in buffer. The following took place at room temperature. Peroxidase activity was developed with a freshly prepared, filtered solution of 0.025% DAB and 0.005%  $H_2O_2$  in buffer. A buffer wash was followed by treatment of pieces with 2% osmium in 0.05 M cacodylate buffer for 1 hour. Two rinses with cacodylate buffer, for 10 minutes each, were followed by a final fixation with 3% GTA, 3% PFA in 0.05 M cacodylate buffer, pH 7.4, for 90 minutes. Pieces were washed twice with buffer, dehydrated through an ascending ethanol concentration gradient and embedded in Araldite resin via propylene oxide, as before. Ultrathin sections were cut, expanded with xylene and mounted on Formvar-coated grids before examination by electron microscopy.

### c. Protein A-colloidal gold

#### i. Ultrathin resin sections

The method of Roth *et al.* (1978) was followed for preparation of protein A-colloidal gold complex (PAG). 1 mg of protein A (Pharmacia Fine Chemicals) dissolved in 0.2 ml of distilled water was placed in a siliconized flask to which 10 ml of colloidal gold solution, pH 6.9 (ICN Pharmaceuticals Inc.), was added. The solution was mixed and then allowed to stand for 2 minutes before adding 0.2 ml of 1% polyethylene glycol (PEG) (Sigma Chemical Co.). Centrifugation at 60,000 g took place for 1 hour at 4°C. The pellet was resuspended in 10 ml PBS, pH 7.4, containing 0.02% PEG. In some cases, sections were treated with 5% or 10% H<sub>2</sub>O<sub>2</sub> for 5-10 minutes before the staining sequence began. Grids with mounted sections were incubated in 0.1 M PBS, pH 7.4, with 0.5% bovine serum albumin (BSA) (Miles Laboratories Inc.) for 5 minutes and blotted. Incubation in drops of immune or normal rabbit sera (diluted 1:100-1:10,000 in 0.1 M PBS, pH 7.4, with 0.5% BSA) for 1-2 hours at 4°C, or 12-72 hours at room temperature was followed by a buffer wash and blotting. Grids were incubated with PAG for 30-60 minutes at room temperature, washed in buffer with several changes of distilled water and dried. Before examining, grids were counterstained with 5% aqueous uranyl acetate for 60 minutes.

The PAG method was used unsuccessfully to examine cat submandibular gland with  $\alpha$ CSK and guinea-pig submandibular gland with  $\alpha$ GPSK. Particles of gold were distributed randomly on Formvar and resin sections. As before, ultrastructure was examined by double staining methods for the electron microscope.

### 3. Specificity and controls

Antibody specificities were tested by double diffusion analysis (Ouchterlony, 1958). Immunodiffusion plates were prepared with 1% agarose (Bio Rad Laboratories) in saline and wells were cut by a template kit (Miles Laboratories Inc.) so that each well held a volume of

35  $\mu$ l (see Figure 2). Antisera were tested against both purified antigen and tissue extracts. Antibody specificity during immunostaining was examined in some cases by absorbing specific rabbit antisera with pure antigen or by tissue extracts before use (Schachter *et al.*, 1983a). High dilutions of antisera and replacement of immune sera with normal rabbit sera were used to illustrate antibody specificity. As controls for method specificity in the PAP method, either FAB, PAP or  $H_2O_2$  were omitted from the staining sequence.

#### E. Microscopy and Photography

A photomicroscope II (Carl Zeiss) was used for light microscopy and tissues were photographed with either Panatomic X black and white or Ektachrome 64 color films (Kodak Canada Inc.). Tissues for electron microscopy were viewed with a Philips 300 electron microscope at 60 or 80 kv and photographed with Kodak fine grain positive 35 mm film. Printing of black and white negatives was on Ilfospeed 4 or 5 medium weight paper and color film was printed using Cibachrome paper and chemicals.

### III. RESULTS

#### A. Summarized Results of Immunostaining

Table 1 summarizes results of immunostaining various tissues which were fixed in several ways. Only wide classes of fixation are mentioned. Details of fixation procedures or exact concentrations of reagents are not included.

Two types of fixation were generally more successful than others in preserving antigenicity and structure. These were freeze-drying and CDI. CDI also appeared to have better success in preserving ultrastructure than freeze-drying. The next most successful type of fixation was 0.5% osmium. Higher concentrations of osmium were much less successful since antigenicity was lost. The last group of fixations, PFA-GTA combinations, were moderately successful, especially in the cat submandibular gland. It is interesting that in some tissues, such as GPCG and GPPG, only freeze-drying or CDI fixation was successful. Even 0.5% osmium, which worked for other tissues on which it was tried, did not work for the male sex glands. In conclusion, some fixations were good for immunostaining all tissues but others were suitable for only some tissues.

#### B. Localization of Kallikrein

##### 1. Submandibular Gland - Light and Electron Microscopy

###### a. Cat

In paraffin-embedded tissue, kallikrein was localized with  $\alpha$ CSK by the PAP method (Figures 5 and 6) Specific staining was seen only in apical regions of striated duct cells, which was similar to the findings of Schachter *et al.* (1980). This tissue was also embedded in Araldite resin and immunostained by the PAP method (Figures 7 and 8). This semi-thin section method was an improvement over the paraffin-embedded system for resolving both histological

TABLE 1. SUMMARIZED RESULTS OF  
IMMUNOSTAINING TISSUES FIXED IN VARIOUS WAYS  
(BY LIGHT MICROSCOPY)

Osmium

<u>Tissue</u>	<u>0.5%</u>	<u>1.0%</u>	<u>Freeze dried</u>	<u>CDI</u>	<u>PFA-GTA</u>
CSG	na	-	++	+++	+++
GPSG	na	na	+	++	++
CC	+	-	++++	++++	+
RC	+	-	++++	+++	++
CS	++++	+	+++	na	-
RS	+++	+	+	+++	-
CRT	++	na	na	+	+
GPCG	-	-	+++	++	++*
GPPG	-	-	++	++	-

+ (range + to ++++): Intensity of specific immunostain

-: No specific immunostain

CC: Cat colon

CRT: Cat respiratory tissue

CS: Cat stomach

CSG: Cat submandibular gland

GPCG: Guinea-pig coagulating gland

GPPG: Guinea-pig prostate gland

GPSG: Guinea-pig submandibular gland

RC: Rat colon

RS: Rat stomach


na: Experiment not attempted

\*: With PVVP treatment




**FIGURE 5.** Kallikrein localization in apical regions of striated duct cells of the cat submandibular gland. Tissue was fixed in 1% CDI, 0.5% GTA and 1% PFA, embedded in paraffin and stained with  $\alpha$ CSK (diluted 1:1000) by PAP method. L, lumen. (magnification X800).

**FIGURE 6.** Control section for Figure 5, incubated with normal rabbit serum (diluted 1:1000). L, lumen. (magnification X800).



**FIGURE 7.** Kallikrein localization in apical regions of striated duct cells of the cat submandibular gland. Tissue was fixed in 1% CDI, 0.5% GTA and 1% PFA. Semi-thin sections of resin-embedded tissue stained with  $\alpha$ CSK (diluted 1:5000) by PAP method. Compare resolution of immunostain to Figure 5. L, lumen. (magnification X800).

**FIGURE 8.** Control section for Figure 7, incubated with normal rabbit serum (diluted 1:1000). L, lumen. (magnification X800).



structure and localization of immunostain. Specific stain was again localized exclusively in apical regions of striated duct cells. Another fixative, used to preserve adequate ultrastructure while retaining antigenicity was 1% GTA-3% PFA (Figures 9 and 10) (Schachter *et al.*, 1983c). The PAP method was used and specific localization was only in apical secretory granules of striated duct cells and not in other cells or organelles such as nuclei, mitochondria or endoplasmic reticulum. Fixation for immunocytochemistry did not preserve ultrastructure as well as routine fixation. Figure 11 shows numerous, well-preserved secretory granules and mitochondria in the apical regions of striated duct cells. In Figure 12, large numbers of apical granules were crenated and the apical cytoplasm possessed many vacuoles, perhaps due to fixation artifact. Immunostaining of this region showed granules stained so as to give a studded appearance (Figure 13) characteristic of PAP complexes (Sternberger, 1979). Some intergranule deposition was observed, indicating the presence of intergranule antigen, reaction product effusion artifact or diffusion of antigen out of disrupted granules which was subsequently stained.

Resin-embedded tissue retained antigenicity more satisfactorily in several cases than did paraffin-embedded tissue. Tissues fixed in 2% CDI, 0.5% GTA and 1% PFA and embedded in either paraffin or resin were immunostained. Surprisingly, when tissues were fixed similarly, except with a 15 minute, 1% osmium postfixation step, only resin-embedded tissue retained antigenicity with light (Figures 14 and 15) or electron microscopy (Figures 16 and 17). Paraffin-embedded tissues did not immunostain specifically. Semi-thin or ultrathin sections, fixed in this way, had similar immunostain patterns to sections fixed by other means. In light microscopy, only apical regions of striated duct cells stained. Also, ultrastructurally, only apical granules of striated duct cells stained specifically.

Diffuse and variable reaction product was seen on occasion in acinar cell granules. This happened with both immune and control sera, which supported the view that it was non-specific. Because sodium ethoxide, used to etch sections, greatly diminished this staining, it was probably related to immunoglobulin-resin interactions. Endogenous peroxidase activity was

**FIGURE 9.** Electron microscopic immunolocalization of kallikrein in apical secretory granules of the cat submandibular gland, striated duct cells. Tissue was fixed in 1% GTA-3% PFA. Ultrathin resin-embedded tissue stained with  $\alpha$ CSK (diluted 1:1000) by PAP method. L, lumen. (magnification X5000).

**FIGURE 10.** Control section for Figure 9, incubated with normal rabbit serum (diluted 1:1000). L, lumen. (magnification X5000).





**FIGURE 11.** Ultrastructure of apical region of striated duct cell fixed by concentrated aldehyde fixative (2.5% GTA, 4% PFA and 1% acrolein). Note characteristic secretion granules (arrow), mitochondria and grainy cytoplasm (magnification X40000).

**FIGURE 12.** Ultrastructure of apical region of striated duct cell fixed by less concentrated immunocytochemical fixative (1% GTA-3% PFA). Note crenated secretion granules (arrow) and numerous vacuoles (V) in cytoplasm compared to Figure 11 (magnification X40000).

**FIGURE 13.** Localization of kallikrein in apical granules stained as in Figure 9, fixed as in Figure 12. Note studded appearance of PAP reaction product over secretion granules (arrow) (magnification X40000).





4)



**FIGURE 14.** Kallikrein localization in apical regions and secretions of striated duct cells of the cat submandibular gland post-fixed with osmium. Tissue was fixed in 2% CDI, 0.5% GTA and 1% PFA with a 1% osmium post-fixation step. Semi-thin resin-embedded tissue stained with  $\alpha$ CSK (diluted 1:2000) by PAP method. L, lumen. Compare to Figure 7 (magnification X800).

**FIGURE 15.** Control section for Figure 14, incubated with normal rabbit serum (diluted 1:2000). L, lumen. (magnification X800).



**FIGURE 16.** Electron microscopic localization of kallikrein in apical granules of cat submandibular gland, striated duct cells, fixed as in Figure 14. Ultrathin resin-embedded tissue stained with  $\alpha$ CSK (diluted 1:2000) by PAP method. L, lumen. Compare to Figure 9 (magnification X8000).

**FIGURE 17.** Control section for Figure 16, incubated with normal rabbit serum (diluted 1:2000). L, lumen. (magnification X8000).



ruled out since methanol-H<sub>2</sub>O<sub>2</sub> treatment did not reduce acinar staining.

### b. Guinea-pig

Specific immunostaining was in apical granules of guinea-pig submandibular gland striated duct cells (Figures 18 and 19) with the ultrastructural PAP method. Apical granules were less susceptible to distortion by fixative osmolarity or concentration and were easier to preserve than those of the cat submandibular gland.

Because the semi-thin resin section method correlated with ultrastructural experiments in the cat submandibular gland and had good resolution, it was used almost exclusively for light microscopy in the guinea-pig submandibular gland. Paraffin-embedded tissue was only used to confirm  $\alpha$ GPSK activity and the PAP system used earlier by Schachter *et al.* (1980). Different fixatives, embedding media, and immunostains developed patterns of immunostain which were similar to the cat submandibular gland. Our concentrated aldehyde-based fixative produced guinea-pig submandibular gland ultrastructure which was "normal" and similar to that of Schachter *et al.* (1983b). When concentrations of cross-linking reagents were reduced, both ultrastructure and antigenicity were retained. Etching of resin for ultrastructural studies was not always done and had no obvious effect. Endogenous peroxidase activity, known to be present in acinar granules (Bloom *et al.*, 1974), was absent in either sodium ethoxide-etched semi-thin resin sections or unetched ultrathin sections. In ultrathin PAP localizations,  $\alpha$ GPSK stained apical granules of striated ducts better than  $\alpha$ CSK.

## 2. Colon

### a. Cat



Specific staining of kallikrein was seen only in mucous (goblet) cells of the cat colon, though not all mucous cells stained. No staining was observed in columnar epithelial cells, lamina propria or non-mucosal structures. Several different immunocytochemical methods were

7

**FIGURE 18.** Electron microscopic localization of kallikrein in apical secretory granules of the guinea-pig submandibular gland striated duct cells. Tissue fixed with 5% CDI and 0.5% GTA, embedded in resin and stained with  $\alpha$ GPSK (diluted 1:1000) by PAP method. L, lumen. (magnification X8000).

**FIGURE 19.** Control section for Figure 18, incubated with normal rabbit serum (diluted 1:1000). L, lumen. (magnification X8000).

6






used.

Paraffin-embedded sections, stained by the PAP method, showed specific immunostaining in mucous cells or in their luminal discharges (Figures 20 and 21). Some mucous cells were discharging or had apparently discharged their contents at the time of fixation, and therefore did not stain. Secretion may have occurred physiologically or may have been due to an artifact of fixation or embedding. Fixation of this tissue was optimal with 1% CDI. Other fixatives for paraffin-embedded tissues, which produced specific staining, were not useful due to weak reactions or poor structure.

Semi-thin resin sections, stained by PAP or ABC methods, also showed specific staining only in mucous cells or in their luminal discharges (Figures 22 and 23). With semi-thin sections, resolution of the immunostain was better than in paraffin sections and stain was in the mucous granule compartment of goblet cells. Strands of immunostaining secretion were also observed. Furthermore, less discharge of cell contents was seen compared to paraffin-embedded tissue. Since both of these tissues were fixed with 1% CDI, the increased secretion in paraffin-embedded tissues was likely an artifact due to this embedding method. Other chemical fixatives did not produce as much specific staining as 1% CDI. PLP, PFA and osmium improved structure but reduced antigenicity.  $H_2O_2$  etching was necessary for osmium-fixed tissue. Ethanol-fixed tissue did not retain antigenicity and saponin treatment did not increase antigenicity.

Freeze-dried tissues showed good localization of kallikrein in semi-thin resin sections (Figures 24 and 25). Specific staining was seen only in mucous cells, whether at the base or neck of crypts or at the surface. Tissues from different regions of the colon showed no difference in staining of mucous cells. Antisera and immunostaining reagents were used at optimal concentrations described before, except that primary sera in the ABC system were usually more dilute.

SDS treatment did not improve antisera or reagent penetration. All systems were strongly stained with  $\alpha$ CSK but in the semi-thin PAP system,  $\alpha$ RPK and  $\alpha$ RIK also produced




**FIGURE 20.** Kallikrein localization in mucous cells (arrows) and secretions of cat colon. Tissue fixed in 1% CDI, embedded in paraffin and stained with  $\alpha$ CSK (diluted 1:5000) by PAP method. L, lumen. (magnification X300).

**FIGURE 21.** Control section for Figure 20, incubated with normal rabbit serum (diluted 1:5000) (magnification X300).



**FIGURE 22.** Kallikrein localization in mucous cells (arrows) and secretions of the cat colon. Tissue fixed with 1% CDI, embedded in resin. Semi-thin sections stained with  $\alpha$ CSK (diluted 1:1000) by PAP method. L, lumen. Compare resolution of immunostain to Figure 20 (magnification X800).

**FIGURE 23.** Control section for Figure 22; incubated with normal rabbit serum (diluted 1:1000) (magnification X800).



**FIGURE 24.** Kallikrein localization in mucous cells (arrows) and secretions of the cat colon. Tissue was freeze-dried, without fixation. Semi-thin resin-embedded sections stained with  $\alpha$ CSK (diluted 1:5000) by PAP method. Compare to Figure 22 (magnification X800).

**FIGURE 25.** Control section for Figure 24, incubated with normal rabbit serum (diluted 1:5000) (magnification X800).



weak staining in mucous cells. Other antisera which produced no staining in this system were  $\alpha$ GPSK,  $\alpha$ HSK and  $\alpha$ CGK. Staining was absent from sections incubated with normal rabbit sera or antisera which were preabsorbed with pure antigen or extract (Schachter *et al.*, 1980).


Ultrastructural immunolocalization of kallikrein was possible in preliminary experiments. The PAP method produced specific staining in goblet cells and in strands of secretions (Figures 26 and 27). Although ultrastructure was poor and mucous granules were completely disrupted, staining patterns supported the idea that kallikrein was associated with mucous granules. Further ultrastructural immunolocalization in the colon may require development of better fixatives than 1% CDI.

#### b. Rat


Kallikrein in the rat colon was localized only in mucous cells, in luminal secretions and on the surface (Figures 28 and 29). Mucous cells of ascending and descending regions of rat colon stained equally to the cat colon. Background staining in controls and immune sections was higher and reaction product often appeared in nuclear regions and in columnar epithelial cells. Mucous granule compartments of goblet cells in control sections were unstained. The ABC method produced less background stain than the PAP method, though specific staining was also reduced.  $\alpha$ CSK primary antisera were used exclusively and at higher dilutions in the ABC method than the PAP method.

Several fixation methods preserved kallikrein antigenicity in rat colon tissue and both ABC and PAP semi-thin section methods successfully localized kallikrein. Immunostaining indicated an identical distribution of kallikrein in both cat colon and rat colon. Light microscopic structure and antigenicity were well preserved by the copper block method of freezing. Surfaces of colon often contacted the copper surface first (Figures 28 and 29) and therefore froze rapidly. Ice crystal artifact was only in areas which were not at the colon surface. Cellular detail in resin-embedded, freeze-dried, unfixed preparations was comparable to chemical-fixed tissues. Other fixatives such as 1% CDI, PLP or 2% osmium showed good structure but only







**FIGURE 26.** Electron microscopic localization of kallikrein in the mucous granule region of cat colon mucous, goblet cells (arrow). Ultrathin resin-embedded tissue, fixed with 1% CDI and stained with  $\alpha$ CSK (diluted 1:5000) by PAP method. L, lumen. (magnification X5000).




**FIGURE 27.** Control section for Figure 26, incubated with normal rabbit serum (diluted 1:5000) (magnification X5000).





A black and white micrograph showing a cross-section of rat colon tissue. The tissue is stained to show kallikrein localization. An arrow points to a specific mucous cell. The lumen is labeled 'L'. The magnification is X800.

**FIGURE 28.** Kallikrein localization in mucous cells (arrow) of the rat colon. Semi-thin, freeze-dried, without fixation, resin-embedded tissue sections stained with  $\alpha$ CSK (diluted 1:1000) by PAP method. L, lumen. (magnification X800).

A black and white micrograph showing a control section of rat colon tissue. The tissue is incubated with normal rabbit serum. The lumen is labeled 'L'. The magnification is X800.

**FIGURE 29.** Control section for Figure 28, incubated with normal rabbit serum (diluted 1:1000). Note high level of background stain with PAP method. L, lumen. (magnification X800).



weak antigenicity. Reducing the osmium concentration to 0.5% increased the immunostaining, making this fixative useful.

### 3. Stomach

#### a. Cat

In liquid-fixed cat stomach tissue, kallikrein was localized in surface epithelial mucous cells and in mucous cells of gastric pits, necks and glands (Figures 30 and 31). Specific staining was further resolved in apical mucous granule compartments of these cells with semi-thin sections. Nuclear or basal regions of epithelial cells did not stain. Staining of mucous cells at the surface was usually interrupted, probably because of sloughing of apical cells.

Osmium treatment, which followed immunostaining, was used in some tissues to enhance staining which was due to peroxidase reaction product. Although osmium treatment increased background staining in control sections (Figure 31), it also enhanced the peroxidase reaction product in immune sections (Figure 30). Specific staining was relatively greater than in immune sections which were not stained with osmium (Figure 32).

Different regions of the stomach stained differently due to the pattern of mucous cell distribution. The antrum, possessing the most mucous cells, stained intensely, the corpus, with fewer mucous cells, stained less, and the fundus showed little specific staining.

The stomach antrum was optimally preserved for light microscopic structure and antigenicity with 0.5% osmium (Figures 30 to 33). Liquid fixation with other combinations of CDI, GTA or PPA were less successful due to antigen masking. Saponin addition or preliminary studies with EAI treatment did not enhance specific stain. PLP and ethanol fixatives caused loss of antigenicity. When osmium concentrations were above 0.5%, structure was improved but antigenicity was reduced.

Freeze-dried cat stomach tissue also showed immunolocalization of kallikrein in epithelial mucous cells (Figures 34 to 41). Specific staining was in the mucous granule region of

**FIGURE 30.** Kallikrein localization in apical regions of mucous cells (arrow) of the cat stomach antrum. Tissue was fixed in 0.5% osmium. Semi-thin resin-embedded tissue stained with  $\alpha$ CSK (diluted 1:1000) by ABC method with osmium post-stain treatment. L, lumen. (magnification X800).


**FIGURE 31.** Control section for Figure 30, incubated with normal rabbit serum (diluted 1:1000). Note background stain in mucous cells. L, lumen. (magnification X800).



L




L



**FIGURE 32.** Kallikrein localization in apical regions of mucous cells (arrow) of the cat stomach antrum. Tissue was fixed as in Figure 30, in resin-embedded tissue stained with  $\alpha$ CSK (diluted 1:1000) by ABC method without gold post-stain treatment. L, lumen. Compare immunostain to that of Figure 30 (magnification X800).


**FIGURE 33.** Control section for Figure 32, incubated with normal rabbit serum (diluted 1:1000). Compare background stain to Figure 31 (magnification X800).





# 2 2

OF / DE






**FIGURE 34.** PAS stain with immunostain for kallikrein on semi-thin, freeze-dried without fixation resin-embedded cat stomach antrum.  $\alpha$ CSK (diluted 1:1000) used in ABC method. Note kallikrein localization in periphery of mucous cells (arrow) and in secretions and PAS stain in central region of mucous cells and in secretions. L, lumen. (magnification X625).

**FIGURE 35.** Kallikrein localization in periphery of mucous cells (arrow) and in secretions of cat stomach antrum. Semi-thin, resin-embedded tissue stained with  $\alpha$ CSK (diluted 1:1000) by ABC method. L, lumen. Compare immunostain to Figure 34 (magnification X625).

**FIGURE 36.** Control section for Figures 34, 35 and 37, incubated with normal rabbit serum (diluted 1:1000). L, lumen. (magnification X250).

**FIGURE 37.** PAS stain with immunostain for kallikrein on cat stomach antrum sections.  $\alpha$ CSK (diluted 1:1000) used in ABC method. Note that distribution of kallikrein and PAS stain were not the same. L, lumen. (magnification X250).



**FIGURE 38.** Kallikrein localization in the mucous cells (arrow) of cat stomach antrum. Semi-thin, freeze-dried without fixation, resin-embedded tissue stained with  $\alpha$ CSK (diluted 1:1000) by PAP method. L, lumen. (magnification X800).

**FIGURE 39.** Control section for Figure 38, incubated with normal rabbit serum (diluted 1:1000). L, lumen. (magnification X800).

**FIGURE 40.** Kallikrein localization in mucous cells (arrow) and surface secretions of cat stomach antrum. Semi-thin, freeze-dried, resin-embedded tissue stained with  $\alpha$ CSK (diluted 1:1000) by ABC method. L, lumen. Compare to Figure 38 (magnification X800).

**FIGURE 41.** Control section for Figure 40, incubated with normal rabbit serum (diluted 1:1000). L, lumen. Compare to Figure 39 (magnification X800).



cells of the surface, pit, neck and gland, and in secretions along the surface. Copper block freezing preserved cell integrity, though tissues were deformed from their regular appearance due to the tissue's impact with the copper surface. Fixation with osmium vapor did not mask antigenicity and improved structure. Figure 35 shows kallikrein localization in surface and pit epithelial mucous cells. Staining by the PAP method was in the peripheral zone of mucous compartments of surface cells. Peripheral staining indicated that either kallikrein antigenicity was altered in the central zone of intact surface cells or secretion by surface cells caused kallikrein to be lost from central regions. Immune sections were stained with PAS to test the second possibility. PAS staining and PAP immunostaining in the same sections revealed central mucous regions of surface cells that were positive for carbohydrate but not for kallikrein (Figures 34 and 37). Some regions also had cells which reacted to either PAS stain or kallikrein immunostain, but not both (Figure 37). Kallikrein antigenicity was altered regionally within cells either as an artifact of fixation or embedding, or by intracellular changes due to chemical processing.

PAP and ABC immunostaining methods were compared with serial semi-thin cat stomach sections. Osmicated freeze-dried tissues, which were frozen by the copper block method, were immunostained for kallikrein by either the PAP or ABC method (Figures 38 to 41). Localization of kallikrein was in surface, pit, neck and gland mucous cells by both methods, but surface mucous staining was only in ABC-stained sections. Background staining in PAP control sections (Figure 39) was higher than in ABC control sections (Figure 41), and a better "signal to noise" ratio (i.e. specific to background staining) was obtained in the ABC method. Higher dilutions of  $\alpha$ CSK primary antisera also produced good staining.  $\alpha$ RPK and  $\alpha$ CGK did not produce specific immunostaining for kallikrein in cat stomach tissue.

#### b. Rat

Paraffin-embedded tissue preserved antigenicity and specific staining, using  $\alpha$ CSK, was found in mucous neck cells of the rat stomach (Figures 42 and 43). Staining was not found in

**FIGURE 42.** Kallikrein localization in mucous neck cells (arrow) of the rat stomach antrum. Tissue was fixed with 1% CDI, embedded in paraffin and stained with  $\alpha$ CSK (diluted 1:1000) by PAP method. L, lumen. (magnification X300).

**FIGURE 43.** Control section for Figure 43, incubated with normal rabbit serum (diluted 1:1000) (magnification X300).



L



surface mucous cells or in mucosal secretions.  $\alpha$ RPK and  $\alpha$ RIK did not produce immunospecific staining. Semi-thin resin tissue, fixed similarly, also preserved antigenicity and with better resolution. Specific staining was in the apical region of surface, pit and neck epithelial mucous cells (Figures 44 and 45). The surface mucous layer also stained specifically using either the PAP or ABC method and  $\alpha$ CSK primary antisera. The lack of staining in surface mucous cells in paraffin-embedded tissue may have been due to loss of antigenic material during fixation or embedding.

Different immunocytochemical techniques were used to stain kallikrein in rat stomach tissue. Regional distribution of kallikrein in the rat stomach was similar to that of the cat stomach. Fixatives using higher concentrations of CDI, PFA mixtures or high concentrations of osmium did not preserve antigenicity. Semi-thin sections, fixed with 0.5% osmium, were successful in localizing kallikrein with the ABC method (Figure 46). Surface, pit and neck mucous cells stained specifically. Sections were also immunostained to compare the effects of various chemical treatments to preserve antigenicity. Enhanced staining resulted when these chemical treatments were used. EAI treatment (Figure 47) resulted in more intense staining than PVPP treatment (Figure 48). Control sections for untreated, EAI or PVPP-treated immune sections did not stain.


#### 4. Respiratory Tract

##### a. Cat

Preliminary results localized kallikrein in cat trachea. Semi-thin resin sections of tissue were used which were fixed with a 1% CDI-3% PFA mixture and treated with EAI. Specific staining was localized in submucosal glands (Figures 49 and 50). Frequently, using osmium-fixed or other tissues, specific staining was in surface mucous, goblet cells. Although cat trachea had a variable staining pattern in immune sections, control sections were always devoid of stain.

**FIGURE 44.** Kallikrein localization in mucous cells (arrow) and surface secretions of the rat stomach antrum. Tissue was fixed in 1% CDI. Semi-thin resin-embedded tissue stained with  $\alpha$ CSK (diluted 1:5000) by PAP method. L, lumen. Compare resolution and distribution of immunostain to Figure 42 (magnification X800).


**FIGURE 45.** Control section for Figure 44, incubated with normal rabbit serum (diluted 1:5000) (magnification X800).



**FIGURE 46.** Kallikrein localization in apical regions of mucous cells (arrow) of the rat stomach antrum. Tissue was fixed in 0.5% osmium. Semi-thin resin-embedded tissue stained with  $\alpha$ CSK (diluted 1:5000) by ABC method. L, lumen. (magnification X800).




**FIGURE 47.** Kallikrein localization in apical regions of mucous cells (arrow) of the rat stomach antrum treated, fixed as in Figure 35, with EAI. Semi-thin resin-embedded tissue stained with  $\alpha$ CSK (diluted 1:5000) by ABC method. L, lumen. Compare immunostain to Figure 46 (magnification X800).

**FIGURE 48.** Kallikrein localization in apical regions of mucous cells (arrow) of the rat stomach antrum, fixed as in Figure 46, treated with PVPP. Semi-thin resin-embedded tissue stained with  $\alpha$ CSK (diluted 1:5000) by ABC method. L, lumen. Compare immunostain to Figures 46 and 47 (magnification X800).



**FIGURE 49.** Kallikrein localization in submucosal glands (arrow) of cat respiratory tissue. Tissue was fixed in 1% CDI-3% PFA. Semi-thin resin-embedded tissue stained with  $\alpha$ CSK (diluted 1:1000) by PAP method. G, goblet cell; L, lumen. (magnification X800).

**FIGURE 50.** Control section for Figure 49, incubated with normal rabbit serum (diluted 1:1000) (magnification X800).





## C. Localization of Kallikrein and Nerve Growth Factor

### 1. Male Sex Glands - Light Microscopy

#### a. Guinea-pig



Kallikrein and NGF were localized by light microscopy in male sex glands. With  $\alpha$ CGK antisera, kallikrein was localized in secretory epithelial cells of the GPCG and GPPG. With  $\alpha$ NGF antisera, NGF was localized in epithelial cells of GPPG tissue. Antigen distribution depended on the fixation method.

Figures 51 and 52 show the localization of kallikrein in secretory epithelial cells. Semi-thin sections stained diffusely in apical cytoplasm and in secretions. This pattern was similar to that described by Schachter *et al.* (1978). Staining was in most cells and predominated in the granular endoplasmic reticulum-Golgi region described by Wong and Tse (1981). Peripheral, but not central staining occurred in most apical blebs, characteristic of apocrine secretory mechanisms. Specific staining was not found in nuclear, basal or non-epithelial regions. Semi-thin sections were also prepared from freon-frozen, freeze-dried GPCG tissue and immunostained for kallikrein (Figures 53 and 54). Distribution of specific stain was slightly different from liquid-fixed tissue. Although secretions and peripheral regions of apical blebs stained intensely, apical cytoplasmic regions of epithelial cells did not. This may have been due to large-scale ice crystal artifact present in all cells which caused a loss of antigenicity.

Antigen preservation was successful using PVPP treatment. GPCG tissue, fixed with 6% PFA and treated with PVPP, stained specifically in secretory regions of epithelial cells (Figure 55). Control sections were devoid of stain (Figure 56). Tissues that were fixed identically but without PVPP treatment did not stain. Although PVPP treatment enhanced antigenicity in tissues fixed by relatively high aldehyde concentrations (6% PFA), neither it nor EAI treatment enhanced staining in tissues which were fixed with 1% CDI-2% PFA. Fixatives containing GTA or osmium caused antigen masking though structure was improved. Other



✓  
FIGURE 51. Kallikrein localization in secretory epithelial cells of the guinea-pig coagulating gland. Semi-thin resin-embedded tissue, fixed with 1% CDI and 1% PFA and stained with  $\alpha$ CGK (diluted 1:5000) by PAP method. L, lumen. (magnification X800).

FIGURE 52. Control section for Figure 51, incubated with normal rabbit serum (diluted 1:5000). L, lumen. (magnification X800).



**FIGURE 53.** Kallikrein localization in secretory epithelial cells of the guinea-pig coagulating gland. Tissue was freeze-dried, without fixation. Semi-thin resin-embedded tissue stained with  $\alpha$ CSK (diluted 1:5000) by PAP method. L, lumen. Compare to Figure 49 (magnification X800).

**FIGURE 54.** Control section for Figure 53, incubated with normal rabbit serum (diluted 1:5000). L, lumen. (magnification X800).



**FIGURE 55.** Kallikrein localization in secretory epithelial cells of the guinea-pig coagulating gland fixed in 6% PFA, treated with PVPP. Semi-thin resin-embedded tissue stained with  $\alpha$ CGK (diluted 1:1000) by ABC method. L, lumen. (magnification X800).


**FIGURE 56.** Control section for Figure 55, incubated with normal rabbit serum (diluted 1:1000) (magnification X800).



fixatives produced poor structure and weak antigenicity. Trypsin treatment did not unmask antigen and saponin did not enhance immunostain.  $\alpha$ CGK produced immunostain with PAP or ABC methods, but  $\alpha$ NGF did not immunostain the GPCG with the ABC semi-thin method. Ultrastructural immunostaining was unsuccessful despite trypsin and saponin treatments and pre-embedding immunocytochemistry. Despite semi-thin sections which immunostained, ultra-thin sections which were treated identically only had non-specific staining in immune or control sections.


Kallikrein and NGF were immunolocalized in copper block-frozen, freeze-dried GPPG tissue. Figure 57 shows specific staining in apical cytoplasm of epithelial secretory cells of the dorsal prostate. Secretions also stained. Epithelial cells of control sections were devoid of stain (Figure 58). The apical cytoplasm and secretions of epithelial cells also stained specifically with  $\alpha$ NGF antisera (Figure 59), which was similar to the results of Shikata *et al.* (1984) using paraffin-embedded tissues. Control sections did not stain (Figure 60).






**FIGURE 57.** Kallikrein localization in apical cytoplasm and secretions of epithelial secretory cells of the guinea-pig prostate gland. Tissue was freeze-dried without fixation. Semi-thin resin-embedded tissue stained with  $\alpha$ CGK (diluted 1:5000) by PAP method. L, lumen. (magnification X800).

**FIGURE 58.** Control section for Figure 57, incubated with normal rabbit serum (diluted 1:5000). L, lumen. (magnification X800).



**FIGURE 59.** NGF localization in apical cytoplasm and secretions of epithelial secretory cells of the guinea-pig prostate gland. Tissue was freeze-dried without fixation. Semi-thin resin-embedded tissue stained with  $\alpha$ NGF (diluted 1:1000) by PAP method. L, lumen. (magnification X800).

**FIGURE 60.** Control section for Figure 59, incubated with normal rabbit serum (diluted 1:1000). L, lumen. (magnification X800).



## IV. DISCUSSION

### A. Kallikrein and Related Proteases in Salivary Glands

Many chemically related proteases exist in salivary glands and saliva. Results in this thesis show the localization of kallikrein, one of these, in the apical granules of submandibular duct cells. Salivary glands from different mammals also contain various esterolytic enzymes and other macromolecules including kallikrein (Brandtzaeg *et al.*, 1976), tonin (Ørstavik *et al.*, 1982a; Zacharatos *et al.*, 1983), sialotinin, a vasopressor substance (Barion *et al.*, 1976), renin (Angeletti *et al.*, 1967; Barka, 1980; Bing *et al.*, 1980; Hiramatsu *et al.*, 1981), esteroproteases A, B, C and D (Boesman *et al.*, 1976) and other variously named serine proteases. Erythropoietin (Fava-de-Moraes *et al.*, 1979) is found in salivary gland duct cells along with AM<sub>1</sub>, a secretory glycoprotein (Nieuw Amerongen *et al.*, 1981). Localization studies indicate that most proteases above are present in cells of the salivary gland ducts.

Many esteroproteases are found to resemble each other chemically but to possess different activities. Protein A (Bennick, 1975), a calcium-binding protein, is probably identical to protein III or IV. Protein C also resembles protein I or II. Esteroproteases A and D from mouse submandibular glands (Boesman *et al.*, 1976) are functionally separate, since protease D is thought to be identical to the epidermal growth factor (EGF)-binding protein. Tonin, identical to esterase A (Brandtzaeg *et al.*, 1976), represents 10% of the protein from rat submandibular gland (Thibault and Genest, 1981) and co-exists in the gland with kallikrein (Ørstavik and Glenner, 1978). Similarly, in mouse submandibular glands, renin and kallikrein co-exist (Barka, 1980).

Among salivary growth factors with profound effects are EGF, which stimulates the proliferation of various epithelia (Gospodarowicz, 1981), and NGF, which affects development of the sympathetic adrenergic system. EGF, isolated by Cohen (1959), was localized in secretory granules of duct cells of rodent submandibular glands (Turkington *et al.*, 1971; Gresik and Barka, 1977; Tanaka *et al.*, 1984; Murphy *et al.*, 1980).

A connection was made between salivary EGF and gastrointestinal function by the observation that this factor stimulated gastric mucosal growth (Oka *et al.*, 1983; Dembinski and Johnson, 1985). EGF from duct cells is secreted both into saliva and into the circulation (Hirata and Orth 1979; Bing *et al.*, 1980; Nexø *et al.*, 1984). Since aggressive behaviour and  $\alpha$ -adrenergic stimulation caused stimulation of salivary secretion of EGF and renin, possibly these factors may be involved in stress and related events, e.g., wound healing.

NGF co-exists with kallikrein in ducts of submandibular glands and is induced by testosterone and thyroxine (Hendry and Iverson, 1973; Ishii and Shooter, 1975). A part of the molecular complex of mouse salivary NGF is chemically similar to tissue kallikrein (Fiedler and Fritz, 1981),  $\beta$ -NGF endopeptidase and EGF-binding protein (Bothwell *et al.*, 1979). NGF is involved in accelerating the rate of wound healing in sialoadrenalectomized animals and the process is thought to be partially via a licking process (Li *et al.*, 1980). Schwab *et al.* (1976) suggested that NGF might function in protease inactivation during storage or protease synthesis, unrelated to effects on nervous tissue. Despite uncertainties in the physiology of NGF and other growth factors, their sources have served as aids in chemical characterization of the complexes.

Concentrations of both kallikrein and arginine esterase A can be increased by testosterone in the rat submandibular gland (Chao *et al.*, 1982). The quantity of submandibular proteases is under androgenic control. A digestive role for these enzymes is difficult to imagine because of the large quantitative difference between glands of male or female mice or the absence of enzymes prior to sexual maturity.

The idea that salivary gland kallikrein is involved in functional hyperaemia is not new (Hilton and Lewis, 1956). Although several counter-arguments have been expressed which indicate that kallikrein is largely secreted into ducts rather than into the circulation (Schachter *et al.*, 1977) (see Results), movement of the enzyme towards the vasculature is also apparent. Ørstavik *et al.* (1980b) showed that kallikrein injected retrogradely into the rat submandibular ducts could pass into the circulation. Enzyme secretion into the interstitium may possibly be via

granule-like structures which immunostain for kallikrein at the base of duct cells (Ørstavik *et al.*, 1980a; Simson *et al.*, 1983). Kallikrein was also located in the interstitium and basement membrane of salivary gland ducts (Ørstavik *et al.*, 1975; Kimura and Moriya, 1984a). A role for kallikrein in local blood flow regulation was also suggested by Rabito *et al.* (1982), who concluded that kallikrein was secreted into interstitial and vascular compartments since there was an arterial-venous difference in kallikrein concentration, and by Ørstavik *et al.* (1982b), who found an increase in blood flow on nerve stimulation after increasing kinin levels with converting enzyme inhibitor.

Schmid and Heidland (1984) found that exocrine salivary kallikrein secretion is elevated in rats with hypertension, though a rise in blood pressure may not be the cause of enhanced kallikrein secretion in hypertensive rats.

From the mouse submandibular glands, a related group of proteases (A, B, C and D) was found by Schenkein *et al.* (1969), who suggested that limited proteolysis converted "A" to "D". These esteroproteases were also shown to cause differentiation in muscle tissue of chick embryos. Furthermore, several other salivary esteroproteases were thought to be involved in processing precursors of NGF and EGF (Berger and Shooter, 1977; Bothwell *et al.*, 1979).

The primary and striking luminal localization of kallikrein in salivary ducts suggests that kallikrein is secreted into saliva. It was suspected that kallikrein at the luminal surface released kinins, which could in turn increase the apical permeability to ions and thus indirectly promote ion transfer either into or out of the duct lumen (Simson *et al.*, 1983). Tonin was also suspected to play a role in ion transport in the submandibular gland via generation of angiotensin II (Zacharatos *et al.*, 1983). Curiously, a selective intracellular secretion mechanism occurs for the release of kallikrein and tonin, since different  $\alpha$  and  $\beta$  agonists cause differential protease release (Zacharatos *et al.*, 1983).

## B. Kallikreins, Nerve Growth Factor and Related Proteases in Male Sex Glands and Their Possible Roles

Results in this thesis show the presence of tissue kallikrein (CGK) in epithelial cells of prostate and coagulating glands and NGF in epithelial cells of the prostate gland. These findings are similar to those of Schachter *et al.* (1978) for kallikrein, and of Shikata *et al.* (1984) for NGF. Other related factors such as EGF (Shikata *et al.*, 1984) or androgen-dependent esterases (Isaacs and Shaper, 1983) are synthesized, stored and secreted by prostatic tissue. Tissue kallikreins are also found in porcine (Fink *et al.*, 1979) and human seminal fluid (Mann *et al.*, 1980; Geiger and Clausnitzer, 1981; Fink *et al.*, 1985), which probably originates from prostate glands.

In the male genital tract, as elsewhere, kallikrein may function physiologically by producing kinin via action on the ubiquitously occurring kininogens (Geiger and Clausnitzer, 1981). A common function for kallikrein in male sex glands based on its ability to release kinin is unlikely for mammals generally since only in the guinea-pig sex glands does the arginine esterase possess the ability to release kinin (Barton *et al.*, 1973).

Seminal plasma esterase has several molecular similarities to tissue kallikreins though some important ones are lacking. For example, canine prostate esterase does not release a kinin (Dubé *et al.*, 1983). The major androgen-dependent protein accounts for over 90% of the total protein present. Absence of the protein in prostates from castrated animals implies androgen dependence for synthesis, and correlates well with the morphological atrophy of the epithelium, measured by loss of secretory granules (Isaacs and Shaper, 1983).

It was proposed that motility of human spermatozoa is regulated by a kallikrein-kinin system (Schill and Haberland, 1974). Systemic kallikrein treatment of human patients improved sperm motility, volume of semen, sperm concentration and spermatozoa motility. Kallikrein also enhances sperm penetration through cervical mucus (Schill *et al.*, 1974). Evidence suggests that during normal capacitation of mammalian spermatozoa, a glycoprotein coat on the sperm plasma membrane is removed or modified, perhaps by protease action, and acceleration of the



acrosome reaction of hamster spermatozoa by proteolytic enzymes such as kallikrein seems to be partly due to alteration of sperm counts (Shinohara *et al.*, 1985).

Possible roles of NGF and other growth factors may be unrelated to, but in addition to, their effects on neuronal tissue. In the prostate gland, since it is an exocrine organ like the salivary glands, it is reasonable to speculate that these substances have some role related to the reproductive system.

### C. Gastrointestinal Kallikreins and Their Possible Roles

A large body of evidence supports the concept of kallikreins playing fundamental roles in transport of electrolytes, glucose, and amino acids in the gastrointestinal tract (Moriwaki *et al.*, 1977).

Cuthbert and Margolius (1982) found that kallidin enhances gut chloride secretion, involving a prostaglandin pathway. Kinin receptors were thought to regulate this process (Manning and Snyder, 1982) despite an apparent location on the basolateral surface of colon epithelial cells and in the lamina propria and despite the fact kinins only stimulated intestinal electrogenic chloride secretion when applied to the basolateral (serosal) side of tissue (Cuthbert and Margolius, 1982; Manning and Snyder, 1982). In the intestine, chloride secretion occurs in crypt cells but not in surface cells (Welsh *et al.*, 1982) and it may be that the inability of kinin to effect transport at the surface is due to its inability to penetrate into the crypts (Kirkland *et al.*, 1985). Crocker and Willavoys (1975) found that bradykinin could either stimulate or inhibit transepithelial water transfer. Different mechanisms were proposed since inhibitory effects depended on whether the kinin was added to the serosal or mucosal bathing solution. Kallikrein administration into the intestine enhances amino acid absorption, which suggests that gastrointestinal kallikrein is related to nutrient absorption (Moriwaki *et al.*, 1977; Moriwaki and Fujimori, 1981). Geiger *et al.* (1977) suggested that kinins may function in absorption, and possibly affect transport of glucose and amino acids in the gut.

In the kidney tubule, there is evidence for kinin-mediated transport at both apical and basolateral sides of the cell (Kauker, 1980). Cuthbert *et al.* (1985) also found kinin influenced transport at both surfaces of cultured tubule cells. In the toad urinary bladder (Orce *et al.*, 1980) and rat renal cortical cells (Margolius and Chao, 1980), kallikrein-like proteases were effective near the mucosal surfaces in stimulating sodium transport.

Our findings of the presence of kallikrein in goblet cells of the gastrointestinal tract raises the possibility that, along with mucus, they may have a cytoprotective role. Kinins produced by colonic kallikrein may also contribute to pathological conditions such as diarrhea associated with carcinoid tumours (Donowitz and Binder, 1975).

The observation of greater changes in whole small intestine blood flow after kallikrein administration when compared with saline injection suggests that tissue kallikrein may regulate intestinal blood flow (Overlack *et al.*, 1983). If, as Zeitlin (1970) suggested, the kinins which are dilators and inhibit peristalsis, are important in intestinal absorption, they may be major factors in controlling normal vasomotor tone, permeability and motility of the intestine. It is interesting that 5-hydroxytryptamine, a vasoconstrictor substance that stimulates peristalsis, is also found along the length of the gastrointestinal tract.

#### **D. Interpretation of Immunocytochemical Findings**

Cytological procedures are used in attempts to discern relationships between structure and function. Immunocytochemical localizations are the most widely used methods and their aim is to identify and characterize proteins in tissue by labelling the antibody-antigen complex, making it visible.

The introduction of fluorescein-labelled immune sera (Coons and Kaplan, 1950) was a breakthrough in investigations of specific cellular components by light microscopy. Analogous techniques are also useful for ultrastructural studies where, in principle, the electron microscope can resolve an antibody molecule which has reacted with its antigen. Singer and Schick (1961), using ferritin-labelled antibody, successfully localized antigen at the electron

microscope level, but found the disadvantage of impaired reagent penetration.

With the advent of enzyme-labelled antibodies and the development of the unlabelled antibody method of Mason *et al.* (1969) and the PAP method of Sternberger and his colleagues (Sternberger, 1979) came increased sensitivity in detecting small amounts of antigen. This method was used in this thesis, and it is widely used in localization of intracellular antigens. A further advancement in antigen detection—also used in this thesis, was made by Hsu *et al.* (1981) with the avidin-biotin complex method. Superior results are generally obtained and are attributed to the formation of a large complex containing multiple peroxidase molecules. The ABC method has several advantages over the PAP method, including lower background stain, easily recognized reaction product, shorter incubation times, and higher dilutions of reagents which can be used.

The pre-embedding immunostaining method, used occasionally in this thesis and found to be unsuccessful, has several disadvantages over the post-embedding method. First, immunoglobulins are unable to adequately penetrate into well-fixed thick sections. Secondly, non-specific globulin binding and randomly oriented reaction product make interpreting positive results difficult. Thirdly, reagents often do not permeate membranes easily. The major advantage of post-embedding immunostaining techniques, apparent in these results, is that immunoperoxidase reactions take place on the surface of routinely prepared ultrathin sections where intracellular antigens are accessible.

A fundamental conclusion which can be drawn from these localization experiments is that for each tissue examined and for a given antigen, specifically tailored methods of preparation must be developed. Although a search for the ideal immunocytochemical fixative produced several good results, no one treatment was universally applicable.

Discrepancies in localizations have occurred for other antigens where different techniques produced different results. In the colon, ethanol fixation produces carcinoembryonic antigen localization in epithelial cells (Rognum *et al.*, 1980; Primus *et al.*, 1981; Lindgren *et al.*, 1982; Kaku *et al.*, 1983). Formaldehyde-fixed, pronase-treated sections also produce similar

results, but without pronase treatment the sections do not produce the expected stain (Rognum *et al.*, 1980). Other studies which used different methods localized the same or very similar antigens in mucous cells (Rogalsky, 1975; Huitric *et al.*, 1976). Localizations of NGF in specific subcellular compartments of duct cells of submandibular glands have also yielded contradictory results, but in this case the discrepancies were related to inexperience with new methods. By immunofluorescence, NGF was first associated with cytoplasm, secretory granules, basal parts of cells and nuclei (Levi-Montalcini and Angeletti, 1961; Goldstein and Burdman, 1965; Kumar, 1972). A different pattern was later observed by Schwab *et al.* (1976), who localized the antigen only in the apical secretory granules. Since basal granules and other organelles did not stain, it was postulated that the NGF antigen is physiologically converted from a proprotein to an immunoreactive NGF, probably under the influence of a protease. Using methods very similar to those used in this thesis, Hofmann and Drenehahn (1981) demonstrated NGF antigenicity uniformly in all secretory granules. Possibly in some studies cell damage caused protease activity to expose antigenicity during tissue fixation as a post-mortem artefact.

Localization of tissue kallikrein to acinar cells of the exocrine pancreas or to  $\beta$  cells of the endocrine pancreas in different studies appears to depend both on the specificity of  $\alpha$  kallikrein antibodies used and the manner in which pancreatic tissue is prepared. Acinar localization of kallikrein in the pancreas was achieved by Ørstavik *et al.* (1981a), who used ethanol as the primary fixative, but ole Moi-Yoi *et al.* (1979a), who used formaldehyde, found antigenicity in endocrine islet cells. Lack of staining in acinar cells may be ascribed to antigenic masking since protease digestion, which was not used in their studies, produces immunostaining in acini (Pinkus *et al.*, 1983). Unmasking of endocrine kallikrein by trypsin or pronase suggests that it is less accessible to antibody than exocrine kallikrein.

### E. Possible Sources of Circulating Tissue Kallikrein

Tissue kallikrein, in an active or inactive form, has been detected in plasma (Geiger *et al.*, 1980; Rabito *et al.*, 1980, 1982; Lawton *et al.*, 1981; Shimamoto *et al.*, 1984). Although its exact role in plasma is unknown, Masferrer *et al.* (1985) suggested that it could influence blood circulation via its ability to generate kinins. In plasma, the enzyme is present but bound to several inhibitors of different molecular weights (Johansen *et al.*, 1984).

Sources of the immunoreactive enzyme in plasma have not been determined, though several organs may contribute kallikrein to the circulation. Pancreatic kallikrein, which is absorbed across the intestinal wall (Moriwaki *et al.*, 1974), may contribute to circulating enzyme, but not significantly, since plasma from animals that had undergone subtotal pancreatectomy did not differ in antigen levels from control plasma (Lawton *et al.*, 1981). The submandibular gland was suggested to be a major contributor to circulating tissue kallikrein when labelled enzyme injected into the gland's main duct was detected in venous blood (Ørstavik *et al.*, 1980b). Support for this idea also came from Lawton *et al.* (1981), who found a great reduction in circulating antigen levels after bilateral submandibular and sublingual gland excision. The presence of residual antigen after removal of these glands suggests that other sources of enzyme such as parotid glands, pancreas, intestine or kidney exist.

The kidney may deliver tissue kallikrein into the circulation since enzyme antigenicity is associated with basolateral infoldings of tubule cells (Figuroa *et al.*, 1984). It is released into renal lymph and it is found in venous effluent of isolated perfused kidneys (Roblero *et al.*, 1976). The kidney may also play a role in clearance or metabolism of tissue kallikrein since an increased enzyme activity in plasma of nephrectomized rats was observed (Masferrer *et al.*, 1985). Renal kallikrein may also arise from circulating tissue kallikrein (Fiedler and Gebhard, 1980).

## V. CONCLUSION

In this thesis, several kallikrein-like enzymes sharing immunological properties were localized in gastrointestinal, respiratory and male reproductive tissues. Nerve growth factor was also localized in male sex glands. Techniques were developed that were highly successful, particularly those using semi-thin resin embedded tissue sections and the avidin-biotin complex method of immunostaining.

In the various tissues examined here, antigen localization only occurred in apical regions of secretory cells and in secretions, indicating that kallikrein is an exported protein, likely playing a role in extracellular secretions. This finding is largely, though not exclusively, supported by other studies. Exceptions include cases of the kidney (Vio *et al.*, 1983), the pancreas (ole Moi-Yoi, 1979) and salivary glands (Kimura and Moriya, 1984a), where antigens were also localized in areas suggestive of endocrine secretion. Indeed, the presence of tissue kallikrein in the circulation indicates that it must be derived from at least one of the tissues known to synthesize it, and likely the mechanism by which the antigen enters the circulation is by endocrine-like secretion. While this thesis does not supply evidence for any endocrine secretion of kallikrein or nerve growth factor in tissues studied, it does not claim that this type of secretion does not exist.

Kallikrein enzyme activity has been implicated by many studies in a variety of essential physiological functions. Gene specific probes have recently isolated 25-30 members of the kallikrein family having 75% amino acid sequence homology (Mason *et al.*, 1983). Since sequences differ only at the active site region, it is apparent that functional differences may exist for all of the enzyme gene family members. Because three of these enzymes, tissue kallikrein, EGF-binding protein and  $\gamma$ -NGF are associated with bioactive peptide processing, it seems likely that other members will be discovered in similar roles.

## VI. REFERENCES

- Abelous, J.-E. and Bardier, E., Les substances hypotensives de l'urine humaine normale. *C.R.S. Soc. Biol.* 66: 511-512, 1909.
- Albano, J., Bhoola, K.D., Heap, P.F. and Lemon, M.J.C. Stimulus-secretion coupling: role of cyclic AMP, cyclic GMP and calcium in mediating enzyme (kallikrein) secretion in the submandibular gland. *J. Physiol.* 258: 631-658, 1976.
- Aloe, L., Cozzari, C. and Levi-Montalcini, R., The submaxillary salivary glands of the African rodent *Praomys (mastomys) natalensis* is the richest available source of the nerve growth factor. *Exp. Cell Res.* 133: 475-480, 1981.
- Amouric, M. and Figarella, C., Characterization and purification of a kallikrein from human pancreatic juice and immunological comparison with other kallikreins. *Hoppe-Seyler's Z. Physiol. Chem.* 360: 457-465, 1979.
- Amundsen, E. and Nustad, K., Kinin-forming and destroying activities of cell homogenates. *J. Physiol.* 179: 479-488, 1965.
- Angeletti, P., Calissano, P., Chen, J.S. and Levi-Montalcini, R., Multiple molecular forms of the nerve growth factor. *Biochim. Biophys. Acta* 147: 180-182, 1967.
- Arnold, W.H., A comparative study on the localization of kallikrein and kallikrein-like activity in different organs. in *Kininogenases, Kinins VI*, pp. 59-73. Ed. Haberland, G.L., Rohen, J.W., Fritz, H. and Huber, P. Stuttgart, Schattauer Verlag, 1983.
- Arnold, W.H., Comparative studies on the localization of esteroproteases and kallikrein-like activity in primate organs. *Histochem. J.* 16: 755-769, 1984.
- Arnold, W.H., Ultrahistochemical localization of kallikrein-like activity in the submandibular gland of *Macaca fasciculatis* monkey. *Histochem. J.*, submitted for publication, 1985.
- Baggio, B., Favaro, S., Antonello, A., Zen, A., Zen, F. and Borsatto, A., Subcellular localization of renin and kallikrein in rat kidney. *Ital. J. Biochem.* 24: 199-206, 1975.
- Barka, T., Biologically active polypeptides in submandibular glands. *J. Histochem. Cytochem.* 28: 836-859, 1980.
- Barton, S., Wimalasena, J. and Schachter, M., Subcellular location of the kininogenase in the coagulating gland of the guinea-pig. *Biochem. Pharmacol.* 22: 1121-1123, 1973.

- Barton, S., Sanders, E.J., Schachter, M. and Uddin, M., Autonomic nerve stimulation, kallikrein content and acinar cell granules of the cat's submandibular gland. *J. Physiol.* **251**: 363-369, 1975.
- Barton, S., Karpinski, E., Moriwaki, C. and Schachter, M., Sialotonin: vasopressor substance in saliva and submandibular gland of the cat. *J. Physiol.* **261**: 523-533, 1976.
- Beilenson, S., Schachter, M. and Smaje, L.H., Secretion of kallikrein and its role in vasodilatation in the submaxillary gland. *J. Physiol.* **199**: 303-317, 1968.
- Bendayan, M. and Ørstavik, T.B., Immunocytochemical localization of kallikrein in the rat exocrine pancreas. *J. Histochem. Cytochem.* **30**: 58-66, 1982.
- Bennick, A., Chemical and physical characteristics of a phosphoprotein from human parotid saliva. *Biochem. J.* **145**: 557-567, 1975.
- Beraldo, W.T., Feldberg, W. and Hilton, S.M., Experiments on the factor in urine forming substance U. *J. Physiol.* **133**: 558-565, 1956.
- Berger, E.A. and Shooter, E.M., Evidence for pro- $\beta$ -nerve growth factor, a biosynthetic precursor to  $\beta$ -nerve growth factor. *Proc. Natl. Acad. Sci. U.S.A.* **74**: 3647-3651, 1977.
- Bhoola, K.D., Comparative study of the subcellular distribution of submaxillary kallikrein. *Biochem. Pharmacol.* **18**: 1252-1254, 1969.
- Bhoola, K.D. and Dorey, G., Isolation of kallikrein-containing granules from the pancreas and submaxillary gland of the cat. *J. Physiol.* **203**: 59-60P, 1969.
- Bhoola, K.D. and Dorey, G., Intracellular localization of kallikrein, trypsin and amylase in dog pancreas. *J. Physiol.* **214**: 553-570, 1971.
- Bhoola, K.D. and Heap, P.F., Properties of kallikrein-containing granules isolated from the submaxillary gland of the guinea-pig. *J. Physiol.* **210**: 421-432, 1970.
- Bhoola, K.D. and Ogle, C.W., The subcellular localization of kallikrein, amylase and acetylcholine in the submaxillary gland of the guinea-pig. *J. Physiol.* **184**: 663-672, 1966.
- Bhoola, K.D., May May Yi, R., Morley, J. and Schachter, M., Release of kinin by an enzyme in the accessory sex glands of the guinea-pig. *J. Physiol.* **163**: 269-280, 1962.



- Bhoola, K.D., Dorey, G. and Jones, C.W., The influence of androgens on enzymes (chymotrypsin- and trypsin-like proteases, renin, kallikrein and amylase) and on cellular structure of the mouse submaxillary gland. *J. Physiol.* **235**: 503-522, 1973.
- Bhoola, K.D., Matthews, R.W. and Roberts, F., A survey of salivary kallikrein and amylase in a population of schoolgirls throughout the menstrual cycle. *Clin. Sci. Molec. Med.* **55**: 561-565, 1978.
- Bing, J., Poulsen, K., Hackenthal, E., Rix, E. and Taugner, R., Renin in the submaxillary gland: a review. *J. Histochem. Cytochem.* **29**: 874-880, 1980.
- Bloom, G.D., Carlsson, Danielsson, A., Marklund, S. and Stigbrand, T., Peroxidase and amylase activity of the guinea pig submandibular gland during the secretory cycle. *Histochemistry* **38**: 271-280, 1974.
- Boesman, M., Levy, M. and Schenkein, I., Esteroproteolytic enzymes from the submaxillary gland. Kinetics and other physicochemical properties. *Arch. Biochem. Biophys.* **175**: 463-476, 1976.
- Bothwell, M.A., Wilson, W.H. and Shooter, E.M., The relationship between glandular kallikrein and growth factor-processing proteases of mouse submaxillary gland. *J. Biol. Chem.* **254**: 7287-7294, 1979.
- Brandtzaeg, P., Gautvik, K.M., Nustad, K. and Pierce, J.V., Rat submandibular gland kallikreins: purification and cellular localization. *Brit. J. Pharmac.* **56**: 155-167, 1976.
- Britos, J. and Nolly, H., Kinin-forming enzyme of rat cardiac tissue. Subcellular distribution and biochemical properties. *Proc. Inter. Am. Soc.* **3**: II-42-II-45, 1981.
- Bueker, E.D., Implantation of tumors in the hind limb field of the embryonic chick and the developmental response of the lumbosacral nervous system. *Anat. Rec.* **102**: 369-390, 1948.
- Burger, A., Lembeck, F. and Wagner, J., Serotonin und Kininogenase-Aktivität in Fraktionen der Dünndarmschleimhaut des Kaninchens. *Naunyn-Schmiedebergs Arch. Pharmak. u. exp. Path.* **261**: 412-421, 1968.
- Calissano, P., Cattaneo, A., Biocca, S., Aloe, L., Mercanti, D. and Levi-Montalcini, R., The nerve growth factor. Established findings and controversial aspects. *Exp. Cell Res.* **154**: 1-9, 1984.

- Campenot, R.B., Local control of neurite development by nerve growth factor. *Proc. Nat. Acad. Sci. U.S.A.* **74**: 4516-4519, 1977.
- Chao, J., Sostek, M., Shimamoto, K., Bank, H.L., Bigelow, J. and Margolius, H.S., Kallikrein content of rat pancreatic acinar cells or islets by direct radioimmunoassay. *Hoppe-Seyler's Z. Physiol. Chem.* **361**: 1805-1810, 1980.
- Chao, J., Buse, J., Shimamoto, K. and Margolius, H.S., Kallikrein-induced uterine contraction independent of kinin formation. *Proc. Natl. Acad. Sci. U.S.A.* **78**: 6154-6157, 1981.
- Chao, J., Ross, C.G., Shimamoto, K., Miller, D.H. and Margolius, H.S., Regulation of kallikrein and arginine esterase A by testosterone in the rat submaxillary gland and kidney. in *Recent Progress in Kinins*. Ed. Fritz, H., Dietze, G., Fiedler, F. and Haberland, G.L. *Agents and Actions, Suppl.* **9**, pp. 332-330, 1982.
- Chao, J., Chao, L. and Margolius, H.S., Identification of a kallikrein-like latent serine protease in human erythrocyte membranes. *Biochem. Biophys. Res. Comm.* **121**: 722-729, 1984a.
- Chao, J., Chao, L. and Margolius, H.S., Isolation of tissue kallikrein in rat spleen by monoclonal antibody-affinity chromatography. *Biochim. Biophys. Acta* **801**: 244-249, 1984b.
- Chapdelaine, P., Dubé, J.Y., Frenette, G. and Tremblay, R.R., Identification of arginine esterase as the major androgen-dependent protein secreted by dog prostate and preliminary molecular characterization in seminal plasma. *J. Androl.* **5**: 206-210, 1984.
- Chapman, C.A., Banks, B.E.C., Vernon, C.A. and Walker, J.M., The isolation and characterisation of nerve growth factor from the prostate gland of the guinea-pig. *Eur. J. Biochem.* **115**: 347-351, 1981.
- Cohen, S., Purification and metabolic effects of a nerve growth-promoting protein from snake venom. *J. Biol. Chem.* **234**: 1129-1137, 1959.
- Colman, R.W. and Bagdasarian, A., Human kallikrein and prekallikrein. in *Methods in Enzymology. XLV. Proteolytic Enzymes*, ed. L. Lorand, 1976.
- Colman, R.W. and Wong, P.Y., Kallikrein-kinin system in pathologic conditions. *Handb. Exp. Pharm.* **25**: 569-607, 1979.

- Coons, A.H. and Kaplan, M.H., Localization of antigen in tissue cells. II. Improvements in a method for the detection of antigen by means of fluorescent antibody. *J. Exp. Med.* **91**: 1-14, 1950.
- Coulter, H.D. and Terracio, L., Preparation of biological tissues for electron microscopy by freeze-drying. *Anat. Rec.* **187**: 477-494, 1977.
- Coulter, H.D. and Terracio, L., An all glass freeze-drier for TEM specimens with an improved design for temperature regulation, fixation, and infiltration. in *Ninth International Congress on Electron Microscopy*, Toronto, Vol. IV, p. 60-61, 1978.
- Crocker, A.D. and Willavoys, P., Effect of bradykinin on transepithelial transfer of sodium and water *in vitro*. *J. Physiol.* **253**: 401-410, 1975.
- Cuthbert, A.W. and Margolius, H.S., Kinins stimulate net chloride secretion by the rat colon. *Brit. J. Pharmacol.* **75**: 587-598, 1982.
- Cuthbert, A.W., George, A.M. and MacVinish, L., Kinin effects on electrogenic ion transport in primary cultures of pig renal papillary collecting tubule cells. *Am. J. Physiol.*, in press, 1985.
- Darling, T.L.J., Petrides, P.E., Beguin, P., Frey, P., Shooter, E.M., Selby, M. and Rutter, W.J., The biosynthesis and processing of proteins in the mouse 7S nerve growth factor complex. *Cold Spring Harbor Symp. Quant. Biol.* **48**: 427-434, 1983.
- DeArmond, S.J., Siegel, M.W., Dixon, R.G. and Eng, L.F., Post-embedding immunoperoxidase staining of glial fibrillary acidic protein for light and electron microscopy. *J. Neuroimmunol.* **1**: 3-15, 1981.
- de Carvalho, I.F. and Diniz, C.R., Kinin-forming enzyme (kininogenin) in homogenates of rat kidney. *Biochim. Biophys. Acta* **128**: 136-148, 1966.
- Dembinski, A.B. and Johnson, L.R., Effect of epidermal growth factor on the development of rat gastric mucosa. *Endocrinology* **116**: 90-94, 1985.
- Dietl, T., Kruck, J. and Fritz, H., Localization of kallikrein in porcine pancreas and submandibular gland as revealed by the indirect immunofluorescence technique. *Hoppe-Seyler's Z. Physiol. Chem.* **359**: 499-505, 1978.
- Donowitz, M. and Binder, H.J., Jejunal fluid and electrolyte secretion in carcinoid syndrome. *Digestive Diseases* **20**: 1115-1122, 1975.

- Dubé, J.Y., Chapdelaine, P. and Tremblay, R.R., Effect of testicular hormones on synthesis of soluble proteins by dog prostate slices. *Can. J. Biochem. Cell Biol.* 61: 756-763, 1983.
- Ekfors, T.O. and Hopsu-Havu, V.K., Immunofluorescent localization of trypsin-like estero-peptidases in the mouse submandibular gland. *Histochem. J.* 3: 415-421, 1971.
- Emmelin, N. and Henriksson, K.G., Depressor activity of saliva after section of the chorda tympani. *Acta Physiol. Scand.* 30: 75-82, 1953.
- Fasth, S., Hultén, L., Johnson, B.J., Nordgren, S. and Zeitlin, I.J., Mobilization of colonic kallikrein following pelvic nerve stimulation in the atropinized cat. *J. Physiol.* 285: 471-478, 1978.
- Fava-de-Moraes, F., Zangheri, E.O. and Doine, A.I., Immunohistochemical localization of erythropoietin in the rat and mouse submandibular gland. *Histochem. J.* 11: 97-102, 1979.
- Feldberg, W. and Guimaraes, J.A., Effects of sympathetic impulses and adrenaline on salivary secretion. *J. Physiol.* 83: 43P-45P, 1935.
- Fiedler, F., Enzymology of glandular kallikreins: in *Bradykinin, Kallidin and Kallikrein*, pp. 103-154. Ed. Erdős, E.G. Springer-Verlag, Berlin, Heidelberg, New York, 1979.
- Fiedler, F. and Fritz, H., Striking structural similarity of pig pancreatic kallikrein, mouse nerve growth factor  $\gamma$ -subunit and rat tonin. *Hoppe-Seyler's Z. Physiol. Chem.* 362: 1171-1175, 1981.
- Fiedler, F. and Gebhard, W., Isolation and characterization of native single-chain porcine pancreatic kallikrein, another possible precursor of urinary kallikrein. *Hoppe-Seyler's Z. Physiol. Chem.* 361: 1661-1671, 1980.
- Fiedler, F., Ehret, W., Dogee, G., Hirschauer, C., Kutzbach, C., Schmidt-Kastner, G. and Tschesche, H., The primary structure of pig pancreatic kallikrein B. in *Kininogenases 4*, pp. 7-14. Ed. Haberland, G.L., Rohen, J.W. and Suzuki, T. Schattauer, New York, Stuttgart, 1977.
- Fiedler, F., Bhoola, K.D., Hirschauer, C., Leysath, G., Matthews, R.W., Lemon, M.J.C. and Whicher, J.T., Purification and properties of guinea-pig submandibular kallikrein. in *Recent Progress in Kinins*, pp. 157-161. Ed. Fritz, H., Dietze, G., Fiedler, F. and Haberland, G.L., 1982.

- Figueroa, C.D., Caorsi, I., Subiabre, J. and Vio, C.P., Immunoreactive kallikrein localization in the rat kidney: an immuno-electron-microscopic study. *J. Histochem. Cytochem.* 32: 117-121, 1984.
- Fink, E., Dietl, T., Seitert, J. and Fritz, H., Studies on the biological function of glandular kallikrein. in *Advances in Experimental Medicine and Biology*, Vol. 120B, pp. 261-273. Ed. Fujii, S., Moriya, H. and Suzuki, T. Plenum Press, New York, London, 1979.
- Fink, E., Schill, W-B. and Fiedler, F., Tissue kallikrein in human seminal fluid. in *Protides of the Biological Fluids*, Vol. 32, pp. 299-303. Ed. H. Peeters, Pergamon Press, 1985.
- Forell, M.M., Pancreatic kallikrein in physiological and pathological conditions. in *Polypeptides Which Affect Smooth Muscles*, pp. 247-252. Ed. M. Schachter, 1960.
- Frankish, N.H. and Zeitlin, I.J., The effect of diet on tissue levels of kinin-forming enzyme in blood-free rat gastro-intestinal tract. *J. Physiol.* 298: 361-370, 1980.
- Freund, J. and Thompson, G.E., Toxic effects of fluid from the coagulating gland of the guinea pig. *Proc. Soc. Exp. Biol. Med.* 94: 350-351, 1957.
- Freund, J., Miles, A.A., Mill, P.J. and Wilhelm, D.L., Vascular permeability factors in the secretion of the guinea pig coagulating gland. *Nature* 182: 174-175, 1958.
- Frey, E.K. and Kraut, H., Über einen von der Niere ausgeschiedenen, die Herztätigkeit anregenden Stoff. *Hoppe-Seyler's Z. Physiol. Chem.* 157: 32-61, 1926.
- Frey, E.K., Kraut, H. and Werle, E., *Das Kallikrein-Kinin-System und seine Inhibitoren*. Enke, Stuttgart, 1968.
- Fujimori, H., Levison, P.R. and Schachter, M., Purification and partial characterisation of cat pancreatic and submandibular gland kallikreins. in *Kinins IV: Advances in Experimental Medicine and Biology*, in press, 1985a.
- Fujimori, H., Levison, P.R. and Schachter, M., Purification and partial characterisation of cat pancreatic and urinary kallikreins. in *Kinins IV: Advances in Experimental Medicine and Biology*, in press, 1985b.
- Fujimoto, Y., Moriwaki, C. and Moriya, H., Studies on human salivary kallikrein. II. Properties of purified salivary kallikrein. *J. Biochem. (Tokyo)* 74: 239-247, 1973.

- Fukuoka, Y., Hojima, Y., Miyaura, S. and Moriwaki, C., Purification of cat submaxillary kallikrein. *J. Biochem. (Tokyo)* 85: 549-557, 1979.
- Fuller, P.J., Clements, J.A., Whitfeld, P.L. and Funder, J.W., Kallikrein gene expression in the rat anterior pituitary. *Mol. Cell. Endocrin.* 39: 99-105, 1985.
- Garrett, J.R. and Kidd, A., Effects of nerve stimulation and denervation on secretory material in submandibular striated duct cells of cats, and the possible role of these cells in the secretions of salivary kallikrein. *Cell Tiss. Res.* 161: 71-84, 1975.
- Garrett, J.R., Smith, R.E., Kidd, A., Kyriacou, K. and Grabske, R.J., Kallikrein-activity in salivary glands using a new tripeptide substrate, including preliminary secretory studies and observations on mast cells. *Histochem. J.* 14: 967-979, 1982.
- Garrett, J.R., Kidd, A., Kyriacou, K. and Smith, R.E., New observations on the localization of kallikrein-like activity in human salivary parenchymal and mast cells by enzyme histochemistry. *Histochem. J.* 16: 789-791, 1984.
- Geiger, R. and Clausnitzer, B., Isolation of an enzymatically active tissue kallikrein from human seminal plasma by immunoaffinity chromatography. *Hoppe-Seyler's Z. Physiol. Chem.* 362: 1279-1283, 1981.
- Geiger, R., Feifel, G. and Haberland, G.L., A precursor of kinins in the gastric mucus. *Hoppe-Seyler's Z. Physiol. Chem.* 358: 931-933, 1977.
- Geiger, R., Clausnitzer, B., Fink, E. and Fritz, H., Isolation of an enzymatically active glandular kallikrein from human plasma by immunoaffinity chromatography. *Hoppe-Seyler's Z. Physiol. Chem.* 362: 1795-1803, 1980.
- Geller, R.G., Margolius, H.S., Pisano, J.J. and Keiser, H.R., Effects of mineralocorticoids, altered sodium intake, and adrenalectomy on urinary kallikrein in rats. *Circ. Res.* 31: 857-863, 1972.
- Gibbs, O.S., On the alleged occurrence of acetylcholine in the saliva. *J. Physiol.* 84: 33-40, 1935.
- Goldstein, L.A., Reynolds, C.P. and Perez-Polo, J.R., Isolation of human nerve growth factor from placental tissue. *Neurochem. Res.* 3: 175-183, 1983.
- Goldstein, M.N. and Burdman, J.A., Studies of the nerve growth factor in submandibular glands of female mice treated with testosterone. *Anat. Rec.* 151: 199-208, 1965.

- Gospodarowicz, D., Epidermal and nerve growth factors in mammalian development. *Ann. Rev. Physiol.* 43: 251-263, 1981.
- Gotterer, G., Banks, J. and Williams-Ashman, H.G., Hydrolysis of arginine esters by male accessory sexual tissues. *Proc. Soc. Exp. Biol. Med.* 82: 58-61, 1956.
- Greene, L.A., Shooter, E.M. and Varon, S., Enzymatic activities of mouse nerve growth factor and its subunits. *Proc. Nat. Acad. Sci. U.S.A.* 60: 1383-1388, 1969.
- Gresik, E. and Barka, T., Immunocytochemical localization of epidermal growth factor in mouse submandibular gland. *J. Histochem. Cytochem.* 25: 1027-1035, 1977.
- Hare, E.R. and Verpoorte, J.A., The purification and partial characterization of human salivary kallikrein. *Biochim. Biophys. Acta* 709: 65-72, 1982.
- Harper, G.P. and Thoenen, H., The distribution of nerve growth factor in the male sex organs of mammals. *J. Neurochem.* 34: 893-903, 1980.
- Harper, G.P. and Thoenen, H., Biochemical, biological and immunological characterization of bovine nerve growth factor. in *Chemistry of Peptides and Proteins*, Vol. 2, pp. 171-177. Walter de Gruyter & Co., Berlin, 1984.
- Harper, G.P., Barde, Y.A., Burnstock, G., Carstairs, J.R., Dennison, M.E., Suda, K. and Vernon, C.A., Guinea pig prostate is a rich source of nerve growth factor. *Nature* 279: 160-162, 1979.
- Hassell, J. and Hand, A.R., Tissue fixation with diimidoesters as an alternative to aldehydes. I. Comparison of cross-linking and ultrastructure obtained with dimethylsuberimide and glutaraldehyde. *J. Histochem. Cytochem.* 22: 223-239, 1974.
- Havez, R., Degand, P., Roussel, P., Voisin, C., Biserte, G. and Gernez-Rieux, C., Isolement et caractérisation immunologique de la kallicréine bronchique humaine. *C.R. Acad. Sci., Paris* 262: 309-311, 1966a.
- Havez, R., Degand, P., Roussel, P., Voisin, C., Biserte, G. and Gernez-Rieux, C., Identification des  $\gamma$ A-globulines, de la kallicréine et de la transferrine dans la muqueuse bronchique humaine. *C.R. Acad. Sci., Paris* 262: 1777-1779, 1966b.
- Hayat, M.A., *Principles and Techniques of Electron Microscopy. Biological Applications*, Vol. 1. Van Nostrand Reinhold Co., New York, 1970.

- Heidrich, H.G. and Geiger, R., Kininogenase activity in plasma membranes and cell organelles from rabbit kidney cortex: subcellular localization of renal kallikrein by free-flow electrophoresis and density-gradient fractionation. *Kidney Int.* **18**: 77-85, 1980.
- Hendry, I.A. and Iversen, L.L., Reduction in the concentration of nerve growth factor in mice after sialectomy and castration. *Nature (London)* **243**: 500-504, 1973.
- Hilton, S.M. and Lewis, G.P., The relationship between glandular activity, bradykinin formation and functional vasodilatation in the submandibular salivary gland. *J. Physiol.* **134**: 471-483, 1956.
- Hiramatsu, M., Hatakeyama, K., Kumegawa, M., Yajima, T. and Minami, N., Genetic variation in esteroproteases in the mouse submandibular gland. *Experientia* **37**: 1068-1069, 1981.
- Hirata, Y. and Orth, D.N., Concentrations of epidermal growth factor, nerve growth factor, and submandibular gland renin in male and female mouse tissue and fluids. *Endocrinology* **105**: 1382, 1979.
- Hofmann, H-D. and Drenckhahn, D., Distribution of nerve growth factor in the submandibular gland of the male and female mouse. *Cell Tiss. Res.* **221**: 77-83, 1981.
- Hojima, Y., Maranda, B., Moriwaki, C. and Schachter, M., Direct evidence for the location of kallikrein in the striated ducts of the cat's submandibular gland by the use of specific antibody. *J. Physiol.* **268**: 793-801, 1977a.
- Hojima, Y., Yamashita, M., Ochi, N., Moriwaki, C. and Moriya, H., Isolation and some properties of dog and rat pancreatic kallikreins. *J. Biochem. (Tokyo)* **81**: 599-610, 1977b.
- Hsu, S-M., Raine, L. and Fanger, H., Use of avidin-biotin-peroxidase complex (ABC) in immunoperoxidase techniques: a comparison between ABC and unlabelled antibody (PAP) procedures. *J. Histochem. Cytochem.* **29**: 577-580, 1981.
- Hsueh, W.A., Carlson, E.J., O'Connor, D. and Warren, S., Renin requires a structural alteration prior to activation by renal kallikrein. *J. Clin. Endocrin. Metab.* **51**: 942-944, 1980.
- Huitric, E., Laumonier, R., Burton, D., von Kleist, S. and Charanel, G., An optical and ultrastructural study of the localization of carcinoembryonic antigen (CEA) in normal and cancerous human rectocolonic mucosa. *Lab. Invest.* **34**: 97-107, 1976.



- Humason, G.L., *Animal Tissue Techniques*, 4th Edition. W.H. Freeman & Co., San Francisco, 1979.
- Isaacs, W.B. and Shaper, J.H., Isolation and characterization of the major androgen-dependent glycoprotein of canine prostatic fluid. *J. Biol. Chem.* **258**: 6610-6615, 1983.
- Ishii, D.N. and Shooter, E.M., Regulation of nerve growth factor synthesis in mouse submaxillary glands by testosterone. *J. Neurochem.* **25**: 843-851, 1975.
- Izumi, H., Release of kallikrein-like esterase and tonin from dispersed cells of the rat submandibular gland. *Brit. J. Pharmac.* **82**: 175-182, 1984.
- Izumi, H. and Aoki, T., Release of benzoyl-arginine-ethylester esterase from dispersed cells of mouse submandibular gland. *Archs. Oral Biol.* **26**: 845-851, 1981.
- Johansen, L., Nustad, K., Berg, T. and Pierce, J.V., Excess antibody immunoassay for rat glandular kallikrein. Measurement of kallikrein complexed with inhibitors and in plasma. *J. Immunol. Methods* **69**: 253-266, 1984.
- Kaizu, T. and Margolius, H.S., Studies on rat renal cortical cell kallikrein: Separation and measurement. *Biochim. Biophys. Acta* **411**: 305-315, 1975.
- Kaku, T., Ekem, J.K., Lindayen, C., Bailey, D.J., van Nostrand, A.W.P. and Farber, E., Comparison of formalin- and acetone-fixation for immunohistochemical detection of carcinoembryonic antigen (CEA) and keratin. *Am. J. Clin. Path.* **80**: 806-815, 1983.
- Kauker, M.L., Bradykinin action on the efflux of luminal  $^{22}\text{Na}$  in the rat nephron. *J. Pharmacol. Exp. Therap.* **214**: 119-123, 1980.
- Kimura, K. and Moriya, H., Enzyme- and immuno-histochemical localization of kallikrein. I. The human parotid gland. *Histochemistry* **80**: 367-372, 1984a.
- Kimura, K. and Moriya, H., Enzyme- and immuno-histochemical localization of kallikrein. II. The human kidney. *Histochemistry* **80**: 443-448, 1984b.
- Kimura, K., Takagi, M., Igari, T., Ishii, M., Ikeda, T., Takeda, T. and Murao, S., Histochemical localization of kallikrein-like Pro-Phe-Arg-Naphthylester esterase activity in the rat kidney. *Histochemistry* **75**: 91-98, 1982.

- Kirkland, S.C., Cuthbert, A.W. and MacVinish, L.J., Kinin effects on ion transport in monolayers of HCA7 cells, a line from a human colonic carcinoma. Submitted for publication, 1985.
- Kobayashi, M. and Ohata, K., Property of kinin-forming enzyme in rat stomach. *Jap. J. Pharmacol.* 31: 369-373, 1981.
- Kobayashi, M., Shikimi, T., Miyata, S. and Ohata, K., Studies on kinin-forming enzyme in rat stomach. *Jap. J. Pharmacol.* 29: 947-950, 1979.
- Kraut, H., Frey, E.K. and Werle, E., Der Nachweis eines Kreislaufhormons in der Pankreasdrüse. IV. Mitteilung über dieses Kreislaufhormon. *Hoppe-Seyler's Z. Physiol. Chem.* 189: 97-106, 1930.
- Kuhlmann, W.D. and Krischan, R., Resin embedment of organs and postembedment localization of antigens by immunoperoxidase methods. *Histochemistry* 72: 377-389, 1981.
- Kumar, S., Stewart, J.K., Taylor, G. and Waghe, M.A., Fluorescence studies using anti-nerve growth factor. *Exptl. Cell Res.* 74: 170-174, 1972.
- Lane, B.P. and Europa, D.L., Differential staining of ultrathin sections of epon-embedded tissues for light microscopy. *J. Histochem. Cytochem.* 13: 579-582, 1965.
- Larson, P.S., On the alleged occurrence of acetylcholine and adrenalin in cat's saliva. *J. Pharmacol. Exp. Therap.* 54: 341-345, 1935.
- Lawton, W.J., Proud, D., Frech, M.E., Pierce, J.V., Keiser, H.R. and Pisano, J.J., Characterization and origin of immunoreactive glandular kallikrein in rat plasma. *Biochem. Pharmacol.* 30: 1731-1737, 1981.
- Lazure, C., Leduc, R., Seidah, N.G., Chrétien, M., Dubé, J.Y., Chapdelaine, P., Frenette, G., Paquin, R. and Tremblay, R.R., The major androgen-dependent protease in dog prostate belongs to the kallikrein family: confirmation by partial amino acid sequencing. *FEBS* 175: 1-7, 1984.
- Lechene de la Porte, M., Amouric, M., and Figarella, C., Immunolocalization of kallikrein in porcine pancreas at the ultrastructural level. *Hoppe-Seyler's Z. Physiol. Chem.* 362: 439-446, 1981.
- Leduc, E.H. and Bernhard, W., Recent modifications of the glycol methacrylate embedding procedure. *J. Ultrastruct. Res.* 19: 196-199, 1967.

- Levi-Montalcini, R. and Angeletti, P.U., Growth control of the sympathetic system by a specific protein-factor. *Quart. Rev. Biol.* 36: 99-108, 1961.
- Levi-Montalcini, R. and Angeletti, P.U., Hormonal control of the nerve growth factor content in the submaxillary glands of mice. in *Salivary Glands and Their Secretion*, pp. 129-141. New York, Pergamon Press, 1964.
- Levi-Montalcini, R. and Booker, B., Excessive growth of the sympathetic ganglia evoked by a protein isolated from mouse salivary glands. *Proc. Nat. Acad. Sci. U.S.A.* 46: 373-384, 1960.
- Levi-Montalcini, R. and Hamburger, V., Selective growth stimulating effects of mouse sarcoma on the sensory and sympathetic nervous system of the chick embryo. *J. Exp. Zool.* 116: 321-362, 1951.
- Lewis, P.R. and Knight, D.P., *Staining Methods for Sectioned Material: Practical Methods in Electron Microscopy*, Ed. A.M. Glauert. North Holland Publ. Co., Amsterdam, New York, Oxford, 1980.
- Li, A.K.C., Koroly, M.J., Schattenkerk, M.E., Malt, R.A. and Young, M., Nerve growth factor: acceleration of the rate of wound healing in mice. *Proc. Natl. Acad. Sci. U.S.A.* 77: 4379-4381, 1980.
- Lindgren, J., Wahlström, T., Bång, B., Hurme, M. and Mäkelä, O., Immunoperoxidase staining of carcinoembryonic antigen with monoclonal antibodies in adenocarcinoma of the colon. *Histochemistry* 74: 223-227, 1982.
- Mann, K., Göring, W., Lipp, W., Keipert, B., Karl, H.J., Geiger, R. and Fink, E., Radio-immunoassay of human urinary kallikrein. Determination of human urinary kallikrein, II. *J. Clin. Chem. Clin. Biochem.* 18: 395-401, 1980.
- Manning, D.C. and Snyder, S.H., Bradykinin receptor-mediated chloride secretion in intestinal function. *Nature* 299: 256-259, 1982.
- Maranda, B., Rodrigues J.A.A., Schachter, M., Shnitka, T.K. and Weinberg, J., Studies on kallikrein in the duct systems of the salivary glands of the cat. *J. Physiol.* 276: 321-328, 1978.
- Marchetti, J., Imbert-Teboul, M., Alhenc-Gelas, F., Allegrini, J., Menard, J. and Morel, F., Kallikrein along the rabbit microdissected nephron: a micromethod for its measurement. *Pflüg. Arch.* 401: 27-33, 1984.

- Margolius, H.S. and Chao, J., Amiloride inhibits mammalian renal kallikrein and a kallikrein-like enzyme from toad bladder and skin. *J. Clin. Invest.* 65: 1343-1350, 1980.
- Margolius, H.S., Geller, R., de Jong, W., Pisano, J.J. and Sjoerdsma, A., Altered urinary kallikrein excretion in rats with hypertension. *Circ. Res.* 30: 358, 1972.
- Margolius, H.S., Horwitz, D., Geller, R.G., Alexander, R.W., Gill, J.R. Jr., Pisano, J.J. and Keiser, H.R., Urinary kallikrein excretion in normal man. Relationships to sodium intake and sodium-retaining steroids. *Circ. Res.* 35: 812-817, 1974.
- Masferrer, J., Albertini, R., Croxatto, H.R., García, P. and Pinto, I., Isolation and characterization of rat plasma glandular kallikrein. *Biochem. Pharmacol.* 34: 51-56, 1985.
- Mason, A.J., Evans, B.A., Cox, D.R., Shine, J. and Richards, R.I., Structure of mouse kallikrein gene family suggests a role in specific processing of biologically active peptides. *Nature* 303: 300-307, 1983.
- Mason, T.E., Phifer, R.F., Spicer, S.S., Swallow, R.A. and Dreskin, R.B., An immunoglobulin-enzyme bridge method for localizing tissue antigens. *J. Histochem. Cytochem.* 17: 563-569, 1969.
- Mattioli, W. and Mattioli, A., Presence of callicrein in the salivary glands of rabbits and the possible correlation between its concentration and the functional activity of the glands. *Chem. Abst.* 41: 508, 1947a.
- Mattioli, W. and Mattioli, A., Callicrein content of salivary glands. Callicrein; supposed internal secretion of the salivary glands. *Chem. Abst.* 41: 801, 1947b.
- McLean, I.W. and Nakane, P.K., Periodate-lysine-paraformaldehyde fixative. A new fixative for immunoelectron microscopy. *J. Histochem. Cytochem.* 22: 1077-1083, 1974.
- Millonig, G., Advantages of a phosphate buffer for OsO<sub>4</sub> solutions in fixation. *J. Appl. Physics* 32: 1637, 1961.
- Mobley, W.C., Server, A.C., Ishii, D.N., Riopelle, R.J. and Shooter, E.M., Nerve growth factors. Part I. *New Engl. J. Med.* 297: 1096-1104, 1977a.
- Mobley, W.C., Server, A.C., Ishii, D.N., Riopelle, R.J. and Shooter, E.M., Nerve growth factors. Part II. *New Engl. J. Med.* 297: 1149-1158, 1977b.

- Mobley, W.C., Server, A.C., Ishii, D.N., Riopelle, R.J. and Shooter, E.M., Nerve growth factors. Part III. *New Engl. J. Med.* 297: 1211-1218, 1977c.
- Moore, B., Jr., Mobley, W.C. and Shooter, E.M., Proteolytic modification of the  $\beta$  nerve growth factor protein. *Biochemistry* 13: 833-840, 1974.
- Moriwaki, C. and Fujimori, H., Studies on kallikreins. VI. Effect of pancreatic kallikrein and kinin on the transport of amino acids and glucose from the rat small intestine. *Chem. Pharm. Bull.* 29: 804-809, 1981.
- Moriwaki, C. and Schachter, M., Kininogenase of the guinea-pig's coagulating gland and the release of bradykinin. *J. Physiol.* 219: 341-353, 1971.
- Moriwaki, C., Watanuki, N., Fujimoto, Y. and Moriya, H., Further purification and properties of kininogenases from the guinea pig's coagulating gland. *Chem. Pharm. Bull.* 22: 628-633, 1974.
- Moriwaki, C., Kiyazaki, K., Matsuda, Y., Moriya, H., Fujimoto, Y. and Ueki, H., Dog renal kallikrein: purification and some properties. *J. Biochem. (Tokyo)* 80: 1277-1285, 1976.
- Moriwaki, C., Fujimori, H., Moriya, H. and Kizuki, K., Studies on kallikreins. IV. Enhancement of valine transport across the rat small intestine. *Chem. Pharm. Bull.* 25: 1174-1178, 1977.
- Moriwaki, C., Fujimori, H., Toyono, Y. and Nagai, T., Studies on kallikreins. V. Purification and characterization of rat intestinal kallikrein. *Chem. Pharm. Bull.* 28: 3612-3620, 1980.
- Moriya, H., Pierce, J.V. and Webster, M.E., Purification and some properties of three kallikreins. *Ann. N.Y. Acad. Sci.* 104: 172-185, 1963.
- Morris, B.J., Catanzaro, D.F. and De Zwart, R.T., Evidence that the arginine esteropeptidase ( $\gamma$ ) subunit of nerve growth factor can activate inactive renin. *Neurosci. Lett.* 24: 87-92, 1981.
- Murphy, R.A., Watson, A.Y., Metz, J. and Forssmann, W.G., The mouse submandibular gland: an exocrine organ for growth factors. *J. Histochem. Cytochem.* 28: 890-902, 1980.

- Nexo, E., Olsen, P.S. and Poulsen, K., Exocrine and endocrine secretion of renin and epidermal growth factor from the mouse submandibular glands. *Reg. Pep.* 8: 327-334, 1984.
- Nieuw Amerongen, A.V., Aarsman, M.E.G., Oderkerk, C.H., Vreugdenhil, A.P. and Roukema, P.A., AM<sub>2</sub>, a submandibular secretory glycoprotein of the mouse. An immunochemical and immunohistochemical study. *J. Biol. Buccale* 9: 41-52, 1981.
- Nolly, H. and Lama, M.C., 'Vascular kallikrein': a kallikrein-like enzyme present in vascular tissue of the rat. *Clin. Sci.* 63: 249s-251s, 1982.
- Nolly, H., Bertini, F. and Lama, M.C., Kinin-forming enzymes in vascular tissue. in *Kinins III, Advances in Experimental Medicine and Biology*, Vol. 156A, pp. 399-407. Ed. H. Fritz, N. Back, G. Dietze and G.L. Haberland. Plenum Press, New York, 1983.
- Nustad, K., The relationship between kidney and urinary kininogenase. *Brit. J. Pharmacol.* 39: 73-86, 1970a.
- Nustad, K., Localization of kininogenase in the rat kidney. *Brit. J. Pharmacol.* 39: 87-98, 1970b.
- Nustad, K. and Pierce, J.V., Purification of rat urinary kallikreins and their specific antibody. *Biochemistry* 13: 2312, 1974.
- Nustad, K. and Rubin, I., Subcellular localization of renin and kininogenase in the rat kidney. *Brit. J. Pharmacol.* 40: 326-333, 1970.
- Nustad, K. and Vaaje, K., Synthesis of kallikreins by rat kidney slices. *Brit. J. Pharmacol.* 53: 229-234, 1975.
- Oka, Y., Ghishan, F.K., Greene, H.L. and Orth, D.N., Effect of mouse epidermal growth factor/urogastrone on the functional maturation of rat intestine. *Endocrinology* 112: 940-944, 1983.
- ole-Moi Yoi, O., Austen, K.F. and Spragg, J., Kinin-generating and esterolytic activity of purified human urinary kallikrein (urokallikrein). *Biochem. Pharmacol.* 26: 1893-1900, 1977.
- ole-Moi Yoi, O., Spragg, J. and Austen, K.F., Inhibition of urinary kallikrein (urokallikrein) by anti-enzyme FAB. *J. Immunol.* 121: 66-71, 1978.

- ole-Moy Yoi O., Pinkus, G.S., Spragg, J. and Austen, K.F., Identification of human glandular kallikrein in the beta cell of the pancreas. *New Engl. J. Med.* 300: 1289-1294, 1979a.
- ole-Moi Yoi, O., Spragg, J. and Austen, K.F., Structural studies of human urinary kallikrein (urokallikrein). *Proc. Natl. Acad. Sci. U.S.A.* 76: 3121-3125, 1979b.
- Omata, K., Carretero, O.A., Scicli, G. and Jackson, B.A., Localization of active and inactive kallikrein (kininogenase activity) in the microdissected-rabbit nephron. *Kidney Int.* 22: 602-607, 1982.
- Orce, G.G., Castillo, G.A. and Margolius, H.S., Inhibition of short-circuit current in toad urinary bladder by inhibitors of glandular kallikrein. *Am. J. Physiol.* 239: F459-F465, 1980.
- Ørstavik, T.B., An immunohistochemical study of kallikrein in the rat parotid and exorbital lacrimal glands. *Archs. Oral Biol.* 23: 1023-1025, 1978a.
- Ørstavik, T.B., The distribution and secretion of kallikrein in some exocrine organs of the rat. *Acta Physiol. Scand.* 104: 431-442, 1978b.
- Ørstavik, T.B. and Gautvik, K.M., Regulation of salivary kallikrein secretion in the rat submandibular gland. *Acta Physiol. Scand.* 100: 33-44, 1977.
- Ørstavik, T.B. and Glenner, G.G., Localization of kallikrein and its relation to other trypsin-like esterases in the rat pancreas. A comparison with the submandibular gland. *Acta Physiol. Scand.* 103: 384-393, 1978.
- Ørstavik, T.B. and Inagami, T., Localization of kallikrein in the rat kidney and its anatomical relationship to renin. *J. Histochem. Cytochem.* 30: 385-390, 1982.
- Ørstavik, T.B., Brandtzaeg, P., Nustad, K. and Halvorsen, K.M., Cellular localization of kallikreins in rat submandibular and sublingual salivary glands. *Acta Histochem.* 54: 183-192, 1975.
- Ørstavik, T.B., Nustad, K., Brandtzaeg, P. and Pierce, J.V., Cellular origin of urinary kallikreins. *J. Histochem. Cytochem.* 24: 1037-1039, 1976.
- Ørstavik, T.B., Brandtzaeg, K., Nustad, K. and Pierce, J.V., Immunohistochemical localization of kallikrein in human pancreas and salivary glands. *J. Histochem. Cytochem.* 28: 557-562, 1980a.

- Ørstavik, T.B., Gautvik, K.M. and Nustad, K., Intraglandular transport of  $^{125}\text{I}$ -glandular kallikrein in the rat submandibular salivary gland. *Acta Physiol. Scand.* 109: 315-323, 1980b.
- Ørstavik, T.B., Brandtzaeg, P., Nustad, K. and Pierce, J.V., Effects of different tissue processing methods on the immunohistochemical localization of kallikrein in the pancreas. *J. Histochem. Cytochem.* 29: 985-988, 1981a.
- Ørstavik, T.B., Brekke, I.B., Alumets, J. and Carretero, O.A., Kallikrein in rat pancreatic tissue after beta cell destruction or acinar cell atrophy. *J. Histochem. Cytochem.* 29: 1431-1436, 1981b.
- Ørstavik, T.B., Carretero, O.A., Hayashi, H., Scicli, G.A. and Johansen, L., Immunohistochemical localization of tonin and its relation to kallikrein in rat salivary glands. *J. Histochem. Cytochem.* 30: 1123-1129, 1982a.
- Ørstavik, T.B., Carretero, O.A. and Scicli, A.G., Kallikrein-kinin system in regulation of submandibular gland blood flow. *Am. J. Physiol.* 242: H1010-H1014, 1982b.
- Ouchterlony, O., Diffusion-in-gel methods for immunological analysis. in *Progress in Allergy*, Vol. 5, pp. 1-78. ed. P. Kallós. Karger, Basel, 1958.
- Overlack, A., Scicli, A.G. and Carretero, O.A., Intestinal absorption of glandular kallikrein in the rat. *Am. J. Physiol.* 245: G689-G-694, 1983.
- Pattison, S.E. and Dunn, M.F., On the relationship of zinc ion to the structure and function of the 7S nerve growth factor protein. *Biochemistry* 14: 2733-2739, 1975.
- Pease, D.C., *Histological Techniques for Electron Microscopy*. Academic Press, New York, 1964.
- Pelliniemi, L.J. and Van Noorden, S., A streamlined technique for on-grid electron immunocytochemistry. *Histochemistry* 53: 271-272, 1977.
- Pignatelli, F., Maurice, M. and Feldmann, G., Immunoperoxidase localization of albumin and fibrinogen in rat liver fixed by perfusion or immersion: effect of saponin on the intracellular penetration of labelled antibodies. *J. Histochem. Cytochem.* 30: 1004-1014, 1982.



- Pinkus, G.S., ole-Moi Yoi, O., Austen, K.F. and Spragg, J., Antigenic separation of a nonkinin-generating TAME esterase from human urinary kallikrein and immunohistochemical comparison of their localization in the kidney. *J. Histochem. Cytochem.* 29: 38-44, 1981.
- Pinkus, G.S., Maier, M., Seldin, D.C., ole-Moi Yoi, O., Austen, K.F. and Spragg, J., Immunohistochemical localization of glandular kallikrein in the endocrine and exocrine human pancreas. *J. Histochem. Cytochem.* 31: 1279-1288, 1983.
- Polivka, E., Maier, M. and Binder, B.R., Purification and characterization of a kallikrein-like kininogenase from pig pituitary glands. *Agents and Actions Suppl.* 9: 153-156, 1982.
- Porcelli, G., Bianchi, G. and Croxatto, H.R., Urinary kallikrein excretion in a spontaneously hypertensive strain of rats (38939). *Proc. Soc. Exp. Biol. Med.* 149: 983-986, 1976.
- Powers, C.A. and Nasjletti, A., A kininogenase resembling glandular kallikrein in the rat pituitary pars intermedia. *Endocrinology* 112: 1194-1200, 1983.
- Powers, C.A. and Nasjletti, A., A major sex difference in kallikrein-like activity in the rat anterior pituitary. *Endocrinology* 114: 1841-1844, 1984.
- Powers, C.A., Baer, P.G. and Nasjletti, A., Reduced glandular kallikrein-like activity in the anterior pituitary of the New Zealand genetically hypertensive rat. *Biochem. Biophys. Res. Comm.* 119: 689-693, 1984.
- Prado, E.S., Prado de Carvalho, L., Araujo-Viel, M.S., Ling, N. and Rossier, J., A met-enkephalin-containing-peptide, BAM 22P, as a novel substrate for glandular kallikreins. *Biochem. Biophys. Res. Comm.* 112: 366-371, 1983.
- Primus, F.J., Clark, C.A. and Goldenberg, D.M., Immunoperoxidase localization of carcino-embryonic antigen in normal human intestinal mucosa. *J. Nat. Canc. Inst.* 67: 1031-1039, 1981.
- Proud, D., Bailey, G.S., Ørstavik, T.B. and Nustad, K., The location of prekallikrein in the rat pancreas. *Biochem. Soc. Trans.* 5: 1402-1404, 1977.
- Proud, D., Knepper, M.A. and Pisano, J.J., Distribution of immunoreactive kallikrein along the rat nephron. *Am. J. Physiol.* 13: F510-F515, 1983.

- Rabito, S.F., Scicli, A.G. and Carretero, O., Immunoreactive glandular kallikrein in plasma. in *Enzymatic Release of Vasoactive Peptides*, pp. 247-258. Ed. Gross, F. and Vogel, G., Raven Press, New York, 1980.
- Rabito, S.F., Scicli, A.G., Kher, V. and Carretero, O.A., Immunoreactive glandular kallikrein in rat plasma: a radioimmunoassay for its determination. *Am. J. Physiol.* **242**: H602-H610, 1982.
- Reissig, M., Bang, B.G. and Bång, F.B., Ultrastructure of the mucociliary interface in the nasal mucosa of the chicken. *Am. Rev. Resp. Disease* **117**: 327-341, 1978.
- Reynolds, E.S., The use of lead citrate at high pH as an electron-opaque stain in electron microscopy. *J. Cell. Biol.* **17**: 208-212, 1963.
- Richardson, K.C., Jarrett, L. and Finke, E.H., Embedding in epoxy resins for ultrathin sectioning in electron microscopy. *Stain Technol.* **35**: 313-323, 1960.
- Roblero, J., Croxatto, H., García, R., Corthorn, J. and De Vito, E., Kallikrein-like activity in perfusates and urine of isolated rat kidneys. *Am. J. Physiol.* **231**: 1383-1389, 1976.
- Rogalsky, V.Y., Variations in carcinoembryonic antigen localization in tumours of the colon. *J. Nat. Canc. Inst.* **54**: 1061-1071, 1975.
- Rognum, T.O., Brandtzaeg, P., Ørjasaeter, H. and Fausa, O., Immunohistochemistry of epithelial cell markers in normal and pathological colon mucosa. *Histochemistry* **67**: 7-21, 1980.
- Roth, J., Bendayan, M. and Orci, L., Ultrastructural localization of intracellular antigens by the use of protein A-gold complex. *J. Histochem. Cytochem.* **26**: 1074-1081, 1978.
- Rush, R.A., Immunohistochemical localization of endogenous nerve growth factor. *Nature* **312**: 364-367, 1984.
- Sallay, C. and Nador, C., Changes in the kallikrein contents of saliva under physiological circumstances and in periodontoclasia. *J. Dent. Res.* **29**: 232, 1950.
- Schachter, M., Kallikreins and kinins. *Physiol. Rev.* **49**: 509-547, 1969.
- Schachter, M., Kallikreins (kininogenases) - a group of serine proteases with bioregulatory actions. *Pharmacol. Rev.* **31**: 1-17, 1980.

- Schachter, M., Barton, S., Uddin, M., Karpinski, E. and Sanders, E.J., Effect of nerve stimulation, denervation, and duct ligation, on kallikrein content and duct cell granules of the cat's submandibular gland. *Experientia* 33: 746-748, 1977.
- Schachter, M., Maranda, B. and Moriwaki, C., Localization of kallikrein in the coagulating and submandibular glands of the guinea pig. *J. Histochem. Cytochem.* 26: 318-321, 1978.
- Schachter, M., Peret, M.W., Moriwaki, C. and Rodrigues, J.A.A., Localization of kallikrein in submandibular gland of cat, guinea pig, dog, and man by the immunoperoxidase method. *J. Histochem. Cytochem.* 28: 1295-1300, 1980.
- Schachter, M., Peret, M.W., Billing, A.G. and Wheeler, G.D., Immunolocalization of the protease kallikrein in the colon. *J. Histochem. Cytochem.* 31: 1255-1260, 1983a.
- Schachter, M., Peret, M.W., Moriwaki, C., Wheeler, G.D., Matthews, R.W., Labeledz, T. and Kaneko, S., Localization of kininogenases in salivary glands, sex glands and spermatozoa. in *Kinins III, Advances in Experimental Medicine and Biology*, Vol. 156A, pp. 377-386, Ed. H. Fritz, N. Back, G. Dietze and G.L. Haberland. Plenum Press, New York, 1983b.
- Schachter, M., Wheeler, G.D., Matthews, R.W., Peret, M.W. and Moriwaki, C., Ultrastructural immunolocalization of kallikrein in apical granules of striated duct cells of cat submandibular gland. *J. Histochem. Cytochem.* 31: 345-347, 1983c.
- Schenkein, I., Boesman, M., Tokarsky, E., Fishman, L. and Levy, M., Proteases from mouse submaxillary gland. *Biochem. Biophys. Res. Comm.* 36: 156-165, 1969.
- Schill, W.B. and Haberland, G.L., Kinin inducement of sperm motility. *Hoppe-Seyler's Z. Physiol. Chem.* 355: 229, 1974.
- Schill, W.B., Braun-Falco, O. and Haberland, G.L., The possible role of kinins in sperm motility. *Int. J. Fert.* 19: 163-167, 1974.
- Schmid, G. and Heidland, A., Cation excretion and kallikrein secretion by the rat parotid gland in various hypertensive models. *J. Cardiovasc. Pharmacol.* 6: S101-S106, 1984.
- Schwab, M.E., Stöckel, K. and Thoenen, H., Immunocytochemical localization of nerve growth factor (NGF) in the submandibular gland of adult mice by light and electron microscopy. *Cell Tiss. Res.* 169: 289-299, 1976.

- Scioli, A.G., Carretero, O.A., Hampton, A., Cortes, P. and Oza, N.B., Site of kininogenase secretion in the dog nephron. *Am. J. Physiol.* 230: 533-536, 1976a.
- Scioli, A.G., Carretero, O.A., Oza, N.B. and Schork, A., Distribution of kidney kininogenases. *Proc. Soc. Exp. Biol. Med.* 151: 57-60, 1976b.
- Scioli, A.G., Forbes, G., Nolly, H., Dujovny, M., and Carretero, O.A., Kallikrein-kinins in the central nervous system. *Clin. Exp. Theory Prac.* A6: 1731-1738, 1984.
- Sealey, J.E., Atlas, S.A. and Laragh, J.H., Linking the kallikrein and renin systems via activation of inactive renin. *Am. J. Med.* 65: 994-1000, 1978.
- Secker, J., The humoral control of the secretion by the submaxillary gland of the cat following chorda stimulation. *J. Physiol.* 81: 81-82, 1934a.
- Secker, J., The humoral control of the secretion of the submaxillary gland of the cat following sympathetic stimulation. *J. Physiol.* 82: 293-304, 1934b.
- Seki, T., Nakajima, T. and Erdős, E.G., Colon kallikrein, its relation to the plasma enzyme. *Biochem. Pharmacol.* 218: 1227-1235, 1972.
- Server, A.C. and Shooter, E.M., Nerve growth factor. *Adv. Protein Chem.* 31: 339-405, 1976.
- Shikata, H., Utsumi, N., Hiramatsu, M., Minami, N., Nemoto, N. and Shikata, T., Immunohistochemical localization of nerve growth factor and epidermal growth factor in guinea pig prostate gland. *Histochemistry* 80: 411-413, 1984.
- Shikimi, T., Kema, R., Matsumoto, M., Yamahata, Y. and Miyata, S., Studies on kinin-like substances in brain. *Biochem. Pharmacol.* 22: 567-573, 1972.
- Shimamoto, K., Mayfield, R.K., Margolius, H.S., Chao, J., Stroud, W. and Kaplan, A.P., Immunoreactive tissue kallikrein in human serum. *J. Lab. Clin. Med.* 103: 731-738, 1984.
- Shinohara, H., Yanagimachi, R. and Srivastava, P.N., Enhancement of the acrosome reaction of hamster spermatozoa by the proteolytic enzymes, kallikrein, trypsin, and chymotrypsin. *Gamete Res.* 11: 19-28, 1985.

- Simson, J.A.V., Hazen, D., Spicer, S.S., Murphy, R.A. and Young, M., Secretagogue-mediated discharge of nerve growth factor from granular tubules of male mouse submandibular glands: an immunocytochemical study. *Anat. Rec.* 192: 375-388, 1978.
- Simson, J.A.V., Spicer, S.S., Chao, J., Grimm, L. and Margolius, H.S., Kallikrein localization in rodent salivary glands and kidney with the immunoglobulin-enzyme bridge technique. *J. Histochem. Cytochem.* 27: 1567-1576, 1979.
- Simson, J.A.V., Fenters, R. and Chao, J., Electron microscopic immunostaining of kallikrein in rat submandibular glands. *J. Histochem. Cytochem.* 31: 301-306, 1983.
- Simson, J.A.V., Chao, J., Dom, R., Chao, L. and Margolius, H.S., Kallikrein quantitation and localization in rat brain with radioimmunoassay and immunohistochemistry. in *Kinin 84, International Congress*, Savannah, Georgia, 1984 (abstract).
- Singer, S.J. and Schick, A.P., The properties of specific stains for electron microscopy prepared by the conjugation of antibody molecules with ferritin. *J. Biophys. Biochem. Cytol.* 9: 519-537, 1961.
- Spurr, A.R., A low-viscosity epoxy resin embedding medium for electron microscopy. *J. Ultrastruct. Res.* 26: 31-43, 1969.
- Sternberger, L.A., *Immunocytochemistry*, 2nd edition. John Wiley & Sons, New York, Chichester, Brisbane, Toronto, 1979.
- Streefkerk, J.G., Inhibition of erythrocyte pseudoperoxidase activity by treatment with hydrogen peroxidase following methanol. *J. Histochem. Cytochem.* 20: 829-931, 1972.
- Tanaka, T., Sakano, A. and Takahashi, A., Immunohistochemical localization of epidermal growth factor in the intercalated duct secretory granules of the submandibular glands of adult female A/J mice. *Acta Histochem. Cytochem.* 17: 565-570, 1984.
- Thibault, G. and Genest, J., Tonin, an esteroprotease from rat submaxillary glands. *Biochim. Biophys. Acta* 660: 23-29, 1981.
- Thoenen, H. and Barde, Y-A., Physiology of nerve growth factor. *Physiol. Rev.* 60: 1284-1335, 1980.
- Thomas, K.A. and Bradshaw, R.A., [46] $\gamma$ -subunit of mouse submaxillary gland 7 S nerve growth factor: an endopeptidase of the serine family. *Methods Enzymol.* 80: 609-620, 1981.

- Tornita, K., Endou, H. and Sakai, F., Localization of kallikrein-like activity along a single nephron in rabbits. *Pflug. Arch.* 389: 91-95, 1981.
- Turkington, R.W., Males, J.L. and Cohen, S., Synthesis and storage of epithelial-epidermal growth factor in submaxillary gland. *Cancer Res.* 31: 252-256, 1971.
- Uchida, K., Niinobe, M., Kato, H. and Fujii, S., Purification and properties of rat stomach kallikrein. *Biochim. Biophys. Acta* 614: 501-510, 1980.
- Uchida, K., Kushiro, H., Kodama, J., Hitomi, Y., Niinobe, M. and Fujii, S., Rat thyroid kallikrein: its purification and properties. *Agents and Actions Suppl.* 9: 167-172, 1982.
- Uddin, M. and Tyler, D.W., Demilune cells of the cat submandibular gland; an unlikely source of kallikrein. *Experientia* 34: 609-610, 1978.
- Uddin, M. and Tyler, D.W., The localization of kallikrein in the dog and guinea-pig submandibular glands. *Experientia* 37: 872-878, 1981.
- Uetsuji, S., Yamamura, M., Yamamoto, M., Uchida, K., Kushiro, H., Kodama, J. and Fujii, S., Human stomach kallikrein. *Agents and Actions Suppl.* 9: 137-142, 1982.
- Ueyama, T., Saito, H. and Yohro, T., *Suncus murinus* submandibular gland and prostate are new sources of nerve growth factor. *Biomed. Res.* 2: 438-441, 1981.
- Van Leeuwen, B.H., Grinblat, S.M. and Johnston, C.I., Tissue-specific control of glandular kallikreins. *Am. J. Physiol.* 246: F760-F764, 1984.
- Varon, S., Nomura, J. and Shooter, E.M., Reversible dissociation of mouse nerve growth factor protein into different subunits. *Biochemistry* 7: 1296-1303, 1968.
- Vio, C.P., Roblero, J.S. and Croxatto, H.R., Dexamethasone, aldosterone and kallikrein release by isolated rat kidney. *Clin. Sci.* 61: 241-243, 1981.
- Vio, C.P., Churchill, L., Terragno, A., McGiff, J.C. and Terragno, N.A., Arachidonic acid stimulates renal kallikrein release in isolated rat kidney. *Clin. Sci.* 63: 235s-237s, 1982.
- Vio, C.P., Churchill, L., Rabito, S.F., Terragno, A., Carretero, O.A. and Terragno, N.A., Renal kallikrein in venous effluent of filtering and non-filtering isolated kidneys. *Adv. Exp. Med. Biol.* 156: 897-905, 1983.

- Ward, P.E., Gedney, C.D., Dowben, R.M. and Erdös, E.G., Isolation of membrane-bound renal kallikrein and kininase. *Biochem. J.* **151**: 755-758, 1975.
- Ward, P.E., Erdös, E.G., Gedney, C.D., Dowben, R.M. and Reynolds, R.C., Isolation of membrane-bound renal enzymes that metabolize kinins and angiotensins. *Biochem. J.* **157**: 643-650, 1976a.
- Ward, P.E., Erdös, E.G., Gedney, C.D., Dowben, R.M. and Reynolds, R.C., Isolation of renal membranes that contain kallikrein, angiotensin I-converting enzyme (kininase II) and angiotensinase in the rat. *Clin. Sci. Mol. Med.* **51**: 267s-270s, 1976b.
- Welsh, M.J., Smith, P.L., Fromm, M. and Frizzell, R.A., Crypts are the site of intestinal fluid and electrolyte secretion. *Science* **218**: 1219-1221, 1982.
- Werle, E., Kallikrein, kallidin and related substances. in *Polypeptides Which Affect Smooth Muscles and Blood Vessels*, pp.199. Ed. Schachter, M., Pergamon Press, London, 1960.
- Werle, E. and Semm, K., Methode zue Schwangerschaftsdiagnose mit Bestimmung des Alters der Frucht. *Arch. Gynäk.* **187**: 106-111, 1955.
- Werle, E. and Trautschold, I., Kallikrein, kallidin, kallikrein inhibitors. *Ann. N.Y. Acad. Sci.* **104**: 117, 1963.
- Werle, E. and Vogel, R., Über die kallikreinausscheidung im harn nach Experimenteller Nierenschädigung. *Arch. Int. Pharmacodyn.* **126**: 1-2, 1960.
- Werle, E. and Vogel, R., Über die Freisetzung einer Kallikreinartigen Substanz aus Extrakten Verscheidener Organe. *Arch. Int. Pharmacodyn.* **131**: 257-261, 1961.
- Werle, E. and von Roden, P., Über das Vorkommen von Kallikrein in den Speicheldrüsen und im Mundspeichel. *Biochemische Z.* **286**: 213-219, 1936.
- Werle, E., Vogel, R. and Tentrodt, J., Über den Kallikreingehalt der Gl. submandibularis während der Entwicklung einiger Säugeracten und nach Abbindung des D. Whartoni. *Arch. Int. Pharmacodyn.* **126**: 187, 1960.
- Willingham, M.C. and Yamada, S.S., Development of a new primary fixative for electron microscopic immunocytochemical localization of intracellular antigens in cultured cells. *J. Histochem. Cytochem.* **27**: 947-960, 1979.

Wong, Y.C. and Tse, M.K.W., Fine structural and functional study of the prostatic complex of the guinea pig. *Acta Anat.* 109: 289-312, 1981.

Zacharatos, D.T., Vasak, E.Z., Cheng, E.S.P., van Lennep, E.W. and Morris, B.J., Immunohistochemistry of tonin in rat submandibular gland during development, lactation and secretion. *Histochemistry* 79: 433-442, 1983.

Zeitlin, I.J., Kinin release associated with the gastro-intestinal tract. *Adv. Exp. Med. Biol.* 8: 329-339, 1970.

Zeitlin, I.J. Pharmacological characterization of kinin-forming activity in rat intestinal tissue. *Brit. J. Pharmacol.* 42: 648-649P, 1971.

Zeitlin, I.J. and Smith, A.N., Mobilization of tissue kallikrein in inflammatory disease of the colon. *Gut* 14: 133-138, 1973.

Zeitlin, I.J., Singh, Y.N., Lembeck, F. and Theiler, M., The molecular weights of plasma and intestinal kallikreins in rats. *Naunyn-Schmiedeberg's Arch. Pharmacol.* 293: 159-161, 1976.

Zimmerman, A., Geiger, R. and Kortmann, H., Similarity between a kininogenase (kallikrein) from human large intestine and human urinary kallikrein. *Hoppe-Seyler's Z. Physiol. Chem.* 360: 1767-1773, 1979.