

30746



National Library  
of Canada

Bibliothèque nationale  
du Canada

CANADIAN THESES  
ON MICROFICHE

THÈSES CANADIENNES  
SUR MICROFICHE

NAME OF AUTHOR/NOM DE L'AUTEUR

ANIL R. LONDHE

TITLE OF THESIS/TITRE DE LA THÈSE

THE LIMITING DISTRIBUTION  
OF THE MEASURE OF A  
RANDOM SET

UNIVERSITY/UNIVERSITÉ

UNIVERSITY OF ALBERTA

DEGREE FOR WHICH THESIS WAS PRESENTED/  
GRADE POUR LEQUEL CETTE THÈSE FUT PRÉSENTÉE

M.Sc.

YEAR THIS DEGREE CONFERRED/ANNÉE D'OBTENTION DE CE DEGRÉ

1976

NAME OF SUPERVISOR/NOM DU DIRECTEUR DE THÈSE

Dr. J. R. McGregor

Permission is hereby granted to the NATIONAL LIBRARY OF  
CANADA to microfilm this thesis and to lend or sell copies  
of the film.

The author reserves other publication rights, and neither the  
thesis nor extensive extracts from it may be printed or otherwise  
reproduced without the author's written permission.

L'autorisation est, par la présente, accordée à la BIBLIOTHÈQUE  
NATIONALE DU CANADA de microfilmer cette thèse et  
de prêter ou de vendre des exemplaires du film.

L'auteur se réserve les autres droits de publication; ni la  
thèse ni de longs extraits de celle-ci ne doivent être imprimés  
ou autrement reproduits sans l'autorisation écrite de l'auteur.

DATED/DATE

Sept. 22, 1976

SIGNED/SIGNÉ

PERMANENT ADDRESS/RÉSIDENCE FIXE

Qrt. No. 12, Type IV, Seminary Hills,  
Nagpur - 440006, INDIA

## INFORMATION TO USERS

THIS DISSERTATION HAS BEEN  
MICROFILMED EXACTLY AS RECEIVED

This copy was produced from a microfiche copy of the original document. The quality of the copy is heavily dependent upon the quality of the original thesis submitted for microfilming. Every effort has been made to ensure the highest quality of reproduction possible.

PLEASE NOTE: Some pages may have  
indistinct print. Filmed as  
received.

Canadian Theses Division  
Cataloguing Branch  
National Library of Canada  
Ottawa, Canada K1A 0N4

## AVIS AUX USAGERS

LA THESE A ETE MICROFILMEE  
TELLE QUE NOUS L'AVONS RECUE

Cette copie a été faite à partir d'une microfiche du document original. La qualité de la copie dépend grandement de la qualité de la thèse soumise pour le microfilmage. Nous avons tout fait pour assurer une qualité supérieure de reproduction.

NOTA BENE: La qualité d'impression de certaines pages peut laisser à désirer. Microfilmee telle que nous l'avons reçue.

Division des thèses canadiennes  
Direction du catalogage  
Bibliothèque nationale du Canada  
Ottawa, Canada K1A 0N4

THE UNIVERSITY OF ALBERTA

THE LIMITING DISTRIBUTION OF THE  
MEASURE OF A RANDOM SET

by

(C)

ANIL R. LONDHE

A THESIS

SUBMITTED TO THE FACULTY OF GRADUATE STUDIES AND RESEARCH  
IN PARTIAL FULFILMENT OF THE REQUIREMENTS FOR THE DEGREE  
OF MASTER OF SCIENCE

IN

STATISTICS

DEPARTMENT OF MATHEMATICS


EDMONTON, ALBERTA


FALL, 1976


THE UNIVERSITY OF ALBERTA


FACULTY OF GRADUATE STUDIES AND RESEARCH

The undersigned certify that they have read, and  
recommend to the Faculty of Graduate Studies and Research, for  
acceptance, a thesis entitled THE LIMITING DISTRIBUTION OF  
THE MEASURE OF A RANDOM SET submitted by ANIL R. LONDHE in  
partial fulfilment of the requirements for the degree of  
Master of Science in Mathematical Statistics.

  
(Supervisor)







Date Sept. 22, 1976

## ABSTRACT

W.L. Stevens initiated research in the general area of random sets by considering arcs of length  $D$  ( $0 < D < 1$ ) at random points on the circumference of a circle of length unity and derived the probability that the circumference is completely covered. D.F. Votaw Jr. considered a somewhat similar but non-circular problem and derived the probability function of the measure of a random linear set. D.D. Beck, using the methods of Votaw, derived the probability function of the measure of a random circular set. The results of these authors are reviewed in the first chapter.

The second chapter is devoted to finding the limiting distribution and the non-central moments of the measure of a random circular set for the case  $D = 1/(n+1)$  where  $n$  is the number of arcs. The last section of this chapter contains discussions of plots and tables of the probability function of the measure of a random circular set for various values of  $n$  and  $D$ .

#### **ACKNOWLEDGEMENTS**

I would like to express my sincere gratitude to Dr. J.R. McGregor for suggesting this topic and for his thoughtful and meticulous guidance throughout the preparation of this thesis. Thanks are also due to June Talpash for typing the thesis.

TABLE OF CONTENTS

CHAPTER	PAGE
I    INTRODUCTION . . . . .	1
II    MAIN RESULTS . . . . .	7
2.1 Preliminary . . . . .	7
2.2 Limiting Distribution . . . . .	9
2.3 Moments of Variable $\{1-\mu(X)\}$ . . . . .	27
2.4 Plots and Tables . . . . .	30
BIBLIOGRAPHY . . . . .	32
APPENDIX I . . . . .	33
APPENDIX II . . . . .	39
APPENDIX III . . . . .	59

## CHAPTER I

### INTRODUCTION



Consider  $n+1$  points  $z_i$  ( $i = 0, 1, \dots, n$ ) selected independently and at random from the interval  $(0, 1)$ ; the distribution of any  $z_i$  being the uniform distribution with distribution function

$$(1.1) \quad F(z) = z, \quad 0 \leq z \leq 1.$$

Order the  $n+1$  points in ascending order of magnitude as  $x_i$  ( $i = 0, 1, \dots, n$ ). The totality of all cases in which two of the  $x_i$ 's are equal has zero probability and such cases can be excluded without affecting the problem. Associate with each  $x_i$  an interval  $I$ , where

$$(1.2) \quad I = \begin{cases} (x_i, x_i + D) & \text{if } x_i + D \leq 1, \\ (x_i, 1) \cup (0, x_i + D - 1) & \text{if } x_i + D > 1; \end{cases}$$

where  $0 < D < 1$  and  $i = 0, 1, \dots, n$ .

Let  $X$  denote the random set which is the point set sum of the  $(n+1)$  intervals  $\{I\}$  and let  $\mu(X)$  be its measure. The range of  $\mu(X)$  is  $D \leq \mu(X) \leq m$ , where  $m$  denotes the minimum of 1 and  $(n+1)D$ .

D.D. Beck in his M. Sc. thesis (1968) derived the probability function of  $\mu(X)$  using the methods of D.F. Votaw Jr. (1946). The probability function,  $f_n(x)$ , of the random variable  $\mu(X)$  is given by

$$(1.3) \quad f_n(x) = n! \sum_{j=1}^q \sum_{r=0}^{q-j} \frac{(-1)^r}{(n-j)!(j-1)!} \binom{n+1}{j} \binom{n-j+1}{r} \\ \cdot (1-x)^{j-1} [x-D(j+r)]^{n-j},$$

where

$$qD \leq x < (q+1)D, \quad q = 1, 2, \dots, M;$$

$$M = \text{minimum } (n, [1/D]),$$

[a] = greatest integer less than or equal to a,

and  $x < m$ .

When  $m = (n+1)D$ , there is a finite probability that the random set  $X$  will consist of  $n+1$  non-overlapping intervals  $\{I\}$  and hence the distribution function,  $F_n(x)$ , of the variable  $\mu(X)$  has a discontinuous jump or saltus point in the case when  $m = (n+1)D$  at  $x = m$ . At such a saltus point, the probability function  $f_n(x)$ , of  $\mu(X)$ , is not defined but at all other points it defines the probability density. The expected value of  $\mu(X)$  is given by

$$(1.4) \quad E(\mu(X)) = 1 - (1-D)^{n+1}$$

and the variance is given by

$$(1.5) \quad \text{Var } (\mu(X)) = \begin{cases} \frac{1}{n+2} \{2(1-D)^{n+2} + n(1-2D)^{n+2}\} - (1-D)^{2n+2}, & 0 < D \leq \frac{1}{2}; \\ \frac{2}{n+2} (1-D)^{n+2} - (1-D)^{2n+2}, & \frac{1}{2} < D < 1. \end{cases}$$

For  $D = 1/(n+1)$ , we show in the second section of

~~Chapter IX~~ that the  $k^{\text{th}}$  non-central moment,  $m_k$ , of the random variable  $[1-\mu(X)]$  is given by

$$(1.6) \quad m_k = \frac{1}{\binom{n+k}{k}} \sum_{i=1}^k \binom{n+1}{k} \binom{k-1}{i-1} \left(1 - \frac{i}{n+1}\right)^{n+k}, \quad k = 1, 2, \dots, n+1.$$

In the last section of Chapter II we plot and tabulate the function  $f_n(x)$  for various values of  $n$  and  $D$ . In fact, we attempted to find the limiting distribution of the standardized variable  $\mu(X)$  in the case  $D = 1/(n+1)$  as  $n$  tends to infinity, which amounts to increasing the number of intervals and at the same time contracting them. Although we were unable to determine it exactly, our results suggest that the limiting distribution of the variable obtained by standardizing  $\mu(X)$  is standard normal. We give the method of analysis we were attempting and indicate where difficulties arose, in the second chapter. It is unfortunate that we have been unable to complete the analysis of the problem, but the partial analysis gives strong evidence to suggest that the limiting distribution is normal.

Research, which one might classify under the general heading of the measure of a random set, was initiated by a paper in 1939 by W.L. Stevens. Beck, in his thesis (1968), gives a good account of the various papers published in this area since then. Consider the following problem: take  $n+1$  arcs of length  $D$ ,  $0 < D < 1$ , at random points on a circle of unit circumference. Let the location of each arc be represented by the point at its clockwise

end. Choose the end of one arc arbitrarily as the origin and measure distances anticlockwise around the circle. If we consider the point set sum of all the arcs then this representation of the random set is equivalent to the linear representation described earlier. The question is what is the probability that every point of the circle is included in at least one of the arcs? Or, in other words, what is the probability that the circle is completely covered? Stevens (1939) solved this problem and proved that the required probability is given by

$$(1.7) \quad 1 - \binom{n+1}{1} (1-D)^n + \binom{n+1}{2} (1-2D)^n + \dots + (-1)^k \binom{n+1}{k} (1-kD)^n,$$

where  $k = [1/D]$ .

He also calculated the frequency distribution of the number of gaps. There is said to be a gap after the  $r^{\text{th}}$  arc if for a distance greater than  $D$  beyond the first point of this arc, there is no other arc.

In the remainder of this chapter we give the results obtained by D.F. Votaw Jr. (1946). Using the methods given in this paper, Beck (1968) derived (1.3).

Votaw considered the following non-circular representation: take a random sample  $z_i$  ( $i = 1, 2, \dots, n$ ) of  $n$  values of a one dimensional random variable  $Z$  having distribution function  $F(z)$ . Arrange the values in increasing order of magnitude as  $x_i$ ,  $i = 1, 2, \dots, n$  and consider the set  $X$  consisting of the point set sum of the following intervals:

$$(1.8) \quad (x_1 - \frac{D}{2}, x_1 + \frac{D}{2}), \quad i = 1, 2, \dots, n, \quad \text{and} \quad 0 < D < 1.$$

This representation of the random set  $X$  is not equivalent to the linear representation described earlier, which was equivalent to the circular representation. The difference arises when we consider the intervals at the end-points. Let  $\mu(X)$  be the measure of  $X$  and define a random variable  $y$  as  $y = \mu(X) - D$ . Then the range of  $y$  is  $0 \leq y \leq m_1$ , where  $m_1$  denotes the minimum of 1 and  $(n-1)D$ . Votaw showed that if  $F(z)$  is given by (1.1) then the probability function  $f_n(y)$  of  $y$  is given by

$$(1.9) \quad f_n(y) = n \sum_{j=0}^q \sum_{r=0}^{q-j} (-1)^r \binom{n-1}{j} \binom{n-1}{j+1} \binom{n-j-1}{r} \cdot (1-y)^{j+1} \{y - D(j+r)\}^{n-j-2},$$

where

$$qD \leq y < (q+1)D,$$

$$q = 0, 1, \dots, M_1;$$

$$M_1 = \min(n-2, [1/D]), \quad \text{and} \quad y < m_1.$$

He also showed that the expected value of  $y$  is given by

$$(1.10) \quad E(y) = \frac{(n-1)}{(n+1)} [1 - (1-D)^{n+1}]$$

Rewriting (1.9) we get

$$(1.11) \quad f_n(y) = n! \sum_{j=0}^q \sum_{r=0}^{q-j} \frac{(-1)^r}{(n-1-j)!j!} \binom{n-1}{j+1} \binom{n-j-1}{r}$$

$$\cdot (1-y)^{j+1} \{y - D(j+r)\}^{n-j-2},$$

with the same conditions as in (1.9). Comparing (1.3) and (1.11), we see a lot of resemblance and the differences may be attributed to the following reasons:

(i) the two representations of the random set are not equivalent and hence we have two different random variables,

(ii) Beck considered  $(n+1)$  intervals instead of the  $n$  considered by Votaw, and

(iii) Beck derived the distribution of the random variable  $\mu(X)$ , whereas Votaw derived the distribution of  $\mu(X)-D$ .

## CHAPTER II

### MAIN RESULTS

In this chapter, we present our attempt to find the limiting distribution of the random variable  $\mu(X)$ . The case  $D = 1/(n+1)$  is of particular interest because this amounts to reducing the size of each interval  $I$  and at the same time increasing the total number of such intervals in such a fashion that it is just possible for the circle to be completely covered by the arcs.

#### 2.1 Preliminary

When  $D = 1/(n+1)$ , the expressions (1.3), (1.4) and (1.5) for the probability function  $f_n(x)$ , the expected value and the variance of  $\mu(X)$  respectively reduce to

$$(2.1.1) \quad f_n(x) = n! \sum_{j=1}^q \sum_{r=0}^{q-j} \frac{(-1)^r}{(n-j)!(j-1)!} \binom{n+1}{j} \binom{n-j+1}{r} \cdot (1-x)^{j-1} [x - \frac{j+r}{n+1}]^{n-j},$$

where

$$\frac{q}{n+1} \leq x < \frac{q+1}{n+1}, \quad q = 1, 2, \dots, n \quad \text{and} \quad x < 1;$$

$$(2.1.2) \quad E(\mu(X)) = \mu_n = 1 - \left(1 - \frac{1}{n+1}\right)^{n+1}$$

and

$$(2.1.3) \quad \text{Var } (\mu(X)) = \sigma_n^2 = \frac{1}{n+2} \left\{ 2 \left(1 - \frac{1}{n+1}\right)^{n+2} + n \left(1 - \frac{2}{n+1}\right)^{n+2} - \left(1 - \frac{1}{n+1}\right)^{2n+2} \right\}$$

as  $0 < \frac{1}{n+1} \leq \frac{1}{2}$ , for all  $n \geq 1$ .

Consider the limiting behaviour of the expected value and the variance of the random variable  $\mu(X)$ . From (2.1.2),

$$(2.1.4) \quad \lim_{n \rightarrow \infty} \mu_n = \lim_{n \rightarrow \infty} \left\{ 1 - \left(1 - \frac{1}{n+1}\right)^{n+1} \right\} = 1 - e^{-1},$$

that is, the expected value tends to a constant as  $n$  goes to infinity. However, from (2.1.3),

$$\begin{aligned} (2.1.5) \quad \lim_{n \rightarrow \infty} \sigma_n^2 &= \lim_{n \rightarrow \infty} \left\{ \frac{1}{n+2} [2\left(1 - \frac{1}{n+1}\right)^{n+2} + n\left(1 - \frac{2}{n+1}\right)^{n+2} \right. \\ &\quad \left. - \left(1 - \frac{1}{n+1}\right)^{2n+2}] \right\} \\ &= \lim_{n \rightarrow \infty} \frac{1}{n+2} \left\{ 2\left[\left(1 - \frac{1}{n+1}\right)^{n+2} - \left(1 - \frac{1}{n+1}\right)^{2n+2}\right] \right. \\ &\quad \left. + n\left[\left(1 - \frac{2}{n+1}\right)^{n+2} - \left(1 - \frac{1}{n+1}\right)^{2n+2}\right] \right\} \\ &= e^{-2} - (e^{-1})^2 = 0. \end{aligned}$$

Since the limiting variance is zero, the limiting distribution of  $\mu(X)$  appears to be concentrated on the limiting mean value  $(1-e^{-1})$ . Let us try to find the rate at which  $\sigma_n^2$  tends to zero.

Lemma 2.1:

$$\lim_{n \rightarrow \infty} (n+2)\sigma_n^2 = 2e^{-1} - 5e^{-2}.$$

Proof: From (2.1.5),

$$\sigma_n^2 = \frac{2}{n+2} \left\{ \left(1 - \frac{1}{n+1}\right)^{n+2} - \left(1 - \frac{1}{n+1}\right)^{2n+2} \right\}$$

$$+ \frac{n}{n+2} \left\{ \left(1 - \frac{2}{n+1}\right)^{n+2} - \left(1 - \frac{1}{n+1}\right)^{2n+2} \right\} .$$

Now

$$\left(1 - \frac{2}{n+1}\right)^{n+2} - \left(1 - \frac{1}{n+1}\right)^{2n+2}$$

$$= \left(1 - \frac{2}{n+1}\right)^{n+2} - \left(1 - \frac{2}{n+1} + \frac{1}{(n+1)^2}\right)^{n+1}$$

$$= \left(1 - \frac{2}{n+1}\right)^{n+1} \left\{ \left(1 - \frac{2}{n+1}\right) - \left(1 + \frac{1}{(n+1)(n-1)}\right)^{n+1} \right\}$$

$$= \left(1 - \frac{2}{n+1}\right)^{n+1} \left\{ 1 - \frac{2}{n+1} - 1 - \frac{n+1}{(n+1)(n-1)} \right\}$$

$$- \frac{(n+1)n}{2(n+1)^2(n-1)^2} - \frac{(n+1)n(n-1)}{6(n+1)^3(n-1)^3} \dots \}$$

Thus

$$\lim_{n \rightarrow \infty} (n+2)\sigma_n^2 = \lim_{n \rightarrow \infty} 2 \left\{ \left(1 - \frac{1}{n+1}\right)^{n+2} - \left(1 - \frac{1}{n+1}\right)^{2n+2} \right\}$$

$$+ \lim_{n \rightarrow \infty} \left\{ -\frac{2n}{n+1} - \frac{n(n+1)}{(n+1)(n-1)} - O\left(\frac{1}{n}\right) \right\}$$

$$\cdot \left(1 - \frac{2}{n+1}\right)^{n+1}$$

$$= 2(e^{-1} - e^{-2}) + (-3)e^{-2} = 2e^{-1} - 5e^{-2} .$$

## 2.2 Limiting Distribution

To find the limiting distribution of a sequence of absolutely continuous distributions it suffices to consider the

limit of their moment generating functions  $\{M_n(t)\}$ , where

$$(2.2.1) \quad M_n(t) = E(e^{tx}) = \int_R e^{tx} f_n(x) dx ,$$

$R$  being the range of  $\mu(X)$ . Thus

$$\begin{aligned} M_n(t) &= \int_{1/(n+1)}^1 e^{tx} f_n(x) dx \\ &= \sum_{l=1}^n \int_{l/(n+1)}^{(l+1)/(n+1)} e^{tx} f_n(x) dx . \end{aligned}$$

Substituting the expression for  $f_n(x)$ , we get

$$\begin{aligned} M_n(t) &= \sum_{l=1}^n \int_{l/(n+1)}^{(l+1)/(n+1)} n! \sum_{j=1}^l \sum_{r=0}^{l-j} \frac{(-1)^r}{(n-j)!(j-1)!} \binom{n+1}{j} \binom{n-j+1}{r} \\ &\quad \cdot (1-x)^{j-1} \left(x - \frac{j+r}{n+1}\right)^{n-j} e^{tx} dx \\ &= n! \sum_{l=1}^n \sum_{j=1}^l \sum_{r=0}^{l-j} \frac{(-1)^r}{(n-j)!(j-1)!} \binom{n+1}{j} \binom{n-j+1}{r} \\ &\quad \cdot \int_{l/(n+1)}^{(l+1)/(n+1)} e^{tx} (1-x)^{j-1} \left(x - \frac{j+r}{n+1}\right)^{n-j} dx . \end{aligned}$$

Let  $(n+1)x = y$ . Then

$$\begin{aligned} (2.2.2) \quad M_n(t) &= \frac{n!}{(n+1)} \sum_{l=1}^n \sum_{j=1}^l \sum_{r=0}^{l-j} \frac{(-1)^r}{(n-j)!(j-1)!} \binom{n+1}{j} \binom{n-j+1}{r} \\ &\quad \cdot \int_l^{l+1} e^{ty/(n+1)} \left(1 - \frac{y}{n+1}\right)^{j-1} \left(\frac{y}{n+1} - \frac{j+r}{n+1}\right)^{n-j} dy \end{aligned}$$

$$= \frac{n!}{n+1} \sum_{l=1}^n \sum_{j=1}^l \sum_{r=0}^{l-j} g_{j,r} \int_l^{l+1} h_{j,r} dy,$$

where

$$g_{j,r} = \frac{(-1)^r}{(n-j)!(j-1)!} \binom{n+1}{j} \binom{n-j+1}{r}$$

and

$$h_{j,r} = e^{ty/(n+1)} \left(1 - \frac{y}{n+1}\right)^{j-1} \left(\frac{y}{n+1} - \frac{j+r}{n+1}\right)^{n-j}.$$

Consider the triple sum over  $l$ ,  $j$  and  $r$ ,

$$S = \sum_{l=1}^n \sum_{j=1}^l \sum_{r=0}^{l-j} g_{j,r} \int_l^{l+1} h_{j,r} dy.$$

When  $l = 1$ , we must have  $j = 1$  and  $r = 0$  and the contribution to  $S$  is therefore  $g_{1,0} \int_1^2 h_{1,0} dy$ . When  $l = 2$ , we must have  $j = 1$  with  $r = 0, 1$ , or  $j = 2$  with  $r = 0$ ; the contributions to  $S$  from these two cases are therefore

$$g_{1,0} \int_2^3 h_{1,0} dy + g_{1,1} \int_2^3 h_{1,1} dy$$

and

$$g_{2,0} \int_2^3 h_{2,0} dy, \text{ respectively.}$$

Continuing in this manner, it is readily verified that the contributions of the various terms resulting from the permitted combinations of  $(l, j, r)$  result in

$$\begin{aligned}
 s &= g_{1,0} \int_1^{n+1} h_{1,0} dy + g_{1,1} \int_2^{n+1} h_{1,1} dy + \dots + g_{1,n-1} \int_n^{n+1} h_{1,n-1} dy \\
 &\quad + g_{2,0} \int_2^{n+1} h_{2,0} dy + \dots + g_{2,n-2} \int_n^{n+1} h_{2,n-2} dy \\
 &\quad + \dots \dots \dots \\
 &\quad + g_{n,0} \int_n^{n+1} h_{n,0} dy \\
 &= \sum_{j=1}^n \sum_{r=0}^{n-j} g_{j,r} \int_{j+r}^{n+1} h_{j,r} dy
 \end{aligned}$$

Thus

$$\begin{aligned}
 (2.2.3) \quad M_n(t) &= \frac{n!}{n+1} \sum_{j=1}^n \sum_{r=0}^{n-j} \frac{(-1)^r}{(n-j)!(j-1)!} \binom{n+1}{j} \binom{n-j+1}{r} \\
 &\quad \cdot \int_{j+r}^{n+1} e^{ty/(n+1)} (1 - \frac{y}{n+1})^{j-1} (\frac{y}{n+1} - \frac{j+r}{n+1})^{n-j} dy
 \end{aligned}$$

Let  $z = \frac{y-(j+r)}{n+1-j-r}$ , so that  $0 \leq z \leq 1$  and  $y = [(n+1)-j-r]z + j+r$ ,  
 $dy = (n+1-j-r)dz$ . Then

$$\begin{aligned}
 M_n(t) &= \frac{n!}{n+1} \sum_{j=1}^n \sum_{r=0}^{n-j} \frac{(-1)^r}{(n-j)!(j-1)!} \binom{n+1}{j} \binom{n-j+1}{r} \\
 &\quad \cdot \int_0^1 e^{t(j+r+(n+1-j-r)z)/(n+1)} (1 - \frac{j+r+(n+1-j-r)z}{n+1})^{j-1} \\
 &\quad \cdot (\frac{j+r+(n+1-j-r)z-j-r}{n+1})^{n-j} \frac{1}{(n+1-j-r)} dz \\
 &= \frac{n!}{n+1} \sum_{j=1}^n \sum_{r=0}^{n-j} \frac{(-1)^r}{(n-j)!(j-1)!} \binom{n+1}{j} \binom{n-j+1}{r} e^{t(j+r)/(n+1)} \\
 &\quad \cdot \frac{(n+1-j-r)^n}{(n+1)^{n-1}} \int_0^1 e^{t(1 - \frac{j+r}{n+1})z} z^{n-j} (1-z)^{j-1} dz
 \end{aligned}$$

$$= n \sum_{j=1}^n \sum_{r=0}^{n-j} (-1)^r \frac{(n-1)!}{(n-j)!(j-1)!} \frac{n!}{j!(n+1-j)!} \frac{(n+1-j)!}{r!(n+1-j-r)!}$$

$$\cdot e^{t(j+r)/(n+1)} (n+1-j-r)(1 - \frac{j+r}{n+1})^{n-1}$$

$$\cdot \int_0^1 e^{t(1 - \frac{j+r}{n+1})z} z^{n-j} (1-z)^{j-1} dz$$

$$= n \sum_{j=1}^n \sum_{r=0}^{n-j} (-1)^r \binom{n-1}{j-1} \binom{n}{j} \binom{n-j}{r} e^{t(j+r)/(n+1)} (1 - \frac{j+r}{n+1})^{n-1}$$

$$\cdot \int_0^1 e^{t(1 - \frac{j+r}{n+1})z} z^{n-j} (1-z)^{j-1} dz.$$

Let  $(1-z) = q, dz = -dq$ . Then

$$(2.2.4) \quad M_n(t) = n \sum_{j=1}^n \sum_{r=0}^{n-j} (-1)^r \binom{n-1}{j-1} \binom{n}{j} \binom{n-j}{r} e^{t(j+r)/(n+1)} (1 - \frac{j+r}{n+1})^{n-1}$$

$$\cdot \int_0^1 e^{t(1 - \frac{j+r}{n+1})(1-q)} (1-q)^{n-j} q^{j-1} dq.$$

Consider the standardized variable  $\frac{\mu(X)-\mu_n}{\sigma_n}$  with moment generating function

$$\begin{aligned} \psi_n(t) &= E[e^{((\mu(X)-\mu_n)/\sigma_n)t}] = e^{-\mu_n t/\sigma_n} E(e^{\mu(X)t/\sigma_n}) \\ &= e^{-\mu_n t/\sigma_n} M_n(t/\sigma_n). \end{aligned}$$

From (2.2.4)

$$\psi_n(t) = n e^{(t/\sigma_n) - (\mu_n t/\sigma_n)} \int_0^1 \sum_{j=1}^n \frac{(n-1)_j}{(j-1)} \frac{\binom{n}{j}}{(1-q)^{n-j}} q^{j-1} \\ \cdot e^{-(tq/\sigma_n) + (tqj/(\sigma_n(n+1)))} \sum_{r=0}^{n-j} (-1)^r \binom{n-j}{r} e^{tqr/((n+1)\sigma_n)} \\ \cdot \left(1 - \frac{j+r}{n+1}\right)^{n-1} dq.$$

Let

$$(2.2.5) \quad \left\{ \begin{array}{l} \left(1 - \frac{y}{n+1}\right)^{n-1} = e^{-y} \sum_{k=0}^{\infty} a_k(n) y^k \quad \text{so that} \\ \sum_{k=0}^{\infty} a_k(n) y^k = e^y \left(1 - \frac{y}{n+1}\right)^{n-1}. \end{array} \right.$$

For  $y = j+r$ , this yields

$$\left(1 - \frac{j+r}{n+1}\right)^{n-1} = e^{-(j+r)} \sum_{k=0}^{\infty} a_k(n) (j+r)^k$$

and, since

$$(j+r)^k = \frac{d^k}{dp^k} (e^{(j+r)p}) \Big|_{p=0}$$

we have

$$\left(1 - \frac{j+r}{n+1}\right)^{n-1} = e^{-(j+r)} \sum_{k=0}^{\infty} a_k(n) \frac{d^k}{dp^k} (e^{(j+r)p}) \Big|_{p=0}.$$

Substituting this into the expression for  $\psi_n(t)$ , we get

\* This method of dealing with the term  $\left(1 - \frac{j+r}{n+1}\right)^{n-1}$  was suggested by Dr. J.R. McGregor.

$$(2.2.6) \quad \psi_n(t) = \sum_{k=0}^{\infty} a_k(n) \psi_n^k(t,0)$$

where

$$\psi_n^k(t,0) = \frac{d^k}{dp^k} \psi_n(t,p) \Big|_{p=0}$$

and

$$(2.2.7) \quad \begin{aligned} \psi_n(t,p) &= ne^{-t(1-\mu_n)/\sigma_n} \int_0^1 e^{-tq/\sigma_n} \\ &\cdot \sum_{j=1}^n \binom{n-1}{j-1} \binom{n}{j} (1-q)^{n-j} q^{j-1} e^{tqj/((n+1)\sigma_n)} \\ &\cdot \sum_{r=0}^{n-j} (-1)^r \binom{n-j}{r} e^{tqr/((n+1)\sigma_n) - (j+r) + p(j+r)} dq \\ &= ne^{-t(1-\mu_n)/\sigma_n} \int_0^1 e^{-tq/\sigma_n} \sum_{j=1}^n \binom{n-1}{j-1} \binom{n}{j} (1-q)^{n-j} q^{j-1} \\ &\cdot \{e^{tq/((n+1)\sigma_n) + p-1} \}^j \\ &\cdot \sum_{r=0}^{n-j} (-1)^r \binom{n-j}{r} \{e^{tq/((n+1)\sigma_n) + p-1} \}^r dq \\ &= ne^{-t(1-\mu_n)/\sigma_n} \int_0^1 e^{-tq/\sigma_n} \\ &\cdot \sum_{j=1}^n \binom{n-1}{j-1} \binom{n}{j} (1-q)^{n-j} q^{j-1} \beta^j (1-\beta)^{n-j} dq, \end{aligned}$$

where

$$(2.2.8) \quad \beta = e^{tq/((n+1)\sigma_n) + p-1}$$

Thus

$$(2.2.9) \quad \psi_n(t, p) = ne^{t(1-\mu_n)/\sigma_n} \int_0^1 e^{-tq/\sigma_n} \beta \cdot \sum_{j=1}^n \binom{n-1}{j-1} \binom{n}{j} (q\beta)^{j-1} \{(1-q)(1-\beta)\}^{n-j} dq.$$

Consider

$$\begin{aligned} s_1 &= \sum_{j=1}^n \binom{n-1}{j-1} \binom{n}{j} (q\beta)^{j-1} \{(1-q)(1-\beta)\}^{n-j} \\ &= \sum_{v=0}^{n-1} \binom{n-1}{v} \binom{n}{v+1} (q\beta)^v \{(1-q)(1-\beta)\}^{n-1-v} \\ &= \{(1-q)(1-\beta)\}^{n-1} \sum_{v=0}^{n-1} \binom{n-1}{v} \binom{n-1+1}{n-1-v} \left\{ \frac{q\beta}{(1-q)(1-\beta)} \right\}^v. \end{aligned}$$

Let

$$\frac{q\beta}{(1-q)(1-\beta)} = \frac{s-1}{s+1} \quad \text{where} \quad s = \frac{q\beta + (1-q)(1-\beta)}{(1-q)(1-\beta) - q\beta}$$

Then

$$\begin{aligned} s_1 &= \{(1-q)(1-\beta)\}^{n-1} \sum_{v=0}^{n-1} \binom{n-1}{v} \binom{n-1+1}{n-1-v} \left\{ \frac{(s-1)/2}{(s+1)/2} \right\}^v \\ &= \frac{\{(1-q)(1-\beta)\}^{n-1}}{(s+1/2)^{n-1}} \sum_{v=0}^{n-1} \binom{n-1}{v} \binom{n-1+1}{n-1-v} \left( \frac{s-1}{2} \right)^v \left( \frac{s+1}{2} \right)^{n-1-v} \\ &= \left\{ \frac{2(1-q)(1-\beta)}{s+1} \right\}^{n-1} P_{n-1}^{(1,0)}(s), \end{aligned}$$

where  $P_{n-1}^{(1,0)}(s)$  is the Jacobi polynomial with parameters  $\alpha = 1$ ,  $\beta = 0$  (see Szegő (1974), equation (4.3.2)). Now

$$s = \frac{q\beta + (1-q)(1-\beta)}{(1-q)(1-\beta) - q\beta} = \frac{q\beta + (1-q)(1-\beta)}{1-q-\beta},$$

so that

$$s+1 = \frac{q\beta + 1 - q - \beta + q\beta + 1 - q - \beta}{1 - q - \beta} = \frac{2(1-q)(1-\beta)}{1-q-\beta}$$

giving

$$(2.2.10) \quad s_1 = (1-q-\beta)^{n-1} P_{n-1}^{(1,0)} \left( \frac{(1-q)(1-\beta)+q\beta}{1-q-\beta} \right).$$

Consider the following theorem of Darboux which is given in Szegő (1974, page 196).

Theorem 2.2.1: Let  $\alpha$  and  $\beta$  be arbitrary real numbers. Then

$$P_n^{(\alpha, \beta)}(x) \sim (x-1)^{-\alpha/2} (x+1)^{-\beta/2} \{ (x+1)^{1/2} + (x-1)^{1/2} \}^{\alpha+\beta} \cdot (2\pi n)^{-1/2} (x^2-1)^{-1/4} \{ x + (x^2-1)^{1/2} \}^{n+1/2},$$

where  $x$  is outside of the closed interval  $[-1,1]$ . This formula holds uniformly in the exterior of an arbitrary closed curve which encloses the segment  $[-1,1]$ , in the sense that the ratio tends uniformly to 1.

Lemma 2.2.1:  $s = \frac{(1-q)(1-\beta)+q\beta}{1-q-\beta}$  lies outside the interval  $[-1,1]$ .

Proof: Recall that  $\beta = e^{-tq/(n+1)\sigma_n^2} + p-1$ ,  $0 < q < 1$  and also  $\beta$  need only be differentiated with respect to  $p$  and  $t$  in a neighbourhood of  $p = 0$ ,  $t = 0$ . Therefore, we may take  $-\epsilon < t < \epsilon$  and  $-\delta < p < \delta$ , and choose  $\epsilon$  and  $\delta$  arbitrarily small. Thus

$$-\frac{\epsilon q}{(n+1)\sigma_n} - \delta - 1 < \frac{tq}{(n+1)\sigma_n} + p - 1 < \frac{\epsilon q}{(n+1)\sigma_n} + \delta - 1$$

or

$$\frac{-1-(\epsilon q/((n+1)\sigma_n)+\delta)}{\epsilon} < \beta < \frac{-1+(\epsilon q/((n+1)\sigma_n)+\delta)}{\epsilon}$$

Thus  $e^{-1} < 1/2$  implies that we can choose  $\epsilon$  and  $\delta$  so that  
 $0 < \beta < 1/2$ . Also, since  $0 < q < 1$ , there exists  $q_0$  such that  
 $q_0 = 1-\beta$ . Now

$$\frac{d}{dq} \left[ \frac{(1-q)(1-\beta)+q\beta}{1-q-\beta} \right] = \frac{2\beta}{(1-\beta-q)^2} [1-\beta + \frac{q(1-q)t}{(n+1)\sigma_n}]$$

By restricting  $t$  to an open interval  $(-\epsilon, \epsilon)$ ,  $\epsilon > 0$ , of arbitrarily small width, the derivative is positive over the range of  $q$  except when the function itself is not defined at  $q = 1-\beta$ .

Now at,

$$q = 0, \frac{(1-q)(1-\beta)+q\beta}{1-q-\beta} = 1 \quad \text{and} \quad q = 1, \frac{(1-q)(1-\beta)+q\beta}{1-q-\beta} = -1$$

Therefore  $s$  lies outside the interval  $[-1, +1]$  when  $q \in (0, 1)$ .

Thus Theorem 2.2.1 is applicable. Note that

$$s-1 = \frac{q\beta+1-q-\beta+q\beta-1+q+\beta}{1-q-\beta} = \frac{2q\beta}{1-q-\beta}$$

so that

$$s^2-1 = \frac{4q\beta(1-q)(1-\beta)}{(1-q-\beta)^2}$$

Then for large  $n$ ,

$$\begin{aligned}
 p_{n-1}^{(1,0)} \left( \frac{(1-q)(1-\beta)+q\beta}{1-q-\beta} \right) &\approx \left( \frac{2q\beta}{1-q-\beta} \right)^{-1/2} \left[ \left( \frac{2(1-q)(1-\beta)}{1-q-\beta} \right)^{1/2} + \left( \frac{2q\beta}{1-q-\beta} \right)^{1/2} \right] \\
 &\cdot [2\pi(n-1)]^{-1/2} \left[ \frac{4q\beta(1-q)(1-\beta)}{(1-q-\beta)^2} \right]^{-1/4} \\
 &\cdot \left\{ \frac{q\beta+(1-q)(1-\beta)}{1-q-\beta} + \frac{2q^{1/2}\beta^{1/2}(1-q)^{1/2}(1-\beta)^{1/2}}{1-q-\beta} \right\}^{n-1+(1/2)} \\
 &= \frac{(1-q-\beta)^{1/2}}{\sqrt{2} q^{1/2} \beta^{1/2}} \cdot \frac{1}{\sqrt{2} \pi^{1/2} (n-1)^{1/2}} \cdot \frac{(1-q-\beta)^{1/2}}{\sqrt{2} q^{1/4} \beta^{1/4} (1-q)^{1/4} (1-\beta)^{1/4}} \\
 &\cdot \frac{\sqrt{2}}{(1-q-\beta)^{1/2}} \left\{ q^{1/2} \beta^{1/2} + (1-q)^{1/2} (1-\beta)^{1/2} \right\} \\
 &\cdot \frac{1}{(1-q-\beta)^{n-1/2}} \left\{ [q^{1/2} \beta^{1/2} + (1-q)^{1/2} (1-\beta)^{1/2}]^2 \right\}^{n-1/2} \\
 &= \frac{[q^{1/2} \beta^{1/2} + (1-q)^{1/2} (1-\beta)^{1/2}]^{2n}}{2\{(n-1)\pi\}^{1/2} (1-q-\beta)^{n-1} q^{3/4} \beta^{3/4} (1-q)^{1/4} (1-\beta)^{1/4}}
 \end{aligned}$$

Substituting in (2.2.10) we get

$$s_1 \approx \frac{[q^{1/2} \beta^{1/2} + (1-q)^{1/2} (1-\beta)^{1/2}]^{2n}}{2\{(n-1)\pi\}^{1/2} q^{3/4} \beta^{3/4} (1-q)^{1/4} (1-\beta)^{1/4}}$$

which when substituted in (2.2.9) yields

$$\begin{aligned}
 (2.2.11) \quad \Psi_n(t, p) &\approx \frac{ne^{t(l-\mu_n)/\sigma_n}}{2\{(n-1)\pi\}^{1/2}} \\
 &\cdot \int_0^1 e^{-tq/\sigma_n} \frac{[q^{1/2} \beta^{1/2} + (1-q)^{1/2} (1-\beta)^{1/2}]^{2n}}{q^{3/4} (1-q)^{1/4}} \left( \frac{\beta}{1-\beta} \right)^{1/4} dq.
 \end{aligned}$$

To evaluate this integral, we use the following generalization of Laplace's method (see Olver, 1974, page 333):

**Theorem 2.2.2:** Let

$$I(x) = \int_0^k \exp \{-xp(t) + x^{\nu/\mu} r(t)\} q(t) dt ;$$

where  $k$  is fixed and positive, and assume that:

(i) In the interval  $(0, k]$ ,  $p(t)$  is continuous and positive, and the real or complex functions  $q(t)$  and  $r(t)$  are continuous.

(ii) As  $t \rightarrow 0+$ ,

$$p(t) = p(0) + P t^\mu + O(t^{\mu_1}) ;$$

$$p'(t) = \mu P t^{\mu-1} + O(t^{\mu_1-1}) ;$$

$$q(t) = Q t^{\lambda-1} + O(t^{\lambda_1-1}) ;$$

$$r(t) = R t^\nu + O(t^{\nu_1}) ; \quad \text{where } P > 0 ,$$

$$\mu_1 > \mu > 0, \quad \lambda_1 > \lambda > 0, \quad \mu > \nu \geq 0, \quad \text{and} \quad \nu_1 > \nu .$$

Then

$$I(x) = \frac{Q}{\mu} F_1 \left( \frac{\nu}{\mu}, \frac{\lambda}{\mu}; \frac{R}{P^{\nu/\mu}} \right) \frac{e^{-xp(0)}}{(Px)^{\lambda/\mu}} \left\{ 1 + O\left(\frac{1}{x^{\tilde{w}/\mu}}\right) \right\}$$

as  $x \rightarrow \infty$ , where

$$\tilde{w} = \min (\lambda_1 - \lambda, \mu_1 - \mu, \nu_1 - \nu) \quad \text{and}$$

$F_1(a, b; y)$  is the Faxen's integral

$$F_1(\alpha, \beta; y) = \int_0^\infty \exp\{-\tau + y\tau^\alpha\} \tau^{\beta-1} d\tau,$$

$$(0 \leq \operatorname{Re} \alpha < 1, \operatorname{Re} \beta > 0)$$

In particular

$$F_1\left(\frac{1}{2}, \frac{1}{2}; y\right) + F_1\left(\frac{1}{2}, \frac{1}{2}; -y\right) = 2\pi^{1/2} \exp\left(\frac{1}{4}y^2\right).$$

To apply this theorem to the integral in (2.2.11), note that

$$(i) \lim_{n \rightarrow \infty} \mu_n = 1 - e^{-1},$$

$$(ii) \lim_{n \rightarrow \infty} (n+2) \sigma_n^2 = 2e^{-1} - 5e^{-2} \text{ so that } \sigma_n^2 = \frac{k_n^2}{n}$$

$$\text{where } k_n^2 = 2e^{-1} - 5e^{-2}, \text{ and}$$

$$(iii) \beta = e^{\frac{tq}{((n+1)\sigma_n)} + p-1} \text{ so that for large } n, \beta \approx e^{p-1}.$$

Then (2.2.11) reduces to

$$\begin{aligned} \Psi_n(t, p) &\approx \frac{ne^{-1}\sqrt{n}/k_n}{2\{(n-1)\pi\}^{1/2}} \int_0^1 e^{-\sqrt{n}tq/k_n} \frac{e^{(p-1)/4}}{q^{3/4}(1-q)^{1/4}(1-e^{p-1})^{1/4}} \\ &\quad \cdot \{q^{1/2} e^{(p-1)/2} + (1-q)^{1/2} (1-e^{p-1})^{1/2}\}^{2n} dq \\ (2.2.12) &= \frac{ne^{(p-1)/4}}{2\{(n-1)\pi\}^{1/2}(1-e^{p-1})^{1/4}} \int_0^1 \frac{e^{(\sqrt{n}t/k_n)(e^{-1}-q)}}{q^{3/4}(1-q)^{1/4}} \theta^{2n} dq, \end{aligned}$$

where

$$\theta(q, p) \equiv \theta = q^{1/2} e^{(p-1)/2} + (1-q)^{1/2} (1-e^{p-1})^{1/2}.$$

Let

$$\begin{aligned} I_n &= \int_0^1 e^{(\sqrt{nt}/k_n)(e^{-1}-q)} \frac{\theta^{2n}}{q^{3/4}(1-q)^{1/4}} dq \\ &= \int_0^1 \exp \{-n(-2\ln \theta) + n^{1/2} \frac{t}{k_n} (e^{-1}-q)\} (q^{3/4}(1-q)^{1/4})^{-1} dq. \end{aligned}$$

Note that  $\theta(0, p) = (1-e^{p-1})^{1/2}$ ,  $\theta(1, p) = e^{(p-1)/2}$  and

$\theta(e^{p-1}, p) = e^{p-1} + 1 - e^{p-1} = 1$ . Also,

$$\frac{d}{dq} \theta(q, p) = \frac{1}{2} q^{-1/2} e^{(p-1)/2} - \frac{1}{2} (1-q)^{-1/2} (1-e^{p-1})^{1/2}$$

so that  $\frac{d}{dq} \theta(e^{p-1}, p) = 0$ . Let  $y = q - e^{p-1}$  so that when  $q = 0$ ,  $y = -e^{p-1}$  and when  $q = 1$ ,  $y = 1 - e^{p-1}$ . Thus

$$\begin{aligned} I_n &= e^{(\sqrt{nt}/k_n)(e^{-1}-e^{p-1})} \left\{ \int_{-e^{p-1}}^0 + \int_0^{1-e^{p-1}} \right\} \\ &\quad \exp \{-n(-2\ln \theta(y+e^{p-1}, p)) + n^{1/2} \frac{t}{k_n} (-y)\} \\ &\quad [(y+e^{p-1})^{3/4} (1-y-e^{p-1})^{1/4}]^{-1} dy \\ &= I_{n_1} + I_{n_2} \end{aligned}$$

Consider  $I_{n_1}$  and put  $w = -y$ . Then

$$I_{n_1} = e^{(\sqrt{nt}/k_n)(e^{-1}-e^{p-1})} \int_0^{e^{p-1}} \exp \{-np(w) + n^{1/2} r(w)\} q(w) dw,$$

where

$$p(w) = -2 \ln \{(1+w-e^{p-1})^{1/2} (1-e^{p-1})^{1/2} + (e^{p-1}-w)^{1/2} e^{(p-1)/2}\}$$

$$r(w) = \frac{t}{k_n} w \quad \text{and} \quad q(w) = [(e^{p-1}-w)^{3/4} (1+w-e^{p-1})^{1/4}]^{-1}$$

Compare with Theorem 2.2.2, then  $\frac{v}{\mu} = \frac{1}{2}$  and also as  $r(w) = \frac{t}{k_n} w$ ,

this implies  $R = \frac{t}{k_n}$  and  $v = 1$  so that  $\mu = 2$ . Now to find

the value of  $P$ , note that  $\frac{p(w) - p(0)}{w^2} \rightarrow P$  as  $w \rightarrow 0+$  and  
also  $p(0) = 0$ . Thus

$$\lim_{w \rightarrow 0+} \frac{p(w) - p(0)}{w^2} = \lim_{w \rightarrow 0+} \frac{p(w)}{w^2} = \lim_{w \rightarrow 0+} \frac{p'(w)}{2w}$$

Now

$$p'(w) = \frac{-2\{\frac{1}{2}(1+w-e^{p-1})^{-1/2}(1-e^{p-1})^{1/2} - \frac{1}{2}(e^{p-1}-w)^{-1/2}e^{(p-1)/2}\}}{(1+w-e^{p-1})^{1/2}(1-e^{p-1})^{1/2} + (e^{p-1}-w)^{1/2}e^{(p-1)/2}}$$

and so  $p'(0) = 0$ . Therefore

$$\begin{aligned} P &= \lim_{w \rightarrow 0+} \frac{p''(w)}{2} = \frac{1}{4} \{(1-e^{p-1})^{-1/2} + e^{-(p-1)/2} \\ &= \frac{1}{4\{e^{p-1}(1-e^{p-1})\}} \end{aligned}$$

Also

$$\lim_{w \rightarrow 0+} \frac{q(w)}{w^{\lambda-1}} = Q$$

Let  $\lambda = 1$ , then  $\lim_{w \rightarrow 0+} q(w) = Q$  which implies

$$Q = [e^{3(p-1)/4} (1-e^{p-1})^{1/4}]^{-1}$$

Now applying Theorem (2.2.2),

$$I_{n_1} \approx e^{\frac{(\sqrt{nt}/k_n)(e^{-1}-e^{p-1})}{2(nP)^{1/2}}} \frac{Q}{F_1(\frac{1}{2}, \frac{1}{2}; -\frac{R}{P^{1/2}})}$$

Similarly,

$$I_{n_2} \approx e^{\frac{(\sqrt{nt}/k_n)(e^{-1}-e^{p-1})}{2(nP)^{1/2}}} \frac{Q}{F_1(\frac{1}{2}, \frac{1}{2}; -\frac{R}{P^{1/2}})}$$

Thus

$$I_n = I_{n_1} + I_{n_2}$$

$$\approx e^{\frac{(\sqrt{nt}/k_n)(e^{-1}-e^{p-1})}{2(nP)^{1/2}}} \cdot 2\pi^{1/2} \exp\left(\frac{R^2}{4P}\right)$$

Substituting this into (2.2.12), we get

$$\begin{aligned} \psi_n(t, p) &\approx \frac{n e^{(p-1)/4}}{2((n-1)\pi)^{1/2} (1-e^{p-1})^{1/4}} e^{\frac{(\sqrt{nt}/k_n)(e^{-1}-e^{p-1})}{2(nP)^{1/2}}} \\ &\quad \cdot \frac{2\pi^{1/2} 2 e^{(p-1)/2} (1-e^{p-1})^{1/2}}{2n^{1/2} e^{3(p-1)/4} (1-e^{p-1})^{1/4}} \exp\left\{\frac{t^2}{k_n^2} e^{p-1} (1-e^{p-1})\right\} \\ &\approx e^{\frac{(\sqrt{nt}/k_n)(e^{-1}-e^{p-1})}{2(nP)^{1/2}}} + \frac{t^2}{k_n^2} e^{p-1} (1-e^{p-1}) \\ &= \exp\{g(p)\}, \end{aligned}$$

where

$$g(p) = \frac{\sqrt{n} t}{k_n} (e^{-1}-e^{p-1}) + \frac{t^2}{k_n^2} e^{p-1} (1-e^{p-1})$$

Now

$$\frac{d}{dp} \psi_n(t, p) \approx e^{g(p)} g'(p),$$

$$\frac{d^2}{dp^2} \psi_n(t, p) \approx e^{g(p)} \{(g'(p))^2 + g''(p)\},$$

$$\frac{d^3}{dp^3} \psi_n(t, p) \approx e^{g(p)} \{(g'(p))^3 + 3g'(p)g''(p) + g'''(p)\}$$

...

...

$$\frac{d^k}{dp^k} \psi_n(t, p) \approx e^{g(p)} \{(g'(p))^k + (2k-3)(g'(p))^{k-2}g''(p) + \dots\}.$$

Now

$$g'(p) = -\frac{\sqrt{n}t e^{p-1}}{k_n} + \frac{t^2}{k_n^2} \{e^{p-1}(1-e^{p-1}) - e^{2(p-1)}\}$$

and

$$g''(p) = -\frac{\sqrt{n}t e^{p-1}}{k_n} + \frac{t^2}{k_n^2} \{e^{p-1}(1-e^{p-1}) - 3e^{2(p-1)}\}.$$

¶

In general, we shall have

$$\frac{d^k}{dp^k} \psi_n(t, 0) \approx e^{(t^2/k_n^2)e^{-1}(1-e^{-1})} \left\{ \left(-\frac{\sqrt{n}t e^{-1}}{k_n}\right)^k \right\} (1 + o(n^{-1/2})).$$

Substituting this into (2.2.6) we get

$$\psi_n(t) \approx \sum_{k=0}^{\infty} a_k(n) e^{(t^2/k_n^2)e^{-1}(1-e^{-1})} \left(-\frac{\sqrt{n}t e^{-1}}{k_n}\right)^k,$$

which by (2.2.5) is

$$\begin{aligned}
 &= e^{(t^2/k_n^2)e^{-1}(1-e^{-1})} e^{-\sqrt{n}te^{-1}/k_n} \left(1 + \frac{\sqrt{n}te^{-1}}{k_n(n+1)}\right)^{n-1} \\
 &\approx e^{(t^2/k_n^2)e^{-1}(1-e^{-1})} e^{-\sqrt{n}te^{-1}/k_n} (n-1) \ln(1 + (\sqrt{n}te^{-1})/(k_n(n+1))) \\
 &\approx e^{(t^2/k_n^2)e^{-1}(1-e^{-1})} e^{-\sqrt{n}te^{-1}/k_n} \\
 &\quad \cdot e^{(n-1)[\sqrt{n}te^{-1}/(k_n(n+1)) - \frac{1}{2}nt^2e^{-1}/(k_n^2(n+2)^2)] + \dots} \\
 &\approx e^{(t^2/k_n^2)e^{-1}(1-e^{-1})} e^{-\sqrt{n}te^{-1}/k_n} \\
 &\approx e^{[\sqrt{n}te^{-1}/k_n - t^2e^{-2}/2k_n^2]\{1+o(n^{-1})\}} \\
 &\approx e^{(t^2/k_n^2)\{e^{-1}-e^{-2} - \frac{1}{2}e^{-2}\}} = e^{(t^2/2k_n^2)(2e^{-1}-3e^{-2})}
 \end{aligned}$$

Note that  $k_n^2 \rightarrow 2e^{-1}-5e^{-2}$  as  $n$  goes to infinity. Hence

$$\lim_{n \rightarrow \infty} \psi_n(t) = e^{(t^2/2)\{(2e^{-1}-3e^{-2})/(2e^{-1}-5e^{-2})\}}$$

Since for every  $n$ ,  $\psi_n(t)$  is the moment generating function of a random variable with zero mean and unit variance the same should be true of  $\lim_{n \rightarrow \infty} \psi_n(t)$ . It is clear, therefore, that at some stage the approximations used have been unsatisfactory since the result obtained above implies that the limiting distribution is normal with zero mean and variance  $\{(2e^{-1}-3e^{-2})/(2e^{-1}-5e^{-2})\}$  instead of unity. We have not been able to refine the approximations to eliminate this discrepancy. It is perhaps of some interest to

note that the asymptotic evaluations of integrals by modifications of the Laplace method, such as we have used here, often result in functions which are exact except for the constants involved. Thus, for example, if one applies the method of Steepest Descents to determine the F-distribution the result is exact except that the factorials in the normalizing constant are replaced by their Stirling's approximations. This may provide a clue of where one should look to refine the analysis to produce the exact result.

### 2.3 Moments of the Variable $\{1-\mu(X)\}$

For  $D = 1/(n+1)$ , this section contains a result about the non-central moments of the random variable  $(1-\mu(X))$ .

Theorem 2.3.1: The  $k^{\text{th}}$  non-central moment of the variable  $\{1-\mu(X)\}$  when  $D = 1/(n+1)$  is given by

$$(2.3.1) \quad m_k = \frac{1}{\binom{n+k}{k}} \sum_{i=1}^k \binom{n+1}{i} \binom{k-1}{i-1} \left(1 - \frac{i}{n+1}\right)^{n+k}$$

$$k = 1, 2, \dots, n+1$$

Proof:

$$m_k = E(1-\mu(X))^k$$

$$= \int_{1/(n+1)}^1 (1-x)^k f_n(x) dx$$

$$= \sum_{i=1}^n \int_{i/(n+1)}^{(i+1)/(n+1)} (1-x)^k f_n(x) dx$$

$$= \sum_{l=1}^n \int_{l/(n+1)}^{(l+1)/(n+1)} n! \sum_{j=1}^l \sum_{r=0}^{l-j} \frac{(-1)^r}{(n-j)!(j-1)!} \binom{n+1}{j} \binom{n-j+1}{r} \\ \cdot (1-x)^{k+j-1} [x - \frac{j+r}{n+1}]^{n-j} dx .$$

Comparing with (2.2.2) and following the same analysis as done there,  
we get

$$(2.3.2) \quad m_k = \frac{n!}{(n+1)} \sum_{j=1}^n \sum_{r=0}^{n-j} \frac{(-1)^r}{(n-j)!(j-1)!} \binom{n+1}{j} \binom{n-j+1}{r} \\ \cdot \int_{j+r}^{n+1} \left(1 - \frac{y}{n+1}\right)^{k+j-1} \left(\frac{y-j-r}{n+1}\right)^{n-j} dy .$$

Let

$$I = \int_{(j+r)}^{(n+1)} \left(1 - \frac{y}{n+1}\right)^{k+j-1} \left(\frac{y-j-r}{n+1}\right)^{n-j} dy$$

and take  $z = \frac{y-j-r}{n+1-j-r}$  so that  $0 \leq z \leq 1$  and  $dy = (n+1-j-r)dz$ .

Then

$$I = \int_0^1 \left(1 - \frac{j+r+(n+1-j-r)z}{n+1}\right)^{k+j-1} \\ \cdot \left(\frac{j+r+(n+1-j-r)z-j-r}{n+1}\right)^{n-j} (n+1-j-r) dz \\ = \frac{(n+1-j-r)^{n+k}}{(n+1)^{n+k-1}} \int_0^1 (1-z)^{k+j-1} z^{n-j} dz \\ = (n+1) \left(1 - \frac{j+r}{n+1}\right)^{n+k} \beta(n-j+1, k+j)$$

$$= (n+1) \left(1 - \frac{j+r}{n+1}\right)^{n+k} \frac{(n-j)! (j+k-1)!}{(n+k)!}$$

Substituting in (2.3.2) we get

$$\begin{aligned} m_k &= \frac{n!}{(n+k)!} \sum_{j=1}^n \sum_{r=0}^{n-j} \frac{(-1)^r}{(n-j)! (j-1)!} \binom{n+1}{j} \binom{n-j+1}{r} \left(1 - \frac{j+r}{n+1}\right)^{n+k} \\ &\quad \cdot (n+1) \frac{(n-j)! (j+k-1)!}{(n+k)!} \end{aligned}$$

Put  $(j+r) = i$ , then

$$\begin{aligned} m_k &= \frac{n!}{(n+k)!} \sum_{j=1}^n \sum_{i=j}^n \frac{(-1)^{i-j}}{(j-1)!} \frac{(n+1)!}{j! (n+1-j)!} \frac{(n+1-j)!}{(i-j)! (n-i+1)!} \\ &\quad \cdot (k+j-1)! \left(1 - \frac{1}{n+1}\right)^{n+k} \\ &= \frac{n! k!}{(n+k)!} \sum_{i=1}^n \sum_{j=1}^i (-1)^{i-j} \left\{ \frac{(k+j-1)!}{(j-1)! k!} \right\} \left\{ \frac{(n+1)!}{i! (n+1-i)!} \right\} \\ &\quad \cdot \left\{ \frac{i!}{j! (i-j)!} \right\} \left(1 - \frac{1}{n+1}\right)^{n+k} \end{aligned}$$

Thus

$$(2.3.3) \quad m_k = \frac{1}{\binom{n+k}{k}} \sum_{i=1}^n \sum_{j=1}^i (-1)^{i-j} \binom{k+j-1}{j-1} \binom{i}{j} \cdot \binom{n+1}{i} \left(1 - \frac{1}{n+1}\right)^{n+k}$$

Consider the sum

$$\sum_{j=1}^i (-1)^{i-j} \binom{k+j-1}{j-1} \binom{i}{j} = \sum_{j=0}^i (-1)^{i+j} \binom{i}{j} \binom{k+j-1}{j-1}$$

Compare this with the identity

$$\binom{n+p}{m+p} = \sum_{k_1=0}^m (-1)^{k_1+m} \binom{m}{k_1} \binom{n+p+k_1}{p+k_1}$$

(see Riordan (1968), page 11). Thus identifying the variables in the two sums, we get

$$\binom{k-1}{i-1} = \sum_{j=0}^i (-1)^{i+j} \binom{i}{j} \binom{k+j-1}{j-1}$$

Thus (2.3.3) reduces to

$$\begin{aligned} m_k &= \frac{1}{\binom{n+k}{k}} \sum_{i=1}^n \binom{n+1}{i} \binom{k-1}{i-1} \left(1 - \frac{1}{n+1}\right)^{n+k} \\ &= \frac{1}{\binom{n+k}{k}} \sum_{i=1}^k \binom{n+1}{i} \binom{k-1}{i-1} \left(1 - \frac{1}{n+1}\right)^{n+k}, \\ k &= 1, 2, \dots, n+1. \end{aligned}$$

#### 2.4 Plots and Tables

The results of the previous section strongly suggest that the limiting distribution of the standardized random variable  $\mu(X)$  is standard normal when  $D = 1/(n+1)$ . In this section we examine the probability function,  $f_n(x)$ , of the random variable  $\mu(X)$  by plotting it for four values of  $n$ . For each value of  $n$ , three values of  $D$  are considered: one less than, one equal to and one greater than  $1/(n+1)$ . Appendix I contains the programs used for generating the data, Appendix II contains the data generated and Appendix III

contains the plots obtained by using that data. In plots 2, 3 and 4 the scaling used for the Y axis is 1 unit = 2.0.

Looking at the plots in Appendix III, it is clear that the function is symmetric only when  $D = 1/(n+1)$  and is skewed in other cases. Also, even for the small values of  $n$  considered, the function tends to be a spike as  $n$  increases indicating that the variance decreases very rapidly.

## BIBLIOGRAPHY

1. BECK, D.D. (1968). "The Probability Distribution of the Measure of a Random Linear Set".  
M. Sc. thesis, Department of Mathematics, University of Alberta.
2. BRONOWSKI, J. and Neyman, J. (1945). "The Variance of Measure of Two-Dimensional Random Set".  
Annals of Mathematical Statistics, 16, pp. 330-341.
3. DOMB, C. (1947). "The Problem of Random Intervals on a Line".  
Proceedings of the Cambridge Philosophical Society, 43,  
pp. 329-341.
4. HALL, P. (1927). "The Distribution of Means for Samples of Size  $n$  Drawn from a Population in which the Variate Takes Values Between 0 and 1, All Such Values Being Equally Probable".  
Biometrika, 19, pp. 240-245.
5. OLVER, F.W.J. (1974). "Asymptotics and Special Functions".  
Academic Press, New York, pp. 333-335.
6. RIORDAN, J. (1968). "Combinatorial Identities".  
Wiley, New York, page 11.
7. ROBBINS, H.E. (1944). "On the Measure of a Random Set".  
Annals of Mathematical Statistics, 15, pp. 70-74.
8. ROBBINS, H.E. (1945). "On the Measure of a Random Set II".  
Annals of Mathematical Statistics, 16, pp. 342-347.
9. SANTALÓ, L.A. (1927). "On the First Two Moments of the Measure of a Random Set".  
Annals of Mathematical Statistics, 18, pp. 37-49.
10. STEVENS, W.L. (1939). "Solution to a Geometric Problem in Probability".  
Annals of Eugenics, 9, pp. 315-320
11. SZEGÖ, G. (1974). "Orthogonal Polynomials".  
American Mathematical Society, Colloquium Publications, Vol. 23,  
page 196.
12. VOTAW, D.F. (1946). "The Probability Distribution of the Measure of a Random Linear Set".  
Annals of Mathematical Statistics, 17, pp. 240-244.

**APPENDIX I**

## PROGRAM I

```

C ****
C THIS PROGRAM FINDS POINTS FOR THE PROBABILITY FUNCTION FNX. *
C GIVEN THE VALUE OF N+2 AND THREE VALUES OF D AS INPUT, IT *
C PRINTS THE VALUES OF FNX FOR EACH VALUE OF D. INCREMENT IN *
C X IS .02.
C ****
C
C
C
C ****
C
C SUBROUTINE COMB USES THE METHOD OF PASCAL'S TRIANGLE TO *
C CALCULATE N CHOOSE M FOR VARIOUS VALUES OF N AND M. A(I,J) *
C GIVES THE VALUE OF I-1 CHOOSE J-1 FOR J < OR = I.
C
C ****
C
C
C
C
SUBROUTINE COMB(N2,A)
DIMENSION A(N2,N2)
DO 5 I=1,N2
A(I,1)=1
A(I,I)=1
5 CONTINUE
N1=N2-1
DO 10 J=2,N1
J1=J+1
DO 10 I=J1,N2
A(I,J)=A(I-1,J)+A(I-1,J-1)
10 CONTINUE
RETURN
END
C
C
C ****
C
C GIVEN THE MATRIX OF COMBINATIONS, VALUE OF N+2 AND A VALUE *
C OF D, THIS SUBROUTINE CALCULATES AND PRINTS THE VALUES OF *
C THE PROBABILITY FUNCTION FNX. INCREMENT IN X IS .02. THE *

```

C LAST VALUE PRINTED IS (0.0,0.0).

C \*\*\*\*

C \*\*\*\*

C \*\*\*\*

C \*\*\*\*

SUBROUTINE POINTS(N2,D,A)

DIMENSION A(N2,N2)

REAL M1

INTEGER Q,R

B1=0

B2=0

M=INT(1/D)

N1=N2-1

N=N1-1

IF (N .LT. M) M=N

M1=N1\*D

IF (M1 .GT. 1) M1=1

DO 15 Q=1,M

X=Q\*D

X2=A MIN1(M1,(Q+1)\*D)

2 IF (X .LT. X2) GOTO 5

GOTO 15

5 FNX=0

DO 10 J=1,Q

JQ1=Q-J+1

DO 11 R=1,JQ1

S=A(N1,J+1)\*A(N2,J+1)\*A(N2-J,R)\*J

\* \*(-1)\*\*(R-1)\*(1-X)\*\*(J-1)

\* IF( ((X-D\*(J+R-1)) .EQ. (-.0)) .AND.

\* (N .EQ. J)) GOTO 10

S=S\*(X-D\*(J+R-1)) \*\*(N-J)

7 FNX=FNX+S

10 CONTINUE

100 WRITE(6,100) X,FNX

FORMAT(100,F12.8,5X,F12.8)

X=X+.02

GOTO 2

15 CONTINUE

WRITE(6,100) B1,B2

RETURN

```
C
C
C
C      END
C
C
C      MAIN PROGRAM STARTS HERE.  DIMENSION OF A IS TO BE CHANGED*
C      EVERY TIME N IS CHANGED.  RIGHT NOW N+2=21.          *
C
C
C
DIMENSION A(21,21)
READ(5,10) N,D1,D2,D3
10 FORMAT(I3,2X,3F4.4)
CALL COMB(N,A)
CALL POINTS(N,D1,A)
CALL POINTS(N,D2,A)
CALL POINTS(N,D3,A)
STOP
END
```

## **PROGRAM II**

```

C*****
C THIS PROGRAM PLOTS THE FUNCTION FNX. THE INPUT*
C IS VALUES OF X AND THE CORRESPONDING VALUES *
C OF THE FUNCTION FNX. DATA FOR A PARTICULAR *
C VALUE OF N AND THREE VALUES OF D IS GIVEN. *
C THE THREE SETS OF DATA ARE SEPARATED BY A *
C VALUE OF (0.0,0.0). SUBROUTINE SMOOTH IS USED*
C TO PLOT A SMOOTH CURVE FOR EACH SET OF DATA.*
C*****
C
C
C CALL PLOTS
CALL PLOT(4.0,4.0,-3)
CALL AXIS(0.0,0.0,'FNX',3,6.0,90.0,0.0,2.0)
CALL AXIS(0.0,0.0,'X',-1,6.0,0.0,0.0,0.2)
C
C
C X AND Y AXES HAVE BEEN PLOTTED. SCALING USED IS:
C X AXIS, 1 UNIT=.2; Y AXIS, 1 UNIT=2.0. LENGTH OF
C THE AXES IS 6 UNITS EACH.
C
C
100 FORMAT(F12.8,5X,F12.8)
DO 30 I=1,3
READ(5,100) X1,Y1
X1=X1*5
Y1=Y1/2.0
CALL SMOOTH(X1,Y1,0)
READ(5,100) X1,Y1
READ(5,100) X2,Y2
IF ((X2.EQ.0.0).AND.(Y2.EQ.0.0))
* GOTO 20
X1=X1*5
Y1=Y1/2.0
CALL SMOOTH(X1,Y1,-2)
X1=X2
Y1=Y2
GOTO 10
20 X1=X1*5
Y1=Y1/2.0

```

30        CALL SMOOTH(X1,Y1,-25)  
CONTINUE  
CALL PLOT(0.0,10.0,999)  
STOP  
END

**APPENDIX II**

$x$	$f_n(x)$
0.14999998	0.0
0.16999996	0.00016000
0.18999994	0.00128000
0.20999992	0.00431998
0.22999990	0.01023997
0.24999988	0.01999994
0.26999986	0.03455990
0.28999984	0.05487984
0.29999995	0.06749994
0.31999993	0.13025975
0.33999991	0.25877947
0.35999990	0.44441915
0.37999988	0.67853862
0.39999986	0.95249844
0.41999984	1.25765705
0.43999982	1.58537579
0.44999993	1.75499821
0.46999991	2.69945676
0.48999989	3.40977669
0.50999987	3.90491581
0.52999985	4.20683765
0.54999983	4.33499622
0.56999981	4.30955601
0.58999979	4.15067863
0.59999990	4.02749538
0.61999989	4.16465569
0.63999987	3.19318008
0.65999985	2.37961674
0.67999983	1.71053791
0.69999981	1.17250061
0.71999979	0.75206834
0.73999977	0.43578565

N+1 = 5

D = .15

POINTS OF THE PROBABILITY FUNCTION  $f_n(x)$

$x$	$f_n(x)$
0.19999999	0.0
0.21999997	0.00016000
0.23999995	0.00128000
0.25999993	0.00431998
0.27999991	0.01023997
0.29999989	0.01999994
0.31999987	0.03455990
0.33999985	0.05487984
0.35999984	0.08191973
0.37999982	0.11663961
0.39999980	0.15999949
0.39999998	0.15999991
0.41999996	0.24015969
0.43999994	0.37887949
0.45999992	0.56751919
0.47999990	0.79743874
0.49999988	1.05999756
0.51999986	1.34655762
0.53999984	1.64847660
0.55999982	1.95711613
0.57999980	2.26383495
0.59999979	2.55999565
0.59999996	2.55999756
0.61999995	3.12975597
0.63999993	3.50847626
0.65999991	3.71631718
0.67999989	3.77343559
0.69999987	3.69999790
0.71999985	3.51615906
0.73999983	3.24207973
0.75999981	2.89792061
0.77999979	2.50384140
0.79999977	2.07999992
0.79999995	2.08000183
0.81999993	1.63296223
0.83999991	1.14687920
0.85999990	0.76831901
0.87999988	0.48284267
0.89999986	0.28000164
0.91999984	0.14336455
0.93999982	0.06048144

continued ...

$x$	$f_n(x)$
0.95999980	0.01792191
0.97999978	0.00224317
0.99999976	0.00001091

$N+1 = 5$

$D = .2$

$x$	$f_n(x)$
0.25000000	0.0
0.26999998	0.00016000
0.28999996	0.00128000
0.30999994	0.00431998
0.32999992	0.01023997
0.34999990	0.01999994
0.36999989	0.03455990
0.38999987	0.05487984
0.40999985	0.08191973
0.42999983	0.11663961
0.44999981	0.15999949
0.46999979	0.21295935
0.48999977	0.27647948
0.50000000	0.31250000
0.51999998	0.41605973
0.53999996	0.57097948
0.55999994	0.76861930
0.57999992	1.00033855
0.59999990	1.25749779
0.61999989	1.53145790
0.63999987	1.81357670
0.65999985	2.09521580
0.67999983	2.36773586
0.69999981	2.62249565
0.71999979	2.85985583
0.73999977	3.04417610
0.75000000	3.12500000
0.76999998	3.34435558
0.78999996	3.41587639
0.80999994	3.35971737
0.82999992	3.1963729
0.84999990	2.94499588
0.86999989	2.62675953
0.88999987	2.26147938
0.90999985	1.86931992
0.92999983	1.47044277
0.94999981	1.08500481
0.96999979	0.73316389
0.98999977	0.43508625

 $N+1 = 5$  $D = .25$

$x$	$f_n(x)$
0.05000000	0.0
0.06999999	0.00000000
0.08999997	0.00000000
0.09999996	0.00000000
0.11999995	0.00000006
0.12999993	0.00000084
0.14999998	0.00000302
0.16999996	0.00002686
0.18999994	0.00021317
0.19999999	0.00056049
0.21999997	0.00308594
0.23999995	0.01525126
0.25001000	0.03141912
0.26999998	0.11402690
0.28999996	0.36735576
0.29999995	0.60745716
0.31999993	1.48858547
0.33999991	3.11749649
0.34999996	4.14516640
0.26999995	6.75558186
0.38999993	8.80857949
0.39999998	9.25242710
0.41999996	9.22718306
0.43999994	6.76822090
0.44999999	5.32125378
0.46999997	2.43540382
0.48999995	0.75956202

N+1 = 10

D = .05

POINTS OF THE PROBABILITY FUNCTION  $f_n(x)$

$x$	$f_n(x)$
0.10000002	0.0
0.12000000	0.00000000
0.13999999	0.00000000
0.15999997	0.00000002
0.17999995	0.00000015
0.19999993	0.00000090
0.20000005	0.00000090
0.22000003	0.00030387
0.24000001	0.0001368
0.25999999	0.70004523
0.27999997	0.00014674
0.29999995	0.00044910
0.30000007	0.00044910
0.32000005	0.00124937
0.34000003	0.00317647
0.36000001	0.00762981
0.38000002	0.01752147
0.39999998	0.03807797
0.40000010	0.03807894
0.42000008	0.27751429
0.44000006	0.14859414
0.46000004	0.27097487
0.48000002	0.46976060
0.50000002	0.77021760
0.50000012	0.77021956
0.52000010	1.19187737
0.54000008	1.74831486
0.56000006	2.43155861
0.58000004	3.19505310
0.60000002	3.95596313
0.60000014	3.95596504
0.62000012	4.61495018
0.64000010	5.06993961
0.66000009	5.22136688
0.68000007	5.02256393
0.70000005	4.49936199
0.70000017	4.49935627
0.72000015	3.74134350
0.74000013	2.86602783
0.76000011	2.00880432
0.78000009	1.27870274

continued ...

$x$	$f_n(x)$
0.80000007	0.73132712
0.80000019	0.73134053
0.82000017	0.37004709
0.84000015	0.16275913
0.86000013	0.06070851
0.88000011	0.01852956
0.90000010	0.00433540
0.90000021	0.00435212
0.92000020	0.00074362
0.94000018	0.00006932
0.96000016	0.00004155
0.98000014	0.00002963

N+1 = 10

D = .1

$x$	$f_n(x)$
0.14999998	0.0
0.16999996	0.000000200
0.18999994	0.000000000
0.20999992	0.00000002
0.22999990	0.00000015
0.24999988	0.00000090
0.26999986	0.000000387
0.28999984	0.00001328
0.29999985	0.00002307
0.31999993	0.0006278
0.33999991	0.00015320
0.35999990	0.00034607
0.37999988	0.00074556
0.39999986	0.00155958
0.41999984	0.00318037
0.43999982	0.00629526
0.44999993	0.00874200
0.46999991	0.01637739
0.48999989	0.02949681
0.50999987	0.05122401
0.52999985	0.08613634
0.54999983	0.14060110
0.56999981	0.22284114
0.58999979	0.34261190
0.59999997	0.41985470
0.61999989	0.61553764
0.63999987	0.87394774
0.65999985	1.20217991
0.67999983	1.60206604
0.69999981	2.06673431
0.71999979	2.57788563
0.73999977	3.10473633
0.74999988	3.36125374
0.76999986	3.83015823
0.78999984	4.19707966
0.80999982	4.41191196
0.82999980	4.43454361
0.84999979	4.24462891
0.86999977	3.84883213
0.88999975	3.28310966
0.89999986	2.95496559

continued ...

x	$f_n(x)$
.91999984	2.25514317
.93999982	1.56464767
.95999980	0.95775485
.97999978	0.48789287
.99999976	0.17744696

N+1 = 10

D = .15

$x$	$f_n(x)$
0.0470000	0.0
0.06699997	0.000000000
0.08699995	0.000000000
0.10399998	0.000000000
0.11399996	0.000000000
0.13399994	0.000000000
0.14499997	0.000000000
0.16099995	0.000000000
0.18099993	0.000000000
0.18799996	0.000000001
0.20799994	0.000000112
0.22799993	0.000000116
0.23499995	0.000000246
0.25499994	0.00001896
0.27499992	0.00012390
0.28199995	0.00023011
0.30199993	0.00122227
0.32199991	0.00558607
0.32899994	0.00918358
0.34899992	0.03464938
0.36899990	0.11329764
0.37599999	0.16598892
0.39599997	0.44957542
0.41599995	1.05748653
0.42299998	1.37983144
0.44299996	2.67675877
0.46299994	4.48677444
0.46999997	5.18875599
0.48999995	7.08026695
0.50999993	8.23807526
0.51699996	8.34980106
0.53699994	7.71578312
0.55699992	5.9233112
0.56399995	5.15394878
0.58399993	3.00089173
0.60399991	1.38145638
0.61099994	0.98982897
0.62099992	0.31209290
0.65099990	0.06821448
0.65799993	0.03635873
0.67799991	0.00498340

 $N+1 = 15$  $D = .047$ POINTS OF THE PROBABILITY FUNCTION  $f_n(x)$

$x$	$f_n(x)$
0.06699997	0.0
0.08699995	0.00000000
0.10699993	0.00000000
0.12699991	0.00000000
0.13399994	0.00000000
0.15399992	0.00000000
0.17399991	0.00000000
0.19399989	0.00000000
0.20099992	0.00000000
0.22099990	0.00000001
0.24099988	0.00000006
0.26099986	0.00000037
0.26799989	0.00000067
0.28799987	0.000001348
0.30799985	0.00001638
0.32799983	0.00006947
0.33499986	0.00011254
0.35499984	0.00042028
0.37499982	0.00143709
0.39499980	0.00450720
0.40199983	0.00657623
0.42199981	0.01838376
0.44199979	0.04732316
0.46199977	0.11221516
0.46899980	0.14892495
0.48899978	0.31654280
0.50899976	0.62017536
0.52999975	1.11945152
0.53599977	1.34994507
0.55599976	2.17830563
0.57599974	3.22858715
0.59599972	4.38687038
0.60299975	4.78222466
0.62299973	5.75334167
0.64299971	6.30015087
0.66299969	6.25321107
0.66999972	6.08820320
0.68999970	5.24847031
0.70999968	4.04568577
0.72999966	2.76617813
0.73699969	2.34974480

continued ...

	$f_n(x)$
0.75699967	1.34495831
0.77699965	0.66483343
0.79699963	0.27903068
0.80399966	0.19713759
0.82399964	0.06425452
0.84399962	0.01611321
0.86399961	0.00323460
0.87399963	0.00134316
0.89199962	0.00036085
0.91099960	0.00007201
0.93099958	0.00073744
0.93799961	0.00046743
0.95799959	0.00026522
0.97799957	0.00075879
0.99799955	0.00034385

N+1 = 15

D = .067

$x$	$f_n(x)$
0.08700001	0.0
0.10699999	0.000000700
0.12699997	0.000000200
0.14699996	0.000000000
0.16699994	0.000000000
0.17400002	0.000000000
0.19400001	0.000000000
0.21399999	0.000000000
0.23399997	0.000000000
0.25399995	0.000000702
0.26100004	0.000000203
0.28100002	0.000000015
0.30100000	0.00000064
0.32099998	0.000000261
0.34099996	0.000001972
0.34800005	0.00001511
0.36800003	0.00005183
0.38800001	0.00016022
0.40799999	0.00047358
0.42799997	0.00131346
0.43500006	0.00184985
0.45500004	0.00472699
0.47500002	0.01139560
0.49500000	0.02593213
0.51499999	0.05573017
0.52200007	0.07187277
0.54200006	0.14311713
0.56200004	0.26937126
0.58200002	0.47906375
0.60200000	0.80468741
0.61900009	0.95179145
0.62900007	1.47828360
0.64900005	2.16368580
0.66900003	2.97978401
0.68900001	3.85405540
0.69600010	4.15341187
0.71600008	4.91685200
0.73600006	5.43187237
0.75600004	5.57915497
0.77600002	5.30327129
0.78300011	5.11030674

continued. ...

x	$f_n(x)$
0.80300009	4.33485031
0.82300007	3.35044616
0.84300105	2.33553410
0.86300004	1.45004177
0.87000012	1.19043541
0.89000010	0.61420840
0.91000009	0.26759511
0.93000007	0.09427696
0.95000005	0.02484014
0.95700014	0.01443111
0.97000012	0.00188907
0.99700010	0.00010137

N+1 = 15

D = .087

$x$	$f_n(x)$
0.03000000	0.0
0.05000000	0.00000000
0.06000000	0.00000000
0.07999998	0.00000000
0.08999997	0.00000000
0.10999995	0.00000000
0.12000000	0.00000000
0.13999999	0.00000000
0.14999998	0.00000000
0.16999996	0.00000000
0.18000001	0.00000000
0.19999999	0.00000000
0.20999998	0.00000001
0.22999996	0.00000031
0.24000001	0.00000134
0.25999999	0.00002022
0.26999998	0.00007120
0.28999996	0.00072930
0.30000001	0.00212499
0.31999999	0.01499309
0.32999998	0.03632654
0.34999996	0.17751634
0.36000001	0.35787213
0.38000000	1.20739174
0.38999999	2.01778603
0.40999997	4.64319311
0.42000002	6.37708350
0.44000000	9.77926826
0.44999999	10.87393570
0.46999997	10.63343281
0.48000002	9.39288330
0.50000000	5.56202221
0.50999999	3.70538807
0.52999997	1.17961025
0.54000002	0.55092895
0.56000000	0.07719499
0.56999999	0.01934515
0.58999997	0.02313974

continued ...

$x$	$f_n(x)$
0.73999995	1.39158344
0.75000000	0.93759954
0.76999998	0.36744744
0.78999996	0.11698687
0.80000001	0.05807549
0.81999999	0.01151742
0.83999997	0.00000341
0.84999996	0.000062970
0.86999995	0.00609926
0.88999993	0.00370306
0.89999998	0.00293845
0.91999996	0.00327434
0.93999994	0.00136712
0.94999999	0.10408846
0.96999997	0.00822474
0.98999995	0.00132161

 $N+1 = 20$  $D = .03$ POINTS OF THE PROBABILITY FUNCTION  $f_n(x)$

$x$	$f_n(x)$
0.05000000	0.0
0.06999999	0.00000000
0.08999997	0.00000000
0.09999996	0.00000000
0.11999995	0.00000000
0.13999993	0.00000000
0.14999998	0.00000000
0.16999996	0.00000000
0.18999994	0.00000000
0.19999999	0.00000000
0.21999997	0.00000000
0.23999995	0.00000000
0.25000000	0.00000000
0.26999998	0.00000001
0.28999996	0.00000007
0.29999995	0.00000019
0.31999993	0.00000131
0.33999991	0.00000819
0.34999996	0.00001954
0.36999995	0.00010148
0.38999993	0.00046891
0.39999998	0.00096560
0.41999996	0.00376274
0.43999994	0.01311893
0.44999999	0.02357425
0.46999997	0.06948674
0.48999995	0.18411052
0.50000000	0.28759397
0.51999998	0.64599752
0.53999996	1.29794502
0.54999995	1.76352024
0.56999993	2.98703003
0.58999991	4.50006390
0.59999996	5.28046227
0.61999995	6.62814331
0.63999993	7.32459641
0.64999998	7.32770157
0.66999996	6.61493206
0.68999994	5.16995907
0.69999999	4.31777668
0.71999997	2.67052364

 $N+1 = 20$  $D = .05$

$x$	$f_n(x)$
0.06999999	0.0
0.08999997	0.00070000
0.10999995	0.00000000
0.12999994	0.00000000
0.13999999	0.00001000
0.15999997	0.00000000
0.17999995	0.00000000
0.19999993	0.00000000
0.20999998	0.00000000
0.22999996	0.00000000
0.24999994	0.00000000
0.26999992	0.00000000
0.27999997	0.00000000
0.29999995	0.00000000
0.31999993	0.00000002
0.33999991	0.00000009
0.34999996	0.00000019
0.36999995	0.00000093
0.38999993	0.00070408
0.40999991	0.00001658
0.41999996	0.00003246
0.43999994	0.00011759
0.45999992	0.00039573
0.47999990	0.00123911
0.48999995	0.00213511
0.50999993	0.00601482
0.52999991	0.01580729
0.54999989	0.03877158
0.55999994	0.05917258
0.57999992	0.13087237
0.59999990	0.27005053
0.61999989	0.51951927
0.62999994	0.70165992
0.64999992	1.21254730
0.66999990	1.94695377
0.68999988	2.89853191
0.69999993	3.43502045
0.71999991	4.54229927
0.73999989	5.52599525
0.75999987	6.15758801
0.76999992	6.27902699

continued ...


$x$	$f_n(x)$
0.78999990	6.06845284
0.80999988	5.28544044
0.82999986	4.10895729
0.83999991	3.45872498
0.85999990	2.21260929
0.87999988	1.21485519
0.89999986	0.55688423
0.90999991	0.34862763
0.92999989	0.11256355
0.94999987	0.02570609
0.96999985	0.00372719
0.97999990	0.00123571
0.99999988	0.00010629

 $N+1 = 20$  $D = .07$

**APPENDIX III**


PROBABILITY FUNCTION  $f_n(x)$  OF THE MEASURE  
OF A RANDOM CIRCULAR SET

PLOT 1




PROBABILITY FUNCTION  $f_n(x)$  OF THE MEASURE  
OF A RANDOM CIRCULAR SET

PLOT 2



PROBABILITY FUNCTION  $f_n(x)$  OF THE MEASURE  
OF A RANDOM CIRCULAR SET

PLOT 3



PROBABILITY FUNCTION  $f_n(x)$  OF THE MEASURE  
OF A RANDOM CIRCULAR SET

PLOT 4

