

University of Alberta

Effect of Beta-Amyloid Protein on Nicotinic Acetylcholine Receptor Function in
Diagonal Band of Broca Neurons

by

James Howe Chin



A thesis submitted to the Faculty of Graduate Studies and Research in partial
fulfillment of the requirements for the degree of Doctor of Philosophy

Centre for Neuroscience

Edmonton, Alberta

Spring 2006



Library and
Archives Canada

Bibliothèque et
Archives Canada

Published Heritage
Branch

Direction du
Patrimoine de l'édition

395 Wellington Street
Ottawa ON K1A 0N4
Canada

395, rue Wellington
Ottawa ON K1A 0N4
Canada

Your file *Votre référence*
ISBN: 0-494-13951-X
Our file *Notre référence*
ISBN: 0-494-13951-X

NOTICE:

The author has granted a non-exclusive license allowing Library and Archives Canada to reproduce, publish, archive, preserve, conserve, communicate to the public by telecommunication or on the Internet, loan, distribute and sell theses worldwide, for commercial or non-commercial purposes, in microform, paper, electronic and/or any other formats.

The author retains copyright ownership and moral rights in this thesis. Neither the thesis nor substantial extracts from it may be printed or otherwise reproduced without the author's permission.

AVIS:

L'auteur a accordé une licence non exclusive permettant à la Bibliothèque et Archives Canada de reproduire, publier, archiver, sauvegarder, conserver, transmettre au public par télécommunication ou par l'Internet, prêter, distribuer et vendre des thèses partout dans le monde, à des fins commerciales ou autres, sur support microforme, papier, électronique et/ou autres formats.

L'auteur conserve la propriété du droit d'auteur et des droits moraux qui protègent cette thèse. Ni la thèse ni des extraits substantiels de celle-ci ne doivent être imprimés ou autrement reproduits sans son autorisation.

In compliance with the Canadian Privacy Act some supporting forms may have been removed from this thesis.

Conformément à la loi canadienne sur la protection de la vie privée, quelques formulaires secondaires ont été enlevés de cette thèse.

While these forms may be included in the document page count, their removal does not represent any loss of content from the thesis.

Bien que ces formulaires aient inclus dans la pagination, il n'y aura aucun contenu manquant.


Canada

ABSTRACT

β -amyloid ($A\beta$), a 39-43 amino acid peptide deposited in the brains of Alzheimer's patients, disrupts synaptic function through mechanisms that are not fully understood. Nicotinic acetylcholine receptors (nAChRs) are potential targets for the actions of $A\beta$ in the brain. This thesis examines 1) the role of nAChRs on basal forebrain neurons in Ca^{2+} signaling and excitatory neurotransmission and 2) if $A\beta$ influences these functions via nAChRs.

Intracellular calcium levels ($[Ca^{2+}]_i$) in acutely dissociated rat basal forebrain neurons were measured with the ratiometric dye fura-2. Focal applications of nicotine, evoked increases of $[Ca^{2+}]_i$ mediated via Ca^{2+} entry via nAChRs and intracellular Ca^{2+} release. Nicotine-evoked $[Ca^{2+}]_i$ rises were inhibited by α -bungarotoxin, and dihydro-beta-erythroidine (DH β E). Nicotine-evoked increases were irreversibly potentiated by $A\beta_{1-42}$, while focal application of $A\beta_{1-42}$ alone did not alter $[Ca^{2+}]_i$. $A\beta_{1-42}$ also potentiated caffeine-mediated, but not KCl-evoked rises of $[Ca^{2+}]_i$. $A\beta_{1-42}$ potentiation of nicotine-mediated rises of $[Ca^{2+}]_i$ was blocked by either the SERCA inhibitor thapsigargin or CGP-37157, an inhibitor of the mitochondrial Na^+/Ca^{2+} exchanger. Thus $A\beta_{1-42}$ potentiation of repeated release of Ca^{2+} from intracellular stores by multiple nicotine challenges is due to overfilling of caffeine sensitive stores by mitochondria.

The effects of $A\beta_{1-42}$, nicotine and acetylcholine (ACh) on cholinergic basal forebrain neurons prelabeled with Cy3-IgG192 or identified by electrophysiological criteria were examined by recording miniature excitatory synaptic currents (mEPSCs) using whole-cell patch clamp recordings in brain slice preparations. In 54% of neurons, $A\beta_{1-42}$ significantly increased mEPSC frequency while in 32% of neurons, $A\beta_{1-42}$

significantly decreased mEPSC frequency. DH β E blocked A β_{1-42} mediated increases of mEPSC frequency, but did not block A β_{1-42} mediated decreases of mEPSC frequency. Nicotine or ACh in the presence of atropine, increased the frequency of mEPSCs in a dose-dependent manner. No changes in peak amplitude were observed in any treatments. These data suggest that in a subset of cholinergic neurons, A β_{1-42} increases mEPSC frequency by activating pre-synaptic $\alpha 4\beta 2$ nAChRs, while another unknown pre-synaptic mechanism mediated the decrease of mEPSC frequency. As the regulation of $[Ca^{2+}]_i$ and synaptic transmission are important for cognitive functioning, these results may be important to understanding the pathophysiological mechanisms underlying Alzheimer's disease.

Acknowledgements

Research is not conducted in a vacuum and many people have enabled me to complete this thesis. First and foremost, I would like to thank Dr. Jack Jhamandas, who has guided me through both joyful and stressful times during my time in his laboratory. He has always provided patient guidance and encouragement for me and I consider him a friend, a great role model and a colleague. I will not forget his generous spirit in creating a supportive environment and providing me with many so many opportunities.

I would also like to thank the members of my supervisory committee for providing guidance. Their constructive criticism was appreciated and always opened my eyes to new avenues that I hadn't seen before. I would like to thank Dr. Fred Tse for always being generous with his time whenever I came around to talk about ideas and for always pointing me in a direction I hadn't seen. Dr. Bill Colmers has always been generous with his ideas by always making suggestions as to how I could improve things and in showing me the finer points of electrophysiology.

I would like to acknowledge Dr. John Greer and Dr. Satya Kar and Dr. Jeff Goldberg for serving as external examiners on my candidacy exam. Also I would like to thank Dr. Staya Kar, Dr. Declan Ali, and Dr. Brian MacVicar for serving on my thesis defense committee.

I would like to extend my thanks to my colleagues in the lab: Kim Harris, David MacTavish, Wen Fu, Monica Ding, Li Ma and Zhong Ming. I appreciate all the help that Kim has given me. He has always been generous with his time and has helped me time and again with the technical aspects of performing experiments. His technical skills are second to none whether it comes to experiments, gardening, painting or basketball. Dave

has been fantastic in teaching me immunohistochemistry and introducing me to running. Wen has always been an excellent source of information and great source for suggestions when things aren't going well. I would like to thank Monica for being a great colleague and for providing me with practical tips on how to complete this thesis and Li Ma, who despite being relatively new to the lab has shared with me many tips on how to improve my recordings. I would like to thank Ms. Cynthia Krys for helping me with paperwork and coordinating everyone's schedule for committee meetings. Additionally, I would like to thank Ms. Nina Pronchuk, one of the kindest people I have ever met, for always sharing her knowledge of electrophysiological techniques with me.

To Dad, Mom and Jen, thank you for always supporting me and encouraging me in my pursuit of knowledge. Without their support and encouragement I could not be where I am today. I know that they are proud of me and I dedicate this thesis to them.

To Haili Wang, the love of my life, I thank her for being there and supporting me throughout graduate school. In the past 4 years she has always been there for me and I feel fortunate to have such a wonderful relationship with such a special person.

Table of Contents

Chapter 1

General Introduction	1
Overview	2
Anatomy of the Basal Forebrain Cholinergic System	3
Neurochemical Characteristics of the DBB.....	4
Acetylcholine.....	4
GABA.....	5
Glutamate.....	6
Other Chemical Phenotypes within the DBB.....	6
Afferent and Efferent Connections and their Relation to Function.....	7
Efferent Connections of the DBB.....	7
Afferent Connections to the DBB.....	9
Physiological Functions of the DBB	10
Theta Rhythm.....	10
Learning and Memory.....	11
DBB Neuronal Loss in AD.....	12
Cellular Properties of DBB Neurons	13
Intrinsic Electrophysiological Properties.....	13
Receptors Expressed on DBB Neurons.....	14
Intracellular Ca ²⁺ Stores.....	15
Nicotinic Acetylcholine Receptors in the Central Nervous System	16
General Properties.....	16

NAChRs in the DBB.....	18
Physiological Functions of nAChRs in the Central Nervous System.....	19
Modulation of Transmitter Release.....	19
NAChRs and Second Messengers.....	20
Age-Dependent Alteration of nAChRs.....	20
NAChRs and AD.....	21
General Properties of Amyloid.....	22
A β Production.....	23
Conformation of A β	24
Effects of A β on Neurons.....	25
A β and nAChRs.....	26
Rational and Objectives.....	29
References.....	32
Chapter 2	
β-Amyloid Enhances Intracellular Calcium Rises Mediated by Intracellular Calcium Stores and Nicotinic Receptors in Acutely Dissociated Rat Basal Forebrain Neurons.....	60
Introduction.....	61
Materials and Methods.....	63
Dissociation Procedures.....	63
Calcium Imaging.....	64
Immunohistochemistry.....	65
Drugs.....	66

Statistical Analysis.....	66
Results.....	66
Nicotine-Evoked Ca ²⁺ Responses in DBB Neurons: Modulation by Aβ.....	66
Aβ Potentiation of nAChR-Induced Elevation of [Ca ²⁺] _i is Not Mediated by Voltage Gated Ca ²⁺ Channels.....	68
Aβ Potentiates Release of Ca ²⁺ from Intracellular Ca ²⁺ Stores.....	68
Pharmacology of Non-α7 and α7 nAChR Receptors on DBB Neurons.....	71
Discussion.....	72
Activation of nAChRs on Basal Forebrain Neurons.....	73
NACHR Coupling to Intracellular Ca ²⁺ Stores.....	74
Aβ and [Ca ²⁺] _i Homeostasis.....	75
Functional Implications.....	77
References.....	105
 Chapter 3	
Synaptic Effects of β-Amyloid on Glutamate Neurotransmission in the Rat	
Cholinergic Basal Forebrain.....	112
Introduction.....	113
Materials and Methods.....	116
Cy3-192 IgG Neuron Labeling.....	116
Diagonal Band of Broca Slice Preparation.....	116
Recordings from DBB Slices.....	117
Statistical Analysis.....	118
Immunohistochemistry.....	119

Drugs.....	120
Results.....	120
Cy3-192 IgG Labeling Colocalizes with vChAT Labeling in DBB Neurons.....	121
CBQX Abolishes mEPSCs in DBB Neurons.....	121
A β Modulates Frequency of mEPSCs in Cholinergic DBB Neurons.....	121
A β Mediated Increase in mEPSC Frequency is Inhibited by DH β E.....	123
A β Mediated Decrease in mEPSC Frequency Persists in the Presence of DH β E.....	124
Nicotine Increases mEPSC Frequency in a Dose-Dependent Manner.....	125
Acetylcholine Increases mEPSC Frequency in a Dose-Dependent Manner.....	125
Discussion.....	126
Soluble A β Modulates Synaptic Transmission onto Cholinergic DBB Neurons.....	127
Activation of nAChRs Modulate Synaptic Transmission onto Cholinergic DBB Neurons.....	128
Future Experiments.....	129
Functional Implications.....	129
References.....	164
Chapter 4	
General Discussion.....	169
Aβ Potentiates Repeated nAChR Mediated [Ca²⁺]_i Rises.....	170
Aβ Modulates Synaptic Transmission in Cholinergic DBB Neurons.....	174
Future Experiments and Conclusions.....	175

References.....181

List of Tables

Table 3-1. Kinetic Parameters of Cholinergic Neurons which Demonstrate an Increase of mEPSCs in the Presence of 100 nM A β ₁₋₄₂	131
Table 3-2. Kinetic Parameters of Cholinergic Neurons which Demonstrate a Decrease of mEPSCs in the presence of 100 nM A β ₁₋₄₂	131
Table 3-3. Kinetic Parameters of Cholinergic Neurons which Demonstrate an Increase in the Frequency of mEPSCs in the Presence of 100 nM A β ₁₋₄₂ that are Subsequently Exposed to 100 nM A β ₁₋₄₂ and 10 μ M DH β E.....	131
Table 3-4. Kinetic parameters of Cholinergic Neurons that Demonstrate a Decrease in the Frequency of mEPSCs in the presence of 100 nM A β ₁₋₄₂ that are Subsequently Exposed to 100 nM A β ₁₋₄₂ and 10 μ M DH β E.....	131
Table 3-5. Kinetic Parameters of Cholinergic Neurons that Demonstrate an Increase in the Frequency of mEPSCs in response to Bath Application of 100 μ M Nicotine.....	132
Table 3-6. Kinetic Parameters of Cholinergic Neurons Demonstrating an Increase of mEPSC Frequency in Response to Bath Application of 10 μ M Nicotine.....	132
Table 3-7. Kinetic Parameters of Cholinergic Neurons Demonstrating an Increase of mEPSC Frequency in Response to Bath Application of 1 μ M Nicotine.....	132
Table 3-8. Kinetic Parameters of Cholinergic Neurons Demonstrating no Change of mEPSC Frequency in Response to Bath Application of 100 μ M Nicotine.....	132

Table 3-9. Kinetic Parameters of Cholinergic Neurons in the Presence of 1 mM ACh.....	132
Table 3-10. Kinetic parameters of Cholinergic Neurons in the Presence of 100 μ M ACh.....	133
Table 3-11. Kinetic Parameters of Cholinergic Neurons in the Presence of 1 μ M ACh.....	133

List of Figures

Figure 1-1. Proteolytic Processing of Amyloid Precursor Protein.....	31
Figure 2-1. Changes in $[Ca^{2+}]_i$ in Response to Focal Application of Nicotine.....	80
Figure 2-2. Nicotine-Evoked Rises in $[Ca^{2+}]_i$ are Not Mediated by Changes in Osmolarity or Cross reaction with Muscarinic Receptors.....	82
Figure 2-3. Effect of $A\beta_{1-42}$ on Nicotine-Evoked $[Ca^{2+}]_i$ Responses in Acutely Dissociated DBB Cells.....	84
Figure 2-4. Dose-Response Curve for $A\beta_{1-42}$ Potentiation of 10 mM Nicotine-Evoked $[Ca^{2+}]_i$ Fitted to a Sigmoidal Dose-Response Curve.....	86
Figure 2-5. Effect of Various Concentrations of $A\beta_{1-42}$ on DBB Neurons.....	88
Figure 2-6. Effect of $A\beta$ on KCl Induced Depolarization in Acutely Dissociated DBB Neurons.....	90
Figure 2-7. Contribution of Intracellular Ca^{2+} Stores to Nicotine-Mediated Rise in $[Ca^{2+}]_i$	92
Figure 2-8. Effect of PLC Blockade on Nicotine-Evoked $[Ca^{2+}]_i$ Rises.....	94
Figure 2-9. Effect of 5 s Focal Application 10 mM Caffeine on DBB Neurons.....	96
Figure 2-10. Effect of Antagonists of Intracellular Stores on $A\beta$ Potentiation of Nicotinic responses.....	98
Figure 2-11. Pharmacology of Nicotine-Mediated $[Ca^{2+}]_i$ Rises in DBB Neurons I....	100
Figure 2-12. Pharmacology of Nicotine-Mediated $[Ca^{2+}]_i$ Rises in DBB Neurons II...	102
Figure 2-13. Confocal Microscopy of Double Labeled Acutely Dissociated DBB Neurons Reveals $\alpha 4$ and $\alpha 7$ nAChR Subunits Present on Cholinergic Neurons.....	104

Figure 3-1. Sample Traces of Current Clamp Identification of Cholinergic Neurons, Voltage Ramp and mEPSC.....	135
Figure 3-2. The Fluorescent Retrograde tracer Cy3-192 IgG Labels vChAT Positive Cells in the DBB.....	137
Figure 3-3. mEPSC Activity in Cholinergic Neurons is Abolished by CNQX.....	139
Figure 3-4. Bath Application of 100 nM A β ₁₋₄₂ Increases mEPSC Frequency in 54% of Cholinergic DBB Neurons I.....	141
Figure 3-5. Bath Application of 100 nM A β ₁₋₄₂ Increases mEPSC Frequency in 54% of Cholinergic DBB Neurons II.....	143
Figure 3-6. Bath Application of 100 nM A β ₁₋₄₂ Decreased mEPSC Frequency in 32% of Cholinergic Neurons I.....	145
Figure 3-7. Bath Application of 100 nM A β ₁₋₄₂ Decreased mEPSC Frequency in 32% of Cholinergic Neurons II.....	147
Figure 3-8. Bath Application of the Reverse Peptide 100 nM A β ₄₂₋₁ did not Alter Frequency or Amplitude when Applied to Cholinergic Neurons, which Responded to 100 nM A β ₁₋₄₂ with an Increased mEPSC Frequency I.....	149
Figure 3-9. Bath Application of the Reverse Peptide 100 nM A β ₄₂₋₁ did not Alter Frequency or Amplitude when Applied to Cholinergic Neurons, which Responded to 100 nM A β ₁₋₄₂ with an Increased mEPSC Frequency II.....	151
Figure 3-10. The A β Mediated Increase of mEPSC Frequency in Cholinergic Neurons is blocked by 10 μ M of the non- α 7 nAChR Antagonist DH β E.....	153
Figure 3-11. Sample Cumulative Frequency Histograms Demonstrating that A β ₁₋₄₂ Mediated Increase of mEPSC Frequency in Cholinergic Neurons	

is Blocked by 10 μ M of the non- α 7 nAChR Antagonist DH β E.....	155
Figure 3-12. A β ₁₋₄₂ Mediated Decreases of mEPSC Frequency in Cholinergic Neurons Persist in the Presence of 10 μ M DH β E.....	157
Figure 3-13. Sample Cumulative Frequency Histograms Showing that A β ₁₋₄₂ Mediated Decrease of mEPSC Frequency in a Cholinergic Neurons is not Blocked by DH β E.....	159
Figure 3-14. Nicotine Increases mEPSC Frequency of Cholinergic DBB Neurons.....	161
Figure 3-15. ACh, the Endogenous Neurotransmitter for nAChRs Increases the Frequency of mEPSCs in Cholinergic DBB Neurons.....	163
Figure 4-1. Effect of A β ₁₋₄₂ on Intracellular Ca ²⁺ Homeostasis in DBB Neurons.....	178
Figure 4-2. A β ₁₋₄₂ Increases or Decreases Glutamatergic Transmission in Cholinergic DBB Neurons.....	180

Abbreviations

A β	β -amyloid peptide
ACh	acetylcholine
AChE	acetylcholinesterase
ACSF	artificial cerebrospinal fluid
AD	Alzheimer's Disease
AHP	afterhyperpolarization
AMPA	α -amino-3-hydroxy-5-methyl-4-isoxazolepropionate
APP	amyloid precursor protein
APV	(-) 2-amino-7-phosphonopentanoate
BHQ	di-(tert-butyl)-1,4-benzohydroquinone
α BTX	α -bungarotoxin
Ca ²⁺	calcium
[Ca ²⁺] _i	intracellular concentration of calcium
CGP-37157	7-chloro-5-(2-chlorophenyl)-1,5-dihydro-4,1-benzothiazepin-2(3H)-one
ChAT	choline acetyltransferase
CNQX	cyano-7-nitroquinoxaline-2,3-dione
CREB	cAMP regulatory element binding protein
DBB	diagonal band of Broca
DH β E	dihydro-beta-erythroidine
EPSC	excitatory post-synaptic current
ER	endoplasmic reticulum
FAHP	fast afterhyperpolarization

GABA	γ -aminobutyric acid
InsR	inositol 1,4,5-phosphate receptor
IP ₃	inositol 1,4,5-triphosphate
KS	Kolmogorov-Smirnov
LTP	long-term potentiation
NMDA	<i>N</i> -Methyl-D-Aspartic Acid
NO	nitric oxide
MAP2	mitogen activation protein kinase
mEPSCs	miniature spontaneous excitatory postsynaptic currents
PKC	protein kinase C
RyR	ryanodine receptor
SERCA	sarco-ER Ca ²⁺ ATPase
SAHP	slow afterhyperpolarization
TG	thapsigargin
VGCC	voltage-gated calcium channel

General Introduction

Chapter 1

Overview

Alzheimer's disease (AD) is a devastating neurodegenerative disease that affects elderly individuals and is characterized by a progressive decline in memory and other cognitive functions such as language and perception (McKhann et al., 1984). The majority of AD occurs in a sporadic form, with an average age of onset of 70 years of age. Apolipoprotein E gene, which regulates the transport of cholesterol, has been identified as a susceptibility gene for the common late-onset form of AD (Poirier et al., 1993; Saunders et al., 1993). In a small subset of AD individuals, the onset of symptoms begin at approximately 50 years of age and these cases are associated with familial autosomal dominant mutations in transmembrane proteins regulated by genes located on chromosomes 21 (Amyloid Precursor Protein, (APP), 14 (Presenilin 1) and 1 (Presenilin 2)(Goate et al., 1991; Levylahad et al., 1995; Sherrington et al., 1995). Key neuropathological findings in individuals with AD include extracellular and neuritic (intracellular) plaques composed of β -amyloid peptide ($A\beta$), neurofibrillary tangles composed of abnormally phosphorylated tau protein, and selective loss of acetylcholine (ACh) synthesizing (cholinergic) neurons of the basal forebrain (Yankner, 1996; Mayeux and Sano, 1999). The loss of cholinergic neurons, which are one of the most vulnerable neuronal populations in AD, is associated with severe memory and learning deficits, which are the hallmarks of the clinical symptom complex of AD. The role of $A\beta$, a 39-43 amino acid peptide) in the pathogenesis of AD is controversial, i.e. whether it is a cause or consequence of the disease. However a large body of experimental evidence indicates that it may be a critical determinant of synaptic dysfunction and cell death that has been observed in many animal models of AD and the human condition (Hardy and

Allsop, 1991; Selkoe, 2002b). For example, A β has also been demonstrated to be neurotoxic to rat hippocampal and human cortical neurons (Mattson et al., 1993). Despite its wide-ranging effects, however, no definitive “receptor” has been identified for A β . Several candidate receptors have been proposed to mediate the biological actions of A β , including plasma membrane receptors for advanced glycation end products (RAGE), tumor necrosis factor receptor, and the p75 neurotrophin receptor (Rabizadeh et al., 1994; Barger et al., 1995; Yan et al., 1996). A β has been demonstrated to bind to nicotinic acetylcholine receptors (nAChRs) in nano- and picomolar concentrations and these receptors are important for mediating many cellular processes such as calcium (Ca²⁺) signaling and neuromodulation. However, the interactions of A β with nAChRs are only beginning to be investigated and have not been studied in the basal forebrain, a region which is at the epicenter of the chemical pathology observed in AD (Castro and Albuquerque, 1995; Wang et al., 2000b; Wang et al., 2000a). **Therefore, the overall objective of my thesis was to 1) examine the role of nAChRs on basal forebrain neurons in Ca²⁺ signaling and neuromodulation 2) examine how A β influences nAChR mediated neuromodulation and Ca²⁺ signaling 3) examine the effects of A β on nAChR modulation of excitatory synaptic transmission in a cholinergic basal forebrain nucleus.**

I. Anatomy of the Basal Forebrain Cholinergic System

The basal forebrain nuclei are the major sources of cholinergic innervation to the brain in rats and humans (Mesulam et al., 1983a; Saper and Chelimsky, 1984). These nuclei are located ventral to the anterior horn of the lateral ventricle and the striatum and extend from the base of where the cerebral hemispheres unite and extend to the cerebral

peduncle. In coronal sections of rat brain, the anterior portion of the basal forebrain resembles an inverted “Y”. The dorsal stalk of the Y is composed of the medial septum and the vertical limb of the diagonal band of Broca (DBB), while the horizontal arms of the Y comprise the horizontal limbs of the DBB (Harkmark et al., 1975). The most posterior sector of the basal forebrain is the nucleus basalis of Meynert, which can be further subdivided into anterior, intermediate, and posterior segments (Mesulam et al., 1983b; Mesulam et al., 1983a).

A. Neurochemical Characteristics of the DBB

DBB neurons are heterogeneous and contain many different neurotransmitters and neuropeptides. The two main populations of neurons in the DBB are the cholinergic and γ -aminobutyric acid (GABA) synthesizing cells (Brashear et al., 1986). There is also a local population of glutamatergic neurons (Sotty et al., 2003; Manseau et al., 2005). The cholinergic and GABAergic neurons can be divided into a population of large and a population of small diameter neurons. The large neurons are believed to be projection neurons, while the small neurons release neurotransmitters locally within the nucleus (interneurons).

1. Acetylcholine

Within the DBB, it is estimated that between 34-45% of neurons are cholinergic using staining methods for detecting either choline acetyltransferase (ChAT), which is an enzyme involved in ACh synthesis, or acetylcholinesterase, an enzyme that breaks down ACh in the synaptic cleft (Eckenstein and Sofroniew, 1983; Mesulam et al., 1983a). The morphology of these ChAT immunoreactive neurons can be divided into two types based upon their cell diameter (Milner, 1991). The first type is the magnocellular neurons,

which are 20-30 μM in diameter and are either elongated or round. They have an abundant cytoplasm with a small indented nucleus and may contain lamellar bodies and are the principal projection neurons of the DBB (Mesulam and Van Hoesen, 1976). The second population of neurons is the parvocellular (15-20 μM) neurons, which are round and do not have lamellar bodies. These neurons are responsible for local release of ACh (Metcalf et al., 1988).

The magnocellular neurons of the DBB provide cholinergic innervation to many CNS regions (Mesulam and Van Hoesen, 1976). Cholinergic DBB neurons project to the olfactory bulb, the hippocampus and to many areas of the neocortex (Mesulam et al., 1983a). They play an important role in a number of physiological functions including memory (Waite et al., 1994) and theta rhythm (Gaztelu and Buno, 1982) which shall be discussed more fully in later sections. In AD, it is the cholinergic projection neurons of the basal forebrain that preferentially undergo neurodegeneration (Whitehouse et al., 1982).

2. *GABA*

Immunohistochemical labeling for GABA and the GABA synthesizing enzyme L-glutamate decarboxylase in the basal forebrain identified a population of GABAergic neurons (Panula et al., 1984; Onteniente et al., 1986). This population of neurons was estimated to comprise between 34-54% of the neuronal population in the basal forebrain (Griffith, 1988; Kiss et al., 1990). The morphology of GABAergic neurons in the DBB is multiform and variable (Onteniente et al., 1986). When compared to other GABAergic neurons of the basal forebrain, GABAergic neurons of the DBB are larger, with a 20 to 30 μm diameter versus 10-20 μm in the lateral septum (Onteniente et al., 1986). The

aforementioned observations of the DBB GABAergic neurons suggest that this population of neurons is distinct from other neurons in the basal forebrain and may consist of mainly projection neurons (Onteniente et al., 1986). Based on anterograde and retrograde tracer studies, GABAergic neurons of the DBB have been identified to project to the neocortex and the hippocampus (Freund, 1989; Freund and Gulyas, 1991).

3. *Glutamate*

Recently, a population of glutamate releasing neurons was identified in the basal forebrain (Gritti et al., 1997; Manns et al., 2001). Based upon retrograde tracer analysis and single cell reverse transcription polymerase reaction for vesicular glutamate transporters, glutamatergic neurons are medium in size with an average diameter of 17 μm and consist of neurons that project to the neocortex and the entorhinal cortex (Sotty et al., 2003; Wu et al., 2003). There are also local glutamatergic interneurons that synapse onto both GABAergic and cholinergic neurons (Wu et al., 2003; Manseau et al., 2005).

4. *Other Chemical Phenotypes within the DBB*

Neurons in the DBB express other neuromodulatory compounds including galanin, nerve growth factor (NGF), nitric oxide (NO), and vasopressin. Galanin is a polypeptide which is co-localized in a subset of cholinergic neurons of the DBB that project to the hippocampus (Melander et al., 1985; Skofitsch and Jacobowitz, 1985; Senut et al., 1989). It has been shown that galanin inhibits the release of ACh onto hippocampal neurons via a presynaptic mechanism (Fisone et al., 1987). However, electrophysiological data show galanin to inhibit a suite of potassium conductances on cholinergic basal forebrain neurons resulting in hyperexcitability of these cells (Jhamandas et al., 2002). NO synthase, an enzyme responsible for the synthesis of NO

has been found to colocalize with the ChAT and nerve growth factor receptor, however, further studies need to be done to examine the role of NO role in the DBB (Kitchener and Diamond, 1993).

B. Afferent and Efferent Connections and their Relation to Function

The DBB has many afferent and efferent connections in the brain, which are important for theta rhythm, arousal, learning and memory. The following sections will be divided into a section outlining the anatomical connectivity of the DBB and another describing the role of DBB in theta rhythm, learning and memory.

1. Efferent Connections of the DBB

Projections from the DBB extend to many areas of the brain. Anatomical studies utilizing anterograde and retrograde tracers have shown that the DBB projects to the hippocampus, olfactory bulb, neocortex, hypothalamus, amygdala, dorsal raphe, and locus coeruleus (Divac, 1975; Conrad and Pfaff, 1976a, 1976b).

Both cholinergic and GABAergic DBB neurons project to the hippocampus, olfactory bulb and neocortex and these projections have been studied in great detail (Mesulam et al., 1983a; Senut et al., 1989; Kiss et al., 1990; Freund and Gulyas, 1991).

The neurons projecting to the hippocampus reside in the vertical and horizontal limbs of the DBB, while neurons projecting to the olfactory bulb and the neocortex mainly reside in the horizontal limb of the DBB (Mesulam et al., 1983a; Gaykema et al., 1990).

Ultrastructural characterization of cholinergic DBB neurons has yielded more information about the synaptic connections of neurons projecting to the hippocampus and the neocortex. Cholinergic fibres projecting to the hippocampus have en passant axon collaterals that synapse with local somata and dendrites in the DBB (Henderson et al.,

2001). Electrophysiological experiments suggest that these collateral projections exert a modulatory effect on DBB neurons (Henderson et al., 2005). The main cholinergic fibres project to the hippocampus via the fornix and synapse onto hippocampal neurons (Nyakas et al., 1987). These hippocampal synapses terminate in the CA1 and CA3 regions onto pyramidal and granule neurons (Nyakas et al., 1987). GABAergic neurons of the DBB project along the fornix and synapse upon GABAergic neurons in the stratum oriens of the CA1 region of the hippocampus (Gulyas et al., 1990).

The cholinergic neurons of the DBB also project to the cingulate, visual, and olfactory cortices (Bigl et al., 1982; Luskin and Price, 1982; McKinney et al., 1983). GABAergic neurons of the DBB mainly project to the cingulate cortex, however, the projection to the visual cortex is sparse (Freund and Gulyas, 1991). The GABAergic terminals from DBB neurons form en passant synapses on GABAergic cell bodies and dendritic spines in all layers of the cortex (Freund and Gulyas, 1991).

The DBB projections that are important for cardiovascular regulation include projections to the supraoptic and paraventricular nuclei of the hypothalamus and the amygdala (Meibach and Siegel, 1977; Oka and Yoshida, 1985; Kovacs and Versteeg, 1993). DBB projections to the amygdala are also involved in memory and learning (Gallagher et al., 1977; Stock et al., 1981; Hostetter et al., 1987).

The DBB also projects to the dorsal raphe and the locus coeruleus which are areas with high concentrations of biogenic amines (Lee et al., 2005). The dorsal raphe nucleus contains somatostatin expressing neurons and receives a non-cholinergic, inhibitory innervation which is believed to be important in REM sleep (Kalen and Wiklund, 1989; Guzman-Marin et al., 2000). Little is known about the projection to the locus coeruleus,

except that a few identified neurons projecting from DBB to the locus coeruleus, also project to the dorsal raphe (Lee et al., 2005). The functional implications of this collateral branching connectivity to these two nuclei remain to be elucidated.

2. Afferent Connections to the DBB

The DBB receives inputs from many of the areas to which it projects. The reciprocal connectivity of these projections has implications for how the DBB functions in physiological processes. Anatomical studies have demonstrated that afferent input to the DBB originates from the cerebral cortex, hypothalamus, amygdala, the brainstem locus coeruleus and dorsal raphe.

Autoradiography and anterograde tracer techniques have shown glutamatergic and GABAergic neurons project from the hippocampus to the DBB (Freund, 1989; Carnes et al., 1990). Most of these projection neurons are GABAergic and form a reciprocal septohippocampal connection with GABAergic neurons of the DBB, however, some of the neurons also synapse onto the cholinergic neurons (Toth et al., 1993). There is also a minor glutamatergic projection from the hippocampus to the DBB (Carnes et al., 1990). The orbital, prefrontal, insular and olfactory cortices also provide some glutamatergic innervation to both the vertical and horizontal limbs of the DBB (Carnes et al., 1990; Zaborszky et al., 1997). These glutamate fibres cannot account for all the cortical projections to the DBB and the remaining fibres contributing to the cortical input to DBB are believed to be GABAergic.

The DBB receives significant input from all areas of the hypothalamus. Retrograde labeling has shown that there is significant glutamatergic innervation of the DBB from the medial and lateral aspects of the hypothalamus (Carnes et al., 1990). In

addition to glutamatergic innervation, vasopressin releasing neurons project to the DBB from the medial hypothalamus, the bed nucleus of the stria terminalis and the amygdala (Sofroniew, 1985; Caffé et al., 1989; Insel et al., 1994).

Catecholaminergic innervation of the DBB has been observed from the locus coeruleus and the dorsal raphe nuclei (Swanson and Cowan, 1979; Semba et al., 1988; Zaborszky and Cullinan, 1996). Although the precise chemical phenotype of these projections has not been definitively identified, they are most likely noradrenergic and serotonergic projections.

II. Physiological Functions of the DBB

Due to its extensive connectivity, the DBB is involved in a wide array of physiological functions in the central nervous system. The involvement of the DBB in theta rhythm, learning and memory are well characterized and important in the context of understanding how the loss of basal forebrain neurons causes cognitive dysfunction in the context of AD.

A. Theta Rhythm

Theta rhythm is a synchronized, low frequency (4-12 Hz) oscillation that is recorded in limbic structures (Leung et al., 1982). Initial studies associated theta rhythm with an arousal state. As theta is associated with a wide variety of physiological functions such as active exploratory sniffing (Forbes and Macrides, 1984), vibrissae movements (Semba and Komisaruk, 1984), eye movements during immobile visual exploration (Kemp and Kaada, 1975), and rapid eye movements in paradoxical sleep (Monmaur, 1981), it has been hypothesized that theta allows the hippocampus to

selectively receive inputs from sensory systems and voluntary motor systems to monitor performance in tasks (Vinogradova, 1995; Bland and Oddie, 2001; Buzsaki, 2002).

The involvement of the medial septum in the generation of theta rhythm was first observed in experiments in which lesioning of the DBB resulted in the loss of theta activity (Petsche et al., 1962). Electrophysiological experiments characterizing DBB neurons demonstrated that DBB neurons switch between firing bursts of action potentials either in phase or out of phase with theta (Gaztelu and Buno, 1982; Sweeney et al., 1992). Bland and Colom (1993) hypothesized that theta frequency modulation is determined by the balance of septal GABAergic and cholinergic inputs to the hippocampus. As theta frequency rhythm decreases below 5 Hz, there is less inhibitory input from the DBB to the interneurons of the hippocampus, so that these interneurons are disinhibited. Consequently, the interneurons decrease the activity of neurons in phase with theta rhythm. As theta rhythm rises above 5 Hz, cholinergic-mediated EPSPs in the pyramidal neurons begin to dominate, resulting in a feed-forward increase in theta rhythm. The role of the DBB in theta suggests that these neurons may be important in maintaining an aroused state.

B. Learning and Memory

A central role for DBB in functions related to learning and memory is supported by several lines of evidence. Lesions to the DBB result in impairments in spatial learning, visual memory tasks and olfactory learning tasks (Hagan et al., 1988; Ridley et al., 1988; Paolini and McKenzie, 1993). Bilateral lesions of both GABAergic and cholinergic projection neurons in the DBB reduce the performance of rats in the Morris water maze suggesting that DBB projections are important in spatial memory (Hagan et

al., 1988). Subsequent studies in which cholinergic neurons were selectively lesioned demonstrated that although the loss of cholinergic neurons does not affect spatial learning when measured by the Morris water maze, rats with cholinergic lesions used different learning strategies to negotiate these tasks (Janis et al., 1998; Janisewicz and Baxter, 2003). The results of non-selective lesioning of the DBB and selective lesioning of cholinergic neurons showed that both GABAergic and cholinergic neurons play a role in spatial learning and memory.

The horizontal DBB connections to the olfactory bulb are involved in memory and learning as bilateral lesions of this region results in significant impairment in odor-reward associations and odor habituation (Paolini and McKenzie, 1993; Roman et al., 1993). These studies suggest that the horizontal DBB acts as a relay station between the olfactory bulb and the olfactory cortex. The DBB also plays a role in visual discrimination as excitotoxic lesions which reduce cholinergic activity in the cingulate cortex, an area where there is significant DBB projections, reduced performance in a visual discrimination test (Marston et al., 1994). These studies demonstrate that the DBB plays an important role in many different modes of learning and that loss of these projection neurons can significantly impair cognitive function.

C. DBB Neuronal Loss in AD

Post-mortem examination of brains from individuals with AD established a link between the preferential loss of basal forebrain cholinergic neurons and cognitive impairment (Whitehouse et al., 1982). The most severely affected regions in the basal forebrain were the medial septum which lost 50% of its neuronal population, and the DBB which lost 65% of its cholinergic neurons (Henke and Lang, 1983). Lesions of the

DBB in rats produce disruptions in spatial memory and these changes can be attributed to the loss of cholinergic neurons (Janis et al., 1998; Janisewicz and Baxter, 2003) In addition, to the depletion of cholinergic neurons, there is also accumulation of soluble A β , the deposition of insoluble amyloid plaques and the presence of neurofibrillary tangles in the basal forebrain (Saper et al., 1985). Given the importance of DBB cholinergic neurons in memory and learning, the significant degeneration of basal forebrain cholinergic neurons in AD may be responsible for the cognitive decline and that treatments designed to protect the loss of cholinergic cells may protect individuals against cognitive decline observed in AD.

III. Cellular Properties of DBB Neurons

A. Intrinsic Electrophysiological Properties

Initial examination of the electrophysiological properties of individual DBB neurons performed in guinea pigs identified 3 populations of neurons of this region (Griffith, 1988). The first neuronal cell type exhibited a slow afterhyperpolarization (AHP) of 600 ms with an amplitude of 10-20 mV which was attributed to a Ca²⁺ – activated potassium (K⁺) conductance. Cells with the 600 ms AHP comprised 40% of the neurons recorded from and were called SAHP neurons. The second cell type identified exhibited a fast AHP (duration 5-50 ms) and comprised 53% of the cells recorded. The final cell type comprised the smallest population of neurons (7%) and fired in a burst pattern. Staining for AChE in cells that had been recorded from demonstrated that only SAHP neurons expressed AChE, suggesting that these were cholinergic neurons, while FAHP neurons were GABAergic. These findings were confirmed later on in slice preparations (Markram and Segal, 1990; Wu et al., 2000).

DBB neurons exhibit many voltage-activated and ligand-activated currents. A transient outward current (A-current) that was blocked by 4-aminopyridine and a suite of Ca^{2+} channels were described in acutely dissociated DBB cells (Griffith and Sim, 1990). Cholinergic neurons possessed both high voltage-activated and low voltage-activated Ca^{2+} channels, while non-cholinergic population only expressed high voltage-activated currents (Griffith and Sim, 1990). Further pharmacological analysis of cholinergic neurons demonstrated that N and L-type channels were the predominant population of Ca^{2+} channels, while the P/Q-type and T-type channels made up a small component of the population of Ca^{2+} channels (Murchison and Griffith, 1995, 1996; Easaw et al., 1999; Chin et al., 2002). These channels are modulated by many different ligands, suggesting that DBB neurons are influenced by a variety of neurotransmitters and neuromodulators (Chin and Jhamandas, 2002; Jhamandas and Mactavish, 2002).

B. Receptors Expressed on DBB Neurons

Slice recordings from the DBB demonstrated that there are many receptors systems involved in the regulation of neurons. Initial whole cell recordings demonstrated the presence of glutamate receptors on horizontal DBB neurons as perfusion of α -amino-3-hydroxy-5-methyl-4-isoxazolepropionate (AMPA) increased inward currents during voltage ramps, which were blocked by the AMPA receptor antagonist 6-cyano-7-nitroquinoxaline-2,3-dione (CNQX) (Easaw et al., 1997). In addition, stimulation of the DBB evoked excitatory postsynaptic currents (EPSCs) that were blocked by CNQX, indicating the presence of a glutamatergic input onto DBB neurons. In the absence of magnesium, application of *N*-methyl-D-aspartate (NMDA) onto the slices also evoked an inward current that was inhibited by (-) 2-amino-7-phosphonopentanoate (APV). Further

examination of EPSCs showed the presence of a NMDA component on some cells (Easaw et al., 1997).

C. Intracellular Ca^{2+} stores

Ca^{2+} is an important intracellular ion, with many physiological functions ranging from involvement in release of neurotransmitters at synapses to regulation of neuronal death. Given the wide variety of functions that are influenced by intracellular Ca^{2+} concentration ($[Ca^{2+}]_i$), it is not surprising that it is very tightly regulated within individual neurons.

As discussed earlier, the basal forebrain neurons possess a suite of voltage gated Ca^{2+} channels and neurotransmitter gated channels that gate extracellular Ca^{2+} entry (Murchison and Griffith, 1995, 1996; Chin et al., 2002). The endoplasmic reticulum (ER) and the mitochondria are intracellular sources of Ca^{2+} release in the DBB (Murchison and Griffith, 1999, 2000).

The ER is a continuous endomembrane structure that extends throughout the neuron from the nuclear envelope to the distal dendrites (Martone et al., 1993). The ER accumulates intraluminal Ca^{2+} via the sarco-ER Ca^{2+} ATPase (SERCA) located on its membrane (Verkhratsky, 2005). Ca^{2+} is released from the ER by the activation of either inositol 1,4,5-phosphate receptor (InsR) or the ryanodine receptor (RyR), which are tetrameric Ca^{2+} channels that look remarkably similar under electron microscopy (Prentki et al., 1984; McPherson et al., 1991; Verkhratsky, 2005). Ca^{2+} is the endogenous ligand for RyRs, and in the DBB, these receptors amplify VGCC mediated $[Ca^{2+}]_i$ increases (Murchison and Griffith, 1999). A useful pharmacological tool used in the study of RyR is caffeine, which is a membrane soluble agonist of the RyR and an antagonist of the

InsR (Verkhatsky, 2005). At millimolar concentrations, caffeine is capable of raising $[Ca^{2+}]_i$, and depleting intracellular stores (Murchison and Griffith, 1999). The signal for release of Ca^{2+} from InsRs is transduced from the activation of a cell surface metabotropic receptor linked to the enzyme phospholipase C. Phospholipase C then cleaves phospholipids to generate inositol 1,4,5-phosphate (IP3), which diffuses to the ER and activates InsR (Berridge, 1993).

In addition to their role in energy production, mitochondria also play an important role in the regulation of Ca^{2+} in both pathophysiological and physiological states (White and Reynolds, 1996; Murchison and Griffith, 2000). Initial studies focusing on the role of mitochondria in excitotoxic neuronal death found that excess accumulation of Ca^{2+} in the mitochondria from influx of Ca^{2+} via glutamate receptors plays a key role in neuronal death (White and Reynolds, 1996; Stout et al., 1998). In DBB neurons, depolarization of the mitochondria has been shown to be capable of releasing Ca^{2+} , which suggests that mitochondria may play a role in the regulation of Ca^{2+} (Murchison and Griffith, 2000). Given the importance of these intracellular Ca^{2+} stores in regulating Ca^{2+} and apoptosis, they may be important targets for $A\beta$ in mediating neurodegeneration in AD.

IV. Nicotinic Acetylcholine Receptor in the Central Nervous System

A. General Properties

Nicotinic acetylcholine receptors (nAChRs) are part of a superfamily of pentameric ligand gated channels that includes the serotonin, GABA, and glycine receptors. The properties of the nAChRs such as ligand affinity, opening probabilities, and cation permeability are dependent on subunit composition. The genes cloned for

neuronal nAChR subunits are divided into two subfamilies. The α subunit subfamily has 9 members ($\alpha 2$ - $\alpha 10$), while the β subunit subfamily has 3 members ($\beta 2$ - $\beta 4$).

The division of neuronal nAChR subunits into 2 subfamilies is based upon the homology of the neuronal α subfamily with the muscle α subunit, which both have consecutive cysteine residues at positions 192 and 193, while the β subunit subfamily does not have consecutive cysteine residues (Boulter et al., 1987). Two of the most commonly expressed nAChRs in the brain are the heteromeric $\alpha 4\beta 2$ and homomeric $\alpha 7$ nAChRs. The subunits which form the $\alpha 4\beta 2$ receptor along with the $\alpha 3$ subunit were initially discovered by low stringency DNA-DNA hybridization (Boulter et al., 1986; Boulter et al., 1987; Goldman et al., 1987). Electrophysiological studies performed on oocytes established that $\alpha 4$ and $\beta 2$ subunits assemble to form functional nAChRs as cotransfection of these subunits yielded a greater current response to ACh than if the $\alpha 4$ subunit was expressed on its own (Boulter et al., 1987).

The initial pharmacological characterization of the $\alpha 4\beta 2$ nAChR revealed that this receptor was not sensitive to blockade by α -bungarotoxin, a well established blocker of muscle nAChRs (Boulter et al., 1987). This suggests that there were additional nAChR subunits to be discovered as it was well established that there were α -bungarotoxin binding sites in the brain (Wang and Schmidt, 1976; Wang et al., 1978; Conti-Tronconi et al., 1985). Eventually, a new α subunit with a significantly different nucleotide sequence and splice sites from the $\alpha 2$ - $\alpha 6$ subunits was cloned and termed the $\alpha 7$ subunit (Couturier et al., 1990; Schoepfer et al., 1990). Expression of the $\alpha 7$ subunit on its own in oocytes resulted in the formation of a nAChR channel with a rapidly desensitizing, α -bungarotoxin-sensitive inward current (Couturier et al., 1990).

The $\alpha 4\beta 2$ and $\alpha 7$ channels display a significant degree of permeability to Ca^{2+} , an important second messenger in intracellular signal transduction pathways. The permeability of the $\alpha 7$ nAChR to Ca^{2+} is comparable to that of the NMDA glutamate receptor (Castro and Albuquerque, 1995). The $\alpha 4\beta 2$ nAChR is not as permeable to Ca^{2+} as the $\alpha 7$ nAChR as the fraction of Ca^{2+} ions passing through $\alpha 4\beta 2$ nAChRs is 2.6%, while the fraction of Ca^{2+} ions passing through $\alpha 7$ nAChR is 11% (Lax et al., 2002; Fucile et al., 2003). The significant Ca^{2+} permeability of nAChRs suggests that they play a role in Ca^{2+} regulation of DBB neurons and second messenger signaling.

B. NACHRs in the DBB

A recent report examining nAChRs in the DBB has shown that focal application of ACh onto medial septum/DBB neurons, elicits inward currents that can be blocked by mecamylamine, a $\alpha 7$ nAChR selective antagonist and dihydro- β -erythroline, a non- $\alpha 7$ selective antagonist (Henderson et al., 2005). Of the neurons recorded from, a mecamylamine sensitive response was present in all cholinergic neurons examined and 10% of GABAergic neurons. However, 50% of GABAergic neurons displayed non- $\alpha 7$ nicotinic responses (Henderson et al., 2005). This data should be interpreted with caution as mecamylamine is not a pure $\alpha 7$ nAChR antagonist as it blocks both $\alpha 7$ and $\alpha \beta$ heteromeric nAChRs in the micromolar range (Chavez-Noriega et al., 1997). This leaves open the possibility that other nAChRs could be expressed on cholinergic neurons. This data suggest that the nAChR receptor system plays an important role in the modulation of neuronal activity in the cholinergic basal forebrain system.

C. Physiological Functions of nAChRs in the Central Nervous System

1. Modulation of Transmitter Release

NAChRs play an important role in modulation of neurotransmitter release in the brain. Dopamine release from substantia nigra nerve terminals in the striatum was one of the first neurotransmitter systems found to be modulated by nAChRs (Westfall, 1974; Rapier et al., 1990). Initially, when brain slices containing the striatum were perfused with nicotine, an increase in dopamine release was observed which was attributed to activation of presynaptic nAChRs (Westfall, 1974). Pharmacological and molecular characterization of dopaminergic synaptic terminals established that activation of presynaptic $\alpha 4\beta 2$ nAChRs increases dopamine release in the striatum (Rapier et al., 1990; Sacaan et al., 1995; Sharples et al., 2000; Zoli et al., 2002). The first electrophysiological evidence for presynaptic modulation of neurotransmitter release was found in the medial habenula nucleus of the hypothalamus (McGehee et al., 1995). Activation of $\alpha 7$ nAChRs was found to enhance glutamate release in a Ca^{2+} dependent manner (McGehee et al., 1995).

Neurons in areas involved in learning and memory such as the cortex and hippocampus are also modulated by nAChRs. In cortical interneurons, $\alpha 4\beta 2$ and $\alpha 7$ nAChRs mediate both post-synaptic and pre-synaptic responses (Alkondon et al., 2000). Additionally, there are presynaptic $\alpha 4\beta 2$ nAChRs that increase the release of GABA onto cortical interneurons (Alkondon et al., 2000). In the hippocampus, $\alpha 4\beta 2$ and $\alpha 7$ nAChRs are found to mediate postsynaptic currents in CA1 pyramidal neurons and interneurons, however, $\alpha 7$ nAChRs are also found to increase presynaptic release of GABA (Alkondon and Albuquerque, 1993; Alkondon et al., 1997).

2. nAChRs and Second Messengers

Given that nAChRs have high permeability to Ca^{2+} , studies have established these receptors to be involved in signal transduction pathways crucial to cell survival. In pheochromocytoma cells, activation of nAChRs depolarizes the cell resulting in Ca^{2+} influx that activates Protein kinase C (PKC). PKC then phosphorylates Raf1, a mitogen activated kinase kinase kinase, starting a cascade that leads to the eventual phosphorylation of CREB which then activates gene transcription (Tang et al., 1998). In the prefrontal cortex, $\alpha 4\beta 2$ nAChR activation leads to activation of intracellular Ca^{2+} stores, which then activates PKC, leading to increased dopamine release through the dopamine transporter (Drew and Werling, 2001). Activation of $\alpha 7$ nAChRs has been shown to cause Ca^{2+} entry via voltage gated Ca^{2+} channels, which leads to activation of intracellular Ca^{2+} stores via activation of InsRs (Dajas-Bailador et al., 2002). Release of Ca^{2+} from intracellular stores has been shown to activate the extracellular signal receptor kinase pathway (ERK1/2) which is important in learning and memory (Dajas-Bailador et al., 2002; Tomizawa and Casida, 2002). There are additional pathways to be discussed below.

3. Age-Dependent Alteration of nAChRs

In both rats and humans, nAChRs are present during the early moments of life. In rats, the $\alpha 2$, $\alpha 3$, $\alpha 4$, $\alpha 5$, $\alpha 6$, $\beta 2$ and $\beta 4$ subunits are localized in the diencephalon, brainstem, spinal cord and telencephalon (Gotti and Clementi, 2004). However as the individual subunits need to combine with other subunits in order to form receptors, variations in the temporal expression of the subunits also varies the expression of functional nAChRs, for example, the $\beta 4$ subunit precedes the $\alpha 3$ subunit by 2 days,

which combine to form $\alpha 3\beta 4$ nAChRs (Zoli et al., 1995). In the human brain, nAChR expression starts at 12 weeks of gestation and reaches an apex at 27 weeks of gestation. During the perinatal period and early infancy, nAChR concentration decreases in all brain areas except for the cerebellum, where concentrations remain constant. Interestingly, the highest concentration of nAChRs during fetal development was observed in the nucleus basalis of Meynert, a basal forebrain nucleus (Cairns and Wonnacott, 1988).

During ageing, nAChR binding sites decrease in both rats and humans. Both $\alpha 7$ and $\alpha 4$ subunit mRNA and protein levels were lower in aged humans (60-90 years of age) in the DBB, medial septum, and frontoparietal cortex which may contribute to cognitive decline (Tohgi et al., 1998). Declines in nAChR receptor expression has also been observed in rats, however, the level of nAChR loss is not associated with decreased performance in learning and memory tasks (Smith et al., 1995).

4. NACHRs and AD

NACHR expression in certain areas of AD brains is significantly reduced when compared to age-matched controls (Flynn and Mash, 1986). Removal of cholinergic input in rats by selectively lesioning cholinergic neurons in the basal forebrain leads to decreased performance in learning and memory task (Steckler et al., 1995). Furthermore, nicotinic agonist treatment improves performance in memory and learning tasks (Levin and Simon, 1998).

Studies measuring nicotinic radioligand binding comparing age-matched control brains with AD brains have established that there is a significant decrease in nAChR binding sites (Whitehouse et al., 1988; Warpman and Nordberg, 1995). These decreases in nAChR have been localized to mainly those areas of the brain associated with

cognition such as the frontal cortex, temporal cortex, and the hippocampus (Rinne et al., 1991; Warpman and Nordberg, 1995). Immunohistochemical examination of the subtypes of nAChRs frontal cortex revealed a 30% decrease in $\alpha 4\beta 2$ and $\alpha 7$ nAChR neurons (Wevers et al., 1999). In the temporal cortex, $\alpha 4\beta 2$ nAChR binding sites and $\alpha 4$ subunit expression were decreased, while there was no change in $\alpha 3$ and $\alpha 7$ subunit expression (Martin-Ruiz et al., 1999). In the hippocampus, there was a decrease in nAChR binding sites in the dentate granular layer, the presubiculum, and the parahippocampal gyrus, however, no changes in nAChR binding sites were observed in the CA1 or CA3 layers (Perry et al., 1995). The loss of specific nAChR binding sites in brain regions involved in memory and learning may play an important role in mediating the cognitive impairment observed in AD.

V. General Properties of Amyloid

A β , a 39-43 amino acid peptide, which was isolated from amyloid plaques, plays an important role in the AD pathophysiological process (Glennner and Wong, 1984; Masters et al., 1985a; Masters et al., 1985b). Initial studies of AD individuals detected no correlation between dementia and insoluble A β load, however, when both soluble and insoluble forms of A β were measured, the A β load in these areas correlated with the extent of cognitive decline (Terry, 1999; Naslund et al., 2000).

The amyloid hypothesis postulates that an increase of A β production is the primary pathophysiological mediator of AD (Selkoe, 2001). Three lines of evidence support the basis for this hypothesis. The deposition of A β in the brains of AD individuals is a major pathological finding. Secondly, application of A β onto a variety of animal and human neuronal preparations is neurotoxic (Mattson et al., 1992; Yan et al.,

1996; Nakagawa et al., 2000; Jhamandas and MacTavish, 2004). Finally, mutations associated with early onset AD increase A β production (Citron et al., 1992; Borchelt et al., 1996). These mutations are located either on proteolytic processing sites of APP, which is the precursor of A β or on the genes of proteins responsible for APP cleavage (Chartierharlin et al., 1991; StGeorgehylop et al., 1992). Individuals afflicted with Down's syndrome, who have an extra chromosome 21, which is the chromosome that APP is located on, exhibit severe mental retardation and developmental delay along with significant deposition of A β in their brains (Masters et al., 1985c).

A. A β production

A β is a cleavage product of a larger precursor protein, the APP (Kang et al., 1987; Kitaguchi et al., 1988; Ponte et al., 1988; Tanzi et al., 1988). There are 8 known isoforms of the APP gene that are formed by differential splicing, however the major isoforms of APP are 695, 751 and 770 residues in length (Sandbrink et al., 1996). The 751 and 770 residue form of APP are different from the 695 residue form in that they have a Kunitz-type serine protease inhibitor domain (Kang et al., 1987; Kitaguchi et al., 1988; Ponte et al., 1988; Tanzi et al., 1988).

APP has multiple cleavage sites and is processed either by a non-amyloidogenic or amyloidogenic pathway (Figure 1)(Selkoe, 2001). α -Secretase cleaves APP to generate a long fragment termed α APP_s and an 83 amino acid carboxy-terminal peptide (Esch et al., 1990; Sisodia et al., 1990). Proteolytic cleavage of APP by α -secretase is the first step of non-amyloidogenic cleavage as it precludes the formation of A β (Buxbaum et al., 1998; Lammich et al., 1999). If APP is not processed by α -secretase, APP is cleaved by an enzyme termed β -secretase to generate β APP_s, a fragment that is slightly

smaller than α APP_s and a 99 amino acid carboxy-terminal fragment that begins at residue 1 of the A β protein (Vassar et al., 1999; Selkoe, 2001). The carboxy-terminal fragments produced by α - and β -secretase cleavage are subsequently cleaved by γ -secretase. γ -Secretase cleavage of the 83 amino acid COOH-terminal fragment yields a NH₂-terminal fragment terminal peptide termed p3, while cleavage of the 99 amino acid carboxy-terminal fragment yields the 39 to 43 amino acid forms of A β (Selkoe, 2001). The proteolytic enzyme responsible for γ -secretase activity is composed of 5 proteins, which are termed presenilin-1, presenilin-2, α 1, nicastrin and pen-2 that form an intramembrane complex to facilitate carboxy-terminal cleavage (Selkoe, 2002a).

B. Conformation of A β

After A β is secreted, it forms different conformations which determine its effects on neurons (Pike et al., 1993; Lambert et al., 1998; Walsh et al., 2002). The simplest forms of A β are the soluble monomeric and small oligomeric forms that are found in the cerebrospinal fluid (Hilbich et al., 1991; Seubert et al., 1992). The small oligomeric forms have been shown to inhibit long term potentiation, while the monomeric forms exhibited no effect (Walsh et al., 2002). In addition, small oligomeric forms are significantly more neurotoxic than either fibrils or monomers and the toxicity is specific to the aggregation of A β (Pike et al., 1993; Lambert et al., 1998; Dahlgren et al., 2002). Over time, oligomeric forms of A β eventually aggregate into non-soluble fibrils that are in equilibrium with soluble forms of A β . These are the major constituent of neuritic plaques (Hilbich et al., 1991). However, prior to the formation of fibrils, a transient proto-fibril conformation is observed (Harper et al., 1997; Walsh et al., 1997). The proto-fibril is 6-10 nm in diameter and between 30 and 500 nm in length, while fibrils

exhibit a beaded appearance, with a maximal diameter that is larger than the proto-fibril and extend for over 1 μM (Harper et al., 1997; Walsh et al., 1997). $\text{A}\beta_{1-42}$ is significantly more fibrillogenic than $\text{A}\beta_{1-40}$ (Harper et al., 1997). The fibrillar form of $\text{A}\beta$ is neurotoxic, although it is significantly less neurotoxic than the soluble oligomeric forms (Dahlgren et al., 2002). The work on the different conformations of $\text{A}\beta$ highlights the importance of soluble conformations of $\text{A}\beta$ in mediating neuronal effects.

C. Effects of $\text{A}\beta$ on neurons

$\text{A}\beta$ has been shown to disrupt neuronal function and mediate neurotoxicity in many neuronal systems, however, its definitive role and receptor that mediates its actions remains unknown. Several pathways have been proposed to mediate $\text{A}\beta$ neurotoxicity. One hypothesis is that $\text{A}\beta$ mediates neurotoxicity by causing oxidative stress on neurons (Behl et al., 1994). Generation of free radicals results neuronal damage and activation of a pro-inflammatory transcription factor NF-kappa-B, which is believed to further contribute to neuronal death (Kaltschmidt et al., 1997). Another possible mediator of $\text{A}\beta$ mediated neurotoxicity is the disruption of Ca^{2+} homeostasis (Mattson et al., 1992). The exact mechanism whereby $\text{A}\beta$ affects Ca^{2+} homeostasis is unknown and many theories have been suggested. Evidence for $\text{A}\beta$ peptides forming plasma membrane channels that pass Ca^{2+} ions have been observed in lipid bilayers, however there is no convincing physiological data in neurons demonstrating the passage of Ca^{2+} ions through these channels (Arispe et al., 1993; Rhee et al., 1998). Aggregated $\text{A}\beta$ inhibits Ca^{2+} current through voltage-gated Ca^{2+} channels, while unaggregated $\text{A}\beta$ inhibits this current (Ramsden et al., 2002). $\text{A}\beta$ has been shown to disrupt the functioning of the ER, an important regulator of Ca^{2+} and results in the activation of caspases, the proteolytic

cascade which leads to apoptosis (Nakagawa et al., 2000). A β has also been shown in slice preparations to enhance glutamate release, which would lead to increased Ca²⁺ entry into neurons via glutamatergic receptors (Kar, 2005). *These studies support the hypothesis that A β disruption of Ca²⁺ homeostasis is an important step in mediating neurodegeneration.*

D. A β and nAChRs

A major conceptual advance in the role of the nAChR in neurodegeneration was the discovery that both the 40 and 42 amino acid forms of the A β protein could bind to nAChRs (Wang et al., 2000b; Wang et al., 2000a). Further studies have shown that A β is capable of functional modulation of the nAChR (Liu et al., 2001; Pettit et al., 2001; Dineley et al., 2002; Fu and Jhamandas, 2003; Grassi et al., 2003). In their original studies on A β -nAChR binding, Wang et al. (2000a) showed that A β were capable of competitively displacing α 7 nAChR selective antagonists from synaptic membranes produced from guinea pig and rat cortex and hippocampus with picomolar affinity. A β was also found to be able to bind to α 4 β 2 nAChRs with nanomolar affinity (Wang et al., 2000a). Additional evidence for the A β interaction with nAChR was obtained from co-immunoprecipitation data (Wang et al., 2000b). A β binding nAChR could potentially activate apoptotic pathways and the A β -nAChR complex may seed the development of amyloid plaques (Wang et al., 2000b). The nAChR-A β binding studies have been very interesting, however, it is problematic that other laboratories have had difficulty reproducing the original findings of Wang and co-workers (2000a, b).

Electrophysiological experiments examining the effect of A β on nAChR in the CNS have focused on two subtypes of nAChR, the homomeric α 7 nAChR and the

heteromeric $\alpha 4\beta 2$ nAChR. Initially, studies showed that in CA1 stratum radiatum interneurons, 100 nM $A\beta_{1-42}$ blocked $\alpha 7$ nAChR responses (Liu et al., 2001; Pettit et al., 2001; Grassi et al., 2003). Studies of the $A\beta$ interaction with $\alpha 7$ nAChRs in *Xenopus* oocytes expression systems suggest that $A\beta_{1-42}$ weakly activates nAChRs resulting in desensitization of the $\alpha 7$ nAChRs (Dineley et al., 2002; Grassi et al., 2003). Dougherty et al. (2003) examined the effect of $A\beta$ on presynaptic terminals and found that $A\beta$ antagonizes homomeric $\alpha 7$ nAChR at picomolar concentrations by blocking agonist binding.

Further examination of $A\beta$ actions at a cellular level also revealed that it has effects on $\alpha 4$ nAChRs (Fu and Jhamandas, 2003; Wu et al., 2004). In basal forebrain neurons, $A\beta_{1-42}$ activated $\alpha 4\beta 2$ nAChRs in a dose dependent manner and this activation lead to depolarization of these neurons (Fu and Jhamandas, 2003). Another study examining the effect of $A\beta$ on $\alpha 4\beta 2$ nAChR expressed in a cell line revealed that $A\beta$ blocks $\alpha 4\beta 2$ nAChR in a non-competitive manner, however, when the cell line was transfected with functional $\alpha 7$ nAChRs, $A\beta_{1-42}$ was unable to block these receptors (Wu et al., 2004). Given the conflicting data with regards to $A\beta_{1-42}$ and nAChRs interactions, two issues remained to be resolved. Firstly, the nAChR subtype that $A\beta$ interacts with is unclear and secondly, whether $A\beta$ acts as an agonist or antagonist at such receptors remains to be resolved.

The consequences of these $A\beta$ actions on nAChR interactions are undoubtedly wide-ranging but from a pathophysiological point of view, the effects of such interactions on Ca^{2+} homeostasis in the cell are perhaps the most intriguing. A potential consequence of the $A\beta$ -nAChR interaction would be the excessive entry of Ca^{2+} into presynaptic


terminals resulting in inappropriate desensitization of nAChR and activation of second messenger pathways that may lead to neurodegeneration (Dougherty et al., 2003). Alternately, the interaction of A β and the nAChR may lead to internalization of the receptor and A β_{1-42} , where A β may mediate additional effects (Nagele et al., 2002).

Activation of the $\alpha 7$ nAChR also activates second messenger systems important in neuroprotection. The effect of A β has been examined on the Janus kinase 2 (JAK2) and the mitogen activation protein kinase (MAP2) pathways, which are two neuroprotective pathways that the $\alpha 7$ nAChR activates (Dineley et al., 2001; Shaw et al., 2002). Activation of the JAK2 pathway results in the JAK2 enzyme phosphorylating itself and activating downstream mediators phosphatidylinositol and akt to prevent activation of apoptotic pathways (Shaw et al., 2002). It was found that $\alpha 7$ nAChR mediated neuroprotection via this pathway in cultured neurons was inhibited by A β_{1-42} (Shaw et al., 2002). Another possible mechanism via which A β_{1-42} mediates its effects is via excessive activation of $\alpha 7$ nAChR, which would down-regulate the MAPK2 pathway and reduce the phosphorylation of its downstream target, the cAMP regulatory element binding protein (CREB). This cascade may lead to derangement of the signaling pathway (Dineley et al., 2001). These data demonstrate the importance of second messenger systems mediated by nAChRs and the ability of A β to modulate these systems. *As A β has been implicated to both disrupt synaptic function and disturb Ca^{2+} homeostasis, the studies on nAChR-A β interactions would shed light on the effect of A β on two important cellular functions; the tight regulation of $[Ca^{2+}]_i$ and synaptic function.*

VI. Objectives and Rationale

Given the recent data showing that A β binds to nAChRs, I examined the effect of A β on basal forebrain neurons in two ways. In the first part of my thesis, I examined nAChR mediated Ca²⁺ rises in acutely dissociated DBB neurons using the fluorescent ratiometric Ca²⁺ dye fura-2 and investigated the mechanisms whereby A β may modulate [Ca²⁺]_i. In the second part of my thesis, I examined the actions of A β on nAChR modulation of excitatory synaptic neurotransmission in cholinergic DBB neurons using brain slices, an experimental preparation where the neuronal connectivity is intact.


A. APP 770



B. Non-amyloidogenic pathway



C. Amyloidogenic pathway



References

- Alkondon M, Albuquerque EX (1993) Diversity of nicotinic acetylcholine receptors in rat hippocampal neurons. I. Pharmacological and functional evidence for distinct structural subtypes. *J Pharmacol Exp Ther* 265:1455-1473.
- Alkondon M, Pereira EF, Barbosa CT, Albuquerque EX (1997) Neuronal nicotinic acetylcholine receptor activation modulates gamma-aminobutyric acid release from CA1 neurons of rat hippocampal slices. *J Pharmacol Exp Ther* 283:1396-1411.
- Alkondon M, Pereira EF, Eisenberg HM, Albuquerque EX (2000) Nicotinic receptor activation in human cerebral cortical interneurons: a mechanism for inhibition and disinhibition of neuronal networks. *J Neurosci* 20:66-75.
- Arispe N, Rojas E, Pollard HB (1993) Alzheimer-Disease Amyloid Beta-Protein Forms Calcium Channels In Bilayer-Membranes - Blockade By Tromethamine And Aluminum. *Proceedings Of The National Academy Of Sciences Of The United States Of America* 90:567-571.
- Barger SW, Horster D, Furukawa K, Goodman Y, Krieglstein J, Mattson MP (1995) Tumor-Necrosis-Factor-Alpha And Tumor-Necrosis-Factor-Beta Protect Neurons Against Amyloid Beta-Peptide Toxicity - Evidence For Involvement Of A Kappa-B-Binding Factor And Attenuation Of Peroxide And Ca²⁺ Accumulation. *Proceedings Of The National Academy Of Sciences Of The United States Of America* 92:9328-9332.
- Behl C, Davis JB, Lesley R, Schubert D (1994) Hydrogen-Peroxide Mediates Amyloid-Beta Protein Toxicity. *Cell* 77:817-827.

- Berridge MJ (1993) Inositol Trisphosphate And Calcium Signaling. *Nature* 361:315-325.
- Bigl V, Woolf NJ, Butcher LL (1982) Cholinergic projections from the basal forebrain to frontal, parietal, temporal, occipital, and cingulate cortices: a combined fluorescent tracer and acetylcholinesterase analysis. *Brain Res Bull* 8:727-749.
- Bland BH, Oddie SD (2001) Theta band oscillation and synchrony in the hippocampal formation and associated structures: the case for its role in sensorimotor integration. *Behav Brain Res* 127:119-136.
- Borchelt DR, Thinakaran G, Eckman CB, Lee MK, Davenport F, Ratovitsky T, Prada CM, Kim G, Seekins S, Yager D, Slunt HH, Wang R, Seeger M, Levey AI, Gandy SE, Copeland NG, Jenkins NA, Price DL, Younkin SG (1996) Familial Alzheimer's disease-linked presenilin 1 variants elevate A beta 1-42/1-40 ratio in vitro and in vivo. *Neuron* 17:1005-1013.
- Boulter J, Connolly J, Deneris E, Goldman D, Heinemann S, Patrick J (1987) Functional expression of two neuronal nicotinic acetylcholine receptors from cDNA clones identifies a gene family. *Proc Natl Acad Sci U S A* 84:7763-7767.
- Boulter J, Evans K, Goldman D, Martin G, Treco D, Heinemann S, Patrick J (1986) Isolation of a cDNA clone coding for a possible neural nicotinic acetylcholine receptor alpha-subunit. *Nature* 319:368-374.
- Brashear HR, Zaborszky L, Heimer L (1986) Distribution of GABAergic and cholinergic neurons in the rat diagonal band. *Neuroscience* 17:439-451.
- Buxbaum JD, Liu KN, Luo Y, Slack JL, Stocking KL, Peschon JJ, Johnson RS, Castner BJ, Cerretti DP, Black RA (1998) Evidence that tumor necrosis factor alpha

- converting enzyme is involved in regulated alpha-secretase cleavage of the Alzheimer amyloid protein precursor. *J Biol Chem* 273:27765-27767.
- Buzsaki G (2002) Theta oscillations in the hippocampus. *Neuron* 33:325-340.
- Caffe AR, Van Ryen PC, Van der Woude TP, Van Leeuwen FW (1989) Vasopressin and oxytocin systems in the brain and upper spinal cord of *Macaca fascicularis*. *J Comp Neurol* 287:302-325.
- Cairns NJ, Wonnacott S (1988) [³H](⁻)nicotine binding sites in fetal human brain. *Brain Res* 475:1-7.
- Carnes KM, Fuller TA, Price JL (1990) Sources of presumptive glutamatergic/aspartatergic afferents to the magnocellular basal forebrain in the rat. *J Comp Neurol* 302:824-852.
- Castro NG, Albuquerque EX (1995) alpha-Bungarotoxin-sensitive hippocampal nicotinic receptor channel has a high calcium permeability. *Biophys J* 68:516-524.
- Chartierharlin MC, Crawford F, Houlden H, Warren A, Hughes D, Fidani L, Goate A, Rossor M, Roques P, Hardy J, Mullan M (1991) Early-Onset Alzheimers-Disease Caused By Mutations At Codon-717 Of The Beta-Amyloid Precursor Protein Gene. *Nature* 353:844-846.
- Chavez-Noriega LE, Crona JH, Washburn MS, Urrutia A, Elliott KJ, Johnson EC (1997) Pharmacological characterization of recombinant human neuronal nicotinic acetylcholine receptors h alpha 2 beta 2, h alpha 2 beta 4, h alpha 3 beta 2, h alpha 3 beta 4, h alpha 4 beta 2, h alpha 4 beta 4 and h alpha 7 expressed in *Xenopus* oocytes. *J Pharmacol Exp Ther* 280:346-356.

- Chin JH, Jhamandas JH (2002) β -amyloid potentiates nicotine-evoked intracellular calcium rises in acutely dissociated rat basal forebrain neurons. In: Meeting of the Society for Neuroscience.
- Chin JH, Harris K, MacTavish D, Jhamandas JH (2002) Nociceptin/orphanin FQ modulation of ionic conductances in rat basal forebrain neurons. *J Pharmacol Exp Ther* 303:188-195.
- Citron M, Oltersdorf T, Haass C, McConlogue L, Hung AY, Seubert P, Vigopelfrey C, Lieberburg I, Selkoe DJ (1992) Mutation Of The Beta-Amyloid Precursor Protein In Familial Alzheimers-Disease Increases Beta-Protein Production. *Nature* 360:672-674.
- Conrad LC, Pfaff DW (1976a) Efferents from medial basal forebrain and hypothalamus in the rat. I. An autoradiographic study of the medial preoptic area. *J Comp Neurol* 169:185-219.
- Conrad LC, Pfaff DW (1976b) Efferents from medial basal forebrain and hypothalamus in the rat. II. An autoradiographic study of the anterior hypothalamus. *J Comp Neurol* 169:221-261.
- Conti-Tronconi BM, Dunn SM, Barnard EA, Dolly JO, Lai FA, Ray N, Raftery MA (1985) Brain and muscle nicotinic acetylcholine receptors are different but homologous proteins. *Proc Natl Acad Sci U S A* 82:5208-5212.
- Couturier S, Bertrand D, Matter JM, Hernandez MC, Bertrand S, Millar N, Valera S, Barkas T, Ballivet M (1990) A neuronal nicotinic acetylcholine receptor subunit (α 7) is developmentally regulated and forms a homo-oligomeric channel blocked by alpha-BTX. *Neuron* 5:847-856.

- Dahlgren KN, Manelli AM, Stine WB, Baker LK, Krafft GA, LaDu MJ (2002) Oligomeric and fibrillar species of amyloid-beta peptides differentially affect neuronal viability. *Journal Of Biological Chemistry* 277:32046-32053.
- Dajas-Bailador FA, Soliakov L, Wonnacott S (2002) Nicotine activates the extracellular signal-regulated kinase 1/2 via the alpha7 nicotinic acetylcholine receptor and protein kinase A, in SH-SY5Y cells and hippocampal neurones. *J Neurochem* 80:520-530.
- Dineley KT, Bell KA, Bui D, Sweatt JD (2002) beta -Amyloid peptide activates alpha 7 nicotinic acetylcholine receptors expressed in *Xenopus* oocytes. *J Biol Chem* 277:25056-25061.
- Dineley KT, Westerman M, Bui D, Bell K, Ashe KH, Sweatt JD (2001) beta-amyloid activates the mitogen-activated protein kinase cascade via hippocampal alpha 7 nicotinic acetylcholine receptors: In vitro and in vivo mechanisms related to Alzheimer's disease. *Journal Of Neuroscience* 21:4125-4133.
- Divac I (1975) Magnocellular nuclei of the basal forebrain project to neocortex, brain stem, and olfactory bulb. Review of some functional correlates. *Brain Res* 93:385-398.
- Dougherty JJ, Wu JL, Nichols RA (2003) beta-amyloid regulation of presynaptic nicotinic receptors in rat hippocampus and neocortex. *Journal Of Neuroscience* 23:6740-6747.
- Drew AE, Werling LL (2001) Protein kinase C regulation of dopamine transporter initiated by nicotinic receptor activation in slices of rat prefrontal cortex. *J Neurochem* 77:839-848.

- Easaw JC, Petrov T, Jhamandas JH (1997) An electrophysiological study of neurons in the horizontal limb of the diagonal band of Broca. *Am J Physiol* 272:C163-172.
- Easaw JC, Jassar BS, Harris KH, Jhamandas JH (1999) Zinc modulation of ionic currents in the horizontal limb of the diagonal band of Broca. *Neuroscience* 94:785-795.
- Eckenstein F, Sofroniew MV (1983) Identification of central cholinergic neurons containing both choline acetyltransferase and acetylcholinesterase and of central neurons containing only acetylcholinesterase. *J Neurosci* 3:2286-2291.
- Esch FS, Keim PS, Beattie EC, Blacher RW, Culwell AR, Oltersdorf T, McClure D, Ward PJ (1990) Cleavage of amyloid beta peptide during constitutive processing of its precursor. *Science* 248:1122-1124.
- Fisone G, Wu CF, Consolo S, Nordstrom O, Brynne N, Bartfai T, Melander T, Hokfelt T (1987) Galanin inhibits acetylcholine release in the ventral hippocampus of the rat: histochemical, autoradiographic, in vivo, and in vitro studies. *Proc Natl Acad Sci U S A* 84:7339-7343.
- Flynn DD, Mash DC (1986) Characterization of L-[3H]nicotine binding in human cerebral cortex: comparison between Alzheimer's disease and the normal. *J Neurochem* 47:1948-1954.
- Forbes WB, Macrides F (1984) Temporal matching of sensory-motor behavior and limbic theta rhythm deteriorates in aging rats. *Neurobiol Aging* 5:7-17.
- Freund TF (1989) GABAergic septohippocampal neurons contain parvalbumin. *Brain Res* 478:375-381.

- Freund TF, Gulyas AI (1991) GABAergic interneurons containing calbindin D28K or somatostatin are major targets of GABAergic basal forebrain afferents in the rat neocortex. *J Comp Neurol* 314:187-199.
- Fu W, Jhamandas JH (2003) Beta-amyloid peptide activates non-alpha7 nicotinic acetylcholine receptors in rat basal forebrain neurons. *J Neurophysiol* 90:3130-3136.
- Fucile S, Renzi M, Lax P, Eusebi F (2003) Fractional Ca(2+) current through human neuronal alpha7 nicotinic acetylcholine receptors. *Cell Calcium* 34:205-209.
- Gallagher M, Kapp BS, Musty RE, Driscoll PA (1977) Memory formation: evidence for a specific neurochemical system in the amygdala. *Science* 198:423-425.
- Gaykema RPA, Luiten PGM, Nyakas C, Traber J (1990) Cortical Projection Patterns Of The Medial Septum-Diagonal Band Complex. *Journal Of Comparative Neurology* 293:103-124.
- Gaztelu JM, Buno W, Jr. (1982) Septo-hippocampal relationships during EEG theta rhythm. *Electroencephalogr Clin Neurophysiol* 54:375-387.
- Glenner GG, Wong CW (1984) Alzheimer's disease: initial report of the purification and characterization of a novel cerebrovascular amyloid protein. *Biochem Biophys Res Commun* 120:885-890.
- Goate A, Chartierharlin MC, Mullan M, Brown J, Crawford F, Fidani L, Giuffra L, Haynes A, Irving N, James L, Mant R, Newton P, Rooke K, Roques P, Talbot C, Pericakvance M, Roses A, Williamson R, Rossor M, Owen M, Hardy J (1991) Segregation Of A Missense Mutation In The Amyloid Precursor Protein Gene With Familial Alzheimers-Disease. *Nature* 349:704-706.

- Goldman D, Deneris E, Luyten W, Kochhar A, Patrick J, Heinemann S (1987) Members of a nicotinic acetylcholine receptor gene family are expressed in different regions of the mammalian central nervous system. *Cell* 48:965-973.
- Gotti C, Clementi F (2004) Neuronal nicotinic receptors: from structure to pathology. *Prog Neurobiol* 74:363-396.
- Grassi F, Palma E, Tonini R, Amici M, Ballivet M, Eusebi F (2003) Amyloid beta(1-42) peptide alters the gating of human and mouse alpha-bungarotoxin-sensitive nicotinic receptors. *Journal Of Physiology-London* 547:147-157.
- Griffith WH (1988) Membrane properties of cell types within guinea pig basal forebrain nuclei in vitro. *J Neurophysiol* 59:1590-1612.
- Griffith WH, Sim JA (1990) Comparison of 4-aminopyridine and tetrahydroaminoacridine on basal forebrain neurons. *J Pharmacol Exp Ther* 255:986-993.
- Gritti I, Mainville L, Mancina M, Jones BE (1997) GABAergic and other noncholinergic basal forebrain neurons, together with cholinergic neurons, project to the mesocortex and isocortex in the rat. *J Comp Neurol* 383:163-177.
- Gulyas AI, Gorcs TJ, Freund TF (1990) Innervation of different peptide-containing neurons in the hippocampus by GABAergic septal afferents. *Neuroscience* 37:31-44.
- Guzman-Marin R, Alam MN, Szymusiak R, Drucker-Colin R, Gong H, McGinty D (2000) Discharge modulation of rat dorsal raphe neurons during sleep and waking: effects of preoptic/basal forebrain warming. *Brain Res* 875:23-34.

- Hagan JJ, Salamone JD, Simpson J, Iversen SD, Morris RG (1988) Place navigation in rats is impaired by lesions of medial septum and diagonal band but not nucleus basalis magnocellularis. *Behav Brain Res* 27:9-20.
- Hardy J, Allsop D (1991) Amyloid Deposition As The Central Event In The Etiology Of Alzheimers-Disease. *Trends In Pharmacological Sciences* 12:383-388.
- Harkmark W, Mellgren SI, Srebro B (1975) Acetylcholinesterase histochemistry of the septal region in rat and human: distribution of enzyme activity. *Brain Res* 95:281-289.
- Harper JD, Wong SS, Lieber CM, Lansbury PT (1997) Observation of metastable A beta amyloid protofibrils by atomic force microscopy. *Chemistry & Biology* 4:119-125.
- Henderson Z, Morris NP, Grimwood P, Fiddler G, Yang HW, Appenteng K (2001) Morphology of local axon collaterals of electrophysiologically characterised neurons in the rat medial septal/ diagonal band complex. *J Comp Neurol* 430:410-432.
- Henderson Z, Boros A, Janzso G, Westwood AJ, Monyer H, Halasy K (2005) Somato-dendritic nicotinic receptor responses recorded in vitro from the medial septal diagonal band complex of the rodent. *J Physiol* 562:165-182.
- Henke H, Lang W (1983) Cholinergic enzymes in neocortex, hippocampus and basal forebrain of non-neurological and senile dementia of Alzheimer-type patients. *Brain Res* 267:281-291.

- Hilbich C, Kisterswoike B, Reed J, Masters CL, Beyreuther K (1991) Aggregation And Secondary Structure Of Synthetic Amyloid Beta-A4 Peptides Of Alzheimers-Disease. *Journal Of Molecular Biology* 218:149-163.
- Hostetter G, Eaton A, Carnes M, Gildner J, Brownfield MS (1987) Immunocytochemical distribution of luteinizing hormone in rat central nervous system. *Neuroendocrinology* 46:185-193.
- Insel TR, Wang ZX, Ferris CF (1994) Patterns of brain vasopressin receptor distribution associated with social organization in microtine rodents. *J Neurosci* 14:5381-5392.
- Janis LS, Glasier MM, Fulop Z, Stein DG (1998) Intraseptal injections of 192 IgG saporin produce deficits for strategy selection in spatial-memory tasks. *Behav Brain Res* 90:23-34.
- Janisewicz AM, Baxter MG (2003) Transfer effects and conditional learning in rats with selective lesions of medial septal/diagonal band cholinergic neurons. *Behav Neurosci* 117:1342-1352.
- Jhamandas JH, Mactavish D (2002) Central administration of neuropeptide FF (NPFF) causes increased neuronal activation and up-regulation of NPFF gene expression in the rat brainstem. *J Comp Neurol* 447:300-307.
- Jhamandas JH, MacTavish D (2004) Antagonist of the amylin receptor blocks beta-amyloid toxicity in rat cholinergic basal forebrain neurons. *J Neurosci* 24:5579-5584.

- Jhamandas JH, Harris KH, MacTavish D, Jassar BS (2002) Novel excitatory actions of galanin on rat cholinergic basal forebrain neurons: implications for its role in Alzheimer's disease. *J Neurophysiol* 87:696-704.
- Kalen P, Wiklund L (1989) Projections from the medial septum and diagonal band of Broca to the dorsal and central superior raphe nuclei: a non-cholinergic pathway. *Exp Brain Res* 75:401-416.
- Kaltschmidt B, Uherek M, Volk B, Baeuerle PA, Kaltschmidt C (1997) Transcription factor NF-kappa B is activated in primary neurons by amyloid beta peptides and in neurons surrounding early plaques from patients with Alzheimer disease. *Proceedings Of The National Academy Of Sciences Of The United States Of America* 94:2642-2647.
- Kang J, Lemaire HG, Unterbeck A, Salbaum JM, Masters CL, Grzeschik KH, Multhaup G, Beyreuther K, Muller-Hill B (1987) The precursor of Alzheimer's disease amyloid A4 protein resembles a cell-surface receptor. *Nature* 325:733-736.
- Kar S (2005) In: Unpublished observation.
- Kemp IR, Kaada BR (1975) The relation of hippocampal theta activity to arousal, attentive behaviour and somato-motor movements in unrestrained cats. *Brain Res* 95:323-342.
- Kiss J, Patel AJ, Freund TF (1990) Distribution of septohippocampal neurons containing parvalbumin or choline acetyltransferase in the rat brain. *J Comp Neurol* 298:362-372.

- Kitaguchi N, Takahashi Y, Tokushima Y, Shiojiri S, Ito H (1988) Novel precursor of Alzheimer's disease amyloid protein shows protease inhibitory activity. *Nature* 331:530-532.
- Kitchener PD, Diamond J (1993) Distribution and colocalization of choline acetyltransferase immunoreactivity and NADPH diaphorase reactivity in neurons within the medial septum and diagonal band of Broca in the rat basal forebrain. *J Comp Neurol* 335:1-15.
- Kovacs GL, Versteeg DH (1993) Neurohypophysial peptides and brain neurochemistry. *Ann N Y Acad Sci* 689:309-319.
- Lambert MP, Barlow AK, Chromy BA, Edwards C, Freed R, Liosatos M, Morgan TE, Rozovsky I, Trommer B, Viola KL, Wals P, Zhang C, Finch CE, Krafft GA, Klein WL (1998) Diffusible, nonfibrillar ligands derived from A beta(1-42) are potent central nervous system neurotoxins. *Proceedings Of The National Academy Of Sciences Of The United States Of America* 95:6448-6453.
- Lammich S, Kojro E, Postina R, Gilbert S, Pfeiffer R, Jasionowski M, Haass C, Fahrenholz F (1999) Constitutive and regulated alpha-secretase cleavage of Alzheimer's amyloid precursor protein by a disintegrin metalloprotease. *Proc Natl Acad Sci U S A* 96:3922-3927.
- Lax P, Fucile S, Eusebi F (2002) Ca(2+) permeability of human heteromeric nAChRs expressed by transfection in human cells. *Cell Calcium* 32:53-58.
- Lee HS, Kim MA, Waterhouse BD (2005) Retrograde double-labeling study of common afferent projections to the dorsal raphe and the nuclear core of the locus coeruleus in the rat. *J Comp Neurol* 481:179-193.

- Leung LW, Lopes da Silva FH, Wadman WJ (1982) Spectral characteristics of the hippocampal EEG in the freely moving rat. *Electroencephalogr Clin Neurophysiol* 54:203-219.
- Levin ED, Simon BB (1998) Nicotinic acetylcholine involvement in cognitive function in animals. *Psychopharmacology* 138:217-230.
- Levy-Lahad E, Wasco W, Poorkaj P, Romano DM, Oshima J, Pettingell WH, Yu CE, Jondro PD, Schmidt SD, Wang K, Crowley AC, Fu YH, Guenette SY, Galas D, Nemens E, Wijsman EM, Bird TD, Schellenberg GD, Tanzi RE (1995) Candidate Gene For The Chromosome-1 Familial Alzheimers-Disease Locus. *Science* 269:973-977.
- Liu QS, Kawai H, Berg DK (2001) beta-Amyloid peptide blocks the response of alpha 7-containing nicotinic receptors on hippocampal neurons. *Proceedings Of The National Academy Of Sciences Of The United States Of America* 98:4734-4739.
- Luskin MB, Price JL (1982) The distribution of axon collaterals from the olfactory bulb and the nucleus of the horizontal limb of the diagonal band to the olfactory cortex, demonstrated by double retrograde labeling techniques. *J Comp Neurol* 209:249-263.
- Manns ID, Mainville L, Jones BE (2001) Evidence for glutamate, in addition to acetylcholine and GABA, neurotransmitter synthesis in basal forebrain neurons projecting to the entorhinal cortex. *Neuroscience* 107:249-263.
- Manseau F, Danik M, Williams S (2005) A functional glutamatergic neurone network in the medial septum and diagonal band area. *J Physiol* 566:865-884.

- Markram H, Segal M (1990) Electrophysiological characteristics of cholinergic and non-cholinergic neurons in the rat medial septum-diagonal band complex. *Brain Res* 513:171-174.
- Marston HM, West HL, Wilkinson LS, Everitt BJ, Robbins TW (1994) Effects of excitotoxic lesions of the septum and vertical limb nucleus of the diagonal band of Broca on conditional visual discrimination: relationship between performance and choline acetyltransferase activity in the cingulate cortex. *J Neurosci* 14:2009-2019.
- Martin-Ruiz CM, Court JA, Molnar E, Lee M, Gotti C, Mamalaki A, Tsouloufis T, Tzartos S, Ballard C, Perry RH, Perry EK (1999) alpha 4 but not alpha 3 and alpha 7 nicotinic acetylcholine receptor subunits are lost from the temporal cortex in Alzheimer' s disease. *Journal Of Neurochemistry* 73:1635-1640.
- Martone ME, Zhang Y, Simpliciano VM, Carragher BO, Ellisman MH (1993) Three-dimensional visualization of the smooth endoplasmic reticulum in Purkinje cell dendrites. *J Neurosci* 13:4636-4646.
- Masters CL, Multhaup G, Simms G, Pottgiesser J, Martins RN, Beyreuther K (1985a) Neuronal origin of a cerebral amyloid: neurofibrillary tangles of Alzheimer's disease contain the same protein as the amyloid of plaque cores and blood vessels. *Embo J* 4:2757-2763.
- Masters CL, Simms G, Weinman NA, Multhaup G, McDonald BL, Beyreuther K (1985b) Amyloid plaque core protein in Alzheimer disease and Down syndrome. *Proc Natl Acad Sci U S A* 82:4245-4249.

- Masters CL, Simms G, Weinman NA, Multhaup G, McDonald BL, Beyreuther K (1985c) Amyloid Plaque Core Protein In Alzheimer-Disease And Down Syndrome. Proceedings Of The National Academy Of Sciences Of The United States Of America 82:4245-4249.
- Mattson MP, Cheng B, Davis D, Bryant K, Lieberburg I, Rydel RE (1992) Beta-Amyloid Peptides Destabilize Calcium Homeostasis And Render Human Cortical-Neurons Vulnerable To Excitotoxicity. Journal Of Neuroscience 12:376-389.
- Mattson MP, Cheng B, Culwell AR, Esch FS, Lieberburg I, Rydel RE (1993) Evidence for excitoprotective and intraneuronal calcium-regulating roles for secreted forms of the beta-amyloid precursor protein. Neuron 10:243-254.
- McGehee DS, Heath MJ, Gelber S, Devay P, Role LW (1995) Nicotine enhancement of fast excitatory synaptic transmission in CNS by presynaptic receptors. Science 269:1692-1696.
- McKinney M, Coyle JT, Hedreen JC (1983) Topographic analysis of the innervation of the rat neocortex and hippocampus by the basal forebrain cholinergic system. J Comp Neurol 217:103-121.
- McPherson PS, Kim YK, Valdivia H, Knudson CM, Takekura H, Franziniarmstrong C, Coronado R, Campbell KP (1991) The Brain Ryanodine Receptor - A Caffeine-Sensitive Calcium Release Channel. Neuron 7:17-25.
- Meibach RC, Siegel A (1977) Efferent connections of the septal area in the rat: an analysis utilizing retrograde and anterograde transport methods. Brain Res 119:1-20.

- Melander T, Staines WA, Hokfelt T, Rokaeus A, Eckenstein F, Salvaterra PM, Wainer BH (1985) Galanin-like immunoreactivity in cholinergic neurons of the septum-basal forebrain complex projecting to the hippocampus of the rat. *Brain Res* 360:130-138.
- Mesulam MM, Van Hoesen GW (1976) Acetylcholinesterase-rich projections from the basal forebrain of the rhesus monkey to neocortex. *Brain Res* 109:152-157.
- Mesulam MM, Mufson EJ, Wainer BH, Levey AI (1983a) Central cholinergic pathways in the rat: an overview based on an alternative nomenclature (Ch1-Ch6). *Neuroscience* 10:1185-1201.
- Mesulam MM, Mufson EJ, Levey AI, Wainer BH (1983b) Cholinergic innervation of cortex by the basal forebrain: cytochemistry and cortical connections of the septal area, diagonal band nuclei, nucleus basalis (substantia innominata), and hypothalamus in the rhesus monkey. *J Comp Neurol* 214:170-197.
- Metcalf RH, Boegman RJ, Riopelle RJ, Ludwin SK (1988) The release of endogenous acetylcholine from the medial septum/diagonal band of rat brain. *Neurosci Lett* 93:85-90.
- Milner TA (1991) Cholinergic neurons in the rat septal complex: ultrastructural characterization and synaptic relations with catecholaminergic terminals. *J Comp Neurol* 314:37-54.
- Monmaur P (1981) Phasic hippocampal activity during paradoxical sleep in the rat: selective suppression after diazepam administration. *Experientia* 37:261-262.
- Murchison D, Griffith WH (1995) Low-voltage activated calcium currents increase in basal forebrain neurons from aged rats. *J Neurophysiol* 74:876-887.

- Murchison D, Griffith WH (1996) High-voltage-activated calcium currents in basal forebrain neurons during aging. *J Neurophysiol* 76:158-174.
- Murchison D, Griffith WH (1999) Age-related alterations in caffeine-sensitive calcium stores and mitochondrial buffering in rat basal forebrain. *Cell Calcium* 25:439-452.
- Murchison D, Griffith WH (2000) Mitochondria buffer non-toxic calcium loads and release calcium through the mitochondrial permeability transition pore and sodium/calcium exchanger in rat basal forebrain neurons. *Brain Res* 854:139-151.
- Nagele RG, D'Andrea MR, Anderson WJ, Wang HY (2002) Intracellular accumulation of beta-amyloid(1-42) in neurons is facilitated by the alpha 7 nicotinic acetylcholine receptor in Alzheimer's disease. *Neuroscience* 110:199-211.
- Nakagawa T, Zhu H, Morishima N, Li E, Xu J, Yankner BA, Yuan JY (2000) Caspase-12 mediates endoplasmic-reticulum-specific apoptosis and cytotoxicity by amyloid-beta. *Nature* 403:98-103.
- Naslund J, Haroutunian V, Mohs R, Davis KL, Davies P, Greengard P, Buxbaum JD (2000) Correlation between elevated levels of amyloid beta-peptide in the brain and cognitive decline. *Jama-Journal Of The American Medical Association* 283:1571-1577.
- Nyakas C, Luiten PG, Spencer DG, Traber J (1987) Detailed projection patterns of septal and diagonal band efferents to the hippocampus in the rat with emphasis on innervation of CA1 and dentate gyrus. *Brain Res Bull* 18:533-545.

- Oka H, Yoshida K (1985) Septohippocampal connections to field CA1 of the rat identified with field potential analysis and retrograde labeling by horseradish peroxidase. *Neurosci Lett* 58:19-24.
- Onteniente B, Tago H, Kimura H, Maeda T (1986) Distribution of gamma-aminobutyric acid-immunoreactive neurons in the septal region of the rat brain. *J Comp Neurol* 248:422-430.
- Panula P, Revuelta AV, Cheney DL, Wu JY, Costa E (1984) An immunohistochemical study on the location of GABAergic neurons in rat septum. *J Comp Neurol* 222:69-80.
- Paolini AG, McKenzie JS (1993) Effects of lesions in the horizontal diagonal band nucleus on olfactory habituation in the rat. *Neuroscience* 57:717-724.
- Perry EK, Morris CM, Court JA, Cheng A, Fairbairn AF, McKeith IG, Irving D, Brown A, Perry RH (1995) Alteration In Nicotine Binding-Sites In Parkinsons-Disease, Lewy Body Dementia And Alzheimers-Disease - Possible Index Of Early Neuropathology. *Neuroscience* 64:385-395.
- Petsche H, Stumpf C, Gogolak G (1962) [The significance of the rabbit's septum as a relay station between the midbrain and the hippocampus. I. The control of hippocampus arousal activity by the septum cells.]. *Electroencephalogr Clin Neurophysiol* 14:202-211.
- Pettit DL, Shao Z, Yakel JL (2001) beta-Amyloid(1-42) peptide directly modulates nicotinic receptors in the rat hippocampal slice. *Journal Of Neuroscience* 21.

- Pike CJ, Burdick D, Walencewicz AJ, Glabe CG, Cotman CW (1993) Neurodegeneration Induced By Beta-Amyloid Peptides In Vitro - The Role Of Peptide Assembly State. *Journal Of Neuroscience* 13:1676-1687.
- Poirier J, Davignon J, Bouthillier D, Kogan S, Bertrand P, Gauthier S (1993) Apolipoprotein-E Polymorphism And Alzheimers-Disease. *Lancet* 342:697-699.
- Ponte P, Gonzalez-DeWhitt P, Schilling J, Miller J, Hsu D, Greenberg B, Davis K, Wallace W, Lieberburg I, Fuller F (1988) A new A4 amyloid mRNA contains a domain homologous to serine proteinase inhibitors. *Nature* 331:525-527.
- Prentki M, Biden TJ, Janjic D, Irvine RF, Berridge MJ, Wollheim CB (1984) Rapid mobilization of Ca²⁺ from rat insulinoma microsomes by inositol-1,4,5-trisphosphate. *Nature* 309:562-564.
- Rabizadeh S, Bitler CM, Butcher LL, Bredesen DE (1994) Expression Of The Low-Affinity Nerve Growth-Factor Receptor Enhances Beta-Amyloid Peptide Toxicity. *Proceedings Of The National Academy Of Sciences Of The United States Of America* 91:10703-10706.
- Ramsden M, Henderson Z, Pearson HA (2002) Modulation of Ca²⁺ channel currents in primary cultures of rat cortical neurones by amyloid beta protein (1-40) is dependent on solubility status. *Brain Research* 956:254-261.
- Rapier C, Lunt GG, Wonnacott S (1990) Nicotinic modulation of [3H]dopamine release from striatal synaptosomes: pharmacological characterisation. *J Neurochem* 54:937-945.
- Rhee SK, Quist AP, Lal R (1998) Amyloid beta protein-(1-42) forms calcium-permeable, Zn²⁺-sensitive channel. *J Biol Chem* 273:13379-13382.

- Ridley RM, Samson NA, Baker HF, Johnson JA (1988) Visuospatial learning impairment following lesion of the cholinergic projection to the hippocampus. *Brain Res* 456:71-87.
- Rinne JO, Myllykyla T, Lonnberg P, Marjamaki P (1991) A postmortem study of brain nicotinic receptors in Parkinson's and Alzheimer's disease. *Brain Res* 547:167-170.
- Roman FS, Simonetto I, Soumireu-Mourat B (1993) Learning and memory of odor-reward association: selective impairment following horizontal diagonal band lesions. *Behav Neurosci* 107:72-81.
- Sacaan AI, Dunlop JL, Lloyd GK (1995) Pharmacological characterization of neuronal acetylcholine gated ion channel receptor-mediated hippocampal norepinephrine and striatal dopamine release from rat brain slices. *J Pharmacol Exp Ther* 274:224-230.
- Sandbrink R, Masters CL, Beyreuther K (1996) APP gene family - Alternative splicing generates functionally related isoforms. In: *Neurobiology Of Alzheimer's Disease*, pp 281-287.
- Saper CB, Chelimsky TC (1984) A cytoarchitectonic and histochemical study of nucleus basalis and associated cell groups in the normal human brain. *Neuroscience* 13:1023-1037.
- Saper CB, German DC, White CL, 3rd (1985) Neuronal pathology in the nucleus basalis and associated cell groups in senile dementia of the Alzheimer's type: possible role in cell loss. *Neurology* 35:1089-1095.

- Saunders AM, Strittmatter WJ, Schmechel D, Georgehyslop PHS, Pericakvance MA, Joo SH, Rosi BL, Gusella JF, Crappermaclachlan DR, Alberts MJ, Hulette C, Crain B, Goldgaber D, Roses AD (1993) Association Of Apolipoprotein-E Allele Epsilon-4 With Late-Onset Familial And Sporadic Alzheimers-Disease. *Neurology* 43:1467-1472.
- Schoepfer R, Conroy WG, Whiting P, Gore M, Lindstrom J (1990) Brain alpha-bungarotoxin binding protein cDNAs and MAbs reveal subtypes of this branch of the ligand-gated ion channel gene superfamily. *Neuron* 5:35-48.
- Selkoe DJ (2001) Alzheimer's disease: genes, proteins, and therapy. *Physiol Rev* 81:741-766.
- Selkoe DJ (2002a) Deciphering the genesis and fate of amyloid beta-protein yields novel therapies for Alzheimer disease. *J Clin Invest* 110:1375-1381.
- Selkoe DJ (2002b) Alzheimer's disease is a synaptic failure. *Science* 298:789-791.
- Semba K, Komisaruk BR (1984) Neural substrates of two different rhythmical vibrissal movements in the rat. *Neuroscience* 12:761-774.
- Semba K, Reiner PB, McGeer EG, Fibiger HC (1988) Brainstem afferents to the magnocellular basal forebrain studied by axonal transport, immunohistochemistry, and electrophysiology in the rat. *J Comp Neurol* 267:433-453.
- Senut MC, Menetrey D, Lamour Y (1989) Cholinergic and peptidergic projections from the medial septum and the nucleus of the diagonal band of Broca to dorsal hippocampus, cingulate cortex and olfactory bulb: a combined wheatgerm

agglutinin-apohorseradish peroxidase-gold immunohistochemical study.

Neuroscience 30:385-403.

Seubert P, Vigo-Pelfrey C, Esch F, Lee M, Dovey H, Davis D, Sinha S, Schlossmacher M, Whaley J, Swindlehurst C, et al. (1992) Isolation and quantification of soluble Alzheimer's beta-peptide from biological fluids. *Nature* 359:325-327.

Sharples CG, Kaiser S, Soliakov L, Marks MJ, Collins AC, Washburn M, Wright E, Spencer JA, Gallagher T, Whiteaker P, Wonnacott S (2000) UB-165: a novel nicotinic agonist with subtype selectivity implicates the alpha4beta2* subtype in the modulation of dopamine release from rat striatal synaptosomes. *J Neurosci* 20:2783-2791.

Shaw S, Bencherif M, Marrero MB (2002) Janus kinase 2, an early target of alpha 7 nicotinic acetylcholine receptor-mediated neuroprotection against Abeta-(1-42) amyloid. *J Biol Chem* 277:44920-44924.

Sherrington R, Rogaev EI, Liang Y, Rogaeva EA, Levesque G, Ikeda M, Chi H, Lin C, Li G, Holman K, Tsuda T, Mar L, Foncin JF, Bruni AC, Montesi MP, Sorbi S, Rainero I, Pinessi L, Nee L, Chumakov I, Pollen D, Brookes A, Sanseau P, Polinsky RJ, Wasco W, Dasilva HAR, Haines JL, Pericakvance MA, Tanzi RE, Roses AD, Fraser PE, Rommens JM, Stgeorgehslop PH (1995) Cloning Of A Gene Bearing Missense Mutations In Early-Onset Familial Alzheimers-Disease. *Nature* 375:754-760.

Sisodia SS, Koo EH, Beyreuther K, Unterbeck A, Price DL (1990) Evidence that beta-amyloid protein in Alzheimer's disease is not derived by normal processing. *Science* 248:492-495.

- Skofitsch G, Jacobowitz DM (1985) Immunohistochemical mapping of galanin-like neurons in the rat central nervous system. *Peptides* 6:509-546.
- Smith TD, Gallagher M, Leslie FM (1995) Cholinergic binding sites in rat brain: analysis by age and cognitive status. *Neurobiol Aging* 16:161-173.
- Sofroniew MV (1985) Vasopressin- and neurophysin-immunoreactive neurons in the septal region, medial amygdala and locus coeruleus in colchicine-treated rats. *Neuroscience* 15:347-358.
- Sotty F, Danik M, Manseau F, Laplante F, Quirion R, Williams S (2003) Distinct electrophysiological properties of glutamatergic, cholinergic and GABAergic rat septohippocampal neurons: novel implications for hippocampal rhythmicity. *Journal Of Physiology-London* 551:927-943.
- Steckler T, Keith AB, Wiley RG, Sahgal A (1995) Cholinergic lesions by 192 IgG-saporin and short-term recognition memory: role of the septohippocampal projection. *Neuroscience* 66:101-114.
- StGeorgehyslop P, Haines J, Rogaev E, Mortilla M, Vaula G, Pericakvance M, Foncin JF, Montesi M, Bruni A, Sorbi S, Rainero I, Pinessi L, Pollen D, Polinsky R, Nee L, Kennedy J, Macciardi F, Rogaeva E, Liang Y, Alexandrova N, Lukiw W, Schlumpf K, Tanzi R, Tsuda T, Farrer L, Cantu JM, Duara R, Amaducci L, Bergamini L, Gusella J, Roses A, McLachlan DC (1992) Genetic-Evidence For A Novel Familial Alzheimers-Disease Locus On Chromosome-14. *Nature Genetics* 2:330-334.

- Stock G, Rupprecht U, Stumpf H, Schlor KH (1981) Cardiovascular changes during arousal elicited by stimulation of amygdala, hypothalamus and locus coeruleus. *J Auton Nerv Syst* 3:503-510.
- Stout AK, Raphael HM, Kanterewicz BI, Klann E, Reynolds IJ (1998) Glutamate-induced neuron death requires mitochondrial calcium uptake. *Nat Neurosci* 1:366-373.
- Swanson LW, Cowan WM (1979) The connections of the septal region in the rat. *J Comp Neurol* 186:621-655.
- Sweeney JE, Lamour Y, Bassant MH (1992) Arousal-dependent properties of medial septal neurons in the unanesthetized rat. *Neuroscience* 48:353-362.
- Tang K, Wu H, Mahata SK, O'Connor DT (1998) A crucial role for the mitogen-activated protein kinase pathway in nicotinic cholinergic signaling to secretory protein transcription in pheochromocytoma cells. *Mol Pharmacol* 54:59-69.
- Tanzi RE, McClatchey AI, Lamperti ED, Villa-Komaroff L, Gusella JF, Neve RL (1988) Protease inhibitor domain encoded by an amyloid protein precursor mRNA associated with Alzheimer's disease. *Nature* 331:528-530.
- Terry R (1999) *The neuropathology of Alzheimer disease and the structural basis of its cognitive alterations*. Philadelphia: Lippincott, Williams and Wilkins.
- Tohgi H, Utsugisawa K, Yoshimura M, Nagane Y, Mihara M (1998) Age-related changes in nicotinic acetylcholine receptor subunits alpha4 and beta2 messenger RNA expression in postmortem human frontal cortex and hippocampus. *Neurosci Lett* 245:139-142.

- Tomizawa M, Casida JE (2002) Desnitro-imidacloprid activates the extracellular signal-regulated kinase cascade via the nicotinic receptor and intracellular calcium mobilization in N1E-115 cells. *Toxicol Appl Pharmacol* 184:180-186.
- Toth K, Borhegyi Z, Freund TF (1993) Postsynaptic targets of GABAergic hippocampal neurons in the medial septum-diagonal band of Broca complex. *J Neurosci* 13:3712-3724.
- Vassar R, Bennett BD, Babu-Khan S, Kahn S, Mendiaz EA, Denis P, Teplow DB, Ross S, Amarante P, Loeloff R, Luo Y, Fisher S, Fuller J, Edenson S, Lile J, Jarosinski MA, Biere AL, Curran E, Burgess T, Louis JC, Collins F, Treanor J, Rogers G, Citron M (1999) Beta-secretase cleavage of Alzheimer's amyloid precursor protein by the transmembrane aspartic protease BACE. *Science* 286:735-741.
- Verkhatsky A (2005) Physiology and pathophysiology of the calcium store in the endoplasmic reticulum of neurons. *Physiol Rev* 85:201-279.
- Vinogradova OS (1995) Expression, control, and probable functional significance of the neuronal theta-rhythm. *Prog Neurobiol* 45:523-583.
- Waite JJ, Chen AD, Wardlow ML, Thal LJ (1994) Behavioral and biochemical consequences of combined lesions of the medial septum/diagonal band and nucleus basalis in the rat when ibotenic acid, quisqualic acid, and AMPA are used. *Exp Neurol* 130:214-229.
- Walsh DM, Lomakin A, Benedek GB, Condron MM, Teplow DB (1997) Amyloid beta-protein fibrillogenesis - Detection of a protofibrillar intermediate. *Journal Of Biological Chemistry* 272:22364-22372.

- Walsh DM, Klyubin I, Fadeeva JV, Cullen WK, Anwyl R, Wolfe MS, Rowan MJ, Selkoe DJ (2002) Naturally secreted oligomers of amyloid beta protein potently inhibit hippocampal long-term potentiation in vivo. *Nature* 416:535-539.
- Wang GK, Schmidt J (1976) Receptors for alpha-bungarotoxin in the developing visual system of the chick. *Brain Res* 114:524-529.
- Wang GK, Molinaro S, Schmidt J (1978) Ligand responses of alpha-bungarotoxin binding sites from skeletal muscle and optic lobe of the chick. *J Biol Chem* 253:8507-8512.
- Wang HY, Lee DHS, Davis CB, Shank RP (2000a) Amyloid peptide A beta(1-42) binds selectively and with picomolar affinity to alpha 7 nicotinic acetylcholine receptors. *Journal Of Neurochemistry* 75:1155-1161.
- Wang HY, Lee DH, D'Andrea MR, Peterson PA, Shank RP, Reitz AB (2000b) beta-Amyloid(1-42) binds to alpha7 nicotinic acetylcholine receptor with high affinity. Implications for Alzheimer's disease pathology. *J Biol Chem* 275:5626-5632.
- Warpman U, Nordberg A (1995) Epibatidine and ABT 418 reveal selective losses of alpha 4 beta 2 nicotinic receptors in Alzheimer brains. *Neuroreport* 6:2419-2423.
- Westfall TC (1974) Effect of nicotine and other drugs on the release of 3H-norepinephrine and 3H-dopamine from rat brain slices. *Neuropharmacology* 13:693-700.
- Wevers A, Monteggia L, Nowacki S, Bloch W, Schutz U, Lindstrom J, Pereira EF, Eisenberg H, Giacobini E, de Vos RA, Steur EN, Maelicke A, Albuquerque EX, Schroder H (1999) Expression of nicotinic acetylcholine receptor subunits in the cerebral cortex in Alzheimer's disease: histotopographical correlation with

- amyloid plaques and hyperphosphorylated-tau protein. *Eur J Neurosci* 11:2551-2565.
- White RJ, Reynolds IJ (1996) Mitochondrial depolarization in glutamate-stimulated neurons: an early signal specific to excitotoxin exposure. *J Neurosci* 16:5688-5697.
- Whitehouse PJ, Price DL, Struble RG, Clark AW, Coyle JT, Delon MR (1982) Alzheimer's disease and senile dementia: loss of neurons in the basal forebrain. *Science* 215:1237-1239.
- Whitehouse PJ, Martino AM, Wagster MV, Price DL, Mayeux R, Atack JR, Kellar KJ (1988) Reductions in [³H]nicotinic acetylcholine binding in Alzheimer's disease and Parkinson's disease: an autoradiographic study. *Neurology* 38:720-723.
- Wu J, Kuo YP, George AA, Xu L, Hu J, Lukas RJ (2004) beta-Amyloid directly inhibits human alpha4beta2-nicotinic acetylcholine receptors heterologously expressed in human SH-EP1 cells. *J Biol Chem* 279:37842-37851.
- Wu M, Shanabrough M, Leranath C, Alreja M (2000) Cholinergic excitation of septohippocampal GABA but not cholinergic neurons: implications for learning and memory. *J Neurosci* 20:3900-3908.
- Wu M, Hajszan T, Leranath C, Alreja M (2003) Nicotine recruits a local glutamatergic circuit to excite septohippocampal GABAergic neurons. *Eur J Neurosci* 18:1155-1168.
- Yan SD, Chen X, Fu J, Chen M, Zhu HJ, Roher A, Slattery T, Zhao L, Nagashima M, Morser J, Migheli A, Nawroth P, Stern D, Schmidt AM (1996) RAGE and amyloid-beta peptide neurotoxicity in Alzheimer's disease. *Nature* 382:685-691.

- Zaborszky L, Cullinan WE (1996) Direct catecholaminergic-cholinergic interactions in the basal forebrain.1. Dopamine-beta-hydroxylase- and tyrosine hydroxylase input to cholinergic neurons. *Journal Of Comparative Neurology* 374:535-554.
- Zaborszky L, Gaykema RP, Swanson DJ, Cullinan WE (1997) Cortical input to the basal forebrain. *Neuroscience* 79:1051-1078.
- Zoli M, Le Novere N, Hill JA, Jr., Changeux JP (1995) Developmental regulation of nicotinic ACh receptor subunit mRNAs in the rat central and peripheral nervous systems. *J Neurosci* 15:1912-1939.
- Zoli M, Moretti M, Zanardi A, McIntosh JM, Clementi F, Gotti C (2002) Identification of the nicotinic receptor subtypes expressed on dopaminergic terminals in the rat striatum. *J Neurosci* 22:8785-8789.

Chapter 2

β -Amyloid Enhances Intracellular Calcium Rises Mediated by Intracellular Calcium Stores and Nicotinic Receptors in Acutely Dissociated Rat Basal Forebrain Neurons

INTRODUCTION

Alzheimer's disease (AD) is a devastating neurodegenerative disease that affects elderly patients and is characterized by a progressive decline in memory and other cognitive functions such as language and perception (McKhann et al., 1984). Key neuropathological findings in patients with AD include extracellular neuritic plaques composed of β -amyloid peptide ($A\beta$), neurofibrillary tangles composed of abnormally phosphorylated tau protein, and loss of cholinergic neurons of the basal forebrain (Yankner, 1996; Mayeux and Sano, 1999). $A\beta$, a 39-43 amino acid peptide, plays an important role in the AD pathophysiological process as total $A\beta$ content in the brain correlates with cognitive decline (Naslund et al., 2000). Although the precise mechanism of $A\beta$ toxicity has not been established, $A\beta$ and other peptide fragments derived from amyloid precursor protein may destabilize intracellular calcium (Ca^{2+}) homeostasis via modulation of specific ion channels (Mattson et al., 1992; Fraser et al., 1997). An intriguing class of candidate receptors for $A\beta$ -mediated actions is the neuronal nicotinic acetylcholine receptors (nAChRs), a family of ligand-gated cation permeable channels with multiple subtypes that are widely distributed in the human brain. nAChRs are involved in a number of physiological and behavioral processes and post-mortem studies of AD individuals demonstrate that reduced numbers of nAChRs in brain regions are associated with AD (Nordberg and Winblad, 1986; Whitehouse et al., 1988; Paterson and Nordberg, 2000).

$A\beta_{1-42}$ binds to $\alpha 7$ and $\alpha 4\beta 2$ subtypes of nAChRs with high affinity in the picomolar and nanomolar range respectively (Wang et al., 2000b; Wang et al., 2000a). In the rat basal forebrain, $A\beta$ directly activates $\alpha 4\beta 2$ or $\alpha 3$ postsynaptic nAChRs in the

diagonal band of Broca (DBB), a basal forebrain nucleus (Fu and Jhamandas, 2003). Based on previous data this A β -non- α 7 nAChR interaction appears to occur on cholinergic rather than GABAergic basal forebrain neurons (Jhamandas et al., 2001).

Interactions of amyloid peptides with cholinergic neurons are considered to be a basis for the modified cholinergic tone observed in AD (Auld et al., 2002). Both α 7 and non- α 7 nAChRs are abundantly located within the rat basal forebrain cholinergic system (Dominguez del Toro et al., 1994; Rogers et al., 1998). Although the exact functional consequences of A β interaction with nAChRs remain unknown, nAChRs regulate many neuronal processes. They play an important role in neurotransmission, synaptic modulation, and second messenger signaling via intracellular Ca²⁺ stores (Tsuneki et al., 2000; Dajas-Bailador et al., 2002; Henderson et al., 2005). Thus, A β disruption of synaptic transmission and Ca²⁺ homeostasis could be mediated via an interaction with nAChRs (Mattson et al., 1992; Kamenetz et al., 2003).

The main sources of intracellular Ca²⁺ stores in the DBB are the endoplasmic reticulum (ER) and the mitochondria (Murchison and Griffith, 2000). These stores have been shown to be important in both the release and buffering of Ca²⁺ (Murchison and Griffith, 2000). Accumulation of Ca²⁺ in the mitochondria and endoplasmic reticulum has been associated with apoptotic cell death in pathophysiological conditions and A β ₁₋₄₂ has been shown to mediate apoptosis via an ER-dependent pathway (White and Reynolds, 1997; Nakagawa et al., 2000). Thus examining Ca²⁺ regulation in these organelles is important to understanding A β neurotoxicity.

I examined the effect of A β ₁₋₄₂ modulation of nAChRs mediated Ca²⁺ rises and the associated intracellular Ca²⁺ stores by performing ratiometric Ca²⁺ imaging in acutely

dissociated DBB neurons. nAChR activation was demonstrated to be associated with activation of intracellular Ca^{2+} release. I also demonstrated that $\text{A}\beta_{1-42}$ modulates the repeated activation of nAChRs by facilitating the release of Ca^{2+} from intracellular Ca^{2+} stores through overfilling of thapsigargin-sensitive stores.

MATERIALS AND METHODS

Dissociation Procedures

Ratiometric fura-2 imaging was performed on acutely dissociated rat DBB neurons. The protocol for preparation of tissue was reviewed and approved by the University of Alberta Health Sciences Laboratory Animal Services (Protocol number 154/04/05). Details of the procedure for acute dissociation of neurons from the DBB have been previously described (Chin et al., 2002). Briefly, brains were quickly removed from decapitated male Sprague Dawley rats (21-25 day postnatal) and placed in cold artificial cerebrospinal fluid (ACSF) that contained (in mM) 140 NaCl, 2.5 KCl, 1.4 CaCl_2 , 5 MgCl_2 , 10 HEPES, and 33 D-glucose (pH 7.4). Brain slices (350 μm thick) were cut on a vibratome, and the area containing the horizontal and vertical limbs of DBB was dissected out. Acutely dissociated neurons were prepared by the enzymatic treatment of slices with trypsin (0.65 mg/ml) at 30 °C for 16 min, followed by mechanical trituration for dispersion of individual cells. Cells were then plated on poly-L-lysine (0.005% wt/vol)-coated cover slips and viewed under an inverted microscope (Zeiss Axiovert 35). Neurons were identified based upon previously established visual criteria (Chin et al., 2002). Basal forebrain neurons could be differentiated from glia based upon the presence of truncated dendrites. In addition, neurons were larger than glia. All solutions were kept oxygenated by continuous bubbling with 100% oxygen.

Calcium Imaging

Neurons were incubated with 10 μ M fura-2 AM (Molecular Probes, Eugene, Oregon, USA) in 1 ml of oxygenated ACSF for 20 min and perfused for 20 min prior to imaging. All experiments were performed at room temperature (20-22 °C). Measurements were collected every 8 seconds for 2 minutes to establish a baseline. During focal application of drugs, the image collection rate was increased to 1 Hz for 15 seconds and then returned to its baseline of rate of 1 image every 8 seconds. Fura-2 loaded neurons were exposed to 340 and 380 nm light emitted by a Polychrome IV monochromator (T.I.L.L. Photonics GMBH, Pleasanton, CA, USA) controlled by Metafluor 5.03 software (Universal Imaging Corporation, Downingtown, PA, USA) and emitted fluorescence was collected by a CoolSnap HQ CCD camera (Roper Scientific, Duluth, GA, USA). A three-point calibration was performed on acutely dissociated cells loaded with 100 μ M fura-2 and EGTA/ Ca^{2+} buffered solutions via patch pipette with a final concentration of 0, 224 nM and 5 mM of Ca^{2+} (Grynkiewicz et al., 1985). $[\text{Ca}^{2+}]_i$ for the pipette solutions was determined with MaxChelator 2.32 (Stanford, Chris Patton).

The values obtained from the calibrations were used to convert background subtracted experimental fluorescent intensity ratios into $[\text{Ca}^{2+}]$ over the physiological range of $[\text{Ca}^{2+}]_i$ using the following equation: $[\text{Ca}^{2+}]_i = K_d(R-R_{\min})/(R_{\max}-R)$, where K_d is the dissociation constant for fura-2, R_{\min} is the background subtracted 340/380 ratio when $[\text{Ca}^{2+}]_i$ was 0, R_{\max} is the background subtracted 340/380 ratio when $[\text{Ca}^{2+}]_i$ was 5 mM and R is the experimentally measured background subtracted 340/380 ratio (Grynkiewicz et al., 1985).

Immunohistochemistry

DBB cells were chemically and mechanically isolated from 14-21 day old Sprague Dawley rats as described above. The DBB cell suspension was centrifuged for 5 minutes at 500 rpm onto glass slides. Cells were not fixed. Cells were first blocked with 1% bovine serum albumin (Sigma, Oakville, CA) for 1 hour and then incubated overnight with primary rabbit antibodies for vesicular choline acetyltransferase (vChAT: 1:2500 dilution) $\alpha 4$ (1:250 dilution) or $\alpha 7$ (1:500 dilution; Santa Cruz Biotechnology Inc., Santa Cruz, USA) and then washed 3 times in 0.1 M phosphate-buffered saline (PBS), pH 7.2. Cells were then incubated with biotinylated goat anti-rabbit antibodies for 1.5 hours and washed again with PBS. Next, cells were incubated with the fluorescent dye Alexa 546 streptavidin to yield red fluorescence (1:400 dilution Molecular Probes, Eugene, Oregon). Cells were then washed 3 times in PBS and incubated with 1% BSA then incubated overnight in a different primary antibody. The following day they were washed 3 times in PBS and incubated with Alexa 488 chicken anti-rabbit antibody to green fluorescence (1:400 dilution) and washed 3 times in PBS. Slides were allowed to air dry and dehydrated by ethanol and xylene before being coverslipped with Cytoseal (Canada Wide Scientific, Ottawa, ON). Slides were examined with a Zeiss Axioplan 2 at 400x magnification and images captured with Zeiss AxioCam MRc camera and MGrab software. For each colocalization study, regions of 1 mm² were randomly counted on 3 slides. Confocal images were collected at 900x magnification on an Olympus FV1000 (Olympus, Canada) and optimized for brightness and contrast in Corel Draw 9 (Corel, Canada).

Drugs

Drugs were applied by a 4-channel gravity feed focal applicator controlled by a Master-8 pulse generator. All chemicals were purchased from Sigma except for the following chemicals. A β_{1-42} and reverse peptide A β_{42-1} , which were purchased from American Peptide Co. (Sunnyvale, USA) and 7-chloro-5-(2-chlorophenyl)-1,5-dihydro-4,1-benzothiazepin-2(3H)-one (CGP-37157) was purchased from Calbiochem (San Diego, USA). A β_{1-42} was prepared from aliquots stored at -80°C and used on the day of the experiments. Nicotine was prepared from 100 mM nicotine in ACSF stock solution stored in the dark at room temperature on the day of the experiment. Nicotine was applied via focal applicator. During Ca^{2+} imaging experiments, A β_{1-42} was applied both through the focal applicator and in the bath solution.

Statistical Analysis

[Ca^{2+}]_i measurements under control conditions and during drug applications were compared using Student's paired t-test to detect differences between treatments.

RESULTS

Nicotine-Evoked Ca^{2+} Responses in DBB Neurons: Modulation by A β

The average basal [Ca^{2+}]_i for all cells measured was 99.4 ± 5.5 nM (n = 115). Doses and duration of applications of nicotine were examined to establish an optimal dose that produced a reproducible response of a sufficient amplitude. Figure 2-1A shows a DBB neuron exposed to 10 and 20 mM nicotine for 5 s and Figure 2-1B shows the dose-response relationship between 5 s applications of varying concentrations of nicotine and the corresponding rise in [Ca^{2+}]_i. Nicotine has previously been shown to desensitize receptor subtypes, especially the $\alpha 7$ nAChR, however we did not observe desensitization

in our experiments. Control experiments were performed to exclude the contribution of osmolarity-related changes in $[Ca^{2+}]_i$ and to ensure that nicotine applications did not cross-react with muscarinic receptors to alter $[Ca^{2+}]_i$. Neither focal application of iso-osmotic sucrose solutions nor of 500 μ M muscarine elicited any change in $[Ca^{2+}]_i$ (Figure 2-2A, B respectively).

Next, the effect of soluble $A\beta_{1-42}$ on the focal application of nicotine was examined. In 16 of 18 cells examined, 100 nM $A\beta_{1-42}$ significantly potentiated nicotinic mediated $[Ca^{2+}]_i$ rises ($p < 0.05$; Figure 2-3A). Control nicotinic responses were 63 ± 12 nM, while responses in the presence of 100 nM $A\beta_{1-42}$ increased to 137 ± 25 nM representing a $98 \pm 12\%$ increase over controls. The potentiation of nicotinic responses by $A\beta_{1-42}$ was maintained during washout of $A\beta_{1-42}$ and was dose dependent (Figure 2-4). The $A\beta_{1-42}$ potentiation of the nicotine evoked $[Ca^{2+}]_i$ had an EC50 of 10 nM when fitted with a sigmoidal dose-response curve, although the maximum response could be at a higher dose. We observed no significant change in nicotine mediated $[Ca^{2+}]_i$ rises in response to $A\beta_{1-42}$ in 2 of 18 cells (Figure 2-3B). 100 nM $A\beta_{1-42}$ also caused a rise in baseline $[Ca^{2+}]_i$ of 54 ± 14 nM when stimulated with nicotine. Application of 2 concentrations of $A\beta_{1-42}$ alone either via rapid 5 s focal application or by bath application (100 nM, $n = 6$ and 1 μ M $n = 5$; $p > 0.05$) did not change basal $[Ca^{2+}]_i$ (Figure 2-5A, B). In the presence of the reverse peptide, $A\beta_{42-1}$, neither nicotine-evoked $[Ca^{2+}]_i$ nor baseline $[Ca^{2+}]_i$ changed significantly ($p < 0.05$; $n = 5$; data not shown).

Aβ Potentiation of nAChR-Induced Elevation of $[Ca^{2+}]_i$ is Not Mediated by Voltage Gated Ca^{2+} Channels

Previous studies have reported that Aβ may modify Ca^{2+} entry via modulation of Ca^{2+} channels (Silei et al., 1999; Ramsden et al., 2002). Therefore, in the next set of experiments, I bypassed the nAChRs by applying depolarizing concentrations of KCl at doses evoking equivalent rises in $[Ca^{2+}]_i$ as nicotine. A dose-response relationship for KCl-induced elevations in $[Ca^{2+}]_i$ was established and on the basis of this, 1 s applications of 8 mM KCl were used to depolarize the cell. Application of 8 mM KCl yielded an average rise of $[Ca^{2+}]_i$ of 53 ± 10 nM ($n = 7$) which is similar to that evoked by nicotine (see Figure 2-3). Concomitant application of Aβ₁₋₄₂, with repeated challenges by 8 mM KCl resulted in no significant change in KCl-induced $[Ca^{2+}]_i$ elevation (59 ± 13 nM; $n = 5$; $p > 0.05$; Figure 2-6).

Aβ Potentiates Release of Ca^{2+} from Intracellular Ca^{2+} Stores

The increases of $[Ca^{2+}]_i$ evoked by nicotine may originate from extracellular Ca^{2+} entry via the nAChRs and/or from the release of intracellular Ca^{2+} from stores (Tsuneki et al., 2000; Dajas-Bailador et al., 2002). Thus in the next set of experiments, I examined the source of the nicotine-mediated $[Ca^{2+}]_i$ rise. When cells were perfused with 0 Ca^{2+} /200 μM EGTA, there was a $79.4 \pm 5.5\%$ reduction in the nicotinic response ($p < 0.05$; $n = 7$; Figure 2-7A), suggesting that the majority of the nicotine-mediated rise in $[Ca^{2+}]_i$ is mediated by extracellular Ca^{2+} entry. This also demonstrates that there is an intracellular Ca^{2+} component to the nicotine-mediated response. Application of 10 μM of di-(tert-butyl)-1,4-benzohydroquinone (BHQ), a SERCA inhibitor, in perfusing solution containing 0 Ca^{2+} /1 mM EGTA resulted in an initial rise in $[Ca^{2+}]_i$ due to a transient net

efflux of Ca^{2+} from intracellular stores and then completely abolished the nicotine-mediated $[\text{Ca}^{2+}]_i$ rise as the intracellular stores are depleted. In fact, at times greater than 1500 s in Figure 2-7B, each application of nicotine caused a small decrease in $[\text{Ca}^{2+}]_i$, which may be due to Ca^{2+} efflux via nAChRs due to the reversal of the Ca^{2+} concentration gradient ($p < 0.05$; $n = 7$; Figure 2-7B). In order to test this possibility I performed the same experiment in the presence of non- $\alpha 7$ nAChR antagonist dihydro-beta-erythroidine (5 μM DH β E) and the $\alpha 7$ nAChR antagonist α -bungarotoxin (α BTX). Application of nAChR antagonists did not block the small decrease in $[\text{Ca}^{2+}]_i$ observed with nicotine in the presence of BHQ and 0 Ca^{2+} /1 mM EGTA (data not shown). Another possible explanation may be due to the fluorescent properties of nicotine. I observed that focal application of nicotine increased background 340 nm and 380 nm fluorescence, which resulted in an overall decrease in the background ratio. This is unlikely an osmotic effect as previous applications of iso-osmotic did not result in any deflection of the baseline $[\text{Ca}^{2+}]_i$.

Previously, nAChR receptors have been shown to be linked to intracellular Ca^{2+} stores via the PLC pathway in SH-SY5Y cells (Dajas-Bailador et al., 2002). In order to determine mechanisms underlying nAChR coupling to intracellular Ca^{2+} stores in DBB neurons, I examined the effect of xestospongins on nAChR responses. Xestospongins have been reported to be antagonists of IP_3 -mediated (inositol 1,4,5-triphosphate) Ca^{2+} release (Reviewed in Verkhratsky, 2005, See Discussion). In the presence of 5 μM xestospongins, nicotine-mediated $[\text{Ca}^{2+}]_i$ rises were reduced from 82.9 ± 13.1 nM to 21.8 ± 12.0 nM, which represents an inhibition of $82.3 \pm 7.1\%$ ($n = 14$; $p < 0.05$; Figure 2-8A, B). Application of xestospongins also increased the basal $[\text{Ca}^{2+}]_i$ from 138.8 ± 13.5 nM to

188.3 ± 19.7 nM, an increase of 38.4 ± 12.7% (n = 14; *p*<0.05). I explored the effect of Aβ₁₋₄₂ on the intracellular Ca²⁺ component of the nicotine-mediated [Ca²⁺]_i rise further by applying caffeine, an agonist of intracellular Ca²⁺ stores. Focal application of caffeine for 5 s caused an average rise in [Ca²⁺]_i of 175 ± 17 nM. In the presence of 400 nM thapsigargin (TG), a SERCA pump inhibitor, the caffeine response was abolished (*p*<0.05; n = 5; Figure 2-9A). As with the other SERCA inhibitor BHQ (Figure 2-7B), the application of TG caused a slow and transient elevation in [Ca²⁺]_i (Figure 2-7A), which can be attributed to transient net efflux from and eventual emptying of intracellular Ca²⁺ stores. In the presence of 100 nM Aβ₁₋₄₂ there was a potentiation of caffeine-mediated [Ca²⁺]_i rises of 137 ± 37 % and a reversible rise in baseline [Ca²⁺]_i of 42.0 ± 8.5 nM. In 3 of 5 neurons, no caffeine-mediated [Ca²⁺]_i rise was observed during washout (*p*<0.05; n = 5; Figure 2-9B).

Application of TG significantly inhibited Aβ mediated potentiation of nAChR mediated [Ca²⁺]_i rises. In the 5 cells examined, Aβ potentiated the nAChR mediated [Ca²⁺]_i rises from 37 ± 8 nM to 89 ± 17 nM (141 ± 22%; *p* < 0.05; Figure 2-10A). Application of TG inhibited this response to 29 ± 7 nM, which represents a 68 ± 4 % inhibition. The next set of experiments examined the effect of a mitochondrial release inhibitor CGP-37157 on Aβ potentiation. In 5 cells, with an average potentiation of 97 ± 18%, 25 μM CGP-37157 significantly inhibited this potentiation by 73 ± 7% (n = 5; *p*<0.05; Figure 2-10B). Previously, CGP-37157 was reported to inhibit voltage gated Ca²⁺ channels (Baron and Thayer, 1997). In our experiments, 25 μM CGP-37157 did not inhibit 8 mM K⁺ mediated [Ca²⁺]_i rises. In 4 cells, 8 mM K⁺ evoked a 64 ± 18 nM increase, while in the presence of 25 μM CGP-37157 8 mM K⁺ evoked a 70 ± 27 nM

[Ca²⁺]_i rise ($p > 0.05$). Furthermore, CGP-37157 did not inhibit 10 mM caffeine mediated responses ($n = 4$; $p > 0.05$). In 4 cells, 10 mM caffeine elicited average responses of 117 ± 37 nM, while in the presence of 25 μ M CGP-37157 10 mM caffeine response was 111 ± 27 nM.

Pharmacology of Non- $\alpha 7$ and $\alpha 7$ nAChR Receptors on DBB Neurons

I then elucidated the pharmacology of the nicotinic response with the non- $\alpha 7$ nAChR antagonist DH β E and the $\alpha 7$ subunit containing competitive nAChR antagonist α BTX. DH β E (5 μ M) inhibited nicotine-mediated rises of [Ca²⁺]_i in 7 of 12 neurons by 42.8 ± 6.0 ($p < 0.05$ Figure 2-11A). In the presence of 100 nM α BTX, the nicotine response was decreased by 35.8 ± 7.7 % ($n = 10$), suggesting that nicotine also activates, in part, $\alpha 7$ nAChRs (Figure 2-11B). In the presence of 5 μ M DH β E and 100 nM α BTX, blocked 70 ± 12 % of the nicotine evoked response ($n = 5$; Figure 2-12A, B). The residual nicotinic response is similar in amplitude as that observed in the experiment when extracellular Ca²⁺ was removed suggesting that the residual response reflects activation of intracellular Ca²⁺ stores that are not activated by Ca²⁺ influx.

Examination of acutely dissociated cells from the DBB using double-labeling immunohistochemistry revealed the presence of $\alpha 4$ or $\alpha 7$ nAChR subunits on vChAT positive neurons (Figure 2-13A-D). Co-localization of $\alpha 4$ and $\alpha 7$ nAChR subunits on the same DBB neurons was also observed (Figure 2-13 E-G). Cell counts of slides double labeled for $\alpha 4$ and $\alpha 7$ subunits revealed that of the neurons counted expressing at least one type of nAChR subunit, 36.0% were labeled only for the $\alpha 4$ subunit (27/75 neurons), 16.0% were labeled only for the $\alpha 7$ subunit (12/75 neurons) and 48.0% co-expressed both $\alpha 4$ and $\alpha 7$ nAChR subunits (36/75 neurons; Figure 2-13H). Furthermore,

46.5% of ChAT labeled neurons expressed the $\alpha 4$ nAChR subunit and 57.0% of ChAT labeled neurons expressed the $\alpha 7$ nAChR subunit.

DISCUSSION

In this study, focal application of nicotine to acutely dissociated rat basal forebrain neurons increased $[Ca^{2+}]_i$ through a combination of enhanced extracellular Ca^{2+} entry via activation of $\alpha 4$ subunit- and $\alpha 7$ -containing nAChRs, and also Ca^{2+} release from intracellular stores with repeated challenges of nicotine. $A\beta$ potentiated the nicotine-mediated $[Ca^{2+}]_i$ rise in the DBB neurons irreversibly, although applications of $A\beta$ alone at concentrations up to 1 μM did not increase $[Ca^{2+}]_i$. $A\beta$ potentiation of nicotine-evoked $[Ca^{2+}]_i$ signals was not due to an enhancement of voltage-dependent Ca^{2+} conductances since $A\beta$ did not potentiate Ca^{2+} entry due to KCl induced depolarization. It appears that the nicotine-evoked rises in $[Ca^{2+}]_i$ involved the release of Ca^{2+} from intracellular stores as such elevations in $[Ca^{2+}]_i$ could be significantly attenuated by both BHQ and xestospongin, a putative IP_3 receptor blocker, which may also inhibit SERCA pumps (Dajas-Bailador et al., 2002; Verkhratsky, 2005). $A\beta$ also potentiated caffeine-mediated rises in $[Ca^{2+}]_i$ suggesting that $A\beta$ increases the release of Ca^{2+} from at least some intracellular stores. Furthermore, $A\beta$ potentiation of caffeine-evoked increases in $[Ca^{2+}]_i$ could be abolished by the SERCA inhibitor TG. TG was also able to block $A\beta$ potentiation of nicotine mediated rises suggesting that $A\beta$ potentiates nicotine and caffeine $[Ca^{2+}]_i$ via overlapping SERCA sensitive stores. Additionally, application of CGP-37157, a mitochondrial Na^+/Ca^{2+} inhibitor also inhibited $A\beta$ potentiation of nicotine mediated rises.

Activation of nAChRs on Basal Forebrain Neurons

nAChRs are present in the basal forebrain and in acutely isolated basal forebrain DBB neurons, and previous studies suggest that nicotinic effects appear to be predominantly mediated via the $\alpha 4\beta 2$ subtype of nAChRs (Fu and Jhamandas, 2003). In the brain slice preparation, application of nicotine increases basal firing rates in GABAergic basal forebrain neurons likely via indirect effects on local circuit glutamate interneurons that express $\alpha 4\beta 2$ nAChRs (Wu et al., 2003). In cells where I studied alterations in $[Ca^{2+}]_i$, the chemical phenotype of the DBB cells (i.e. cholinergic versus GABAergic) was not determined, but a significant number of such neurons showed A β potentiation of nicotine-evoked increases in $[Ca^{2+}]_i$. Based on previous data showing the effect of A β to be limited to cholinergic DBB neurons, I surmise that the nicotinic effects are likely predominantly on cholinergic cells of the basal forebrain (Jhamandas et al., 2001). Our pharmacological and immunohistochemical experiments demonstrated colocalization of $\alpha 4$ and $\alpha 7$ nAChR subunits in a majority of DBB neurons. Additionally, there was significant colocalization of these subunits with the cholinergic marker ChAT. These data correspond well with a recent study examining electrophysiological responses following activation of these receptors in basal forebrain neurons, which demonstrated the co-localization of both non- $\alpha 7$ and $\alpha 7$ nAChR mediated responses (Henderson et al., 2005). Henderson et al. (2005) also performed immunohistochemistry to demonstrate that the $\alpha 7$ nAChRs were correlated with ChAT positive neurons. Given the high proportion of neurons in the present study that displayed sensitivity to α BTX, this suggests that a significant number of DBB cells that were studied were cholinergic neurons. The doses of nicotine that I used were higher than

other studies. However, the use of acutely dissociated cells (where dendritic processes containing nAChRs may be lost) would require a higher concentration of nicotine to activate a sufficient number of nAChRs on the soma. Additionally, trypsin treatment may decrease the binding affinity of nAChRs on the soma. The high doses were blocked by the nicotinic receptor antagonists. The binding affinity reported for α BTX in competition binding assays for the α 7 nAChR is between 0.35-3.5 nM and the binding affinity values reported for DH β E binding to the α 4 β 2 nAChR is between 13.9 and 1900 (Sharples and Wonnacott, 2001). However, in my experiments, these antagonists were capable of inhibiting 10 mM nicotine at 100 nM for α BTX and 10 μ M for DH β E. The discrepancy between the binding data and the Ca²⁺ data may be explained the differences in application. In binding studies, the antagonists were applied for hours of incubation and then the nicotine ligand is applied over the course of minutes. However, in my experiments, the antagonists were applied for minutes and the agonists were applied over the course of seconds. These differences in application technique could explain why the observations in my experiments differ from what is expected from the binding affinity of the antagonists.

NACHR Coupling to Intracellular Ca²⁺ Stores

The effect of A β on nicotinic receptors has been highly controversial as some studies have shown that A β directly activate nAChRs, while others have shown that A β inhibits nAChR (Liu et al., 2001; Pettit et al., 2001; Dineley et al., 2002). In addition, whether A β affects α 7 or non- α 7 nAChR subtypes has also been a subject of debate, since initially the effects of A β on nAChRs were attributed mostly to α 7 nAChRs (Liu et al., 2001), but more recently, non- α 7, probably, α 4 β 2 nAChRs, have been shown to be

affected by A β (Fu and Jhamandas, 2003; Wu et al., 2004). These studies exclusively examined the effect of A β on nAChR currents. In synaptosomes, pretreatment with A β inhibited nicotine-mediated Ca²⁺ entry via both non- α 7 and α 7 nAChRs (Dougherty et al., 2003). In acutely dissociated DBB neurons, both α 4 and α 7 nAChR subunits have been functionally identified to be present on DBB neurons (Fu and Jhamandas, 2003). This observation was further confirmed in the present study where I identified, using immunohistochemical double labeling, the presence of α 4 and α 7 subunits on cholinergic DBB neurons and the colocalization of these subunits. The source of [Ca²⁺]_i rises evoked by nicotine comes from a combination of extracellular activation of both moieties of nAChRs, and also from an interaction between nAChR receptors and intracellular Ca²⁺ stores. The effect of 100 nM A β ₁₋₄₂ in potentiating nicotine-induced responses is specific to nAChRs as no potentiation was observed with KCl-mediated rises in [Ca²⁺]_i. Activation of VGCC by KCl in DBB has previously been shown not to activate any TG-sensitive stores as simultaneous voltage clamp and Ca²⁺ imaging experiments have demonstrated that [Ca²⁺]_i rises are not changed in the presence of TG (Murchison and Griffith, 1998).

A β and [Ca²⁺]_i Homeostasis

Previous studies have shown that A β alters Ca²⁺ homeostasis (Mattson et al., 1992). Mattson et al. (1992) reported that A β fragments enhance glutamate-mediated [Ca²⁺]_i rise and excitotoxic neuronal death, however, sole application of A β fragments did not raise [Ca²⁺]_i entry. On the other hand, Brorson et al. (1994) also showed that A β fragments enhances glutamate-mediated Ca²⁺ and A β fragments evoked [Ca²⁺]_i rises. In our preparation, focal applications of A β alone at concentrations up to 1 μ M did not alter

[Ca²⁺]_i as some other studies have reported (Brorson et al., 1995; Kawahara et al., 2000). However, neither of the aforementioned studies examined actions of the isoform Aβ₁₋₄₂, which is the more pathophysiologically relevant of the Aβ species. Moreover, the study by Kawahara et al. (2000) utilized an immortal hypothalamic cell line where Aβ formed *de novo* Ca²⁺-permeable pores, a finding not observed in preparations other than planar lipid bilayers (Arispe et al., 1996). Recently, application of Aβ₁₋₄₂ for up to six hours was shown not alter [Ca²⁺]_i in cultured hippocampal neurons (Abramov et al., 2003).

Although I did not observe that application of Aβ-induced Ca²⁺ rises, I did observe that Aβ was able to potentiate nAChR and caffeine-evoked increases in [Ca²⁺]_i which involves the repeated release of Ca²⁺ from intracellular stores. Moreover, the potentiation with repeated nicotine applications became larger as the baseline [Ca²⁺]_i rose gradually during and after Aβ₁₋₄₂ application. Aβ₁₋₄₂ application also potentiated repeat activation of intracellular Ca²⁺ stores by caffeine, however, during washout of Aβ₁₋₄₂, caffeine no longer mediated [Ca²⁺]_i rises. Aβ potentiation of both caffeine and nicotine-induced [Ca²⁺]_i rises could be blocked by thapsigargin. Furthermore, Aβ potentiation of nicotine responses could be blocked by the mitochondrial Na⁺/Ca²⁺ inhibitor CGP-37157.

The phenomena that nicotinic responses are potentiated irreversibly by Aβ whereas caffeine responses are transiently potentiated may be related to differences in the total load of cytoplasmic Ca²⁺ and saturation of Ca²⁺ buffers. Repeated application of nicotine results in a significant entry of extracellular Ca²⁺, which would result in an increase in baseline [Ca²⁺]_i and transient [Ca²⁺]_i elevations from repeated activation of nAChRs. Additionally, each activation of nAChRs by nicotine also results in the release of Ca²⁺ from intracellular stores (Dajas-Bailador et al., 2002). When intracellular stores

of $[Ca^{2+}]_i$ are repeatedly released and eventually depleted, a mechanism whereby $A\beta$ can potentiate nicotine-evoked $[Ca^{2+}]_i$ rises could result from progressive overfilling of the nicotine-sensitive stores. Our study shows that $A\beta$ is mediating the overfilling of TG-sensitive stores by increasing Ca^{2+} release from the Na^+/Ca^{2+} exchanger in mitochondria located in close proximity to TG-sensitive stores. The exact mechanism of $A\beta$ action is not known, however, I propose that $A\beta$ could be acting upon mitochondria to cause overfilling resulting in the excess release of Ca^{2+} via the Na^+/Ca^{2+} exchanger, resulting in overfilling of TG sensitive stores. Saturation of the buffers may also play a role in maintaining the elevation. Previous studies have demonstrated that ER-mitochondrial interactions are important for continued filling of the ER (Arnaudeau et al., 2001; Csordas and Hajnoczky, 2001; Smaili et al., 2001).


Repeated application of caffeine activates release of Ca^{2+} from caffeine-sensitive stores via ryanodine receptors. In the absence of any additional entry of extracellular Ca^{2+} , the overall result of repeatedly activating Ca^{2+} stores would be the gradual overall loss of cytoplasmic Ca^{2+} via plasma membrane pumps or exchangers. The transient enhancement of the caffeine response by $A\beta$ is consistent with the idea that there is only a transient overfilling of caffeine-sensitive stores and no net increase in the total cytoplasmic Ca^{2+} load (which also suggested that there was no persistent activation of a Ca^{2+} release activated -like channel).

Functional Implications


In our study, I showed that $A\beta$ potentiates nicotine-evoked $[Ca^{2+}]_i$ rises via overfilling of mitochondria. This effect is only prominent with repeated activation of intracellular Ca^{2+} stores and on a large cytoplasmic Ca^{2+} load via extracellular Ca^{2+} entry.

Given the tight regulation of Ca^{2+} signaling in neurons, the increase in $[\text{Ca}^{2+}]_i$ triggered by nAChR activation may cause dysfunction in the basal forebrain neurons prior to deposition, but not necessarily in the absence of amyloid plaques due to the presence of soluble forms of $\text{A}\beta$. There is general agreement that the neurotoxic effects of $\text{A}\beta$ may, in part, occur via its dysregulation of intracellular Ca^{2+} homeostasis, thus $\text{A}\beta$ -induced enhancement of $[\text{Ca}^{2+}]_i$ via nAChRs on basal forebrain neurons may render them vulnerable to neurodegeneration in AD. Further studies examining the role of overloading endoplasmic reticulum Ca^{2+} stores and its role in neurotoxicity in the basal forebrain may lead to treatments to protect against the degeneration of basal forebrain in AD.


A.




B.




A.




B.




A.




B.







A.




B.







A.




B.




A.




B.




A.




B.




A.




B.




A.




B.




A.




B.





H.



REFERENCES

- Abramov AY, Canevari L, Duchen MR (2003) Changes in intracellular calcium and glutathione in astrocytes as the primary mechanism of amyloid neurotoxicity. *J Neurosci* 23:5088-5095.
- Arispe N, Pollard HB, Rojas E (1996) Zn²⁺ interaction with Alzheimer amyloid beta protein calcium channels. *Proc Natl Acad Sci U S A* 93:1710-1715.
- Arnaudeau S, Kelley WL, Walsh JV, Demaurex N (2001) Mitochondria recycle Ca²⁺ to the endoplasmic reticulum and prevent the depletion of neighboring endoplasmic reticulum regions. *Journal Of Biological Chemistry* 276:29430-29439.
- Auld DS, Kornecook TJ, Bastianetto S, Quirion R (2002) Alzheimer's disease and the basal forebrain cholinergic system: relations to beta-amyloid peptides, cognition, and treatment strategies. *Prog Neurobiol* 68:209-245.
- Baron KT, Thayer SA (1997) CGP37157 modulates mitochondrial Ca²⁺ homeostasis in cultured rat dorsal root ganglion neurons. *Eur J Pharmacol* 340:295-300.
- Brorson JR, Bindokas VP, Iwama T, Marcuccilli CJ, Chisholm JC, Miller RJ (1995) The Ca²⁺ influx induced by beta-amyloid peptide 25-35 in cultured hippocampal neurons results from network excitation. *J Neurobiol* 26:325-338.
- Chin JH, Harris K, MacTavish D, Jhamandas JH (2002) Nociceptin/orphanin FQ modulation of ionic conductances in rat basal forebrain neurons. *J Pharmacol Exp Ther* 303:188-195.

- Csordas G, Hajnoczky G (2001) Sorting of calcium signals at the junctions of endoplasmic reticulum and mitochondria. *Cell Calcium* 29:249-262.
- Dajas-Bailador FA, Mogg AJ, Wonnacott S (2002) Intracellular Ca²⁺ signals evoked by stimulation of nicotinic acetylcholine receptors in SH-SY5Y cells: contribution of voltage-operated Ca²⁺ channels and Ca²⁺ stores. *J Neurochem* 81:606-614.
- Dineley KT, Bell KA, Bui D, Sweatt JD (2002) beta -Amyloid peptide activates alpha 7 nicotinic acetylcholine receptors expressed in *Xenopus* oocytes. *J Biol Chem* 277:25056-25061.
- Dominguez del Toro E, Juiz JM, Peng X, Lindstrom J, Criado M (1994) Immunocytochemical localization of the alpha 7 subunit of the nicotinic acetylcholine receptor in the rat central nervous system. *J Comp Neurol* 349:325-342.
- Dougherty JJ, Wu JL, Nichols RA (2003) beta-amyloid regulation of presynaptic nicotinic receptors in rat hippocampus and neocortex. *Journal Of Neuroscience* 23:6740-6747.
- Fraser SP, Suh YH, Djamgoz MB (1997) Ionic effects of the Alzheimer's disease beta-amyloid precursor protein and its metabolic fragments. *Trends Neurosci* 20:67-72.
- Fu W, Jhamandas JH (2003) Beta-amyloid peptide activates non-alpha7 nicotinic acetylcholine receptors in rat basal forebrain neurons. *J Neurophysiol* 90:3130-3136.

- Grynkiewicz G, Poenie M, Tsien RY (1985) A new generation of Ca²⁺ indicators with greatly improved fluorescence properties. *J Biol Chem* 260:3440-3450.
- Henderson Z, Boros A, Janzso G, Westwood AJ, Monyer H, Halasy K (2005) Somato-dendritic nicotinic receptor responses recorded in vitro from the medial septal diagonal band complex of the rodent. *J Physiol* 562:165-182.
- Jhamandas JH, Cho C, Jassar B, Harris K, MacTavish D, Easaw J (2001) Cellular mechanisms for amyloid beta-protein activation of rat cholinergic basal forebrain neurons. *J Neurophysiol* 86:1312-1320.
- Kamenetz F, Tomita T, Hsieh H, Seabrook G, Borchelt D, Iwatsubo T, Sisodia S, Malinow R (2003) APP processing and synaptic function. *Neuron* 37:925-937.
- Kawahara M, Kuroda Y, Arispe N, Rojas E (2000) Alzheimer's beta-amyloid, human islet amylin, and prion protein fragment evoke intracellular free calcium elevations by a common mechanism in a hypothalamic GnRH neuronal cell line. *J Biol Chem* 275:14077-14083.
- Liu QS, Kawai H, Berg DK (2001) beta-Amyloid peptide blocks the response of alpha 7-containing nicotinic receptors on hippocampal neurons. *Proceedings Of The National Academy Of Sciences Of The United States Of America* 98:4734-4739.
- Mattson MP, Cheng B, Davis D, Bryant K, Lieberburg I, Rydel RE (1992) beta-Amyloid peptides destabilize calcium homeostasis and render human cortical neurons vulnerable to excitotoxicity. *J Neurosci* 12:376-389.

- Mayeux R, Sano M (1999) Drug therapy - Treatment of Alzheimer's disease. New England Journal Of Medicine 341:1670-1679.
- McKhann G, Drachman D, Folstein M, Katzman R, Price D, Stadlan EM (1984) Clinical-Diagnosis Of Alzheimers-Disease - Report Of The Nincds-Adrda Work Group Under The Auspices Of Department-Of-Health-And-Human-Services Task-Force On Alzheimers-Disease. Neurology 34:939-944.
- Murchison D, Griffith WH (1998) Increased calcium buffering in basal forebrain neurons during aging. J Neurophysiol 80:350-364.
- Murchison D, Griffith WH (2000) Mitochondria buffer non-toxic calcium loads and release calcium through the mitochondrial permeability transition pore and sodium/calcium exchanger in rat basal forebrain neurons. Brain Res 854:139-151.
- Nakagawa T, Zhu H, Morishima N, Li E, Xu J, Yankner BA, Yuan JY (2000) Caspase-12 mediates endoplasmic-reticulum-specific apoptosis and cytotoxicity by amyloid-beta. Nature 403:98-103.
- Naslund J, Haroutunian V, Mohs R, Davis KL, Davies P, Greengard P, Buxbaum JD (2000) Correlation between elevated levels of amyloid beta-peptide in the brain and cognitive decline. Jama-Journal Of The American Medical Association 283:1571-1577.
- Nordberg A, Winblad B (1986) Reduced number of [3H]nicotine and [3H]acetylcholine binding sites in the frontal cortex of Alzheimer brains. Neurosci Lett 72:115-119.

- Paterson D, Nordberg A (2000) Neuronal nicotinic receptors in the human brain. *Progress In Neurobiology* 61:75-111.
- Pettit DL, Shao Z, Yakel JL (2001) beta-Amyloid(1-42) peptide directly modulates nicotinic receptors in the rat hippocampal slice. *Journal Of Neuroscience* 21.
- Ramsden M, Henderson Z, Pearson HA (2002) Modulation of Ca²⁺ channel currents in primary cultures of rat cortical neurones by amyloid beta protein (1-40) is dependent on solubility status. *Brain Research* 956:254-261.
- Rogers SW, Gahring LC, Collins AC, Marks M (1998) Age-related changes in neuronal nicotinic acetylcholine receptor subunit alpha4 expression are modified by long-term nicotine administration. *J Neurosci* 18:4825-4832.
- Sharples C, Wonnacott S (2001) Neuronal Nicotinic Receptors. *Toctris Reviews* 19:1-12.
- Silei V, Fabrizi C, Venturini G, Salmona M, Bugiani O, Tagliavini F, Lauro GM (1999) Activation of microglial cells by PrP and beta-amyloid fragments raises intracellular calcium through L-type voltage sensitive calcium channels. *Brain Res* 818:168-170.
- Smaili SS, Stellato KA, Burnett P, Thomas AP, Gaspers LD (2001) Cyclosporin A inhibits inositol 1,4,5-trisphosphate-dependent Ca²⁺ signals by enhancing Ca²⁺ uptake into the endoplasmic reticulum and mitochondria. *Journal Of Biological Chemistry* 276:23329-23340.

- Tsuneki H, Klink R, Lena C, Korn H, Changeux JP (2000) Calcium mobilization elicited by two types of nicotinic acetylcholine receptors in mouse substantia nigra pars compacta. *Eur J Neurosci* 12:2475-2485.
- Verkhatsky A (2005) Physiology and pathophysiology of the calcium store in the endoplasmic reticulum of neurons. *Physiol Rev* 85:201-279.
- Wang HY, Lee DHS, Davis CB, Shank RP (2000a) Amyloid peptide A beta(1-42) binds selectively and with picomolar affinity to alpha 7 nicotinic acetylcholine receptors. *Journal Of Neurochemistry* 75:1155-1161.
- Wang HY, Lee DH, D'Andrea MR, Peterson PA, Shank RP, Reitz AB (2000b) beta-Amyloid(1-42) binds to alpha7 nicotinic acetylcholine receptor with high affinity. Implications for Alzheimer's disease pathology. *J Biol Chem* 275:5626-5632.
- White RJ, Reynolds IJ (1997) Mitochondria accumulate Ca²⁺ following intense glutamate stimulation of cultured rat forebrain neurones. *J Physiol* 498 (Pt 1):31-47.
- Whitehouse PJ, Martino AM, Wagster MV, Price DL, Mayeux R, Atack JR, Kellar KJ (1988) Reductions in [3H]nicotinic acetylcholine binding in Alzheimer's disease and Parkinson's disease: an autoradiographic study. *Neurology* 38:720-723.
- Wu J, Kuo YP, George AA, Xu L, Hu J, Lukas RJ (2004) beta-Amyloid directly inhibits human alpha4beta2-nicotinic acetylcholine receptors heterologously expressed in human SH-EP1 cells. *J Biol Chem* 279:37842-37851.

Wu M, Hajszan T, Leranth C, Alreja M (2003) Nicotine recruits a local glutamatergic circuit to excite septohippocampal GABAergic neurons. *Eur J Neurosci* 18:1155-1168.

Yankner BA (1996) Mechanisms of neuronal degeneration in Alzheimer's disease. *Neuron* 16:921-932.

Chapter 3

**Synaptic Effects of β -Amyloid on Glutamate
Neurotransmission in the Rat Cholinergic Basal
Forebrain**

INTRODUCTION

Alzheimer's disease (AD) is a disease characterized by a progressive decline in memory and other cognitive functions such as language and perception in elderly individuals (McKhann et al., 1984). β -amyloid peptide ($A\beta$), a 39-43 amino acid peptide, is a critical pathological mediator of the synaptic dysfunction, synaptic loss and neuronal death observed in AD (Small et al., 2001; Selkoe, 2002). The pathology of AD starts with mild synaptic dysfunction and eventually progresses to neuronal death, however no receptor has yet been identified to mediate the effects of $A\beta$ on synaptic activity (Small et al., 2001). A potential candidate is the nicotinic acetylcholine receptor (nAChR) as studies have suggested that $A\beta$ binds to this receptor and mediates electrophysiological effects (Wang et al., 2000a; Dineley et al., 2001; Pettit et al., 2001; Fu and Jhamandas, 2003)

Several lines of evidence suggest that the cognitive impairment observed early on in AD is mediated by $A\beta$ induced synaptic dysfunction. Pathological examination of AD brains has revealed that specific neurotransmitter systems are targeted in AD. The most well studied system is the cholinergic basal forebrain neurons, which are especially prone to neurodegeneration (Whitehouse et al., 1982). Additionally, examination of brains from AD individuals 2-4 years after diagnosis reveal significant reductions of synaptic density in the temporal and frontal lobe (Davies et al., 1987; Hamos et al., 1989; Masliah et al., 1991). In individuals with mild cognitive impairment or very mild AD, there was a significant reduction in synaptic density when compared with control individuals (Masliah et al., 2001). Furthermore, studies examining the cognitive decline in AD and neuropathology reveal that the reduction of synaptic density is more closely correlated

with the decline in cognitive function of AD individuals than the deposition of A β plaques (Terry et al., 1991). Studies examining the level of insoluble A β demonstrated either a weak correlation or no correlation between A β plaque load and cognitive decline (Katzman et al., 1988; Morris et al., 1991; Terry et al., 1991). However, a correlation between A β and the degree of cognitive decline was found when both the insoluble and soluble forms of A β were measured (Lue et al., 1999; Naslund et al., 2000). These studies demonstrate that both the accumulation of A β and synaptic loss occur early on AD and may be responsible for the initial mild cognitive decline observed.

Deficits in basal synaptic transmission and long term potentiation (LTP), a form a synaptic plasticity, prior to the development of A β plaques have also been observed in transgenic mice that overexpress A β . These mice express mutant forms of the human amyloid precursor protein that result in overexpression of A β . In the V717F transgenic model, there is a significant reduction in synaptic excitatory postsynaptic potentials and rapid decay of LTP compared with control animals at 4-5 months of age prior to the formation of A β plaques, suggesting that soluble A β isoforms mediate this action (Chen et al., 2000). Although such transgenic mice do not display all the features of AD pathology such as neurofibrillary tangles consisting of hyperphosphorylated tau protein, the synaptic loss observed early on in these animals fits well with the pathological observations made in brains from mildly cognitive impaired individuals. Furthermore, *in vitro* experiments have shown that soluble oligomers of A β are capable of inhibiting LTP (Walsh et al., 2002). Taken together, these data demonstrate the importance of soluble forms of A β in mediating synaptic dysfunction. However, there is at present little information on how A β may influence normal synaptic transmission in the brain,

particularly in structures such as the cholinergic basal forebrain, that are at the epicenter of the chemical pathology seen in AD.

Currently, no receptor has been definitively identified to mediate A β actions on synaptic function, although many have been proposed. One possible candidate receptor is the nicotinic acetylcholine receptor (nAChR), which has been demonstrated to be involved in neuromodulation (McGehee et al., 1995). There are very few examples of primary nicotinic mediated EPSC in the brain and these include hippocampal interneurons, the medial vestibular nucleus and the nucleus ambiguous (Phelan and Gallagher, 1992; Zhang et al., 1993; Frazier et al., 1998). In this project, I examined the DBB for nAChR mediated EPSCs, but did not find any and thus I focused on nicotinic modulation of glutamate EPSCs. A β has been shown to exhibit significant affinity to these receptors and electrophysiological experiments have shown that A β is capable of influencing the function of these receptors (Pettit et al., 2001; Fu and Jhamandas, 2003). I will examine the effect of A β_{1-42} on synaptic transmission in cholinergic neurons of the basal forebrain identified with the fluorescent dye Cy3 192IgG or with electrophysiological criteria. The aims of this project are 1) to examine the effect of soluble A β_{1-42} on spontaneous miniature post-synaptic excitatory events (mEPSCs) in cholinergic basal forebrain neurons and 2) to determine if A β effects on spontaneous excitatory events are mediated by nAChRs, and if so, what specific subtypes of nAChRs are involved.

MATERIALS AND METHODS

Cy3-192 IgG Neuron Labeling

The majority of neurons used in this study (43/50) were identified with Cy3-192 IgG (Advanced Targeting Systems, San Diego, USA) which is an inert fluorescent dye conjugated to an antibody that binds to the p75 neurotrophin receptor expressed only in cholinergic neurons of the basal forebrain. Following intracerebroventricular injection, Cy3-192 IgG retrogradely labels only cholinergic neurons of the basal forebrain that project to the hippocampus (Hartig et al., 1998a; Wu et al., 2000). Injection of Cy3-192 IgG was performed based upon a previously described protocol (Wu et al., 2000). 22-27 day post-natal Sprague-Dawley rats (50-70 g) were anaesthetized with an intraperitoneal injection of sodium pentobarbital (50 mg/kg; 0.05% Somnotol, MTC Pharmaceuticals, Hamilton, Canada) and then injected subcutaneously with 0.02% of atropine. The rats were then placed in a stereotaxic frame (Narishige, Tokyo, Japan) and 5 μ l of 1:1 diluted Cy3-192 IgG was injected into the left and right ventricles (1.1 mm posterior to Bregma, 1.2 mm lateral from the midline, and 2.6-3.7 mm below the dura). All procedures were approved by University of Alberta Health Sciences Laboratory Animal Services (Protocol number 154/04/05).

Diagonal Band of Broca Slice Preparation

Brain tissue was removed from Sprague-Dawley rats previously injected with Cy3-192 IgG between three and seven days after injections based on previously described procedures (Easaw et al., 1997). Briefly, animals were anaesthetized with halothane and decapitated. The brain was quickly removed and placed in a 3-5 °C bicarbonate buffered solution that contained (in mM): 140 NaCl, 2.5 KCl, 12 MgCl₂, 1.2 NaH₂PO₄ and 2.4

CaCl₂, 25 mM HCO₃ and 11 mM D-glucose (pH 7.4). All solutions were oxygenated by bubbling with a mixture of 95% O₂ and 5% CO₂. 300 μM thick longitudinal brain slices were cut with a vibratome (Slicer HR2, Sigmund Elektronik, Germany) and then incubated for one hour at 32 °C in artificial cerebrospinal fluid (ACSF) prior to recording (in mM; NaCl 126, 2.5 KCl, 1.2 MgCl₂, 1.2 NaH₂PO₄, 2.4 CaCl₂, 25 NaHCO₃ and 11 mM D-glucose; pH 7.4).

Recordings from DBB Slices

DBB slices were bath perfused with ACSF (23-25°C) at a rate of 1 ml/min and visualized under an Axioscope 2 Fs microscope (Zeiss, Germany) at 60x magnification. Cy3-192 IgG labeled neurons were selected using the appropriate filter for Cy3 (546 nm excitation and 575-640 nm emission). Individual neurons were then visualized under 60x magnification using differential infrared contrast optics. Cells were recorded using whole cell patch clamp technique. The internal pipette solution was composed of the following (in mM): 140 K⁺-gluconate, 2 KCl, 5 HEPES, 5 MgATP, 0.5 NaGTP, 10 EGTA and pH was raised to between 7.2 and 7.3 with potassium hydroxide. The internal solution had an osmolarity of 280 mM. Patch clamp electrodes (World Precision Instruments, Sarasota, USA – Thin Wall with Filament, 1.5 mm) were pulled with a Narishige (PP-83) puller to yield electrodes with resistances of 4-8 MΩ. Recordings were made from the DBB, where seals of 1 GΩ or greater were obtained using a Siskiyou Design Instruments 4 axis motorized micromanipulator (MX 831; Grants Pass, Oregon). Using an Axopatch 200B amplifier (Molecular Devices, Sunnyvale CA), Cy3-192IgG labeled cells were initially held in voltage-clamp mode at -60 mV. Recordings were made at a bandwidth of 10 kHz and filtered with a 2 kHz low pass Bessel filter.

Prior to performing voltage-clamp recordings, in 7 of 50 neurons, neurons were held in current clamp and 2 second current injections of 0.2 nA steps from -0.6 nA to +0.6 nA were performed in the absence of tetrodotoxin (TTX; Figure-3-1A). The average resting membrane potential for these cells was -53.9 ± 5.1 mV ($n = 7$). Cholinergic neurons were identified based upon previously described criteria such as lack of a hyperpolarizing sag and the presence of burst firing upon depolarization (Wu et al., 2000). A gap free protocol was used to hold cells at either -60 or -80 mV and recordings were performed in the presence of 1 μ M TTX and either 10 μ M of bicuculline or 50 μ M picrotoxin, GABA_A receptor antagonists. Baseline recordings for 5 minutes were observed to ensure stability of the cell. Each recorded treatment was between 5 and 7 minutes, with a 2 minute interval between treatments so that at least 100 events were recorded. Seal and voltage ramp protocols were run between treatments. Seal tests were performed between treatments to monitor cell access was maintained. Voltage ramps consisted of hyperpolarizing the cells to -110 mV, and then depolarizing them to +30 mV over the course of 8 seconds (Figure 3-1B).

Statistical Analysis

Miniature excitatory post-synaptic currents were recorded using pClamp 9.0 and analyzed with Clampfit 9.2 using a template search for events (Clements and Bekkers, 1997). A template was created based upon the average of 10 recorded events. The template search parameter was set between 4.0 and 5.0 and events were then inspected visually for removal of extraneous events. Student's paired and unpaired t-test was used to compare treatments and parameters between cell populations respectively. The significance level for the t-tests was set at $p < 0.05$. Kolmogorov-Smirnov (KS) testing

was also used to examine differences and the significance level of this test was set at $p < 0.0005$. The mEPSC parameters illustrated in Figure 3-1 were measured. Peak amplitude of the mEPSC was measured (Figure 3-1(a)). The rise time from 10% of the peak amplitude to 90% of the peak amplitude was measured (Figure 3-1(b)) and the decay time from 90% of the peak amplitude to 10% of the peak amplitude was measured (Figure 3-1(c)) and the area between the baseline and the event was also measured (Figure 3-1(d)). The conductance of was calculated from voltage ramps by taking performing linear regression of the current-voltage relationship between -110 mV and -60 mV.

Immunohistochemistry

Rats previously injected with Cy3-192 IgG were anaesthetized with halothane and intraperitoneal urethane was administered. Animals were then exsanguinated with phosphate buffered saline (PBS) and infused with 4% formaldehyde in 0.1% PBS that was kept at 3-5 °C. Brains removed from the animals were incubated for 1 hour in 4% formaldehyde in PBS and then placed in 10% sucrose solution. 50 µm coronal DBB brain slices were cut using a cryostat and then blocked with 1% bovine serum albumin for 1 hour. Slices were then incubated overnight with primary rabbit antibodies for vesicular choline acetyltransferase at 3-5 °C (vChAT; 1:2500 dilution; Santa Cruz Biotechnology Inc., Santa Cruz, USA). Slices were then washed 3 times in PBS and incubated with Alexa 488 chicken anti-rabbit antibodies for 1.5 hours to yield green fluorescence (1:400 dilution) and washed 3 times in PBS. Slides were allowed to air dry and dehydrated with ethanol and xylene before being coverslipped with Cytoseal (Canada

Wide Scientific, Ottawa, Canada). Slides were examined with a Zeiss Axioplan 2 and images captured with Zeiss AxioCam MRc camera and MGrab software.

Drugs

A β ₁₋₄₂ and A β ₄₂₋₁ were purchased from American Peptide Co. (Sunnyvale, USA). A β ₁₋₄₂ was freshly prepared on the day of the experiment from aliquots stored at – 80 °C. All voltage clamp experiments were conducted in the presence of 1 μ M tetrodotoxin (TTX) and either 10 μ M bicuculline or 50 μ M picrotoxin. All chemicals were purchased from Sigma except for the following. TTX was purchased from Alamone Lab (Jerusalem, Israel). All drugs were bath applied through four-way valve system and the average time to the onset of drug action was 30 sec.

RESULTS

Cy3-192 IgG Labeling Colocalizes with vChAT Labeling in DBB Neurons

Immunohistochemical experiments demonstrated significant co-localization of Cy3-192 IgG labeling with vChAT labeling (Figures 3-2A, B), confirming previous studies suggesting Cy3-192 IgG-labeled cells are indeed cholinergic (Hartig et al., 1998b; Wu et al., 2000). Figure 3-2C shows a Cy3-192 IgG labeled cell viewed under a water immersion lens at 60x magnification and Figure 3-2D shows the same neuron being patched under differential infrared contrast imaging. The average capacitance of neurons was 18.9 ± 0.5 pF and the average access resistance was 6.6 ± 0.2 m Ω (n = 50). The average conductance of the cells was 2.85 ± 0.25 nS (n = 50). No significant change was observed in the conductance in any of the treatments recorded.

CNQX Abolishes mEPSCs in DBB Neurons

The character of these mEPSCs was examined with the AMPA/kainate receptor antagonist 6-cyano-7-nitroquinoxaline-2,3-dione (CNQX). In the presence of 1 μ M TTX and 50 μ M picrotoxin, 2 μ M CNQX abolished all synaptic activity in 3 electrophysiologically identified cholinergic neurons (Fig. 3-3). These data demonstrate that the recorded events were glutamatergic events mediated by AMPA and kainate receptors.

A β Modulates Frequency of mEPSCs in Cholinergic DBB Neurons

In 12 Cy3-IgG 192 labeled neurons and 3 electrophysiologically identified cholinergic neurons, bath application of 100 nM A β ₁₋₄₂ increased mEPSC frequency in 54% of neurons tested (15/28 neurons tested; Figure 3-4A; $p < 0.05$). KS testing of individual cells demonstrated that in all 15 neurons, there was no significant change in amplitude, while the interevent interval was significantly decreased ($p < 0.0005$; Figure 3-4B, C). The average mEPSC frequency was 0.89 ± 0.16 Hz, which increased to 1.21 ± 0.28 Hz in the presence of 100 nM A β ₁₋₄₂ and recovered to 0.92 ± 0.28 Hz which represents a $29.6 \pm 5.4\%$ increase in frequency ($p < 0.05$; Fig. 3-4A, 3-5). The peak amplitude of mEPSCs did not exhibit any significant change in these neurons as the control peak amplitude was -37.1 ± 1.7 pA and the peak amplitude in the presence of 100 nM A β ₁₋₄₂ was -36.8 ± 1.9 pA (Fig. 3-4B, 3-5). Examination of specific kinetic parameters of the mEPSCs revealed that there were no significant changes in rise time, decay time, and area between control conditions and 100 nM A β ₁₋₄₂ treatment (Table 3-1). Linear regression of the current from -110 mV to -60 mV revealed that there were no significant difference between control conductance, which was 3.08 ± 0.32 nS and

conductance in the presence of 100 nM A β_{1-42} , which was 3.19 ± 0.45 nS ($p > 0.05$). In 7 Cy3-192IgG labeled neurons and 2 electrophysiologically identified cholinergic neurons, bath application of 100 nM A β_{1-42} significantly reduced mEPSC frequency from 0.97 ± 0.21 to 0.68 ± 0.18 Hz, which represents a $31.8 \pm 4.8\%$ decrease of frequency (Figure 3-6 and 3-7; $p < 0.05$; 9/28 neurons). In all 9 neurons, no change in peak amplitude was observed with KS' testing, however, there was a significant decrease in the interevent interval ($p < 0.0005$). The peak amplitude was -34.7 ± 2.7 pA under control conditions, while in the presence of 100 nM A β_{1-42} the peak amplitude was -31.4 ± 2.3 pA (Figure 3-7B; $p > 0.05$; $n = 9$). Examination of specific kinetic parameters of the mEPSCs revealed that there were no significant changes in rise time (10%-90%), decay time (90%-10%), and area between control conditions and 100 nM A β_{1-42} treatment of cholinergic neurons (Table 3-2). In 14% of neurons ($n = 4$), no change in peak amplitude or frequency of 100 nM A β_{1-42} was observed. Comparison of neurons did not reveal any difference in amplitude, frequency, or kinetic parameters between neurons that demonstrated an increase in mEPSC frequency and those that demonstrated a decrease in mEPSC frequency.

Application of the reverse peptide A β_{42-1} did not significantly change either the peak amplitude or mEPSC frequency in Cy3 IgG192 labeled neurons ($p < 0.05$; $n = 4$; Figure 3-8). In these 4 neurons, KS testing did not detect a change in either peak amplitude or interevent interval during the application of A β_{42-1} . In neurons that exhibited increased mEPSC frequency in response to A β_{1-42} , application of 100 nM A β_{42-1} did not elicit any change in either frequency or peak amplitude. In the presence of the reverse peptide 100 nM A β_{42-1} , the frequency was 0.69 ± 0.27 Hz, which was not

significantly different from the frequency under control conditions, which was 0.71 ± 0.29 Hz ($p > 0.05$; Figure 3-9A; $n = 4$). The average peak amplitude under control conditions was -41.1 ± 3.9 pA, while in the presence of $A\beta_{42-1}$, the peak amplitude was -40.0 ± 5.6 pA ($p > 0.05$; Figure 3-9B; $n = 4$). No significant differences in kinetic parameters were observed between control conditions and application of 100 nM $A\beta_{42-1}$.

A β Mediated Increase in mEPSC Frequency is Inhibited by DH β E

The effect of the $\alpha 4\beta 2$ nAChR selective antagonist DH β E on $A\beta_{1-42}$ mediated increases in mEPSC frequency was examined. Application of 100 nM $A\beta_{1-42}$ to 5 Cy3 IgG 192 labeled neurons resulted in an increase in frequency from 0.64 ± 0.13 Hz under control conditions to 0.84 ± 0.15 Hz in the presence of 100 nM $A\beta_{1-42}$, which is a $25.4 \pm 7.9\%$ increase (Figure 3-10A; $p < 0.05$; $n = 5$). As previously observed, there was no significant change in peak amplitude which was -37.4 ± 1.7 pA under control conditions and -36.2 ± 0.09 pA in the presence of 100 nM $A\beta_{1-42}$. Subsequent application of 100 nM $A\beta_{1-42}$ in the presence of 10 μ M DH β E blocked the increase by 100 nM $A\beta_{1-42}$. Prior to co-application of 100 nM $A\beta_{1-42}$ and 10 μ M DH β E, the frequency was 0.65 Hz ± 0.17 Hz. In the presence of 100 nM $A\beta_{1-42}$ and 10 μ M DH β E, the frequency was 0.67 ± 0.17 Hz, which represents a $2.4 \pm 5.6\%$ change ($p > 0.05$; $n = 5$; Figure 3-10B). Under control conditions, the average peak amplitude was -36.0 ± 0.07 pA, while in the presence of 100 nM $A\beta_{1-42}$ and 10 μ M DH β E, the average peak amplitude was -35.6 ± 1.8 pA ($p > 0.05$; $n = 5$). Additionally, no significant change in kinetic parameters was observed (Table 3-3). KS testing of the individual neurons also demonstrated that $A\beta_{1-42}$ mediated increases of mEPSCs were inhibited by DH β E (Figure 3-11).

Control experiments demonstrated that $A\beta_{1-42}$ mediated increases of frequency in electrophysiologically identified cholinergic neurons were not time dependent as in 3 cholinergic neurons, consecutive application of 100 nM $A\beta_{1-42}$ 7 minutes apart yielded significant increases of mEPSC frequency of $27.3 \pm 4.5\%$ and $29.8 \pm 3.5\%$ ($p > 0.05$). There was no significant change in peak amplitude. KS testing of the individual neurons also showed that the effect of $A\beta_{1-42}$ was not time dependent. Application of 10 μ M DH β E on electrophysiologically identified cholinergic neurons did not significantly change mEPSC frequency or peak amplitude when examined with KS testing ($p > 0.0005$; $n = 2$).

A β Mediated Decrease in mEPSC Frequency Persists in the Presence of DH β E

In Cy3-192 IgG labeled neurons, 10 μ M DH β E did not block the $A\beta_{1-42}$ mediated decrease in frequency. In the neurons tested, application of $A\beta_{1-42}$ on its own significantly decreased frequency from 1.75 ± 0.51 Hz to 1.25 ± 0.36 Hz, which represents a $27.5 \pm 3.7\%$ decrease (Figure 3-12; $n = 5$, $p < 0.05$). As previously described, there was no significant change in the average peak amplitude in the presence of 100 nM $A\beta_{1-42}$. Subsequent application of 100 nM $A\beta_{1-42}$ and 10 μ M DH β E did not significantly inhibit the decreased frequency observed. The average frequency under control conditions was 1.50 ± 0.45 Hz and in the presence of 100 nM $A\beta_{1-42}$ and 10 μ M DH β E, the frequency decreased to 1.15 ± 0.37 Hz, representing a $24.5 \pm 2.8\%$ decrease ($p < 0.05$; $n = 5$). After washout, the frequency recovered to 1.46 ± 0.46 Hz. The peak amplitude was -31.5 ± 4.2 pA under control conditions and -32.4 ± 3.5 pA in the presence of 100 nM $A\beta_{1-42}$ and 10 μ M DH β E ($p < 0.05$; $n = 5$). Additionally, no changes to kinetic parameters were observed between treatments (Table 3-4). KS testing of the individual

neurons also demonstrated that A β_{1-42} mediated decreases of mEPSCs were not inhibited by DH β E (Figure 3-13).

Nicotine Increases mEPSC Frequency in a Dose-Dependent Manner

To further examine nicotinic receptors on cholinergic basal forebrain neurons, various doses of nicotine were bath applied to Cy3-192IgG labeled neurons. Application of 1 μ M (n = 7), 10 μ M (n = 5), and 100 μ M (n = 5) nicotine revealed that Cy3-192IgG labeled neurons exhibited a dose-dependent increase in frequency in the majority of neurons (19/22) (Fig. 3-14A, B). mEPSC frequency increased from 0.66 ± 0.19 Hz to 1.02 ± 0.25 Hz in the presence of 100 μ M nicotine and recovered back to 0.71 ± 0.23 Hz after washout. No significant changes in amplitude, rise time, decay time or area were observed (Tables 3-5 to 3-7). In 3 of 9 neurons, bath application of 100 μ M nicotine did not significantly change mEPSC frequency. No significant changes in amplitude, decay time or area were observed, however there was a significant increase in rise time (Table 3-8). KS testing of individual neurons, demonstrated that in all 9 neurons, there was no change in peak amplitude during the application of 100 μ M nicotine (Figure 3-15). In 6 of 9 neurons, KS testing demonstrated that 100 μ M nicotine significantly decreased the interevent interval ($p < 0.0005$), while in 3 of 9 neurons, KS testing did not show any change in the interval event interval ($p < 0.0005$).

Acetylcholine Increases mEPSC Frequency in a Dose-Dependent Manner

In the next set of experiments, the endogenous neurotransmitter ACh in the presence of 1 μ M atropine, a muscarinic receptor blocker, was applied to Cy3-192 IgG labeled neurons. Application of ACh demonstrated, in a dose-dependent manner, an increase in mEPSC frequency (Figure 3-15). For 1 mM of ACh, mEPSC frequency

increased from 0.49 ± 0.03 Hz to 0.76 ± 0.05 Hz and then recovered back to 0.44 ± 0.05 Hz, which represents an average increase of $54 \pm 12\%$. No significant changes to peak amplitude, 90%-10% decay time or area were observed (Tables 3-9 to 3-11). KS testing of individual neurons demonstrated that 1 mM ACh significantly decreases interevent interval in cholinergic neurons ($p < 0.0005$) without significantly changing peak amplitude.

DISCUSSION

In this study, the effect of $A\beta_{1-42}$ and nicotine on fluorescently labeled cholinergic DBB neurons were examined. In the majority of cholinergic neurons, bath applications of $A\beta_{1-42}$ significantly increased the frequency of mEPSCs without any significant change in amplitude or kinetic parameters of EPSCs. Furthermore, mEPSCs recorded from DBB neurons were mediated by the AMPA and kainate subtypes of glutamate receptors as application of CNQX, an AMPA/kainate receptor antagonist, completely abolished the spontaneous mEPSCs. This effect of $A\beta_{1-42}$ could be blocked by a non- $\alpha 7$, $\alpha 4\beta 2$ selective nAChR antagonist, suggesting that soluble $A\beta_{1-42}$ is able to modulate presynaptic glutamatergic neurotransmission via activation of $\alpha 4\beta 2$ nAChRs.

In another subset of cholinergic neurons, $A\beta_{1-42}$ decreased the frequency of mEPSCs without changing any kinetic parameters. The decrease in frequency was not inhibited by DH β E suggesting that $A\beta_{1-42}$ is mediating a presynaptic effect via a different receptor. No differences in kinetic parameters could be observed between the two populations of neurons.

Applications of nicotine and ACh (in the presence of atropine) resulted in a dose-dependent increase in frequency, without a significant change in amplitude of the mEPSCs that mimicked the effects of A β_{1-42} .

Soluble A β Modulates Synaptic Transmission onto Cholinergic DBB Neurons

There is significant controversy regarding the interaction of A β_{1-42} and nAChRs. Initial binding studies demonstrated that A β_{1-42} binds to both $\alpha 7$ nAChRs with picomolar affinity and $\alpha 4\beta 2$ nAChRs with nanomolar affinity, although these studies have been difficult to replicate in other laboratories (Wang et al., 2000b; Wang et al., 2000a). Nonetheless, a number of electrophysiological studies have demonstrated that A β_{1-42} can exert its effects through both $\alpha 7$ and $\alpha 4\beta 2$ nAChRs in a variety of *in vitro* preparations (Pettit et al., 2001; Dineley et al., 2002; Fu and Jhamandas, 2003). However, the effect of A β_{1-42} interaction with the nAChR remains controversial as some studies suggest that A β_{1-42} activates $\alpha 4\beta 2$ nAChRs, while other studies suggest that A β_{1-42} inhibits nAChRs (Fu and Jhamandas, 2003; Wu et al., 2004). In this study, A β_{1-42} was demonstrated to mediate an increase in the frequency of spontaneous glutamatergic mEPSCs in a subset of cholinergic DBB neurons via activation of non- $\alpha 7$ nAChR, which is probably of the $\alpha 4\beta 2$ nAChR subtype. This A β_{1-42} activation of $\alpha 4\beta 2$ nAChR is in agreement with single channel recordings from DBB neurons which demonstrated that A β_{1-42} was able to act as an agonist at $\alpha 4\beta 2$ nAChR in DBB neurons (Fu and Jhamandas, 2003). However, the effects of A β_{1-42} observed in this study were localized to a presynaptic locus of action, whereas Fu and Jhamandas (2003) reported post-synaptic $\alpha 4\beta 2$ nAChR-mediated responses. This difference may reflect the topographical differences in the location of glutamate terminals and nAChRs in that post-synaptic nAChRs may not be in close

proximity to glutamatergic presynaptic terminals in order to mediate a recordable effect. Alternately, mEPSC recordings may not be sensitive enough to detect the level of change in the kinetic parameters that could result in subtle post-synaptic modulation mediated by nAChRs.

In another subset of cholinergic neurons, DH β E did not affect the A β_{1-42} -induced inhibition of mEPSC frequency, suggesting that such effects of A β_{1-42} are mediated by another mechanism. As there are a plethora of receptors in the basal forebrain and the cholinergic neurons receive innervation from many different areas of the brain, further experiments need to be conducted in order to elucidate the possible receptor that could mediate this aspect of synaptic modulation.

Activation of nAChRs Modulate Synaptic Transmission onto Cholinergic DBB Neurons

These experiments demonstrated that both nicotine and ACh in the presence of atropine were able to modulate spontaneous mEPSCs via a presynaptic mechanism in a dose dependent manner. The increase in mEPSC frequency without any change in peak amplitude observed is similar to that observed to the effect that A β_{1-42} had on a population of neurons. Studies in other areas of the brain such as the cortex, hippocampus, and striatum have demonstrated that presynaptic nAChRs can modulate synaptic transmission (Alkondon et al., 1997; Girod et al., 2000; Cao et al., 2005). These studies have examined nAChR modulation of dopamine, GABA, and glutamate release and found that activation of both $\alpha 7$ and $\alpha 4\beta 2$ nAChR on presynaptic terminals can increase neurotransmitter release.

Studies examining nicotine in the basal forebrain neurons did not reveal any changes in cholinergic neuron firing in the presence of ACh or nicotine (Wu et al., 2000;

Wu et al., 2003). However, one recent study demonstrated that nAChRs could modulate the excitability of cholinergic DBB neurons via $\alpha 4\beta 2$ and $\alpha 7$ nAChRs (Henderson et al., 2005). We were unable to observe any changes in conductance or in the kinetic parameters of mEPSCs to suggest that there was post-synaptic modulation of the glutamate response by nicotine or ACh.

Future Experiments

Two key issues regarding nAChRs need to be resolved. First, the identity of the receptor involved in $A\beta_{1-42}$ decreases of mEPSCs needs to be examined. This decrease may be mediated by $A\beta_{1-42}$ effects on another neurotransmitter system in the DBB or possibly by the $\alpha 7$ nAChR. An experiment the $\alpha 7$ nAChR antagonist α BTX was applied in the presence of $A\beta$ would resolve this issue. Secondly, pharmacological experiments need to be performed on DBB neurons to elucidate if $\alpha 4\beta 2$ or $\alpha 7$ nAChRs are involved in mediating the effects nicotine and ACh. Furthermore, it would be of interest to examine the link between $A\beta$ and nicotine effects on synaptic function. Experiments in which $A\beta$ and nicotine were applied concurrently after application of a single dose would elicit whether $A\beta$ and nicotine were acting on the same receptor to mediate their effects on mEPSCs.

Functional Implications

This study demonstrates that $A\beta_{1-42}$ is capable of modulating synaptic transmission in septo-hippocampal neurons of the basal forebrain, an important pathway for learning and memory via $\alpha 4\beta 2$ nAChRs in cholinergic neurons of the basal forebrain. Thus the implications of this research are twofold. First, this study suggests that modulation of synaptic transmission by soluble forms of $A\beta_{1-42}$ may be a potential

pathological mechanism for the cognitive decline observed early on in AD. $A\beta_{1-42}$ augmentation of glutamate neurotransmission could lead to excitotoxicity over the long term, while $A\beta_{1-42}$ reduction of glutamate neurotransmission, would weaken cholinergic tone and lead to impaired cognition. Secondly, as the detection of AD becomes more sensitive, the nAChR system is a potential therapeutic target in the treatment of AD.

Table 3-1. Kinetic Parameters of Cholinergic Neurons that Demonstrate an Increase of mEPSCs in the Presence of 100 nM A β ₁₋₄₂.

Statistic	Control	100 nM A β ₁₋₄₂	Recovery
Peak Amplitude (pA)	-37.1 \pm 1.7	-36.8 \pm 1.9	-35.9 \pm 1.7
10%-90% Rise time (ms)	0.73 \pm 0.09	0.80 \pm 0.10	0.81 \pm 0.10
90%-10% Decay time (ms)	2.84 \pm 0.80	2.88 \pm 0.31	2.86 \pm .32
Area (nA ms)	-0.068 \pm 0.007	-0.068 \pm 0.007	-0.068 \pm 0.007

Table 3-2. Kinetic Parameters of Cholinergic Neurons that Demonstrate a Decrease of mEPSCs in the presence of 100 nM A β ₁₋₄₂.

Statistic	Control	100 nM A β ₁₋₄₂	Recovery
Peak Amplitude (pA)	-35.0 \pm 2.9	-31.4 \pm 2.3	-33.3 \pm 2.7
10%-90% Rise time (ms)	0.65 \pm 0.04	0.64 \pm 0.06	0.63 \pm 0.06
90%-10% Decay time (ms)	2.56 \pm 0.34	2.53 \pm 0.34	2.51 \pm 0.31
Area (nA ms)	-0.063 \pm 0.006	-0.056 \pm 0.006	-0.058 \pm 0.006

Table 3-3. Kinetic Parameters of Cholinergic Neurons that Demonstrate an Increase in the Frequency of mEPSCs in the Presence of 100 nM A β ₁₋₄₂ that are Subsequently Exposed to 100 nM A β ₁₋₄₂ and 10 μ M DH β E

Statistic	Control	100 nM A β ₁₋₄₂ + 10 μ M DH β E	Recovery
Peak Amplitude (pA)	-36.0 \pm 0.7	-35.6 \pm 0.2	-34.8 \pm 0.9
10%-90% Rise time (ms)	0.77 \pm 0.06	0.80 \pm 0.07	0.78 \pm 0.06
90%-10% Decay time (ms)	2.27 \pm 0.17	2.28 \pm 0.23	2.26 \pm 0.20
Area (nA ms)	-0.052 \pm 0.007	-0.053 \pm 0.007	-0.053 \pm 0.007

Table 3-4. Kinetic parameters of Cholinergic Neurons that Demonstrate a Decrease in the Frequency of mEPSCs in the presence of 100 nM A β ₁₋₄₂ that are Subsequently Exposed to 100 nM A β ₁₋₄₂ and 10 μ M DH β E

Statistic	Control	100 nM A β ₁₋₄₂ + 10 μ M DH β E	Recovery
Peak Amplitude (pA)	-33.3 \pm 4.5	-32.7 \pm 4.1	-31.5 \pm 4.2
10%-90% Rise time (ms)	0.67 \pm 0.04	0.66 \pm 0.05	0.69 \pm 0.05
90%-10% Decay time (ms)	1.77 \pm 0.36	1.69 \pm 0.39	2.08 \pm 0.17
Area (nA ms)	-0.054 \pm 0.007	-0.051 \pm 0.006	-0.051 \pm 0.007

Table 3-5. Kinetic Parameters of Cholinergic Neurons that Demonstrate an Increase in the Frequency of mEPSCs in response to Bath Application of 100 μ M Nicotine.

Statistic	Control	100 μ M Nicotine	Recovery
Peak Amplitude (pA)	-27.3 \pm 4.6	-22.6 \pm 2.5	-24.0 \pm 4.0
10%-90% Rise time (ms)	0.62 \pm 0.07	0.72 \pm 0.07	0.74 \pm 0.08
90%-10% Decay time (ms)	2.12 \pm 0.19	1.99 \pm 0.17	2.31 \pm 0.22
Area (nA ms)	-0.047 \pm 0.006	-0.041 \pm 0.04	-0.045 \pm 0.006

Table 3-6. Kinetic Parameters of Cholinergic Neurons Demonstrating an Increase of mEPSC Frequency in Response to Bath Application of 10 μ M Nicotine. * denotes ($p < 0.05$)

Statistic	Control	10 μ M Nicotine	Recovery
Peak Amplitude (pA)	-27.4 \pm 5.7	-27.2 \pm 4.5	-25.7 \pm 5.4
10%-90% Rise time (ms)	0.61 \pm 0.05	0.93 \pm 0.20*	0.98 \pm 0.23
90%-10% Decay time (ms)	2.03 \pm 0.16	1.99 \pm 0.19	2.07 \pm 0.16
Area (nA ms)	-0.054 \pm 0.001	-0.051 \pm 0.010	-0.048 \pm 0.010

Table 3-7. Kinetic Parameters of Cholinergic Neurons Demonstrating an Increase of mEPSC Frequency in Response to Bath Application of 1 μ M Nicotine. * denotes ($p < 0.05$)

Statistic	Control	10 μ M Nicotine	Recovery
Peak Amplitude (pA)	-27.3 \pm 3.2	-26.8 \pm 2.6	-27.6 \pm 2.6
10%-90% Rise time (ms)	0.60 \pm 0.07	0.62 \pm 0.06	0.61 \pm 0.05
90%-10% Decay time (ms)	2.06 \pm 0.16	1.91 \pm 0.11	1.96 \pm 0.10
Area (nA ms)	-0.050 \pm 0.006	-0.049 \pm 0.005	-0.05 \pm 0.005

Table 3-8. Kinetic Parameters of Cholinergic Neurons Demonstrating no Change of mEPSC Frequency in Response to Bath Application of 100 μ M Nicotine. * denotes ($p < 0.05$)

Statistic	Control	100 μ M Nicotine	Recovery
Peak Amplitude (pA)	-29.9 \pm 4.6	-29.7 \pm 5.0	-26.9 \pm 2.2
10%-90% Rise time (ms)	0.79 \pm 0.18	0.93 \pm 0.20*	0.98 \pm 0.23
90%-10% Decay time (ms)	2.25 \pm 0.40	2.53 \pm 0.38	2.49 \pm 0.45
Area (nA ms)	-0.048 \pm 0.03	-0.057 \pm 0.02	-0.053 \pm 0.02

Table 3-9. Kinetic Parameters of Cholinergic Neurons in the Presence of 1 mM ACh. * denotes ($p < 0.05$)

Statistic	Control	1 mM ACh	Recovery
Peak Amplitude (pA)	-29.4 \pm 3.5	-22.7 \pm 2.8	-23.3 \pm 2.4
10%-90% Rise time (ms)	0.51 \pm 0.05	0.67 \pm 0.08	0.70 \pm 0.07
90%-10% Decay time (ms)	1.62 \pm 0.12	1.60 \pm 0.10	1.68 \pm 0.10
Area (nA ms)	-0.043 \pm 0.003	-0.034 \pm 0.002	-0.036 \pm 0.003


Table 3-10. Kinetic parameters of Cholinergic Neurons in the Presence of 100 μM ACh. * denotes ($p < 0.05$)

Statistic	Control	100 μM ACh	Recovery
Peak Amplitude (pA)	-23.0 ± 4.4	-20.5 ± 4.8	-22.1 ± 5.8
10%-90% Rise time (ms)	0.56 ± 0.04	0.63 ± 0.05	0.57 ± 0.07
90%-10% Decay time (ms)	1.60 ± 0.17	1.52 ± 0.18	1.54 ± 0.17
Area (nA ms)	-0.035 ± 0.008	-0.032 ± 0.009	-0.032 ± 0.008


Table 3-11. Kinetic Parameters of Cholinergic Neurons in the Presence of 1 μM ACh. * denotes ($p < 0.05$)

Statistic	Control	1 μM ACh	Recovery
Peak Amplitude (pA)	-25.5 ± 3.4	-23.4 ± 1.1	-24.3 ± 2.0
10%-90% Rise time (ms)	0.67 ± 0.11	0.66 ± 0.06	0.62 ± 0.03
90%-10% Decay time (ms)	$1.65 \pm .010$	1.63 ± 0.07	1.82 ± 0.10
Area (nA ms)	-0.042 ± 0.006	-0.039 ± 0.002	-0.043 ± 0.004


A.




B.




C.






Control



2 μ M CNQX




Wash




40 pA
5 sec


A.




B.




C.



A.



B.



A.

Control



100 nM A β ₁₋₄₂




Wash




50 pA
5 sec


B.




C.



A.



B.



A.

Control



100 nM A β_{42-1}




Wash




100 pA

5 sec


B.




C.




A.




B.




A.



B.





A.

Control



100 nM A β_{1-42}



Wash



100 nM A β_{1-42} + 10 μ M DH β E








Wash



50 pA
5 sec


B.






A.

Control




100 μ M Nicotine




25 pA
5 sec

Wash




B.




A.

Control




100 μ M ACh




25 pA
5 sec

Wash



B.



REFERENCES

- Alkondon M, Pereira EF, Barbosa CT, Albuquerque EX (1997) Neuronal nicotinic acetylcholine receptor activation modulates gamma-aminobutyric acid release from CA1 neurons of rat hippocampal slices. *J Pharmacol Exp Ther* 283:1396-1411.
- Cao YJ, Surowy CS, Puttfarcken PS (2005) Different nicotinic acetylcholine receptor subtypes mediating striatal and prefrontal cortical [3H]dopamine release. *Neuropharmacology* 48:72-79.
- Chen GQ, Chen KS, Knox J, Inglis J, Bernard A, Martin SJ, Justice A, McConlogue L, Games D, Freedman SB, Morris RGM (2000) A learning deficit related to age and beta-amyloid plaques in a mouse model of Alzheimer's disease. *Nature* 408:975-979.
- Clements JD, Bekkers JM (1997) Detection of spontaneous synaptic events with an optimally scaled template. *Biophysical Journal* 73:220-229.
- Davies CA, Mann DMA, Sumpter PQ, Yates PO (1987) A Quantitative Morphometric Analysis Of The Neuronal And Synaptic Content Of The Frontal And Temporal Cortex In Patients With Alzheimers-Disease. *Journal Of The Neurological Sciences* 78:151-164.
- Dineley KT, Bell KA, Bui D, Sweatt JD (2002) beta -Amyloid peptide activates alpha 7 nicotinic acetylcholine receptors expressed in *Xenopus* oocytes. *J Biol Chem* 277:25056-25061.
- Dineley KT, Westerman M, Bui D, Bell K, Ashe KH, Sweatt JD (2001) beta-amyloid activates the mitogen-activated protein kinase cascade via hippocampal alpha 7

- nicotinic acetylcholine receptors: In vitro and in vivo mechanisms related to Alzheimer's disease. *Journal Of Neuroscience* 21:4125-4133.
- Easaw JC, Petrov T, Jhamandas JH (1997) An electrophysiological study of neurons in the horizontal limb of the diagonal band of Broca. *Am J Physiol* 272:C163-172.
- Frazier CJ, Buhler AV, Weiner JL, Dunwiddie TV (1998) Synaptic potentials mediated via alpha-bungarotoxin-sensitive nicotinic acetylcholine receptors in rat hippocampal interneurons. *Journal Of Neuroscience* 18:8228-8235.
- Fu W, Jhamandas JH (2003) Beta-amyloid peptide activates non-alpha7 nicotinic acetylcholine receptors in rat basal forebrain neurons. *J Neurophysiol* 90:3130-3136.
- Girod R, Barazangi N, McGehee D, Role LW (2000) Facilitation of glutamatergic neurotransmission by presynaptic nicotinic acetylcholine receptors. *Neuropharmacology* 39:2715-2725.
- Hamos JE, Degennaro LJ, Drachman DA (1989) Synaptic Loss In Alzheimers-Disease And Other Dementias. *Neurology* 39:355-361.
- Hartig W, Seeger J, Naumann T, Brauer K, Bruckner G (1998a) Selective in vivo fluorescence labelling of cholinergic neurons containing p75(NTR) in the rat basal forebrain. *Brain Res* 808:155-165.
- Hartig W, Seeger J, Naumann T, Brauer K, Bruckner G (1998b) Selective in vivo fluorescence labelling of cholinergic neurons containing p75(NTR) in the rat basal forebrain. *Brain Research* 808:155-165.
- Henderson Z, Boros A, Janzso GE, Westwood AJ, Monyer H, Halasy K (2005) Somato-dendritic nicotinic receptor responses recorded in vitro from the medial septal

diagonal band complex of the rodent. *Journal Of Physiology-London* 562:165-182.

Katzman R, Terry R, Deteresa R, Brown T, Davies P, Fuld P, Xiong RB, Peck A (1988) Clinical, Pathological, And Neurochemical Changes In Dementia - A Subgroup With Preserved Mental Status And Numerous Neocortical Plaques. *Annals Of Neurology* 23:138-144.

Lue LF, Kuo YM, Roher AE, Brachova L, Shen Y, Sue L, Beach T, Kurth JH, Rydel RE, Rogers J (1999) Soluble amyloid beta peptide concentration as a predictor of synaptic change in Alzheimer's disease. *American Journal Of Pathology* 155:853-862.

Masliah E, Terry RD, Alford M, Deteresa R, Hansen LA (1991) Cortical And Subcortical Patterns Of Synaptophysinlike Immunoreactivity In Alzheimers-Disease. *American Journal Of Pathology* 138:235-246.

Masliah E, Mallory M, Alford M, DeTeresa R, Hansen LA, McKeel DW, Morris JC (2001) Altered expression of synaptic proteins occurs early during progression of Alzheimer's disease. *Neurology* 56:127-129.

McGehee DS, Heath MJ, Gelber S, Devay P, Role LW (1995) Nicotine enhancement of fast excitatory synaptic transmission in CNS by presynaptic receptors. *Science* 269:1692-1696.

Morris JC, McKeel DW, Storandt M, Rubin EH, Price JL, Grant EA, Ball MJ, Berg L (1991) Very Mild Alzheimers-Disease - Informant-Based Clinical, Psychometric, And Pathological Distinction From Normal Aging. *Neurology* 41:469-478.

- Naslund J, Haroutunian V, Mohs R, Davis KL, Davies P, Greengard P, Buxbaum JD (2000) Correlation between elevated levels of amyloid beta-peptide in the brain and cognitive decline. *Jama-Journal Of The American Medical Association* 283:1571-1577.
- Pettit DL, Shao Z, Yakel JL (2001) beta-Amyloid(1-42) peptide directly modulates nicotinic receptors in the rat hippocampal slice. *Journal Of Neuroscience* 21.
- Phelan KD, Gallagher JP (1992) Direct Muscarinic And Nicotinic Receptor-Mediated Excitation Of Rat Medial Vestibular Nucleus Neurons In Vitro. *Synapse* 10:349-358.
- Selkoe DJ (2002) Alzheimer's disease is a synaptic failure. *Science* 298:789-791.
- Small DH, Mok SS, Bornstein JC (2001) Alzheimer's disease and AP toxicity: from top to bottom. *Nature Reviews Neuroscience* 2:595-598.
- Terry RD, Masliah E, Salmon DP, Butters N, Deteresa R, Hill R, Hansen LA, Katzman R (1991) Physical Basis Of Cognitive Alterations In Alzheimers-Disease - Synapse Loss Is The Major Correlate Of Cognitive Impairment. *Annals Of Neurology* 30:572-580.
- Walsh DM, Klyubin I, Fadeeva JV, Cullen WK, Anwyl R, Wolfe MS, Rowan MJ, Selkoe DJ (2002) Naturally secreted oligomers of amyloid beta protein potently inhibit hippocampal long-term potentiation in vivo. *Nature* 416:535-539.
- Wang HY, Lee DHS, Davis CB, Shank RP (2000a) Amyloid peptide A beta(1-42) binds selectively and with picomolar affinity to alpha 7 nicotinic acetylcholine receptors. *Journal Of Neurochemistry* 75:1155-1161.

- Wang HY, Lee DH, D'Andrea MR, Peterson PA, Shank RP, Reitz AB (2000b) beta-Amyloid(1-42) binds to alpha7 nicotinic acetylcholine receptor with high affinity. Implications for Alzheimer's disease pathology. *J Biol Chem* 275:5626-5632.
- Whitehouse PJ, Price DL, Struble RG, Clark AW, Coyle JT, Delon MR (1982) Alzheimer's disease and senile dementia: loss of neurons in the basal forebrain. *Science* 215:1237-1239.
- Wu J, Kuo YP, George AA, Xu L, Hu J, Lukas RJ (2004) beta-Amyloid directly inhibits human alpha4beta2-nicotinic acetylcholine receptors heterologously expressed in human SH-EP1 cells. *J Biol Chem* 279:37842-37851.
- Wu M, Shanabrough M, Leranath C, Alreja M (2000) Cholinergic excitation of septohippocampal GABA but not cholinergic neurons: implications for learning and memory. *J Neurosci* 20:3900-3908.
- Wu M, Hajszan T, Leranath C, Alreja M (2003) Nicotine recruits a local glutamatergic circuit to excite septohippocampal GABAergic neurons. *Eur J Neurosci* 18:1155-1168.
- Zhang M, Wang YT, Vyas DM, Neuman RS, Bieger D (1993) Nicotinic Cholinoceptor-Mediated Excitatory Postsynaptic Potentials In Rat Nucleus Ambiguus. *Experimental Brain Research* 96:83-88.

Chapter 4

General Discussion

This thesis demonstrates that the sum of the actions of A β in the cholinergic basal forebrain are expressed through nAChRs. In the first part of this study, the effect of A β_{1-42} potentiation on repeated activation of nAChR mediated [Ca²⁺]_i rises was examined. In the second part, A β_{1-42} was demonstrated to modulate glutamate-mediated synaptic transmission in fluorescently labeled cholinergic DBB neurons of the basal forebrain via nAChRs. As the cholinergic neurons in this area of the brain play an important role in memory and learning and undergo early neurodegeneration in AD, A β -nAChR interactions that result in changes to Ca²⁺ homeostasis and alteration of synaptic function in these cells may provide insights into the pathology responsible for the cognitive impairment in AD (Ridley et al., 1988; Paolini and McKenzie, 1993). The findings reported here illustrate the role of nAChRs in DBB neurons and suggests that nAChRs are an important receptor target for mediating the effects of A β_{1-42} that may have important implications in processes such as neurodegeneration and synaptic dysfunction.

A β Potentiates Repeated nAChR Mediated [Ca²⁺]_i Rises

A β_{1-42} potentiated nicotine-mediated [Ca²⁺]_i rises in acutely dissociated DBB neurons and was associated with a rise in basal Ca²⁺. Some studies have suggested that in certain preparations, A β_{1-42} is capable of forming Ca²⁺ ion channels, however the majority of these studies have either been in membrane bilayer systems or have only demonstrated the presence of A β_{1-42} insertion onto the plasma membrane (Arispe et al., 1993; Rhee et al., 1998; Kawahara et al., 2000). In DBB neurons focal application of A β_{1-42} did not cause any observable change in intracellular [Ca²⁺]_i. Previous studies have demonstrated that A β_{1-42} is capable of modulating activity of voltage gated Ca²⁺ channels (Price et al., 1998; Ramsden et al., 2002). The potentiation of nicotine-mediated [Ca²⁺]_i

rises in acutely dissociated DBB neurons was not mediated by $A\beta_{1-42}$ effects on voltage-gated Ca^{2+} channels as $[Ca^{2+}]_i$ rises evoked by focal application of KCl were not potentiated by $A\beta_{1-42}$. Both extracellular and intracellular sources of Ca^{2+} contributed to nicotine-mediated $[Ca^{2+}]_i$ rises. Pharmacological characterization of the nicotine-induced Ca^{2+} rises revealed that both $\alpha 7$ and $\alpha 4\beta 2$ nAChR subtypes mediated Ca^{2+} entry, however, there was a persistent component of Ca^{2+} rise that was not blocked by nAChR antagonists. The distribution of $\alpha 7$ and $\alpha 4$ nAChR subunits on DBB neurons was heterogeneous, which corresponded with the pharmacological data, with some cells displaying co-localization of the subunits on the same cell whereas others displayed labeling for either the $\alpha 7$ or the $\alpha 4$ -subunit. These results are in line with previous studies examining nicotinic modulation of DBB neuron properties (Henderson et al., 2005). Additionally, anatomical evidence suggests that there are collateral ACh terminals from cholinergic neurons projecting to the hippocampus that mediate synapses onto DBB neurons (Henderson et al., 2001).

Further examination of the nicotine induced $[Ca^{2+}]_i$ rise revealed that emptying of intracellular Ca^{2+} stores by inhibiting the SERCA pump, which is responsible for filling of Ca^{2+} stores, inhibits nicotine mediated $[Ca^{2+}]_i$ rises. The nicotine response could also be inhibited by xestospongine, an IP₃ receptor blocker. This antagonist was previously used to demonstrate that $\alpha 7$ nAChRs were coupled to IP₃-sensitive stores, although given the data that xestospongine may also serve to inhibit SERCA pumps, this conclusion should be viewed with caution (De Smet et al., 1999; Castonguay and Robitaille, 2002; Dajas-Bailador et al., 2002; Solovyova et al., 2002).

$A\beta_{1-42}$ was also able to modulate Ca^{2+} -induced Ca^{2+} entry. Repeated application of caffeine to DBB neurons activated intracellular Ca^{2+} stores that could be inhibited by TG, which has been previously reported (Murchison and Griffith, 1999). In the presence of $A\beta_{1-42}$, the release of Ca^{2+} from these caffeine-sensitive stores were temporarily potentiated. This suggests that $A\beta_{1-42}$ was overfilling these caffeine-sensitive stores to the point where caffeine-sensitive intracellular stores were unable to buffer $[Ca^{2+}]_i$. However, unlike in the case of repeated nicotine-activation of intracellular Ca^{2+} stores, caffeine does not allow extracellular Ca^{2+} entry and thus there was no refilling of caffeine-sensitive stores. With no exogenous source of Ca^{2+} to replenish caffeine sensitive stores, the caffeine induced Ca^{2+} rise did not remain potentiated as with nicotine-evoked Ca^{2+} responses. It appears somewhat paradoxical that $A\beta_{1-42}$ was able to potentiate caffeine mediated Ca^{2+} rises, given that it could not potentiate Ca^{2+} rises mediated by KCl activation of voltage-gated Ca^{2+} channels. However, previous studies examining voltage-gated Ca^{2+} channels in DBB neurons revealed that Ca^{2+} entry via these channels is significantly buffered by the ER which prevents Ca^{2+} from activating Ca^{2+} release channels on intracellular Ca^{2+} stores (Murchison and Griffith, 1998).

$A\beta_{1-42}$ potentiation of nicotine-evoked $[Ca^{2+}]_i$ was mediated by its actions on TG sensitive intracellular Ca^{2+} stores. Blockade of these stores by CGP-37157, an inhibitor of the mitochondrial Na^+/Ca^{2+} exchanger after $A\beta_{1-42}$ potentiation of nicotine-mediated Ca^{2+} resulted in the blockade of the $A\beta_{1-42}$ potentiation. The decrease in the potentiation by Na^+/Ca^{2+} exchanger blockade was of similar magnitude to that observed with TG inhibition of the SERCA ATPase, which suggests that most of $A\beta_{1-42}$ potentiation of intracellular Ca^{2+} stores originates from overfilling of the TG-sensitive stores by

mitochondria. Thus we propose that A β_{1-42} acts on mitochondria by causing the mitochondria to overflow with Ca²⁺, which results in the excess release of Ca²⁺ via the Na⁺/Ca²⁺ exchanger, which, in turn is taken up into the TG-sensitive stores by the action of the SERCA pumps (Figure 4-1). It has been demonstrated that mitochondria positioned close to these ER stores are responsible for buffering and refilling intracellular Ca²⁺ stores (Rizzuto et al., 1998). These observations are in line with previous studies which have demonstrated that blocking the Na⁺/Ca²⁺ exchanger impairs Ca²⁺ release and that the Na⁺/Ca²⁺ exchanger is localized in close proximity to SERCA pumps on the TG-sensitive stores (Arnaudeau et al., 2001; Csordas and Hajnoczky, 2001).

In this portion of experimental studies we have demonstrated that A β_{1-42} alters Ca²⁺ homeostasis in DBB neurons via its effects on mitochondria (Figure 4-1). The effect of A β_{1-42} on these mitochondria may be due to A β_{1-42} interaction with a surface receptor such as the P75 or RAGE receptor (Yan et al., 1996; Yaar et al., 1997) or occur in a two step process with A β_{1-42} entering the neuron by endocytosis, perhaps via the $\alpha 7$ nAChR (Wang et al., 2002) and then binding to an intracellular receptor. Modulation of mitochondrial Ca²⁺ is important in the regulation of neuronal apoptosis. A β modulation of mitochondria by A β_{1-42} has previously been shown to potentiate glutamate excitotoxicity and to trigger an apoptotic pathway involving the ER and caspases associated with this cellular organelle (Mattson et al., 1992; Nakagawa et al., 2000). Additionally, A β_{1-42} modulation of the mitochondrial enzyme, A β -binding alcohol dehydrogenase, has been shown to impair the function of this enzyme in AD patients and to increase oxidative stress in AD transgenic mice (Lustbader et al., 2004).

A β Modulates Synaptic Transmission in Cholinergic DBB Neurons

In the second part of my project, the effect of A β_{1-42} on spontaneous synaptic transmission in cholinergic neurons of the DBB was examined. Recordings of glutamate-mediated mEPSCs from fluorescently labeled cholinergic neurons in the DBB were examined. Application of A β_{1-42} revealed cholinergic neurons of the DBB respond to A β_{1-42} in two different ways. In the majority of neurons, A β_{1-42} increased the frequency of mEPSCs. In a second population of neurons, A β_{1-42} decreased the frequency of mEPSCs. In both groups of A β -responsive cells, A β_{1-42} did not significantly change kinetic parameters such as rise time, decay time or amplitude, suggesting that A β_{1-42} modulation of synaptic activity occurs via a presynaptic effect (Figure 4-2).

The increase in mEPSC frequency induced by A β_{1-42} could be blocked in the presence of DH β E, a non- $\alpha 7$ nAChR antagonist, suggesting that A β_{1-42} mediates this effect on synaptic transmission via a non- $\alpha 7$, $\alpha 4\beta 2$ mechanism as the dose of DH β E used is in the low enough range to affect $\alpha 4\beta 2$ nAChRs and not the other subtypes (Sharples and Wonnacott, 2001; Figure 4-2A). Previous electrophysiological studies examining non- $\alpha 7$ and $\alpha 7$ nAChR-A β_{1-42} effects have been performed on expression systems and acutely dissociated neurons, and thus unable to assess the effect of modulation of synaptic activity. This study is the first to demonstrate that A β_{1-42} is capable of altering spontaneous synaptic activity. As the basal forebrain plays an important role in learning and memory, alteration of synaptic function would interfere with cognition (Ridley et al., 1988; Paolini and McKenzie, 1993). Based upon this electrophysiology data, A β_{1-42} can increase glutamate neurotransmission in a population of cholinergic neurons, while decreasing glutamate neurotransmission in another population of cholinergic neurons.


Although this would lead to an increase of ACh release from some cholinergic neurons, the increase in ACh would not necessarily be able to compensate for the loss of cholinergic tone from $A\beta_{1-42}$ inhibition of the other population of cholinergic neurons, as these neurons would synapse on different hippocampal neurons. Thus alteration of synaptic function in cholinergic neurons of the DBB by soluble $A\beta_{1-42}$ may provide the pathological basis for the mild cognitive dysfunction observed at the beginning of AD prior to the deposition of $A\beta_{1-42}$ plaques.

Application of other nicotinic agonists onto cholinergic neurons demonstrated that both nicotine and acetylcholine were able to modulate synaptic transmission in a manner similar to that observed with $A\beta_{1-42}$. This modulation is presynaptic in nature as only increases in mEPSC frequency were observed in this group of cells. It has been previously reported that nAChRs mediate post-synaptic actions, although these studies did not specifically rule out pre-synaptic modulation. One study examined the effect of nicotine application on acutely dissociated DBB neurons and whereas another examined single channel currents, thus these studies were not specifically looking for presynaptic modulation of neurotransmitter release (Fu and Jhamandas, 2003; Henderson et al., 2005).


Future Experiments and Conclusions

In my thesis, two separate $A\beta_{1-42}$ effects were demonstrated. Firstly, $A\beta_{1-42}$ was shown to alter and possibly disrupt Ca^{2+} homeostasis via mediating overfilling of SERCA stores through the release of Ca^{2+} via the Na^+/Ca^{2+} exchanger. Secondly, $A\beta_{1-42}$ was able to modulate excitatory synaptic function in the basal forebrain. With regards to the first set of experiments, significant questions remain as to the role of experiments measuring


mitochondrial membrane potential and mitochondrial Ca^{2+} may provide more information about the effect of $\text{A}\beta_{1-42}$ on mitochondria and the relationship of this to ER stress which has been postulated to be a major player in neuronal death evoked by $\text{A}\beta$ and other similarly misfolded proteins. Further experiments examining mitochondria will provide additional insights into how $\text{A}\beta_{1-42}$ may overfill mitochondria Ca^{2+} . Examining potential targets through which $\text{A}\beta_{1-42}$ mediates a decrease in mEPSC frequency also need to be performed. Furthermore, pharmacological characterization of the nAChR responses in the DBB needs to be examined as targeting specific subtypes of nAChRs may become important to AD therapy.



A.



B.



REFERENCES

- Arispe N, Rojas E, Pollard HB (1993) Alzheimer disease amyloid beta protein forms calcium channels in bilayer membranes: blockade by tromethamine and aluminum. *Proc Natl Acad Sci U S A* 90:567-571.
- Arnaudeau S, Kelley WL, Walsh JV, Demaurex N (2001) Mitochondria recycle Ca^{2+} to the endoplasmic reticulum and prevent the depletion of neighboring endoplasmic reticulum regions. *Journal Of Biological Chemistry* 276:29430-29439.
- Castonguay A, Robitaille R (2002) Xestospongine C is a potent inhibitor of SERCA at a vertebrate synapse. *Cell Calcium* 32:39-47.
- Csordas G, Hajnoczky G (2001) Sorting of calcium signals at the junctions of endoplasmic reticulum and mitochondria. *Cell Calcium* 29:249-262.
- Dajas-Bailador FA, Mogg AJ, Wonnacott S (2002) Intracellular Ca^{2+} signals evoked by stimulation of nicotinic acetylcholine receptors in SH-SY5Y cells: contribution of voltage-operated Ca^{2+} channels and Ca^{2+} stores. *J Neurochem* 81:606-614.
- De Smet P, Parys JB, Callewaert G, Weidema AF, Hill E, De Smedt H, Erneux C, Sorrentino V, Missiaen L (1999) Xestospongine C is an equally potent inhibitor of the inositol 1,4,5-trisphosphate receptor and the endoplasmic-reticulum Ca^{2+} pumps. *Cell Calcium* 26:9-13.
- Fu W, Jhamandas JH (2003) Beta-amyloid peptide activates non- $\alpha 7$ nicotinic acetylcholine receptors in rat basal forebrain neurons. *J Neurophysiol* 90:3130-3136.

- Henderson Z, Morris NP, Grimwood P, Fiddler G, Yang HW, Appenteng K (2001) Morphology of local axon collaterals of electrophysiologically characterised neurons in the rat medial septal/ diagonal band complex. *J Comp Neurol* 430:410-432.
- Henderson Z, Boros A, Janzso GE, Westwood AJ, Monyer H, Halasy K (2005) Somato-dendritic nicotinic receptor responses recorded in vitro from the medial septal diagonal band complex of the rodent. *Journal Of Physiology-London* 562:165-182.
- Kawahara M, Kuroda Y, Arispe N, Rojas E (2000) Alzheimer's beta-amyloid, human islet amylin, and prion protein fragment evoke intracellular free calcium elevations by a common mechanism in a hypothalamic GnRH neuronal cell line. *J Biol Chem* 275:14077-14083.
- Lustbader JW, Cirilli M, Lin C, Xu HW, Takuma K, Wang N, Caspersen C, Chen X, Pollak S, Chaney M, Trinchese F, Liu SM, Gunn-Moore F, Lue LF, Walker DG, Kuppusamy P, Zewier ZL, Arancio O, Stern D, Yan SSD, Wu H (2004) ABAD directly links A beta to mitochondrial toxicity in Alzheimer's disease. *Science* 304:448-452.
- Mattson MP, Cheng B, Davis D, Bryant K, Lieberburg I, Rydel RE (1992) Beta-Amyloid Peptides Destabilize Calcium Homeostasis And Render Human Cortical-Neurons Vulnerable To Excitotoxicity. *Journal Of Neuroscience* 12:376-389.
- Murchison D, Griffith WH (1998) Increased calcium buffering in basal forebrain neurons during aging. *J Neurophysiol* 80:350-364.

- Murchison D, Griffith WH (1999) Age-related alterations in caffeine-sensitive calcium stores and mitochondrial buffering in rat basal forebrain. *Cell Calcium* 25:439-452.
- Nakagawa T, Zhu H, Morishima N, Li E, Xu J, Yankner BA, Yuan JY (2000) Caspase-12 mediates endoplasmic-reticulum-specific apoptosis and cytotoxicity by amyloid-beta. *Nature* 403:98-103.
- Paolini AG, McKenzie JS (1993) Effects of lesions in the horizontal diagonal band nucleus on olfactory habituation in the rat. *Neuroscience* 57:717-724.
- Price SA, Held B, Pearson HA (1998) Amyloid beta protein increases Ca²⁺ currents in rat cerebellar granule neurones. *Neuroreport* 9:539-545.
- Ramsden M, Henderson Z, Pearson HA (2002) Modulation of Ca²⁺ channel currents in primary cultures of rat cortical neurones by amyloid beta protein (1-40) is dependent on solubility status. *Brain Research* 956:254-261.
- Rhee SK, Quist AP, Lal R (1998) Amyloid beta protein-(1-42) forms calcium-permeable, Zn²⁺-sensitive channel. *J Biol Chem* 273:13379-13382.
- Ridley RM, Samson NA, Baker HF, Johnson JA (1988) Visuospatial learning impairment following lesion of the cholinergic projection to the hippocampus. *Brain Res* 456:71-87.
- Rizzuto R, Pinton P, Carrington W, Fay FS, Fogarty KE, Lifshitz LM, Tuft RA, Pozzan T (1998) Close contacts with the endoplasmic reticulum as determinants of mitochondrial Ca²⁺ responses. *Science* 280:1763-1766.
- Sharples C, Wonnacott S (2001) Neuronal Nicotinic Receptors. *Toctris Reviews* 19:1-12.

- Solovyova N, Fernyhough P, Glazner G, Verkhratsky A (2002) Xestospongins empty the ER calcium store but do not inhibit InsP₃-induced Ca²⁺ release in cultured dorsal root ganglia neurones. *Cell Calcium* 32:49-52.
- Wang HY, D'Andrea MR, Nagele RG (2002) Cerebellar diffuse amyloid plaques are derived from dendritic A β 42 accumulations in Purkinje cells. *Neurobiology Of Aging* 23:213-223.
- Yaar M, Zhai S, Pilch PF, Doyle SM, Eisenhauer PB, Fine RE, Gilchrist BA (1997) Binding of beta-amyloid to the p75 neurotrophin receptor induces apoptosis - A possible mechanism for Alzheimer's disease. *Journal Of Clinical Investigation* 100:2333-2340.
- Yan SD, Chen X, Fu J, Chen M, Zhu HJ, Roher A, Slattery T, Zhao L, Nagashima M, Morser J, Migheli A, Nawroth P, Stern D, Schmidt AM (1996) RAGE and amyloid-beta peptide neurotoxicity in Alzheimer's disease. *Nature* 382:685-691.