

WILDLIFE HABITAT
REQUIREMENT SUMMARIES
FOR
SELECTED WILDLIFE SPECIES
IN ALBERTA

Prepared by
M. Nietfeld, J. Wilk, K. Woolnough
and B. Hoskin

WILDLIFE RESOURCE INVENTORY UNIT

1985
Edmonton

Alberta
ENERGY AND
NATURAL RESOURCES
Fish and Wildlife Division

ENR Technical Report Number: T/73
International Standard Book Number: 0-86499-985-2

FOR ADDITIONAL COPIES OF THIS REPORT, CONTACT:

Information Centre
Alberta Energy and Natural Resources
Main Floor, Bramalea Building
9920 - 108 Street
Edmonton, Alberta, Canada T5K 2M4

Telephone: (403) 427-5590

3590

TABLE OF CONTENTS

	Page
1. SPECIES HABITAT REQUIREMENT SUMMARIES: AN INFORMATION BASE	1
2. KEY HABITAT REQUIREMENTS FOR WHITE-TAILED DEER:	
2.1 GENERAL	7
2.2 COVER	7
2.2.1 Vegetation	7
2.2.2 Land Forms and Topography	8
2.2.3 Climate	8
2.3 FOOD	8
2.3.1 Vegetation	8
Grassland	8
Parkland	9
Forest	9
2.3.2 Land Forms and Topography	9
2.3.3 Aquatic Forms	9
2.3.4 Climate	9
2.4 SPACE	10
2.4.1 Territory/Home Range	10
2.4.2 Population Density	10
2.5 SPECIAL CONSIDERATIONS	10
2.5.1 Size, Shape and Juxtaposition of Habitat Components	10
2.5.2 Significance of Disturbance Phenomena	10
2.6 LIMITING FACTORS	11
2.7 REGIONAL VARIATIONS	11
3. KEY HABITAT REQUIREMENTS FOR MULE DEER:	
3.1 GENERAL	19
3.2 COVER	19
3.2.1 Vegetation	19
3.2.2 Land Forms and Topography	19
3.2.3 Climate	20
3.3 FOOD	20
3.3.1 Vegetation	20
3.3.2 Land Forms and Topography	20
3.3.3 Aquatic Forms	21
3.3.4 Climate	21
3.3.5 Trace Elements	21

	Page
3.4 SPACE	21
3.4.1 Territory/Home Range	21
3.4.2 Population Densities	21
3.5 SPECIAL CONSIDERATIONS	22
3.5.1 Size, Shape and Juxtaposition of Habitat Components	22
3.5.2 Significance of Disturbance Phenomena	22
3.6 LIMITING FACTORS	23
3.7 REGIONAL VARIATIONS	23
4. KEY HABITAT REQUIREMENTS FOR MOOSE:	
4.1 GENERAL	31
4.2 COVER	31
4.2.1 Vegetation	31
4.2.2 Land Forms and Topography	31
4.3 FOOD	32
4.3.1 Vegetation	32
4.3.2 Aquatic Forms	32
4.3.3 Trace Elements	32
4.4 SPACE	33
4.4.1 Territory/Home Range	33
4.4.2 Population Density	33
4.5 SPECIAL CONSIDERATIONS	33
4.5.1 Size, Shape and Juxtaposition of Habitat Components	33
4.5.2 Significance of Disturbance Phenomena	33
4.6 LIMITING FACTORS	34
4.7 REGIONAL VARIATIONS	34
5. KEY HABITAT REQUIREMENTS FOR ELK:	
5.1 GENERAL	43
5.2 COVER	43
5.2.1 Vegetation	43
5.2.2 Land Forms and Topography	43
5.2.3 Climate	44
5.3 FOOD	44
5.3.1 Vegetation	44
5.3.2 Land Forms and Topography	45
5.3.3 Aquatic Forms	45
5.3.4 Climate	45
5.3.5 Trace Elements	45
5.4 SPACE	45
5.4.1 Territory/Home Range	45
5.4.2 Seasonal Population Densities	45

	Page
5.5 SPECIAL CONSIDERATIONS	46
5.5.1 Size, Shape and Juxtaposition of Habitat Components	46
5.5.2 Significance of Disturbance Phenomena	46
5.6 LIMITING FACTORS	47
5.7 REGIONAL VARIATIONS	47
6. KEY HABITAT REQUIREMENTS FOR WOODLAND/MOUNTAIN CARIBOU:	
6.1 GENERAL	53
6.2 COVER	53
6.2.1 Vegetation	53
6.2.2 Land Forms and Topography	53
6.3 FOOD	54
6.3.1 Vegetation	54
6.3.2 Land Forms and Topography	54
6.3.3 Aquatic Forms	54
6.3.4 Climate	54
6.4 SPACE	54
6.4.1 Territory/Home Range	54
6.4.2 Population Densities	55
6.5 SPECIAL CONSIDERATIONS	55
6.5.1 Significance of Disturbance Phenomena	55
6.6 LIMITING FACTORS	56
7. KEY HABITAT REQUIREMENTS FOR PRONGHORN ANTELOPE:	
7.1 GENERAL	63
7.2 COVER	63
7.2.1 Vegetation	63
7.2.2 Land Forms and Topography	63
7.2.3 Climate	64
7.3 FOOD	64
7.3.1 Vegetation	64
7.3.2 Aquatic Forms	65
7.3.3 Climate	65
7.3.4 Trace Elements	65
7.4 SPACE	65
7.4.1 Territory/Home Range	65
7.4.2 Population Densities	65
7.5 SPECIAL CONSIDERATIONS	66
7.5.1 Size, Shape and Juxtaposition of Habitat Components	66
7.5.2 Significance of Disturbance Phenomena	66
7.6 LIMITING FACTORS	66

	Page
8. KEY HABITAT REQUIREMENTS FOR ROCKY MOUNTAIN BIGHORN SHEEP:	
8.1 GENERAL	75
8.2 COVER	75
8.2.1 Vegetation	75
8.2.2 Land Forms and Topography	75
8.3 FOOD	76
8.3.1 Vegetation	76
8.3.2 Land Forms and Topography	76
8.3.3 Aquatic Forms	77
8.3.4 Climate	77
8.3.5 Trace Elements	77
8.4 SPACE	77
8.4.1 Territory/Home Range	77
8.4.2 Population Density	78
8.5 SPECIAL CONSIDERATIONS	78
8.5.1 Size, Shape and Juxtaposition of Habitat Components	78
8.5.2 Significance of Disturbance Phenomena	78
8.6 LIMITING FACTORS	79
9. KEY HABITAT REQUIREMENTS FOR MOUNTAIN GOAT:	
9.1 GENERAL	87
9.2 COVER	87
9.2.1 Vegetation	87
9.2.2 Land Forms and Topography	87
9.2.3 Climate	88
9.3 FOOD	88
9.3.1 Vegetation	88
9.3.2 Land Forms and Topography	88
9.3.3 Aquatic Forms	89
9.3.4 Climate	89
9.3.5 Trace Elements	89
9.4 SPACE	89
9.4.1 Territory/Home Range	89
9.5 SPECIAL CONSIDERATION	90
9.5.1 Size, Shape and Juxtaposition of Habitat Components	90
9.5.2 Significance of Disturbance Phenomena	90
9.6 LIMITING FACTORS	90
10. KEY HABITAT REQUIREMENTS FOR GRIZZLY BEAR:	
10.1 GENERAL	97
10.2 COVER	97
10.2.1 Vegetation	97
10.2.2 Land Forms and Topography	98
10.2.3 Aquatic Forms	98
10.2.4 Climate	98

	Page
10.3 FOOD	98
10.3.1 Vegetation/Animal Matter	98
10.3.2 Land Forms and Topography	99
10.3.3 Aquatic Forms	99
10.3.4 Climate	100
10.4 SPACE	100
10.4.1 Territory/Home Range	100
10.4.2 Population Densities	100
10.5 SPECIAL CONSIDERATIONS	101
10.5.1 Size, Shape and Juxtaposition of Habitat Components	101
10.5.2 Significance of Disturbance Phenomena	101
10.6 LIMITING FACTORS	102
10.7 REGIONAL VARIATION	102
 11. KEY HABITAT REQUIREMENT FOR BEAVER	
11.1 GENERAL	111
11.2 COVER	111
11.2.1 Vegetation	111
11.2.2 Land Forms and Topography	111
11.2.3 Aquatic Forms	111
11.2.4 Climate	112
11.3 FOOD	112
11.3.1 Vegetation	112
11.3.2 Land Forms and Topography	113
11.3.3 Aquatic Forms	113
11.4 SPACE	113
11.4.1 Population Densities	113
11.5 SPECIAL CONSIDERATIONS	113
11.5.1 Size, Shape and Juxtaposition of Habitat Components	113
11.5.2 Significance of Disturbance Phenomena	114
11.6 SIGNIFICANT LIMITNG FACTORS	114
11.7 REGIONAL VARIATIONS	115
 12. KEY HABITAT REQUIREMENTS FOR RIVER OTTER:	
12.1 GENERAL	123
12.2 COVER	123
12.2.1 Vegetation	123
12.2.2 Land Forms and Topography	123
12.2.3 Climate	124
12.2.4 Aquatic Forms	124
12.3 FOOD	125
12.3.1 Vegetation	125
12.3.2 Aquatic Forms	125
12.3.3 Climate	126

	Page
12.4 SPACE	126
12.4.1 Territory/Home Range	126
12.4.2 Population Densities	127
12.5 SPECIAL CONSIDERATIONS	127
12.5.1 Size, Shape and Juxtaposition of Habitat Components	127
12.5.2 Significance of Disturbance Phenomena	127
12.6 LIMITING FACTORS	128
12.7 REGIONAL VARIATION	128
 13. KEY HABITAT REQUIREMENTS FOR MARTEN:	
13.1 GENERAL	139
13.2 COVER	139
13.2.1 Vegetation	139
13.2.2 Land Forms and Topography	140
13.2.3 Climate	140
13.3 FOOD	140
13.3.1 Vegetation/Animal Matter	140
13.3.2 Climate	141
13.4 SPACE	141
13.4.1 Territory/Home Range	141
13.4.2 Population Densities	142
13.5 SPECIAL CONSIDERATIONS	142
13.5.1 Size, Shape and Juxtaposition of Habitat Components	142
13.5.2 Significance of Disturbance Phenomena	143
13.6 LIMITING FACTORS	143
13.7 REGIONAL VARIATION	143
 14. KEY HABITAT REQUIREMENTS FOR SHARP-TAILED GROUSE:	
14.1 GENERAL	155
14.2 COVER	155
14.2.1 Vegetation	155
14.2.2 Land Forms and Topography	157
14.2.3 Aquatic Forms	157
14.2.4 Climate	157
14.3 FOOD	157
14.3.1 Vegetation	157
14.3.2 Trace Elements	158
14.4 SPACE	159
14.4.1 Territory/Home Range	159
14.5 SPECIAL CONSIDERATIONS	159
14.5.1 Size, Shape and Juxtaposition of Habitat Components	159
14.5.2 Significance of Disturbance Phenomena	160
14.6 LIMITING FACTORS	160
14.7 REGIONAL VARIATION	161

	Page
15. KEY HABITAT REQUIREMENTS FOR RING-NECKED PHEASANT:	
15.1 GENERAL	169
15.2 COVER	169
15.2.1 Vegetation	169
15.2.2 Land Forms and Topography	170
15.2.3 Aquatic Forms	170
15.2.4 Climate	171
15.3 FOOD	171
15.3.1 Vegetation	171
15.3.2 Aquatic Forms	171
15.3.3 Climate	171
15.3.4 Trace Elements	172
15.4 SPACE	172
15.4.1 Territory/Home Range	172
15.5 SPECIAL CONSIDERATIONS	172
15.5.1 Size, Shape and Juxtaposition of Habitat Components	172
15.5.2 Significance of Disturbance Phenomena	173
15.6 LIMITING FACTORS	173
15.7 REGIONAL VARIATIONS	173
16. KEY HABITAT REQUIREMENTS FOR SAGE GROUSE:	
16.1 GENERAL	181
16.2 COVER	181
16.2.1 Vegetation	181
16.2.2 Land Forms and Topography	182
16.2.3 Climate	182
16.3 FOOD	183
16.3.1 Vegetation	183
16.3.2 Aquatic Forms	183
16.3.3 Climate	184
16.4 SPACE	184
16.4.1 Territory/Home Range	184
16.4.2 Population Densities	185
16.5 SPECIAL CONSIDERATIONS	185
16.5.1 Size, Shape and Juxtaposition of Habitat Components	185
16.5.2 Significance of Disturbance Phenomena	185
16.6 LIMITING FACTORS	186
16.7 REGIONAL VARIATION	187
17. KEY HABITAT REQUIREMENTS FOR RESIDENT BREEDING DABBING DUCKS:	
17.1 GENERAL	195

	Page
17.2 COVER	195
17.2.1 Vegetation	195
17.2.2 Land Forms and Topography	196
17.2.3 Aquatic Forms	196
17.2.4 Climate	196
17.3 FOOD	197
17.3.1 Vegetation	197
17.3.2 Land Forms and Topography	197
17.3.3 Aquatic Forms	197
17.4 SPACE	198
17.4.1 Territory/Home Range	198
17.4.2 Population Densities	198
17.5 SPECIAL CONSIDERATIONS	198
17.5.1 Size, Shape and Juxtaposition of Habitat Components	198
17.5.2 Significance of Disturbance Phenomena	199
17.6 LIMITING FACTORS	200
17.7 REGIONAL VARIATIONS	200
18. KEY HABITAT REQUIREMENTS FOR BREEDING CANADA GEESE:	
18.1 GENERAL	215
18.2 COVER	215
18.2.1 Vegetation	215
18.2.2 Land Forms and Topography	216
18.2.3 Aquatic Forms	216
18.2.4 Climate	217
18.3 FOOD	217
18.3.1 Vegetation	217
18.3.2 Land Forms and Topography	218
18.3.3 Aquatic Forms	218
18.3.4 Climate	218
18.3.5 Trace Elements	218
18.4 SPACE	218
18.4.1 Territory/Home Range	218
18.4.2 Population Densities	219
18.5 SPECIAL CONSIDERATIONS	219
18.5.1 Size, Shape and Juxtaposition of Habitat Components	219
18.5.2 Significance of Disturbance Phenomena	220
18.6 LIMITING FACTORS	220
18.7 REGIONAL VARIATION	221
19. KEY HABITAT REQUIREMENTS FOR MOULTING, STAGING, AND MIGRANT WATERFOWL:	
19.1 GENERAL	231

	Page
19.2 COVER	231
19.2.1 Vegetation	231
19.2.2 Land Forms and Topography	231
19.2.3 Aquatic Forms	232
19.2.4 Climate	232
19.3 FOOD	232
19.3.1 Vegetation	232
19.3.2 Land Forms and Topography	233
19.3.3 Aquatic Forms	233
19.3.4 Climate	233
19.3.5 Trace Elements	233
19.4 SPACE	234
19.4.1 Territory/Home Range	234
19.4.2 Population Densities	234
19.5 SPECIAL CONSIDERATIONS	235
19.5.1 Size, Shape and Juxtaposition of Habitat Components	235
19.5.2 Significance of Disturbance Phenomena	235
19.6 LIMITING FACTORS	235
19.7 REGIONAL VARIATIONS	236

APPENDICES

1. Common and scientific names of mammals referred to in the text	255
2. Common and scientific names of birds referred to in the text	256
3. Common and scientific names of plants referred to in the text	257

LIST OF ILLUSTRATIONS

Figures

Figure	Page
1. Wildlife Habitat Regions of Alberta	3
2. General Distribution of Habitat Quality for White-Tailed Deer in Alberta	15
3. General Distribution of Habitat Quality for Mule Deer in Alberta	27
4. General Distribution of Habitat Quality for Moose in Alberta	39
5. General Distribution of Habitat Quality for Elk in Alberta	49
6. General Distribution of Habitat Quality for Caribou in Alberta	59
7. General Distribution of Habitat Quality for Pronghorn Antelope in Alberta	71
8. General Distribution of Habitat Quality for Bighorn Sheep in Alberta	83
9. General Distribution of Habitat Quality for Mountain Goat in Alberta	93
10. General Population Distribution for Grizzly Bear in Alberta	107
11.a General Distribution of Habitat Quality for Beaver in Alberta	117
11.b Distribution of Beaver Harvest, 1970-1975	119
12.a General Distribution of Habitat Quality for River Otter in Alberta	133
12.b Distribution of River Otter Harvest	135

	Page
13.a General Distribution of Habitat Quality for Marten in Alberta	149
13.b Distribution of Marten Harvest, 1970-1975	151
14. General Distribution of Habitat Quality for Sharp-tailed Grouse in Alberta	165
15. General Distribution of Habitat Quality for Ring-Necked Pheasant in Alberta	177
16. General Distribution of Habitat Quality for Sage Grouse in Alberta	191
17.a Distribution and Relative Abundance of Waterfowl in Alberta (1982)	205
17.b Priority Duck Production Habitat in Alberta	207
17.c Relative Total Duck Harvest Areas in Alberta (1977-80)	211
18. Priority Goose Production Habitat in Alberta	225
19.a Priority Duck Staging Habitat in Alberta	239
19.b Priority Duck Moulting Habitat in Alberta	243
19.c Priority Goose Staging Habitat in Alberta	247
19.d Priority Goose Moulting Habitat in Alberta	251

LIST OF TABLES

Table	Page
1. Mean Densities of Dabbler Ducks for Various Ecological Zones in Alberta.	198
2. Priority Duck Production Habitat in Alberta.	209
3. Priority Goose Production Habitat in Alberta.	227
4. Priority Duck Staging Habitat in Alberta.	241
5. Priority Duck Moulting Habitat in Alberta.	245
6. Priority Goose Staging Habitat in Alberta.	249
7. Priority Goose Moulting Habitat in Alberta.	253

ACKNOWLEDGMENTS

Numerous species experts provided critical comment and review of summaries for particular wildlife species or species groups - these individuals are specifically acknowledged at the end of the summaries they contributed to. Harry Stelfox (Wildlife Branch) and Marilyn Brown (ENR Public Affairs) provided editorial advice. Cartographic Services (ENR Resource Evaluation and Planning Division) provided assistance in preparing most of the figures.

1. SPECIES HABITAT REQUIREMENT SUMMARIES: AN INFORMATION BASE

This manual attempts to summarize and highlight information to assist in defining species/habitat relationships relevant to Alberta environments and for a number of Alberta wildlife species. The degree of specificity of the summaries, in part or whole, varies according to the background information available. Information must often be extrapolated from studies outside Alberta, or from specific regions in Alberta; this should be taken into consideration when applying this information. The summaries have been reviewed by species experts and their comments have been incorporated. Habitat distribution maps follow the summaries; a comparative Wildlife Habitat Region map for Alberta (Figure 1) is provided.

It is hoped this manual will serve as an information base, and as a reference guide for habitat mapping and evaluation activities. The summaries are being used as an integral part in the development of habitat interpretation models which attempt to establish quantifiable value relationships between landscape features and species life requisites. These models will allow the classification of habitat suitability for a particular region on a species basis. The background information provided in the summaries supports the relative values assigned within the models for a particular habitat region.

Wildlife requires a number of life support requisites for survival and reproduction. Habitat features that will supply food, cover, and space will determine a species' presence or absence from an area, as well as influencing the relative abundance of a species. Food (vegetative or animal matter, water, and trace minerals) should provide all the nutritional requirements necessary for growth and development, maintenance and reproduction. Cover may be required for thermal, reproductive, escape, resting and roosting purposes. A number of landscape components, such as vegetation, land forms, topography, and aquatic forms, may serve these functions. Space is the area or range an animal requires to satisfy these basic requirements during its lifetime.

The habitat requirements of a species vary seasonally, and with sex, age, and reproductive status. As well, many habitat components (vegetation, snow cover) undergo dramatic seasonal changes or gradual successional changes that will temporally influence the suitability of a region. Disturbance phenomena, natural as wildfires or human-induced as timber harvest, may also have a significant effect on habitat suitability. Effects may be species-specific or species-general, and harmful or beneficial to varying degrees.

Species characteristics will determine the components of the habitat essential to its existence. Species plasticity influences the range of habitat conditions that a species can adapt to. Sage grouse, for example, are dependent on expanses of sagebrush and cannot exist elsewhere. Elk, in contrast, are a generalistic species able to utilize a wide variety of habitats under varying conditions. Before accurate evaluations of wildlife habitat of such species can be made, an adequate information base of their habitat needs, and the factors influencing these needs, is required.




Figure 1. Wildlife Habitat Regions of Alberta.

WHITE-TAILED DEER



2. KEY HABITAT REQUIREMENTS FOR WHITE-TAILED DEER

2.1 General

White-tailed deer (*Odocoileus virginianus*), a very adaptable species, is found in three major ecological zones: grassland, aspen parkland and boreal mixedwood forest. Its optimum habitat is the aspen parkland where woody cover, diverse in height and species composition, and containing a high frequency of edge, occurs in proximity to farmland so both food and cover requirements are adequately met.^{9,11} In the grassland the white-tailed deer is mainly confined to brushy and wooded river flats because of cover requirements, and in the boreal mixedwood severe winters with deep snow restrict deer to its southern regions.

The distribution and relative quality of white-tailed deer habitat in Alberta are depicted in Figure 2.

2.2 Cover

2.2.1 Vegetation

Trees, shrubs and tall herbaceous vegetation are all relied upon heavily for shelter and security.¹³

In the agricultural portions of the province, winter cover is best provided by native aspen (*Populus tremuloides*) parkland and groveland in parcels of 0.4 km² or larger and along major wooded river valleys. Smaller tracts of cover may be sufficient if the climate is mild, as in the prairie region.¹³

Dense stands of snowberry (*Symphoricarpos albus*), rose (*Rosa* sp.), silverberry (*Elaeagnus commutata*), choke cherry (*Prunus virginiana*), willow (*Salix* sp.) and tall herbaceous vegetation, either alone, complexed, or in association with treed areas, provide excellent cover.¹³

In the boreal mixed-wood on spring-fall and summer range, deciduous, coniferous and mixed-wood trees and shrubs ≥ 1.5 m tall with $> 75\%$ crown closure are best for thermal cover. Stands < 1.5 m can be used if the stand is at least pole size and has $\geq 60\%$ crown closure.

Thermal cover requirements are the same on winter range except that coniferous stands, developed at least to the pole-sapling stage, are used more heavily.¹⁵

Well developed understory vegetation is preferred in winter but not on summer range.¹⁶

The optimum size of stands for thermal cover is 0.8-2.0 ha, for hiding cover is 2.6-10.5 ha and for fawning cover is 0.4-2.0 ha.¹⁶

Optimum fawning habitat in forested regions includes low shrubs or small trees from 0.6-1.8 m under a tree overstory of 50 per cent crown cover closure.¹⁴ Aspen groves and willow-aspen rings are ideal in the parkland.¹³

2.2.2 Land Forms and Topography

Though topography is significant as cover, it cannot substitute for vegetation, only increase its value.

Wooded river flats, coulees and floodplains provide protection and are especially important in the grassland regions.¹¹

Valley slopes and eroded and hummocky terrain provide important thermal cover in winter.⁶

Rugged and steeply sloped terrain provides shelter and visual obscurity.¹³

Woody draws, drainages and basins with slope < 15 per cent are used for fawning cover.¹⁶

North-facing slopes are preferred in summer and fall; south-facing slopes in spring and winter.¹⁶

The deer prefer lower altitudes in winter; higher slopes and ridge tops in summer.¹¹

2.2.3 Climate

Sites protected by topography or dense vegetative cover are sought out in winter.

In areas where chinooks are frequent and snow depths are shallow, smaller tracts of cover are required.

Snow depths \geq 40 cm cause movement off summer ranges.^{7,11}

If winters are severe and snow depth exceeds 50 cm, deer will aggregate and remain in dense stands of conifer or mixed-wood;¹⁵ the need for adequate thermal cover (i.e. energy conservation) is very important and availability of forage becomes a secondary consideration.

2.3 Food

2.3.1 Vegetation

Food utilization reflects the availability of vegetation, therefore it is highly variable depending on the geographical region and season.

Grassland

Grasses and agricultural crops are consumed in spring and summer, but deer will also browse at this time if shrubs are available.⁶

Browse is the predominant food in fall and winter, but deer will also paw out waste grains and shoots from agricultural fields.⁶

Preferred browse include: choke cherry, rose, snowberry, silverberry, bearberry (Arctostaphylos uva-ursi), juniper (Juniperus sp.), red-osier dogwood (Cornus stolonifera) and aspen.⁶

Parkland

Grasses and agricultural crops such as alfalfa (Medicago sativa), barley (Hordeum sp.), and wheat (Triticum sp.) are primarily consumed in fall and spring.⁶

Browse predominates in winter but waste grains and shoots are also utilized.

In summer deer feed on browse species such as rose and aspen as well as herbaceous forage.⁶

Forest

Deer are primarily browsers of deciduous shrubs and trees in summer and winter, utilizing coniferous trees only if starving.^{6,16}

The preferred seasonal browse is aspen and willow in winter; rose, snowberry, aspen and balsam poplar (Populus balsamifera) in summer.⁶

Browsing also occurs in spring and fall, but forb consumption is usually greater.⁶

2.3.2 Land Forms and Topography

Valley slopes, as well as hummocky and eroded terrain, promote vegetation diversity.⁵

South-facing slopes are important feeding areas in winter and spring since they have the least snow.

2.3.3 Aquatic Forms

Lakes, ponds and streams in close proximity to cover provide water plus succulent grasses and water plants.

The deer require available water within 180 m, especially during fawning.¹⁶

2.3.4 Climate

Deep snow restricts feeding on ground vegetation.⁹ It cuts down movement, restricting access to browse.^{11,15} Snow-free areas associated with chinooks and southerly slopes, are important to deer survival.

2.4 Space

2.4.1 Territory/Home Range

Deer usually have very small home ranges if adequate food and cover are available.¹¹

In good habitat a deer may remain in an area 500-800 ha.^{4,5}

Migration takes place from summer to winter range only if cold weather conditions and deep snow force the move.¹¹

Home ranges are reported to be in the order of 70-190 ha during the summer, and 160-480 ha during the winter.¹⁰

2.4.2 Population Density

The mean densities of white-tailed deer for various ecological zones are measured in number of animals per linear kilometre: prairie - 0.87, parkland 1.24, mixed-wood - 0.22, foothills - 0.18 and mountain - 0.04. In the prairie/parkland densities were recorded at 2 deer/km².¹

2.5 Special Considerations

2.5.1 Size, Shape and Juxtaposition of Habitat Components

Woody cover types which are diverse in height and species composition and occur in close proximity to farmland are ideal white-tailed deer habitat.^{9,11}

Most use of land occurs within 180 m of the edge between cover and feeding areas,¹⁶ therefore a good interspersed and juxtaposition of native and agricultural land is important.

Small openings in forested areas plus the presence of fresh water ponds, streams and rivers enhances interspersed of food and cover, and increases habitat quality.¹⁴

Edge or ecotonal areas associated with transition zones between treed and non-treed areas also brings suitable food and cover types into close association, and are productive of good browse species.¹⁷

A wide variety of natural vegetation cover types within an area is more likely to provide all habitat requirements.²

Small rivers, streams, meltwater channels and stringer forests provide continuous native cover for travel corridors.¹²

2.5.2 Significance of Disturbance Phenomena

Agriculture. Given adequate nutritional status, deer likely prefer natural habitats dominated by natural food.¹⁴

Though cultivated crops provide deer with an important supplement to their browsing, the loss of natural habitat to agriculture reduces the cover required by the deer.

Intensive agriculture improvement reduces the interspersions of native and agricultural land, isolating blocks of suitable native winter habitat and eliminating bluffs of summer habitat.¹⁴

The reduction of natural habitat promotes overcrowding and overutilization of existing suitable lands. This increases animal mortality and reduces productivity.¹⁴

Fire. Controlled burning provides more browse for deer by stimulating sprouting from understory plants and permitting more light to aid growth.³

Fire also increases the amount of suitable food and cover by encouraging edge, and improving nutrient recycling.³

Fire control allows forest maturation which removes suitable browse from the reach of deer.

Human Disturbance. If sufficient cover is available, deer will show a high tolerance to human activity. They have been highly successful in coping with hunting pressure and intensive land management.⁸

Logging. Logging creates earlier stages of vegetation succession which favors forage production, but cutting patterns must be small and irregular to enhance the interspersions of food, cover and the development of edge habitats. These edge habitats are usually narrower and more productive than natural edges.⁸

Deer will use open areas produced by clearcut logging, but prefer natural openings and burned areas.¹²

2.6 Limiting Factors

It is difficult to pinpoint factors that limit white-tailed deer in some areas and not in others.

Snow depth is probably the factor that limits deer movement most, and also limits distribution in the province; deer mobility is restricted by snow depths greater than 38 cm, thus in areas such as the northern boreal forest traditional winter ranges with shallow snow are critically important.¹⁵

In southern grassland regions deer are mainly restricted to large habitat blocks because of the scarcity of escape cover.

2.7 Regional Variations

In the grassland winters are mild. There is less need for thermal cover, but because of intensive human activity, protective cover is still required. The main deer concentrations occur in close proximity to sheltered, wooded locations, which often occur along river valleys.

Treed areas are scarce, so less browse is available, and deer rely heavily on grasses and agricultural crops.

The parkland region is ideal white-tailed deer habitat because the mixture of native and agricultural lands allows cover in close proximity with a good food supply.¹⁸ Aspen groves alone are an excellent supply of food and cover.

The boreal mixed-wood is characterized by long winters with deep snow. Deep snow conditions may cause deer to yard up under heavy cover which decreases their access to browse, and may lead to starvation.

ACKNOWLEDGMENTS

Critical reviews provided by L. Fitch (Alberta Fish and Wildlife, Lethbridge) and E. Bruns (Alberta Fish and Wildlife, Rocky Mountain House).

SELECTED REFERENCES

1. Cook, A. 1980. Stratified random deer census design and trial for W.M.U.'s P208, P210, and P220 using the double observer (front-back) quadrat procedure, January 14-22, 1980. Alberta Fish and Wildlife. 66pp.
2. Dahlberg, B. L. and R. C. Guettinger. 1956. The white-tailed deer in Wisconsin. Wis. Cons. Dept., Tech. Wild. Bull. No. 14. 282pp.
3. Dills, G. D. 1970. Effects of prescribed burning on deer browse. J. Wildl. Manage. 34(3):540-545.
4. Goulden, H. D. 1981. The white-tailed deer of Manitoba. Manitoba Dept. of Nat. Resour.
5. Goulden, R. C. and H. D. Goulden. 1969. White-tailed deer capability, Antler River - Gainsborough Creek area of south western Manitoba. Man. Dept. of Mines and Nat. Resour., Cdn. Land Inventory Proj. Rep. No. 5. 50pp.
6. Harmoning, A. K. 1976. White-tailed deer dispersion and habitat utilization in central North Dakota. North Dakota Fish and Game Dept., Proj. No. W-67-R-13, 14 and 15. Unpubl. report. 37pp.
7. Kelsall, J.P. 1969. Structural adaptations of moose and deer for snow. J. Mammal. 50: 302-310.
8. Kramer, A. 1972. A review of the ecological relationships between mule and white-tailed deer. Alberta Fish and Wildlife Div., Wildlife Section. Occasional paper #3. 54pp.
9. McCaffery, K. R. and W. A. Creed. 1969. Significance of forest openings to deer in northern Wisconsin. Wis. Dept. of Nat. Resources. Tech. Bull. 44. 104pp.
10. Rongstad, O. J. and J. R. Tester. 1969. Movements and habitat use of white-tailed deer in Minnesota. J. Wildl. Manage. 33(2): 366-379.
11. Severinghaus, C. W. and E. L. Cheatum. 1956. Life and times of the white-tailed deer. Pages 57-186, IN W. P. Taylor, ed. The Deer of North America. Stackpole, Harrisburg, P.A.
12. Sparrowe, R. D. and P. F. Springer. 1970. Seasonal activity patterns of the white-tailed deer in eastern South Dakota. J. Wildl. Manag. 34(2):420- 430.
13. Stelfox, H. A. 1979. Terrestrial wildlife habitat inventory of the Weyburn (62E) - Varden (62F) map area. Wildl. Tech. Report 79-6. Sask. Tour. and Renew. Resources.

14. Stewart, R. R. 1981. Seasonal diets of white-tailed deer in three broad ecological zones. Sask. Wildl. Tech. Report 81-4. Sask. Tour and Renew. Resources.
15. Telfer, E. S. 1970. Winter habitat selection by moose and white-tailed deer. J. Wildl. Manage. 34(3): 553-559.
16. Thomas, J. W. 1979. Wildlife Habitats in Managed Forests in the Blue Mts. of Oregon and Washington. U.S. Dept. of Agr. Forest Service. Agr. Handbook 553.
17. Wishart, W. 1979. Regional management of white-tailed deer and their habitat. Unpubl. report. Alta. Fish and Wildl. Div., Edmonton.
18. Saskatchewan, White-tailed deer in Saskatchewan. Conservation Bulletin No. 2. Department of Natural Resources, Regina.




Figure 2: General Distribution of Habitat Quality for White-tailed Deer in Alberta.

MULE DEER



3. KEY HABITAT REQUIREMENTS FOR MULE DEER

3.1 General

Although mule deer (Odocoileus hemionus) is generally considered an open country or prairie species, they are found over much of Alberta, primarily on rugged sloping terrain with native cover. Much of the best mule deer habitat occurs along river banks and breaks where there is shelter from wind, good drainage, good escape terrain and a wide variety of browse and forb species. Adequate food and cover requirements can be met by a variety of vegetation types: low shrubland, deciduous and mixed deciduous-coniferous forests.

The distribution and relative quality of mule deer habitat in Alberta is depicted in Figure 3.

3.2 Cover

3.2.1 Vegetation

In forested areas coniferous or deciduous trees and shrubs approximately 1.5 m tall with 75 per cent crown closure satisfy thermal cover requirements on summer and spring-fall range; requirements on winter range are the same but stands should be coniferous.²

In the foothill and mountain regions, pine-Douglas fir (Pinus-Pseudotsuga menziesii) cover types are important at lower altitudes. The spruce-fir (Picea-Abies) forest is mainly used at high elevation.¹⁵

In these regions the optimum size of mule deer thermal cover is 0.8-2.0 ha with a minimum width of 91 m.²

In prairie and parkland, shrubby and hardwood draws and wooded river valleys are available cover areas; shelter belts are used if they are on or near rough terrain.¹⁵

Optimum favoring habitat in forested regions consists of regenerating low shrubs or small trees between 0.6-1.8 m tall under approximately 50 per cent overstory cover²; in the grasslands mule deer show a preference to fawn in deciduous thickets.¹⁵

3.2.2 Land Forms and Topography

A strong association usually exists between mule deer distribution and rugged steep-sloping terrain.

In the prairie grassland, areas along valleys and eroded slopes are frequently used by mule deer.⁷

Deer use densely wooded valley bottoms to some extent in the fall, but prefer uncultivated terrain in fringe agricultural areas in the winter and spring.

Slopes of < 5 per cent are optimum for fawning habitat.²

3.2.3 Climate

Sites protected by topography or dense vegetative cover are sought in winter.

In foothill and mountainous regions, mule deer winter at the highest altitude that snow will allow; they will descend to lower elevations as snow depth increases¹⁶ and becomes restricting (> 30 cm).¹²

Southern exposures with low snow depths are most heavily utilized in late winter and early spring.

3.3. Food

3.3.1 Vegetation

The food habits of mule deer vary greatly with season and geographic location.

In Alberta during the fall and winter seasons, mule deer paw for dried leaves of deciduous trees, especially balsam poplar (Populus balsamifera). Cured herbaceous vegetation browse becomes important¹⁵ during winters of deep snow.

Mule deer prefer particular browse species: sagebrush (Artemisia cana), bearberry (Arctostaphylos uva-ursi), buckbrush (Symphoricarpos occidentalis), juniper (Juniperus sp.), poplar, choke cherry (Prunus virginiana), saskatoon (Amelanchier alnifolia) and red osier dogwood (Cornus stolonifera);¹⁵ in the foothills willow (Salix sp.) and Douglas fir are important.

Grass is eaten in all seasons but is especially important in spring.¹⁵ Grass composition likely increases in association with agriculture, i.e. Alberta prairie mule deer.

In summer new leaves of shrubs and forbs are used extensively; berries are also preferred.¹⁵

In autumn, shrubby vegetation is most important though forbs may make up 25 per cent of the diet.⁵

Mule deer will feed on agricultural crops, especially alfalfa (Medicago sativa), winter wheat (Triticum sp.), fall rye (Secale cereale), oats (Avena sativa) and on garden produce.

3.3.2 Land Forms and Topography

Southern exposures with the least snow allow best access to grasses and forbs in early spring.¹⁰

3.3.3 Aquatic Forms

Optimum mule deer habitat has open water within 0.8 km of any point; heaviest use of an area occurs within 300 m of water.²

Free water is less important in areas of succulent forage.¹⁴

Riparian and wetland zones are heavily used because of their palatable vegetation.²

3.3.4 Climate

Deer can remain in areas with up to 45 cm of snow³, but the amount of forage available is drastically reduced, particularly herbaceous species.

Deep snow cover during the winter season restricts foraging so that 75 per cent or more of the diet is browse.⁵

Snow of high density, almost regardless of depth, eliminates the availability of herbaceous forage.

3.3.5 Trace Elements

Natural mineral licks are used by deer, particularly in summer months.¹⁵

3.4 Space

3.4.1 Territory/Home Range

Mule deer in the mountains may make an annual migration from high elevation summer range to lower winter range.¹⁶

In Montana, mule deer travelled on average 9.4 km to summer range and 11.7 km back to winter range.

They will usually remain on the same range all year if snow depths are not restrictive, and high quality forage is present.

In Montana, summer and winter home ranges were approximately 3 km² and 13 km², respectively.¹¹

3.4.2 Population Densities

The following mean densities of mule deer for 1965 to 1974 are shown as number of animals per linear kilometre: prairie - 1.25, mixed-wood - 0.11, parkland - 0.43, foothills - 0.36, and montane - 0.11.¹

3.5 Special Considerations

3.5.1 Size, Shape and Juxtaposition of Habitat Components

Interspersion of food and cover is essential to mule deer habitat. Unbroken areas of climax boreal forest and open grassland devoid of shrubs and trees limit use by deer.¹⁵

In forested areas mule deer show a heavy use of natural openings with adjacent escape cover; for optimum use, openings should have no point farther than 180 m from the edge of cover.²

Open areas will be used readily if they are screened from roads by vegetation and topography.⁴

The optimum size of openings is 10.5 ha.²

For adequate cover mule deer usually make use of the added effect of rugged, steep-sloping topography and native shrub and tree vegetation.

Travel lanes with protective cover between timbered drainages are important in forested regions; the optimum width is 130-180 m.²

3.5.2 Significance of Disturbance Phenomena

Logging. Clearcutting of lodgepole pine (*Pinus contorta*) and spruce-fir forests in narrow strips increases forage production by opening the canopy and allowing growth of understory vegetation; this also creates prime fawning habitat.¹⁶

Road building that accompanies timber harvest in many areas is detrimental because it destroys habitat and opens new access to the public.¹⁶

The open canopy creates greater snow accumulations which may restrict deer mobility (if > 30 cm).¹⁵

Fire. Decades of fire suppression has altered ecosystems and created vast stands of extremely dense timber unfavorable for mule deer.¹⁴

Small burns improve deer habitat by creating temporary openings.

Depending on the fire intensity and the season when the fire occurs, burns can increase the diversity and abundance of understory plants.⁸

Controlled burns have increased forage on mule deer ranges in Northeastern Washington.⁴

Agriculture. The expansion of cultivated land decreases mule deer range because of the removal of the native vegetation required for food and cover.

Programs, such as Buck for Wildlife, try to retain critical prairie habitat for deer; this is especially important in southeastern Alberta and along river breaks in the south.

Overgrazing by livestock alters native plant composition by decreasing the abundance of almost all woody species important to mule deer.⁶

Human Disturbance. Mule deer prefer areas of reduced human activity and are less adaptable than white-tailed deer, particularly in regards to hunting pressure.

Roads and highways often disrupt normal migratory and daily travel routes.¹⁵

Forage along roadsides and salt on highways often attracts deer. The results are increased mortality.¹⁵

3.6 Limiting Factors

Winter weather conditions, especially snow depth, are a major limiting factor for mule deer in northern regions.^{5,9}

Studies in Alberta indicate that 30 cm of snow is restrictive to deer movement, while deer avoid areas with snow depths greater than 45 cm.^{12,13}

Because of deep snow conditions in some regions such as the foothills and northern regions, traditional wintering areas which have little snow accumulation are critically important.^{12,13,14}

In areas of intense human activity, escape cover, such as found in river valleys, is very important.

Predation, except by man, has not restricted the geographic distribution of mule deer and has never been the principal cause of a population decline.¹⁴

3.7 Regional Variations

Mule deer are ideally suited to the semi-open natural vegetation cover of prairie and parkland areas. Intensive agricultural development and human disturbance, however, have largely restricted their distribution to localized areas of rough and broken topography associated with river valleys, eroded escarpments and sand hill complexes.

Mule deer are abundant and widely distributed throughout the foothills and mountains of the Eastern Slopes where they prefer steep topography and partially treed lands associated with valley slopes and bottom-lands, south-facing slopes, tree-line subalpine and montane environments. Conifer stands are utilized for thermal and hiding cover.

Mule deer are more thinly distributed through the southern portions of the boreal mixed-wood forest, and again prefer the more rugged valley complexes and

south-facing slopelands where browse conditions are good and snow depths are not restrictive during winter.

ACKNOWLEDGMENTS


Critical reviews provided by E. Bruns (Alberta Fish and Wildlife, Rocky Mountain House), M. Watson (Alberta Fish and Wildlife, Edmonton) and L. Russell (Alberta Fish and Wildlife, Lethbridge).

SELECTED REFERENCES

1. Alberta Fish and Wildlife Division. Undated. Critical Areas Mapping File. Wildlife Services Branch, Edmonton.
2. Black, H., R. J. Scherzinger and J. W. Thomas. 1976. Relationships of Rocky Mountain elk and Rocky Mountain mule deer habitat to timber management in the Blue Mountains of Oregon and Washington. Elk-Logging-Roads Symposium. Univ. of Idaho.
3. Gilbert, P.F., O.C. Wallmo, and R.B. Gill. 1970. Effect of snow depth on Colorado deer. J. Wildl. Manage. 34:15-23.
4. Guenther, S. E. 1964. Annual biological summary, District No. 1 Unpubl. Report. IN file. Wash. State Dept. Game, Olympia. June. 2pp.
5. Hill, R. R. 1956. Forage, food habits and range management of the mule deer. pp. 394-448 IN W. P. Taylor, ed. The Deer of North America. The Stackpole Co., Harrisburg. 668pp.
6. Julander, O. 1955. Deer and cattle range relations in Utah. Forest Science. Vol. 1, No. 2.
7. Kramer, A. 1972. A review of the ecological relationships between mule and white-tailed deer. Alberta Fish and Wildlife Division. Wildlife Section. Occasional paper #3. 54pp.
8. Lewis, C. E. and T. J. Harshbarger. 1976. Shrub and herbaceous vegetation after 20 years of prescribed burning in the South Carolina coastal plain. J. Range Manage. 29(1):13-18.
9. Loveless, C. M. 1967. Ecological characteristics of a mule deer winter range. Tech. Publ. No. 20. Colo. Game, Fish and Parks Dept. 124pp.
10. Martinka, C. J. 1967. Habitat relationships of white-tailed deer and mule deer in Northern Montana. J. Wildl. Manage. 32(3):558-565.
11. Nyberg, H.E. 1980. Population ecology and habitat relationships of mule deer in mountain-foothill habitats of southwestern Montana. Montana Wildl. Div. Proj. No. W-120-R-9 & 10, Final Job Report. 106pp.
12. Telfer, E. S. 1970. Winter habitat selection by moose and white-tailed deer. J. Wildl. Manage. 34(3):553-559.
13. Telfer, E. S. and J. P. Kelsall. 1971. Morphological parameters for mammal locomotion in snow. Paper presented at 51st Ann. Meeting Am. Soc. Mamm., Vancouver, B.C. 6pp. and append. typescript.
14. Telfer, E. S. 1978. Cervid distribution, browse and snow cover in Alberta. J. Wildl. Manage. 42(2):352-361.

15. Wallmo, O. C. 1978. Mule and black-tailed deer in North America. Wildl. Manage. Instit. University of Nebraska Press.
16. Zeigler, D. L. 1978. The Okanagan mule deer. Wash. Dept. of Game. Biol. Bull. No. 15.

MOOSE



4. KEY HABITAT REQUIREMENTS FOR MOOSE

4.1 General

Moose (Alces alces) occupy a range of habitat types within forested areas. They are primarily browsers and favour immature forest shrubland for food while dense woody forest areas are utilized for cover. Preferred habitat is the early stages of mixed-wood forests with a great diversity of deciduous woody vegetation. Fire-induced forest regeneration usually provides ideal forage. The presence of aquatic environments creates edge-type vegetation communities which are diverse in composition and provide important food items during summer.

The distribution and relative quality of moose habitat in Alberta is depicted in Figure 4.

4.2 Cover

4.2.1 Vegetation

Dense, mature, coniferous forest is utilized as shelter from severe winter conditions such as low temperatures and wind, as well as an escape from harassment by insects during summer.^{6,7,12}

Mixed-wood and deciduous forests can be utilized throughout the year though moose tend to abandon deciduous forests during winters of deep snow.

During summer moose select tree muskegs and immature aspen stands > 10 m in height for cover.¹⁴

Both deciduous and coniferous cover should have closed canopies for optimum hiding cover and winter shelter.^{14,17}

The cover should be at least 200 m wide to meet hiding and shelter requirements.¹⁵

4.2.2 Land Forms and Topography

In northeastern Alberta moose occupy upland cover in summer.⁵

Lowlands are heavily utilized in late fall and early winter;^{8,9} as snow depths increase, moose shift to densely forested uplands, preferably coniferous.^{9,14}

4.3 Food

4.3.1 Vegetation

Moose are primarily a browsing species which feed extensively on the foliage and twigs of deciduous shrubs and trees. Forbs and aquatic vegetation may become an important food when available during spring and summer.¹¹

Saskatoon (Amelanchier alnifolia), willow (Salix sp.), aspen (Populus tremuloides), red-osier dogwood (Cornus stolonifera), paper birch (Betula papyrifera), balsam poplar (Populus balsamifera), pin cherry (Prunus pensylvanica) and choke cherry (Prunus virginiana) are the most important browse species for moose in Alberta.^{2,3,10,13} Rose (Rosa sp.), high-bush cranberry (Viburnum trilobum), clematis (Clematis sp.), beaked hazelnut (Corylus cornuta) and honeysuckle (Lonicera sp.) are of lesser importance.^{3,10,13}

The composition of rumen samples usually reflects the relative availability and abundance of the browse species consumed. The exceptions are saskatoon which is a highly palatable and preferred species, and red-osier dogwood which is selected preferentially in the fall when the stems become red and are higher in sugar content.^{10,11}

During the spring moose move into open areas where early green-up allows feeding on forbs and other newly emerged succulent vegetation.^{5,11}

During summer, moose in Alberta feed extensively on deciduous leaves. Aquatic vegetation has limited importance.

Grasses and other grass-like herbaceous material may be utilized but never make up a significant part of the diet.^{11,13}

Bark may be stripped off larger trees, especially in late winter and early spring when food is in short supply.¹³

Recent burns and early successional stages of the forest are most productive for moose. As forest succession advances, the overstory trees grow beyond the reach of moose and shade out the understory browse so neither are available.⁷

4.3.2 Aquatic Forms

Riparian habitats are an important source of palatable forage.

Moose are attracted to weedy lakes, marshes and sluggish streams where they can feed on aquatic vegetation.

4.3.3 Trace Elements

Aspen bark supplies moose with an important source of sodium.

4.4 Space

4.4.1 Territory/Home Range

Moose generally make seasonal movements between winter range (December to March) and summer range (April to November), coinciding with the spring thaw and freeze-up.⁵

The seasonal home range is approximately 20 km², ranging from 10-28 km², depending upon the age and sex of the animal.⁸

4.4.2 Population Density

The mean densities for various ecological zones are measured in number of animals per square kilometre: parkland - 1.25, mixed-wood - 1.10 and foothills - 1.31.¹

4.5 Special Considerations

4.5.1 Size, Shape and Juxtaposition of Habitat Components

Ideal moose habitat requires an interspersed between deciduous shrubland for food and heavy coniferous stands for cover.

A high density and interspersed of aquatic and wetland habitats (i.e. lakes, rivers, deltas and organic terrain) provides increased habitat heterogeneity and foraging opportunities.^{3,15}

Small clearcuts and burns scattered among more mature forest also create an ideal interspersed of food and cover.

These burn/clearcut areas should be no larger than 16-80 ha to ensure close proximity to heavy escape cover.¹⁷

For optimum use by moose, long strips of cover should be no wider than 200 m.¹⁷

4.5.2 Significance of Disturbance Phenomena

Fire. Wildfire is the most important natural factor that has influenced moose distribution and habitat selection for at least several hundred years. Because of recent fire control, logging is becoming an increasingly important factor in creating moose habitat.⁷

Burn areas usually provide the most suitable moose browse after 10-15 years, the length of time varying with the time of year of the burn and its intensity.¹⁷

Logging. Clearcut areas peak in production of suitable browse for moose after 15-20 years of regeneration. This may vary depending on the type of logging method used and the type of stand removed.¹⁶

Three to 10 years after logging/burning, the developing vegetation will contain a much greater variety and quality of available food.⁷

Agriculture. Agriculture, of course, decreases moose habitat by removing natural food supplies and cover.⁴

Agriculture creates increased access to moose habitat resulting in greater human disturbance.

Human Disturbance. Depending on specific experience with the disturbance, moose may avoid agricultural clearings, roads and dwellings¹⁴, or be attracted to the edge habitats created by them.

Hunting increases the need for large tracts of protective cover.

4.6 Limiting Factors

In Alberta, moose are not generally restricted by snow, though moose will tend to avoid areas with > 65-75 cm of snow.¹⁸

Distribution of moose is decided more by food requirements than tradition or shelter factors.

Moose are limited in numbers in the grassland and aspen parkland agricultural areas where tree and shrub vegetation is either non-existent or too fragmented and localized.

Moose are generally restricted to areas with large tracts of cover because of their susceptibility to human disturbance pressures.¹⁵

4.7 Regional Variations

In the Peace-Athabasca delta, willow and other deciduous types are heavily used for food and cover. Despite abundant marsh vegetation, aquatic plants are not an important food source. Saskatoon is uncommon, but is heavily browsed where it occurs.²

In the northeast region, conifers and aspen are most heavily used for cover; saskatoon and willow are the preferred browse.¹⁰

In central Alberta, deciduous or mixed-wood forests may be used for cover. Conifers are used during harsh winters. The predominant browse species are saskatoon and willow.¹⁰

In foothill and mountain regions, various deciduous shrubs are used for food though willow is usually the preferred species. In the Porcupine Hills, moose may winter in the open or under coniferous cover.¹⁸

ACKNOWLEDGMENTS

G. Lynch (Alberta Fish and Wildlife, Edmonton) reviewed earlier drafts and provided useful comments.

SELECTED REFERENCES

1. Alberta Fish and Wildlife Division. Undated. Moose Density Map for Alberta. Wildlife Services Branch, Edmonton.
2. Allison, L. 1972. The status of moose on the Peace-Athabasca Delta. Canadian Wildl. Serv. Rep. 35pp. Unpubl.
3. Barrett, M. W. 1972. A review of the diet, condition, diseases, and parasites of the Cypress Hills moose. Alberta Lands and Forests. Typewritten. 22pp.
4. Berg, W. E. and R. L. Phillips. 1974. Habitat use by moose in northwestern Minnesota with reference to other heavily willowed areas. Naturaliste Can. 101:101-116
5. Hauge, T.M. and L. B. Keith. 1981. Dynamics of moose populations in northeastern Alberta. J. Wildl. Manage. 45(3):573-597.
6. Knowlton, F. F. 1960. Food habits, movements and populations of moose in the Gravelly Mountains, Montana, J. Wildl. Manage. 24(4):162-170.
7. Krefting, L. W. 1974. Moose distribution and habitat selection in northcentral North America. Naturaliste Can. 101:81-100.
8. Lynch, G. M. 1976. Best timing of moose surveys in Alberta. Proc. N. Am. Moose Conf. 11:154-180.
9. Mytton, W. R. and L. B. Keith. 1981. Dynamics of moose populations near Rochester, Alberta. 1975-78. Can. Field Nat. 95:39-49.
10. Nowlin, R. A. 1978. Habitat selection and food habits of moose in northeastern Alberta. Colorado Coop. Wildl. Research Unit, Colorado State Univ., Fort Collins.
11. Peek, J. M. 1974. A review of moose food habit studies in North America. Naturaliste Can. 101(1):195-215.
12. _____, D. L. Ulrich and R. J. Mackie. 1976. Moose habitat selection and relationships to forest management in northeastern Minnesota. Wildl. Monogr. 48, 65pp.
13. Peterson, R. L. 1955. North American Moose. Univ. of Toronto Press. 280pp.
14. Rolley, R. E. and L. B. Keith. 1980. Moose population dynamics and winter habitat use at Rochester, Alberta. 1965-79. Can. Field Nat. 94(1):9-18.
15. Soper, J. D. 1964. The mammals of Alberta. Queen's Printer, Edmonton. 402pp.

16. Stelfox, J. G. 1983. Logging - Wildlife interactions. Canadian Wildlife Service Report. 26pp.
17. Stelfox, J. G. Personal communication.
18. Telfer, E. S. Personal communication.





Figure 4: General Distribution of Habitat Quality for Moose in Alberta.

ELK



5. KEY HABITAT REQUIREMENTS FOR ELK

5.1 General

Elk (Cervus elaphus) is a migratory species whose seasonal movements depend primarily upon vegetation availability in relation to snow depth. Forest edge and riparian vegetation associated with post-fire succession or logging provides excellent habitat for elk in both food and cover requirements. Valley bottoms are especially important for elk as travelways because of the elks' dependence upon the diverse vegetation species found in these unique areas. As well, proximity to sufficient hiding and thermal cover is available.

The distribution and relative quality of elk habitat in Alberta is depicted in Figure 5.

5.2 Cover

5.2.1 Vegetation

Closely stocked stands of coniferous forest, 12 m or greater with high stem densities and an average canopy closure exceeding 70 per cent, are used in winters characterized by very deep snow cover.^{1,10}

Mixed-wood and deciduous forest, with trees less than 12 m in height and a canopy closure less than 70 per cent are used in spring, summer and mild winters.^{1,10}

On spring-fall and summer ranges, the optimum size of stands for thermal cover is 12-24 ha, and for hiding cover is 2.6-10.5 ha.¹

Well-developed shrub strata are preferred in winter but not in summer.¹⁰

Coniferous, mixed-wood and deciduous forests, with a well-developed understory capable of hiding 90 per cent of a standing elk from the view of a hunter at a distance of 61 m, provide effective hiding cover.¹

Dense timber or heavy brush thickets are preferred sites for calving¹⁰ though grassland-shrub. Shrub (> 1 m) communities are also used.

5.2.2 Land Forms and Topography

Elk prefer the upper third of valley slopes in winter; high alpine and subalpine terrain, especially ridge tops, are utilized in summer.

North-facing slopes are preferred in summer and fall; south-facing slopes in spring and winter.¹⁰

Elk prefer 15-30 per cent slopes, diminishing use at > 40-50 per cent and have little use at 90 per cent.^{3,8} They prefer < 15 per cent slopes and dips in steeper slopes during calving.¹

Small benches, basins, draws and stream and valley bottoms are used for calving sites.

Irregular and broken topography are important as visual barriers, particularly in non-forested areas.¹

5.2.3 Climate

Warm, southern exposures and relatively low snow pack areas (< 0.5 m) are sought in winter and early spring.

Sites which are protected by topography or dense vegetation are often sought out during winter or early spring because these areas provide a refuge from strong winds, crusting or drifting snow.

Wind swept sites are preferred in summer to reduce harassment by insects.

5.3 Food

5.3.1 Vegetation

Grassland, parkland and shrubland cover types are most productive.

Grasses are preferred throughout the year and are consumed in spring, summer and fall, as well as winter if available.

Poa sp. (bluegrass), Bromus sp. (brome), Agropyron sp. (wheatgrass), Melica sp. (melic grass), Festuca sp. (fescue) and Phleum sp. (timothy)¹¹ are preferred grasses.

Sedges (Carex sp.) are used extensively, particularly in the summer when they are green, but also in the fall and winter when the plants are brown and dried.¹¹

Forbs are used in early and mid summer and fall.⁷

Deciduous shrubs and saplings are browsed mainly in late summer, fall and winter.⁹ Saskatoon (Amelanchier alnifolia), water birch (Betula occidentalis) and aspen (Populus tremuloides) are all utilized proportionally greater than their availability, indicating preference. Other browse species consumed are willow (Salix sp.), rose (Rosa sp.), red osier dogwood (Cornus stolonifera), dwarf birch (Betula glandulosa) and low-bush cranberry (Viburnum edule).¹¹

In agricultural areas cultivated crops may provide significant quantities of forage, particularly during the fall and winter.⁶

Open forage areas should be no wider than 366 m to allow close access to cover.¹

5.3.2 Land Forms and Topography

Valley bottom floodplains and drainages with fertile soils have high food productivity and diversity.

North-facing slopes in summer and fall provide high quality forage; south-facing slopes in spring and winter are first to be snow-free and give access to forage.¹⁰

5.3.3 Aquatic Forms

Riparian and wetland zones are very productive of palatable forage.¹

Elk prefer habitats within 0.8 km of permanent water,¹ primarily lower perennial rivers and streams.

5.3.4 Climate

Snow-free areas associated with southerly aspects and periodic chinook weather provide the greatest access to forage in winter and spring.⁴

Snow depth limits forage availability in winter, and at depths > 61 cm, browsing will replace grazing.¹⁰

5.3.5 Trace Elements

Elk are attracted to salt licks but they can do well in their absence on suitable soil types.¹⁰

5.4 Space

5.4.1 Territory/Home Range

Migratory elk herds generally are found in mountainous regions where they may move vertically in response to seasonal changes.¹⁰

In the same region some herds may migrate while others do not.¹⁰

In Wyoming, the distance travelled between summer and winter range varies from 32-97 km.⁸

5.4.2 Seasonal Population Densities

Densities of elk are < 28.5/km² on alpine ranges in summer, and for brief periods may be > 180/km² on winter ranges at Ya-Ha-Tinda Ranch along the upper Red Deer River.

5.5 Special Considerations

5.5.1 Size, Shape and Juxtaposition of Habitat Components

Valley and riparian habitats are important as travel corridors between high elevation summer range and low elevation winter range;¹⁰ stringer forest stands also provide protected travel lanes.¹

The most productive foraging areas are in the open or under fairly open canopies since herbaceous understory forage varies inversely with overstory.¹⁰

A good interspersion and juxtaposition of food and cover components is important and is provided by irregular topography and parkland or forest/meadow vegetation cover.¹

Winter ranges are characterized by a high percentage of grassland; the largest wintering area in Alberta, near Bob's Creek, consists of 70 per cent grassland and 30 per cent forest.

Edge or ecotonal areas are high quality elk habitat because they provide great abundance and diversity of forage with close proximity to cover.¹⁰

Most use by elk occurs within 183 m of the edge between cover and forage areas.¹

Ecotonal areas are preferred sites for calving grounds.¹⁰

5.5.2 Significance of Disturbance Phenomena

Fire. Fire encourages the development of edge habitat and recently burned areas are often very productive of preferred forage species on otherwise forested lands.¹⁰

Fire control permits an increased survival of seedling trees producing overstocked tree stands. When the timber stands mature, the closed canopies cause a reduction of understory forage supplies.¹⁰

Control of wildfire also causes a decrease in edge.¹⁰

Logging. Logging puts much of the forest accessible at present (used by elk for summer range) in different stages of succession, providing a constant supply of ecotones and forage areas.¹⁰

Clearcut logging increases forage production and species diversity, but if the distance to available cover is too great these forage areas will not be used.⁵

Roads created for logging have provided greater accessibility to recreationalists and result in a greater need for available escape cover.¹⁰

Small amounts of logging slash can be used for hiding and calving cover, but if the depth of slash becomes greater than 2 m, the area becomes unsuitable for these purposes.¹

Agriculture. Cultivation of land for agriculture may provide elk with food in fall and winter but it generally limits elk range.¹⁰

Livestock that share range with elk may not be in direct competition for the same forage species² but they may alter the native plant composition so that preferred forage species are reduced.¹⁰

The presence of cattle discourages elk use of common water sources and is usually associated with increased human disturbance.

5.6 Limiting Factors

Snow depth is the factor most limiting to elk movement and distribution in the province. Depths of 46 cm begin to restrict elk movement while depths > 76 cm seriously curtail elk escape and foraging opportunities.¹⁰

Elk are frequently precluded from utilizing areas where there is continual disturbance due to human activity (hunting, clearcut logging, snowmobiling etc.).⁸

Blocks of continuous habitat in excess of 1 000 ha which are relatively free from human disturbance are required to maintain a year-round resident population.⁷

Elk show a decreased use of areas adjacent to roads for distances ranging from 0.4-0.8 km.¹

5.7 Regional Variations

Elk in the boreal mixed-wood rely more heavily on browse particularly during the winter. This is in contrast to the boreal foothills and mountain regions where semi-open and broken forest cover provide accessible grassland foraging areas at most times of the year.

ACKNOWLEDGMENTS

Critical reviews were provided by L. Fitch (Alberta Fish and Wildlife, Lethbridge), E. Bruns (Alberta Fish and Wildlife, Rocky Mountain House), and Harold Carr (Alberta Fish and Wildlife, Calgary).

SELECTED REFERENCES

1. Black, H., R. J. Pederson, R. J. Scherzinger and J. W. Thomas. 1979. Deer and Elk IN Wildlife Habitats in Managed Forests in the Blue Mountains of Oregon and Washington. U.S. Dept. of Agr. pp. 104-127.
2. Blood, D. A. 1969. Range relationships of elk and cattle in Riding Mountain National Park, Manitoba. Can. Wildl. Ser., Wildl. Mgmt. Bull., Ser. 1, No. 19. 62pp.
3. Bruns, E. 1983. personal communication.
4. Carr, H. D. 1972. The status of elk in Alberta in 1972. Alberta Fish and Wildlife Division. Unpublished report. 19pp.
5. Hammond, G. R. 1980. Effects of logging on elk populations in Montana. Montana Research Project No. W-120-R-11. Study No. BG-3.1. 81pp.
6. Hunt, H. M. 1978. Foods of elk in the boreal forest of Saskatchewan. Sask. Dept. Tour. and Renew. Resour., Wildl. Tech. Rep. 78-9. 27pp.
7. Hunt, H. M. 1979. The ecology and status of elk in Saskatchewan. Sask. Dept. Tour. and Renew. Resour., Wildl. Resear. Div., Saskatoon. Unpubl. mimeo. 68pp.
8. Martinka, C. J. 1969. Population ecology of summer resident elk in Jackson Hole, Wyoming. J. Wildl. Manage. 32:558-565.
9. Morgantini, L. E. 1979. Habitat selection and resource division among bighorn sheep, elk and mule deer in western Alberta. M. Sc. Thesis. University of Alberta, Edmonton.
10. Skovlin, J. M. 1982. Habitat requirements and evaluations. Pages 369-413 IN Jack Ward Thomas and Dale E. Towell, ed. Elk of North America: ecology and management. Stackpole Books. 698pp.
11. Stelfox, J. G. 1980. Nutritive value and preference ratings of common big game browse plants in Alberta. Unpubl. report. 8pp mimeo.





Figure 5: General Distribution of Habitat Quality for Elk in Alberta.

CARIBOU



6. KEY HABITAT REQUIREMENTS FOR WOODLAND/MOUNTAIN CARIBOU

6.1 General

Woodland caribou (Rangifer tarandus caribou) is a climax forest species closely associated with the coniferous boreal zones. Throughout Alberta's northern region, woodland caribou use remote forested lands which produces an abundant supply of arboreal lichens and are in close proximity to muskeg.⁸ Mountain caribou winter in forested areas but spend summer on open alpine sites. Caribou show much variation in use of habitat components from region to region.

The distribution and relative quality of caribou habitat in Alberta is depicted in Figure 6.

6.2 Cover

6.2.1 Vegetation

Woodland caribou require extensive areas of mature coniferous forest during all seasons.⁴ Mixed-wood forest may be sufficient under certain conditions, i.e. when there is no competition from other ungulates. Mountain caribou winter in forested habitat but spend summer in alpine regions.

In winter, caribou are associated with unevenly-aged climax stands, especially jackpine/lodgepole pine (Pinus banksiana/Pinus contorta) and spruce/tamarack muskegs (Picea sp./Larix laricina).⁵

In northeastern Alberta and the Peace River district, black-spruce (Picea mariana) muskegs were occupied most heavily at all times whereas aspen (Populus tremuloides) or aspen-conifer mixes were seldom used;^{7,8} in summer jackpine ridges are also used.⁸

Protective cover is important on migration routes and ideally should be 400 m wide.³

Stands of 400 m diameter are optimal for wintering areas.³

There is considerable variation in areas used as calving and breeding sites. Mountain caribou usually calve in untreed alpine areas while woodland caribou tend to utilize open muskeg sites with close proximity to forest cover.⁵

6.2.2 Land Forms and Topography

In northeastern Alberta, fairly level lowlands appeared to be selected in all months.⁷

In mountainous regions, glaciers, snowfields and windy ridges are important for relief from summer heat and insect harassment.¹¹

6.3 Food

6.3.1 Vegetation

Caribou use forested habitat year-round, feeding mainly on terrestrial and arboreal lichens in winter and on a variety of vascular plants in summer.⁴

Terrestrial lichens are the main winter forage, however arboreal lichens increase in importance in winters of deep snow and in late winter when snow crusting develops. Lichen/grass meadows and south-facing meadow slopes are used when snow depth or hardness is suitable.⁵

Arboreal lichen development is best in cool, humid coniferous forests, particularly in moderately stocked stands greater than 100 years old.⁵

In spring, caribou paw for shoots of herbs, mosses, lichens and fungi. In the summer they eat leaves of deciduous shrubs, forbs and herbs, lichens, grasses and sedges, and some aquatic plants.⁴

6.3.2 Land Forms and Topography

As winter progresses, accompanied by increasing snow depths, caribou increase use of upland areas where snow is soft and more shallow, and the availability of lichens is possibly greater.^{7,9}

Use of lowland areas, especially open muskegs, occurs as snow disappears in early spring and new growth appears.⁷

Ridges provide avenues of travel.⁶

6.3.3 Aquatic Forms

Wetlands, muskegs and lake shores provide caribou with a rich and abundant summer diet. When frozen in the winter these aquatic forms are frequently used as travel corridors.

6.3.4 Climate

Woodland caribou are adapted for a cold climate and are very tolerant to snow depth. In Alberta, caribou movements are rarely restricted by snow, but deep snow does limit the availability of terrestrial lichens forcing them to utilize arboreal lichens.

6.4 Space

6.4.1 Territory/Home Range

Caribou require large, flexible ranges. Seasonal movements are dictated by food supply, suitable calving and breeding areas, and rotation of winter range to allow the slow regeneration of lichens.⁵

Mountain caribou generally make seasonal altitudinal movements from summer range in open alpine areas to winter forested areas at lower elevations.

6.4.2 Population Densities

In general, densities are low throughout the caribou range in Alberta with the highest densities occurring on late winter ranges and breeding grounds.

In the Berland area, overall annual densities for woodland caribou are $1/83 \text{ km}^2$, and $1/48 \text{ km}^2$ for mountain caribou. Winter densities are $1/27 \text{ km}^2$ for both subspecies.⁵

In the Birch mountains of northeastern Alberta 1 animal/ 24 km^2 is an estimate of the density on late winter range; the adjusted average annual density is $1/32 \text{ km}^2$.⁷

6.5 Special Considerations

6.5.1 Significance of Disturbance Phenomena

Fire. Fire removes climax forest essential to caribou for cover and the production of lichens for food.²

In Alberta, where fire suppression has been in effect for many decades, extensive fires can be a significant threat to existing caribou populations.

Caribou avoid recently burned areas which affect their movements and fragments their range.¹⁰

Fire creates evenly-aged stands and it might take centuries for the even-aged climax forest essential to sustained caribou populations to be reestablished.

In a recent burn, snow depths are higher because fire removes windbreak vegetation and a canopy that traps snow.¹⁰

Logging. Conventional, clearcut timber harvesting removes large tracts of mature and valuable lichen-producing forests³, eliminating wintering habitat.

Clearcut logging methods also create barriers to movement, range discontinuity and increased human access.³

Agriculture. Clearing of land and drainage of swampy lowlands removes the key habitat components required for caribou.⁵

Human Disturbance. Roads through caribou habitat create disturbance and also provide greater accessibility to hunters and recreationists.³

Woodland caribou are more vulnerable to hunting than any other North American cervid.¹

It is most important to minimize disturbance along travel corridors, critical late winter range, mineral licks, calving sites and breeding locations.³

6.6 Limiting Factors

Caribou require extensive areas of mature coniferous forest for cover and lichen production.

Disease and parasites are rarely a limiting factor. However predation, primarily wolves, can have a severe impact on recruitment in a population.

Initial human disturbance in caribou range tends to restrict caribou activity, i.e. abandonment of critical range, barriers to movement, loss of access to calving grounds, etc.

Caribou possess the lowest productivity of the deer family and therefore are slow to recover from population declines.

ACKNOWLEDGMENTS

Critical review provided by J. Edmonds (Alberta Fish and Wildlife, Edmonton).

SELECTED REFERENCES

1. Bergerud, A. T. 1974. Decline of caribou in North America following settlement. J. Wildl. Manage. 38(4):757-770
2. Bloomfield, M. I. 1979. The ecology and status of mountain caribou and caribou range in central British Columbia, M. Sc. Thesis. Univ. Alberta. 318pp.
3. _____. 1979. The impact of development, settlement and associated activities on mountain caribou in central British Columbia, Canada. Proc. of the 2nd International Reindeer and Caribou Symposium. Roros, Norway.
4. Cringan, A. T. 1957. History, food habits and range requirements of continental North America. 22nd North Amer. Wildl. Conf. 485-501.
5. Edmonds, J. 1983. Personal communication.
6. Evans, H. F. 1964. Woodland caribou in northwestern United States. 29th North Amer. Wildl. Conf. 445-453.
7. Fuller, T. K. and L. B. Keith. 1981. Woodland caribou population dynamics in northeastern Alberta. J. Wildl. Manage. 45(1):197-213.
8. McFetridge, R. 1983. Personal communication.
9. Stardom, R. P. 1972. Woodland caribou and snow conditions in southeast Manitoba. Proc. of the 1st International Reindeer and Caribou Symposium. Fairbanks, Alaska.
10. _____. 1977. Winter ecology of woodland caribou, Rangifer tarandus caribou, and some aspects of the winter ecology of moose, Alces alces andersoni and white-tailed deer, Odocoileum virginianus dacotensis in southwest Manitoba. M. Sc. Thesis, Univ. Manitoba.
11. Stelfox, J. G., P. Kuchar and J. A. Bindernagel. 1978. Range ecology of mountain caribou (Rangifer tarandus caribou) in Jasper National Park, 1971-74. Canadian Wildlife Service, Edmonton, Alberta.




Figure 6: General Distribution of Habitat Quality for Caribou in Alberta.

PRONGHORN ANTELOPE



7. KEY HABITAT REQUIREMENTS FOR PRONGHORN ANTELOPE

7.1 General

Pronghorn (*Antilocapra americana*) are restricted to the grassland region, the northern limit of their North American range. Highest antelope densities are found where suitable range, forage, cover requirements, water and abrupt topography occur in juxtaposition.¹⁵ Pronghorns are closely associated with sagebrush (*Artemisia cana*) during winter months, as it is an important source of both food and cover.⁵ Pronghorn antelope are sensitive to the conversion of native grassland to cultivated crops, and highest densities are found mainly where such disturbance is minimal.

The distribution and relative quality of pronghorn habitat is depicted in Figure 7.

7.2 Cover

7.2.1 Vegetation

During all seasons of the year, approximately 85-90 per cent of pronghorns are distributed on native grassland and native grassland/sagebrush cover.⁵

During summer, 70 per cent of pronghorns use native grassland prairie with no shrub cover; in winter, > 40 per cent of pronghorns range on grassland with heavy sagebrush.⁵

Low vegetative structure (averaging 40 cm high) is preferred; vegetation > 60 cm is used less because of reduced visibility and decreased mobility.¹²

Silver sagebrush and areas of grasses or forbs > 25 cm high provide the most successful fawning sites.²

Fawn bedding sites in Alberta had mean canopy coverage of 2.3 per cent shrubs, 10 per cent forbs and 87.7 per cent grass and sedges.¹⁶

7.2.2 Land Forms and Topography

Pronghorns most often use land forms typified by low rolling, open terrain. They are seldom found on slopes exceeding 30 percent.¹⁸

Local hilly areas, pre- and post-glacial meltwater channels and eroded coulees associated with major river channels all provide protective cover for antelope.

Slopes or crests of hills are common fawning areas; some fawns bed in small depressions on slopes or flat land.³

Slough bottoms are important summer habitat use areas.

7.2.3 Climate

A warm chinook wind, which periodically reduces or eliminates snow cover during winter, is the most important climatic event enabling pronghorns to survive in Alberta.

During very occasional cold winters with deep snow in southern Alberta, pronghorns were observed to utilize silverberry (Elaeagnus commutata), quaking aspen (Populus tremuloides) and willow (Salix sp.) as cover. Upper river breaks, summerfallow or harvested fields were occupied by most animals since wind action resulted in less snow in these areas.⁴

7.3 Food

7.3.1 Vegetation

Pronghorns consume a variety of plants with the largest percentage volume recorded: grasses and sedges (Carex sp.) (13 per cent), silver sagebrush (Artemisia cana) (29 per cent), and pasture sage (Artemisia frigida) (26 per cent). These species comprised over 65 per cent of the diet during the eight-month September-April period and accounted for more than 50 per cent of the total food consumed in all months except June, July and August.¹⁵

The winter diet is frequently made up of 75 per cent browse, predominantly sagebrush. Sagebrush is extensively used because of its availability and palatability.¹⁵

There is a marked increase in use of grass and sedges in spring which coincides with new growth.¹⁵ Use of improved pasture planted to crested wheat grass (Agropyron cristatum) is high at this time of year.

The pronghorns' summer diet was characterized by a decreased use of grasses and sedges and an increased consumption of forbs and browse plants. Preferred forb species include: pasture sage, wild tomato (Solanum triflorum), doorweed (Polygonum aviculare), yellow goat's beard (Tragopogon dubius), butterfly-weed (Gaura coccinea), Colorado rubber-plant (Hymenoxys richardsonii), golden aster (Chrysopsis villosa) and wooly yarrow (Achillea millefolium), while silver sagebrush is the most important browse species.¹⁵

Cacti is utilized in late summer and fall.

In the fall, browse plants, especially silver sagebrush, are highly preferred and comprise the largest proportion of the diet; crops, especially winter wheat (Triticum sp.) and alfalfa (Medicago sativa), are also used.¹⁵

Preferred late summer and fall food items have a relatively high moisture content.

7.3.2 Aquatic Forms

Antelope require free water distributed at 1.5-8.0 km intervals during all seasons of the year.¹²

Fawns bed at a mean distance of 586 ± 30 m from water.³

Irrigation ditches, springs, streams, lakes, sloughs and snow, all meet water requirements.

7.3.3 Climate

Frequent chinooks greatly enhance winter forage availability.

Antelope have difficulty obtaining the necessary quality and quantity of forage if snow depth exceeds 25-30 cm.¹⁸

7.3.4 Trace Elements

A deficiency of trace minerals, especially selenium, can lower the survival rate of fawns and result in poor health of adults.¹⁷

7.4 Space

7.4.1 Territory/Home Range

Pronghorns often range over an area not exceeding 250-500 ha in season^{6,10,13} but will make extensive movements from summer to winter ranges in response to snow depth and food availability.

Summer ranges may extend 96 km from winter range.

When undisturbed, the daily summer range of pronghorns in Wyoming was 16-65 ha;¹⁰ in Montana pronghorns moved average daily distances of approximately 3.5 - 4.0 km.⁷

Pronghorns are most widely and evenly spaced during the spring fawning period.

Winter ranges should not exceed 37 pronghorn days per ha.¹

7.4.2 Population Density

The mean density of pronghorn antelope in the grassland region during summer is $0.7/\text{km}^2$ (1972-1982).¹¹

Mean fall densities were 1.2 antelope per linear mile.⁸ Similar numbers of animals were utilizing cultivated lands (47 per cent) and grasslands (44 per cent).

7.5 Special Considerations

7.5.1 Size, Shape and Juxtaposition of Habitat Components

Preferred winter ranges contain an average of 88 per cent native vegetation consisting primarily of grasslands with varying densities of silver sagebrush cover.

The optimum vegetative composition is 40-60 per cent grasses, 10-30 per cent forbs and 5-20 per cent browse.¹⁹

Cultivated lands may be used for food in the fall if within close proximity to native vegetative cover.

Important summer habitat is provided by slough bottoms and associated forb communities.⁹

7.5.2 Significance of Disturbance Phenomena

Agriculture. Pronghorns make some use of agricultural crops, preferably alfalfa and winter wheat during fall, but these are never a major component of their diet.

Improved pastures can provide important spring range.

Irrigation ditches may provide permanent or temporary water supplies for antelope.

Fences constructed to control livestock movements can be barriers to antelope movements.¹²

Fire. Pronghorns are well suited to sub-climax vegetation created by early spring fire; summer and fall fires may seriously retard regrowth.

Human Disturbance. In antelope habitat nearly all recreational vehicles create some harassment; disturbance is especially detrimental during fawning (May 15-June 15).¹²

Regular use of established roads produces minimal disturbance.

7.6 Limiting Factors

Pronghorns in Alberta are at the northern limit of their range in North America.

The extent and quality of winter range is the most limiting factor. Antelope are restricted by snow greater than 25-30 cm therefore areas of low snowfall and frequent chinooks are critical to their survival.¹¹

Antelope require sufficient cover for shelter in severe winter yet need open areas to allow visibility and mobility.

They are limited by large tracts of intensively cultivated land where protective cover and suitable forage is scarce and fences impede movements.

ACKNOWLEDGMENTS

Critical review of the summary was provided by H. Vriend (Alberta Fish and Wildlife, Lethbridge).

SELECTED REFERENCES

1. Barrett, M. W. 1974. Importance, utilization and quality of Artemisia cana on pronghorn winter ranges in Alberta. Proc. Bienn. Pronghorn Antelope Workshop 6:26-57.
2. Barrett, M. W. 1980. Seasonal habitat associations of pronghorns in Alberta. Proc. Bienn. Pronghorn Antelope Workshop 9:174-195.
3. _____. 1981. Environmental characteristics and functional significance of pronghorn fawn bedding sites in Alberta. J. Wildl. Manage. 45(1):120-131.
4. _____. 1982. Distribution, behavior and mortality of pronghorns during a severe winter in Alberta. J. Wildl. Manage. 46(4):991-1002.
5. _____. 1982. Ranges, habitat and mortality of pronghorns at the northern limits of their range. Ph.D. Thesis Univ. Alberta. 226pp.
6. Buechner, H. K. 1950. Life history, ecology and range use of the pronghorn antelope in Trans-Pecos Texas. Am. Midl. Nat. 43:257-354.
7. Cole, G. F. 1956. The pronghorn antelope. Its range use and food habits in central Montana with special reference to alfalfa. Montana Fish and Game Dept. and Montana State Coll. Agr. Expt. Sta. Tech. Bull. 56. 63pp.
8. Froggatt, K. 1983. Fall antelope aerial surveys in hunting zone G (pre and post antelope hunting season) October 19 and 20, November 8 and 15. Alberta Fish and Wildlife. 14pp.
9. Good, J. R. and J. A. Crawford. 1978. Factors influencing use of playas in South-Central Oregon. Proc. Bienn. Pronghorn Antelope Workshop. 8:182-205.
10. Gregg, P. A. 1955. Summer habits of Wyoming antelope. Unpubl. Ph.D. Thesis, Cornell Univ. Ithaca, New York. 163pp.
11. Gudmundson, L. 1982. Antelope aerial survey and population analysis (July 15-22, 1982). Alberta Fish and Wildlife. 119pp.
12. Kindschy, R., C. Sundstrom and J. Yoakum. 1978. Range/wildlife interrelationships -- pronghorn antelope. Proc. Bienn. Pronghorn Antelope Workshop 8:216-269.
13. Leopold, A. 1947. Game management. Charles Scribner's Sons, New York. 481pp.
14. Martinka, C.J. 1967. Mortality of northern Montana pronghorns in a severe winter. J. Wildl. Manage. 31:159-164.

15. Mitchell, G. J. 1980. The pronghorn antelope in Alberta. Univ. of Regina, Saskatchewan. Monogr. 165pp.
16. _____ and S. Smoliak. 1971. Pronghorn antelope range characteristics and food habits in Alberta. J. Wildl. Manage. 35:238-250.
17. Stoszek, M. J., W. B. Kessler and H. Willmes. 1978. Trace mineral content of antelope tissues. Proc. Bienn. Pronghorn Antelope Workshop 8:156-161.
18. Yoakum, J. D. 1978. Managing rangelands for the American pronghorn antelope. Proc. Bienn. Pronghorn Antelope Workshop 8:321-336.
19. _____. 1974. Pronghorn habitat requirements for sagebrush - grasslands. Antelope States Workshop Proc. 6:16-25.





Figure 7: General Distribution of Habitat Quality for Pronghorn Antelope in Alberta.

BIGHORN SHEEP



8. KEY HABITAT REQUIREMENTS FOR ROCKY MOUNTAIN BIGHORN SHEEP

8.1 General

Bighorn sheep (*Ovis canadensis*) inhabit the rugged mountains of the Eastern Slopes. They are generally restricted to semi-open grassy slopes adjacent to precipitous terrain with rocky slopes, ridges and cliffs. This terrain offers abundant forage, accessible escape cover and little competition from other ungulates.⁷ Lungworm-pneumonia complex, along with adverse range and climate conditions, can dramatically decrease populations.^{3,15} Sheep can be very sensitive to human disturbance, particularly hunting pressure.

The distribution of bighorn sheep habitat in Alberta is shown in Figure 8.

8.2 Cover

8.2.1 Vegetation

Bighorn sheep usually occupy areas with few trees, some low growing shrubs and either natural or burned grassland,⁷ where visibility is unobstructed and escape terrain occurs nearby.¹

Bighorn sheep avoid heavily timbered areas although they may temporarily seek shelter in treed stands in adverse weather or to escape predators.⁷

The most productive habitat is range that supports native plants.²⁰

8.2.2 Land Forms and Topography

Sheep normally occupy rugged steep slopes in the range of 1 300-2 625 m elevation.⁵

During winter, preferred slopes have south or southwest exposures and are windswept with shallow snow.^{11,14} Valleys, benches and old burns at low elevations are also preferred winter sites when adequate escape terrain is available nearby.^{10,12}

During severe winter, if snow becomes crusted or deep, sheep will move to slightly higher elevations where wind and sun have cleared the more exposed slopes and ridges;⁵ caves provide excellent shelter in adverse weather.⁵

Favorite bedding sites are usually along ridges or on high knolls from where predators can be readily detected. During the day, beds are made in the open near feeding sites while night beds are found in more rugged terrain.⁷

Preferred lambing range occurs on precipitous, secluded cliffs, near foraging areas^{5,14} which generally are on dry, southern exposures.⁵

8.3 Food

8.3.1 Vegetation

Sheep are very specialized grazers which can live on hard, abrasive, dry plants,⁷ although they prefer succulent grasses, sedges and forbs.¹⁴

They consume a wide variety of plant species and vegetation use throughout the year will depend on its availability.^{8,11}

In summer, grasses are most important; forbs are more palatable but are only consumed when readily available.¹⁸

In winter, sheep paw through snow to obtain grass but if snow is too deep or crusted, they will browse; browse species are predominantly deciduous though conifers may be eaten.¹⁸

Some browsing occurs throughout the year.¹⁴

Preferred graminoids include sedge (Carex sp.), bluegrass (Poa sp.), wheatgrass (Agropyron sp.) and fescue (Festuca sp.); preferred browse species include sagebrush (Artemisia cana), willow (Salix sp.) and bitterbrush (Purshia tridentata).^{14,18}

Food habits vary significantly with availability of forage species, season, climate and habitat sites.¹⁸

Optimum sheep range has a good mixture of grasses, forbs, and shrubs.⁶

Sheep populations may increase until winter ranges become overgrazed.¹⁴

8.3.2 Land Forms and Topography

In spring, sheep forage along south and east-facing slopes until they reach subalpine and alpine areas.⁶

The highest vegetated elevations above timberline, primarily grassland meadows and plateaus, are utilized in summer.⁸

During fall the sheep shift towards north-facing grasslands and semi-open forest where snow recedes last, and the vegetation is thus more succulent and nutritious.¹⁶

South and west-facing slopes are preferred in winter.^{14,16}

Valley-bottoms and low elevation, south-facing slopes are evidently important to sheep in late pregnancy and early lactation as they provide high-protein forage.¹⁶

8.3.3 Aquatic Forms

Snow and succulent vegetation provide a year-round source of water for bighorn sheep^{5,18} however, they often supplement this with open water.

Sheep can go for long periods without drinking.⁸

8.3.4 Climate

Climatic factors affecting the distribution of sheep, especially during winter, include snow depth, snow resistance, wind velocity and temperature.¹⁶

Sheep are not well adapted to deep and crusted snow, therefore they are forced to winter where snow is shallow (< 30 cm) and soft enough for them to forage.⁸

If strong winds sweep snow from cliffs and ridges the animals will extend their winter range to where foraging is easier.⁵

Extended drought is very detrimental to bighorn sheep range, especially if it is overgrazed as well.³

8.3.5 Trace Elements

Sheep are attracted to mineral licks particularly during spring and early summer⁸ and may traverse long distances to reach them.¹⁶

It is unknown whether salt is a necessity or a desired supplement,⁷ though it is suspected that potassium uptake from succulent forages may cause excessive sodium loss.²⁰

Artificial salt causes lip irritations and contagious ecthyma.^{9,15}

8.4 Space

8.4.1 Territory/Home Range

Generally, sheep have distinct, separate summer and winter ranges with corresponding spring and fall migrations.⁷

The migration is altitudinal with lower elevational ranges occupied in winter.

Distances of seasonal migrations vary from 8-60 km.¹³

Sheep migrate by following the best forage conditions, especially the green-up of succulent grasses and forbs in the spring.^{6,7}

Sheep are predictable in their movements and loyal to their home ranges.⁵ Sheep will deviate from an accustomed range usually only because of extreme habitat deterioration, or prolonged human disturbance.

Slope, distance to escape terrain, salt availability, elevation, depth and hardness of snow, aspect, forest cover, shrub production and biomass of grasses all affect local seasonal distributions.⁷

Minimum size for home ranges is about .8 km in diameter in midwinter, and the maximum is about 5.9 km in spring and fall.⁵

8.4.2 Population Density

On winter ranges in the Rocky Mountains densities range up to 19-23 sheep/km².²

8.5 Special Considerations

8.5.1 Size, Shape, and Juxtaposition of Habitat Components

The interspersed of grassland and alpine meadow areas with accessible escape terrain is important for suitable sheep habitat. Seasonal ranges connected by open habitat provide accessible migration routes.

Wintering ranges are often 1/10 the size of summer ranges⁴, can be a part of a summer range or located some distance from it, and are located where snow accumulation is low (windswept ridge tops, steep south and west-facing slopes).^{5,16}

8.5.2 Significance of Disturbance Phenomena

Fire is an important factor in range regeneration, a deterrent to forest encroachment on grass, and a possible aid in parasite control.⁷

Fire also eliminates elk escape cover thus reducing elk use on critical bighorn range.⁷

Selective logging does not contribute to bighorn habitat development, although clearcut logging can create additional range.

Livestock compete for forage especially on critical winter range.¹⁹

Sheep have a fear and aversion to cattle and may be forced on to less preferred habitat by their presence.⁸

Hunted sheep populations are disturbed by human activity; hiking, camping, recreational vehicles and road construction, etc., all place some degree of stress on the animals.^{8,19}

It is especially important to minimize human activity along seasonal migration routes,¹⁹ and on constricted winter ranges that often comprise 10 per cent of the yearlong range.¹⁶

8.6 Limiting Factors

Human activities and predation restrict sheep to rough, secluded terrain.

Exploration, development and recreation on or adjacent to critical sheep ranges may disturb sheep and result in range loss, or abandonment.

Predation somewhat restricts sheep movements, but does not prevent sheep numbers from exceeding range carrying capacity.¹⁴

Winter conditions, especially deep snow > 30 cm, restrict sheep mobility.¹⁷

Competition with livestock and with other wild ungulates (elk, mule deer) for range can be a major factor in their population densities.⁸

Parasites and diseases, particularly pneumonia and lungworms, also limit sheep numbers.^{3,15} There is a definite relationship between disease-parasite burdens and winter forage supplies, stocking rates, winter weather and the duration of time sheep are exposed to overgrazed ranges heavily infested with parasites.¹⁵

ACKNOWLEDGMENTS

Summary was reviewed by J. Stelfox (Can. Wildl. Ser.), W. Wishart (Alberta Fish and Wildlife, Edmonton) and T. Nette (Alberta Fish and Wildlife, Rocky Mountain House).

SELECTED REFERENCES

1. Adams, L.G., K. L. Risenhoover, J.A. Bailey. 1982. Ecological relationship of mountain goats and Rocky Mountain bighorn sheep. IN: Eds, J. A. Bailey and G.G. Schounveld. Proceedings of the Biennial Symp. of the Northern Wild Sheep and Goat Council. Fort Collins, Colo.
2. Blood, D.A. 1963. Some aspects of behavior of a bighorn herd. Can. Field Nat. 77:77-94.
3. Buechner, H.K. 1960. The bighorn sheep in the United States, its past, present, and future. Wildlife Manage. 4. 174.pp.
4. Cowan, I. McT. 1950. Some vital statistics of big game on overstocked mountain range. Trans. N. Amer. Wildl. Conf. 15:581-588.
5. Geist, V. 1971. Mountain sheep - a study of behavior and evolution. The Univ. of Chicago Press.
6. Hebert, D.M. 1973. Altitudinal migrations as a factor in the nutrition of bighorn sheep. Ph.D. Thesis. Univ. B.C., Vancouver. 148pp.
7. Lawson, B. and R. Johnston. 1982. Mountain sheep IN: Wild Mammals of North America - Biology, Management and Economics. The John Hopkins Univ. Press, Baltimore.
8. McCann, L.J. 1956. Ecology of mountain sheep. Amer. Midl. Nat. 56(2):297-324.
9. Samuel, W.M., G.A. Chalmers, J.G. Stelfox, A. Luewen, and J. J. Thomsen. 1974. Contagious ecthyma in bighorn sheep and mountain goats from Western Canada. J. Wild. Dis. 11:26-31.
10. Savage, A. and C. Savage. 1981. Wild mammals of western Canada. Western Producers Prairie Books. Saskatoon, Saskatchewan.
11. Shannon, N.J., R.J. Hudson, V.C. Brink and W.D. Kitts. 1975. Determinants of California bighorn sheep in North America. J. Wildl. Manage. 34:473- 475.
12. Soper, J.D. 1964. The mammals of Alberta. Queen's Printer, Edmonton. 402pp.
13. Spalding, D.J. and H.B. Mitchell. 1970. Abundance and distribution of California bighorn sheep in North America. J. Wildl. Manage. 34:473-475.
14. Stelfox, J.G. 1976. Range ecology of Rocky Mountain bighorn sheep in Canadian National Parks, Can. Wildl. Rep. Ser. No. 39.

15. Stelfox, J.G. 1976. Diseases and parasites of bighorn sheep in Canadian National Parks, 1966-1972. Can. Wildl. Ser. Rep. 132pp.
16. Stelfox, J.G. 1978. Seasonal distributions of Rocky Mountain bighorn sheep in Canadian National Parks, 1966-1972. Can. Wildl. Ser. Rep. 149pp.
17. Telfer, E.S. and J.P. Kelsall. 1971. Morphological parameters for mammal locomotion in snow. Paper presented at 51st Annual Meeting Am. Soc. Mamm. Vancouver, B.C. 6pp and appendix, typescript.
18. Todd, J.W. 1972. A literature review on bighorn sheep food habits. Colo. Div. Wildl. Spec. Rep. No. 27. 21pp.
19. Wishart, W. 1975. Report and recommendations of the Rocky Mountain bighorn workshop group. Proc. of the Workshop on the Manage. Biol. of North Amer. Wild Sheep. pp. 165-207.
20. Wishart, W. 1978. Bighorn sheep. IN: J.L. Schmidt and D.L. Gilbert, eds. Big game of North America - ecology and management. Stockpole Books Harrisburg, PA. 494pp.




Figure 8: General Distribution of Habitat Quality for Bighorn Sheep in Alberta.

MOUNTAIN GOAT



9. KEY HABITAT REQUIREMENTS FOR MOUNTAIN GOATS

9.1 General

Mountain goat (Oreamnos americanus) ranges are very localized within alpine tundra and subalpine areas of the Rocky Mountains. These areas are associated with low temperatures and heavy snowfall, thus lower elevation winter range with accessible food is critical to their survival. Goats prefer steep cliffs and slopes which are inaccessible to humans and other predators.⁷

The distribution of mountain goat habitat in Alberta is depicted in Figure 9.

9.2 Cover

9.2.1 Vegetation

In Colorado, alpine tundra including meadows and marshes, sparse conifers, and mountain shrubs are preferred cover. Areas with willows (Salix sp.), aspen (Populus tremuloides) and dense conifers that accumulate snow are avoided.¹

Seventy-six per cent of goat herds in west-central Alberta were found on areas where shrubs and/or coniferous trees were present.⁸

The greatest use of forested areas in Montana occurred during July, August and October.¹⁰

Unobstructed visibility is not an essential habitat requirement therefore secure habitats may be heavily forested.^{6,14}

9.2.2 Land Forms and Topography

Goats are closely associated with steep, grassy alpine areas, talus slopes and precipitous, bed-rock faces which are relatively inaccessible.

They prefer steep slopes or cliffs which offer protection from predators.^{11,13}

In the Grande Cache area, 50 per cent of the total goat observations occurred on rock-talus areas with most bedding and escape activity occurring within 400 m of these sites.⁹

Nearly 50 per cent of all observations occurred on areas with 45-50° slope.^{5,9}

In summer months goats appear at elevations greater than 2 250 m; goats are found at lower levels in winter.¹⁰

In Alberta, goats below treeline selected steep, south-facing slopes which are the first to lose snow¹ (as was also found in Colorado and Alaska⁷); above treeline, windswept ridges and north-facing slopes are utilized.

9.2.3 Climate

During severe weather, caves, overhanging ledges and the sheltered side of cliffs provide refuge.¹⁵

During cold winters with deep snow, goats use lower elevations and more southerly exposures.¹³

Deep crusted snow on lower altitude ranges may cause goats to move above timberline.⁷

9.3 Food

9.3.1 Vegetation

Preferred habitats of mountain goats generally have a low abundance of forage, requiring them to use a greater diversity of species to meet intake requirements.^{4,2}

They consume a highly varied diet of alpine grasses and forbs as well as twigs and leaves of several trees and shrubs.

In Montana, grasses, sedges (Carex sp.) and rushes were most commonly eaten year round; coniferous trees were second in importance during winter and spring.¹²

During summer, goats in Alberta consume mostly grass and herbaceous vegetation.^{3,9} In one study, willow made up approximately one quarter of the diet.³

Forbs were found to be important in summer diets in Alaska.⁷

Browse, especially coniferous, is the primary winter food of goats in west-central Alberta and grasses, herbs and forbs play minor roles.⁸

Major browse species are saskatoon (Amelanchier alnifolia) and silverberry (Elaeagnus commutata); secondary species include snowberry (Symphoricarpos albus), alder (Alnus sp.) and alpine fir (Abies lasiocarpa); minor species include willow and wild gooseberry (Ribes sp.).⁸

Saskatoon is especially important because of its high palatability and availability.

Cliff terrain has forage which is often patchy and/or sparse so goats must forage alone or in small groups.²

9.3.2 Land Forms and Topography

Forage activity favors the better vegetated, relatively level, high elevation areas away from the rock-gravel cover types.⁹

During winter, sun and wind limit snow depths on south and west exposures where food availability is greatest.^{11,16}

North and east-facing slopes have the greatest supply of snow, water and succulent forage in summer.¹¹

9.3.3 Aquatic Forms

The presence of water does not affect the distribution of goats.¹³

In most ranges free water is not a significant limiting factor.¹¹

Goats obtain most of their moisture from vegetation but have been seen eating snow year round at high altitudes; in most ranges, free water is usually only used periodically.

9.3.4 Climate

Goats can forage in relatively deep snow providing it is not heavily crusted.⁴

To avoid areas of deep or crusted snow they may migrate long distances to lower elevations and/or to areas where wind or aspect limit snow accumulation.⁴

9.3.5 Trace Element

Salt licks, to which goats frequently travel are used extensively, particularly in late summer and fall when sodium is low in their diet.⁵

Licks are heavily used when available but may not be a necessity except possibly where the soil is deficient in essential elements.¹³

9.4 Space

9.4.1 Territory/Home Range

Mountain goats tend to exist in small, isolated populations.²

They are relatively sedentary, occupying small home ranges.

Summer ranges can be widely dispersed, though winter ranges are more restricted.⁷

In winter, goats may merely concentrate on areas within or adjacent to their summer range.²

The mean summer range for males in Montana was 17.6 km² while their yearly range was 21.5 km²; the mean summer range for females was 18.8 km², while their yearly range was 24.0 km²; yearlings generally have larger home ranges.¹⁰

9.5 Special Considerations

9.5.1 Size, Shape and Juxtaposition of Habitat Components

Optimum habitat for mountain goats should contain quality forage areas well interspersed with escape terrain such as rocky cliffs.

On Colorado winter ranges, goats preferred areas without persistent or melt-crusting snow where cliffs were interspersed with tundra above treeline or with shrubs or sparse coniferous habitats below treeline.¹

Usually goats winter on the highest south-facing cliffs available, where wind action is greatest, escape terrain is available, and both shrubs and conifers are present.

9.5.2 Significance of Disturbance Phenomena

Mountain goats tend to abandon areas undergoing logging activity;¹⁰ most herds in Alberta, however, are located in protected areas ie. National Parks.

Road construction and traffic both disturb mountain goats.¹⁰

The existence of roads through goat ranges increases poaching and harassment by the public.¹⁰

9.6 Limiting Factors

The distribution of mountain goats lacks a predictable pattern.

Some ranges are well populated while others are completely ignored. The factors regulating preference for certain areas are not fully understood but include suitable forage, character of the topography and slope exposure.

Because good winter range for goats is scarce, it is the habitat feature most limiting their distribution; their survival depends on the availability of low elevation range where snow depths are not > 45 cm and forage is accessible.

During winters of below normal snow depths, vegetation composition, biomass and/or palatability play a larger role in habitat use.

Since goats live in relatively inaccessible terrain, predators (other than man) do not seriously restrict the population.

Reproductive success has been correlated with spring snow depth at high elevations.²

ACKNOWLEDGMENTS

R. McFetridge, (Alberta Fish and Wildlife - Peace River) reviewed this summary.

SELECTED REFERENCES


1. Adams, L.G. and J.A. Bailey. 1980. Winter habitat selection and group size of mountain goats in Sheep Mountain Gladstone Ridge, Colorado. Proc. of the Biennial Symp. of the Northern Wild Sheep and Goat Conf.
2. Adams, L.G., K.L. Risenhoover, and J.A. Bailey. 1982. Ecological relationships of mountain goats and Rocky Mountain bighorn sheep. IN: J.A. Bailey and G.G. Schoonveld, eds. Proceedings of the Biennial Symposium of the Northern Wild Sheep and Goat Council. Fort Collins, Colorado.
3. Cowan, I. McT. 1944. Report of wildlife studies in Jasper, Banff and Yoho National Parks in 1944 and parasites, diseases and injuries of game animals in the Rocky Mountain National Parks, 1942-44 Wildl. Ser., Dept. Mines and Resour., Ottawa. 83pp.
4. Geist, V. 1971. Mountain sheep: a study in behavior and evolution. Univ of Chicago Press, Chicago, Ill. 383pp.
5. Hebert, D.M. and I. McT. Cowan. 1971. Natural salt licks as a part of the ecology of the mountain goat. Can. J. Zool. 49:605-610.
6. Hebert, D.M., and W.G. Turnbull. 1977. A description of southern interior and coastal mountain goat ecotypes in British Columbia. Pages 126-146 IN: W. Samuel and W.O. MacGregor, eds. Proc. First Int. Mountain Goat Symp. 243pp.
7. Hjeljord, O.G. 1973. Mountain goat forage and habitat preferences in Alaska. J. Wildl. Manage. 37:353-362.
8. Kerr, G.R. 1965. The ecology of mountain goats in West Central Alberta. M.Sc. Thesis. Univ. of Alberta, Edmonton.
9. McFetridge, R.J. 1977. Strategy of resource use by mountain goats in Alberta. M. Sc. Thesis. Univ of Alberta, Edmonton.
10. Rideout, C.B. 1974. A radio telemetry study of the ecology and behavior of the Rocky Mountain goat in Western Montana. Ph.D. Thesis. Univ. of Kansas, Lawrence. 145pp.
11. _____. 1978. Mountain goat. IN: J.L. Schmidt and D.L. Gilbert, eds. Big Game of North America-ecology and management. Stockpile Books, Harrisburg, PA. 494pp.
12. Saunders J.K. 1955. Food habits and range use of the Rocky Mountain goats in the Crazy Mountains, Montana. J. Wildl. Manage. 19:429-437.
13. Smith, B.L. 1972. Influence of snow conditions on winter distribution habitat use and group size of mountain goats. Proc. of the 1st Inter. Mountain Goat Symp. Kalispell, Montana.

14. Smith, K.G. 1982. Winter studies of the forest-dwelling mountain goats of Pinto Creek, Alberta. Proc. Bien. Symp. Northern Wild Sheep and Goat Counc.
15. Soper, J.D. 1970. The mammals of Jasper National Park, Alberta. Can. Wildl. Serv. Rep. Serv. No. 10.



Figure 9: General Distribution of Habitat Quality for Mountain Goat in Alberta.

GRIZZLY BEAR



10. KEY HABITAT REQUIREMENTS FOR GRIZZLY BEAR

10.1 General

The grizzly bear (*Ursus arctos*) is a highly mobile omnivore with large spatial requirements. It inhabits mainly subalpine and alpine regions, though it also occupies areas of the northern boreal forest and boreal-subalpine transitional forest in Alberta.^{10,24} Grasslands and shrublands integrated with forests, subalpine meadows and forests, and alpine communities are typical grizzly habitat. Vegetative matter is the primary food source, though animal prey are taken opportunistically. Viable populations of grizzlies require areas of low human disturbance due to human-related bear mortality and the low reproductive potential of the grizzly.⁶

The distribution and relative quality of grizzly bear habitat in Alberta is shown in Figure 10.

10.2 Cover

10.2.1 Vegetation

In the foothills/mountainous regions, dominant cover includes lodgepole pine (*Pinus contorta*), spruce (*Picea glauca* x *Picea engelmannii*), subalpine fir (*Abies lasiocarpa*), alpine larch (*Larix lyallii*), willow (*Salix* sp.), dwarf birch (*Betula glandulosa*), buffalo berry (*Shepherdia canadensis*), shrublands, grasslands, subalpine and alpine meadows; seasonally important foraging areas include shrub fields, old burns, grasslands, meadows, riparian habitat, open woodlands and old cut blocks.^{6,7,8,15,35}

In boreal and boreal mixed-wood areas, dominant cover includes lodgepole pine, white spruce, with aspen (*Populus tremuloides*) and black spruce (*Picea mariana*) as important constituents in some areas¹; seasonally important foraging areas include shrub fields, old burns, grasslands riparian areas, open woodlands and old cut blocks.³⁵

Security cover is most likely not a limiting factor in wilderness areas except in areas of resource development such as logging operations, and oil and gas exploration with associated road construction, all of which may increase hunting pressure.¹⁹

Adequate security cover to reduce visual contact by man is vegetation and/or topography which hides 90 per cent of a grizzly from view of a person 122 m away. Cover should have a diameter of at least 91 m.³⁵

Alder (*Alnus*) thickets, lodgepole blowdown and other dense vegetation are preferred as bedding sites in Montana.⁶

Den sites vary in location from alpine/subalpine talus slopes, shrub-fields, knumholtz areas to variously-timbered subalpine and lowland areas.³⁴

Surface vegetation covering den sites varies from grass and/or forbs to shrubs (willow, alder) or trees. Aerial vegetation acts as a snow catchment, while roots provide surface soil stability.^{4,26,34}

10.2.2 Land Forms and Topography

Valley bottoms and ridgetops serve as travel corridors throughout home ranges.^{31,36}

Summit ridges, with associated rock outcrops and rubble, served as mating areas in Banff.⁹

Though the exact combination of elevation, aspect, and soil type vary from one geographic area to another, most dens are located to ensure an early and long-lasting snow cover for insulation.³³

Dens were found at an average altitude of 2 280 m asl in Banff,³⁰ and between 2 100-2 380 m asl in Jasper.²⁸

Dens tend to be located on slopes allowing for ease of digging, mostly ranging from 25°-40°.18,25,34

Soil orders of dens in Banff consist of Podzolic, Brunisolic, Chernozemic and Regosolic.³⁴ These soils are well drained (prevent internal flooding of dens during late winter/early spring melts) and cohesive enough to maintain the physical stability of the den during the first winter.

No relationship was found between den sites and bedrock geology.³⁴

No geographical/ecological differences were found between the denning sites of males and females.²⁶

10.2.3 Aquatic Forms

River courses are used as corridors for movement.³¹

10.2.4 Climate

Low temperatures and snow influence the time grizzlies enter their dens.^{4,11}

Den exposure may be related to prevailing wind/radiation patterns which determine area of snow accumulation and possible soil freeze-up.^{4,25,29,22,23}

10.3 Food

10.3.1 Vegetation/Animal Matter

Grizzlies are omnivorous; they have high energy requirements but are physically, biologically and socially adapted to exploit a wide range of foods.⁵

The grizzly is primarily a herbivore and secondarily a carnivore.¹⁷

In spring, grizzlies feed on roots (mainly hedysarum, Hedysarum sp in Banff), green grasses, and bearberry (Arctostaphylos uva-ursi).^{5,8,16,17,18,26,28}

Carion from winter killed animals may also be an important food source upon den emergence.^{2,23}

During the growing season, forbs such as horsetail (Equisetum arvense), cow parsnip (Heracleum lanatum), glacier lily corms (Erythronium grandiflorum), grasses and sedges (Carex), and other green vegetation are primary components in the diet.^{5,8,16,17,18} Moist fens and streamsides produce high densities of prime summer vegetation.¹⁶

In late summer and fall, berries (huckleberry - Vaccinium membranaceum, blueberry - V. myrtilloides, buffaloberry, currants - Ribes, and saskatoons - Amelanchier alnifolia) are an important component of the diet.^{5,8,17}

Roots, grasses and forbs continue to be consumed, especially roots during fall.^{8,16,17}

Animal matter such as ants, ground squirrels (Spermophilus spp.), and young, weak or old ungulates are taken opportunistically.^{5,17,18,8,34}

When available, mammals are priority food items in Banff; scavenged or killed ungulates such as elk (Cervus elaphus) calves are common in the diet.⁸

Grizzlies locate and learn to use specific locales where plant food are abundant; the most productive sites become centres of activity within the home range.⁶

Within years, there are often seasonal concentrations of bears at feeding sites offering unlimited or abundant food supplies (e.g. berry patches).³³

10.3.2 Land Forms and Topography

Ridgetops, talus slopes, snowchutes, creek/river bottoms, fluvial and alluvial flood plains, and river/stream sides are seasonally important foraging areas.^{6,7,15,36,35}

Colluvial slopes are preferred digging areas for hedysarum roots in Banff.⁸

Anthropogenic sites such as reclaimed well sites, pipelines and road sites are also utilized seasonally.¹⁹ Most of these sites are subject to frequent or recent disturbance and therefore support early succession vegetation forms.

10.3.3 Aquatic Forms

Riparian habitats (rivers, streams, creeks, fens, bogs and seeps) provide aquatic vegetation types such as sedges and horse-tails.^{6,7,15,36,32}

10.3.4 Climate

Snow and frozen ground during the fall prehibernation period or upon spring emergence renders many food resources unavailable to bears, resulting in a negative energy balance until green-up occurs.^{22,23}

10.4 Space

10.4.1 Territory/Home Range

The grizzly bear is a mobile, wide-ranging animal that establishes seasonal home ranges.⁵

Adult females exhibit behavior suggestive of territoriality.²⁵

Home range sizes recorded in Alberta boreal forest are 126 km² for males and 179 km² for females^{20,27}, Alberta Rocky Mountain ranges are 535 km² for males and 179 km² for females.^{9,28} Adult males may range over 1 000 km².

Ranges vary greatly in area depending on the sex and age of the animal, seasonal and annual food availability, reproductive condition of females, as well as habitat type and population densities.^{3,5,25}

Seasonal ranges may be widely separated and distributed by altitude.²⁵

Males tend to have larger home ranges than females. Their ranges overlap extensively with both females and other males. Home ranges of females overlap each other, but not to the extent they overlap male ranges.^{3,12,13,25,29,22,23}

Core areas represent the most intensively used sites within a home range. In Alaska, females depend on exclusive occupancy of feeding territories to breed.²⁹

Home range sizes of females are related to reproductive status and age of young accompanying them.

In Banff, females decreased home range size when accompanied by young cubs, though they maintained a larger home range when with mobile offspring than when solitary.⁹

Females with offspring of the year generally occupy small areas in rugged terrain.

Offspring may range with the female for the first two-three years.⁶

10.4.2 Population Densities

Grizzly bear densities in the boreal forest region of Alberta are reported at 118 km²/bear²¹; in mountainous habitat (Glacier National Park, B.C.), densities are 21.2 km²/bear.³³

10.5 Special Considerations

10.5.1 Size, Shape and Juxtaposition of Habitat Components

Grizzlies require relatively large wilderness areas, preferring high habitat diversity.^{7,14}

Combinations of terrain and vegetation forming mosaics of forests, shrublands, grasslands and meadows, and riparian regions provide an interspersed array of habitats for the grizzly bear.¹⁵

Distribution of forests relative to grass-forb shrublands creates an edge-effect, enabling bears to feed on a variety of items within relatively small ranges.⁶

Seral plant communities are important feeding habitat.⁸

10.5.2 Significance of Disturbance Phenomena

Fire/Slides. Fire suppression may decrease available habitat by reducing the amount of open habitat types used for foraging.⁷ It also reduces the availability of foraging areas for ungulates, thus reducing the availability of carrion and the potential for opportunistic kills.¹⁹

Wildfires, snowslides and rockslides provide habitat diversity within coniferous forests by maintaining seral shrub and conifer communities.¹⁵

Logging. Logging of mature timber stands may enhance attractiveness as bear habitat. Earlier successional stages provide an abundance of food.⁷

Selective cuts and untreated clearcuts produce forbs and fruit-bearing shrubs, attracting grizzly use.³⁶

Clearcuts with extensive post-logging treatment reduce grizzly foods and cover.³⁶

Clearcuts should be irregularly shaped, small (<20 ha) with one or more leave patches <1 ha, minimum soil scarification (<20 per cent) and slash disposal by broadcast burning or no slash disposal.³⁶

Timber removals increase access into areas and may result in displacement of bears due to disturbance, through removal of suitable habitat, or losses due to human-bear interactions.²¹

Human Disturbance. There is competition between people and grizzly bear for space and habitat.⁶

Recreation, petrochemical exploration and industrial development modify the habitat and increase accessibility, increasing the chance of human-bear interactions and exploitation of the grizzly by legal and illegal harvest.^{1,6,20,26,30}

Artificial food sites (garbage dumps) affect dispersion and movement of the bear population, often concentrating bears for prolonged periods and may bring bears into close contact with people if placed near areas of human activity.^{6,11}

Overhunting may result in loss or suppression of grizzly bear population.²⁴

Recognition of the grizzly bear's niche can reduce conflict between humans and bears while allowing controlled development to occur.⁸

Agriculture. Conflicts between grizzlies and cattle have generally been associated with bear-caused deaths, resulting in termination or removal of bears from some grazing ranges.³²

Cattle may compete for the forage resources of the grizzly.³⁶

10.6 Limiting Factors

Wilderness, though not essential, is where grizzly bears do best.⁸

Grizzlies do adapt to the presence of humans, but cannot adapt to people's intensive use and modification of its habitat.⁶

Where grizzlies and people compete for the same habitat, people-caused bear deaths rise. The grizzly has a low reproductive rate which cannot offset heavy and persistent mortality.⁶

Preservation of especially high quality grizzly bear habitat in remote areas would reduce human-bear conflicts.⁸

Habitat quality limits the carrying capacity of an area for grizzlies.¹⁹

Failures in berry crops, an important prehibernation food, may reduce reproductive success.¹⁹

10.7 Regional Variation

Grizzly bear densities in mountainous habitat are much higher than those recorded for the boreal forest, suggesting higher habitat quality (food, diversity of habitat components, seclusion).

Prairie grasslands serve as a foraging area for grizzly bear from the Waterton Parks region.

ACKNOWLEDGMENTS

Critical reviews of this summary were provided by S. Herrero (Univ. of Calgary), J. Nagy (Alberta Environmental Centre, Vegreville), and B. Pelchat (Alberta Fish and Wildlife, Calgary).

SELECTED REFERENCES

1. Alberta Environment Centre. 1981. The impact of industrial activity on the population ecology of grizzly and black bears in the boreal forest of northwestern Alberta. 40pp.
2. Cole, G.F. 1972. Preservation and management of grizzly bears in Yellowstone National Park. in: S. Herrero, ed. Bears - their biology and management. I.U.C.N. New Series 23 (Morges):274-288.
3. Craighead, F.C., Jr. 1976. Movements and ranges of grizzly bears (Ursus arctos horribilis) as determined by radiotracking. Pro Intern. Conf. Bear Res. Manage. 3:97-109.
4. Craighead, F.C., Jr. and J.J. Craighead. 1972. Grizzly bear prehibernation and denning activities as determined by radiotracking. Wildl. Monog. No. 32. 35pp.
5. Craighead, J.J. 1977. A proposed delineation of critical grizzly bear habitat in the Yellowstone region. Bear Biology Association Monog. Ser. No. 1. 20pp.
6. Craighead, J.J., J.J. Sumner, and G.B. Scaggs. 1982. A definitive system for analysis of grizzly bear habitat and other wilderness resources. Wildlife-Wildlands Inst. Monog. No. 1. Univ. Montana Missoula. 279pp.
7. Erickson, A.W. 1976. Grizzly bear management in the Thompson Falls area and adjacent environments. U.S. Dept. Agric. Forest Ser. Missoula, Montana. 58pp.
8. Humer, D. and S. Herrero. 1983. Ecological studies of the grizzly bear in Banff National Park. Final Report. Parks Canada. 303pp.
9. Hammer, D., S. Herrero, R.T. Ogilvie and T. Toth. 1979. Ecological studies of the Banff National Park grizzly bear: Cascade/Panther Region. Parks Canada Contract WR 96-78. 86pp.
10. Herrero, S. 1972. Aspects of evolution and adaptation in American black bears (Ursus americana Pallas) and brown and grizzly bear (Ursus arctos Linne) of North America, pages 221-231 in: S. Herrero, ed. Bears - their biology and management. I.U.C.N. New Series 40, Morges.
11. Hornocker, M.G. 1962. Population characteristics and reproductive behavior of the grizzly bear in Yellowstone National Park. M.Sc. Thesis. Univ. Montana, Missoula. 94pp.
12. Knight, R.R. 1976. Yellowstone grizzly bear investigations, 1975. U.S. Dept. Int., Nat. Park Ser. Misc. Rep. 9. 46pp.
13. Knight, R.R. 1977. Yellowstone grizzly bear investigations, 1976. U.S. Dept. Int., Nat. Park Ser. Mis. Rep. 10. 75pp.

14. Martinka, C.J. 1972. Habitat relationships of grizzly bears in Glacier National Park, Montana. U.S. Dept. Int., Nat. Park Ser. Prog. Rep. 19pp.
15. Martinka, C.J. 1976. Ecological role and management of grizzly bears in Glacier National Park, Montana. in: M. Pelton, J. Lentfer, and G. Folks, eds. Bears - their biology and management. I.U.C.N. New Series 40:147-156.
16. McCrory, W. and S. Herrero. 1983. The capability and use of grizzly bear habitats in the Kananaskis and Spray Lakes Valleys and areas from Canmore to Mount Allan, 1982. BIOS Environmental Research and Planning Associated Ltd. 20pp.
17. Mealey, S.P. 1975. The natural food habitats of free-ranging grizzly bears in Yellowstone National Park, 1973-1974. M.Sc. Thesis, Montana State Univ., Bozeman. 158pp.
18. Miller, S.J., N. Barichello, and D. Tait. 1982. The grizzly bear of the Mackenzie Mountains, Northwest Territories. NWT Wildl. Ser. Com. Rep. No. 3., Yellowknife. 118pp.
19. Nagy, J.A. 1984. (per. comm.) Alberta Environmental Centre.
20. Nagy, J.A. and R.H. Russell. 1978. Ecological studies of the boreal forest grizzly bear (Ursus arctos L.) Can. Wildl. Ser. Rep. 72pp.
21. Nagy, J.A., R.H. Russell, A. Pearson, and G. Kemp. 1976. The boreal forest grizzly bear annual report for 1976. Can. Wildl. Ser. Rep. 37pp.
22. Nagy, J.A., R.H. Russell, A.M. Pearson, M.C.S. Kingsley and C.B. Larsen. 1983. A study of grizzly bears on the barren - grounds of Tuktoyaktuk Peninsula and Richards Island, Northwest Territories, 1974-78. Can. Wildl. Serv., Edmonton. 136pp.
23. Nagy, J.A., R.H. Russell, A.M. Pearson, M.C.S. Kingsley and B.C. Goski. 1982. Ecological studies of grizzly bears in the Arctic Mountains, northern Yukon Territory, 1972-1975. Can. Wildl. Serv., Edmonton. 106pp.
24. Nielson, P.L. 1975. The past and present status of the plains and boreal forest grizzly bear in Alberta. Can. Wildl. Ser. Rep. 65pp.
25. Pearson, A.M. 1975. The northern interior grizzly bear Ursus arctos L. Can. Wildl. Ser. Rep. No. 34. 86pp.
26. Pearson, A.M. 1975. Northern pipeline development and its implications to wildlife - grizzly bears. Can. Wildl. Ser. Rep. 40pp.
27. Pearson, A.M. 1976. The boreal forest grizzly bear. Annual Rep. Can. Wildl. Ser. 18pp.

28. Pearson, A.M. and J. Nolan. 1976. The ecology of the grizzly bear (Ursus arctos L.) in Jasper National Park - Report for 1975. Can. Wildl. Ser. Rep. 15pp.
29. Reynolds, H.V., J.A. Curatolo and R. Quimby. 1976. Denning ecology of grizzly bears in northeastern Alaska, pages 403-410. In: M.R. Pelton, J.W. Lentfer, and G.E. Folk, eds. Bears - their biology and management. I.U.C.N. New Ser. 40.
30. Rogers, L.L. 1974. Movement patterns and social organization of blackbears in Minnesota. Ph.D. Thesis, Univ. Minnesota.
31. Russell, R.H., J.W. Nolan, N.A. Woody, G. Anderson and A.M. Pearson. 1978. A study of the grizzly bear (Ursus arctos L.) in Jasper National Park. Can. Wildl. Ser. Prog. Rep. 1976-1977. 95pp.
32. Schallenger, A. 1977. Review of oil and gas exploitation impacts on grizzly bears. in: M.R. Pelton, J.W. Lentfer and G.E. Folk, eds. Bears - their biology and management. I.U.C.N. New Ser. 40.
33. Shaffer, M.L. 1978. Determining minimum viable population sizes: a case study of the grizzly bear (Ursus arctos L.). Ph.D. Thesis, Duke Univ.
34. Vroom, G.W., S. Herrero and R.T. Ogilvie. 1977. The ecology of winter den sites of grizzly bears in Banff National Park, Alberta. pages 321-330. in: M.R. Pelton, J.W. Lentfer and G.E. Folk, eds. Bears - their biology and management. I.U.C.N. New Series. Morges.
35. Zager, P., C. Jonkel and R. Mace. 1980. Grizzly bear habitat terminology. BGP Spec. Rpt. No. 41 - 10pp.
36. Zager, P., R. Mace, L. Lee, C. Jonkel and C. Servheen. 1980. Guidelines for occupied grizzly bear habitat in northwestern Montana. BGP Spec. Rtp. No. 51 - 17pp.




Figure 10: General Population Distribution for Grizzly Bear in Alberta.

BEAVER



11. KEY HABITAT REQUIREMENTS FOR BEAVER

11.1 General

Water is the prerequisite of beaver habitat (Castor canadensis). This semi-aquatic furbearer can utilize a wide range of aquatic environments. Ideal habitat conditions are likely to be found where an adequately deep body of water (0.9-1.5 m minimum) occurs in association with stable, fine-textured shoreline substrata needed for lodge or bank den construction and abundant supplies of upland deciduous woody vegetation for food and lodge building. The beaver occupies a niche that is transitional between aquatic and terrestrial environments and therefore prefers high-density wetland and stream areas.

The distribution and relative quality of beaver habitat in Alberta is shown in Figure 11a. Distribution of beaver harvest is displayed in Figure 11b.

11.2 Cover

11.2.1 Vegetation

Aspen (Populus tremuloides) is the species most commonly used for building material; in its absence, beaver will rely on willow (Salix sp.).¹

Alder (Alnus sp.) is another preferred construction material for lodges and dams and is frequently found in beaver food caches.⁶

Alder may also be used as a capping material for sinking preferred foods so they do not freeze into the ice.⁴

Although aspen and willow are most frequently cut by beaver, almost every woody plant is cut to some extent.¹

11.2.2 Land Forms and Topography

Beavers require bank forms of sufficiently fine texture, with stability to support excavation (burrowing) or dam and lodge construction.⁶

Soils containing clay are best suited for construction; coarse sands and gravels are unsuitable⁶; shales are subject to faulting and slipping.⁸

V-shaped valleys are more subject to dam breakage than U-shaped valleys with broader flood plains and lower gradients.⁷

The majority of beaver colonies in Alberta are built in valleys with gradients of < 2 per cent. Dams tend not to be built on gradients > 15°.⁸

11.2.3 Aquatic Forms

A wide variety of aquatic environments can be utilized by beaver providing that they are of sufficient depth to accommodate bank dens or lodges and permit free movement from the dwelling to the food cache during winter.⁶

Throughout most beaver range in Alberta, minimum water depths required are 0.9-1.5 m.⁶

They prefer natural lakes or streams with seasonally stable water levels, or a watercourse whose level can be controlled by construction of dams. The most stable water level conditions are provided by low gradient meandering streams and rivers with flat bottomed U-shaped valleys.⁶

High densities of permanent wetlands, such as occur in strongly undulating and hummocky moraine areas, provide ideal habitat.⁶

Large lakes with convoluted shorelines and small sheltered bays provide good potential habitat and also reduce wave action and provide refuge.⁶

Deltas subject to large water level fluctuations and active erosion and deposition result in very poor beaver habitat.³

In northern regions, beaver utilize organic wetlands, thermokarst lakes, floodplain oxbows and stabilized deltaic areas because of their availability.³

11.2.4 Climate

The increasing severity of climatic conditions limits areas of suitable habitat.

Snow depth is the main factor regulating ice thickness,³ and in harsh areas with cold temperatures and little snow, the greater ice depths will require deeper ponds.

Longer winter periods will require larger accessible food supplies.

11.3 Food

11.3.1 Vegetation

Deciduous trees and shrubs constitute the main source of food though herbaceous upland vegetation may also be utilized. Grain crops are sometimes used in spring and summer.^{2,6}

Both aspen poplar and willow are heavily used by beaver, but aspen is the preferred food; balsam poplar (Populus balsamifera) and birch (Betula sp.) may be used depending on their availability.⁶

Aquatic herbaceous vegetation is important in the beaver's diet during the spring and summer months.⁶

Yellow pond-lily (Nuphar variegatum) rhizomes may be used by beaver year-round if they are available, especially in more northern organic wetland environments.⁶

Under stress of necessity, almost any type of deciduous species can be used for food;² black spruce (Picea mariana) has been found in food caches though it could be there for capping.

11.3.2 Land Forms and Topography

Steep slopes adjacent to suitable wetland areas restrict upland foraging.⁶

If the relief is shallow large tracts of vegetation will be covered by water as a result of damming, whereas in steep topography a smaller area will be accessible by damming water.¹

11.3.3 Aquatic Forms

Aquatic and riparian communities assume primary importance in feeding.⁶

Meandering streams and rivers with cut-off channels and slow-flowing backwater areas provide beaver with an abundance of accessible food supplies, particularly willow, balsam poplar and water birch.

Small, intermittent streams are easily dammed if well defined, firm banks border the drainage, thus increasing the foraging area for beaver.

11.4 Space

11.4.1 Population Densities

Beaver densities in northern Alberta and southwestern portions of Northwest Territories range from 0-1 colonies per survey kilometre of streams or lake shoreline with an average density of 0.03 lodges per kilometre.^{3,5}

Densities of over one active lodge per kilometre indicates a population of maximum density in the most favorable habitat in northern regions.³

Beaver densities are much higher in parklands where habitat conditions are optimum.

11.5 Special Considerations

11.5.1 Size, Shape and Juxtaposition of Habitat Components

The range of the beaver is remarkably similar to that of aspen and willow; beaver will occupy areas where aspen is absent but very few are found outside of the range of willow, leading some to believe that willow, not aspen, is the key plant in beaver habitat.¹

Aspen stands are regarded as temporary use areas, because selective aspen cutting by beaver often hastens succession of the stand to a more advanced stage, whereas willow/alder stands may produce climax stands which can be cut

on a sustained yield basis. Once a willow/alder population is exhausted, beaver will move to a new area.¹

A high interspersion of land and water provides ideal beaver habitat; beaver will seldom forage more than 200 m from water and most cutting is done within 30 m of water.⁶

Wetland configurations which increase shoreline length relative to surface area generally provide a greater amount of potential habitat.⁶

11.5.2 Significance of Disturbance Phenomena

Agriculture. Each year there is a loss of beaver habitat due to the increase of cultivated land through draining wetlands and removing deciduous tree and shrub cover.

Fire. Small, local burns interrupt plant succession patterns and enhance beaver habitat since the pioneer woody species on burns are aspen and other deciduous trees and shrubs.

Large burns over entire watersheds may make the area very susceptible to flash flooding and erosion.⁷

The magnitude of the effect depends on the severity and extent of the fire, the rate of regeneration and the amount of subsequent erosion caused by the fire.⁷

Beaver populations are likely to be fully recovered 5-6 years after a large burn.

Flooding. Temporary, seasonal flooding adds organic nutrients to the system and enhances the growth of willow and the development of aquatic plants such as sedges (*Carex* sp.) and water lily (*Nuphar*). This may benefit the habitat initially, but flooding can be harmful if it occurs during winter freeze-up.

Human Disturbance. Drainage and decreased water levels resulting from human activities may increase the susceptibility of beaver to predation, and may cause the food cache to freeze in winter making it inaccessible.

The influx of sediment into standing water hastens the filling-in of the basin and also reduces the productivity of aquatic vegetation.

Beaver are a highly resilient species able to exist in close association with human activity.⁷

11.6 Significant Limiting Factors

Beaver are limited in distribution by the presence of aspen and willow for food and building materials and the occurrence of suitable bodies of water for protection, shelter and supplementary aquatic food sources.

Beaver are not greatly affected by human disturbance and show only limited negative response toward highways and pipeline operations except where channelization and siltation occur.

Very coarse-textured substrata limit damming capabilities and infertile substrata reduce food productivity.

Very flat topography limits the availability of suitably deep and permanent aquatic environments.

Steep-sloping topography restricts upland foraging.

Fast-flowing aquatic environments reduce vegetation productivity, restrict the ability of beaver to build dams and lodges and reduce the stability of aquatic cover.

Widely fluctuating water levels reduce dam and lodge construction capabilities and may cause freezing of the food cache during winter.

11.7 Regional Variations

In the northern boreal mixed-wood, willow is the predominant species of beaver habitat because aspen occurs only in small islands and rarely at water's edge.³

Aspen does occur along the Peace and Athabasca River valleys and is utilized more in these areas.³

In agricultural areas where natural water and food requirements may not be present, beaver will inhabit dugouts and utilize grain crops as part of their diet.

ACKNOWLEDGMENTS

A. Todd (Alberta Fish and Wildlife, Edmonton) reviewed earlier drafts and provided useful comments.

SELECTED REFERENCES

1. Hall, J.G. 1960. Willow and aspen in the ecology of beaver on Sagehen Creek, Calif. Ecology 41:484-494.
 2. Nash, J.B. 1951. An investigation of some problems of ecology of the beaver (Castor canadensis canadensis Kuhl) in northern Manitoba. Unpublished M.Sc. thesis. 53pp.
 3. Novakowski, N.S. 1965. Population dynamics of a beaver population in northern latitudes. Unpublished Doctorate theses.
 4. Slough, B.G. 1976. A land capability classification system for beaver. Unpublished M.Sc. Thesis. Simon Fraser Univ. Burnaby, B.C. 106pp.
 5. Stelfox, H.A. 1972. Aerial Beaver Food Cache Surveys in the Fort Simpson region, N.W.T. Canadian Wildlife Service, Edmonton. Unpublished manuscript. 9 p. and 3 append.
 6. Todd, A. 1978. Methodology used for Alberta Land Inventory of furbearers. Alberta Fish and Wildlife Division. 68pp.
 7. Wooly, D.E. 1972. Beaver (Castor canadensis) studies in the MacKenzie Valley IN studies of furbearers associated with proposed pipeline routes in the Yukon and Northwest Territories. Arctic Gas Biological Report Series No. 9.
 8. Yeager, L.E. and W.H. Rutherford. 1956. An ecological basis for beaver management in the Rocky Mountain region. Twenty-second North American Wildlife Conference.
- Boyd, M. 1977. Analysis of fur production records by individual fur-bearing species for registered trapping areas in Alberta, 1970-1975. Unpublished Alberta Fish and Wildlife Report. 72 pp.




Figure 11a: General Distribution of Habitat Quality for Beaver in Alberta.





Figure 11.b: Distribution of Beaver Harvest, 1970-1975; taken from Boyd, 1977.

RIVER OTTER



12. KEY HABITAT REQUIREMENTS FOR THE RIVER OTTER

12.1 General

The river otter (Lutra canadensis) is a semi-aquatic furbearer with commercial value.¹³ Being a mobile species, it requires large tracts of suitable habitat. Its main habitat requirements include access to water year-round, adequate food supply, den sites and shoreline cover.^{7,8,19,16} River otter exhibit a low reproductive potential and exist at low densities over their range with northeastern Alberta supporting the largest population.⁸ The disappearance of river otter over much of their former range in Alberta has been attributed to overtrapping.^{8,17}

The distribution and relative quality of river otter habitat in Alberta is depicted in Figure 12a. Distribution of river otter harvests is shown in Figure 12b.

12.2 Cover

12.2.1 Vegetation

Vegetation types used are highly variable.

Otters use riparian habitat on a localized basis; adequate shoreline cover (trees, shrubs, grasses, forbs) is required in the vicinity of dens and at latrine sites.^{8,17,16.}

Dense, riparian vegetation supported on undercut banks and upper banks provides excellent cover for otter, both in water and on land.²⁵

Latrine sites, though typically well vegetated, may be more open than most of the contiguous bank.²⁵ In Montana, riparian habitat dominated by shrubs and marsh grass was preferred over sparse sagebrush, hayfields, early successional regions and non-vegetated banks.²⁵

Cavities among roots of trees and in thickets of vegetation may be used as den sites or resting sites.^{12,16}

Log jams may receive frequent use as resting sites, offering cover and a site from which to hunt.¹⁶

12.2.2 Land Forms and Topography

Otters use a variety of sites for temporary dens and resting areas. Temporary sites are often used by a number of unrelated otter.¹⁴

Otters utilize natural formations (e.g. talus rock, undercut banks) and dens made by other animals (fox, muskrat, beaver).^{8,14,18, 19,16} Otters will re-excavate or fashion these den sites.^{16,17}

Beaver lodges, often in association with some form of bank burrow, are frequently used as dens.^{8,18,19}

Winter dens are often the route by which otters gain water and above-ground access.¹⁷

Beaver ponds and small bog lakes with leveed shorelines provide a bank for den sites and underwater access below the ice; the substratum generally is silt or detritus (can be dug through). Lake shorelines with gradually sloping sand, gravel or rock substratum are generally unsuitable winter habitat.¹⁷

Natal dens tend to be located well above high water levels and a considerable distance from permanent water (e.g. fox burrow).¹⁹

Distinct points/peninsulas on lakeshores, sandbars, large boulders and elevated banks may serve as latrine sites.^{17,16}

In west central Idaho, otter preferred valley to mountain habitats; topography alone appeared to have little influence on movement.¹⁶

12.2.3 Climate

Frequency of above-ground movement between water bodies declines severely with cold winter conditions; movements tend to be under the ice and otter remain long periods in dens.¹⁹

Above-ground movements between habitat patches in mid-winter are energetically costly, and generally coincide with relatively mild weather conditions.¹⁹

Cavities in the snow, or snow caves made by otter, are used as temporary shelter and as feeding stations.¹³

Spring flooding of natal dens creates potential danger to helpless young.¹⁹

12.2.4 Aquatic Forms

Otters live in association with a wide variety of water sources; habitat composition and seasonal accessibility are important factors in determining water type used.¹⁶

Flowing water and interconnected aquatic forms are more valuable as river otter habitat than isolated standing water aquatic forms.

Medium-sized eutrophic and bog lakes provide good winter habitat.¹⁹

Otter may also use rapid areas in rivers, which remain exposed during winter;^{1,16,17} streams which freeze over and cave-in due to a reduction in water flow, producing extensive air cavities, serve as potential winter habitat.¹⁷

In west central Idaho, otter preferred stream-associated habitats to lakes, ponds and reservoirs.¹⁶ Use of lakes, reservoirs and ponds, as well as

unobstructed portions of forest streams, was greatest during winter; mudflats and associated open marshes and swamps and backwater sloughs were used most often by otter during summer as rearing habitat.¹⁶

Turbid water and low flow volume may discourage otter use.²⁵

Otter movement is generally along watercourses, though they will cross country from one water source to another using the shortest route possible.^{16,19}

In streams impounded by beavers, otters may dig on the dam (rifting) to maintain an overflow of water which provides an access area in winter.¹⁹

Water acts as an escape medium for otter, and as a predator barrier for young in beaver lodges.¹⁹

Latrine sites are often located at river/stream confluences or mouths into lakes.¹⁷

12.3 Food

12.3.1 Vegetation

Vegetation is not a significant component of the diet.

12.3.2 Aquatic Forms

Beaver ponds, lakes, bogs, rivers and streams serve as feeding areas.

River otter are opportunistic feeders, consuming mainly forage fishes (e.g. minnows, sticklebacks) as well as some game species (e.g. trout, whitefish).^{4,10,11,14,21,22,24}

Local abundance of fish, lake stratification, winter ice and individual fish species' swimming ability all may affect the availability of fish to otters.^{21,24}

In Saskatchewan, the greatest densities of prey-sized fish (> 10 cm in length) were found in the back-eddies of rapids.¹

Rapids or fast water regions may provide open access to food in winter.¹

Rifting of beaver dams enables more frequent interchange and restocking of fish between successive ponds along a drainage, especially during high spring water levels.¹⁹

Crayfish are also a main component of the diet.²⁴

Other items in the diet include aquatic invertebrates, aquatic insects and beetles, freshwater clams and amphibians (mainly frogs).^{10,11,14,21,22,24}

Seasonal concentrations of invertebrate nymphs, moulting waterfowl and spawning fish represent a highly abundant food source for otter and is reflected in their diet.^{17,4}

Mammals (as muskrat; Ondatra zibethicus and beaver; Castor canadensis) may be included in the diet, though they are relatively unimportant components.^{10,14,24}

12.3.3. Climate

In winter, ice cover is often a barrier to otter trying to access water from the shore.^{8,19,16}

Deep snow may also reduce access to potential food.¹⁶

12.4 Space

12.4.1 Territory/Home Range

Home ranges generally encompass large lengths of shoreline; drainage pattern determines home range shape.¹⁶

Prey availability, habitat, weather conditions, topography, reproductive status and population densities influence home range use and length.¹⁶

Seasonal variation in movement frequencies occurs, with increased movements in spring and autumn for both sexes.^{18,19,20}

Portions of home range receive repetitive use during crucial periods such as winter.^{18,19}

In northeastern Alberta, average home range size for females was 48.2 km² and 111.4 km² for males. In late winter, home ranges were < 2.5 km² for an adult male and < 0.5 km² for adult females.¹⁹

In the Kananaskis area, an adult male's range was 178 km² from May to November, while a female and her pups travelled in a 2 km² area from May to September and ranged 7.1 km² from October to mid-November.⁹

In Idaho, a female confined her activities to a 11.2 km² section of rivers and lakes, while adult males occupied 20-30 km² of water- way.^{14,15}

There is extensive adult male/adult male and adult male/adult female overlap in home ranges. Adult female home ranges overlap less extensively.¹⁷

Young may restrain the movements of the female; they spend the first winter with the female and may move inside the female's range for the first two years.^{18,19}

Animals in suboptimal habitat may have larger home ranges.¹⁹

12.4.2 Population Densities

In west central Idaho, population densities ranged from 1 otter/2.7-5.8 (x = 3.9) km of waterway.¹⁶ Density based on sex and social status was one breeding female/20 km of waterway, one adult male/53 km, and one yearling or nonbreeding adult/14 km.¹⁶

12.5 Special Considerations

12.5.1 Size, Shape and Juxtaposition of Habitat Components

Characteristic features of studied otter habitat include⁸ waterways and marshes bordered by extensive forests, waterways containing slow moving pools with soft bottoms (as beaver ponds), high water quality with little turbidity or pollution and accessible water, year round. However, otter populations do exist in areas where all these characteristics are not present.

A sizeable area of suitable habitat must be available since otters are a very mobile species, especially during the breeding season.¹²

Regions supporting a complex of water bodies and drainages in close proximity with abundant food provide good otter habitat.

Bog lakes, with adequate food resources and den sites along the shoreline, provide a less costly means of accessing food in winter than do a series of beaver ponds.¹⁹

Size and complexity of rapids contribute to their importance.¹ High densities of rapids may allow open, year-round access. Rivers with a strong current, small islands and complex shoreline may also provide good otter habitat.¹

Locations of den sites are such that it is unnecessary for otter to travel any great distance to seek shelter while moving through the area.^{13,19}

12.5.2 Significance of Disturbance Phenomena

Trapping. Over-trapping is the most common cause of otter elimination from areas in Alberta.^{3,8}

Open trapping seasons on otters until 1961²³ may have prevented repopulation of former habitat in Alberta.⁸

Otters and beaver's use of water bodies overlap substantially,¹⁸ thus otters are often accidentally trapped in beaver sets.^{2,8,18}

Recreation. Recreational use of an area is not necessarily incompatible with river otters; stable populations can exist where continually confronted with human presence.⁸

However, if the major area of open water to which the otters require winter access coincide with areas of regular human activity otters may leave the area.⁸

Human activity which is intense and uncontrolled along stream banks may cause excessive erosion and siltation which may decrease food resources.⁸

Hydroelectric Development. Hydroelectric development which would inundate rapids and change water regimes may have a significant impact on otter.¹

Other. Potential habitat disturbances which eliminate otter by destroying its habitat and/or removal of its prey base include; siltation and infilling, pollution, channelization, damming and draining wetlands, urbanization, industrialization, agricultural developments and timber harvest.²

12.6 Limiting Factors

River otter require aquatic habitat accessible year-round, with an adequate food supply.¹⁶

The river otter occupies only portions of its former range, although areas of uninhabited suitable habitat occur. A low reproductive potential and low population densities may limit otter expansion into former ranges, as may uncontrolled trapping.⁸

12.7 Regional Variation

River otter are uncommon in Alberta.

Areas of relatively high otter production are restricted to the eastern portion of the province in the boreal forest zone.³ The distribution of higher producing areas in the northeast corner appears to be related to the Peace-Athabasca Delta complex and its adjacent river systems. High fish populations, which provide a main source of food for otter, may be an important factor in this area. Other high otter producing areas either include or are adjacent to the Department of National Defence Air Weapons Range near Cold Lake, and are currently closed to trapping.³

In northeastern Alberta, severe winters reduce the number of large open water areas to stretches of rapids on the larger rivers.⁸

Few otter reside in mountainous or prairie habitat.¹⁷

Regional differences in habitat use may be due to differences in habitat composition (water body morphology and drainage patterns).¹⁶

ACKNOWLEDGMENTS

Don Reid (University of Calgary) provided helpful comments on this summary.

SELECTED REFERENCES

1. Barber, S.R., H.A. Stelfox, and J.D. Boden. 1975. Wildlife (Saskatchewan). Fisheries and Wildlife Branch, Saskatoon. Churchill River Study Final Report 28. 272 pp.
2. Bottorff, J.A. 1976. The feasibility of river otter reintroduction in West Virginia. West Virginia Dept. of Natural Resources. Div. of Wildl. Spec. Rep. 11pp.
3. Boyd, M. 1977. Analysis of fur production records by individual fur-bearing species for registered trapping areas in Alberta, 1970-1975. Alberta Fish and Wildlife. 72pp.
4. Gilbert, F.F. and E.G. Nancekirell. 1982. Food habits of mink and otter in northeastern Alberta. Can. J. Zool. 60:1282-1288.
5. Greer, K.R. 1955. Yearly food habits of the river otter in the Thompson Lakes region, northwestern Montana as indicated by scat analysis. Am. Midl. Nat. 54:299-313.
6. Guitar, W. 1975. A literature review of the ecology and management of the river otter (Lutra canadensis). Wildlife Ecology and Management, Colorado State Univ. 10pp.
7. Hall, R.E. and K.R. Kelson. 1959. The Mammals of North America, Vol. II. The Ronald Press Cop., New York. 616pp.
8. Jalkotzy, M. 1980. River otter reintroduction feasibility study Kananaskis Country, Alberta. Faculty of Environmental Design, Univ. Calgary. 56pp.
9. Jalkotzy, M. 1982. River otter reintroduction project. Pilot Project Summary and Evaluation: June to October, 1981. Alberta Energy and Natural Resources, Fish and Wildlife Division. Kananaskis Country. 37pp.
10. Knudsen, J.F. and J.B. Hale. 1968. Food habits of otters in the Great Lakes region. J. Wildl. Manage. 32:89-93.
11. Lagler, J.F. and B.T. Ostenson. 1942. Early spring food of the otter in Michigan. J. Wildl. Manage. 6:244-254.
12. Liers, E.E. 1951. Notes on the river otter (Lutra canadensis) J. Mammal. 32:1-9.
13. Melquist, W.E. 1977. Population characteristics and dynamics of river otters (Lutra canadensis) in west-central Idaho. Idaho Cooperative Wildlife Research Unit Semi-Annual Report. Project WU-110. Univ. Idaho. pp 45-51.

14. Melquist, W.E. 1978. Population characteristics and dynamics of river otters (Lutra canadensis) in west-central Idaho. Idaho Cooperative Wildlife Research Unit Semi-Annual Report. Project WU-110. Univ. Idaho. pp 12-20.
15. Melquist, W.E. and W.G. Hornocker. 1979. Development and use of telemetry technique for studying river otter. in: F.M. Long, ed. Proceedings of the Second Annual International Conference on Wildlife Biotelemetry. Laramie, Wyoming.
16. Melquist, W.E. and M.G. Hornocker. 1983. Ecology of river otters in west central Idaho. Wildlife Monograph #83. 60pp.
17. Reid, D. 1984. pers. comm. Univ. Calgary.
18. Reid, D. 1982. River otter ecology in northeastern Alberta; Annual Report 1981-1982. Univ. Calgary. 22pp.
19. Reid, D. 1983. River otter ecology in northeastern Alberta; Annual Report 1982-1983. Univ. Calgary. 37pp.
20. Reid, D. and S. Herrero. 1982. Behavioural Ecology of the river otter in northeastern Alberta. Progress Report, Univ. Calgary. 9pp.
21. Ryder, R.A 1955. Fish predation by the otter in Michigan. J. Wildl. Manage. 19:497-498.
22. Sheldon, W.G. and W.G. Toll. 1964. Feeding habits of a river otter in a reservoir in Central Massachusetts, U.S.A. J. Mammal. 49:449-455.
23. Todd, A. and L.C. Geisbrecht. 1979. A review of Alberta fur production and management, 1920-21 to 1977-78. Alberta Energy and Natural Resources, Fish and Wildlife Division. 64pp.
24. Towell, D.E. 1974. Winter food habits of river otters in western Oregon. J. Wildl. Manage. 38:107-111.
25. Zackheim, H.S. 1982. Population status and distribution of the river otter in southwestern Montana. Montana Wildlife Div. Proj. No. W-120-R-13, Study No. FB-2.0. 103 pp.




Figure 12a: General Distribution of Habitat Quality for River Otter in Alberta.





Figure 12.b: Distribution of River Otter Harvest; taken from Boyd, 1977.

MARTEN



13. KEY HABITAT REQUIREMENTS FOR MARTEN

13.1 General

Marten (Martes americana) inhabits late successional forest communities throughout North America.²² The species is most abundant in mature coniferous forests, but also inhabits mixedwood forests. Adequate cover and food availability are important factors influencing habitat selection, especially during winter. Open habitats may be used from spring through fall. Clearcut logging reduces the habitat suitability for marten.²⁵

The distribution and relative quality of marten habitat in Alberta is shown in Figure 13a. Distribution of marten harvest in Alberta is shown in Figure 13b.

13.2 Cover

13.2.1 Vegetation

Pine marten rely on closed, coniferous-dominated, mixedwood forests which are mature or over-mature.^{4,10,14,19, 22,23,24,29}

In the Eastern Slopes Region, fir (Abies Spp.), Engelmann spruce (Picea engelmannii) and white spruce (Picea glauca) dominated stands of moderate to high density provide good marten habitat; mature lodgepole pine (Pinus contorta) stands will also be utilized if they include spruce and fir or adjoin such stands.²⁸

In boreal forest habitat, mesic coniferous and mixed coniferous-deciduous stands of moderate to high density appear to be used most frequently; upland white spruce dominated stands are of particular importance.^{8,24}

At Grande Cache, marten used lodgepole pine and spruce stands in proportion to their availability.⁴

Streamside reserves protected from forest cutting could be important as movement/dispersal corridors and reservoirs for marten re-establishment in areas of extensive timber removal.²⁸

Marten have a greater dependence on treed areas in winter; a canopy cover of >30 per cent (preferably 40-60 per cent) has been suggested for optimum winter habitat.^{1,19}

Marten avoid extremely dense stands which suppress herbaceous cover.

Marten use more open habitats during spring through fall if they provide adequate cover and food.^{13,19,27}

Cover is important as protection against aerial predation.³

Fallen trees which protrude above the snow are important in providing subnivean access to active small mammal prey.⁵

Summer resting sites include the crowns of conifers in uncut softwood and mixedwood stands.²⁶

Winter resting sites are generally beneath the snow in uncut coniferous stands, at edges of clearcuts and residual stands; most sites are in or under hummocks associated with stumps or hollow logs, and under standing snags; maternal dens also occur in such areas.²⁶

13.2.2 Land Forms and Topography

In mountainous areas of Alaska, marten were found at altitudes of 1 219-1 524 m as on summer ranges and <1 036 m on winter ranges.²¹

Ground burrows, rock piles and crevices are used as refuge sites.²³

Maternal dens are often underground or in boulder areas.²⁷

13.2.3 Climate

Closed coniferous or mixedwood stands provide more moderate micro- climate in winter for marten than other stands as temperature differentials and radiation fluxes are reduced.¹⁷

Greater snow depth in openings is a factor influencing selection of more dense cover.¹⁹ Once snow depths exceed 30 cm, marten were found to avoid open areas >100 m in width.¹⁸

Marten remain active during periods of extreme cold (-40°C) and heavy snowfall, as well as during light and moderate rainfall,¹⁶ though movements decrease under these conditions.²¹

13.3 Food

13.3.1 Vegetation/Animal Matter

Marten consume a wide variety of food materials if available.

Marten feed extensively, year round, upon small mammals,^{3,7,9,10,21,25,30} the main food items in Alberta being meadow voles (Microtus pennsylvanicus) and red-backed voles (Clethrionomys gapperi).¹¹

Other mammals in the diet include snowshoe hare (Lepus americanus), red squirrels (Tamiasciurus hudsonicus), ground squirrels (Spermophilus Sp.), weasel (Mustela Sp.), mink (Mustela vison), marten and carrion.^{3,11,21}

Passerine birds and their eggs may be taken,¹⁰ as well as grouse species;¹¹ these have a greater importance in the summer diet.

In summer, berries such as wild strawberry (Fragaria virginiana), huckleberry (Vaccinium membranaceum), and raspberry (Ribes Sp.) may be common components in the diet.^{3,9,10,21,25,30}

Insects, when abundantly available, may also be included in spring and summer diets.^{3,30}

In winter, marten hunt by investigating the bases of trees where rodents often surface and where snow depths are shallow enough to provide ready access to the subnivean environment.³

Habitat use by marten was found to closely parallel relative abundance of red-backed voles,^{17,24}.

Changes in abundance or availability of food may be reflected strongly in the marten's movements, its productivity, and habitat selection.²¹

13.3.2 Climate

Excessive snow depth (>30 cm) limits access to subnivean prey.^{3, 4,19}

In open areas, snow further reduces cover resulting in little use of these areas for foraging.^{3,19}

13.4 Space

13.4.1 Territory/Home Range

Marten populations are structured around male territories, which are rigidly defended during spring and summer months.⁶

Home ranges of males are distinct, but female home ranges often overlap those of other females and males.⁶

Home range sizes for males and females were, Wyoming, 2.4 km² and 0.9 km²;⁵ Ontario, 3.6 km² and 1.1 km²;⁹ Montana, 2.4 km² and 0.7km².¹⁵

Males and juvenile females tend to move greater distances than adult females.²¹

When food is very abundant, marten movements are greatly reduced.²¹

Marten appear to utilize shifting foraging areas within a larger home range area;^{3,10} males in Ontario had larger average foraging areas than females (1.68 and .75 km², respectively) and tended to shift their ranges more frequently than females.¹⁰

Vertical migrations are known in Alaska; movements reflect seasonal shifts in abundance of food at different altitudes, related to the depth of snowfall.²⁰

13.4.2 Population Densities

Marten densities fluctuate in relation to habitat types²⁵ and changes in the availability of food.³⁰

In the Northwest Territories, highest densities were found in unbroken and slightly broken tracts of upland, coniferous-dominated stands of mixedwood forest. Riparian conifer-dominated forest types also received substantial use.^{31,32}

Marten densities recorded in Ontario and Montana were 0.6-0.9/km² and 0.5-1.7/km², respectively.³⁰

In Maine, partial timber harvesting had little effect on the density of marten, though commercial clearcutting reduced densities to about 25 per cent of original values.²⁵

13.5 Special Considerations

13.5.1 Size, Shape and Juxtaposition of Habitat Components

Large tracts of unbroken mature forest with a well-developed understory support high marten populations.

Marten select for closed canopies +/- high density shrub cover.⁸

Stands of mature to over-mature (> 100 years) coniferous forest, comprised of 40 per cent fir or spruce, with a total canopy closure greater than 50 per cent should provide near optimal winter habitat.¹

Marten reliance on dense mature mixed or coniferous stands may be related to prey abundance, evasion of predators, winter availability of prey, thermoregulation and availability of den sites.²⁸

Windfalls, associated with old stands, are important as den sites as well as providing subnivean access.⁵ Windfalls covering 20-50 per cent of the ground surface is considered optimum.¹

Homogenous, mature spruce forest may support more marten over a given area, although a diversity of forest communities may support more marten over time.²⁸ A diverse forest is less susceptible to large-scale perturbations, provides a broader year-round food base, and continuous marten habitat over time through succession.

Marten require at least 2.6 km² of suitable habitat before an area will be occupied.¹ Corridors of adequate canopy and width are required to provide marten with cover when moving between suitable habitat patches.

Open areas >100 m in width are generally avoided during winter¹⁹, though other cover (slash piles) may permit the use of these areas.²⁸

Clearcuts will receive use if small islands of old stands are present.¹²

13.5.2 Significance of Disturbance Phenomena

Fire. Fire has been an important agent in establishing and maintaining a diversity of forest communities; it creates and maintains openings where abundant fruits, insects and small mammals provide food items for the marten during the summer.¹⁹

Trapping. Marten populations under sustained trapping pressure may not be capable of increase; a relatively large protected population of marten may be required to ensure population increase and expansion into adjacent trapped areas.³

Logging. Timber harvest can contribute to the decimation of marten populations.³³ In Wyoming, marten did not utilize harvested timber stands for at least a year; in Maine, clearcuts <15 years old were rarely used though marten occurred in partially harvested stands.²⁵

Clear-cutting lowers the carrying capacity of an area for marten, resulting in larger home range size and lower population densities.²⁵ Freshly logged areas with slash piles, cull trees and other debris may receive notable use, even in winter, especially if small islands of old growth are scattered through the cutover.¹²

13.6 Limiting Factors

The per cent tree canopy closure (>50 per cent) and successional stage of the stand (> 100 years) are the two most limiting variables for determining the suitability of marten winter habitat.¹

The presence of little or no spruce or fir in a forest stand will lower the value of the habitat for marten.¹

Timber harvest can greatly reduce habitat quality for marten.

Excessive snow depth limits access to subnivean prey; prey availability affects population densities.²¹

Downfall is essential to provide optimum winter habitat.¹ Inadequate amounts may decrease the availability and accessibility of prey for marten.

13.7 Regional Variation

Regional importance of marten varies significantly throughout the province.

In terms of numbers harvested, the northwest corner and along the southwestern border of the province are more productive marten areas.² These areas correspond to the Hay River, lower foothills and upper foothills regions which are predominantly spruce and pine habitats with limited mixedwood elements.

ACKNOWLEDGMENTS

Pat Dwyer and Hugh Wollis (Alberta Fish and Wildlife, Edmonton) reviewed and provided useful comments on this summary.

SELECTED REFERENCES

1. Allen, A.W. 1982. Habitat suitability index model: marten. Fish and Wildlife Service. U.S. Dept. Int. 9pp.
2. Boyd, M. 1977. Analysis of fur production records by individual fur-bearing species for registered trapping areas in Alberta, 1970-1975. Alberta Fish and Wildlife. 72pp.
3. Boyd, M.G. 1978. Management of marten, fisher and lynx in Saskatchewan with special reference to the effects of forest harvesting in the mixedwood boreal forest. M.Sc. Thesis. Univ. Calgary. 126pp.
4. Boyd, M.G., C.R. Fox and J. Fox. 1976. Recordings of marten, fisher and lynx along a winter trapline-Grande Cache, Alberta, 1975. Unpub. Rep. Alberta Fish and Wildlife Division. 6pp.
5. Clark, T.W. 1977. Analysis of pine marten population organization and regulatory mechanisms in Jackson Hole, Wyoming. Nat. Geog. Soc. Rept. 46.
6. Clark, T.W. and T.M. Campbell. 1976. Population organization and regulatory mechanisms of martens in Grand Teton National Park, Wyoming. Proceedings of the First Conference on Scientific Research in the National Parks, U.S.D.I. Natl. Park Serv., Trans. Proc. Series 5. Vol. I pp. 293-295.
7. Cowan, I.M. and R.H. Mackay. 1950. Food habits of the marten (Martes americana) in the Rocky Mountain region of Canada. Can. Field-Nat. 64:100-104.
8. Douglass, R.J., L. Fisher and M. Mair. 1976. A study of furbearers with particular reference to marten (Martes americana) at Chick Lake, N.W.T. Unpubl. Report. Northern Engineering Services, Calgary. 38pp. and appendices.
9. Douglass, R.J., L.G. Fisher and M. Mair. 1983. Habitat selection and food habits of marten, Martes americana, in the Northwest Territories. The Can. Field-Nat. 97:71-74.
10. Francis, G.R. and A.B. Stephenson. 1972. Marten ranges and food habits in Algonquin Provincial Park, Ontario. Min. Nat. Res. Rep. No. 91. 54pp.
11. Giesbrecht, L. and A. Todd. 1979. A study on the winter food habits of Alberta marten. Alberta Fish and Wildlife Division. Edmonton. 12pp.
12. Grakov, N.N. 1972. The effect of extensive clear felling on the abundance of the pine marten. Bull. Mosk. Obshchest. Ispyt. Prir. Otd. Biol. 77:14-23.

13. Hagmeier, E.M. 1956. Distribution of marten and fisher in North America. Can. Field-Nat. 70:149-168.
14. Hawley, V.D. 1955. The ecology of the marten in Glacier National Park. M.Sc. Thesis, Montana State Univ. 131pp.
15. Hawley, V.D. and F.E. Newby. 1957. Marten home range and population fluctuations. J. Mammal. 38:174-184.
16. Herman, T. and K. Fuller. 1974. Observations of the marten, Martes americana in the Mackenzie District, Northwest Territories. Can. Field-Nat. 88:501-503.
17. Kelly, J.P. 1977. Pine marten-forest interaction. in: A. Todd, ed. Proc. 1977 Western Fur Managers Conference. p. 124-134.
18. Koehler, G.M. 1975. The effects of fire on marten distribution and abundance in the Selway-Bitterroot and Wilderness, Idaho. Idaho Co-operative Wildlife Research Unit. U. of Idaho.
19. Koehler, G.M. and M.G. Hornocker. 1977. Fire effects on marten habitat in the Selway-Bitterroot Wilderness. J. Wildl. Manage. 41:500-505.
20. Lensink, C.J. 1953. The home range of marten and its significance in management. Fish and Wildlife Service, Fairbanks, Alaska.
21. Lensink, C.J., R.O. Skoog, and J.L. Puckley. 1955. Food habits of marten in interior Alaska and their significance. J. Wildl. Manage. 19:364-368.
22. Marshall, W.H. 1951. Pine marten as a forest product. J. For. 49:899-905.
23. Mech, L.D. and L.L. Rogers. 1977. Status, distribution and movement of martens in northeastern Minnesota. USDA For. Ser. Paper NC-143. 7pp.
24. More, G. 1978. Ecological aspects of food selection in pine marten (Martes americana). M.Sc. Thesis. Univ. Alberta. 84pp.
25. Soutiere, E.C. 1979. Effects of timber harvesting on marten in Maine. J. Wildl. Manage. 43:850-860.
26. Steventon, J.D. and J.T. Major. 1982. Marten use of habitat in commercially clear-cut forest. J. Wildl. Manage. 46:175-182.
27. Streeter, R.G. and C.E. Braun. 1968. Occurrence of marten, Martes americana, (Carnivora:Mustelidae) in Colorado alpine areas. Southwest. Nat. 13:449-451.
28. Todd, A. 1978. Guidelines to habitat and land management for furbearers in the East Slopes Region. Alberta Fish and Wildlife Division. 23pp.

29. de Vos, A. 1952. The ecology and management of fish and marten in Ontario. Ont. Dept. Lands Forests Tech. Bull. 90pp.
30. Weckwerth, R.P. and V.D. Hawley. 1962. Marten food habits and population fluctuations in Montana. J. Wildl. Manage. 26:55-74.
31. Wooley, D.R. 1972. A study of marten (Martes americana) in the Mackenzie District, N.W.T. Chapter III in: R.A. Ruttan and D.R. Wooley, eds. studies of Furbearers Associated with Proposed Pipeline Routes in the Yukon and Northwest Territories. Arctic Gas Biological Report Series, Vol. 9.
32. Wooley, D.R. and R.J. Douglass. 1974. A study of furbearers with particular reference to marten (Martes americana) in: D. Reid, ed. A preliminary report of monitoring studies conducted at Chick Lake, N.W.T. Arctic Gas Biol. Rept. Ser.
33. Yeager, L.E. 1950. Implications of some harvest and habitat factors on pine marten management. Trans. N. Am. Wildl. Conf. 15:319-334.





Figure 13.b: Distribution of Marten Harvest, 1970-1975; taken from Boyd, 1977.

SHARP-TAILED GROUSE



14. KEY HABITAT REQUIREMENTS FOR SHARP-TAILED GROUSE

14.1 General

Sharp-tailed grouse (Pedioecetes phasianellus) occupy a variety of habitats in North America from open grasslands and sagebrush (Artemisia cana) through parklands, to early successional stages and open portions of mixedwoods and boreal forests.² They have also adapted to areas where agriculture has created a mosaic of native vegetation and to cultivated lands which provide sharptails with both food and cover requirements.^{13,31}

Sharp-tailed grouse are scattered throughout Alberta in areas of suitable habitat. The distribution and relative quality of habitat is shown in Figure 14.

14.2 Cover

14.2.1 Vegetation

Sharptails require large areas of grass and shrubby cover during the spring and summer.^{26,1,17,10,3}

Large acreages of grassy and/or shrubby vegetation is required in close proximity of a dancing ground if that area is to be maintained permanently.³¹

Sharptails favor edge habitats created by brush and aspen groves in the parkland zone and forest clearings near grain fields.

Male sharp-tailed grouse attending spring dancing grounds roost in nearby low shrub communities of snowberry (Symphoricarpos albus), rose (Rosa sp.) and silverberry (Elaeagnus commutata)^{11,30} for protection from predators.

Females usually nest within 0.8 km of dancing grounds. A lush and dense residual growth of grass-sedges associated with short shrubs such as snowberry or rose is preferred nesting cover.^{16,31,11,34}

Nesting habitat varies but is usually located in native grass and shrub, tame hay fields or cultivated stubble.³¹

Sharp-tails do not nest in treed bluffs or groves and nests are rarely found in vegetation taller than 6 m. In addition, nesting sites are rarely ever found in vegetation where the dominant species are shorter than 24.5 cm.^{31,16,5,12}

In aspen parkland, nests are in close proximity to stands of woody vegetation, either aspen (Populus tremuloides) forest or dense shrub regrowth aspen. Distances to these vegetation types range from 4.7 to 14.0 m^{32,31}

The physiognomy of cover is more important than species composition, generally heavy cover is chosen for the nest site.^{20,31}

Sharp-tails nesting in undisturbed natural grass-shrub vegetation have significantly greater success than in areas under cultivation.³¹

Immediately after hatching, brood mobility is limited and broods are often found near their nest site.^{4,12} Habitat selection is toward low grassy openings in brush. Use of shrubby vegetation increases as the broods become more mobile at four to six weeks of age.⁴

Broods use a greater variety of plants for cover than do nesting hens. Broods utilize cover composed of a greater proportion of forbs and a smaller proportion of grasses and shrubs than that used by nesting hens.^{31,6}

Sharptails utilize plant invaders more for brooding than nesting and broods can generally tolerate more utilization of cover by domestic stock than do nesting hens.

Broods use light cover in early morning and evening but prefer heavier cover at midday.³¹

During the summer, broodless females spend much of the day in open forest cover or along the edge of a forested area.³²

As summer progresses, hens move their broods to areas of heavy cover, either aspen forest edge or dense aspen regrowth. Such areas are used extensively for midday resting. Morning and evening movements from these areas are only for short distances.³²

During summer, females with broods use grassland and grassland-low shrub transition zones significantly more than other grouse including both males and females without broods.²⁹

Sharptails make extensive use of woody vegetation for cover once herbaceous vegetation becomes desiccated. The extreme dryness may cause females and broods to seek heavier cover.⁸

In summer, grouse without broods select taller vegetative types, particularly tall shrubs. The heavier vegetation probably provides better cover from avian predators as well as cooler microclimate during the hottest part of the day.²⁹

In autumn, sharptails move increasingly to woody cover for shelter from cold winds and predators.^{17,70}

With the onset of winter and sub-zero weather, sharptails roost by burrowing into soft snow. The lee of aspen bluffs and clumps of low shrub also provide the birds with a degree of protective cover during adverse winter conditions.

In winter, grouse move into more open cover including (frozen) marsh during midday and roost predominantly in low cover in the evening.²⁹

14.2.2 Land Forms and Topography

Native, grass-shrub vegetation on washed moraine, fluvial, fluvial lacustrine and high relief moraine land forms is likely to provide very good habitat for sharp-tails.

The most important characteristic of leks (dancing grounds) is their location on a slight rise or open flat area which affords the displaying grouse a wide horizontal field of vision in all directions.

14.2.3 Aquatic Forms

During winter, tall wetland sedges and cattails provide suitable roosting conditions.

14.2.4 Climate

Grouse often burrow under the snow to spend cold winter nights in protected tunnels.

The relationship between weather and plant community selected by grouse is not especially strong. Other factors such as food requirements, presence of conspecifics, familiarity with an area and avoidance of predators seem to have more importance in influencing choice of vegetation by grouse.²⁹

14.3 Food

14.3.1 Vegetation

Sharptails are omnivorous and their diet includes fruits, green leaves, buds and insects.^{31,19,34,23}

In spring, grouse feed mainly in grassland-low shrub transition zones and around the periphery of treed areas during early morning, and then seek taller vegetation for cover throughout midday. Grouse return to these grassland-low shrub communities after midday to again feed and roost.²⁹

Favorite spring foods include aspen catkins, the first succulent leaves of forbs such as dandelion (Taraxacum sp.) and fruits which remain from the previous year.

Sharptail broods feed extensively in natural grass-shrub, cultivated crops and tame haylands but usually shun pastures grazed by livestock.³¹

All food taken by chicks until two weeks of age consists of animal material and other foods make up only 1.4 per cent by volume of the three to four week-old's diet.³¹

For chicks from five to 10 weeks of age, animal material still composes approximately 60-70 per cent of total food by volume.³¹

Plant material, largely composed of cultivated crops and weedy species, constitutes a rapidly increasing percentage of chick food in the older age classes.³¹

Chicks in the 13-16 week age group consume 2.5-6.5 per cent insect matter. High energy food such as domestic grains, seeds, berries, rosehips and green leaves are utilized exclusively by both juvenile and adult birds in late summer and fall.³¹

Sharptails exhibit mainly vegetarian food habits except for the chicks, and all food is largely composed of seeds and berries of wild fruits (most notable being rose), green leaves and flower heads and domestic grains where available.³¹

During the summer, sharptails feed in grasslands during early morning, and in the evening return to these open vegetation types to again feed and roost.²⁹

In autumn, grouse use open vegetation communities in early morning for feeding, and move into taller, heavier cover by mid-morning. From midday to dark, grouse select grassland - low shrub transition for food and roosting.²⁹

As fall foods become less available with the onset of winter, sharptails utilize buds of aspen, chokecherry (Prunus virginiana), birch (Betula sp.) and willow (Salix sp.). They still take fruits from rose, silverberry (Elaeagnus commutata) and snowberry (Symphoricarpos albus) as well as domestic grain where available.¹⁸

Favorable winter habitat consists of wooded areas where tree buds are available and grassy areas which supply seeds.

In winter, sharptails often perch in aspen trees to eat buds during morning and midday.²⁹

Sharptails have never been shown to depend upon drinking water in their natural habitat, apparently meeting moisture requirements from succulent foods.

14.3.2 Trace Elements

Unlike some gallinaceous birds, adult sharptails do not appear dependent on gravel for digestive grit because of the hard grinding surfaces provided by the seeds of fruits commonly eaten.¹⁵

Chicks, on the other hand, being almost totally insectivorous, consume large amounts of grit.³¹

14.4 Space

14.4.1 Territory/Home Range

On the dancing ground, the resident males establish small territories adjacent to each other. Each territory is usually only 1.8 to 2.7 metres across and variable in shape.³¹

Established males appear to be faithful to one dancing ground and frequent that particular ground during breeding seasons, and during the autumn and winter, for as long as they are capable of defending a territory.^{30,31,14}

The location of most summer observations of sharptails is found to be within a 1.6 km radius of a dancing ground.³¹

Sharptail nesting sites are located within 1.6 km of the nearest dancing ground and on average the distance is approximately 900 metres.³¹

Non-brooding females make short, daily movements within small, definable areas. Females with broods moved over larger areas and do not appear to have a constant home range for any length of time.³²

Tracking data indicate that movement by broods between one and two weeks of age were quite limited, generally less than 0.2 km per day. By three to four weeks of age, movements averaged 0.5 km per day. Older broods also averaged 0.5 km per day but some long distance movements typical of the four to 10 week-plus age groups were as far as 1.5 km in an eight hour-period.³²

14.5 Special Considerations

14.5.1 Size, Shape and Juxtaposition of Habitat Components

Sharptails attain highest densities in habitat mosaics of grass, shrubs and trees. Shelter from extremes of climate and predators is important for survival.

It is believed that grouse have specific preferences for different plant communities at different times of the year and of the day. Thus a mosaic of plant communities, particularly grassland and grassland-shrub mixtures with extensive ecotones, provides optimum habitat.²⁹

The presence of suitable nesting and brooding cover within 1.6 km of the dancing ground strongly influenced the breeding population.^{31,11}

Grouse in the Prairies prefer large acreages of grassland, preferably in various stages of succession if maximum populations are to be maintained.²²

Female sharptail grouse require a diversity of habitat types. Open areas are required for feeding, shrubby areas for nesting, and areas of tall aspen regrowth or forested stands for escape cover and midday resting sites for broods.³²

Native grass and shrub areas are essential in maintaining year-round sharptail populations.^{31,8,19}

14.5.2 Significance of Disturbance Phenomena

Fire. Because of increasing agricultural use and fire prevention, trees are invading the grasslands in aspen parkland areas. This is leading to a reduction in the number of arenas and consequently lower productivity.³⁵

Fire is known to be an important factor in creating and maintaining seral and subclimax grass-shrub conditions favored by sharp-tailed grouse.^{3,21,33}

Fire-disturbed areas have large numbers of insects which supply high protein animal matter required as food by newly hatched grouse for the first four to six weeks following hatching.³⁵

Agriculture. Intensive grazing by over-stocking of a grassland ecosystem leads to destruction of vegetation used by nesting sharptail hens and also damages the woody cover used by sharptail broods.²⁷ Decline of sharp-tailed grouse populations caused by intensive grazing have been noted by many researchers.^{9,31,24,34,37}

Pastures grazed with a season-long grazing system produced better grassland habitat for sharp-tailed grouse than pastures with a deferred rotation grazing system.²⁷

Initially, agricultural development was beneficial to grouse distribution and numbers as large tracts of native vegetation were still interspersed with cropland areas. However, in recent times, populations have declined as this diversity has been destroyed with the advent of modern, intensive clean-farming practices.

Weather/Predation. Damp, windy weather and harassment by humans or predators depresses breeding activity.³¹ Hens seldom return the same morning after having been disturbed on a lek.³⁰

14.6 Limiting Factors

Carrying capacity for sharptails is limited by the availability of large acreages of ungrazed grass-shrub and hayland within a 1.6 km radius of a dancing ground. This ungrazed herbaceous vegetation is very important to the habitat requirements of sharptails during both the spring breeding season and the summer nesting and brooding season.³¹

The amount of continuous aspen cover within a radius of 0.8 km of a display site or arena is inversely related to the number of grouse which frequent the area. This apparent inverse relationship suggests that with increasing percentages of aspen cover, fewer sharptails will be present.²⁹

Homogeneous grassland vegetation lacking a shrub component is unattractive to sharptails because it does not offer sufficient protective cover and forage capabilities.

Light to moderate grazing in tall to medium grasslands of the Prairies, reduced, and in some cases eliminated, sharptails.²²

Agricultural activities such as haying or summer fallowing cause significant nest destruction or abandonment.³¹

Sharptail chicks feed heavily on insects, consequently they may be subjected to possible hazards from chemical pesticide control used in association with agricultural practices.³¹

Applications of sublethal doses of dieldrin, an insecticide used to control agricultural insect pests, caused sharptails to be more prone to predation for up to 12 days after application.²⁸

The elimination of wild prairie fires which destroy woody vegetation and favor the growth of forbs and grasses, results in tree and tall shrub-dominating growth and reduces habitat quality.^{22,25,36}

14.7 Regional Variation

The distribution of prime sharptail habitat in the province is located mainly in the Short Grass, Mixed Grass, Aspen Parkland, and southern-southwestern Boreal Mixed-wood Ecoregions.

ACKNOWLEDGMENTS

D. Moyles (Alberta Fish and Wildlife, Edmonton) critically reviewed this summary.

SELECTED REFERENCES

1. Aldous, E. 1943. Sharp-tailed grouse in the sand dune country of north-central North Dakota. J. Wildl. Manage. 7(1):23-31.
2. Aldrich, J. W. 1963. Geographical orientation of American Tetraonidae. Wild. Manage. 27:529-545.
3. Ammann, G. A. 1957. Prairie grouse in Michigan. Game Division, Dept. of Conservation, Lansing. 200pp.
4. Artmann, J. W. and J. R. Beer. 1970. Spring-summer movements and habitat usage by female sharp-tailed grouse. Paper presented at Midwest Fish and Wildlife Conference, December 6 to 9, 1970, Winnipeg, Manitoba.
5. Bernhoft, L. S. and G. Kobriger. 1969. Habitat preferences of the sharp-tailed grouse. North Dakota State Game and Fish Dept. Mimeo. Report. No. A-362. 37pp.
6. Blus, Lawrence J. 1964. Studies in the ecology and management of prairie grouse. Job completion report, P.R. proj. W-33-R-6. Job 1. Nebraska Game, Forestation and Parks Commission. 6pp.
7. Brown, R. L. 1968. Sharp-tailed grouse population study. Montana Fish and Game Dept. Proj. W-91-R9, Job 11-E. Unpubl. mimeo. 18pp.
8. Brown, R. L. 1966. Response of sharptail breeding populations to annual changes in residual grassland cover. Paper presented at 46th Annual Conf. of Western Assoc. of State Game and Fish Commissioners, Butte, Montana, July 12-14. 1966.
9. Brown, R. L. 1963. Progress of a sharp-tailed grouse population study in two mixed prairie regions of northcentral and southwestern Montana. Montana Game and Fish Dept. Helena. 5pp.
10. Buss, I. O. and E. S. Dziedzic. 1955. Relation of cultivation to the disappearance of the Columbian sharp-tailed grouse from Southeastern Washington. Condor 57(3):185-187.
11. Caldwell, P. J. 1976. Energetic and population considerations of sharp-tailed grouse in the aspen parkland of Canada. Kansas State Univ. Manhattan. Unpubl. Ph.D. thesis. 109pp.
12. Christenson, C. D. 1971. Habitat preference of the sharp-tailed grouse. North Dakota State Game and Fish Dept. P.R. Div. Proj. W-67-R-11. Phase B. Job No. 13. Mimeo. 53pp.
13. Evans, K. E. 1968. Characteristics and habitat requirements of the Greater Prairie Chicken and Sharp-tailed Grouse - a review of the literature. U.S. Department of Agriculture Forest Service Conservation Research Report Number 12.

14. Evans, R. M. 1961. Territorial stability in sharp-tailed grouse. *Wilson Bull.* 81(1):75-78.
15. Farnes, R. E. 1951. The sharp-tailed grouse. *The Conservation Volunteer.* July-August: 10-14.
16. Folker, R. V. 1964. A study of sharp-tailed grouse on the Gibbs Study Area. *Sask. Dept. Nat. Resour., Regina.* Unpubl. Mimeo. 30pp.
17. Hamerstrom, F. N. Jr. and F. Hamerstrom. 1951. Mobility of the sharp-tailed grouse in relation to its ecology and distribution. *The American Midl. Nat.* 46(1):174-226.
18. Harris, S. W. 1967. Fall foods of sharp-tailed grouse in Minnesota. *J. Wildl. Manage.* 31(3):585-587.
19. Hillman, C. and W. W. Jackson. 1973. The sharp-tailed grouse in South Dakota. *S. Dak. Dept. of Game, Fish and Parks, Tech. Bull. No. 3.* 62pp.
20. Jones, R. F. 1968. A board to measure cover used by prairie grouse. *J. Wildl. Mgmt.* 32(1):28-31.
21. Kirsch, L. M. and A. D. Kruse. 1972. Prairie fires and wildlife. *Proceedings of the Tall Timbers Fire Ecology Conference* 12:289-303
22. Kirsch, L. 1969. Prairie grouse and land use. Paper presented at 8th Conf. Prairie Grouse Technical Council, Sept. 9-10-11, 1969. Woodward, Okla.
23. Kobriger, G. D. 1965. Status, movements, habits and foods of prairie grouse on a sandhills refuge. *J. Wildl. Manage.* 29(4):788-800
24. Kohn, S. C. 1976. Sharp-tailed grouse nesting and brooding habitat in southwestern North Dakota. M.S. Thesis. South Dakota State Univ., Brookings. 123pp.
25. Kruse, A. 1971. Land-use and prairie grouse populations on arrowwood National Wildlife Refuge. Paper presented at 9th Conference of the Prairie Grouse Technical Council, Sept. 14, 15, 16, 1971. Dickinson, North Dakota.
26. Ligon, J. Stokley. 1927. *Wildlife of New Mexico, its conservation and management.* State Game Comm., Santa Fe, New Mexico. 212pp.
27. Mattise, S. N. 1978. Effects of pasture management on sharp-tailed grouse nesting and brooding habitat in southwestern North Dakota. *North Dakota State Game and Fish Dept., Project no. W-67-R-18.* 18pp.
28. McEwen, L. C. and R. L. Brown. 1966. Acute toxicity of dieldrin and malathion to wild sharp-tailed grouse. *J. Wild. Manage.* 30(3):604-611.

29. Moyles, D. L. J. 1981. Seasonal and daily use of plant communities by Sharp-tailed Grouse (Pedioecetes phasianellus) in the parklands of Alberta. Canadian Field-Naturalist 95(3):287-291.
30. Moyles, D. L. J. 1977. A study of territory established by and movements of male sharp-tailed grouse (Pedioecetes phasianellus) relative to the area. M.Sc. thesis. Univ. of Alberta, Edmonton. 91pp.
31. Pepper, G. W. 1972. The ecology of sharp-tailed grouse during spring and summer in the aspen parklands of Saskatchewan. Saskatchewan Department of Natural Resources, Wildlife Report Number One. 56pp.
32. Sealy, S. G. and D. A. Sexton. 1976. Movements and habitat use by female sharp-tailed grouse. Manitoba-Wildlife Research Progress Report.
33. Sexton, D. A. and M. M. Gillespie. 1979. Effects of fire on the location of a sharp-tailed grouse arena. Canadian Field-Naturalist 93(1):74-76.
34. Sisson, L. 1976. The sharp-tailed grouse in Nebraska. Nebraska Game and Parks Commission, Lincoln. 88pp.
35. Sklebar, H. T. 1982. The effects of fire on the sharp-tailed grouse habitat of the Aspen Parkland in Alberta. Forestry Seminar, Dept. of Forest Science, Univ. of Alberta, Feb. 11, 1982.
36. Stewart, O. C. 1956. Fire as the first great force employed by man. In: Man's role in changing the face of the earth, Ed. by W. L. Thomas, Jr. Univ. Chicago Press, Chicago, III pp. 115-133.
37. Yde, C. A. 1977. Distribution and movements of sharp-tailed grouse during spring and summer in relation to rest-rotation grazing. Montana Game and Fish Dept. P-R Proj. Rep., W-120-R. 70pp.





Figure 14: General Distribution of Habitat Quality for Sharp-tailed Grouse in Alberta.

RING-NECKED PHEASANT



15. KEY HABITAT REQUIREMENTS FOR RING-NECKED PHEASANT

15.1 General

In Alberta, ring-necked pheasants (Phasianus colchicus) exist at the northern limit of their North American range.¹⁴ Dense winter cover, such as cattail marshes or shelterbelts, in close proximity to spring nesting sites and cultivated cropland, is the essential habitat requirement. Wetlands are an extremely important element of pheasant habitat and their retention is of utmost importance in ensuring continued pheasant populations in Alberta.

The distribution and relative quality of ring-necked pheasant habitat in Alberta is depicted in Figure 15.

15.2 Cover

15.2.1 Vegetation

Cattail-willow (Typha latifolia-Salix sp.) complexes and other marshy vegetation associated with wetlands represent the best available winter cover for pheasants.

Dense, low, shrubby cover can be important but its value deteriorates with even moderate snowfall and drifting.¹⁰

In areas of deeper snow such as the parkland region, thicker, more substantial cover such as multi-row shelterbelts are required.

In cold weather, woodlots, low lying forest pockets and narrow strip cover are usually avoided but will be used if other suitable cover is not available.¹⁰

In late summer and fall, pheasants basically prefer to roost in relatively open vegetation types such as grass and stubble fields with close proximity to escape cover;¹⁷ during wind, snow or rain, brushy areas or dense marshy vegetation are used.¹⁷

Loafing areas usually contain dusting grounds and pheasants will choose sunny or shady sites depending on their thermal requirements.

A variety of cover types are used for nesting but undisturbed residual cover (>25cm), such as dry grasses and weedy forbs, are most important for early nesting attempts. These have the best chances of successful hatching.²⁵

As the nesting season progresses, alfalfa (Medicago sativa) becomes increasingly preferred for nesting sites, however, haying usually occurs during peak hatch and as a result nesting success is low.

In South Dakota, alfalfa was the most preferred of all cover, even over natural vegetation.¹³

Stands of cover with more than 50 per cent shrub component have low nesting value.⁶

Wetlands, strip cover along roadsides, fencelines and railways, and fields of hay, fall rye (Secale cereale) or winter wheat (Triticum sp.), which still retain vegetation from the previous year, are optimum areas for nesting.⁹

Hay and strip cover usually carry higher nesting densities than heavily grazed pastures, woodlands or small grazing fields.

Very narrow strip vegetation may have limited value because ground predators can more efficiently search these habitats;²² predators may also use these areas as travel lanes.¹

Other locations often used for nesting cover include: field borders, shelterbelts, marsh edges, stream and ditch banks and abandoned farmsteads.²⁵

In Wisconsin, early nesting attempts are located primarily in hayfields where the vegetation exhibits early growth.⁷

There is a high use of hay, grain and herbaceous vegetation for brooding,¹¹ which is dense enough for predator protection, but not so dense as to prevent chicks from moving about and staying together.¹⁸

15.2.2 Land Forms and Topography

The best pheasant populations in Alberta are associated with irrigation districts.

Preferred vegetation types are found along drainages and in wetland basins where natural cover has not been substantially altered by agricultural land use.

Small localized areas of moraine, fluvial and eroded lands, unsuitable for agricultural development, support remnant pheasant populations; areas suitable for agriculture are more productive.

Pheasants prefer valleys and lowlands in late fall and winter (probably because uplands are cropped); spring dispersal tends to be in the direction of the uplands.²¹

15.2.3 Aquatic Forms

Wetland areas which provide dense emergent vegetation (eg. cattails) and riparian herbaceous/shrub vegetation are most valuable as cover, especially in late fall and winter.

Wetlands can provide escape cover in fall during hunting if water depth is < 12 cm.

Pheasants show a strong tendency toward stream bottoms and coulees,²² especially in southern Alberta where it is their only source of cover and water.

15.2.4 Climate

Snowfall and drifting decrease the effectiveness of cover and make many cover areas unavailable for use.^{10,17}

Severe winter conditions create a need for increased cover requirements (larger and more dense stands of woody and herbaceous cover). Survival may depend on the proximity of food to dense cover.¹⁶

During periods of wind, snow, or rain, pheasants utilize brushy areas or dense, marshy vegetation to a larger degree as roosting sites.¹⁷

15.3 Food

15.3.1 Vegetation

Adult pheasants utilize farm crops more heavily than native vegetation or animal matter.

Seeds, leaves and fruit of domestic crops including corn (Zea mays), wheat (Triticum sp.), barley (Hordeum sp.), oats (Avena sativa), flax (Linum usitatissimum), rye, alfalfa and clover (Trifolium sp.) are the principal foods of pheasants; seeds from native vegetation can be emergency food stuffs and are the sole food supply in some areas.¹⁵

Adult pheasants in South Dakota consumed: 82 per cent farm crops, 7 per cent weed seed, 5 per cent insects, 5 per cent foliage and 1 per cent minerals; juveniles consume the same food items but greater amounts of insects which are a rich source of protein needed for growth.²³

The seeds of farm crops are utilized most heavily in mid-winter while insects and green foliage are important as food items in the spring and early summer.²³

15.3.2 Aquatic Forms

Water in free forms such as ponds, creeks and irrigation ditches is not usually necessary, as dew, insects and succulent vegetation generally provide a sufficient supply.²⁵

In dry areas, free water is important to survival.

15.3.3 Climate

15 cm of snow, especially if packed and covering waste grains, can be sufficient to create food stress unless alternate food is available.¹⁰

Pheasants can scratch through 38 cm of snow, though energy expenditure may be greater than that obtained.¹⁰

In restrictive snow conditions, farmsteads and roadsides may be the final sources of food for pheasants.¹⁰

15.3.4 Trace Elements

Calcareous grit, used as a grinding agent in the gizzard, is also an important source of calcium for pheasants whose normal diet of cereal grains is low in this element.^{4,15}

15.4 Space

15.4.1 Territory/Home Range

Pheasants generally have very small home ranges.^{3,20}

Most literature suggests that daily movements are usually within one kilometer.¹⁶

Seasonal movements usually range from 1-3.5 km and very seldom exceed 5.5 km.¹⁶

15.5 Special Considerations

15.5.1 Size, Shape and Juxtaposition of Habitat Components

The type of land use which is especially productive for pheasants in the plains and prairie is 25 per cent row crops, 50 per cent cereal crops, 15 per cent hay and pasture and 10 per cent idle lands supporting shelterbelts, slough margins, willow thickets and weeds.⁶ This interspersation provides sufficient waste grain for food and meets cover requirements for nesting, brooding and winter shelter.

Pheasants show preference for areas with 1) high hay/non-hay ratio; 2) a high cereal crop/row crop ratio; 3) the greatest possible field crop diversity; 4) an abundance of wetland cover types.⁶

Proper juxtaposition of cover types, with adequate wintering areas linked to feeding and nesting sites by well covered travel lanes, is most important.

Nesting areas should be within a 2.5 km² block (a minimum of 8 ha in size; which also provides the minimum winter cover size requirements). On each of these 2.5 km² blocks, at least 5 per cent of the land and optimally 20 per cent of the land must be left as undisturbed nesting cover, preferably with a high proportion of wetlands.⁸

Densities of territorial males are usually highest in areas of greatest cover interspersation, where food supplies occur in close proximity to roosting and loafing cover.²

Dusting and grit-picking sites tend to be more exposed areas, adjacent to denser escape cover.²⁵

Some studies noted preference for the edge area for nesting while others found the distribution to be random.¹⁶

Prime pheasant habitat usually lies in fertile black or brown soil areas (grassland region) which have the most favorable land use patterns for pheasants.³

15.5.2 Significance of Disturbance Phenomena

Agriculture. Cultivated lands provide pheasants with their prime food supply; more advanced farming practices tend to leave fields virtually bare of waste grains.⁶

The best pheasant habitat exists where 85-90 per cent of the land is under cultivation but sufficient cover may be found on lands with up to 95 per cent cultivation.

High nest and hen mortalities result from mowing in hayfield nesting cover at the time of nesting.¹⁹

Human Activity. Pheasants are bothered little by human activity if sufficient cover is available.

Lanes and roads are often used as dusting and loafing areas.⁵

Pheasant-rearing programs establish and supplement wild pheasant populations in Alberta.

15.6 Limiting Factors

Severe winter conditions in most years may be a primary limiting factor; pheasants are generally restricted to locations where January mean temperatures are above -15°C and mid-winter snow depths average less than 20 cm.²²

They are limited where more than 95 per cent of the land is cultivated, where few wetland acreages occur, where progressively more of the landscape is wooded, where the soils are progressively less fertile and where the growing season is shorter.²⁴

The available habitat of pheasants is decreasing as more intensive agriculture produces larger, cleaner fields reducing both food and cover supplies.

15.7 Regional Variations

Habitat quality and pheasant population numbers decline from the Short Grass and Mixed Grass Ecoregions to the Aspen Parkland Ecoregion, primarily because of increasing winter severity, deeper snow conditions, colder winter temperatures, infrequent chinook events and shorter growing seasons.

Suitable habitat and climate does not exist in other portions of the province.

ACKNOWLEDGMENTS

Critical reviews were provided by D. Moyles (Alberta Fish and Wildlife, Edmonton) and H. Vriend (Alberta Fish and Wildlife, Lethbridge).

SELECTED REFERENCES

1. Baskett, T. S. 1947. Nesting and production of the ring-necked pheasant in North Central Iowa. *Ecological Monographs* 17:1-30.
2. Burger, G. V. 1966. Observations on aggressive behavior of male ring-necked pheasants in Wisconsin. *J. Wildl. Manage.* 30(1):57-64.
3. Christisen, D. M. 1951. History and status of the ring-necked pheasant in Missouri. *Missouri Conserv. Commission.*
4. Dale, F. H. 1954. Influence of calcium on the distribution of the pheasant in North America. *Trans. N. Am. Wildl. Conf.* 19:316-323
5. Duebbert, H. F. 1959. Pheasant populations in relation to land use, soils and weather in southeastern North Dakota. *N. Dak. State Game and Fish Dept. Pittman-Robertson Proj. W-35-R06, Job No. 11.* 49pp + 9 appendices.
6. Fish and Wildlife Division - Southern Region. 1981. Habitat management proposal for the ring-necked pheasant in southern Alberta.
7. Gates, J. M. 1966. Renesting behavior in the ring-necked pheasant. *Wilson Bull.* 78(3):307-315.
8. Gates, J. M. 1970. Recommendations for a scattered wetlands program of pheasant habitat preservation in southeast Wisconsin. *Wisconsin Dept. of Natural Resources. Research Rpt. No. 63.* 24p.
9. Gates, J. M., E. J. Frank and E. E. Woehler. 1970. Management of pheasant nesting cover on upland sites in relation to cropland diversion programs. *Wisc. Dept. Nat. Resour. Resear. Report. No. 48.*
10. Gates, J. M. and J. B. Hale. 1974. Seasonal movement, winter habitat use and population distribution of an east-central Wisconsin pheasant population. *Wisc. Dept. Nat. Resour. Tech. Bull. No. 76.* 55pp.
11. Ginn, W. E. 1962. The ring-necked pheasant in Indiana. *P-R Bulletin No. 6, Indiana Div. Fish and Game.*
12. Godfrey, W. E. 1966. The birds of Canada. *Nat. Mus. Can. Bull. No. 203, Biol. Series No. 73.* Ottawa. 428pp.
13. Hanson, L. E. and D. R. Progulski. 1973. Movements and cover preferences of pheasants in South Dakota. *J. Wildl. Manage.* 37(4):454-461.
14. Kimball, J. W., E. L. Kozicky and B. A. Nelson. 1956. Pheasants of the plains and prairies. In D. L. Allen. 1956. *Pheasants in North America.* The Stackpole Co., Harrisburg, P.A. and Wildlife Management Institute, Washington, D. C. 490pp.

15. McCann, L. J. 1961. Grit as an ecological factor. Am. Midl. Nat. 65(1):187-192.
16. Olsen, D. W. 1977. A literature review of pheasant habitat requirements and improved methods. Utah Div. Wildl. Resour., Publ. No. 77-7. 144pp.
17. Olsen, D. W. and J. P. Leatham. 1980. A manual for pheasant habitat management on private lands in Utah. Utah Division of Wildlife Resources. Publication No. 80-4.
18. Pearce, J. 1945. The pheasant in the Northeast. In W. L. McAtee. Ed. The ring-necked pheasant. The Amer. Wildl. Instit., Wash., D.C.
19. Randall, P. E. 1939. Nesting and causes of nest mortality of the ring-necked pheasant. Penn. Game News 10(9):6, 7, 30.
20. Robertson, W. B. 1958. Investigations of ring-necked pheasants in Illinois. Illinois Dept. Conserv. Div. of Game Manage. Tech. Bull. No. 1.
21. Sharp, W. M. and H. W. McClure. 1945. The pheasant in the sandhill region of Nebraska. In W. L. McAtee. 1945. The ring-necked pheasant. Ed. The Amer. Wildl. Instit., Wash. D.C.
22. Stelfox, H. A. 1979. Terrestrial wildlife habitat inventory of the Weyburn (62E) - Virden (62F) map area. Wildl. Tech. Report. 79-6. Sask. Dept. of Tour. and Nat. Resour.
23. Trautman, C. G. 1952. Pheasant food habits in South Dakota. Tech. Bull. No. 1, South Dakota Dept. Game Fish and Parks. 89pp.
24. Wagner, F. H., C. D. Beadry and C. Kabat. 1965. Population ecology and management of Wisconsin pheasants. Wisc. Conserv. Dept. Tech. Bull. No. 34. Madison. 168pp.
25. Weigand, J. P. and R. G. Janson. 1976. Montana's ring-necked pheasant. Mont. Dept. Fish and Game. 178pp.





Figure 15: General Distribution of Habitat Quality for Ring-necked Pheasant in Alberta.

SAGE GROUSE



16. KEY HABITAT REQUIREMENTS OF SAGE GROUSE

16.1 General

Sage grouse (Centrocercus urophasianus) are limited to the extreme southeastern part of Alberta, where semi-arid grasslands and sagebrush plains form their only habitat.³ Few other species are so totally dependent on and restricted to one habitat type. Sagebrush (Artemisia cana) is essential to sage grouse for food and cover; the birds' diet is primarily sagebrush throughout the year, especially in winter.^{3,27} Thus, sage grouse populations are determined by availability and distribution of sagebrush.

The distribution of sage grouse habitat in Alberta is depicted on Figure 16.

16.2 Cover

16.2.1 Vegetation

Sagebrush communities form the main cover for shelter, nesting, loafing, roosting and feeding.

Vegetation associated with sagebrush varies widely. Western wheatgrass (Agropyron smithii) is the most frequently associated grass at all densities of sagebrush.⁶ Sandberg's bluegrass (Poa secunda) and June grass (Koeleria cristata) are closely associated with Western wheatgrass; pasture sage (Artemisia frigida) is the most frequently associated forb.

Wintering areas are dominated by dense stands of sagebrush cover.⁷

Loafing and roosting areas invariably coincide with the heaviest and most dense sagebrush cover.¹⁸

Winter feeding and loafing sites averaged 28 per cent sagebrush cover in Montana.²⁹

Fifteen percent canopy coverage of sagebrush was classified as the minimum coverage acceptable for sage grouse winter and nesting habitat.^{7,28,29}

In Colorado, sagebrush canopy cover of spring habitat averaged 33 per cent for male-use sites and 28 percent for female-use sites; average sagebrush height at feeding-loafing sites was 28 cm for males and 22 cm for females.²²

Sage grouse prefer relatively open areas as strutting grounds,^{18,23} though these must be surrounded by adequate sagebrush required for food and cover.^{9,27}

In Montana, 80 per cent of the feeding and loafing locations of strutting cocks were in sagebrush canopy of 20-50 per cent (average 32 per cent); no cocks were observed in < 10 per cent cover.²⁹

Sagebrush seems to be preferred for nesting cover, though other vegetation is occasionally used.²

In Montana, average height of sagebrush cover over all nests was 40.4 cm as compared with 23.4 cm in surrounding areas.²⁸ Successful nests were located in sagebrush stands with a higher average canopy coverage than those of unsuccessful nests, and had a significantly greater sagebrush cover within 60 cm of the nest and within a 9 m radius.

The productivity of grasses and forbs in an area may also influence the number of nests and nesting success.¹¹

Unsuccessful breeding hen flocks were found in areas of dense sagebrush throughout the summer in Montana.²⁶

Sage grouse broods in Montana preferred relatively open stands of sagebrush compared to those selected at other times of the year.²⁶

On summering areas in Wyoming, meadowlands and hayfields were used as feeding sites, willow (*Salix* sp.) and sagebrush were used for resting and shading, and areas of sparse, low-growing sagebrush were used as night roosting areas.^{11,18}

16.2.2 Land Forms and Topography

Native sagebrush - grass vegetation confined to primarily heavy clay bottoms, coulees, creeks and rivers provide habitat for sage grouse in Alberta.³

Strutting grounds tend to be in open areas with little slope.¹⁸

In Wyoming, most strutting areas occurred on windswept ridges or exposed knolls, flat, open sagebrush areas, and bare openings on relatively level land.¹⁸

In Montana, most roosting, feeding, and loafing sites were on terrain with little if any slope; birds were never encountered on steep terrain although dense stands of sagebrush occurred there.⁷

In Idaho, where birds undergo elevational migrations, sage grouse exhibited little preference for slope.⁴

Sage grouse may use topography as protection from wind.⁷

16.2.3 Climate

Except for shade from summer heat, adult sage grouse require little protection from the elements.²

During severe winters, dense sagebrush may provide shelter.^{7,26}

Inclement weather may inhibit strutting activities and ground attendance.²⁴

Cold, rainy weather at hatching time may result in lowered productivity.²

Winds greater than 16-24 kmph were found to cause movement off flat ridges.⁷

16.3 Food

16.3.1 Vegetation

Sage grouse do not possess a muscular gizzard and therefore lack the capability of grinding and digesting seeds.

The year-round diet consists of leafy vegetation with the exception of some insects taken during summer.

During winter, sage grouse depend entirely on sagebrush for food, which also forms a staple item in diets during other seasons.^{18,20,27} As an evergreen shrub, sagebrush provides a nutritious, available food source throughout the year.

Feeding primarily occurs in stands of sagebrush of intermediate height and density.¹⁸

Meadows and haylands surrounded by sagebrush are used as summer feeding sites.³

Juvenile grouse rely on insects and forbs during the summer period.^{14,19,20}

Grasshoppers, beetles, and ants are the most common animal matter items in the diet, which may comprise > 40 per cent of the chicks diet.^{19,11}

Forbs become the most important component of the juvenile diet after one to two weeks of age.¹⁴

Succulent forbs, dandelion (Taraxacum sp.) and alfalfa (Medicago sativa), preferred foods of sage grouse broods, are an important element of the summer habitat.^{9,15,26}

Sagebrush begins to form a significant part of the diet only after 12 weeks of age.¹⁴

Sagebrush and forbs are the major components of adult summer diets; availability may influence the proportion in the diet.^{18,27}

16.3.2 Aquatic Forms

Loafing and roosting areas are usually in close association to a water supply.¹⁶

Birds congregate around water during dry periods, but disperse after rain.²⁵

Nesting areas have been found to occur within short distances of water sources, and migrations of broods to summering areas may follow accessible waterways.^{10,11,18}

During years of above average rainfall, sage grouse chicks obtain most of their water from succulent vegetation and dew.²⁵

16.3.3 Climate

Sage grouse may occupy windswept sagebrush ridges during winter where accessible, though lower and scantier growth sagebrush occurs.²¹

When snow depth exceeds approximately 30.5 cm, sage grouse are restricted to taller sagebrush stands, which may represent only a small portion of a normal wintering area.²⁷

Small areas of sagebrush may not be adequate for sage grouse to survive the sometimes severe winters.³

16.4 Space

16.4.1 Territory/Home Range

Sage grouse in Alberta are not known to exhibit any major seasonal movements³, possibly due to interspersions of wintering, nesting and brood habitats as was found for eastern Montana birds.⁷

Winter range, summer range, breeding, nesting and brood rearing may all occur in separate areas,² or a number of activities may occur on one habitat complex.⁷

With snow, sage grouse move to ranges where exposed sagebrush provides food.¹

Winter ranges in Montana varied from 1059 - 3143 ha; 75 per cent of the minimum daily movements were < 1.2 km.⁷

In spring, sage grouse begin movement to strutting grounds which may be on winter or summer range.^{5,9}

Thirty-one strutting grounds have been located in Alberta.¹²

Birds show an affinity for particular strutting grounds.¹⁸

Strutting grounds may vary in size from 0.4 - 16 ha.²³

Total area defended by master cocks rarely exceeds 6.1 m in radius from the primary mating spot (rarely exceeds 1.5 m radius).⁹

After being bred, hens select a nesting site which may be located near the strutting grounds⁵, or at some distance from them, usually in sagebrush.^{23,18}

In Montana, most nests were found within a 2.4 km radius of the strutting ground.²⁸

Males move to summer ranges after they abandon the strutting grounds,⁴ and were found to remain within 3.2-4.8 km of the strutting grounds in Montana.²⁷

After hatching, broods move to summer range,^{8,9} which may be restricted to areas supporting succulent vegetation (40 - 80 ha ranges in Montana).²⁶

16.4.2 Population Densities

Sage grouse are gregarious throughout the year, with the exception of nesting hens.²⁷

Sagebrush density is a major factor influencing sage grouse distributions.⁷

Considerable variability in size of winter flocks was found in Montana; hen flocks varied from 4-50 birds except during severe weather when concentrations of > 200 birds were recorded, winter cock flocks ranged from 5-15 birds and occurred near hen flocks.^{7,27}

Male sage grouse densities on strutting grounds averaged 26.6 cocks/lek in Alberta during 1968-1983.¹²

Hens begin to cluster during breeding, with groups ranging in size from a few to 100 females in Montana.²⁷

Male flocks remain segregated from broods and hen flocks in summer, but may remain in close proximity to them.²⁹

16.5 Special Considerations

16.5.1 Size, Shape and Juxtaposition of Habitat Components

Large expanses of sagebrush, encompassing a water source, and interspersed with meadows and haylands provide sage grouse habitat. Dispersed, small openings (a couple of hectares) on flat ground surrounded by 3-4 km² of dense sagebrush provide good lekking areas with adjacent nesting habitat.

16.5.2 Significance of Disturbance Phenomena

Agriculture. Potential conflicts exist between sage grouse and agricultural activities.

Agriculturalists continue to advocate re-seeding native grasslands to domestic forages to increase livestock production, posing a potential threat to sagebrush prairie.^{3,15}

Although sage grouse may tolerate drastic physical changes in their strutting areas before abandoning them,²⁵ elimination of sagebrush around these may result in abandonment or termination of breeding activities at an earlier date.⁸ A 3.2 km radius buffer zone around the strutting grounds should be protected.⁷

Most of Alberta's sage grouse habitat is used as rangeland for domestic livestock, mainly cattle.³ Cattle may compete with sage grouse for sagebrush food, particularly when too many cattle are wintered on sagebrush ranges.

Overgrazing can cause serious decline in sage grouse populations.¹⁸

A sagebrush climax, with a good balance of grasses and forbs as understory species, through proper grazing practices, is recommended.¹⁸

Human Activity. Strutting grounds which are readily accessible by road may be subjected to disturbance which could seriously affect the breeding of the population.¹²

New strutting grounds may be constructed by removing 0.4-0.8 ha of sagebrush within a sagebrush expanse.¹³

Fire. Natural fires can reduce sagebrush ranges.³

16.6 Limiting Factors

Sage grouse populations are limited by the availability and distribution of sagebrush.¹⁷ Sagebrush is an important year-round food as well as providing cover and shelter. Adequate sagebrush must surround strutting grounds for breeding to occur.^{7,29} Agricultural demands may pose a threat to sagebrush habitat.

Nest destruction is probably the most significant factor controlling sage grouse production.³

Severe weather conditions, unless snow completely covers the sagebrush, have little effect on the birds.²⁷ An exception may be cold, rainy weather during hatching, decreasing productivity.²

16.7 Regional Variation

The sage grouse occurs only in the extreme southeast portion of the province where adequate expanses of sagebrush are found.

ACKNOWLEDGMENTS

J. Clark (Alberta Fish and Wildlife, Lethbridge) reviewed this summary.

SELECTED REFERENCES

1. Batterson, W. M. and W. B. Morse. 1948. Oregon sage grouse. Ore. Fauna Sec. 1. Oregon Game Comm., Portland. 29 pp.
2. Carr, H. D. 1968. A literature review on effects of herbicides on sage grouse. Colorado Div. Game, Fish, and Parks Comm. Sep. Rept. No. 13. 16 pp.
3. Carr, H. D. and D. R. Carruthers. 1973. Status of sage grouse in Alberta in 1972. Alberta Fish and Wildlife Div., Calgary. 22 pp.
4. Crawford, J. E., Jr. 1960. The movements, productivity, and management of sage grouse in Clark and Fremont Counties, Idaho. M.S. Thesis. Univ. Idaho, Moscow. 85 pp.
5. Dalke, P. D., D. B. Pyrah, D. C. Stanton, J. E. Crawford and E. F. Schatterer. 1963. Ecology, productivity, and management of sage grouse in Idaho. J. Wildl. Manage. 27:811-841.
6. Dickinson, D. 1969. Sagebrush range study. Alberta Dept. Lands and Forests, Fish and Wildlife Division, Wildlife Investigations Rept. W-1-60.
7. Eng, R. L. and P. Schladweiler. 1972. Sage grouse winter movements and habitat use in central Montana. J. Wildl. Manage. 36:141-146.
8. Enyeart, G. 1956. Responses of sage grouse to grass reseeding in the Pine areas. Garfield County, Utah. M.S. Thesis. Utah State Univ., Logan. 55 pp.
9. Gill, R. B. 1966. A literature review on the sage grouse. Colo. Div. of Wildl., Spec. Rept. 6. 41 pp.
10. Girard, G. L. 1935. Life history, habits, and food of sage grouse, Centrocercus urophasianus. M.S. Thesis. Univ. Wyoming, Laramie. 153 pp.
11. Griner, L. A. 1939. A study of the sage grouse(Centrocercus urophasianus), with special reference to life history, habitat requirements, and numbers and distribution. M.S. Thesis. Utah State Agr. Coll. Logan. 83 pp.
12. Gudmundson, L. 1981. Provincial sage grouse population trend counts. Fish and Wildlife Division, Energy and Natural Resources. 14 pp.
13. Johnsgard, P. A. 1973. Grouse and quails of North America. Univ. of Nebraska Press, Lincoln. 553 pp.
14. Klebenow, D. A. and G. M. Gray. 1968. Food habits of juvenile sage grouse. J. Range Manage. 21:80-83.

15. Martin, N. S. 1970. Sagebrush control related to habitat and sage grouse occurrence. J. Wildl. Manage. 34:313-320.
16. Mitchell, G. J. 1959. Alberta's upland game bird resource. Dept. Lands and Forests, Alberta.
17. Patterson, R. L. 1952. Sage grouse hunting seasons. Wyo. Wildl. 16:10-13.
18. Patterson, R. L. 1952. The sage grouse in the Upper Green River basin of Wyoming. Ph.D. Thesis. Univ. of Michigan, Ann Arbor. 474 pp.
19. Peterson, J. G. 1970. The food habits and summer distribution of juvenile sage grouse in central Montana. J. Wildl. Manage. 34:147-155.
20. Rasmussen, D. I. and L. A. Griner. 1938. Life history and management studies of the sage grouse in Utah, with special reference to nesting and feeding habits. Trans. N. Am. Wildl. Conf. 3:852-864.
21. Rogers, G. E. 1964. Sage grouse investigations in Colorado. Colorado Game, Fish and Parks Dept., Tech. Publ. No. 16. 132 pp.
22. Schoener, T. J. and C. E. Braun. 1982. Spring habitat use by male and female sage grouse. Colo. State Univ. and Colo. Div. of Wildl., Fort Collins.
23. Scott, J. W. 1942. Mating behavior of the sage grouse. Auk. 59:477-498.
24. Stanton, D. C. 1956. A study of breeding and reproduction in a sage grouse population in southeastern Idaho. M.S. Thesis. Univ. Idaho, Moscow. 87 pp.
25. Trueblood, R. W. 1954. The effect of grass reseeding in sagebrush lands on sage grouse populations. M.S. Thesis. Utah State Univ., Logan. 77 pp.
26. Wallestad, R. O. 1971. Summer movements and habitat use by sage grouse broods in central Montana. J. Wildl. Manage. 35:129-136.
27. Wallestad, R. O. 1975. Life history and habitat requirements of sage grouse in central Montana. Mont. Dept. Fish and Game, Helena. 65 pp.
28. Wallestad, R. O. and D. Pyrah. 1974. Movement and nesting of sage grouse hens in central Montana. J. Wildl. Manage. 38:630-633.
29. Wallestad, R. O., and P. Schladweiler. 1974. Breeding season movements and habitat selection of male sage grouse. J. Wildl. Manage. 38:634-637.



Figure 16: General Distribution of Habitat Quality for Sage Grouse in Alberta.

BREEDING DABBLER DUCKS



17. KEY HABITAT REQUIREMENTS FOR RESIDENT BREEDING DABBING DUCKS

Alberta's breeding dabbling ducks include the mallard (Anas platyrhynchos), pintail (Anas acuta), northern shoveler (Anas clypeata), blue-winged teal (Anas discors), green-winged teal (Anas crecca), gadwall (Anas strepera) and American widgeon (Anas americana).

17.1 General

Water is essential to waterfowl habitat. Areas supporting high densities of shallow wetlands in fertile prairie-parkland regions provide the most productive breeding habitat in Alberta, habitat that ranks among the best in North America. Within these areas, largely devoted to agriculture, the less intensive and more diversified farming practices are most likely to result in the retention of higher quality habitat.

Distribution, relative abundances, production areas and harvest regions are depicted in Figures 17a - 17c. Priority duck production habitat is listed in Table 2.

17.2 Cover

17.2.1 Vegetation

Nest site selection is influenced both by environmental factors and by interaction among the nesting ducks.

Nest cover is provided by sedges (Carex sp.), mixed prairie grasses, hardstem bulrush (Scirpus sp.), great bulrush (Scirpus acutus) and rushes (Juncus sp.); forbes such as whitetop (Erigeron sp.), nettles (Urtica sp.), thistles (Cirsium sp.), fireweed (Epilobium angustifolium) and lambsquarter (Chenopodium album), as well as shrubs such as buckbrush (Symphoricarpos occidentalis) and gooseberry (Ribes sp.).^{1,16,18,26}

Selection of nest sites is influenced by the degree of vegetation cover; in central Alberta ducks preferred vegetation in the 1.5 - 3.4 dm range.¹⁸

Residual vegetation from the previous year is used by early nesting species.^{28,32}

Pioneer vegetation, sparse forbs, forest and pure stands of tall reed grass (Calamagrostis sp.) were avoided as nesting areas in central Alberta.¹⁸

Hayfields, stubble fields, grain fields, pastures and roadsides also serve as nesting sites, especially for pintail.²⁰

Broods often use potholes ringed with emergent vegetation and surrounded by haylands or ungrazed woodland.²⁶

Broods show a strong tendency to stay in or near concealing vegetation such as sedges and bulrushes.⁹

17.2.2 Land Forms and Topography

Moraine, especially higher relief moraine, supports the greatest density of wetlands and the highest densities of breeding ducks.²³

In Saskatchewan, lacustrine and solonchic soil land systems were associated with a low density of wetlands, lack of permanency of wetlands due to shallow basins and intensive cultivation (lack of upland nesting cover), resulting in relatively poor areas for duck production.²³ Fluvial, alluvium, aeolian, saline, eroded drainage and meltwater channel land systems tended to support low wetland densities.

Islands, dikes, levees,³³ fields and roadside ditches serve as nesting sites.

17.2.3 Aquatic Forms

Interspersion of shallow marshes, open water marshes, and open water wetlands provide good breeding habitat.²³

The most preferred of all habitats by resident dabblers in Alberta are semi-permanent potholes of up to 1 m deep in summer.²⁶

Northern shoveler pairs whose core areas were small water areas (< 0.4 ha) made use of several ponds centered near the nest site.

Mallard broods favored potholes 0.04-0.2 ha at Lousana, Alberta² and 0.2-0.4 ha in Saskatchewan parklands.²⁹ Blue-winged teal broods often occupy ponds 0.04-0.2 ha in size²⁹; American wigeon prefer semi-permanent areas of 0.2-0.4 ha, though at Lousana, broods favored larger potholes (0.8-2.0 ha).²⁶ Stockponds are also used by blue-winged teal and mallard.²⁹

Broods frequently move from pond to pond.¹

During years of low precipitation ducks require some permanent water bodies which last into late summer for brood salvage.

Creeks and rivers may form key duck production habitat if they contain water into mid-summer, support good emergent growth and meander extensively or have many oxbow lakes within their flood plain.²³

17.2.4 Climate

Cold spring conditions delay nesting.²⁵

Emergent vegetation traps drifting snow, increasing the insulation which causes a thinner sheet of ice to form. These areas are first to melt in the spring and are therefore most attractive to early nesting pairs.¹⁴

The amount of growing season precipitation and spring runoff determines the number of potholes available for duck use.

Drought seriously affects upland cover, reducing growth and thus the protective nature of the vegetation. Danger to broods is also intensified by the need for more frequent and longer excursions overland in search of water.²⁵

Drought decreases the number of potholes available; ducks may be forced to fly further north to boreal habitat.¹⁰

Ducks exhibit a strong tendency to choose loafing sites on the leeward side when on an island.¹⁸

17.3 Food

17.3.1 Vegetation

Dabbling ducks are primarily herbivorous, with the exception of brooding hens and young ducklings.^{21,26}

Various aquatic plants, including duckweed (Lemna sp.), pondweeds (Potamogeton sp.), bulrushes, sedges and smartweed (Polygonum sp.) provide important food for both adult and juvenile ducks.^{16,26,27}

Dabbling ducks, particularly mallards and pintails, make extensive use of cereal grains in spring and fall.^{2,27}

Hens require a diet high in aquatic and terrestrial invertebrates to supply them with the increased protein demand during laying.²

The diet of the newly hatched duckling is dominated by pond surface invertebrates. As the duckling ages these are replaced by aquatic invertebrates and by three weeks the diet is largely herbivorous, with less than 10 per cent animal food.³¹

17.3.2 Land Forms and Topography

Wetlands developed on saline soils generally produce fewer ducks than fresh wetlands as they lack dense emergent growth and support fewer invertebrates.²³

17.3.3 Aquatic Forms

Stagnant sloughs, ponds and other wetland areas provide the various aquatic plants used by dabbling ducks.

Seasonal wetlands are important feeding sites in spring when invertebrate populations are high. As these basins dry out, the ducks shift to semi-permanent ponds and lakes and feed on adult aquatic insects.²³

Wetlands surrounded by undisturbed plant communities have higher production of invertebrate food than areas disturbed through agriculture.²³

17.4 Space

17.4.1 Territory/Home Range

The summer ranges of dabbling ducks vary in size according to each species, population density and habitat conditions; in Manitoba, the home range of mallards exceeded 280 ha though the core area was smaller, the territory of blue-winged teals averaged 7-100 ha.^{6,19} In Utah, the territory of the gadwall ranged from 14-36 ha.⁸ In Alberta, the home range of the northern shoveler averaged 30.4 ha; parts of most home ranges overlapped and some were completely within boundaries of others.²²

On islands, mallards nest in close proximity to geese (within 30m);¹⁸ pintails nest in association with terns.³³

In the fall, most dabbling ducks migrate to winter range mainly in the southern United States and Mexico, and return to northern regions in early spring.¹

17.4.2 Population Densities

Population densities of dabbling ducks are presented in Table 1.

Table 1
MEAN DENSITIES OF DABBLER DUCKS FOR VARIOUS ECOLOGICAL ZONES IN
ALBERTA (MEASURED IN BREEDING PAIRS PER KM²)¹

	<u>Short Grass Prairie</u>	<u>Mixed Grass Prairie</u>	<u>Aspen Parkland</u>	<u>Boreal Forest</u>
Mallard	--	32.2	70.7	2.8
Pintail	40.4	60.1	28.5-38.1	3.1
Northern Shoveler	3.6	17.4	18.8	very low
Blue-winged Teal	--	19.9	42.0	3.4
Green-winged Teal	2.1	4.9	13.0	5.4
Gadwall	3.9	10.9	17.6	very low
American Wigeon	6.5	8.0	19.2	3.4

17.5 Special Considerations

17.5.1 Size, Shape and Juxtaposition of Habitat Components

Dabbling duck breeding pair densities and species diversity are greatest on wetland areas with a high interspersion of land and water.¹⁵

Wetland complexes within 1.5-3.0 km of fields with undisturbed grass and legume cover provide good habitat for breeding pairs and broods.⁷

The broken pattern of emergent aquatic growth in parkland potholes is often used for brood cover.²¹ The extent of nesting is proportionate to the

quality and quantity of marsh vegetation occurring along the shoreline and in the shallow water.⁹

The degree of difference in height and density of vegetation at an interface is an important factor in nest site selection; there was a tendency for nesting ducks to situate within 3 m of plant association interfaces in the denser vegetation in central Alberta.¹⁸

The majority of dabbling nests are found 30.5 - 45.5m from the water's edge, with some nests as far as 135 m from the shoreline;^{9,16} nests close to water decrease the time for the ducks to reach the security of water.²⁶

Diversified land use patterns in the nesting area contribute to better duck production than areas under intensively tilled grain monocultures.⁵

17.5.2 Significance of Disturbance Phenomena

Managed Flooding/Drainage. Temporary drainage of an area, maintains a high level of marsh productivity; after flooding the area peaks in use by ducks because of the abundance of pioneer plants such as duckweed.⁴

The changes in marsh and aquatic vegetation create an increased interspersed of cover and water.^{11,13}

Drainage of marsh areas considerably reduces the invertebrate population for a few years until slow repopulation takes place.¹⁴

Progressive succession of an area from pioneer plants to more stable rooted aquatics such as pondweeds reduces the productivity for ducks.⁴

Drought decreases the number of potholes available; ducks may be forced to fly further north to boreal habitat.¹⁰

Agriculture. Land clearing can open up tree and shrub-ringed potholes, making them more accessible and attractive to ducks.¹²

The draining of wetlands for agricultural expansion destroys good waterfowl habitat.

Cereal grain fields provide a supplementary food supply particularly during early spring and fall.

Cultivation of steep morainal hillsides exposes them to wind and water erosion and speeds up the siltation and filling-in of nearby potholes.¹²

Road construction frequently creates new habitat for ducks by dividing large, deep potholes into two separate bodies of water; ditches created by roads collect runoff and are attractive to ducks.¹²

Nests are destroyed each year by agricultural activities such as cultivation, mowing, burning of stubble and pasture lands, fence building and road construction.¹²

Agriculture creates long narrow bands of cover which are easily searched by nest predators.²⁴

Human Disturbance. Human activity can affect the use of an area by waterfowl in all seasons,¹⁸ but particularly during the nesting season.

Hunting pressure may cause changes in the use of areas.

17.6 Limiting Factors

Ducks are limited in areas with excessive water fluctuations; insufficient nutrients in the water and soil for optimum plant growth; permeable soils which adversely affect the formation and permanency of wetlands; deep, open water with steep shorelines and poorly developed marsh areas; shallow potholes which dry up quickly in the spring.³

Considerable wind and wave action on large water bodies reduces shoreline productivity and cover, and is particularly hazardous to young ducklings.²⁹

Type and density of nesting cover will affect nesting suitability.

17.7 Regional Variations

Numerous ponds and potholes of the aspen parkland and prairie grassland regions provide optimum habitat for large populations of dabbling ducks. Gently to moderately-sloping ground moraine promotes the development of high densities of wetlands with well-developed marsh edges. Higher average precipitation in the parkland region results in a higher proportion of permanent wetlands as compared to the prairie region. The parkland provides a wide range of cover, water and nesting and brooding sites, and the clearing of tree cover in central Alberta has increased the use of potholes that were once unattractive to ducks.²⁷

Draining of wetlands for agricultural use in the grasslands and parkland regions has destroyed much prime duck habitat. Nest sites are often found in stubble, fallow and crop fields which leads to their destruction.

On average, ducks exist in lower numbers in the northern mixed wood, but in times of drought in southern regions, the north provides important stable habitat for ducks.

The large expanse of land in the boreal region with numerous lakes and streams make possible a substantial contribution to waterfowl habitat. Water forms do not undergo as wide fluctuations in water levels as in the more southern regions.

Slow, meandering streams, ox-bow lakes, deltas and beaver ponds which support emergent vegetation and aquatic food resources gain importance as duck

production habitat in the boreal region. Lakes surrounded by dense tree cover and lacking a shoreline discourage dabbler use.

ACKNOWLEDGMENTS

E. Hofman (Alberta Fish and Wildlife, Brooks), W. Wishart (Alberta Fish and Wildlife, Edmonton) and L. Shandruk (Can. Wild. Ser.) provided critical reviews and useful comments.

SELECTED REFERENCES

1. Bellrose, F.C. 1976. Ducks, geese and swans of North America. Stackpole Books.
2. Bellrose, F.C. and J.B. Low. 1978. Advances in waterfowl management research. Wildl. Soc. Bull. 6:63-72.
3. Canadian Wildlife Service. 1967. Outline of the Canadian land capability classification for wildlife. Environment Canada. Ottawa.
4. Di Angelo, S. 1953. Aquatic plant succession at certain waterfowl flooding projects in Michigan. M.S. Thesis, University of Michigan, Ann Arbor, Michigan. 112p.
5. Duebbert, H.F. and J.T. Lokemoen. 1976. Ducks nesting in fields of undisturbed grass-legume cover. J. Wildl. Manage. 40(1): 39-49.
6. Dzubin, A. 1955. Some evidence of home range in waterfowl. Trans. 20th N. Am. Wildl. Conf. :278-298.
7. Dzubin, A. and J.B. Gollop. 1972. Aspects of mallard breeding ecology in Canadian parkland and grassland. Population Ecology of Migratory Birds: A Symposium. U.S. Dept. of Int., Wildl. Res. Rep. 2. 278 p.
8. Gates, J.M. 1962. Breeding biology of the gadwall in northern Utah. Wilson Bull. 74:43-67.
9. Griffith, R. 1948. Improving waterfowl habitat. Trans. N. Am. Wildl. Conf. 13:609-618.
10. Hansen, H.A. and D.E. McKnight. 1964. Emigration of drought-displaced ducks to the Arctic. Trans. 29th North Am. Wildl. and Nat. Resour. Conf. :119-126.
11. Harris, S.W. 1957. Ecological effects of drawdown operations for the purpose of improving waterfowl habitat. Ph.D. Thesis. University of Minnesota, St. Paul, Minnesota. 209p.
12. Johnsgard, P.A. 1956. Effects of water fluctuation and vegetation change on bird populations, particularly waterfowl. Ecology. Vol. 37, No. 4.
13. Kadlec, J.A. 1956. The effects of water level management on two central Wisconsin marshes. M.S. Thesis. University of Michigan. Ann Arbor, Michigan. 84p.
14. Kadlec, J.A. 1962. Effects of a drawdown on waterfowl impoundment. Ecology, Vol. 43, No. 2.

15. Kaminski, R.M. and H.H. Prince. 1981. Dabbling duck and aquatic macroinvertebrates responses to manipulated wetland habitat. J. Wildl. Manage. 45(1):1-15.
16. Keith, L.B. 1961. A study of waterfowl ecology on small impoundments in S.E. Alberta. Wildl. Soc. Wildl. Mono. No. 6. pp. 1-88.
17. Labisky, R.F. 1957. Relation of day harvesting to duck nesting under a refuge - permittee system. J. Wildl. Manage. 21:194-200.
18. Long, R.J. 1970. A study of nest-site selection by island-nesting Anatids in central Alberta. M.Sc. Thesis. University of Alberta. 123pp.
19. McHenry, M.G. 1971. Breeding and post-breeding movements of blue-winged teal (Anas discors) in southwestern Manitoba. Ph.D. Thesis. University of Oklahoma, Norman. 67 pp.
20. Milonski, M. 1958. The significance of farmland for waterfowl nesting and techniques for reducing losses due to agricultural practices. Trans. 23rd. North Am. Wildl. Conf.: 215-227.
21. Munro, D.A. 1967. The prairies and the ducks. Can. Geog. J. 75(1): 2-13.
22. Poston, H.J. 1966. Home range and breeding biology of shovelers (Anas clypeata) in the Alberta grassland. C.W.S. Ann. Rep. 25pp.
23. Schmidt, A.P. 1981. Wetland habitat inventory of the Kindersley (72-N) map area. Wildlife Technical Report 81-12. Wildlife Branch. Depart. Tour. and Renew. Res. 107pp.
24. Shandruk, L. (per. comm.) 1983. Biologist. Canadian Wildlife Service, Edmonton.
25. Smith, A.G. 1961. The 1961 waterfowl survey on the Lousana study area Alberta, Canada. Progress Rep. U.S. Dept. Int. Wildlife Research, Colorado. 33 pp.
26. Smith, A.G. 1971. Ecological factors affecting waterfowl production in the Alberta parklands. U.S. Dept. of the Interior. Fish and Wildl. Serv. Res. Publ. No. 98. 49 pp.
27. Smith, A.G., J.H. Stoudt and J.B. Gollop. 1964. Prairie potholes and marshes/northern watersheds and deltas. pp. 39-66 IN Waterfowl Tomorrow. U.S. Bur. Sport. Fish and Wildl. Wash. D.C.
28. Sowls, L.K. 1955. Prairie Ducks: A study of their behavior, ecology and management. The Stackpole Co., Harrisburg and the Wildl. Manage. Instit. Washington, D.C. 193 pp.

29. Stelfox, H.A. 1977. Waterfowl food ecology and habitat use in northeastern Saskatchewan. M.Sc. Thesis. University of Saskatchewan. 133 pp.
30. Stoudt, J.H. 1971. Ecological factors affecting waterfowl production in the Saskatchewan parklands. U.S. Fish and Wildl. Serv. Resourc. Publ. 99. 58 pp.
31. Sugden, L.G. 1973. Feeding ecology of pintail, gadwall, American widgeon and lesser scaup ducklings in southern Alberta. C.W.S. Report Series No. 24. 43 pp.
32. Townsend, G.H. 1966. A study of waterfowl nesting on the Saskatchewan River Delta. Can. Field Naturalist. 80(2): 74-88.
33. Vermeer, K. 1970. Some aspects of the nesting of ducks on islands in Lake Newell, Alberta. J. Wildl. Manage. 34:126-129.




Figure 17.a: Distribution and Relative Abundance of Waterfowl in Alberta, 1982.
 SOURCE: Adapted from Technical Discussion Paper for Waterfowl Management in Alberta. Unpublished draft, May/85.
 Fish and Wildlife Division, ENR. Edmonton. 120 pp.




Figure 17.b: Priority Duck Production Habitat in Alberta.

SOURCE: Adapted from Migratory Bird Habitat Priorities; Prairie Provinces. Unpublished Report, June 1979. Canadian Wildlife Service. Edmonton. 104 pp.

Table 2
PRIORITY DUCK PRODUCTION HABITAT IN ALBERTA
(Refer to Figure 17b)

AREA NO.	NAME	PRIORITY	VEGETATION ZONE	SIZE (hectares)
1.	Milk River Ridge	High	Grassland	
2.	Stirling-Etzikom-Cypress Hills	Medium	Grassland	312 000
3.	Innisfail-Strathmore-Keho Lake	Medium	Grassland	580 000
4.	Drumheller-Bassano-Brooks	Medium	Grassland	1 480 000
5.	Compeer-Acadia Valley-Berry Creek	Medium	Grassland	530 000
6.	Kneehills-Buffalo Lake	High	Aspen Parkland	720 000
7.	Beaverhill-Mundare	High	Aspen Parkland	285 200
8.	Viking Moraine	High	Aspen Parkland	224 000
9.	Forestburg-Battle River	High	Aspen Parkland	371 200
10.	Lloydminster-Ribstone	High	Aspen Parkland	276 400
11.	Peace River Parkland	High	Aspen Parkland	1 322 400
12.	Peace-Athabasca Delta	High	Boreal Forest	377 600
13.	Gordon Lake	High	Boreal Forest	11 380
14.	Jessie Lake	High	Boreal Forest	324
15.	Hay-Zama lakes	High	Boreal Forest	11 216
16.	Manawan Lake	High	Boreal Forest	742
17.	Smoky Lake	High	Boreal Forest	1 715
18.	Lesser Slave Lake	High	Boreal Forest	3 312
19.	Big Lake	Medium	Boreal Forest	1 715
20.	Lubicon Lake	Medium	Boreal Forest	3 584
21.	Loon Lake	Medium	Boreal Forest	1 460
22.	Cache Lake	Medium	Boreal Forest	312
23.	Carroll lakes	Medium	Boreal Forest	11 400
24.	Forsyth Lake	Medium	Boreal Forest	380
25.	Cold Lake (S. Bays)	Medium	Boreal Forest	2 160
26.	Little Buffalo Lake	Medium	Boreal Forest	464
27.	Chip Lake	Medium	Boreal Forest	7 347
28.	Utikuma Lake	Medium	Boreal Forest	28 416
29.	Missawawi Lake	Medium	Boreal Forest	2 560
30.	Therien lakes	Medium	Boreal Forest	2 246
31.	Flat Lake	Medium	Boreal Forest	3 072
32.	McClelland Lake	Medium	Boreal Forest	2 745
33.	Bison Lake	Low	Boreal Forest	3 226
34.	Lac la Biche	Low	Boreal Forest	24 166
35.	Bistcho Lake	Low	Boreal Forest	40 448
	Grassland Total			3 246 000
	Aspen Parkland Total			3 199 200
	Boreal Forest Total			540 815
	Grand Total			6 985 815

SOURCE: Adapted from: Migratory Bird Habitat Priorities, Prairie Provinces.
Unpublished Report, June, 1979. Habitat Management Section,
Canadian Wildlife Service, Edmonton. 104 pp.





Figure 17.c: Relative Total Duck Harvest Areas in Alberta, 1977-1980.
 SOURCE: Adapted from Technical Discussion Paper For Waterfowl Management in Alberta. Unpublished draft, May/85.
 Fish and Wildlife Division, ENR. Edmonton. 120 pp.

CANADA GEESE



18. KEY HABITAT REQUIREMENTS FOR BREEDING CANADA GEESE

18.1 General

Canada geese (*Branta canadensis*) are the most adaptable of the goose species and breed locally in all parts of Alberta. They will nest wherever they can find a territory near water with a secure nest site, be it in northern forests, mountain valleys, or across prairies and parklands³, though the most productive goose habitat is in the southeastern grassland zone.⁵ Good breeding habitat provides a nesting site, feeding and brooding sites with adequate food and cover, all being in close proximity to an adequate water area.

The major Canada geese production areas and priority goose production habitat in Alberta are depicted in Figure 18 and Table 3.

18.2 Cover

18.2.1 Vegetation

Nest cover types include cattail (*Typha latifolia*), bulrushes (*Scirpus* sp.), grasses, annuals, shrubs⁶ or some obstruction such as fallen trees.²

The type of cover is immaterial so long as the basic requirements for a nesting site (protection and isolation) are met.^{26,30}

Nesting sites which are virtually inaccessible to mammalian predators (i.e. islands) may alleviate the necessity for nest concealment by emergent cover.^{6,15}

Most nesting sites allow some visibility of the surrounding territory and easy access to water.¹⁰

Geese nesting in denser vegetation may be less vulnerable to competitive interactions with each other as this can lead to nest abandonment.⁷

Geese in southeastern Alberta selected artificial islands with a greater coverage of forbs and grasses, and islands with successful nests had denser vegetation than those with unsuccessful nests.¹¹ Tall or very dense growth, however, is not preferred.³⁰

In southeastern Michigan, the density of vegetation at the immediate nest site was significantly lower than the average vegetation density on these same islands.¹⁵

Many forms of artificial nesting sites are accepted by Canada geese: artificial islands,¹¹ various nesting tubs,⁴ cones, and platforms either in trees and poles, on the shore or in the water,² and on flax straw nesting bales.¹⁴

Nests are not elaborate. They are made from weeds, broken sticks, grass, moss, pine needles and other available materials; they are often a hollow scratched out of grasses and soil.²⁹

Geese covering their nests with dead vegetation and down when leaving, may minimize the loss of eggs due to predation.²⁰

Geese typically roost at night in marshes or open water in close proximity to grazing areas.^{17,31}

Vegetation on grazing areas may be used as resting/loafing cover.¹⁰

During the moult, geese may seek safety in heavy, emergent shoreline vegetation such as bulrushes and cattails.³⁰

18.2.2 Land Forms and Topography

Geese exhibit a preference for nesting on islands in rivers or lakes, though other sites such as isolated peninsulas, old heron and osprey nests, cliffs along rivers. Muskrat houses and beaver lodges are also used.^{10,2,24} In marshy areas, muskrat houses may be the predominant nest site.^{6,26}

Most nests on the Churchill River occurred on low profile islands.²³ Islands having more relief render nests less vulnerable to fluctuating water levels but facilitate nest vigilance due to decreased visibility.¹⁵

Islands within 46 m of the mainland or connected to the mainland at any time during the breeding season often attract predators.²

Smaller islands with a larger water-island interface tend to be preferred.^{11,29}

Sandbars, mudflats, a clear shoreline or open water may be used for resting and loafing.³⁰

Broods may return to nesting islands at night to roost.¹²

18.2.3 Aquatic Forms

Nesting pairs are frequently observed in association with large lakes or along streams.²¹ Irrigation reservoirs and stock dams also are used if adequate nest sites are present.

Area of permanent, open water is a factor affecting the use of wetlands; in southeastern Michigan, > 90 per cent of all nests located had ≥ 2 ha of open water.¹⁵

Most nest sites are within a short distance of open water, < 2 m in southeastern Manitoba.⁶

In southeastern Alberta, geese selected islands with deeper surrounding water and those which were further from shore, thus reducing potential predation. These islands had higher nesting success than other areas.¹¹

During moult, geese are rarely seen any distance from water, which is used for escape.¹⁴

18.2.4 Climate

Arrival of breeding geese in southern Alberta is related to the severity of winter and onset of spring snow melt.¹²

Extremely cold springs that delay the thawing of ice can dramatically decrease birds nesting in an area.²¹

Deep snow may force geese to abandon traditional nesting areas.¹²

Air temperature influences the rate of snow and ice melting, and thus may have an indirect as well as direct effect on the beginning of nesting.⁶

Shelter from prevailing winds may influence nest site selection.²²

During cold periods, geese remain at roosting sites for much of the time.¹⁷

Extremely hot dry weather during nesting and cool wet weather during hatching and rearing may contribute to population decreases.⁹

18.3 Food

18.3.1 Vegetation

Geese are essentially grazing birds,² utilizing pastures, fields, marshes and lakes as feeding sites.

They consume a wide variety of grasses, rushes, cattail, forbs and submergent aquatic growth, ingesting leaves, shoots, roots and seeds.^{3,6,19,30}

Geese also make use of grain crops such as wheat (Triticum sp.), barley (Hordeum sp.) and oats (Avena sativa) when available, during nesting, brooding and migration.^{13,17}

Incubating females choose feeding sites having an abundance of new plant growth.⁶

Brood areas have a plentiful supply of aquatic or land vegetation.¹⁰

Goslings may consume insect and insect larvae but this lessens with age and constitutes a small part of the overall diet.³⁰

18.3.2 Land Forms and Topography

Lake shores, mudflats, pastures and agricultural fields are used as feeding sites.

18.3.3 Aquatic Forms

Lakes, marshes, ponds and reservoirs provide aquatic feeding areas for geese.

Forage species, associated with aquatic areas include, spike rush (Eleocharis sp.), manna grass (Glyceria sp.), slough grass (Beckmannia sp.), cattail, arrowhead (Sagittaria sp.) and water plantain (Alisma sp.)^{16,19}

Lowered water levels may adversely affect these feeding areas due to promotion of plant succession.²³

18.3.4 Climate

Geese may not feed for several days during certain periods of cold weather, especially when accompanied by snow cover.¹⁷

Heavy snow which covers food resources, and cold temperatures which freeze roosting lakes in fall initiates southward migrations.¹³

18.3.5 Trace Elements

Geese consume small amounts of sand and gravel to assist the muscular gizzard in grinding hard foods.³⁰

18.4 Space

18.4.1 Territory/Home Range

The Canada goose has a breeding ground home range.⁶

The territory of the Canada goose appears to be a moving territory centered around the female.⁶ The size of the defended site varies with the density of the breeding population, age of the bird, the nature of the surrounding cover, available nesting space and the chronology of breeding.^{3,1} It is smallest in the prelaying period and reaches a peak at clutch completion.⁶

Territoriality may act to space geese in an even pattern, though the heterogeneity of the vegetation may modify the pattern.²⁹

On southeastern Alberta islands, territories located near the shore, where expanses of short grass and mud flats were present, were significantly larger than those located inland on the island where shrubs dominated.⁷

Larger territories may be less subjected to desertion as they may fulfil all the requirements of the nest site and also food and water.⁷

Both female and male actively protect nests.³

During nesting, geese generally have at least two feeding and two bathing sites.⁶

Close proximity between nesting and brooding areas is desirable.^{2,28}

Movements of family groups may be inversely affected by group size.¹⁷

Once broods establish in a rearing area, they rarely move to other brood grounds.¹⁰

18.4.2 Population Densities

Factors affecting nesting density include: territoriality, available habitat, nest site tenacity,⁶ age structure of the population²⁵ and vegetation density.

Nesting density is much higher on preferred sites (river and lake islands) than adjacent areas.²³

On artificially created islands in southern Alberta, an average of 1.6 nests/ha (0.2-7.1 nests/ha) were supported.¹¹

Small islands support a much greater nesting density than larger islands,^{10,11,29} possibly due to a larger water-land interface.

In southeastern Alberta, greater nesting success was recorded for geese nesting singly on small islands.¹¹

Large islands usually have more than one nesting pair.¹¹

When islands are in short supply, high densities occur which may result in high losses through desertion.⁷

Distance between nest sites affects nesting success. In southeastern Alberta, successful nests were found to be an average of 14.9 m from neighboring nests.⁷

18.5 Special Considerations

18.5.1 Size, Shape and Juxtaposition of Habitat Components

Good breeding area should have grazing area available to nesting birds and to paired breeders prior to nesting, nesting sites which are isolated, have firm foundations and afford good visibility, a brooding area adjacent to open water with a grazing and/or an aquatic feeding area, and a cover of emergent plants for use during the moult and for roosting.^{26,30}

Geese prefer to nest on smaller islands in relatively deep water.¹¹

The degree of isolation of the nesting area determines the necessity for cover.¹⁵

The length of shoreline is considered important to nesting geese.⁶

Large territories which include feeding sites and water area, as well as nest site, are less likely to be deserted.⁷

18.5.2 Significance of Disturbance Phenomena

Flooding/Drainage. Flooding may result in nest destruction on a local basis.⁶

Potential hydroelectric developments in northern Alberta could pose a threat to boreal forest production habitat by flooding nesting islands and shorelines.⁵

Low water levels may also adversely affect the number of birds nesting.²¹

Human Disturbance. Canada geese will tolerate some human habitation when selecting nesting wetlands.¹³

Increased industrialization and recreation in northern Alberta may be a potential threat to some breeding areas.⁵

Brood shifts may occur in exceptionally dry years or in response to harassment by people and domestic animals.²⁸

Agriculture. Much of the goose production habitat in southern Alberta is surrounded by agricultural pastures and fields. These areas are used as feeding sites during nesting and brooding.¹⁸

The development of artificial water bodies with suitable nest sites in southern Alberta may enhance goose utilization within the area.¹⁸

The drainage of wetland areas for agricultural expansion reduces waterfowl habitat.

18.6 Limiting Factors

Breeding Canada geese are limited by the availability of safe and isolated nesting sites, preferably on islands in lakes and rivers, with adequate feeding sites, cover and water area for nesting birds and broods.

The availability of nest sites is probably the main limiting factor on the prairies of Alberta as artificial nesting structures increase the breeding population.¹¹

The strong homing instinct of Canada geese to particular breeding grounds is a limiting factor on their distribution.¹

Low water levels during drought years may significantly decrease the number of nesting geese.²¹

Predation and nest desertion can cause substantial losses on a local basis.⁶

People's intolerance of crop depredation may also be a factor influencing the upper limits on Alberta's nesting flocks.¹⁸

18.7 Regional Variation

The major goose production habitat in Alberta occurs in the grassland zone and is designated as the Brooks-Lethbridge-Medicine Hat region.⁵ The Stettler-Coronation-Drumheller region, in the southern part of the parkland vegetation zone, and the Peace River-parklands are the next most important goose production areas in the province.

Nesting concentrations of large Canada geese in southern Alberta have been located along major rivers and on islands in large, permanent water bodies.⁸ Artificial water bodies and nesting sites are also used. Agricultural fields, which surround most of the goose nesting sites in southern Alberta, provide feeding areas. In the northern part of the province, most geese appear to nest on islands in major river systems.⁸

ACKNOWLEDGMENTS

E. Hofman (Alberta Fish and Wildlife, Brooks), W. Wishart (Alberta Fish and Wildlife, Edmonton), and L. Shandruk (Can. Wildl. Ser., Edmonton) provided critical reviews and useful comments.

SELECTED REFERENCES

1. Balham, R.W. 1954. The behavior of the Canada goose (Branta canadensis) in Manitoba. Ph.D. Thesis. Univeristy of Missouri. 22pp.
2. Ball, I.J., E.L. Bowhay, and C.F. Yocom. 1981. Ecology and management of Western Canada goose in Washington. Wash. Dept. Game Biol. Bull. No. 17. 68pp.
3. Bellrose, F.C. 1976. Ducks, geese, and swans of North America. Stackpole Books, Harrisburg, Penn. 543 pp.
4. Brakhage, G.KI. 1965. Biology and behavior of the tub-nesting Canada geese. J. Wildl. Manage. 29:751-771.
5. Canadian Wildlife Serivce. 1979. Migratory bird habitat priorities; prairie provinces. Habitat Management Section, Western and Northern Region, Can. Wildl. Ser. Rep. 104pp.
6. Cooper, J.A. 1978. The history and breeding biology of the Canada geese of Marshy Point, Manitoba. Wildl. Monogr. 61. 87pp.
7. Ewaschuk, E. and D.A. Boag. 1972. Factors affecting hatching success of densely nesting Canada geese. J. Wildl. Manage. 36:1097-1106.
8. Ewaschuk, E. and D.J. Neave. 1970. The status of the large Canada goose in Alberta. Alberta Fish and Wildlife Division, Edmonton. 46 pp.
9. Froggatt, K. 1980. Canada goose brood survey of southern Alberta. Alberta Energy and Natural Resources, Fish and Wildlife Division, Wildlife Branch. 57pp.
10. Geis, M. 1956. Productivity of Canada geese in the Flathead Valley, Montana. J. Wildl. Manage. 20:409-420.
11. Giroux, J. 1981. Use of artificial islands by nesting waterfowl in southeastern Alberta. J. Wildl. Manage. 45:669-679.
12. Glasgow, W.M. 1977. Brood mixing behavior and population dynamics of Canada Geese at Dowling Lake, Alberta. M.Sc. Thesis. University of Alberta. 149 pp.
13. Grieb, J.R. 1970. The shortgrass prairie Canada goose population. Wildl. Monogr. 22. 49 pp.
14. Hofman, D.E. (per comm.) 1983. Wildlife Biologist - Bird Game Unit. Brooks Wildlife Centre.
15. Kaminski, R.M. and H.H. Prince. 1977. Nesting habitat of Canada geese in southeastern Michigan.

16. Kelland, C.D. 1967. Local factors affecting waterfowl use of the Hay-Zama Lakes complex, with particular reference to migrating geese. Annual Job Report 1066-67. CWS-52-67. Canadian Wildlife Service.
17. Koerner, J.W., T.A. Bookhout, and K.E. Bednarik. 1974. Movements of Canada geese color-marked near southwestern Lake Erie. J. Wildl. Manage. 38:275-289.
18. Krohn, W.B. and E.G. Bizeau. 1980. The Rocky Mountain population of the western Canada goose: its distribution, habitats, and management. U.S. Dept. Int., Fish and Wildl. Ser. Spec. Sci. Rep., Wildlife No. 229. 93 pp.
19. Macaulay, A.J. Hay-Zama Lakes goose project. Can. Wildl. Ser. Job Completion Report. Project No. 82-4-5-078.
20. MacInnes, C.D. and R.K. Misra. 1972. Predation on Canada geese nests at McConnel River, Northwest Territories. J. Wildl. Manage. 36:414-422.
21. Malechi, R.A., F.D. Caswell, R.A. Bishop, K.M. Babcock, M.M. Gillespie. 1981. A breeding-ground survey of CPP Canada geese in northern Manitoba. J. Wildl. Manage. 45:46-53.
22. McCabe, T.R. 1979. Productivity and nesting habitat of great basin Canada geese, Umatilla, Oregon. Pages 117-129 in R.L. Jarvis and J.C. Bartonek, eds. Management and biology of Pacific Flyway geese. Oregon State University Book Store, Inc., Corvallis. 346 pp.
23. Raveling, D.G. 1977. Canada geese of the Churchill River basin in North-central Manitoba. J. Wildl. Manage. 41:35-47.
24. Salt, W.R. and J.R. Salt. 1976. The birds of Alberta. Hurtig Publishers, Edmonton. 498 pp.
25. Shandruk, L. (per comm.) 1983. Biologist. Canadian Wildlife Service, Edmonton.
26. Steel, P.E., P.D. Dalke and E.G. Bizeau. 1957. Canada goose production at Gray's Lake, Idaho, 1949-1951. J. Wildl. Manage. 21:38-41.
27. Surrendi, D.C. 1970. The mortality, behavior, and homing of transplanted juvenile Canada geese. J. Wildl. Manage. 34:719-733.
28. Szymczak, M.R. 1975. Canada geese restoration along the foothills of Colorado. Colo. Div. Wildl. Migratory Birds Investigations, Project W-8-8-R. Tech. Publ. No. 31. 64 pp.
29. Vemeer, K. 1970. A study of Canada geese, Branta canadensis, nesting on islands in southeastern Alberta. Can. J. Zool. 48:235-240.
30. Wormer, J.V. 1968. The World of the Canada Goose. J.B. Lippincott Company, N.Y. 192 pp.


31. Zicus, M.C. 1981. Flock behavior and vulnerability to hunting of Canada geese nesting at Crex Meadows, Wisconsin. J. Wildl. Manage. 45:830-841.

Table 3
PRIORITY GOOSE PRODUCITON HABITAT IN ALBERTA
(refer to Figure 18)

AREA NO.	NAME	PRIORITY	VEGETATION ZONE	SIZE (hectares)
1.	Brooks-Lethbridge-Medicine Hat	High	Grasslands	1 452 000
2.	Milk River Ridge	High	Grasslands	344 000
3.	Red Deer River	High	Grasslands	38 800
4.	South Saskatchewan River	High	Grasslands	92 000
5.	Bow River	High	Grasslands	39 080
6.	Oldman River	High	Grasslands	19 000
7.	Stettler-Coronation-Drumheller	Medium	Aspen Parkland	808 000
8.	Cooking Lake Moraine	Medium	Aspen Parkland	72 000
9.	Peace River Parkland	Medium	Aspen Parkland	1 322 500
10.	Wapiti River	Medium	Boreal Forest	4 800
11.	Little Smoky River	Medium	Boreal Forest	3 400
12.	Smoky River	medium	Boreal Forest	1 500
13.	Peace River	Medium	Boreal Forest	154 000
14.	Athabasca River	Medium	Boreal Forest	111 200
15.	Chinchaga River	Medium	Boreal Forest	1 640
16.	Hay River	Medium	Boreal Forest	3 440
17.	Stebing-Forsyth-Manatokan lakes	Low	Boreal Forest	74 000
Grassland Total				1 984 880
Aspen Parkland Total				2 202 400
Boreal Forest Total				353 980
Grand Total				4 541 260

SOURCE: Adapted from: Migratory Bird Habitat Priorities, Prairie Provinces.
Unpublished Report, June, 1979. Habitat Management Section,
Canadian Wildlife Service, Edmonton. 104pp.

MOULTING, STAGING, AND MIGRANT WATERFOWL



19. KEY HABITAT REQUIREMENTS FOR MOULTING, STAGING, AND MIGRANT WATERFOWL

19.1 General

Alberta serves as a moulting, stopover, and staging area for vast numbers of migrating waterfowl primarily from the Pacific and Central flyways, thus having national and international significance as a migratory corridor. Larger marshes and lakes with stable water levels, providing abundant food, adequate cover, isolation, and resting sites are preferred areas for staging and moulting. A greater variety of water bodies are associated with stopover sites, their use being less predictable. The magnitude of the migratory population and seasonal phenology, as well as habitat suitability, influence the use of these areas.

The distribution of priority moulting and staging areas for ducks and geese in Alberta are given on the accompanying maps (Maps 19.a - 19.d) and Tables 4 - 7.

19.2 Cover

19.2.1 Vegetation

On spring-staging areas at the Peace-Athabasca Delta edge, types preferred by dabblers were mud flats, meadow, immature fen and emergent vegetation; divers preferred emergent, tall shrub, meadow and low shrub edges, and geese preferred mud flats and immature fen.⁹

On fall-staging areas, dabblers preferred mud flats, immature fen and emergent shorelines; divers preferred emergent areas of restricted-drainage lakes and open, deep streams; geese preferred emergents and open drainage immature fen, and mud flats.⁹

Geese and swans do not prefer lakes that are heavily forested.¹²

Shoreline resting areas were found to be close to dense expanses of taller emergents such as hard-stem bulrush (Scirpus sp.) and cattail (Typha latifolia) in southeastern Manitoba.⁵

Emergent vegetation is used as roosting cover by dabblers and geese. 16,20

Marginal-emergent shorelines of open-drainage lakes and marshes appear to be the preferred moulting areas for dabblers;⁹ when disturbed they favor the zone of emergent vegetation.^{16,22}

Diving ducks, during the moult, may use marshy inlets for security though they seldom move onto land.¹⁶

19.2.2 Land Forms and Topography

Shoreline resting areas include mud or gravel points, mud bars, or the shores of small bays, inlets, and islands.⁵

When on water, birds may use landforms such as islands as visual obstructions to potential danger.¹¹

19.2.3 Aquatic Forms

Large lakes serve primarily as staging, moulting and migration areas.

Shallow lakes and sloughs with stable water levels provide good stopover habitat for dabbling ducks; larger and deeper lakes are suitable stopover habitat for diving ducks.^{15,16,22}

During the moult, diving ducks and geese use water as an escape medium.¹⁶

As lakes freeze-up, a greater number of waterfowl use rivers.²²

19.2.4 Climate

Length of stay at stopover and staging areas may depend on local weather conditions.

Advance of spring migration progresses fairly slowly in response to northern climates^{3,22} and early use of water areas generally follows the sequence of spring break-up.²²

Fall migration is more rapid, with many birds waiting until freeze-up or low temperatures before suddenly moving south.³

Dry weather may decrease the areas available as stopover sites.

19.3 Food

19.3.1 Vegetation

Dabbling ducks, geese, and swans feed on a variety of emergent, and submergent plants including pondweeds (Potamogeton sp.), duckweed (Lemna sp.), bulrushes, sedges (Carex sp.) and smartweed (Polygonum sp.).^{1, 2,7,14,17}

Diving ducks feed more frequently on submerged plants (pondweeds), and animal foods are more important to diving ducks than most dabblers.^{1,2,7,14}

Geese also utilize a variety of pasture grasses and grains.^{2,13}

Moulting and staging birds may be attracted to sites with an extensive food resource.^{8,10,18}

Movement of males and nonbreeders to moulting areas may be an adaption to reduce competition for food on breeding areas.¹⁸

Although large concentrations of different species of birds may use a given moulting or staging area, competition for food can be reduced by differences between species in foods eaten, feeding habitat and feeding behavior.⁸

When wing moult is complete, mallards, pintails and geese may feed extensively in grain stubble fields, utilizing wheat, barley, oats and corn.^{5,13,3}

19.3.2 Land Forms and Topography

Mud flats, lake shores, marsh shores, pastures and fields provide potential foraging areas.

19.3.3 Aquatic Forms

Lakes, marshes, sloughs and rivers are aquatic feeding sites.

During spring migrations, freshly thawed lakes with their abundance of food are preferred over rivers.²²

Waterfowl tend to feed on the shore and along the shore during pre-moult and post-moult, and in open water during moulting where they are less accessible to predation.¹⁶

Aquatic feeding may be concentrated where local abundance of food occurs.⁴

Animal matter in the diets of divers and dabblers include aquatic insects, amphipods and gastropods.^{1,2,7,14,17}

Mergansers consume mainly fish.²³

19.3.4 Climate

The progression of spring break-up and fall freeze-up influences the use of aquatic feeding areas.²²

Weather significantly influences distance flown to feeding areas. During periods of low temperature, precipitation or cloudy conditions with moderate to heavy winds, ducks feed in close proximity to the staging water body;⁵ geese tend to be more tolerant to weather conditions and continue to feed at preferred sites (e.g. grain fields).⁵

When snow covers waste grain in harvested fields to the extent that it makes for difficulty in the feeding activity of mallards and pintails, migrations may occur.³

19.3.5 Trace Elements

Grit particles are usually consumed to assist the grinding of food particles in the gizzard.¹⁶

19.4 Space

19.4.1 Territory/Home Range

Alberta's wetlands act as a corridor for waterfowl migrating to northern breeding grounds and southern wintering areas.³

Large bodies of water attract migrating waterfowl, influencing their flight pattern.^{22,24}

Birds may move short distances, or hundreds of miles to moulting and staging areas.^{3,16,24}

The distance waterfowl move between stopover and staging areas is variable with species, available habitat, and weather conditions as well as condition of the bird.^{3,22}

Stopover and staging areas may serve as extensions of the wintering areas; abundant food supplies may hold large populations at northern latitudes during mild weather.⁸

19.4.2 Population Densities

Moulting populations include post-breeding and non-breeding resident birds, immigrants from surrounding lakes and marshes, and immigrants from considerable distances.⁹

Moulting densities on the Peace-Athabasca Delta were 0-74.0/ha for dabblers and 0-32.0/ha for divers.⁹

Population densities for spring and fall-staging waterfowl are influenced by the availability of specific habitat types, the magnitude of the continental waterfowl population, the nesting success of the northern breeding population and the seasonal phenology.⁹

Spacial distribution of ducks, geese and swans is variable depending on the water body, its size and depth, food resources and weather conditions, often resulting in concentrations on sections of some lakes.¹²

The density of spring-staging waterfowl in the Peace-Athabasca Delta area ranged from 0 to 140 ducks/km of sampled shoreline in 1971 (means = 1.8 for divers, 17.5 for dabblers).⁹

On sampled areas of the Peace-Athabasca Delta for 1971, the mean densities of fall-staging waterfowl ranged from 1.2 to 709.6/km of shoreline for dabblers, 0 to 65.2/km for geese and swans, and 0.07 to 0.49/ha for divers.⁹

19.5 Special Considerations

19.5.1 Size, Shape and Juxtaposition of Habitat Components

Large bodies of water or water body complexes with abundant food resources attract migrant, moulting and staging waterfowl.

The extent of shoreline development and the diversity of shoreline vegetation types, as well as water depth, will influence the types and numbers of waterfowl accommodated.¹⁰

Local topography and juxtaposition to other water bodies are features which may attract waterfowl.⁹

Moulting and staging areas may or may not be synonymous.

19.5.2 Significance of Disturbance Phenomena

Agriculture. Intensified agriculture and drainage are depleting staging and stopover habitat in the parkland.⁶

A decline in natural wetlands along migration corridors results in a greater concentration of waterfowl on many stopover areas. Concentrating waterfowl may result in a more rapid depletion of food resources.⁸

Human Disturbance. Hunting pressure influences use of feeding sites and water areas.⁵

Increased recreational use, hydroelectric developments and increased industrialization may threaten staging habitat in northern Alberta.⁶

Floods/Water Levels. Flood conditions can have a marked effect upon waterfowl use. Traditional feeding or loafing areas may be temporarily lost, causing earlier than usual departure.²¹

Low fall water levels may provide extensive, mudflat-type shoreline, especially on larger lakes; this may attract a large, fall-staging population.¹⁰

19.6 Limiting Factors

Staging and moulting areas are limited by the distribution of larger water bodies with stable water levels, adequate shoreline types, food supply and cover.^{11,16,18}

Available dry edge may be a limiting factor of staging areas.⁹

Food resource and number of birds using it influences length of stay.⁸

The timing of weather conditions, spring thaw and fall freeze-up also limits the use of staging and stopover areas.²²

Agricultural, recreational, and industrial developments may decrease habitat available for moulting, staging and migrant waterfowl.⁶

19.7 Regional Variations

Staging habitat for ducks is concentrated mainly in the central regions of both the grassland and the aspen parkland zones of the province.⁶ These areas serve as staging sites for birds produced locally and for those from northern areas. Sites scattered throughout the boreal forest zone are also important, especially for waterfowl moving down from the Arctic. The majority of known duck moulting areas are located in the boreal forest vegetation zone, though several lakes in the parklands are also important. Boreal forest areas may provide isolation and abundant food resources required for moulting.

The major goose staging habitat is located in the aspen parkland and grassland zones, the southern sections of these zones being utilized the longest by staging Canada geese as water areas remain open the longest here.⁶ Arctic migrants such as the lesser Canada geese, snow geese, Ross' geese, and white-fronted geese also make partial use of these areas for fall-staging. In the boreal forest vegetation zone, the Peace-Athabasca Delta and the Hay-Zama lakes are important staging areas for arctic migrants.^{6,19}

ACKNOWLEDGMENTS

E. Hofman (Alberta Fish and Wildlife, Brooks), W. Wishart (Alberta Fish and Wildlife, Edmonton), and L. Shandruk (Can. Wildl. Ser., Edmonton) reviewed the summary and provided useful comments.

SELECTED REFERENCES

1. Anderson, H.G. 1959. Food habits of migratory ducks. Illinois Nat. Hist. Surv. Bull. Vol. 27, Art. 4.
2. Bartonek, J.C. and J.J. Hickey. 1969. Food habits of canvasback, redheads, and lesser scaup in Manitoba. Condor 71:280-290.
3. Bellrose, F.C. 1976. Ducks, geese and swans of North America. Stackpole Books. 543 pp.
4. Bergman, R.D. 1973. Use of southern boreal lakes by postbreeding canvasbacks and redheads. J. Wildl. Manage. 37:160-170.
5. Bossenmaier, E.F. and W.H. Marshall, 1958. Field-feeding by waterfowl in southeastern Manitoba. Wildl. Monogr. 1. 32 pp.
6. Canadian Wildlife Service. 1979. Migratory bird habitat priorities; prairie provinces. Habitat Management Section, Western and Northern Region, Can. Wildl. Ser. Rep. 104 pp.
7. Cottam, C. 1939. Food habits of North American diving ducks. United States Dept. Agr. Tech. Bull. No. 643. 138 pp.
8. Fredrickson, L.H. and R.D. Drobney. 1979. Habitat utilization by postbreeding waterfowl. pages 119-131 In: T.A. Bookhout, ed., Waterfowl and Wetlands, an integrated review. La Crosse Printing Co. Inc., Washington. 152 pp.
9. Hennan, E. 1972. Peace-Athabasca delta. Ducks Unlimited (Canada).
10. Hennan, E. and K.R. Ambrock. 1977. A summary of waterfowl investigations in the Peace-Athabasca delta, 1971-1976. Can. Wildl. Ser. Rep. prepared for Parks Canada. 20pp.
11. Hochbaum, H.A. 1944. The canvasback on a prairie marsh. North Am. Wildl. Inst., Washington, D.C. 201 pp.
12. Kemper, J.B. 1976. Waterfowl and migratory birds. Appendix Seven In: Cooking Lake Area Study. Vol. IV Ecology. Planning Division Alberta Environment. 129 pp.
13. Krohn, W.B. and E.G. Rizeua. 1980. The Rocky Mountain population of the western Canada goose: its distribution, habitats, and management. U.S. Dept. Int., Fish and Wildl. Ser., Spec. Sci. Rep. Wildl. No. 229. 93 pp.
14. Lewis, H.F. 1958. Food plants for waterfowl. Hunting and fishing in Canada.

15. Nieman, D.J. and H.J. Dirschl. 1971. Waterfowl populations on the Peace-Athabasca delta, 1969 and 1970. Unpubl. Rep., Can. Wildl. Ser., Saskatoon. 30 pp.
16. Oring, L.W. 1964. Behavior and ecology of certain ducks during the postbreeding period. J. Wildl. Manage. 28:223-233.
17. Rogers, J.P. and L.J. Korshgen. 1966. Foods of lesser scaups on breeding, migration, and wintering areas. J. Wildl. Manage. 30:258-264.
18. Salomonsen, F. 1968. The moult migration. Wildfowl 19:5-24.
19. Smith, R.H., F. Dufrensne, and H.A. Hansen. 1964. Northern watersheds and deltas, pages 51-66 In: J.P. Linduska and A.L. Nelson, eds. Waterfowl Tomorrow. U.S. Dept. Int., Gov. Printing Office. 770 pp.
20. Sowls, L.K. 1955. Prairie duck: a study of their behavior, ecology, and management. Stackpole Books, Harrisburg, Pa. & Wildl. Manage. Inst., Washington, D.C. 193 pp.
21. Sugden, L.G. 1963. Biology, distribution and productivity of Arctic geese migrating through northern Alberta. Can. Wildl. Ser. Proj. No. 01-1-1. 9 pp.
22. Syncrude Canada Ltd. 1973. Migratory waterfowl and the Syncrude tar sands lease. Env. Res. Monogr. 1973-3. 99 pp.
23. Timken, R.L. and B.M. Anderson. 1969. Food habits of common mergansers in the north central United States. J. Wildl. Manage. 33:87-91.
24. Weller, M.W. 1964. Distribution and migration of the redhead. J. Wildl. Manage. 28:65-103.

Table 4
PRIORITY DUCK STAGING HABITAT IN ALBERTA
(refer to Figure 19a)

AREA NO .	NAME	PRIORITY	VEGETATION ZONE	SIZE (hectares)
1.	Murray Lake	High	Grassland	1 690
2.	Frank Lake	High	Grassland	1 200
3.	Pakowki Lake	High	Grassland	13 184
4.	Louisiana Lakes	High	Grassland	1 490
5.	Many Island Lake	Medium	Grassland	8 960
6.	McGregor Reservoir	Medium	Grassland	5 043
7.	Stirling Lake	Medium	Grassland	102
8.	Keho Lake	Medium	Grassland	1 843
9.	Bassano Reservoir	Medium	Grassland	640
10.	Stobart Lake	Medium	Grassland	496
11.	Crow Indian Lake	Medium	Grassland	1 000
12.	Red Deer River	Medium	Grassland	38 800
13.	South Saskatchewan River	Medium	Grassland	92 000
14.	Oldman River	Medium	Grassland	19 000
15.	Bow River	Medium	Grassland	39 080
16.	Travers Reservoir	Low	Grassland	2 227
17.	Lake Newell	Low	Grassland	5 990
18.	Beaverhill Lake	High	Aspen Parkland	13 670
19.	Whitford Lake	High	Aspen Parkland	1 820
20.	Big Hay Lake	High	Aspen Parkland	1 075
21.	Bittern Lake	High	Aspen Parkland	2 714
22.	Demay Lake	High	Aspen Parkland	717
23.	Dusty Lake	High	Aspen Parkland	384
24.	Driedmeat Lake	High	Aspen Parkland	1 075
25.	Buffalo Lake	High	Aspen Parkland	10 354
26.	Cooking Lake	High	Aspen Parkland	3 456
27.	Bear Lake	High	Aspen Parkland	3 098
28.	Saskatoon Lake	High	Aspen Parkland	589
29.	Cardinal Lake	High	Aspen Parkland	4 378
30.	Erskine Lake	High	Aspen Parkland	1 500
31.	Bens-Watt lakes	Medium	Aspen Parkland	256
32.	Kenilworth Lake	Medium	Aspen Parkland	658
33.	Lac La Glace	Medium	Aspen Parkland	896
34.	Winagami Lake	Medium	Aspen Parkland	4 248
35.	Kimiwan Lake	Medium	Aspen Parkland	3 854
36.	Lac Maglorie	Medium	Aspen Parkland	818

Continued next page ...

Table 4
Continued ...

AREA NO.	NAME	PRIORITY	VEGETATION ZONE	SIZE (hectares)
37.	Gordon Lake	High	Boreal Forest	11 392
38.	Peace-Athabasca Delta	High	Boreal Forest	377 600
39.	Manawan Lake	High	Boreal Forest	742
40.	Hay-Zama lakes	High	Boreal Forest	11 216
41.	Utikuma Lake	Medium	Boreal Forest	28 416
42.	Flat Lake	Medium	Boreal Forest	3 072
43.	Lac La Biche	Medium	Boreal Forest	24 166
44.	Lesser Slave Lake	Medium	Boreal Forest	45 312
45.	Chip Lake	Medium	Boreal Forest	7 347
46.	Missawawi Lake	Medium	Boreal Forest	2 560
47.	Smoky Lake	Medium	Boreal Forest	540
48.	Big Lake	Medium	Boreal Forest	1 715
49.	McClelland Lake	Low	Boreal Forest	2 745
Grassland Total				232 745
Aspen Parkland Total				55 560
Boreal Forest Total				516 823
Grand Total				805 128

SOURCE: Adapted from: Migratory Bird Habitat Priorities, Prairie Provinces.
Unpublished Report, June, 1979. Habitat Management Section,
Canadian Wildlife Service. Edmonton. 104 pp.

Table 5
PRIORITY DUCK MOULTING HABITAT IN ALBERTA
(refer to Figure 19b)

AREA NO.	NAME	PRIORITY	VEGETATION ZONE	SIZE (hectares)
1.	Murray Lake	Medium	Grassland	1 690
2.	Beaverhill Lake	High	Aspen Parkland	14 800
3.	Kenilworth Lake	Medium	Aspen Parkland	666
4.	Bittern Lake	Medium	Aspen Parkland	2 714
5.	Buffalo Lake	Medium	Aspen Parkland	10 354
6.	Peace-Athabasca Delta	High	Boreal Forest	377 600
7.	Gordon Lake	High	Boreal Forest	11 392
8.	Jessie Lake	High	Boreal Forest	296
9.	Utikima Lake	Medium	Boreal Forest	28 416
10.	Hay-Zama lakes	Medium	Boreal Forest	11 216
11.	Loon Lake	Medium	Boreal Forest	1 472
12.	Lubicon Lake	Medium	Boreal Forest	3 506
13.	Muskwa Lake	Medium	Boreal Forest	4 813
14.	McClelland Lake	Low	Boreal Forest	2 745
Grassland Total				1 690
Aspen Parkland Total				28 534
Boreal Forest Total				441 456
Grand Total				471 680

SOURCE: Adapted from: Migratory Bird Habitat Priorities; Prairie Provinces. Unpublished Report, June, 1979. Habitat Management Section, Canadian Wildlife Service. Edmonton. 104 pp.

Table 6
PRIORITY GOOSE STAGING HABITAT IN ALBERTA
(refer to Figure 19c)

AREA NO.	NAME	PRIORITY	VEGETATION ZONE	SIZE (hectares)
1.	Berry Creek Reservoir	High	Grassland	589
2.	McGregor Reservoir	High	Grassland	5 043
3.	Murray Lake	High	Grassland	1 690
4.	Lake Newell	High	Grassland	5 990
5.	Louisiana Lakes	High	Grassland	1 490
6.	Plover Lake	High	Grassland	256
7.	Birkshire Reservoir	High	Grassland	240
8.	Milk River Ridge Reservoir	High	Grassland	1 797
9.	Crow Indian Lake	Medium	Grassland	1 000
10.	Deadhorse Lake	Medium	Grassland	1 229
11.	Frank lake	Medium	Grassland	1 997
12.	Grassy Lake	Medium	Grassland	230
13.	Keho Lake	Medium	Grassland	1 843
14.	Pakowki Lake	Medium	Grassland	13 076
15.	Scots Reservoir	Medium	Grassland	205
16.	St. Mary's Reservoir	Medium	Grassland	4 608
17.	Stobart Lake	Medium	Grassland	496
18.	Taber Lake	Medium	Grassland	461
19.	Travers Reservoir	Medium	Grassland	2 227
20.	Verdigris Lake	Medium	Grassland	691
21.	Colemen Lake	Medium	Grassland	1 070
22.	Many Island Lake	Medium	Grassland	8 960
23.	South Saskatchewan River	Low	Grassland	92 000
24.	Bow River	Low	Grassland	39 080
25.	Oldman River	Low	Grassland	19 000
26.	Red Deer River	Low	Grassland	38 800
27.	BeaverHill Lake	High	Aspen Parkland	13 670
28.	Buffalo Lake	High	Aspen Parkland	10 354
29.	Sullivan Lake	High	Aspen Parkland	10 854
30.	Cardinal Lake	High	Aspen Parkland	4 378
31.	Bear Lake	High	Aspen Parkland	3 098
32.	Dowling Lake	High	Aspen Parkland	2 918
33.	Grassy Island Lake	High	Aspen Parkland	1 101
34.	Kirkpatrick lake	High	Aspen Parkland	1 638
35.	Kenilworth Lake	Medium	Aspen Parkland	438
36.	Sounding Lake	Medium	Aspen Parkland	3 789
37.	Lac La Glace	Medium	Aspen Parkland	896
38.	Whitford Lake	Medium	Aspen Parkland	1 920
39.	Antelope Lake	Medium	Aspen Parkland	282
40.	Eagle Lake	Medium	Aspen Parkland	1 306

Continued next page ...

Table 6
Continued ...

AREA NO.	NAME	PRIORITY	VEGETATION ZONE	SIZE (hectares)
41.	Gough Lake	Medium	Aspen Parkland	4 864
42.	Shooting Lake	Medium	Aspen Parkland	794
43.	Birch Lake	Medium	Aspen Parkland	2 473
44.	Winagami Lake	Medium	Aspen Parkland	4 248
45.	Kimiwan Lake	Medium	Aspen Parkland	3 854
46.	Handhills Lake	Medium	Aspen Parkland	803
47.	Contracosta Lake	Medium	Aspen Parkland	695
48.	Marion Lake	Low	Aspen Parkland	1 971
49.	Peace-Athabasca Delta	High	Boreal Forest	377 600
50.	Hay-Zama lakes	High	Boreal Forest	11 216
51.	Bison Lake	High	Boreal Forest	3 226
52.	Bistcho Lake	Medium	Boreal Forest	40 448
Grassland Total				244 068
Aspen Parkland Total				76 344
Boreal Forest Total				432 490
Grand Total				752 902

SOURCE: Adapted from: Migratory Bird Habitat Priorities, Prairie Provinces.
Unpublished Report, June, 1979. Habitat Management Section, Canadian
Wildlife Service. Edmonton. 104pp.




Figure 19.d: Priority Goose Moulting Habitat in Alberta.

SOURCE: Adapted from Migratory Bird Habitat Priorities: Prairie Provinces. Unpublished Report, June 1979. Canadian Wildlife Service. Edmonton. 104 pp.

Table 7
PRIORITY GOOSE MOULTING HABITAT IN ALBERTA
(refer to Figure 19d)

AREA NO.	NAME	PRIORITY	VEGETATION ZONE	SIZE (hectares)
1.	Knight Ranch Reservoir	High	Grassland	188
2.	Ross Lake	High	Grassland	235
Other major goose production lakes have resident moulting birds.				
TOTAL				423

SOURCE: Adapted from: Migratory Bird Habitat Priorities, Prairie Provinces.
Unpublished Report, June, 1979. Habitat Management Section,
Canadian Wildlife Service. Edmonton. 104 pp.

APPENDIX 1
COMMON AND SCIENTIFIC NAMES OF MAMMALS REFERRED TO IN TEXT

COMMON NAME	SCIENTIFIC NAME
bighorn sheep	<u>Ovis canadensis</u>
elk	<u>Cervus elaphus</u>
moose	<u>Alces alces</u>
mountain goat	<u>Oreamnos americanus</u>
mule deer	<u>Odocoileus hemionus</u>
pronghorn antelope	<u>Antilocapra americana</u>
white-tailed deer	<u>Odocoileus virginianus</u>
woodland/mountain caribou	<u>Rangifer tarandus caribou</u>
beaver	<u>Castor canadensis</u>
grizzly bear	<u>Ursus arctos</u>
ground squirrel	<u>Spermophilus sp.</u>
marten	<u>Martes americana</u>
meadow vole	<u>Microtus pennsylvanicus</u>
mink	<u>Mustela vison</u>
muskrat	<u>Ondatra zibethicus</u>
river otter	<u>Lutra canadensis</u>
red squirrel	<u>Tamiasciurus hudsonicus</u>
red-backed vole	<u>Clethrionomys gapperi</u>
snowshoe hare	<u>Lepus americanus</u>
weasel	<u>Mustela sp.</u>

APPENDIX 2
COMMON AND SCIENTIFIC NAMES OF BIRDS REFERRED TO IN TEXT

COMMON NAMES	SCIENTIFIC NAMES
<hr/>	
Waterfowl	
American wigeon	<u>Mareca americana</u>
blue-winged teal	<u>Anas discors</u>
Canada goose	<u>Branta canadensis</u>
gadwall	<u>Anas streperus</u>
green-winged teal	<u>Anas crecca</u>
mallard	<u>Anas platyrhynchos</u>
northern shoveler	<u>Spatula clypeata</u>
pintail	<u>Anas acuta</u>
Upland Game Birds	
ring-necked pheasant	<u>Phasianus colchicus</u>
sage grouse	<u>Centrocercus urophasianus urophasianus</u>
sharp-tailed grouse	<u>Pedioecetes phasianellus</u>

APPENDIX 3
COMMON AND SCIENTIFIC NAMES OF PLANTS REFERRED TO IN TEXT
(ALBERTA SPECIES TAKEN FROM E.H. MOSS, FLORA OF ALBERTA)

COMMON NAMES

SCIENTIFIC NAMES

Trees/Shrubs

alder	<u>Alnus</u> sp.
alpine fir	<u>Abies lasiocarpa</u>
alpine larch	<u>Larix lyallii</u>
aspen	<u>Populus tremuloides</u>
balsam poplar	<u>Populus balsamifera</u>
bearberry	<u>Arctostaphylos uva-ursi</u>
beaked hazelnut	<u>Corylus cornuta</u>
birch	<u>Betula</u> sp.
bitterbrush	<u>Purshia tridentata</u>
black spruce	<u>Picea mariana</u>
blueberry	<u>Vaccinium</u>
bog birch	<u>Betula pumila</u>
buckbrush	<u>Symphoricarpos occidentalis</u>
buffaloberry	<u>Shepherdia canadensis</u>
choke cherry	<u>Prunus virginiana</u>
clematis	<u>Clematis</u> sp.
cottonwood	<u>Populus balsamifera</u>
creeping juniper	<u>Juniperus horizontalis</u>
currant	<u>Ribes</u> sp.
Douglas fir	<u>Pseudotsuga menziesii</u>
dwarf birch	<u>Betula glandulosa</u>
Engelmann spruce	<u>Picea engelmannii</u>
fir	<u>Abies</u> sp.
gooseberry	<u>Ribes</u> sp.
hawthorn	<u>Crataegus</u> sp.
high-bush cranberry	<u>Viburnum trilobum</u>
honeysuckle	<u>Lonicera</u> sp.
huckleberry	<u>Vaccinium membranaceum</u>
jackpine	<u>Pinus banksiana</u>
juniper	<u>Juniperus</u> sp.

APPENDIX 3
Continued ...

COMMON NAMES

SCIENTIFIC NAMES

Trees/Shrubs

lodgepole pine	<u>Pinus contorta</u>
low-bush cranberry	<u>Viburnum edule</u>
mountain ash	<u>Sorbus aucuparia</u>
mountain maple	<u>Acer glabrum</u>
paper birch	<u>Betula papyrifera</u>
pin cherry	<u>Prunus pensylvanica</u>
pine	<u>Pinus sp.</u>
rabbit brush	<u>Chrysothamnos nauseosus</u>
raspberry	<u>Ribes sp.</u>
red osier dogwood	<u>Cornus stolonifera</u>
rose	<u>Rosa sp.</u>
sage	<u>Artemisia sp.</u>
sagebrush	<u>Artemisia cana</u>
saskatoon	<u>Amelanchier alnifolia</u>
silverberry	<u>Elaeagnus commutata</u>
silver buffaloberry	<u>Shepherdia argentea</u>
silver sagebrush	<u>Artemisia cana</u>
snowberry	<u>Symphoricarpos albus</u>
spruce	<u>Picea sp.</u>
subalpine fir	<u>Abies lasiocarpa</u>
tamarack	<u>Larix laricina</u>
tea bush	<u>Ceanothus velutinus</u>
water birch	<u>Betula occidentalis</u>
wild gooseberry	<u>Ribes sp.</u>
willow	<u>Salix sp.</u>
white spruce	<u>Picea glauca</u>

APPENDIX 3
Continued ...

COMMON NAMES

SCIENTIFIC NAMES

Graminoids

barley	<u>Hordeum</u> sp.
bluegrass	<u>Poa</u> sp.
brome	<u>Bromus</u> sp.
bulrush	<u>Scirpus</u> sp.
cattail	<u>Typha latifolia</u>
corn	<u>Zea mays</u>
crested wheatgrass	<u>Agropyron cristatum</u>
fall rye	<u>Secale cereale</u>
fescue	<u>Festuca</u> sp.
flax	<u>Linum usitatissimum</u>
great bulrush	<u>Scirpus acutus</u>
hard-stem bulrush	<u>Scirpus</u> sp.
June grass	<u>Koeleria cristata</u>
manna grass	<u>Glyceria</u> sp.
melic grass	<u>Melica</u> sp.
oats	<u>Avena sativa</u>
reed grass	<u>Calamagrostis</u> sp.
Sandberg's bluegrass	<u>Poa secunda</u>
sedge	<u>Carex</u> sp.
slough grass	<u>Beckmannia</u> sp.
spike rush	<u>Eleocharis</u> sp.
timothy	<u>Phleum</u> sp.
western wheatgrass	<u>Agropyron smithii</u>
wheat	<u>Triticum</u> sp.
wheatgrass	<u>Agropyron</u> sp.
winter wheat	<u>Triticum</u>

Forbs

alfalfa	<u>Medicago sativa</u>
arrowhead	<u>Sagittaria</u> sp.
butterfly weed	<u>Gaura coccinea</u>

APPENDIX 3
Continued ...

COMMON NAMES

SCIENTIFIC NAMES

Forbs (cont'd)

clover	<u>Trifolium</u> sp.
Colorado rubber-plant	<u>Hymenoxys richardsonii</u>
cow-parsnip	<u>Heracleum lanatum</u>
dandelion	<u>Taraxacum</u> sp.
doorweed	<u>Polygonum aviculare</u>
duckweed	<u>Lemna</u> sp.
fireweed	<u>Epilobium angustifolium</u>
glacier lily	<u>Erythronium grandiflorum</u>
golden aster	<u>Chrysopsis villosa</u>
hedysarum	<u>Hedysarum</u> sp.
horsetail	<u>Equisetum arvense</u>
lambsquarters	<u>Chenopodium album</u>
nettle	<u>Urtica</u> sp.
pasture sage	<u>Artemisia frigida</u>
pond-weed	<u>Potamogeton</u> sp.
smartweed	<u>Polygonum</u> sp.
sow thistle	<u>Sonchus</u> sp.
thistle	<u>Cirsium</u> sp.
waterlily	<u>Nuphar</u> sp.
water plantain	<u>Alisma</u> sp.
wild strawberry	<u>Fragaria virginiana</u>
wild tomato	<u>Solanum triflorum</u>
white top	<u>Erigeron</u> sp.
wooly yarrow	<u>Achillea millefolium</u>
yellow goat's beard	<u>Tragopogon dubius</u>
yellow pond lily	<u>Nuphar variegatum</u>

This material is provided under educational reproduction permissions included in Alberta Environment and Sustainable Resource Development's Copyright and Disclosure Statement, see terms at <http://www.environment.alberta.ca/copyright.html>. This Statement requires the following identification:

"The source of the materials is Alberta Environment and Sustainable Resource Development <http://www.environment.gov.ab.ca/>. The use of these materials by the end user is done without any affiliation with or endorsement by the Government of Alberta. Reliance upon the end user's use of these materials is at the risk of the end user.