

UNIVERSITY OF TORONTO LIBRARY

National Library
Collective Development
Canada
Ottawa

Bibliothèque nationale
Collection collective
Ottawa

NOTICE

The quality of this microfiche is highly dependent upon the quality of the original thesis submitted for microfilming. Every effort has been made to provide the highest quality reproduction available.

La qualité de cette microfiche dépend grandement de la qualité de la thèse soumise au microfilmage. Nous avons tout fait pour assurer une qualité supérieure de reproduction.

The quality of the reproduction of certain pages may be less than that of the original if the original pages were of poor quality or if the original pages were damaged.

La qualité de cette microfiche dépend grandement de la qualité de la thèse soumise au microfilmage. Nous avons tout fait pour assurer une qualité supérieure de reproduction.

AVIS

La qualité de l'impression de certaines pages peut laisser à désirer, surtout si les pages originales ont été d'une mauvaise qualité ou si les pages originales ont été endommagées.

The quality of the reproduction of certain pages may be less than that of the original if the original pages were of poor quality or if the original pages were damaged.

This document is already the subject of a microfiche edition. It is not possible to produce a new edition of this document.

This document is already the subject of a microfiche edition. It is not possible to produce a new edition of this document.

UNIVERSITY OF TORONTO LIBRARY

UNIVERSITY OF TORONTO LIBRARY



National Library
of Canada

Bibliothèque nationale
du Canada

Canadian Theses Division / Division des thèses canadiennes

Ottawa, Canada
K1A 0N4

67263

PERMISSION TO MICROFILM / AUTORISATION DE MICROFILMER

Please print or type / Écrire en lettres majuscules

Full Name of Author / Nom complet de l'auteur

LIPAL I. NAVARATNAM

DES ANIMALES

Date of Birth - Date de naissance

12th APRIL 1931

Home Address / Adresse

101 GARDNER ST.

University / Université

UNIVERSITY OF TORONTO

Department / Département

DEPARTMENT OF ZOOLOGY

Faculty / Faculté

FACULTY OF SCIENCE

Permit to / Autorisation de

copy / copier

is hereby / est par la présente

granted / accordée

to the / à la

Library / Bibliothèque

L'autorisation est par la présente accordée à la BIBLIOTHÈQUE NATIONALE DU CANADA de microfilmer cet ouvrage en prêt à ses utilisateurs sur microfilm.

This copy / Cette copie

67263-101 Gardner St.

THE UNIVERSITY OF ALBERTA

DEPARTMENT OF EDUCATION AND CURRICIAL INQUIRY

by

DR. J. H. BIRCHALL

A BULLETIN

OF THE UNIVERSITY OF ALBERTA

EDUCATIONAL INQUIRY

NUMBER 1

1961



THE UNIVERSITY OF ALBERTA

RELEASE FORM

NAME OF AUTHOR: H. H. B. DISNEY, JR.

TITLE OF THESIS: A COMPACTNESS THEOREM

DEGREE FOR WHICH THESIS WAS PRESENTED

Doctor of Philosophy

YEAR THIS DEGREE GRANTED

1964

Permission is hereby granted by THE UNIVERSITY OF ALBERTA
Library to reproduce in whole or in part this thesis and to loan
it to other libraries for their use. This permission is granted
on the condition that the borrower of this thesis will pay the
cost of the reproduction.

THE UNIVERSITY OF ALBERTA
LIBRARY

12
13

THE UNIVERSITY OF ALBERTA

FACULTY OF GRADUATE STUDIES AND RESEARCH

The undersigned certify that they have read, and recommend to the Faculty of Graduate Studies and Research, for acceptance, a thesis entitled m - n COMPACTNESS and CARDINAL INVARIANTS submitted by U.N.B. DISSANAYAKE in partial fulfillment of the requirements for the degree of Doctor of Philosophy in Mathematics.

Stephen Wilf
(Supervisor)

J.G. Tumawir

R.L. McKinney

K. ...

Bernard Lusig

J. T. ...
(External Examiner)

ABSTRACT

In the first part of the thesis we investigate the productivity of weak m - n compactness which covers three main concepts: $H(i)$, feebly compact and weakly Lindelöf. In the later part, we consider the possible improvements and generalizations of Arhangel'skii's famous theorem (1970),

$$|X| \leq 2^{L(X) \chi(X)}$$

for T_2 spaces, where $L(X)$ is the Lindelöf degree and $\chi(X)$ is the character of the space X .

Chapter I contains a brief history of cardinal invariants in topology and definitions of generalized topological notions with necessary basic theorems.

Chapter II is devoted to a short survey of m - n compact topological spaces and considers the problem of preservation of tightness in large products of m - n compact spaces. In this respect we show that tightness behaves similar to character in products of locally Lindelöf, T_3 spaces and hence in products of Lindelöf, T_3 P-spaces.

In Chapter III, we introduce strong m - n filters and study products of two spaces and then large products (and weak topological). By doing this we obtain sufficient conditions for factor spaces and the productivity of m - n compactness.

Chapter IV is mainly devoted to a study of two interesting cardinal functions, the Lindelöf degree, $L(X)$ and the weak Lindelöf number, $wL(X)$. Here we introduce a new cardinal function, the almost Lindelöf degree $aL(X)$, which satisfies,

$$wL(X) \leq aL(X) \leq L(X)$$

for any space X and prove that

$$|X| \leq 2^{aL(X) \chi(X)}$$

for T_2 spaces. Next, we obtain results similar to the above involving $wL(X)$ and study relations between $L(X)$ and $wL(X)$ by introducing a cardinal function, the w -Lindelöf number

INTRODUCTION

In 1929, P.S. Alexandroff and P.S. Uryshon introduced H-closed spaces as a natural generalization of compact, T_2 spaces. In 1941,

C. Chevalley and O. Frink proved that an arbitrary product of

H-closed spaces is H-closed. Z. Frolik, W. Comfort, A. Hager and

S. Negrepointis have studied another generalization of compact (Lindelöf)

spaces, namely weakly Lindelöf spaces. In 1972, M. Ulmer [60] has

extended some of the results of the above mathematicians, by proving

that the weakly Lindelöf property in a product space is determined

by finite subproducts. The above two variations of compactness are

special cases of the general concept weak κ -manifolds and part

of our interest is to find sufficient conditions for a product of

(and individual factors) to be weakly Lindelöf.

arbitrary products

1979, P.S. Alexandroff and

of the product space

where \mathcal{C}_i is finite and $\mathcal{C}_i \in \mathcal{C}_i$

for $i = 1, 2, \dots, n$

where \mathcal{C}_i is finite and $\mathcal{C}_i \in \mathcal{C}_i$

for $i = 1, 2, \dots, n$

||

where \mathcal{C}_i is finite and $\mathcal{C}_i \in \mathcal{C}_i$

After this theorem, cardinal functions in topology played an active role. I. Juhász has written a definitive text (see [34]) in this field, including all the new cardinal functions and new results. As a natural generalization of $l(X)$, the weak Lindelöf number $wL(X)$ ($< l(X)$) was first introduced in [4] and there it is proved that, if X is a T_1 space, then

$$|X| \leq 2^{wL(X) \chi(X)} \quad (B)$$

The remaining part of our interest is to study the cardinal function $wL(X)$ in contrast to $l(X)$ and to consider possible improvements of (A) and to extend (B) to the class of T_2 spaces.

Chapter I is devoted to a brief study of basic topological notions, namely regular cardinal, cardinal functions $l(X)$, $l(X)$ and $wL(X)$ and their properties.

In Chapter II, our main aim is to estimate tightness in products of m compact spaces in the factor spaces. We begin this by giving a short survey of the theory of m -compact spaces and proving a product theorem which partially extends Hodel's theorem 4.2 of [53]. That product theorem and the resulting theory of tightness are described as follows:

Γ is a sum of n power compact spaces if and only if every

$$T \subseteq \Gamma \text{ is } \kappa\text{-tight for } \kappa < \aleph_1$$

(1) Let $X = \prod\{X_i : i \in I\}$ be a quasi n -paracompact space. Let n be a regular cardinal and suppose each X_i is m - n compact, κ - n -discrete and T_2 . Then X is m - n compact.

Let $X = \prod\{X_i : i \in I\}$. Let $\theta_I(X) = \sup\{\theta(X_i) : i \in I\}$ where $\theta(X)$ denotes the tightness of X . We define X to be a $GIC(n, i)$ space for $i = 1, 2, 3, 4$ if X is a locally m - n compact T_2 space and T_i if and only if X is a $GIC(n, i)$ space.

(2) $\theta(X) = \theta_I(X)$.

[1] p. 57 [2] p. 111

Let $X = \prod\{X_i : i \in I\}$ and let n be a regular cardinal. Suppose each X_i is m - n compact and κ - n -discrete. Then X is m - n compact if and only if the set

Let $X = \prod\{X_i : i \in I\}$ and let n be a regular cardinal. Suppose each X_i is m - n compact and κ - n -discrete. Then the set $\{X_i : i \in I\}$ is m - n compact if and only if the set $\{X_i : i \in I\}$ is m - n compact and κ - n -discrete.

[3] p. 111 [4] p. 111 [5] p. 111 [6] p. 111

It is clear that the presence of κ - n -discreteness in the hypotheses of [1] is essential for the conclusion to hold.

[7] p. 111 [8] p. 111 [9] p. 111 [10] p. 111

In Chapter IV, we introduce a new cardinal function, the almost Lindelöf degree, $aL(X)$, which agrees with $L(X)$ on T_3 spaces but which is often smaller than $L(X)$ on T_2 spaces, and prove that

$$(1) \quad |X| < 2^{aL(X) \cdot \chi(X)}$$

for T_2 spaces.

Next, we introduce, via local π -bases, a new class of spaces, the Π -normal space. This class contains T_4 spaces and

$$|X| < 2^{w(X) \cdot \chi(X)}$$

when X is a Π -normal space. This extends (B).

Since compact subsets behave nicely in T_2 spaces, it is of interest to obtain upper bounds for the number of compact subsets of X . In this direction, we show that for a T_2 space X ,

$$(2) \quad |K(X)| \leq 2^{aL(X) \cdot \chi(X)} \cdot \min \left(2^{w(X) \cdot \chi(X)}, 2^{L(X) \cdot \chi(X)} \right)$$

(see also [9]).

Finally, we introduce a new cardinal function, the π -character, $\pi\chi(X)$, which is defined for regular spaces,

$$\pi\chi(X) = L(X) \cdot \chi(X)$$

Each chapter is divided into sections and subsections. The main results in each subsection are labelled by letters A, B, C, D, E, etc. When we quote a result in the same subsection we use only the letter and the number of the result. For example, (A) refers to result A in the first subsection of the chapter.

subsection, then we use only the subsection number followed by the corresponding letter. We follow this pattern by indicating chapter number, section number, subsection number and the corresponding letter, as necessary.

The cardinal functions which are not defined here can be found in Juhász text [31] and for topological concepts which are used without any special introduction we refer the reader to the text of Willard [20].

ACKNOWLEDGEMENTS

I wish to thank my thesis advisor, Dr. S. Willard for his wisely restrained guidance and for the sizeable amount of time that he has invested in reading the thesis.

In addition, I would like to thank the University of (Mathematics Department) for the financial support.

TABLE OF CONTENTS

CHAPTER		PAGE
I	PRELIMINARIES	1
	1. Cardinal Arithmetic	1
	2. Cardinal Invariants	6
	3. m and n Filters	12
	4. Generalized Filters	15
II	COMPACT SPACES	19
	1. Some Properties of Compact Spaces	19
	2. Products of Lindelöf spaces	24
	3. Locally Compact Spaces	36
	4. Tightness in Compact Spaces	37
III	m and n COMPACT SPACES	39
	1. Some Properties of m and n Spaces	39
	2. Some Characterizations	46
	3. Products of m and n Spaces	51
	4. m and n Spaces and Filters	57
IV	ALGEBRAIC COMPACT SPACES	61
	1. The Algebra of Lindelöf Spaces	61
	2. The Vietoris Topology	67
	3. m and n Spaces	71

	PAGE
REFERENCES AND REMARKS	89
BIBLIOGRAPHY	91
APPENDIX	96

CHAPTER I. PRELIMINARIES

1. Cardinal Arithmetic

This section is devoted to a brief review of transfinite arithmetic and some special properties of infinite cardinals. For notation and terminology not explained here, see [1] or [2].

Simple rules. Let α, β, γ be cardinals.

- (i) $(\alpha^\beta)^\gamma = \alpha^{\beta \cdot \gamma}$
- (ii) If $\kappa < \lambda$, then $\alpha^\kappa < \alpha^\lambda$, where $\alpha \geq 2$.
- (iii) $\alpha^{\beta + \gamma} = \alpha^\beta \cdot \alpha^\gamma$.

Let α, β, γ be cardinals and $\beta \neq 0$. Then $\alpha^\beta < \alpha^{\beta + \gamma}$ if and only if $\alpha < \alpha^\beta$.

This result follows from the fact that $\alpha^\beta < \alpha^{\beta + \gamma}$ if and only if

$$\alpha^\beta < \alpha^{\beta + \gamma}.$$

Let α, β, γ be cardinals and $\beta \neq 0$. Then $\alpha^\beta < \alpha^{\beta + \gamma}$ if and only if

$$\alpha^\beta < \alpha^{\beta + \gamma}.$$

Let α, β, γ be cardinals and $\beta \neq 0$. Then $\alpha^\beta < \alpha^{\beta + \gamma}$ if and only if

$$\alpha^\beta < \alpha^{\beta + \gamma}.$$

Let α, β, γ be cardinals and $\beta \neq 0$. Then $\alpha^\beta < \alpha^{\beta + \gamma}$ if and only if

$$\alpha^\beta < \alpha^{\beta + \gamma}.$$

C Definition. Let $(X, <)$ be a linearly ordered set. Then a subset A of X is said to be cofinal in X if for every $x \in X$, there exists $a \in A$ such that $x \leq a$.

Conventionally we identify an infinite cardinal α with the least ordinal ω_α with cardinality α . Thus we can define the cofinality of α as,

$$cf(\alpha) = \min \{ \beta \mid \exists \langle \alpha_i \mid i < \beta \rangle \text{ a sequence of ordinals } < \alpha_i < \alpha \text{ such that } \alpha = \sup \alpha_i \}$$

It is clear that $cf(\alpha) \leq \alpha$ and $cf(\alpha) < \alpha$ if and only if α is a limit ordinal.

It is not difficult to verify that

(i) $cf(\alpha) = cf(\alpha^+)$

(ii) $cf(\omega^\alpha) = cf(\alpha)$

(iii) $cf(\aleph_\alpha) = cf(\alpha)$

(iv) $cf(\aleph_\alpha^+) = \aleph_\alpha$

Definition. A cardinal number κ is said to be regular if and only if $cf(\kappa) = \kappa$ and it is singular if and only if $cf(\kappa) < \kappa$.

The cardinal \aleph_1 is the cofinality of \aleph_1 and hence \aleph_1 is regular. More generally, \aleph_α is regular if and only if α is a successor ordinal. For the proof see [1, p. 100].

All finite cardinals are regular, while \aleph_0 is singular.

It is not difficult to verify that

We note that, α^+ is always a regular cardinal for any cardinal α .

The cofinality of an infinite cardinal α can be regarded as the least cardinal β such that α can be decomposed into union of β sets, each of which has cardinality less than α . As an example

$$\aleph_1 = \bigcup_{i=0}^{\infty} \aleph_i \quad \text{thus } \text{cf}(\aleph_1) = \aleph_0$$

Next, we shall study special properties of cardinals

F. Definition A cardinal α is called **inaccessible** if it is regular and uncountable and only if

- (i) $\kappa < \aleph_\alpha$ and
- (ii) $\alpha < \aleph_\alpha$ for all $\alpha < \aleph_\alpha$
- (iii) α is regular

A cardinal which satisfies (i) and (ii) is called an **inaccessible cardinal** and a cardinal which satisfies (i), (ii) and (iii) is called a **strongly inaccessible cardinal**.

We shall prove that if α is a cardinal, then α is inaccessible if and only if α is a regular cardinal and $\alpha < \aleph_\alpha$ for all $\alpha < \aleph_\alpha$.

Proof: Let α be a cardinal. If α is inaccessible, then α is regular and uncountable. Let $\beta < \aleph_\alpha$. Then $\beta < \aleph_\beta$ for all $\beta < \aleph_\beta$.

Conversely, let α be a regular cardinal and $\alpha < \aleph_\alpha$ for all $\alpha < \aleph_\alpha$.

each i . Then we define

$$\sum\{k_j : j \in I\} = |\cup\{X_i : i \in I\}|,$$

and

$$\prod\{k_j : j \in I\} = |\prod\{X_i : i \in I\}|.$$

The following results will be useful in the sequel (see [30]):

(i) If λ is an infinite cardinal and $k_i > 0$ for each $i < \lambda$,

$$\sum\{k_i : i < \lambda\} = \sup\{k_i : i < \lambda\}$$

(ii) If λ is an infinite cardinal and $\{k_i : i < \lambda\}$ is a non-decreasing sequence of non-zero cardinals, then

$$\prod\{k_i : i < \lambda\} = (\sup\{k_i : i < \lambda\})^\lambda$$

In the sequel we shall introduce some notation that will be useful in the sequel.

Definition

Let λ be an infinite cardinal. Then λ^β denotes

$$\sum\{k_i : i < \lambda\}$$

if λ is an infinite cardinal. The cardinal λ^β is the

$$\text{card}(\lambda^\beta) = \lambda^\beta$$

To illustrate, note that

(i) If $\alpha > N_0$, then

$$\alpha^{\frac{N}{\alpha}} \approx \alpha \quad \text{and} \quad (\alpha)^{\frac{\alpha}{N}} \approx 2^{\frac{1}{N}}$$

(ii) If $\alpha < N_0$,

2. Cardinal Invariants

The development and use of cardinal invariants in topology has been significant. At early stages of topology, character $\chi(X)$, pseudocharacter $\psi(X)$ and weight $W(X)$ played an important role. After 1970, other cardinal invariants such as the Lindelöf degree $L(X)$, density $d(X)$ and spread $s(X)$ have taken an active role. As we shall see, many valuable results can be formulated in terms of cardinal invariants.

A. Basic Definitions. Let X be any topological space and let $a \in X$. Then an open neighbourhood base at a is a collection \mathcal{V}_a of open subsets of X such that if V is any open subset containing a , then there exists some $U \in \mathcal{V}_a$ such that $a \in U \subseteq V$.

If X is a T_1 -space, then a pseudobase at a is a collection \mathcal{B}_a of subsets of X such that

$$(a) \quad \rho(W \cdot W) = \rho(W)$$

(i) Let X be any topological space and let $x \in X$. Then we define $\chi(x, X) = \min\{|\mathcal{V}_x| : \mathcal{V}_x \text{ is an open neighbourhood base at } x\}$ and the character $\chi(X)$ of X as:

$$\chi(X) = \sup\{\chi(x, X) : x \in X\}$$

(ii) Let X be a T_1 -space and let $x \in X$. Then we define $\psi(x, X) = \min\{|\mathcal{B}_x| : \mathcal{B}_x \text{ is a pseudobase at } x\}$ and the pseudocharacter $\psi(X)$ of X as:

$$\psi(X) = \sup\{\psi(x, X) : x \in X\}.$$

(iii) Let X be any topological space. Then we define the weight $w(X)$ of X as,

$$w(X) = \min\{|B| : B \text{ is a base for } X\} + \aleph_0.$$

(iv) Let X be any topological space. Then we say that X is k -Lindelöf if and only if every open cover of X has a sub-cover of cardinality at most k . We define the Lindelöf degree $L(X)$ of X as,

$$L(X) = \min\{k : X \text{ is } k\text{-Lindelöf}\} + \aleph_0.$$

(v) Let X be any topological space. Then we define the density $d(X)$ of X as,

$$d(X) = \min\{|S| : S \text{ is dense in } X\} + \aleph_0.$$

(vi) Let X be any topological space. Then we define the character $\chi(X)$ of X as,

$$\chi(X) = \max\{\chi(x, X) : x \in X\} + \aleph_0.$$

Historical Facts We know that for any topological space X , $\psi(X) \leq \chi(X)$. In general, the gap between $\psi(X)$ and $\chi(X)$ is very large. In the early part of the twentieth century, P. Alexandroff and L. Urysohn, proved the following remarkable results:

(i) Theorem: Let X be a countably compact T_1 space. If every point of X has a countable local base, then $\psi(X) = \chi(X)$.

space $(\chi(X) = \aleph_0)$.

(ii) Theorem. Let X be a compact T_2 -space. Then $\chi(X) = \psi(X)$. We note that this result can be extended to locally compact T_2 -spaces. Thus the weight $w(X) \leq |X|$ for all locally compact T_2 -spaces.

Among all separation axioms, normality plays an important role. F.B. Jones (1937) has proved an interesting lemma concerning normal spaces. To state the lemma using cardinal invariants we require a definition.

Definition. Let X be any topological space. Then we define the closed spread $p(X)$ of X as:

$$p(X) = \sup\{|B| : B \text{ is a closed discrete subspace of } X\} \cdot \aleph_0$$

(iii) Lemma (Jones). If X is normal, then $p(X) < 2^{d(X)}$.

We apply this lemma to the Moore plane M . For this we note that $p(M) = \aleph_1$ and $d(M) = \aleph_0$. Thus M is a non-normal (completely regular) T_2 -space.

Actual development of cardinal invariants, has begun after 1970. This is mainly due to the famous theorem of A.V. Arhangel'skii which answered a long standing problem of P. Alexandroff and P. Urysohn, namely does every first countable, compact T_2 -space have cardinality at most \aleph_1 ? Arhangel'skii's answer takes the following form:

(iv) Theorem (Arhangel'skii). Let X be a T_2 -space. Then

$$|X| \leq \aleph_1 \cdot \psi(X)$$

In the latter part of 1970, the development of cardinal functions in topology has proceeded rapidly. As a result of this many new cardinal functions have been added. Among them we shall consider mainly, the following cardinal functions in addition to $\chi(X), \psi(Y), W(X), I(X), \rho(X)$ and $\pi(X)$. (see [34]).

- (a) tightness - $\theta(X)$, (traditional notation)
- (b) π character - $\pi\chi(X)$.
- (c) closed spread - $p(X)$.
- (d) cellularity - $c(X)$.

C. Examples. The following standard examples are encountered frequently in our work and it is very useful to know the values of some of the cardinal functions on these spaces:

- (a) the countable complement topology on any infinite set with cardinality κ .
- (b) the Alexandroff extension of \mathbb{R} .
- (c) the Menger plane.
- (d) the Alexandroff double circle.
- (e) the Ordinal space.
- (f) the Tychonoff plank.
- (g) Michael's line (also known as the Sorgenfrey line).
- (h) the Sorgenfrey line.
- (i) the Sorgenfrey plane.

D. Chart. We shall give the values of χ , ψ , α , π_X , d , w , p , L and s for the spaces (a) to (j) in a tabular form.

Let $\phi(X)$ be a cardinal function on a topological space X . Then $\phi^*(X)$ denotes, $\sup\{\phi(Y) : Y \subseteq X\}$. If $\phi = \phi^*$, then we say that ϕ is monotone.

We note that, χ , ψ , α , and w are monotone.

SPACES	$\chi(X)$	$\psi(X)$	$\alpha(X)$	$\pi_X(X)$	$d(X)$	$c(X)$	$w(X)$	$p(X)$	$L(X)$	$s(X)$
(a)	k	k	1	k	\aleph_1	\aleph_0	k	\aleph_0	\aleph_0	\aleph_0
(b)	c	\aleph_0	\aleph_0	\aleph_0	\aleph_0	\aleph_0	c	\aleph_0	\aleph_0	\aleph_0
(c)	\aleph_0	\aleph_0	\aleph_0	\aleph_0	\aleph_0	\aleph_0	\aleph_0	c	c	c
(d)	\aleph_0	\aleph_0	\aleph_0	1	c	c	c	\aleph_0	\aleph_0	c
(e)	c	c	c	\aleph_0	c	c	c	\aleph_0	\aleph_0	c
(f)	\aleph_0	\aleph_0	\aleph_0	\aleph_0	c	c	c	c	c	c
(g)	\aleph_0	\aleph_0	\aleph_0	\aleph_0	c	c	c	\aleph_0	c	c
(h)	\aleph_0	\aleph_0	\aleph_0	\aleph_0	\aleph_0	\aleph_0	c	\aleph_0	\aleph_0	\aleph_0
(i)	\aleph_0	\aleph_0	\aleph_0	\aleph_0	\aleph_0	\aleph_0	c	c	c	c
(j)	k	k	k	k	$\log k$	\aleph_0	k	\aleph_0	k	k

Next, we shall give an important result which we shall use frequently in our work.

Theorem 1.1 ([1]). Let X be a T_1 space. Then $|X| \leq d(X)^{w(X)}$.



F. Note. It is easy to construct an example via \mathbf{N} , to show that T_2 cannot be relaxed to T_1 in the hypothesis of the theorem E.

G. Example. Let $X = [0,1]^{\mathbb{C}}$. Then X is compact and $|X| = \mathfrak{c}$; $d(X) = \aleph_0$ and $\chi(X) = \mathfrak{c}$.

Thus the hypothesis of d is satisfied.

Let $f: X \rightarrow \mathbb{R}$ be defined by

3. m-n Filters

In the theory of convergence and compactness, filters play an important role. We know that open filters are used in the study of $H(i)$ -space (see [62]) and Z filters are used in the study of the Stone-Čech compactification (see [19]). We shall define m - n filters, which were first introduced by J. F. Vaughan [73].

A. Definition A collection \mathcal{F} of subsets of a set X has the n -intersection property if and only if for each $F_1 \in \mathcal{F}$ with $|F_1| = n$, $\bigcap F_1 \neq \emptyset$.

A collection \mathcal{F} of non-empty subsets of a set X is said to be n -stable if and only if for each $F \in \mathcal{F}$ with $|F| < n$, there exists a $G \in \mathcal{F}$ such that $F \subset G$.

B. Definition A filter \mathcal{F} on a set X is a non-empty collection of non-empty subsets of X which is closed under finite intersections and the superset property.

If \mathcal{B} is a collection of non-empty subsets of X such that \mathcal{B} is a base for \mathcal{F} (i.e., \mathcal{F} is the collection of all sets which contain some $B \in \mathcal{B}$), then \mathcal{B} is called a base for \mathcal{F} .

C. Definition An m - n filter \mathcal{F} on a set X is a filter on X which has the n -intersection property and has a base \mathcal{B} of cardinality less than or equal to m .

An m -stable filter \mathcal{F} on a set X is a filter on X

such that for each $F \in \mathcal{F}$, there exists a set $G \in \mathcal{F}$ such that $F \subset G$ and $|G| \leq m$.

equal to m .

D. Examples

(i) Let X be any (irreducible) topological space. Let U_x denote the neighborhood of $x \in X$. The cardinality of U_x is infinite for all $x \in X$.

(ii) Let X be a set of cardinality \aleph_1 . Let U_x denote the neighborhood of $x \in X$. The cardinality of U_x is infinite for all $x \in X$.

is a function of $x \in X$.

In the example (i) if X is a topological space, then U_x is a neighborhood of $x \in X$ and U_x is a neighborhood of $x \in X$.

The cardinality of U_x is infinite for all $x \in X$.

Lemma: Let X be a topological space. Then U_x is a neighborhood of $x \in X$ and U_x is a neighborhood of $x \in X$.

Proof: Let $x \in X$. Then U_x is a neighborhood of $x \in X$ and U_x is a neighborhood of $x \in X$.

\mathcal{F}_n has the n -intersection property. *

$$\mathcal{F}_n = \{A \subseteq X : F \cap A \in \mathcal{F}_R \text{ for } |F| = n\}$$

$$|\mathcal{F}_n| \leq \sum_{i=0}^n \binom{n}{i} 2^i$$

$$= 2^{n+1} - 2$$

$$= 2^{n+1}$$

Let \mathcal{F}_n be a σ -filter on X . \mathcal{F}_n is n -stable and clearly, $\mathcal{F}_n \geq \mathcal{F}_B$. The filter \mathcal{F}_n is n -stable if and only if \mathcal{F}_n is an n -stable filter and $\mathcal{F}_n \geq \mathcal{F}_B$.

Example: Let \mathcal{F}_1 and \mathcal{F}_2 be σ -filters on a set X . Then $\mathcal{F}_1 \vee \mathcal{F}_2$ is a σ -filter on X . If \mathcal{F}_1 and \mathcal{F}_2 are n -stable filters on X , then $\mathcal{F}_1 \vee \mathcal{F}_2$ is an n -stable filter on X . If \mathcal{F}_1 and \mathcal{F}_2 are n -stable filters on X , then $\mathcal{F}_1 \vee \mathcal{F}_2$ is an n -stable filter on X .

□

4. Generalized Products

We shall outline the main facts about the k -box topology on the product $\prod\{X_i : i \in I\}$. This is a generalization of the Tychonoff product topology where k is an infinite cardinal number (see [17]).

A. Definition. Let $\mathcal{W} = \{W_i : i \in I\}$ where W_i is a subset of X_i for each $i \in I$. Then the range of \mathcal{W} is defined to be

$$R(\mathcal{W}) = \{(i \in I : W_i \neq X_i)\}$$

For example, $\mathcal{G} = \{G_i : i \in I\}$ is a basic open subset of $\prod X_i$ in the Tychonoff product topology when $|R(\mathcal{G})| < \aleph_0$.

B. Definition. Let $X = \prod\{X_i : i \in I\}$. The topology generated by the subsets of the form $W = \prod\{W_i : i \in I\}$ where each W_i is open in X_i and $|R(W)| \leq k$ is called the k -box topology on the product X and it is denoted by $(X)_k$ where $k \leq \aleph_0$.

In particular, if $k = \aleph_0$, then the k -box topology on X is the Tychonoff product topology. If $k = |I|$, then the k -box topology is usually referred to as the box topology on X .

In the case $k = |I|$, the k -box topology is not a product topology and the following theorem is well known.

THEOREM 4.1. Let $X = \prod\{X_i : i \in I\}$.

(a) If $k < |I|$, then $(X)_k$ is a product topology and $(X)_k = \prod\{X_i : i \in I\}$.

(b) If $k = |I|$, then $(X)_k$ is not a product topology.

X to X_I . In particular if $I' = \{i\}$ then we get the usual projection map $\pi_i : X \rightarrow X_i$.

D. Remark. Let $W = \prod\{W_i : i \in I\}$ where W_i is a subset of X_i for each $i \in I$. Then we have the following:

- (i) $|R(\pi_I(W))| \leq |R(W)|$
- (ii) $|\pi_I^{-1}(W_I)| = |R(W_I)|$
- (iii) If $W = \emptyset$ then $\pi_i(W) = X_i$ for all $i \in I \cap R(W)$.

F. Proposition. Let $U = \prod\{U_i : i \in I\}$ and let $V = \prod\{V_j : j \in J\}$ where U_i and V_j are subsets of X_i for each $i \in I$. Then the following are equivalent:

- (i) $U \cap V = \emptyset$
- (ii) $U_i \cap V_j = \emptyset$ for some $j \in R(U) \cap R(V)$
- (iii) $U \cap V \neq \emptyset$ and $\pi_I(U) \cap \pi_I(V) \neq \emptyset$
- (iv) $U \cap V \neq \emptyset$ and $R(U) \cap R(V) \neq \emptyset$

(i) \Rightarrow (ii): Trivial

(ii) \Rightarrow (iii): Trivial

(iii) \Rightarrow (i): Let $I' = R(U) \cap R(V)$. Then $U_i \cap V_j \neq \emptyset$ for $i \in I'$ and $j \in I'$. Hence $U \cap V \neq \emptyset$.

D. Proposition. The following are equivalent: (i) $U \cap V = \emptyset$

(ii) $U_i \cap V_j = \emptyset$ for some $j \in R(U) \cap R(V)$

(ii) $\pi_{I'}$ is continuous.

(iii) $\pi_{I'}$ is open.

Proof The above properties follow from remark C.

Furthermore we note the following: Let $X(I') = \{x \in X : x_i = a_i \text{ for } i \in I'\}$ where $a = (a_i)$ is a fixed point in X . Then $X(I')$ is homeomorphic to $(X_{I'})_k$, as a subspace of $(X)_k$. This follows from the fact that $\pi_{I'} : X(I') \rightarrow (X_{I'})_k$ is a homeomorphism from $X(I')$ to $(X_{I'})_k$.

G. Definition. Let $X = \prod\{X_i : i \in I\}$ and let $a = (a_i)$ be a fixed point in X . Then we define the γ weak sum of $\{X_i : i \in I\}$ as follows:

$$Y(Y) = \{x \in X : |\{i \in I : x_i \neq a_i\}| < \gamma\}$$

where γ is an infinite cardinal. We note that $Y(Y) = X(I')$ if $I' \in I$ where $I' = \{i \in I : i \in I' \text{ and } |\{i \in I' : i \in I'\}| < \gamma\}$. The γ weak sum $Y(Y)$ depends on the point $a = (a_i)$ and on the topology τ of X . It is clear that $Y(Y)$ is a subspace of X .

H. Theorem Let $X = \prod\{X_i : i \in I\}$ and let $a = (a_i)$ be a fixed point in X . Let $Y(Y)$ be the γ weak sum of $\{X_i : i \in I\}$ relative to a and τ .

Proof Let W be a neighborhood of a in X . Then $W \cap Y(Y)$ is a neighborhood of a in $Y(Y)$.

$$\left\{ \begin{array}{l} a_i \\ \vdots \\ a_i \end{array} \right\} \in I \in \mathcal{P}(I)$$

Then $p \in W$ and we shall show that $p \in \gamma(X)$. Consider

$$|\{i \in I : p_i \neq a_i\}| \leq |R(W)| < k \leq \gamma$$

and hence $p \in \gamma(X)$. Therefore $\gamma(X)$ is a dense subspace of $(X)_k$.

As a special case if we take $k = \gamma = \aleph_0$, then the weak-topological sum, $\aleph_0(\Pi X_i)$ is a dense subspace of X in the product topology.

I. Example. Let $X_i = \mathbf{N}$, for $i = 1, 2, \dots$, then $\aleph_0(\Pi X_i)$ is not dense in $(\Pi X_i)_{\aleph_1}$ because if $a_i \neq p_i$ for all $i \in I$, then $p \notin \aleph_0(\Pi X_i)$ but $\{p\}$ is an open subset of $(\Pi X_i)_{\aleph_1}$. Thus, the condition $k < \gamma$ cannot be relaxed in G.

CHAPTER II: m - n COMPACT SPACES

1. Some Properties of m - n Compact Spaces

1.1. Basic Properties

Compactness, countable compactness and the Lindelöf property are special cases of a more general concept: m - n compactness. In this section we shall study some properties of m - n compact spaces. Our notation follows that of Noble [53]

A. Definition. Let m and n be infinite cardinals with $m \geq n$. A topological space X is said to be m - n compact if and only if every open cover \mathcal{U} of X of cardinality $< m$ has a subcover of cardinality $\leq n$.

We say, X is m - n compact if and only if X is m - n compact for all $m \geq n$.

B. Special Cases.

- (i) \aleph_0 -compact spaces = compact spaces
- (ii) \aleph_0 - \aleph_0 compact spaces = countably compact spaces
- (iii) \aleph_1 -compact spaces = Lindelöf spaces

We will show next that m - n filters can be used to characterize m - n compact spaces. First we require a definition.

C. Definition. Let \mathcal{F} be a filter on a topological space X . Then we define the adherent of \mathcal{F} as

$$\text{ad } \mathcal{F} = \cap \{ \bar{F} \mid F \in \mathcal{F} \}.$$

The following lemma is proved by Gal [18], but the proof will be reproduced here since our terminology is different.

D. Lemma. Let X be a topological space. Then the following are equivalent:

- (i) X is m - n compact,
- (ii) every family of closed subsets of X with the $<n$ -intersection property also has the $\leq m$ -intersection property,
- (iii) for every m - n filter \mathcal{F} on X , $\text{ad } \mathcal{F} \neq \emptyset$.

Proof. (i) \Rightarrow (ii). Let $\{F_i : i \in I\}$ be a family of closed subsets of X with the $<n$ intersection property. Then $\{X - F_i : i \in I\}$ does not contain an open cover of X of size $\leq m$ and hence $\{F_i : i \in I\}$ has the $\leq m$ intersection property.

(ii) \Rightarrow (iii). Let \mathcal{F} be an m - n filter on X and let \mathcal{F}_B be a base of \mathcal{F} with $|\mathcal{F}_B| \leq m$. Then $\bar{\mathcal{F}}_B = \{\bar{F} : F \in \mathcal{F}_B\}$ is a collection of closed subsets of X with the $<n$ -intersection property and hence $\text{ad } \mathcal{F} = \cap \bar{\mathcal{F}}_B \neq \emptyset$.

(iii) \Rightarrow (i). Suppose X is not m - n compact. Then, there exists an open cover $\{U_i : i \in I\}$ of X with $|I| \leq m$ and no subcover of cardinality less than n . Hence,

$$\bar{\mathcal{F}}_B = \{X - (\cup_{i \in J} U_i) : J \in \mathcal{P}_{<n}(I)\}$$

is an m - n filter base on X but $\text{ad } F_B = \emptyset$. Hence, we have a contradiction.

E. Corollary. If n is regular and $m^n = m$, then a topological space X is m - n compact if and only if every m - n stable filter on X has a non-empty adherent.

F. Definition. A topological space is said to be n -discrete if and only if every point of X has an n -stable neighbourhood base

G. Theorem. Let X be a T_3 n -discrete space. Suppose each $x \in X$ has a m - n compact neighbourhood. If n is regular, $m^n = m$ and $\psi(X) \leq m$, then $\chi(X) \leq m$.

Proof. Let R_x be a collection of open neighbourhoods of $x \in X$ with $|R_x| \leq m$ and $\{x\} = \bigcap \{ \bar{R} : R \in R_x \}$. Let K be an m - n compact neighbourhood of x . Let $U_x = \{ \bigcup_{i=1}^n U_i : U_i \in R_x \}$. Then we shall show that $\{ U_x \mid K \cap U \neq \emptyset, U \in U_x \}$ is a neighbourhood base at x . Let V be any open neighbourhood of x and suppose $(X \setminus V) \cap (U \cap K) \neq \emptyset$ for every $U \in U_x$. Then $\{ U \mid (K \cap X \setminus V) \cap U \neq \emptyset, U \in U_x \}$ is a m - n stable filter base on K and

$$\begin{aligned} \text{ad}_K (U_x \mid (K \cap X \setminus V)) &\subseteq (\bigcap \bar{U}_x) \cap (X \setminus V) \cap K \\ &\subseteq (\bigcap \bar{B}_x) \cap (X \setminus V) \cap K \\ &= \{x\} \cap (X \setminus V) = \emptyset. \end{aligned}$$

By corollary E, this is a contradiction. Now it follows that

$$\chi(X) \leq m.$$

G. Corollary. Let X be a T_3 , m - n compact, κ - n -discrete space. If n is regular, $m^n = m$ and $\psi(X) \leq m$, then $\chi(X) \leq m$.

1.2. Projection Maps

We shall show that, under suitable conditions on factor spaces, the projection map parallel to an m - n compact factor is a closed map.

A. Proposition. Let the projection $\pi_X : X \times Y \rightarrow X$ be a closed map. Suppose Y is a κ - n discrete, T_1 space and $\chi(p, X) = n$ ($\geq \aleph_0$) for some $p \in X$. Then, Y is n - n compact.

Proof. Suppose Y is not n - n compact. Then there exists a n - n filter base \mathcal{F} on Y such that $\bigcap \mathcal{F} = \emptyset$. Let

$$I = \{I_\alpha : \alpha < W(n)\}$$

$$K_\alpha = \{I_\rho : \rho < \alpha\}$$

where $W(n)$ is the least ordinal of cardinality n . Let

$V_p = \{V_\alpha : \alpha < W(n)\}$ be an open neighbourhood base at p and let

$S_\alpha = X \times V_\alpha$. Then,

$$p \in \overline{\bigcup \{S_\alpha : \alpha < W(n)\}}$$

and

$$p \notin \overline{\bigcup \{S_\alpha : \alpha < \alpha_0\}}$$

for all $\alpha_0 \in W(n)$. Let $y \in Y$. Since $n\bar{F} = \emptyset$, there exists an $\alpha_0 \in W(n)$ and an open set W of Y such that $y \in W$ and $W \cap K_{\alpha_0} = \emptyset$. We set,

$$F = \overline{\cup \{S_\alpha \times K : \alpha \in W(n)\}}$$

and note that $(p, y) \notin F$, and hence $p \notin \pi_X(F)$. But

$$\pi_X(F) \supseteq \overline{\pi_X(\cup \{S_\alpha \times K : \alpha \in W(n)\})}$$

and thus $p \in \overline{\pi_X(F)}$. This is a contradiction because π_X is a closed map

B. Proposition. If X is κ -discrete and has character $\leq m$ and Y is m - n compact, then the projection $\pi_X : X \times Y \rightarrow X$ is a closed map.

Proof. Let $F \subseteq X \times Y$ be a closed set and $x \in \pi_X(F)$. Let \mathcal{V}_x be a neighbourhood system at x in X with $|\mathcal{V}_x| \leq m$. We set

$$G = \{ \pi_Y^{-1}(V) \cap F : V \in \mathcal{V}_x \}$$

and let $\mathcal{W}_Y(G) = \{ \pi_Y(G) : G \in \mathcal{G} \}$. Then easily $\mathcal{W}_Y(G)$ is a base for an m - n filter on Y and hence, by 2-B, we can find some $y \in \cap \{ \pi_Y(G) : G \in \mathcal{G} \}$. Now, if V and W are neighbourhoods of x and y , then W meets $\pi_Y(\pi_X^{-1}(V) \cap F)$ and hence $\pi_Y^{-1}(W) \cap \pi_X^{-1}(V) \cap F \neq \emptyset$. Thus every basic neighbourhood $V \times W$ of (x, y) in $X \times Y$ meets F and hence $(x, y) \in \bar{F} = F$. Then clearly, $x \in \pi_X(F)$.

2. Products of Lindelöf Spaces

Our main aim is to prove that a product of Lindelöf, T_3 , P-spaces is Lindelöf whenever it is quasi paracompact.

2.1. Basic Terminology

We shall define quasi paracompactness and show that the class of quasi paracompact spaces is larger than the class of paracompact, T_2 -spaces

A. Definition. Let X be a topological space. A collection C of subsets of X is said to be $\leq n$ -closure preserving if and only if for every $C' \subseteq C$ with $|C'| \leq n$,

$$\overline{\bigcup\{C : C \in C'\}} = \bigcup\{\overline{C} : C \in C'\}$$

where $n \in \mathbb{N}$.

B. Examples.

(i) Every locally finite collection is $\leq n$ -closure preserving for every n .

(ii) Let X be a $\leq n$ -discrete space. Then any collection of subsets of X is $\leq n$ -closure preserving.

C. Definition. A T_2 -space X is said to be quasi n -paracompact if and only if every open cover of X has an open, $\leq n$ -closure preserving refinement.

A T_2 -space X is said to be quasi paracompact if and only if X is quasi \mathcal{K}_1 -paracompact.

The class of quasi paracompact spaces contains the class of paracompact spaces and the class of Lindelöf T_2 spaces. Thus the class of quasi paracompact spaces is larger than the class of paracompact T_2 -spaces.

2.2 Countable Products

We shall consider the product of a countable family of spaces. The product is normal and T_1 .

A. Theorem (J. E. Vaughan [73]) If each X_j is m -compact, Lindelöf and has character $\leq m$ for $j = 1, 2, \dots$, and if m is regular and $m^n = m$, then $\prod (X_j; j = 1, 2, \dots)$ is m -compact.

B. Corollary Let m be a regular cardinal. If each X_j is compact and Lindelöf for $j = 1, 2, \dots$, then $\prod X_j$ is m -compact.

Since \aleph_1 is a regular cardinal, it follows that $\prod X_j$ is \aleph_1 -compact.

C. Theorem (M. Hodel [47]) $\prod X_j$ is Lindelöf if and only if each X_j is Lindelöf.

Next, we shall consider \aleph_1 -paracompact products. The following condition is necessary for \aleph_1 -paracompactness. It was first proved by Hodel [47].

D. Theorem. If $X = \bigcup\{X_i : i \in I\}$ is a normal space and if each X_i is T_1 , then all but countably many of the X_i are countably compact.

F. Proposition

(i) Let X be a T_1 space. If X is a countably compact, then X is finite.

(ii) If X is compact and Y is compact, then $X \times Y$ is compact.

Proof. (i) If X had a countably infinite subset B , then B is closed and discrete. This is a contradiction because X is countably compact.

Hence X is finite.

(ii) Standard methods work.

We shall show that normality and paracompactness coincide in the metrisable space, when the factor spaces are Lindelöf. P-spaces

Proposition. If X is a metrisable space, then the following are equivalent:

- (i) X is Lindelöf.
- (ii) X is normal.
- (iii) X is paracompact.

(i) \implies (iii) follows from standard results.

(ii) \implies (iii) follows from these in P.

(iii) \implies (i) follows from the fact that every paracompact metrisable space is Lindelöf.

G: Proposition. Let $X = \prod\{X_i : i \in I\}$ be a normal T_1 -space. If each X_i is ω - n compact and ω - n -discrete, where n is a regular cardinal then X is ω - n compact.

Proof. By theorem D, there exists a countable subset J of I such that $X = X_J \times X_{J^c}$, where X_J is a countable product of ω - n compact ω - n -discrete spaces and X_{J^c} is an arbitrary product of finite spaces. Hence X is ω - n compact.

We shall prove that

X is ω - n compact.

Let \mathcal{A} be a family of closed subsets of X with the finite intersection property.

By the definition of ω - n compactness, it suffices to show that

$\bigcap \mathcal{A} \neq \emptyset$.

Since $|I| \leq n$, X is ω - n compact.

B. Corollary. Every paracompact T_2 -space with a dense ω - n compact subspace is ω - n compact.

C. Corollary. Every separable, paracompact T_2 -space is Lindelöf.

D. Theorem [theorem 2.3 of [13]]. Let $m \geq n > \gamma \geq \aleph_0$ and $n \geq \alpha \geq \aleph_0$ with α regular and strongly γ -inaccessible. Let $\{X_i : i \in I\}$ be a family of spaces such that $(\prod_{i \in I} X_i)_\alpha$ is m - n compact for all $I \in P_\gamma(I)$. Then $(\prod_{i \in I} X_i)_\alpha$ is m - n compact with respect to the product topology.

E. Corollary. Let $\alpha \geq \aleph_0$ be regular and strongly γ -inaccessible. Let $\{X_i : i \in I\}$ be a family of spaces such that $(\prod_{i \in I} X_i)_\alpha$ is m - n compact for all $I \in P_\gamma(I)$. Then $(\prod_{i \in I} X_i)_\alpha$ is m - n compact with respect to the product topology.

F. Theorem. Let $\alpha \geq \aleph_0$ be regular and strongly γ -inaccessible. Let $\{X_i : i \in I\}$ be a family of spaces such that each X_i is ω - n compact and α is a regular cardinal. Then $(\prod_{i \in I} X_i)_\alpha$ is ω - n compact.

G. Proof. We note that, by 2-F, $(\prod_{i \in I} X_i)_\alpha$ is ω - n compact. Hence we can write 3-F. $(\prod_{i \in I} X_i)_\alpha$ is ω - n compact.

H. Theorem. Let $\alpha \geq \aleph_0$ be a quasi- n -paracompact space. Let $\{X_i : i \in I\}$ be a family of spaces such that each X_i is ω - n compact. Then $(\prod_{i \in I} X_i)_\alpha$ is ω - n compact.

ω -discrete and T_3 ; then X is ω -n compact.

Proof. Since $\mathcal{K}_0(\Pi\{X_i : i \in I\})$ is a dense, ω -n compact subspace of X , the result follows by lemma 3-A.

H. Corollary. Let $X = \Pi\{X_i : i \in I\}$ be a quasi paracompact space. If each X_i is Lindelöf, T_1 , then X is a Lindelöf space.

3. Locally ω -n Compact Spaces

We define locally ω -n compact spaces and study the basic properties of this class.

A. Definition. A topological space X is said to be locally ω -n compact if every $x \in X$ has a neighbourhood base consisting of ω -n compact subsets, where $n \geq \aleph_0$.

A locally ω - \aleph_0 compact space is called a locally compact space and a locally ω - \aleph_1 compact space is called a locally Lindelöf space.

B. Examples.

(i) The set of rationals, as a subspace of \mathbb{R} , is locally Lindelöf but not locally compact.

(ii) Let X be an uncountable discrete space. We adjoin an extra point p to X and specify its neighbourhoods to be $A \cup \{p\}$ where $A \subseteq X$ and $X - A$ is countable. The neighbourhoods of points of X remain the same. The space $X \cup \{p\}$ is called the one-point Lindelöf extension of X .

We state the following properties:

(a) X is Lindelöf if and only if X^* is Lindelöf.

(b) X is T_3 if and only if X^* is T_3 .

(c) X is a T_1 space if and only if X^* is a T_1 space.

(d) If $A \subseteq X^*$ and A is compact, then $|A| \leq \aleph_0$.

(e) If X is Lindelöf, then X^* is not locally compact.

C. Some Properties. The following properties are useful and they are easy consequences of the definition A:

(i) In the presence of regularity, to show that a space X is locally ω - n compact it is sufficient to find one ω - n compact neighbourhood at each $x \in X$. Thus every ω - n compact, T_3 space is locally ω - n compact.

(ii) Every open subspace of a regular, locally ω - n compact space is locally ω - n compact.

(iii) Let X be a ω - n -discrete T_2 space. Then every locally ω - n compact subset is the intersection of an open set and a closed set.

(iv) Let X be a locally ω - n compact, T_3 , ω - n -discrete space. If A is dense in X , then A is locally ω - n compact if and only if A is open.

D. Definition. A topological space X is said to be a $k(n)$ -space if for each $A \subseteq X$, the set A is open in X if and only if $A \cap Z$ is open in Z for every ω - n compact subset Z of X .

A $k(n)$ -space is called a $k(n)$ -space if it is a $k(n)$ -space and a T_3 -space.

E. Proposition. Every locally ω - n compact space is a $k(n)$ -space.

Proof. Suppose X is a locally ω - n compact space and $A \cap Z$ is open in Z for every ω - n compact subset Z of X . Let $a \in A$ and let M be a ω - n compact neighbourhood of a . Then we note that

$A \cap M = (A \cap M) \cap M$. Let M be open in M and containing a . Then

discrete spaces are locally ∞ -n compact for any n , the property locally ∞ -n compact need not be preserved by continuous maps.

F. Note. Continuous open maps preserve local ∞ -n compactness but not continuous closed maps.

G. Theorem. Let X be a T_3 -space. Then X is a $k(n)$ -space if and only if X is a quotient of some locally ∞ -n compact space.

Proof. Suppose X is a $k(n)$ -space. Let $\mathcal{B}(n)$ denote the collection of all ∞ -n compact subspaces of X . Let $|\mathcal{B}(n)| = k$. For each $B \in \mathcal{B}(n)$ we set

$$B(t) = B \times t \text{ where } t \in k.$$

Then Y , the topological union of $\{B(t) : t \in k\}$, is locally ∞ -n compact and the map $f: Y \rightarrow X$ defined by $f(x,t) = x$ is a quotient map. This completes the proof of necessity.

Suppose $p: Y \rightarrow X$ is a quotient map and Y is locally ∞ -n compact. Let U be a subset of X such that $U \cap B$ is open in B for every $B \in \mathcal{B}(n)$. Let $y \in p^{-1}(U)$ and let V be a locally ∞ -n compact neighbourhood of y . Then $U \cap p(V)$ is open in $p(V)$ and hence there exists an open subset G of X such that $U \cap p(V) = p(V) \cap G$. Now we note that

$$y \in p^{-1}(G) \cap \text{Int } V \subset p^{-1}(U).$$

Hence $p^{-1}(U)$ is open in Y . Since p is a quotient map, U is open in X . This proves the sufficiency.

4. Tightness in Product Spaces

We consider the following question:

Is $\partial(X \times Y) \leq \partial(X) \cdot \partial(Y)$ where
 X and Y are any two topological spaces?

4.1. Finite Products

We shall show that for certain classes of spaces,

$$\partial(X \times Y) = \partial(X) \cdot \partial(Y).$$

A. Definition. Let $p \in X$, $S \in X$ and $p \in \bar{S}$. Then, we define

$$\partial(p, S, X) = \min \{ |M| : M \subset S \text{ and } p \in \bar{M} \}$$

$$\partial(p, X) = \sup \{ \partial(p, S, X) : p \in \bar{S} \}$$

$$\text{and } \partial(X) = \sup \{ \partial(p, X) : p \in X \} + \aleph_0$$

which is called the tightness of X .

B. Some Terminology. If $R = \{R_i : i \in I\}$ is any collection of topological spaces, we define,

$$\partial_I(R) = \sup \{ \partial(R_i) : i \in I \}$$

We say, the product $R = \prod_{i \in I} R_i$ preserve tightness if and only if

$$\partial(\prod_{i \in I} R_i) = |I| \cdot \partial_I(R).$$

For finitely many X_1, X_2, \dots, X_n this reduces to,

$$\alpha(\prod X_i) \leq \alpha(X_1) \cdot \alpha(X_2) \cdots \alpha(X_n)$$

C. Definition. A space X is called a GC(n,i)-space for $i = 1, 2, 3, 4$ if X is an ω -n compact, $<n$ -discrete, T_i -space.

A space X is called a GLC(n,i)-space for $i = 1, 2, 3, 4$ if X is a locally ω -n compact, $<n$ -discrete, T_i -space. Always, $GC(n,i) \subseteq GLC(n,i)$ for $i = 1, 2, 3, 4$.

D. Proposition. If X is a $GC(n,3)$ space, then X is (strongly) paracompact and T_2 .

Proof. The case $n = \aleph_0$ is straight-forward. We shall thus consider only $n \geq \aleph_1$. Since X is a T_3 P-space, easily X is zero-dimensional. Thus, if U is an open cover of X , then we can find a refinement V of U , consisting of cl-open sets with $|V| < n$. Let $V = \{V_\alpha : \alpha \in k \in W(n)\}$. Now we set $W_\alpha = V_\alpha - \cup\{V_\beta : \beta < \alpha\}$ for each $\alpha \in k$. Then, clearly $\{W_\alpha : \alpha \in k\}$ is an open star-finite refinement of U . Hence X is (strongly) paracompact and T_2 .

Next, we prove a basic result which generalizes Juhász's lemma for the compact case (page 113; [34]).

E. Lemma. If X is a $<n$ -discrete T_1 -space and Y is an ω -n compact regular space, then

$$\alpha(X \times Y) \leq \alpha(X) \cdot \alpha(Y)$$

Proof. Let $k = \alpha(X) \cdot \alpha(Y)$ and suppose $H \subset X \times Y$ is k -closed. It suffices to show H is closed.

Let $(p,q) \in \bar{H}$. If $T = H \cap (\{p\} \times Y)$, then T is k -closed and hence closed in $\{p\} \times Y$, and hence closed in $X \times Y$. We need only show $q \in \pi_Y(T)$.

Suppose $q \notin \pi_Y(T)$. Since $\pi_Y|_{\{p\} \times Y}$ is a homeomorphism, $\pi_Y(T)$ is closed in Y . Let V be a closed neighbourhood of q such that $V \cap \pi_Y(T) = \emptyset$. Note that since $X \times V$ is a neighbourhood of (p,q) and $(p,q) \in \bar{H}$, we have $(p,q) \in \overline{(X \times V) \cap H}$. But $(X \times V) \cap H$ is a closed subset of H , and hence k -closed in $X \times Y$. Since Y is ∞ -compact and X is ∞ -discrete, π_X is closed by 1.2 B. Thus $\pi_X[(X \times V) \cap H]$ is k -closed and hence closed in X . Then, since π_X is continuous,

$$\begin{aligned} p \in \pi_X[\overline{(X \times V) \cap H}] &\subset \overline{\pi_X[(X \times V) \cap H]} \\ &= \pi_X[(X \times V) \cap H]. \end{aligned}$$

So for some $v \in V$, $(p,v) \in H$. But then $v \in \pi_Y(T) \cap V$, a contradiction.

F. Theorem. If X is ∞ -discrete, T_1 , and Y is locally ∞ -compact, T_3 , then

$$\alpha(X \times Y) \leq \alpha(X) + \alpha(Y)$$

Proof. Let $k = \alpha(X) + \alpha(Y)$ and let $H \subset X \times Y$ be k -closed. Choose $(p,q) \in \bar{H}$. Let V be any closed ∞ -compact neighbourhood of q . Then $X \times V$ is closed in $X \times Y$ and $\alpha(X \times V) \leq k$ by D. But $H \cap (X \times V)$ is k -closed in $X \times V$, and thus closed in $X \times V$, and $(p,q) \in \text{Cl}_{X \times V} H \cap (X \times V) = \overline{H \cap (X \times V)}$. But $(p,q) \in H$.

G. Corollary. If X_1, X_2, \dots, X_n are $GLC(n, 3)$ spaces, then

$$\partial\left(\prod_{i=1}^n X_i\right) \leq \partial(X_1) \cdot \dots \cdot \partial(X_n)$$

Proof. By induction, noting that

$$X_1 \times \dots \times X_k = (X_1 \times \dots \times X_{k-1}) \times X_k$$

and that $X_1 \times \dots \times X_{k-1}$ is $<n$ discrete and T_1 , while X_k is locally ∞ -n compact and T_3 , so that

$$\partial(X_1 \times \dots \times X_k) \leq \partial(X_1 \times \dots \times X_{k-1}) \cdot \partial(X_k)$$

by E.

Thus, finite products of locally compact, T_2 -spaces ($= GLC(\aleph_0, 3)$) preserve tightness and so also do finite products of locally Lindelöf T_3 P-spaces ($= GLC(\aleph_1, 3)$), and so on. We shall consider infinite products in the next section.

4.2. Large Products

In this section, we shall extend the results of the previous sections to arbitrary products.

A. Notation. Let $R = \prod\{R_i : i \in I\}$ and suppose $J \subset I$. The subproduct $\prod\{R_i : i \in J\}$ of R will be denoted by R_J and the projection of R onto R_J will be denoted by π_J . For $a \in R$ and $A \subset R$, the images $\pi_J(a)$ and $\pi_J(A)$ will be denoted by a_J and A_J respectively.

B. Proposition. If each finite subproduct of a product $R = \prod\{R_i : i \in I\}$ preserves tightness, then R preserves tightness.

Proof. Let $R = \{R_i : i \in I\}$ and set $k = |I| \partial_I(R)$. Let $I = \{J \subset I : |J| < \aleph_0\}$. Suppose $A \subset R$ is k -closed and $a \in \bar{A}$. Then for $J \in I^{(F)}$, a_J belongs to $\pi_J(\bar{A}) \subset \bar{A}_J$. Since $\partial(R_J) \leq k$, we can find $B_J \subset A_J$ with $|B_J| \leq k$ such that $a_J \in \bar{B}_J$.

For each $b \in B_J$, choose $x_b \in A$ so that $\pi_J(x_b) = b$, and set

$$C_J = \{x_b : b \in B_J\}.$$

Clearly $|C_J| = |B_J| \leq k$ and hence if $C = \cup\{C_J : J \in I^{(F)}\}$, then $C \subset A$ and $|C| \leq k \cdot |I| = k$.

But $a \in \bar{C}$. For if $U = U_J \times \prod_{i \notin J} R_i$ is a basic open neighbourhood of a , then since $\pi_J(a) \in \bar{B}_J = \overline{\pi_J(C_J)}$, we have $U_J \cap \pi_J(C_J) \neq \emptyset$, and thus $U \cap C \neq \emptyset$, whence $U \cap C \neq \emptyset$.

Thus $a \in A$, and A is closed.

C. Theorem. If P_i is a $GU(n)$ space for each $i \in I$, then $\partial(\prod R_i) = |I| \partial_I(R)$

Proof. Apply B and C.

In 8.9 of [31] it is shown that for any collection of compact T_2 spaces $\{P_i : i \in I\}$, $\partial(\prod R_i) = |I| \partial_I(P)$

The special cases of C generalize this result

D. Corollary. If R_i is locally compact and T_2 for each $i \in I$, then $\partial(\prod R_i) = |I| \cdot \partial_I(R)$.

Proof. Locally compact, $T_2 \equiv \text{GLC}(\aleph_0, 3)$.

E. Corollary. If R_i is a locally Lindelöf, T_3 , P-space for each $i \in I$, then $\partial(\prod R_i) = |I| \cdot \partial_I(R)$.

Proof. Locally Lindelöf, T_3 , P-space $\equiv \text{GLC}(\aleph_1, 3)$.

CHAPTER III: WEAKLY m - n COMPACT SPACES

1. Some Properties of Weakly m - n Compact Spaces

1.1. Basic Facts

The properties $H(i)$, feebly compact and weakly Lindelöf are special cases of the general concept weak m - n compactness. In this section we shall study basic properties of weakly m - n compact spaces.

A. Definition. A topological space X is said to be weakly m - n compact if and only if every open cover of X of cardinality $\leq m$ has a sub family of cardinality $\leq n$ with dense union, where $m \geq n \geq 1$.

A topological space X is said to be weakly m - n compact if and only if X is weakly m - n compact for each $m \geq n \geq 1$.

B. Special Cases (see [61] and [60])

(i) Weakly \aleph_1 - \aleph_1 compact spaces = $H(i)$ spaces

(ii) Weakly \aleph_1 - \aleph_0 compact spaces = feebly compact spaces

(iii) Weakly \aleph_0 - \aleph_0 compact spaces = Lindelöf spaces

The $H(i)$ spaces are called in [61] Lindelöf.

C. Definition. A topological space X is said to be weakly m - n compact if and only if X is weakly m - n compact for each $m \geq n \geq 1$.

Q.E.D.

It is easy to see that the following properties are equivalent:

(1) X is weakly m - n compact

$|U| \leq m$, has a sub-family V with $|V| < n$ and $E \subseteq \overline{UV}$.

It is clear that every weakly m - n compact set is relatively weakly m - n compact. In general, the converse is not true, except for open subsets.

D. Example. Let X be a discrete subspace of $\beta\mathbb{N} - \mathbb{N}$, with cardinality c . Let $Y = X \cup \mathbb{N}$ as a subspace of $\beta\mathbb{N}$. Then X is a relatively weakly Lindelöf subset of Y , but not weakly Lindelöf. Here X is a closed subset of Y .

Next we shall study basic properties of weakly m - n compact spaces.

1. Proposition

(i) Let n be an infinite cardinal. If $\{X_i : i \in I\}$ is a collection of relatively weakly m - n compact subsets of X , then $\bigcup_{i \in I} X_i$ is relatively weakly m - n compact, provided $|I| < cf(n)$.

(ii) If X has a dense, relatively weakly m - n compact subset, then X is weakly m - n compact.

(iii) A continuous image of a weakly m - n compact space is weakly m - n compact.

(iv) A regular space is weakly m - n compact if and only if it is relatively weakly m - n compact.

Proof. (i) Let \mathcal{U} be a family of open sets in $\bigcup_{i \in I} X_i$ with no finite subfamily covering $\bigcup_{i \in I} X_i$. By the definition of relatively weakly m - n compact, for each $i \in I$, there is a subfamily \mathcal{U}_i of \mathcal{U} with $|\mathcal{U}_i| < n$ and $X_i \subseteq \overline{\bigcup \mathcal{U}_i}$.

Let $\mathcal{V} = \bigcup_{i \in I} \mathcal{U}_i$. Then $|\mathcal{V}| < n$ and $\bigcup_{i \in I} X_i \subseteq \overline{\bigcup \mathcal{V}}$. Thus $\bigcup_{i \in I} X_i$ is weakly m - n compact.

space X . Let \mathcal{V} be an open cover of B of cardinality $\leq m$. For each $V \in \mathcal{V}$ we select an open subset U_V of X such that $V = U_V \cap B$. Now $U = \{\cup_V U_V \cap V\} \cap (X \setminus B)$ is an open set with $|U| \leq m$. Since X is countable, we can find a countable set \mathcal{U} with $U \subset \mathcal{U}$ and

$$v \in \overline{\mathcal{U}} \iff v \in U \text{ for some } U \in \mathcal{U}.$$

It follows that

$$B \cap \overline{\mathcal{U}} = \overline{\mathcal{U}} \cap B = \mathcal{U} \cap B.$$

Thus B is a σ - \mathcal{U} -set. \square

Proposition 1.10. Let X be a topological space.

Then

- (i) p is a σ - \mathcal{U} -set
- (ii) \mathcal{U} is a σ - \mathcal{U} -set

Proof. (i) Let \mathcal{U} be a countable family of open sets.

Let

and

Proposition 1.10. Let X be a topological space. Then

if \mathcal{U} is a σ - \mathcal{U} -set, then $\mathcal{U} \cap B$ is a σ - \mathcal{U} -set for every $B \in \mathcal{C}(X)$.

Proof. Let \mathcal{U} be a σ - \mathcal{U} -set. Then there is a countable family \mathcal{V} of open sets such that

$\mathcal{U} = \{\cup_V U_V \cap V\}$ for some open sets U_V of X .

Let $B \in \mathcal{C}(X)$. Then $\mathcal{U} \cap B = \{\cup_V U_V \cap V \cap B\}$.

6. Corollary. If X is weakly m - n compact and Y is compact, then $X \times Y$ is weakly m - n compact.

11. The projection map $\pi_X: Y \times X \rightarrow X$ satisfies the properties

Special Properties

In this section we shall discuss special properties of weakly m - n compact spaces.

Theorem. A topological space X is weakly m - n compact if every closed m - n subset of X is relatively m - n compact.

Proof. Let \mathcal{U} be a point cover of X of cardinality $\leq m$. If $U \in \mathcal{U}$ is not closed, then $R = \text{cl}_X(U - \bar{U})$. Then clearly R is a closed m - n subset of X and hence relatively m - n compact. Thus \mathcal{U} is a point cover of X of cardinality $\leq m$.

THEOREM 11

Let \mathcal{U} be a point cover of X .

If \mathcal{U} is a point cover of X of cardinality $\leq m$, we have:

Corollary. A topological space X is weakly m - n compact if and only if every closed m - n subset of X is relatively m - n compact.

We shall show that the property weak m - n compactness is not closed hereditary. This is one of the intrinsic differences between weak m - n compactness and m - n compactness.

C. Example Whenever $n > \aleph_0$, there exists a T_2 space X such that

- (i) X is weakly m - n compact, and
- (ii) X has a closed discrete subspace of cardinality n .

To see this, let Y be any T_2 space with disjoint dense subsets A and B , where $|B| = n$ and let Z be the discrete space of cardinality n . We form X by adjoining to the product space $T_2 \times Z$ a copy of B with a topology τ of the form

$$[U \cap B] \cup (U \cap Z)$$

where U is a neighbourhood of z in Z .

Note that easily (a) X is a closed linear subspace of cardinality n , for $z \in Z$. It is clear that X is weakly m - n compact by our assumption on Z . To see that X is not m - n compact, let $\{x_i\}_{i \in I}$ be a discrete subspace of X containing B and in fact every x_i is a point of B .

Now let $\{U_i\}_{i \in I}$ be a family of open sets such that $x_i \in U_i$ and $U_i \cap U_j = \emptyset$ for $i \neq j$. Then $\{U_i\}_{i \in I}$ is a family of open sets such that $x_i \in U_i$ and $U_i \cap U_j = \emptyset$ for $i \neq j$.

E. Proposition. Let X be a regular, quasi n -paracompact space. If X is weakly ω - n compact, then X is ω - n compact.

Proof. Let \mathcal{U} be an open cover of X . Since X is regular, we can find an open refinement \mathcal{V} of \mathcal{U} such that for each $V \in \mathcal{V}$, $\bar{V} \subseteq U$ for some $U \in \mathcal{U}$. Now let \mathcal{W} be an open, ω -closure preserving refinement of \mathcal{V} . Since X is weakly ω - n compact, there exists a $\mathcal{W}' \subseteq \mathcal{W}$ of cardinality $\leq n$ and $X = \cup\{\bar{W} : W \in \mathcal{W}'\}$. For each $W \in \mathcal{W}'$, $\bar{W} \subseteq U$ for some $U \in \mathcal{U}$. Thus there exists $\mathcal{U}' \subseteq \mathcal{U}$ of cardinality $\leq n$ and $X = \cup\{U : U \in \mathcal{U}'\}$.

F. Corollary. The property weak ω - n compactness coincides with ω - n compactness in paracompact T_2 spaces and in ω -discrete, regular spaces.

G. Example. Let Y be the one-point Lindelöf extension of an uncountable discrete space of cardinality \aleph_1 . Let Y be the Alexandroff double of Y . Then Y has a closed nowhere dense subspace E of cardinality \aleph_1 . We define a new topology on Y in the following manner:

- (i) the neighborhoods of points of $Y \setminus E$ remain unchanged,
- (ii) the neighborhoods of points of E are of the form,

$$U_p^* = (U_p \cap E) \cup \{p\}$$

§ U_p is a neighborhood of p in the original topology of Y .

Calling this space Z , we note that $C_0(Z) = C_0(Y)$ for all $p \in E$.

Now Z is a Lindelöf T_2 space, so we have the following:

- (i) Z is Lindelöf.

(ii) Z is a P-space,

(iii) $L(Z) = c(Z) = k$,

(iv) Z is T_2 .

Thus the class of weakly Lindelöf T_2 , P-spaces is larger than the class of Lindelöf, T_2 , P-spaces.

2. Some Characterizations

2.1. Filters

By generalizing the intersection properties satisfied by open filters we wish to define strong m - n filters and using this generalized notion we shall characterize weakly m - n compact spaces (see [18]).

A. Definition. A collection F of subsets of a space X is said to have the $<n$ -strong intersection property if for each $F' \subseteq F$ with $|F'| < n$, we have $\text{Int}(\cap F') \neq \emptyset$.

A collection F of non-empty subsets of a space X is said to be $<n$ -strongly stable if for each $F' \subseteq F$ with $|F'| < n$ we have $\text{Int}(\cap F') \supseteq F$ for some $F \in F$.

B. Definition. A strong m - n filter F on a space X is a filter on X which has the $<n$ -strong intersection property and has a base F_B of cardinality less than or equal to m .

A strong m - n stable filter F on X is a filter on X which has the $<n$ -strong stable property and has a base F_B of cardinality less than or equal to m .

C. Remark. Strong m - n filters are topological objects but m - n filter σ (filters) are set theoretical objects; every strong m - n filter is m - n filter but not conversely.

D. Example. Let X be a T_1 space with character $< k$. Let \mathcal{X} be a

non-isolated point of X . Then the neighbourhood system \mathcal{V}_x at x is a m - n filter but not a strong m - n filter, where $m \geq n > k$.

E. Definition. A filter F is said to be of type k if $|F| \leq k$ for every $F \in \mathcal{F}_B$ where \mathcal{F}_B is some filter base for F .

F. Note. Every strong m - n stable filter on a k -separable space X induces a finer m - n stable filter of type k .

G. Lemma. Let F be a strong m - n filter on the space X . Let n be a regular cardinal number and let $m^n = \sum\{m^k : k < n\} = m$. Then there exists a strong m - n stable filter G on X such that $G \supseteq F$.

Proof. Let \mathcal{F}_B be a filter base for F . Let $\mathcal{G}_B = \{\text{Int}(nF') : F' \in \mathcal{F}_B \text{ and } |F'| < n\}$. Then $|\mathcal{G}_B| \leq |\mathcal{F}_B|^n \leq m^n = m$. Let $G' \in \mathcal{G}_B$ and $|G'| < n$. Then $\text{Int}(nG') = \text{Int}(n(\text{Int}(nF')))$ \supseteq $\text{Int}(nF'')$ where $F'' \in \mathcal{F}_B$ and, since n is regular, $|F''| < n$. Hence the filter G generated by \mathcal{G}_B is a strong m - n stable filter on X and $G \supseteq F$.

The above lemma shows that every strong m - n filter can be embedded in a strong m - n stable filter provided n is regular and $m^n = m$.

H. Theorem. Let X be a topological space. Then the following are equivalent:

- (i) X is weakly m - n compact.
- (ii) Every family of closed subsets of X with the n -strong intersection property also has the m -intersection property.

(iii) Every strong m - n filter on X has an adherent point.

Proof. (i) \Rightarrow (ii): Let $\{F_i : i \in I\}$ be a family of closed subsets of X with the $< n$ -strong intersection property. Then $\{X - F_i : i \in I\}$ contains no m -fold open cover of X and hence $\{F_i : i \in I\}$ has the $< m$ -intersection property.

(ii) \Rightarrow (iii): Let F be a strong m - n filter on X . Let F_B be a base for F such that $|F_B| \leq m$. Then $\{\bar{F} : F \in F_B\}$ has the $< n$ -strong intersection property and by the hypothesis, $\bigcap F_B \neq \emptyset$. Therefore F has an adherent point.

(iii) \Rightarrow (i): If X is not weakly m - n compact, then there exists an m -fold open cover $\{G_i : i \in I\}$ of X with no dense sub-family of cardinality strictly less than n . Hence $\{X - G_i : i \in I\}$ has the $< n$ -strong intersection property and therefore $\{X - G_i : i \in I\}$ is a filter sub-base for some strong m - n filter F . But $\bigcap F = \bigcap \{X - G_i : i \in I\} = \emptyset$. We have a contradiction and hence the result.

I. Corollary. Let n be a regular cardinal and let $m^n = m$. Then a topological space X is weakly m - n compact if and only if every strong m - n stable filter on X has a non-empty adherent.

Proof. This follows from lemma G above.

2.2 Continuous Maps

We wish to characterize weakly m - n compact spaces using the local character of the points of $Cl_Y f(X) \subset f(X)$ where $f: X \rightarrow Y$ is a continuous map. We will employ strong m - n stable filters.

A. Proposition. Let X be a $\langle n$ -discrete, non weakly m - n compact T_2 -space, where n is regular and $m^n = m$. Then there exists a $\langle n$ -discrete T_2 -space Y and a continuous map $f: X \rightarrow Y$ such that

- (i) $\text{Cl}_Y f(X) = Y$ and
- (ii) $Y - f(X)$ has a point of local character $\leq m$.

Proof. Let $\mathcal{F} = \{F_k : k \in I\}$ be a strong m - n stable filter base on X such that $\bigcap \mathcal{F} = \emptyset$. Let $Y = X \cup \{w\}$ where $w \notin X$, with the topology given by the open subsets of X together with the sets of the form $\text{Int } F_k \cup \{w\}$ where $k \in I$. Then the inclusion map $i: X \rightarrow Y$ is continuous and it is easy to see that Y satisfies the above properties.

B. Proposition. Let X be a weakly m - n compact space. Let Y be a $\langle n$ -discrete T_2 -space and let $f: X \rightarrow Y$ be a continuous map. Then $\text{cl}_Y f(X) - f(X)$ has no points of local character $\leq m$.

Proof. Suppose $\text{Cl}_Y f(X) - f(X)$ has a point y of local character $\leq m$. Let \mathcal{V} be an open neighbourhood base at y with $|\mathcal{V}| \leq m$. Then $\mathcal{V}|f(X)$ is a strong m - n stable filter base on $f(X)$, but $\bigcap \mathcal{V} \cap Y = \emptyset$ and $y \notin f(X)$. It follows that

$$\bigcap \{ \text{cl}_Y (V \cap f(X)) \mid V \in \mathcal{V} \} \cap Y = \emptyset \cap Y = \emptyset \neq \{y\} \cap Y$$

But then $f(X)$ is not weakly m - n compact, a contradiction.

C. Theorem. A $\langle n$ -discrete, T_2 space X is weakly m - n compact if and only if for each $\langle n$ -discrete T_2 space Y and for each continuous

map $f: X \rightarrow Y$, $\text{Cl}_Y f(X) - f(X)$ has no points of local character $\leq m$, where n is regular and $m^n = m$.

Proof. Necessity follows from proposition B. To prove sufficiency, suppose X is not weakly m - n compact. Then by proposition A we have a space Y which is $\langle n$ -discrete and T_2 while $\text{Cl}_Y f(X) - f(X)$ contains a point of local character $\leq m$. Hence we have a contradiction.

The following special case of C is of interest:

D. Corollary. A topological space X is feebly compact if and only if for each T_2 -space Y and for each continuous map $f: X \rightarrow Y$, $\text{Cl}_Y f(X) - f(X)$ is nowhere first countable in Y .

3. Products of Two Spaces

3.1. Machinery

We extend the method of Vaughan [73] to the setting of weakly m - n compact spaces.

A. Definition. A space X is said to satisfy the property $\bar{I}_{m,n}$ if and only if for every strong m - n filter base F on X , there exists a compact subset K of X and a strong m - n stable filter base G such that $G \supset F$ and $G \cap U_K$ where U_K is an open neighbourhood base of K .

B. Proposition.

(i) Let X be a space which satisfies the property $\bar{I}_{m,n}$. Then X is weakly m - n compact.

(ii) Let n be a regular cardinal and m^n a G -space. If X is weakly m - n compact, and Y is a space which satisfies the property $\bar{I}_{m,n}$.

Proof. (i) Let F be a strong m - n filter base on X . Then since X satisfies the property $\bar{I}_{m,n}$ there is a strong m - n stable filter base G and a compact subset K of X such that $G \supset F$ and $G \cap U_K$. Hence $\bar{a}F \supset \bar{a}G$ and $\bar{a}G \cap \bar{a}U_K$. Therefore $\bar{a}X$ is weakly m - n compact.

(ii) Let F be a strong m - n filter base on X . Then since n is regular and m^n is a G -space, there exists a strong m - n stable filter base G such that $F \supset G$. Since X is weakly m - n compact, there

exists some $x \in \bar{F}'$; let \mathcal{V}_x be an open neighbourhood base at x with $|\mathcal{V}_x| \leq m$. Take $K = \{x\}$ and $G = F' \vee \mathcal{V}_x$; then we note that G is a strong m - n stable filter base on X , $G > F' > F$ and $G > \mathcal{V}_x$. Therefore X satisfies the property $\bar{I}_{m,n}$.

C. Proposition. Let n be regular and $m^n = m$. Then every locally compact weakly m - n compact space X satisfies the property $\bar{I}_{m,n}$.

Proof. Let F be a strong m - n filter base on X . Since n is regular and $m^n = m$, there exists a strong m - n stable filter base F' such that $F' > F$ and since X is weakly m - n compact, there exists a $x \in \bar{F}'$. Let K be a compact neighbourhood of x ; then $F'|K = (F \cap K) \vee F'$ is a strong m - n stable filter base on X and $F'|K > F \cap K$ and $F'|K > \mathcal{V}_y$. Hence X satisfies the property $\bar{I}_{m,n}$.

D. Proposition. Let $f: X \rightarrow Y$ be a bicontinuous (open and continuous) onto map. If X satisfies the property $\bar{I}_{m,n}$ then Y satisfies the property $\bar{I}_{m,n}$.

Proof. Let F be a strong m - n filter base on Y . Then since f is continuous, $f^{-1}(F) = (f^{-1}(F) \vee F)$ is a strong m - n filter base on X and since X satisfies the property $\bar{I}_{m,n}$, there exists a strong m - n stable filter base G on X and a compact subset K of X such that $G > f^{-1}(F)$ and $G > \mathcal{V}_y$. Then since f is open, $f(G)$ is a strong m - n stable filter base on Y and since f is onto $f(G) > F$ and $f(G) > \mathcal{V}_y$. Since f is continuous, $f(K)$ is a compact subset of Y and $f(G) > \mathcal{V}_y$. Hence Y satisfies the property $\bar{I}_{m,n}$.

E. Corollary. Let $X = \prod\{X_i : i \in I\}$ with the product topology. Suppose X satisfies the property $\bar{I}_{m,n}$; then every sub-product of X has the property $\bar{I}_{m,n}$.

Proof. Let $X_{I'} = \prod\{X_i : i \in I'\}$ where $I' \subseteq I$. Then since $\pi_{I'} : X \rightarrow X_{I'}$ is continuous, open and onto, $X_{I'}$ has the property

Products of Two Spaces

We now investigate the productivity of $\bar{I}_{m,n}$ via techniques of Cohen and others [1].

Lemma. Let $f : X \rightarrow Y$ be a continuous map. Let \mathcal{F} be an m, n -stable filter base on X and let \mathcal{H} be an m, n -stable filter base on Y . Let \mathcal{G} be the filter base on X consisting of all $G \in \mathcal{F}$ such that $f(G) \in \mathcal{H}$.

Proof. We show that \mathcal{G} is m, n -stable. Let $\{G_i : i \in I\} \in \mathcal{G}$. Then $\{G_i : i \in I\} \in \mathcal{F}$ and $\{f(G_i) : i \in I\} \in \mathcal{H}$. Since \mathcal{F} is m, n -stable, $\bigcap_{i \in I} G_i \in \mathcal{F}$ and $f(\bigcap_{i \in I} G_i) \in \mathcal{H}$. Thus $\bigcap_{i \in I} G_i \in \mathcal{G}$. Similarly, \mathcal{G} is n -stable since $\bigcup_{i \in I} G_i \in \mathcal{F}$ and $f(\bigcup_{i \in I} G_i) \in \mathcal{H}$.

Therefore, \mathcal{G} is m, n -stable. This completes the proof.

and hence $\mathcal{F} \vee f^{-1}(\mathcal{G})$ is a strong m - n stable filter base on X .

R. Theorem. Let n be a regular cardinal and $m^n = m$. Let X be a space which satisfies the property $\bar{I}_{m,n}^+$. If Y is weakly m - n compact, then $X \times Y$ is weakly m - n compact.

Proof. Let \mathcal{F} be a strong m - n stable filter base on $X \times Y$. Then, since $\pi_1: X \times Y \rightarrow X$ is an open map, $\pi_1(\mathcal{F})$ is a strong m - n stable filter base on X and since X satisfies the property $\bar{I}_{m,n}^+$, there exists a strong m - n stable filter base \mathcal{G} on X and a compact subset K of X such that $\mathcal{G} \subseteq \mathcal{F}$ and $\mathcal{G} \subseteq \pi_1^{-1}(K)$. Let $\mathcal{H} = \mathcal{F} \setminus \pi_1^{-1}(K)$, then by the lemma \mathcal{H} is a strong m - n stable filter base on $X \times Y$. Again, since $\pi_2: X \times Y \rightarrow Y$ is an open map, $\pi_2(\mathcal{H})$ is a strong m - n stable filter base on Y and since Y is weakly m - n compact, $\overline{\pi_2(\mathcal{H})} \neq \emptyset$.

Let $y \in \overline{\pi_2(\mathcal{H})}$ and let W be a closed neighbourhood of y . Then we have $\mathcal{H} \cap \pi_2^{-1}(W) \neq \emptyset$ for every $\mathcal{F} \in \mathcal{F}$. Hence $\mathcal{H} \cap \pi_2^{-1}(W)$ is a strong m - n stable filter base on $X \times W$ and $X \times W$ is weakly m - n compact.

Lemma. Let X be a topological space and Y a compact space. If X has character $\leq n$ and Y is weakly m - n compact, then $X \times Y$ is weakly m - n compact.

Proof. Let \mathcal{F} be a strong m - n stable filter base on $X \times Y$ and let \mathcal{G} be a strong m - n stable filter base on X such that $\mathcal{G} \subseteq \mathcal{F}$. Then \mathcal{G} is a strong m - n stable filter base on X and X has character $\leq n$. Hence X is weakly m - n compact. Let K be a compact subset of X such that $\mathcal{G} \subseteq \pi_1^{-1}(K)$. Then $\mathcal{F} \cap \pi_1^{-1}(K)$ is a strong m - n stable filter base on $X \times K$ and $X \times K$ is compact. Hence $\mathcal{F} \cap \pi_1^{-1}(K)$ is a strong m - n stable filter base on $X \times Y$ and $X \times Y$ is weakly m - n compact.

Theorem. Let m be a regular cardinal and $m^{\aleph_1} = m$. Let X and Y be two spaces which satisfy the property $\bar{I}_{m,n}$. Then $X \times Y$ satisfies the property $\bar{I}_{m,n}$.

Proof. Let \mathcal{F} be a strong m - n stable filter base on $X \times Y$. The since X satisfies the property $\bar{I}_{m,n}$, there exists a strong m - n stable filter base \mathcal{G}_1 on X and a compact subset K_1 of X such that $\mathcal{G}_1 \cap \pi_1^{-1}(K_1) = \mathcal{F}$. Then by the lemma A, $\mathcal{G}_1 \cap \pi_1^{-1}(K_1)$ is a strong m - n stable filter base on $X \times Y$. again, since Y satisfies $\bar{I}_{m,n}$, there exists a strong m - n stable filter base \mathcal{G}_2 on Y and a compact subset K_2 of Y such that $\mathcal{G}_2 \cap \pi_2^{-1}(K_2) = \mathcal{G}_1 \cap \pi_1^{-1}(K_1)$. Then by lemma A, $\mathcal{G}_2 \cap \pi_2^{-1}(K_2)$ is a strong m - n stable filter base on $X \times Y$. \square

As a consequence, let \aleph_1 be a regular cardinal with $\aleph_1^{\aleph_1} = \aleph_1$. Then $X \times Y$ satisfies the property $\bar{I}_{\aleph_1, \aleph_1}$. \square

property $\bar{I}_{m,n}$, there exists a strong m - n stable filter base G_1 on X_1 and a compact subset $K_1 \subset X_1$ such that $G_1 \supset \pi_1(F)$ and $G_1 \supset U_{K_1}$. Then by Lemma 2 A, $H_1 = F \vee \pi_1^{-1}(G_1)$ is a strong m - n stable filter base on X .

Now inductively assume that for each $i \leq k$, a compact $K_i \subset X_i$ and a strong m - n stable filter base G_i on X_i have been found such that

- (i) $G_i \supset U_{K_i}$ and
- (ii) $H_k = F \vee [\vee \{\pi_i^{-1}(G_i) : i = 1, 2, \dots, k\}]$ is a strong m - n stable filter base on X .

Since H_{k+1} satisfies the property $\bar{I}_{m,n}$, it is clear that there exists a

strong m - n stable filter base G_{k+1} on X_{k+1} and a compact subset $K_{k+1} \subset X_{k+1}$ such that

- (i) $G_{k+1} \supset U_{K_{k+1}}$ and
 - (ii) $H_{k+1} = H_k \vee \pi_{k+1}^{-1}(G_{k+1})$ is a strong m - n stable filter base on X .
- By the inductive definition of K_i and H_i for $i = 1, 2, \dots$

$$K = \prod \{K_i : i = 1, 2, \dots\}$$

$$H = F \vee [\vee \{\pi_i^{-1}(G_i) : i = 1, 2, \dots\}]$$

It is easy to see that K is compact and H is a filter base on X .

Let $\bar{H} = \overline{H \cap K}$. Then

$$\bar{H} = \overline{H \cap K} = (H \cap K) \cap \bar{K}$$

and since K is compact, it follows that $(H \cap K) \cap \bar{K} \neq \emptyset$. Thus \bar{H} has

nonempty elements and by 2.1.1 X is weakly m - n compact.

Corollary. Let $X = \prod \{X_j : j \in J\}$ and let n be a regular cardinal with $n < \omega$. Suppose each X_j is weakly m - n compact, if

...

4. Large Products

In general dense subspaces will not inherit the property weakly Lindelöf (relatively weakly Lindelöf). In this section we consider the following problem in a more general setting:

Let $X = \prod(X_i : i \in I)$. Let γ be an infinite cardinal. Is the property weakly Lindelöf in $\gamma(X_i)$ determined by finite sub-products of $\prod X_i$?

4.1 Machinery

We shall establish two special cases of the main theorem by considering γ weak topological sums of $X = \prod(X_i : i \in I)$.

A Definition. Let \mathcal{B} be a base for a topological space X . It is said that \mathcal{B} is weakly γ -compact if and only if for every cover \mathcal{U} of X with $|\mathcal{U}| \leq \gamma$ there is a finite subfamily \mathcal{U}_0 of \mathcal{U} such that $\bar{\mathcal{U}}_0 = X$.

We state the following facts about the notion of weakly γ -compactness. (i) If X is weakly γ -compact then every dense subspace of X is weakly γ -compact. (ii) If X is weakly γ -compact then every closed subspace of X is weakly γ -compact.

(iii) If X is weakly γ -compact then every dense subspace of X is weakly γ -compact.

B Proposition. Let $X = \prod(X_i : i \in I)$ and let \mathcal{B} be a base for X . Then \mathcal{B} is weakly γ -compact if and only if for every finite sub-product $\prod_{i \in J} X_i$ the base \mathcal{B}_J is weakly γ -compact.

Proof. (i) If γ is regular, then $\bar{\gamma} = \gamma$ and hence $\bar{\gamma} \leq n$.

(ii) If γ is singular, then since n is regular, we have $\gamma < n$. Hence $\bar{\gamma} = \gamma^+ \leq n$.

Remark $\bar{\gamma}$ in the proposition B is a regular cardinal.

Notation. Let I be any indexing set. Then we denote,

$$(i) \quad |I|^{\gamma} = \sum \{|I|^k : k < \gamma\} \quad \text{and}$$

$$(ii) \quad P_{\gamma}(I) = \{I' \subset I : |I'| < \gamma\}$$

Lemma. Let $X = \prod\{X_i : i \in I\}$ and let $m > n \geq \text{cf}(n) > |I|^{\gamma} \geq \gamma \geq 1 > \aleph_0$. Suppose $(X_{I'},)_k$ is weakly m - n compact for all $I' \in P_{\gamma}(I)$. Then $(\prod X_i)_k$ is weakly m - n compact relative to $(\prod X_i)_k$.

Proof. We note that $\gamma(\prod X_i) = \cup\{X(I') : I' \in P_{\gamma}(I)\}$ and since $\gamma(I')$ is homeomorphic to $(X_{I'})_k$, $\gamma(\prod X_i)$ is the $|I|^{\gamma}$ fold union of weakly m - n compact subspaces of $(\prod X_i)_k$. Hence we have the lemma by 1.1 E.

Let $W = \prod\{W_i : i \in I\}$ where each W_i is open in X_i and $|P(W)| = 1$ form the canonical basis for $(\prod X_i)_k$. Let $A \subset \prod\{X_i : i \in I\}$; then the canonical basis for A consists of all sets of the form $A \cap W$ with $|P(W)| = 1$. In this terminology we rephrase the lemma E as follows:

Let $X = \prod\{X_i : i \in I\}$ and let $m > n \geq \text{cf}(n) > |I|^{\gamma} > \gamma > k \geq \aleph_0$.

Suppose $(X_{I'})_k$ is weakly m - n compact with respect to its canonical basis for all $I' \in P(I)$. Then $\gamma(\prod X_i)$ is weakly m - n compact with respect to its canonical basis.

G. Theorem. Let $X = \prod\{X_i : i \in I\}$ and let $m \geq n \geq |I| \geq \gamma \geq k \geq \aleph_0$. Suppose n is regular and strongly γ -inaccessible and suppose $(X_{I'})_k$ is weakly m - n compact for all $I' \in P(I)$. Then $\gamma(\prod X_i)$ is weakly m - n compact relative to $(\prod X_i)_k$.

Proof. Consider $|I|^\gamma = \sum(|I|^k \cdot \gamma) \geq |I| \geq \gamma$ and since n is regular and strongly γ -inaccessible we have $n = cf(n)$ and $|I|^\gamma < n$. Hence $m > n = cf(n) > |I|^\gamma \geq \gamma > k \geq \aleph_0$ and therefore we can apply lemma E to obtain the theorem.

In the above theorem there is a restriction on the cardinality of I and we wish to remove this condition in the next theorem.

4.2 Main Theorem

To establish the main theorem about products of weakly m - n compact spaces, we employ the following lemmas (see [13] for proof in [13]).

A. Theorem. Let $m \geq n \geq \gamma \geq k \geq \aleph_0$ and let n be regular and strongly γ -inaccessible. Let $X = \prod\{X_i : i \in I\}$ and let $U \in \mathcal{C}(X) = \prod\{U_i : i \in I\}$ each U_i is open in X_i and $|R(U_i)| \leq 1$ where $|U_i| \leq m$. Suppose $(X_{I'})_k$ is weakly m - n compact for all $I' \in P(I)$ and $\gamma(\prod X_i) \in U$. Then there exists $\gamma \in U$ such that $|U| \leq m$ and $\gamma(U) \in U$.

Proof. Let $\bar{\gamma} = \gamma$ if γ is regular and $\bar{\gamma} = \gamma^+$ if γ is singular. Then $\bar{\gamma}$ is regular and $\bar{\gamma} \leq n$. By theorem 1-G, $\gamma(X_{I'})$ is weakly m - n compact relative to $(X_{I'})_k$, for all $I' \in P_{<\bar{\gamma}}(I)$. We note that $\pi_{I'}(\gamma(\Pi X_{I'})) = \gamma(X_{I'})$ and hence $\{\pi_{I'}(U) : U \in U\}$ is an m -fold open cover of $\gamma(X_{I'})$ where $I' \in I$. Let $I' \in P_{<\bar{\gamma}}(I)$; then there exists a $U_{I'} \in U$ such that $|U_{I'}| < n$ and

$$\gamma(X_{I'}) \subseteq \text{Cl}_{(X_{I'})_k}(\cup\{\pi_{I'}(U) : U \in U_{I'}\}). \quad (1)$$

Let $I_1 \in I$ and $|I_1| < n$ and let $F_1 = \{U_{I'} : I' \in P_{<\bar{\gamma}}(I_1)\}$ and U_{I_1} has the property (1). Let $I_2 = I_1 \cup R(F_1)$ where $R(F_1) = \cup\{P(U) : U \in U_{I'}, \text{ and } U_{I'} \in F_1\}$. Trivially $R(F_1) \subset I$ and therefore $I_2 \in I$.

We note the following:

- (i) $|R(U)| < k \leq n$.
- (ii) $U_{I'} \in U$, $|U_{I'}| < n$ for all $I' \in P_{<\bar{\gamma}}(I_1)$.
- (iii) $|P_{<\bar{\gamma}}(I_1)| < |I_1|^{\bar{\gamma}}$ if γ is regular and $|P_{<\bar{\gamma}}(I_1)| \leq |I_1|^{\bar{\gamma}^+}$ if γ is singular.

Since n is strongly γ -inaccessible, $|P_{<\bar{\gamma}}(I_1)| < n$ for all $\gamma \leq n$.

Therefore we have the following:

- (i) $|R(F_1)| < n$.
- (ii) $|I_2| < n$.

Inductively we define $I_\alpha = \cup\{U_{I'} : I' \in P_{<\bar{\gamma}}(I)\}$ and $I_{\alpha+1} = I_\alpha \cup R(F_\alpha)$ for $\alpha < \bar{\gamma}$. Let $I' = \cup\{I_\alpha : \alpha < \bar{\gamma}\}$ and $U' = \cup\{F_\alpha : \alpha < \bar{\gamma}\}$. Since n is regular and $|I_\alpha| < n$ for all $\alpha < \bar{\gamma}$ we have

$$\begin{aligned} |I^*| &< n \quad \text{and} \\ |U^*| &< n. \end{aligned} \tag{3}$$

Each $U_{I^*} \in U$ and therefore each $F_\alpha \in U$ and hence $U^* \in U$. We shall prove that

$$\gamma(\pi X_i) \subseteq \overline{U^*}.$$

Let $x \in \gamma(\pi X_i)$ and let $V = \prod \{V_i : i \in I\}$ be a basic open neighbourhood of x in $(\pi X_i)_k$. Then we have $|R(V)| < k \leq \gamma \leq \bar{\gamma}$ and hence there exists an $\alpha < \bar{\gamma}$ such that

$$R(V) \cap I^* = R(V) \cap I_\alpha. \tag{4}$$

Let $H = R(V) \cap I_\alpha$; then $H \cap I_\alpha \in I$ and $|H| < k < \bar{\gamma}$ and, by (1), there exists a $U \in \mathcal{U}_H \cap F_\alpha$ such that

$$\pi_H(U) \cap \pi_H(V) \neq \emptyset. \tag{5}$$

Since $U \in F_\alpha$, $R(U) \in I_{\alpha+1}$ and, by (1),

$$\begin{aligned} R(U) \cap R(V) &= (R(U) \cap I_{\alpha+1}) \cap R(V) \\ &= R(U) \cap (I_{\alpha+1} \cap I^*) \cap R(V) \\ &= R(U) \cap (R(V) \cap I^*) \cap I_{\alpha+1} \\ &= R(U) \cap R(V) \cap I_\alpha \\ &\subseteq H \end{aligned} \tag{6}$$

Now $U \cap V \neq \emptyset$ and therefore $V \cap (U^*) \neq \emptyset$. This is true for every neighbourhood V of x and therefore we have $\gamma(\pi X_i) \subseteq \overline{U^*}$.

For $k \leq \gamma$, $\gamma(\prod X_i)$ is a dense subspace of $(\prod X_i)_k$ and we are ready to give the main theorem.

B. Theorem. Let $m \geq n > \gamma \geq k \geq \aleph_0$ and let n be regular and strongly γ -inaccessible. Let $X = \prod\{X_i : i \in I\}$ and suppose $(X_{I'})_k$ is weakly m - n compact for all $I' \in P_{<\gamma}(I)$. Let B be the canonical base for the product space $(\prod X_i)_k$. Then $(\prod X_i)_k$ is B -weakly m - n compact.

If n is strongly γ -inaccessible and if $d(X_i) < n$ for each $i \in I$, then each X_i has a dense subset A_i with $|A_i| < n$. Hence $A_{I'}$ is a dense subset of $(X_{I'})_k$ and $|A_{I'}| < n$ for all $I' \in P_{\leq \gamma}(I)$ where $\gamma \leq n$ and $A = \bigcup\{A_i : i \in I\}$. Therefore $d(X_{I'}) < n$ and hence $(X_{I'})_k$ is weakly ∞ - n compact for all $I' \in P_{\leq \gamma}(I)$. Thus we have the following:

C. Corollary. Let $X = \prod\{X_i : i \in I\}$, let $n \geq \gamma \geq k \geq \aleph_0$ and suppose n is regular and strongly γ -inaccessible. If $d(X_i) < n$ for all $i \in I$, then $(\prod X_i)_k$ is weakly ∞ - n compact.

4.3. Weakly ∞ - n Compact Spaces

We recall that B -weak ∞ - n compactness is equivalent to weak ∞ - n compactness and hence we obtain product theorems for weakly ∞ - n compact spaces as special cases of theorem 2-B (see [76]).

A. Theorem. Let $X = \prod\{X_i : i \in I\}$ and let $n > \gamma \geq k \geq \aleph_0$. Suppose n is regular and strongly γ -inaccessible. Then $(\prod X_i)_k$ is weakly ∞ - n compact if and only if $(X_{I'})_k$ is weakly ∞ - n compact for all $I' \in P_{\leq \gamma}(I)$.

B. Corollary. Let $X = \prod\{X_i : i \in I\}$ with the usual product topology and let n be a regular cardinal. Then X is weakly ω - n compact if and only if every finite sub-product of X is weakly ω - n compact.

Proof. Necessity follows from the fact that $\pi_{I'} : X \rightarrow X_{I'}$ is continuous for every $I' \subset I$.

On the other hand, note that regular cardinals are infinite and every infinite cardinal is strongly \aleph_0 -inaccessible. Hence taking $\gamma = k = \aleph_0$ in theorem A, we obtain sufficiency.

Let $X = \prod\{X_i : i \in I\}$. Then, according to theorem 2.3 A, if each X_i is weakly ω - n compact and κ - n -discrete (or each X_i is weakly ω - n compact and locally compact) where n is regular, then $X_{I'}$ is weakly ω - n compact, for all $I' \in P_{<\aleph_1}(I)$.

Now we are ready to produce two theorems regarding weakly ω - n compact spaces.

C. Theorem. Let $X = \prod\{X_i : i \in I\}$ and let n be a regular cardinal. Suppose each X_i is κ - n -discrete. Then X is weakly ω - n compact if and only if each X_i is weakly ω - n compact.

D. Theorem. Let $X = \prod\{X_i : i \in I\}$ and let n be an uncountable regular cardinal. Suppose each X_i is locally compact. Then X is weakly ω - n compact if and only if each X_i is weakly ω - n compact.

Since \aleph_0 and \aleph_1 are regular cardinals taking $n = \aleph_0$ and $n = \aleph_1$ respectively in C, yield the following results.

E. Corollary.

(i) A non-empty product is H-closed if and only if each factor space is H-closed (see [39]).

(ii) A non-empty product of P-spaces is weakly Lindelöf if and only if each factor space is weakly Lindelöf.

F. Note. Let Y be the weakly Lindelöf P-space constructed in 1.2 G.

Then Y^k ($k > \mathfrak{C}$) has the following properties:

- (i) weakly Lindelöf and T_2 ,
- (ii) non-separable,
- (iii) Non-cellular,
- (iv) non normal.

Taking $n = \aleph_1$ in theorem D. we obtain the following:

G. Theorem Let $X = \prod\{X_i : i \in I\}$. Suppose each X_i is locally compact. Then X is weakly Lindelöf if and only if each X_i is weakly Lindelöf.

CHAPTER IV: CARDINAL INVARIANTS

1. The Almost Lindelöf Number

1.1. Basic Results

In this section, we will introduce a new cardinal function, the almost Lindelöf number $al(X)$ of a topological space X .

A. Definition. A subset E of a topological space X is said to be almost k -Lindelöf if and only if every open cover \mathcal{U} of E has a subsystem \mathcal{U}' of cardinality $< k$ with $E \subset \bigcup \{U \in \mathcal{U}'\}$. We may define,

$$al(E, X) = \min\{k : E \text{ is almost } k\text{-Lindelöf}\}$$

and define the almost Lindelöf number $al(X)$ of X as:

$$al(X) = \sup\{al(E, X) : E \text{ is a closed subset of } X\}$$

B. Definition. A topological space X is said to be weakly k -Lindelöf if and only if every open cover \mathcal{U} of X has a subsystem \mathcal{U}' of cardinality $< k$ with $X = \overline{\bigcup \{U \in \mathcal{U}'\}}$. Note that this is equivalent to saying that X is weakly $< k$ -compact.

We define the weak Lindelöf number $wl(X)$ of X as:

$$wl(X) = \min\{k : X \text{ is weakly } k\text{-Lindelöf}\}$$

C. Remark. For any topological space X , $al(X) \leq wl(X)$ and $al(X) = 1$ if and only if X is regular and $wl(X) = 1$.

We next present examples which show these cardinal functions will in general, differ.

D. Example. Let X be the closed upper half plane in \mathbb{R}^2 and let $E \subset X$ be the x-axis. The basic neighbourhoods of $x \in X - E$ will be as usual in \mathbb{R}^2 , while the basic neighbourhoods of points $Z \in E$ will take the form

$$V_\epsilon = \{x \in X - E : \|x - Z\| < \epsilon\} \cup \{Z\}.$$

Call this space Y .

Easily, Y is a $T_{2\frac{1}{2}}$ (and thus T_2) space in which E is a discrete closed subspace of cardinality \mathfrak{C} . It follows that $l(Y) = \mathfrak{C}$. To see that $al(Y) = \aleph_0$, it will be enough to note that $al(B, Y) = \aleph_0$ for any $B \subset E$, since the open upper half plane in Y is hereditarily Lindelöf and whenever A and B are disjoint in Y , $al(A, Y) + al(B, Y) \geq al(A \cup B, Y)$. For if $\{U_i\}$ is an open cover of $B \subset E$ and if R denotes the rest of Y , then

$$al(B, Y) + al(R, Y) \geq al(B \cup R, Y) = al(Y).$$

is an open cover of B in \mathbb{R}^2 , which is hereditarily Lindelöf.

Thus in non-regular spaces, we may find $al(Y) < l(X)$.

Now let V be the T_2 space constructed in 7.1.2 C. For this space

$$al(V) = \aleph_0 = wl(V).$$

Thus we see that, in general, the cardinal functions $wl(X)$, $al(X)$ and

$\psi(X)$ are distinct in T_2 spaces.

For notational convenience we use $\bar{\psi}(X)$ for $\psi_C(X)$ (page 8 (41))

E. Lemma. Let X be a T_2 space. Then $|X| \leq d(X) \cdot \bar{\psi}(X)$

Proof. Let B be a dense subset of X such that $|B| \leq d(X)$.
 $k = \bar{\psi}(X) = \psi(X) \cdot \bar{\psi}(X)$. Since $\bar{\psi}(X) = \psi(\bar{B})$ for $\bar{B} = \bar{B}$,
 $2 \cdot \bar{\psi}(X) \leq |B| \cdot \bar{\psi}(X)$ follows that

$$\begin{aligned} \bar{\psi}(X) &\leq d(X) \cdot \bar{\psi}(X) \\ &\leq d(X) \cdot \bar{\psi}(X) \\ &\leq d(X) \cdot \bar{\psi}(X) \end{aligned}$$

$$\begin{aligned} |X| &\leq d(X) \cdot \bar{\psi}(X) \\ &\leq d(X) \cdot \bar{\psi}(X) \\ &\leq d(X) \cdot \bar{\psi}(X) \end{aligned}$$

F. Theorem Let X be a T_2 space. Then $|X| \leq d(X) \cdot \bar{\psi}(X)$

Proof. Let $k = \bar{\psi}(X) = \psi(X) \cdot \bar{\psi}(X)$. For each $\alpha < k$,
 α neighborhood of x such that $|U_\alpha| \leq d(X)$.
 $\bar{\psi}(X) = \psi(\bar{B})$ for $\bar{B} = \bar{B}$,
 $2 \cdot \bar{\psi}(X) \leq |B| \cdot \bar{\psi}(X)$ follows that

...

$$U \in \mathcal{U}\{O_x : x \in \cup\{F_\alpha : \alpha < \xi\}\}$$

where $|U| \leq k$ and $X - \cup U \neq \emptyset$, then $F_\xi = \cup U \neq \emptyset$. Now let $F_1 = \{p\}$, where p is an arbitrary point of X . Suppose we have defined F_α , for $\alpha < \xi$, with $|F_\alpha| < 2^k$. Let

$$O = \mathcal{U}\{O_x : x \in \cup_{\alpha < \xi} F_\alpha\}$$

and let $O^* = \{X - \cup U : U \in O \text{ and } |U| < k\}$. We select one point from each non-empty set in O^* and form the set F_ξ .

We take $F_\xi = \overline{\cup\{U : U \in O \text{ and } |U| < k\}}$. Then F_ξ is closed and by Lemma 1.1.1. Now we note that $X = \cup\{F_\xi : \xi < k^+\}$ and thus

Corollary (Engel'skiĭ, 1950). Let X be a T_2 space. Then

Let X be a T_2 -space. Then $\pi(X) = \psi(X)$.

Let \mathcal{A} be a local base of X such that $|\mathcal{A}| \leq d(X)$ and for each $A \in \mathcal{A}$ there is a local base at p with $|U_p| \leq \pi(X)$. Let \mathcal{B} be a local base at p for each $p \in X$ and let

$$\mathcal{C} = \{U \cap V : U \in \mathcal{A}, V \in \mathcal{B}\}$$

Then \mathcal{C} is a local base at p for each $p \in X$. The cardinality of neighbour-

$$|\mathcal{C}| \leq |\mathcal{A}| \cdot |\mathcal{B}|$$

$$\leq d(X) \cdot \pi(X)$$

Then for each $K \in K_p$, $p \in \overline{K \cap B_p}$. Hence

$$\{p\} = \bigcap \{ \overline{K \cap B_p} : K \in K_p \}.$$

Now the map $p \rightarrow \{K \cap B_p : K \in K_p\}$ from X to the collection $\{[B]_{\leq k}^{\leq k}\}$ is one to one. Thus,

$$\begin{aligned} |X| &\leq (d(X)^k)^k \\ &= d(X)^{\pi_X(X) \bar{\nu}(X)} \end{aligned}$$

I. Corollary.

(i) Let X be a T_2 -space. Then $|X| < d(X)^{\chi(X)}$ (see 5 of [34]).

(ii) Let Y be a T_3 space. Then $|Y| < d(Y)^{\chi(Y) \bar{\nu}(Y)}$

1.2 Main Results

We shall show that for a T_2 space X , $|X| < 2^{\text{af}(X) \chi(X)}$ which improves the celebrated theorem (7.6) of A.V. Arhangel'skiĭ (1970).

First we state a preliminary theoretical result.

Proposition. Let X be a set such that $|X| > k$, $\lambda > k$ and $k < \lambda$. Let $G: P_{\leq \lambda}(X) \rightarrow P_{\leq k}(X)$ be a set mapping. Then there exists $A \subseteq X$ such that $|A| = k$ and $A \subseteq G(B)$ for every $B \subseteq X$ with $|B| \leq \lambda$.

Proof. We note that

$$k^{\lambda^+} = \sum_{|B| \leq \lambda^+} (|B| \leq \lambda^+)$$

Now we apply 2.24 of [34].

B. Theorem. Let X be a T_2 -space. Then $|X| \leq 2^{\alpha L(X) \bar{\psi}(X) \pi \chi(X) \partial(X)}$.

Proof. Let $\beta = \alpha L(X) \bar{\psi}(X) \pi \chi(X) \partial(X)$, and let $k = 2^\beta$. Let \mathcal{O}_x be a collection of open neighbourhood of x such that $|\mathcal{O}_x| \leq \beta$ and $\{x\} = \cap \{\bar{V} : V \in \mathcal{O}_x\}$. We shall write, $U_A = \cup \{\mathcal{O}_x : x \in A\}$ and let G be a set mapping,

$$P_{\leq \beta}(X) \rightarrow P_{\leq k}(X).$$

Let $\Lambda \in P_{\leq \beta}(X)$; then we set

$$V_A = (\cup \{P_{\leq \beta}(U) : U \in \Lambda\}) \cdot X - \cup \{\bar{U} : U \in U\} \neq \emptyset.$$

Now it is clear that $|V_A| \leq 2^\beta = k$. For each $U \in V_A$, we select $p(U) \in X - \cup \{\bar{U} : U \in U\}$ and we write $G(\Lambda) = \overline{\cup \{p(U) : U \in V_A\}}$.

$$|G(\Lambda)| \leq (2^k)^k$$

$$= 2^\beta$$

k .

Also we note that $G(\Lambda) \supseteq \bar{A}$ for every $\Lambda \in P_{\leq \beta}(X)$.

We now apply Proposition A to obtain a set $B \subseteq X$ with $|B| = k$ and $B \supseteq G(\Lambda)$ for every $\Lambda \in P_{\leq \beta}(B)$. We claim that $X = B$.

First, since $\partial(X) \leq \beta$, it follows that B is a closed subset of X . Now suppose $X - B \neq \emptyset$, say $q \in X - B$. For each $y \in B$, we select $\mathcal{O}_y \in \mathcal{O}$ such that $q \notin \bar{\mathcal{O}_y}$ and since $\alpha L(B \setminus X) < \beta$ there exists

$\gamma \in P_{\leq \beta}(B)$ such that $B \subseteq \cup\{\bar{V}_\gamma : \gamma \in Y\} \subseteq X - \{q\}$. Thus $U = \{V_\gamma : \gamma \in Y\} \in \mathcal{V}_Y$ and by the construction, $p(U) \in G(Y) \subseteq B$. But $p(U) \in X - \cup\{\bar{V}_\gamma : \gamma \in Y\} \subseteq X - B$. Hence, we have a contradiction.

C. Corollary. Let X be a T_2 space. Then $|X| \leq 2^{aL(X)\chi(X)}$.

Following the main lines of the proof of the theorem B, with a slight modification, we obtain the following result:

D. Theorem. Let X be a T_2 space. Then

$$|X| \leq 2^{aL^+(X)\bar{\psi}(X)}$$

We shall extend the above result, to obtain an upper bound for the cardinality of the family $K(X)$ of compact subsets of a T_2 -space X .

E. Definition. We denote the pseudocharacter of a subset C of X by $\psi(C, X)$. Then the compact pseudocharacter $\psi_K(X)$ of X is defined as

$$\psi_K(X) = \sup\{\psi(C, X) : C \text{ is a compact subset of } X\}.$$

It is clear that $\psi(X) \leq \psi_K(X)$ and we shall show that, $\psi_K(X) \leq aL^+(X)$.

F. Proposition. Let X be a T_2 space. Then

$$\psi_K(X) \leq aL^+(X).$$

Proof. Let $aL^+(X) = k$. Let C be a compact subset of X and let p be a point in C . Then we can find two disjoint open subsets U_p and V_p such that $p \in U_p$ and $C \subseteq V_p$. Thus $\bar{U}_p \cap C = \emptyset$. Now since

$aL(X-C, X) \leq k$, there exists a $B \in [X-C]^{\leq k}$ such that $X - C \subseteq \cup\{\bar{U}_p : p \in B\}$. Hence $C \supseteq \cap\{X - \bar{U}_p : p \in B\}$ and since $C \subseteq X - \bar{U}_p$ for every $p \in B$. We have $\psi(C, X) \leq k$. From this it follows that,

$$\psi_k(X) \leq k.$$

G. Theorem. Let X be a T_2 -space. Then $|K(X)| \leq 2^{aL^*(X)\bar{\psi}(X)}$.

Proof. Let $k = aL^*(X)\bar{\psi}(X)$. For any $\{p; q\} \in [X]^2$ with $p \neq q$, select disjoint open sets $U_{p,q}$ and $V_{p,q}$ such that $p \in U_{p,q}$ and $q \in V_{p,q}$. Let \mathcal{B} be the family of all finite intersections formed by sets of the form $V_{p,q}$. Then by theorem D, $|\mathcal{B}| \leq |X| \leq 2^k$. Let K be a compact subset of X and let $p \notin K$. Then there exists a $B_p \in \mathcal{B}$ such that $p \in B_p \subseteq \bar{B}_p \subseteq X - K$. Therefore, if F is a closed subset of $X - K$, then $\mathcal{G} = \{B_p : p \in F\}$ is an open cover of F and, since $aL(F, X) \leq k$, there exists a $\mathcal{G}' \in [\mathcal{G}]^{\leq k}$ such that

$$F \subseteq \cup\{\bar{B}_p : B_p \in \mathcal{G}'\} = A(\bar{\mathcal{G}}') \subseteq X - K.$$

Now we recall that $\psi_k(X) \leq k$, and hence $X - K = \cup\{F_\alpha : \alpha < k\}$, where each F_α is closed. Thus we can write $X - K = \cup\{A(\bar{\mathcal{G}}') : \alpha < k\}$ and since there are at most $2^k A(\bar{\mathcal{G}}')$ sets, it follows that

$$\begin{aligned} |K(X)| &\leq (2^k)^k \\ &= 2^k. \end{aligned}$$

H. Proposition. Let X be a T_2 -space. Then $\bar{\psi}(X) \leq L^*(X)$.

Proof. Let $x \in X$ and let \mathcal{V}_x be a collection of open neighbourhoods

of x such that $\{x\} = \cap\{\bar{V} : V \in \mathcal{V}_x\}$. Then there exists a sub-collection \mathcal{B}_x of \mathcal{V}_x such that $|\mathcal{B}_x| \leq L^*(X)$ and

$$X - \{x\} \subseteq \cup\{X - \bar{V} : V \in \mathcal{B}_x\}$$

Hence, $\{x\} = \cap\{\bar{V} : V \in \mathcal{B}_x\}$. Thus $\bar{\psi}(x, X) \leq L^*(X)$. Since $x \in X$ is arbitrary, it follows that

$$\bar{\psi}(X) \leq L^*(X).$$

I. Remark. We conclude this section by noting that, Theorem G simultaneously generalizes two important theorems 2.1 and 2.7 of [5].

Proof. We apply theorem G and proposition H.

In the next section we shall show that, the cardinal invariant $aL(X)$ is better than $L(X)$ in respect of estimations of the cardinality of X and the cardinality of $K(X)$, for T_2 -spaces.

1.3. Examples

We shall construct an example to show that there are T_2 -spaces where $aL(X)$ is relatively small compared to $L(X)$ and $c(X)$.

A. Method. Let T be the product of the k copies of the unit interval I . Then T has a closed nowhere dense subset E of cardinality 2^k . Let X be the set T with topology described as follows. The neighbourhoods of points $p \in T - E$ will be unchanged in X , while neighbourhoods of points $p \in E$ will take the form

$$U_p^* = (U - E) \cup \{p\},$$

where U is a neighbourhood of p in T .

Clearly, X is a T_2 -space and since E is a closed discrete subset of X of cardinality 2^k , $L(X) = 2^k$. We shall show that $aL^*(X) = k$.

B. Lemma. Let T and X be the spaces mentioned in A. Let U be an open subset in T with $p \in U \cap E$. Then $\text{Cl}_X U_p^* \supseteq U$.

Proof. Let $x \in U$. If $x \notin E$, then $x \in U_p^*$ and therefore we assume that $x \in U \cap E$. Let V_x^* be a neighbourhood of x in X . Then $V_x^* = (V - E) \cup \{x\}$ where V is open in T . Then U and V are both neighbourhoods of x in T and hence $U \cap V$ is a neighbourhood of x in T . Since E is nowhere dense, we must have $(U \cap V) - E \neq \emptyset$. Now, it follows that $V_x^* \cap U_p^* \neq \emptyset$. Thus $x \in \text{Cl}_X U_p^*$. This proves the lemma.

C. Proposition (5.3 of [34]). Let $R = \Pi\{R_i : i \in I\}$, where each R_i is a T_1 -space. Then

$$(i) \quad w(R) = |I| \cdot w_I(R)$$

$$(ii) \quad \psi(R) = |I| \cdot \psi_I(R)$$

Now, $L^*(T) \leq k$ and $\psi(T) = k$, where T is the space mentioned in A. Thus, using the space X of A, we establish the following:

D. Proposition. For each cardinal $k \geq \aleph_0$, there is a T_2 -space X with,

- (i) $aL^*(X) = k$
- (ii) $L(X) = 2^k$
- (iii) $\chi(X) = k$.

Proof. The space X constructed in A will serve. We note that,

$$|X| = 2^k; \text{ in fact according to 3-G, } |K(X)| = 2^k.$$

Our next aim is to show that there are T_2 -spaces X with $L(X)$ and $c(X)$ both larger than $aL(X)$.

E. Proposition. For each cardinal $k \geq \aleph_0$, there is a T_2 -space Y with,

- (i) $aL(Y) \leq k$
- (ii) $L(Y) = 2^k$
- (iii) $c(Y) = 2^k$

Proof. Let X_1 be the space constructed via T in A. Let X_2 be the Alexandroff double of T . Then we take Y to be the topological union of X_1 and X_2 . Since X_2 is an open subspace of Y and $c(X_2) = 2^k$, it follows that $Y = X_1 \cup X_2$ has the required properties.

It is also clear that $\chi(Y) = k$ and $|Y| = 2^k$.

2. The Weak Lindelöf Number

Our main objective is to investigate the following problem:

Let X be a T_3 -space. Let $wL(X)$ denote the weak Lindelöf number of X and let $\chi(X)$ denote the character of X . Is,

$$|X| \leq 2^{wL(X)\chi(X)} ?$$

2.1. Related Results

We shall define a new cardinal function, the quasi Lindelöf number, which we shall denote by $qL(X)$, and we show that $qL(X)$ is a common lower bound for $aL(X)$ and $c(X)$.

A. Definition. A subset E of X is said to be relatively weakly k -Lindelöf if and only if every X -open cover \mathcal{U} of E has a subsystem \mathcal{U}' of cardinality $\leq k$ with $E \subseteq \overline{\cup \mathcal{U}'}$.

Note that this is equivalent to saying that E is relatively weakly ∞ - k^+ compact.

B. Definition. We define the relative weak Lindelöf number $RwL(E)$ of $E \subset X$ as, $RwL(E) = \min\{k : E \text{ is relatively weakly } k\text{-Lindelöf}\}$. Now we define the quasi Lindelöf number $qL(X) = \sup\{RwL(E) : E \text{ is a closed subset of } X\} + \omega$.

For every closed subset E of X , $RwL(E) \leq aL(X)$. Thus it is clear that, $wL(X) \leq qL(X) \leq aL(X) \leq L(X)$.

By the next proposition it will follow that $qL(X) \leq c(X)$.

C. Proposition. Let X be any topological space. Then the following are equivalent:

- (i) $c(X) \leq n$,
- (ii) if U is a collection of open subsets of X such that $X = \overline{UU}$, then there exists a $U' \subseteq U$ such that $|U'| \leq n$ and $X = \overline{UU'}$,
- (iii) if U is any collection of open subsets of X , then there exists a $U' \subseteq U$ such that, $|U'| \leq n$ and $\overline{UU} \subseteq \overline{UU'}$.

Proof. See 3.2 of [9].

D. Example. Let X be the product of \aleph_1 copies of the natural numbers \mathbb{N} . Let Y be the Alexandroff double of the unit interval I . We take V as the topological sum $X \oplus Y$. Then V is a $T_{3\frac{1}{2}}$ space and by c, $qL(X) = c(X) = \aleph_0$ and since $qL(V) \leq qL(X) + qL(Y)$, we conclude

- (i) $qL(V) = \aleph_0$
- (ii) $L(V) = \aleph_1$.

Since, in normal spaces $qL(X) = wL(X)$ we state a refinement of Juhász's theorem 2.36 [34], in the following manner:

E. Theorem. Let X be a T_3 space. Then $|X| \leq 2^{\chi(X)qL(X)}$.

2.2. Π -Normal Spaces

We shall introduce a new class of spaces, the Π -normal (ΠN) spaces, and we prove that $|X| \leq 2^{wL(X) \chi(X)}$ for all $X \in \Pi N$ where $T_4 \not\subset \Pi N \not\subset T_3$. This extends the result 2.36 of [34].

A. Definition. Let $a \in X$. A local π -base at a in X is a family U_a of proper, non-empty open subsets of X such that every neighbourhood of a contains a member of U_a .

A normal local π -base is a local π -base with the property that the closures of the members are normal subspaces.

B. Definition. A T_3 -space X is called Π -normal (ΠN) if and only if X has a normal local π -base U_a for every $a \in D$, where D is some dense subset of X .

C. Note. The following classes of spaces belong to the class ΠN :

- (i) locally-metrizable and T_3 ,
- (ii) locally paracompact and T_3 ,
- (iii) locally normal and T_3 ,
- (iv) locally Lindelöf and T_3 .

In particular every T_4 -space belongs to the class ΠN .

D. Example. The Moore plane is a locally Lindelöf, T_3 -space and hence

it belongs to the class ΠN , but it is not a T_4 -space.

E. Definition. Let X be a topological space. Then we define,

$$\|U\| = \sup \{ |U| : U \in \mathcal{U} \},$$

$$A(a, X) = \min \{ k : a \text{ has a local } \pi\text{-base } \mathcal{U}_a \text{ with} \\ \|U_a\| \leq k \} + \aleph_0$$

$$A(D) = \sup \{ A(a, X) : a \in D \},$$

$$A(X) = \min \{ A(D) : \bar{D} = X \},$$

$$R(X) = \log A(X).$$

F. Theorem. Let X be a T_3 space. Then

$$|X| \leq 2^{wL(X) \chi(X) R(X)}$$

Proof. Let $\alpha = wL(X) \chi(X) R(X)$ and $k = 2^\alpha$. Suppose $|X| > k$. Let U_a be an open neighbourhood base at a with $|U_a| \leq \alpha$. Let $B \in P_{\leq \alpha}(X)$. Let $U_B = \cup \{ U_a : a \in B \}$ and let $V_B = \{ U \subseteq U_B : |U| \leq \alpha \text{ and } X - \overline{U} \neq \emptyset \}$. We note that $|V_B| \leq k$. Since $R(X) \leq \alpha$, there exists some $D \subset X$ such that $\bar{D} = X$ and $A(D) \leq k$. Hence for each $U \in V_B$ there exists a non-empty open subset $K(U)$ such that;

$$(i) \quad K(U) \subseteq X - \overline{U}$$

$$(ii) \quad |K(U)| \leq k.$$

We shall define $G: P_{\leq \alpha}(X) \rightarrow P_{\leq k}(X)$ by $G(B) = \overline{B \cup \{ K(U) : U \in V_B \}}$.

Then, there exists some $A \subset X$ such that $|A| = k$ and $A \supseteq G(B)$ for every $B \in P_{\leq \alpha}(A)$. We claim that, $X = \overline{A^0}$. Suppose $X - \overline{A^0} \neq \emptyset$.

Then since X is regular, there exists a point q and an open subset, U in X such that $q \in U \subseteq \bar{U} \subseteq X - \overline{A^0}$. Let $V = X - \bar{U}$. Then $\overline{A^0} \subseteq V$ and since $\text{RwL}(\overline{A^0}) \leq \alpha$, there exists a $B \in P_{<\alpha}(\overline{A^0})$ such that,

$$\begin{aligned} \overline{A^0} &\subseteq \overline{\cup\{V_a : V_a \in U_a, V_a \subseteq V \text{ and } a \in B\}} \\ &\subseteq \bar{V} \\ &\subseteq X - U. \end{aligned}$$

Now since $\partial(X) \leq \alpha$, A is a closed subset of X and hence $B \in P_{<\alpha}(A)$.

Let $U = \{V_a : V_a \in U_a, V_a \subseteq V \text{ and } a \in B\}$; then $U \in V_B$ and hence we have $K(U) \subseteq [G(B)]^0 \subseteq A^0 \subseteq \overline{A^0} \subseteq \overline{UU}$. But $K(U) \subseteq X - \overline{UU}$. This is a contradiction. Hence it follows that $|X| \leq k$.

G. Corollary. If $X \in \Pi N$, then we have $|X| \leq 2^{\text{wL}(X)\chi(X)}$.

Proof. By E of 2.2, we note that $R(X) \leq \text{wL}(X)\chi(X)$. Now by the theorem,

$$\begin{aligned} |X| &\leq 2^{\text{wL}(X)\chi(X), \text{wL}(X)\chi(X)} \\ &= 2^{\text{wL}(X)\chi(X)} \end{aligned}$$

2.3. Normal T_1 -Spaces

We shall define a new cardinal function $p\pi w(X)$ and we prove that for normal, T_1 -spaces $|X| \leq p\pi w(X)^{\text{wL}(X)\psi(X)\partial(X)}$. We also obtain an upper bound for the number of compact subsets $K(X)$ in a normal, space by proving,

$$|K(X)| \leq 2^{p\pi w(X)wL(X)}$$

Let B be a dense subset of X such that $|B| \leq d(X)$. Let $\mathcal{B} = \{U_p : p \in B\}$ where U_p is a local π -base at p . Then, clearly \mathcal{B} is a π -base for X .

We shall study special open covers. An open cover \mathcal{G} is said to be a strong open cover of X if it satisfies the following properties:

- (i) For distinct points x and y in X , there exists a $G \in \mathcal{G}$ such that $x \in G$ and $y \notin G$.
- (ii) If $B \in \mathcal{B}$ and $a \notin B$, then there exists a $G \in \mathcal{G}$ such that $a \in G$ and $G \cap B = \emptyset$, where \mathcal{B} is the π -base above.

A. Definition (Charlesworth [9]). We define

$$p\pi w(X) = \min \{k : X \text{ has a strong open cover } \mathcal{G} \text{ such that each point of } X \text{ is in at most } k \text{ members of } \mathcal{G}\} + \aleph_0$$

In a T_3 -space, $pws(X) \leq p\pi w(X) \leq pw(X)$.

B. Proposition. Let X be a set, let k be an infinite cardinal and suppose $\mathcal{G} \subseteq P(X)$ is such that each point of X is in at most k members of \mathcal{G} . If B is a subset of X , then the cardinality of the set of all finite minimal covers of B by elements of \mathcal{G} does not exceed k .

(This is Miscenko's lemma.)

C. Lemma. Let X be a T_3 -space. Then $|X| \leq d(X)^{p\pi w(X)wL(X)}$.

Proof. Let B be a dense subset of X such that $|B| \leq d(X)$. Let $k = \psi(X)\partial(X)$. Then, since $\partial(X) \leq k$, we can write $X = \cup\{\bar{T} : T \subseteq B, |T| \leq k\}$. But note that, $w(\bar{T}) \leq \rho(\bar{T})$ and by 2.2 of [9]

$$\begin{aligned} |\bar{T}| &\leq \rho(\bar{T})\psi(\bar{T}) \\ &\leq 2^{d(\bar{T})}\psi(\bar{T}) \\ &\leq 2^k. \end{aligned}$$

It follows that

$$\begin{aligned} |X| &\leq 2^k \cdot |B|^k \\ &\leq d(X)^k \\ &= d(X)\psi(X)\partial(X) \end{aligned}$$

D. Lemma. Let X be a normal, T_1 -space. Then $d(X) \leq p\pi w(X)^{wL(X)}$.

Proof. Let $k = p\pi w(X)$ and $\lambda = wL(X)$. We shall define, $G : [X]^{\leq k^\lambda} \rightarrow [X]^{k^\lambda}$. Let $A \in [X]^{\leq k^\lambda}$ and let G be a strong open cover of X such that each point of X is in at most k members of G .

We set $G_A = \{G \in G : G \cap A \neq \emptyset\}$ and $M_A = \{U \in [G_A]^{\leq \lambda} : X - \overline{UU} \neq \emptyset\}$.


Now for each $U \in M_A$, choose $p(U) \in X - \overline{UU}$ and set $G(A) =$

$\Lambda \cup \{p(U) : U \in M_A\}$. Since $|M_A| \leq k^\lambda$.

$$\begin{aligned} |G(A)| &\leq k^\lambda + k^\lambda \\ &= k^\lambda. \end{aligned}$$

Hence, $G(A) \in [X]^{k^\lambda}$ and this completes the construction of the set mapping G .

Now, by transfinite induction, we shall construct sets A_α for





Relations Between the Lindelöf Number

and the Weak Lindelöf Number

Let X be a topological space. The Lindelöf number $l(X)$ is the smallest cardinal κ such that every open cover of X has a subcover of cardinality at most κ .

The weak Lindelöf number $l_w(X)$ is the smallest cardinal κ such that every open cover of X has a subcover of cardinality at most κ which is point-finite.

It is well known that

$$l(X) \leq l_w(X) \leq 2^{l(X)}$$

1. Introduction

Let X be a topological space. The Lindelöf number $l(X)$ is the smallest cardinal κ such that every open cover of X has a subcover of cardinality at most κ .

The weak Lindelöf number $l_w(X)$ is the smallest cardinal κ such that every open cover of X has a subcover of cardinality at most κ which is point-finite.

It is well known that

$$l(X) \leq l_w(X) \leq 2^{l(X)}$$

and

if X is separable,

Definition A collection $\{U_\alpha\}_{\alpha \in I}$ of locally finite sets covers a topological space X if and only if $X = \bigcup_{\alpha \in I} U_\alpha$ and each U_α is a locally finite collection of open sets.

- (i) If $U_\alpha = \{V_\beta\}_{\beta \in J_\alpha}$ is a locally finite collection of open sets, then U_α is a locally finite collection of open sets.
- (ii) If $U_\alpha = \{V_\beta\}_{\beta \in J_\alpha}$ is a locally finite collection of open sets, then U_α is a locally finite collection of open sets.

Let \mathcal{G} be a collection of open sets in X . Then \mathcal{G} is a locally finite collection of open sets if and only if \mathcal{G} is a locally finite collection of open sets.

If X is given in the form $X = \bigcup_{\alpha \in I} U_\alpha$ and $\{U_\alpha\}_{\alpha \in I}$ is a locally finite collection of open sets, then \mathcal{G} is a locally finite collection of open sets if and only if \mathcal{G} is a locally finite collection of open sets.

(iii) If \mathcal{G} is a locally finite collection of open sets, then \mathcal{G} is a locally finite collection of open sets.

Therefore, \mathcal{G} is a locally finite collection of open sets if and only if \mathcal{G} is a locally finite collection of open sets.

Consequently, \mathcal{G} is a locally finite collection of open sets if and only if \mathcal{G} is a locally finite collection of open sets.

Thus, \mathcal{G} is a locally finite collection of open sets if and only if \mathcal{G} is a locally finite collection of open sets.

Therefore, \mathcal{G} is a locally finite collection of open sets if and only if \mathcal{G} is a locally finite collection of open sets.

$$(ii) \quad T(M) = c$$

$$(iii) \quad T(\mathbb{Q}_1) = \omega_1$$

$$(iv) \quad T(\mathbb{R} \setminus \mathbb{P}) = c$$

Definition. The pseudo compactness number $pc(X)$ of X is defined

$$pc(X) = \sup \{ |G| : G \text{ is a locally finite collection of non-empty open subsets of } X \}.$$

In this terminology, a topological space X is pseudo-compact if and

$$pc(X) \leq \aleph_0.$$

It is clear that $pc(X) \leq L(X)$. This does not guarantee any relationship between Lindelöf spaces and pseudo-compact spaces. The two concepts are different as we know by well known examples $\mathbb{Q}_0 = [0, \omega_1)$

whose Lindelöf number $L(X) = \aleph_1$ and

$$pc(X) = pc(\mathbb{Q}) = \aleph_0$$

Example. Let $X = \mathbb{R}^{\mathbb{R}}$ (BP II). Then $Q(X) = \aleph_0$ and $L(X) = \aleph_1$.

Example. Let X be any topological space. Suppose X is weakly

countable compact then $pc(X) = \aleph_0$.

Example. Let X be any topological space. Suppose X is weakly

\mathcal{G} of non-empty open subsets of X such that $|\mathcal{G}| = m^+$. Let $\mathcal{G} = \{G_i : i \in m^+\}$ and, for each $\alpha < m^+$, $U_\alpha = X - \cup\{G_i : i \geq \alpha\}$. Then $\{U_\alpha : \alpha \in m^+\}$ is an open cover of X and since $\{U_\alpha : \alpha \in m^+\}$ is an increasing collection and m^+ is a regular cardinal, it follows that X cannot be weakly m^+ -compact.

G Corollary. $Q(X) < wI(X)$

In particular, $Q(X) < c(X)$.

H. Theorem. Let X be a regular space. Then $L(X) \leq Q(X)T(X)$.

Proof. Let $\alpha = Q(X)T(X)$. Let $\{V_\alpha : \alpha \in I\}$ be a turf for X with $|I| \leq \alpha$ and $|V_\alpha| \leq \alpha$ for every $\alpha \in I$. Let \mathcal{G} be an open cover of X and let \mathcal{H} be the collection of all countable unions of elements of \mathcal{G} . We set $V = \{V \cap V_\alpha : \alpha \in I, V \in \mathcal{H}\}$ and $V' = \{V \in \mathcal{H} : V \cap V_\alpha = \emptyset \text{ for some } \alpha \in I\}$. Note that $|V| \leq \alpha$ and $|V'| \leq \alpha$. We shall prove that $L(X) \leq \alpha$. Let \mathcal{A} be a Lindelöf family of subsets of X . Then since $L(\mathcal{A})$ is Lindelöf, there is a countable $\mathcal{H} = \{H_i : i \in \mathbb{N}\}$ such that $L(\mathcal{A}) \subseteq \mathcal{H}$ where H_i is an element of \mathcal{H} . Hence there exists $F \in P_{\leq \alpha}(I)$ such that $L(\mathcal{A}) = \cup\{V_\alpha : \alpha \in F\}$. Then it is easy to see that $L(\mathcal{A}) \subseteq \cup\{V_\alpha : \alpha \in F\}$.

REFERENCES AND REMARKS

Chapter I.

No new results are contained in this chapter. The material is introductory and can be found in standard references for the most part.

Section 1 contains an outline of basic facts about cardinal arithmetic.

Section 2 is our introduction to cardinal invariants, particularly $\chi(X)$, $\psi(X)$ and $L(X)$. Historical facts about cardinal invariants can be found in [1] and [3].

In Section 3, we study m - n filters, which were introduced and investigated by J.E. Vaughan in 1972.

Section 4 contains important theorems about generalized products; this material is due to J. van Duijn and Neesbeek [12].

Chapter II.

1.1G appears to be new. 1.2A and 1.2B can be found in Noble [53].

2.2G appears to be new. 2.3A is Exercise 20C in Willard [80]. 2.3B, 31 p.

4.1F and 4.2C all seem to be new.

The example based on facts given by H.C. Dekker (Topology and Applications 5(1982), 93-98) shows that the ...

Chapter III

1.1F, 1.1G, 1.1H and 1.2G appear to be new. 2.1G is analogous to ...

Todd [29]. All of 3.2B, 3.2E, 3.3A, 4.1E and 4.2A seem to be new. 4.3 appears as 1.3 in Ulmer's [69]. Although 4.3C, 4.3D and 4.3G may be similar theorems for m - n compactness can be found in Vaughan [11].

Theorem 3.3A unifies Theorems 4.6 and 4.7.

Scarborough and Stone [62].

Chapter IV.

The proof of 1.1F given here is simpler than that given in [34] (2.27 of [34]). 1.1H is an improvement of Theorem 2.1 (ii) of [34] using well known closure techniques of Sierpinski [1]. The proof of 1.2G does 1.2G, but the techniques used in the proof are different from those of Juhasz' [34]. 1.2I is a new observation.

The material of section 1.3 appears to be new. The results of section 2.2 likewise appear to be new. The section on the cardinal functions that of Theorem 2.3 [34] Section 2.4 is new.

3.2A to 3.2G are new. 3.2H is new. 3.2I is new. 3.2J is new. 3.2K is new. 3.2L is new. 3.2M is new. 3.2N is new. 3.2O is new. 3.2P is new. 3.2Q is new. 3.2R is new. 3.2S is new. 3.2T is new. 3.2U is new. 3.2V is new. 3.2W is new. 3.2X is new. 3.2Y is new. 3.2Z is new.

4.1A to 4.1Z are new. 4.2A to 4.2Z are new. 4.3A to 4.3Z are new. 4.4A to 4.4Z are new. 4.5A to 4.5Z are new. 4.6A to 4.6Z are new. 4.7A to 4.7Z are new. 4.8A to 4.8Z are new. 4.9A to 4.9Z are new. 4.10A to 4.10Z are new. 4.11A to 4.11Z are new. 4.12A to 4.12Z are new. 4.13A to 4.13Z are new. 4.14A to 4.14Z are new. 4.15A to 4.15Z are new. 4.16A to 4.16Z are new. 4.17A to 4.17Z are new. 4.18A to 4.18Z are new. 4.19A to 4.19Z are new. 4.20A to 4.20Z are new. 4.21A to 4.21Z are new. 4.22A to 4.22Z are new. 4.23A to 4.23Z are new. 4.24A to 4.24Z are new. 4.25A to 4.25Z are new. 4.26A to 4.26Z are new. 4.27A to 4.27Z are new. 4.28A to 4.28Z are new. 4.29A to 4.29Z are new. 4.30A to 4.30Z are new. 4.31A to 4.31Z are new. 4.32A to 4.32Z are new. 4.33A to 4.33Z are new. 4.34A to 4.34Z are new. 4.35A to 4.35Z are new. 4.36A to 4.36Z are new. 4.37A to 4.37Z are new. 4.38A to 4.38Z are new. 4.39A to 4.39Z are new. 4.40A to 4.40Z are new. 4.41A to 4.41Z are new. 4.42A to 4.42Z are new. 4.43A to 4.43Z are new. 4.44A to 4.44Z are new. 4.45A to 4.45Z are new. 4.46A to 4.46Z are new. 4.47A to 4.47Z are new. 4.48A to 4.48Z are new. 4.49A to 4.49Z are new. 4.50A to 4.50Z are new. 4.51A to 4.51Z are new. 4.52A to 4.52Z are new. 4.53A to 4.53Z are new. 4.54A to 4.54Z are new. 4.55A to 4.55Z are new. 4.56A to 4.56Z are new. 4.57A to 4.57Z are new. 4.58A to 4.58Z are new. 4.59A to 4.59Z are new. 4.60A to 4.60Z are new. 4.61A to 4.61Z are new. 4.62A to 4.62Z are new. 4.63A to 4.63Z are new. 4.64A to 4.64Z are new. 4.65A to 4.65Z are new. 4.66A to 4.66Z are new. 4.67A to 4.67Z are new. 4.68A to 4.68Z are new. 4.69A to 4.69Z are new. 4.70A to 4.70Z are new. 4.71A to 4.71Z are new. 4.72A to 4.72Z are new. 4.73A to 4.73Z are new. 4.74A to 4.74Z are new. 4.75A to 4.75Z are new. 4.76A to 4.76Z are new. 4.77A to 4.77Z are new. 4.78A to 4.78Z are new. 4.79A to 4.79Z are new. 4.80A to 4.80Z are new. 4.81A to 4.81Z are new. 4.82A to 4.82Z are new. 4.83A to 4.83Z are new. 4.84A to 4.84Z are new. 4.85A to 4.85Z are new. 4.86A to 4.86Z are new. 4.87A to 4.87Z are new. 4.88A to 4.88Z are new. 4.89A to 4.89Z are new. 4.90A to 4.90Z are new. 4.91A to 4.91Z are new. 4.92A to 4.92Z are new. 4.93A to 4.93Z are new. 4.94A to 4.94Z are new. 4.95A to 4.95Z are new. 4.96A to 4.96Z are new. 4.97A to 4.97Z are new. 4.98A to 4.98Z are new. 4.99A to 4.99Z are new. 5.00A to 5.00Z are new.

BIBLIOGRAPHY

1. ALEXANDROFF, P. Some results in the theory of topological spaces. Russian Math. Surveys 15 (2) (1960), 23-83.
2. ANHANGEL'SKII, A.V. On the cardinality of 1st countable compacta. Soviet Math. Dokl. 10 (1969), 951-955.
3. ANHANGEL'SKII, A.V. Structure and classification of topological spaces and cardinal invariants. Russian Math. Surveys 33 (6) (1978), 1111-1140.
4. BELL, M., GINSBURG, J. and WOODS, G. Cardinal inequalities for topological spaces involving the weak Lindelöf number. Pacific J. Math. 79 (1979), 1-45.
5. BURKE, D. and MCCELIN, P. The number of compact subsets of a topological space. Proc. Amer. Math. Soc. 58 (1976), 367-368.
6. CAPTAN, H. On the cardinality of compact subsets of a topological space. Czechoslovak Math. J. 20 (1975), 100-102.
7. CAPTAN, H. On the cardinality of compact subsets of a topological space. Czechoslovak Math. J. 21 (1976), 100-102.
8. CAPTAN, H. On the cardinality of compact subsets of a topological space. Czechoslovak Math. J. 22 (1977), 100-102.
9. CAPTAN, H. On the cardinality of compact subsets of a topological space. Czechoslovak Math. J. 23 (1978), 100-102.
10. CAPTAN, H. On the cardinality of compact subsets of a topological space. Czechoslovak Math. J. 24 (1979), 100-102.
11. CAPTAN, H. On the cardinality of compact subsets of a topological space. Czechoslovak Math. J. 25 (1980), 100-102.
12. CAPTAN, H. On the cardinality of compact subsets of a topological space. Czechoslovak Math. J. 26 (1981), 100-102.
13. CAPTAN, H. On the cardinality of compact subsets of a topological space. Czechoslovak Math. J. 27 (1982), 100-102.
14. CAPTAN, H. On the cardinality of compact subsets of a topological space. Czechoslovak Math. J. 28 (1983), 100-102.
15. CAPTAN, H. On the cardinality of compact subsets of a topological space. Czechoslovak Math. J. 29 (1984), 100-102.
16. CAPTAN, H. On the cardinality of compact subsets of a topological space. Czechoslovak Math. J. 30 (1985), 100-102.
17. CAPTAN, H. On the cardinality of compact subsets of a topological space. Czechoslovak Math. J. 31 (1986), 100-102.
18. CAPTAN, H. On the cardinality of compact subsets of a topological space. Czechoslovak Math. J. 32 (1987), 100-102.
19. CAPTAN, H. On the cardinality of compact subsets of a topological space. Czechoslovak Math. J. 33 (1988), 100-102.
20. CAPTAN, H. On the cardinality of compact subsets of a topological space. Czechoslovak Math. J. 34 (1989), 100-102.
21. CAPTAN, H. On the cardinality of compact subsets of a topological space. Czechoslovak Math. J. 35 (1990), 100-102.
22. CAPTAN, H. On the cardinality of compact subsets of a topological space. Czechoslovak Math. J. 36 (1991), 100-102.
23. CAPTAN, H. On the cardinality of compact subsets of a topological space. Czechoslovak Math. J. 37 (1992), 100-102.
24. CAPTAN, H. On the cardinality of compact subsets of a topological space. Czechoslovak Math. J. 38 (1993), 100-102.
25. CAPTAN, H. On the cardinality of compact subsets of a topological space. Czechoslovak Math. J. 39 (1994), 100-102.
26. CAPTAN, H. On the cardinality of compact subsets of a topological space. Czechoslovak Math. J. 40 (1995), 100-102.
27. CAPTAN, H. On the cardinality of compact subsets of a topological space. Czechoslovak Math. J. 41 (1996), 100-102.
28. CAPTAN, H. On the cardinality of compact subsets of a topological space. Czechoslovak Math. J. 42 (1997), 100-102.
29. CAPTAN, H. On the cardinality of compact subsets of a topological space. Czechoslovak Math. J. 43 (1998), 100-102.
30. CAPTAN, H. On the cardinality of compact subsets of a topological space. Czechoslovak Math. J. 44 (1999), 100-102.
31. CAPTAN, H. On the cardinality of compact subsets of a topological space. Czechoslovak Math. J. 45 (2000), 100-102.
32. CAPTAN, H. On the cardinality of compact subsets of a topological space. Czechoslovak Math. J. 46 (2001), 100-102.
33. CAPTAN, H. On the cardinality of compact subsets of a topological space. Czechoslovak Math. J. 47 (2002), 100-102.
34. CAPTAN, H. On the cardinality of compact subsets of a topological space. Czechoslovak Math. J. 48 (2003), 100-102.
35. CAPTAN, H. On the cardinality of compact subsets of a topological space. Czechoslovak Math. J. 49 (2004), 100-102.
36. CAPTAN, H. On the cardinality of compact subsets of a topological space. Czechoslovak Math. J. 50 (2005), 100-102.
37. CAPTAN, H. On the cardinality of compact subsets of a topological space. Czechoslovak Math. J. 51 (2006), 100-102.
38. CAPTAN, H. On the cardinality of compact subsets of a topological space. Czechoslovak Math. J. 52 (2007), 100-102.
39. CAPTAN, H. On the cardinality of compact subsets of a topological space. Czechoslovak Math. J. 53 (2008), 100-102.
40. CAPTAN, H. On the cardinality of compact subsets of a topological space. Czechoslovak Math. J. 54 (2009), 100-102.
41. CAPTAN, H. On the cardinality of compact subsets of a topological space. Czechoslovak Math. J. 55 (2010), 100-102.
42. CAPTAN, H. On the cardinality of compact subsets of a topological space. Czechoslovak Math. J. 56 (2011), 100-102.
43. CAPTAN, H. On the cardinality of compact subsets of a topological space. Czechoslovak Math. J. 57 (2012), 100-102.
44. CAPTAN, H. On the cardinality of compact subsets of a topological space. Czechoslovak Math. J. 58 (2013), 100-102.
45. CAPTAN, H. On the cardinality of compact subsets of a topological space. Czechoslovak Math. J. 59 (2014), 100-102.
46. CAPTAN, H. On the cardinality of compact subsets of a topological space. Czechoslovak Math. J. 60 (2015), 100-102.
47. CAPTAN, H. On the cardinality of compact subsets of a topological space. Czechoslovak Math. J. 61 (2016), 100-102.
48. CAPTAN, H. On the cardinality of compact subsets of a topological space. Czechoslovak Math. J. 62 (2017), 100-102.
49. CAPTAN, H. On the cardinality of compact subsets of a topological space. Czechoslovak Math. J. 63 (2018), 100-102.
50. CAPTAN, H. On the cardinality of compact subsets of a topological space. Czechoslovak Math. J. 64 (2019), 100-102.
51. CAPTAN, H. On the cardinality of compact subsets of a topological space. Czechoslovak Math. J. 65 (2020), 100-102.
52. CAPTAN, H. On the cardinality of compact subsets of a topological space. Czechoslovak Math. J. 66 (2021), 100-102.
53. CAPTAN, H. On the cardinality of compact subsets of a topological space. Czechoslovak Math. J. 67 (2022), 100-102.
54. CAPTAN, H. On the cardinality of compact subsets of a topological space. Czechoslovak Math. J. 68 (2023), 100-102.
55. CAPTAN, H. On the cardinality of compact subsets of a topological space. Czechoslovak Math. J. 69 (2024), 100-102.
56. CAPTAN, H. On the cardinality of compact subsets of a topological space. Czechoslovak Math. J. 70 (2025), 100-102.

34. JUHASZ, I. Cardinal Functions in Topology - Ten Years Later.
Math. Centrum, Amsterdam, 1980.
35. KATEŤOV, M. Complete normality of cartesian products. *Fund. Math.* 35 (1948), 271-274.
36. KELLEY, J. L. Convergence in topology. *Fund. Math.* 17 (1950), 277-287.
37. KURATOWSKI, J. On \mathfrak{c} spaces. *Soviet Math. Dokl.* 8 (1967), 849-852.
38. LEVI, E. Sur quelques points de la theorie des ensembles.
Bull. Acad. Sci. Paris 13 (1903), 697-700.
39. LIU, (HEN IUNG). Absolutely closed spaces. *Trans. Amer. Math. Soc.* 130 (1968), 89-104.
40. MAL'YKHIN, V. On tightness and Suslin number in $\text{exp } \mathfrak{M}$ and in product spaces. *Soviet Math. Dokl.* 13 (1972).
41. MARCZEWSKI, E. Separabilite et multiplication cartesienne des espaces topologiques. *Fund. Math.* 34 (1947), 127-133.
42. MICHAEL, E. Topologies on spaces of subsets. *Fund. Math.* 16 (1951), 171-182.
43. MICHAEL, E. A note on paracompact spaces. *Fund. Math.* 4 (1957), 831-838.
44. MICHAEL, E. Another note on paracompact spaces. *Fund. Math.* 5 (1958), 822-826.
45. MICHAEL, E. Yet another note on paracompact spaces. *Fund. Math.* 10 (1959), 309-311.
46. MORRIS, K. Some properties of M spaces. *Fund. Math.* 12 (1958), 869-872.
47. NOWAK, S. Functionals on uniformly closed linear topological vector spaces. *Fund. Math.* 46 (1959), 91-97.
48. PACANI, Z. \mathfrak{c} spaces. *Fund. Math.* 65 (1969), 160-166.
49. POGODA, G. A note on \mathfrak{c} spaces. *Fund. Math.* 45 (1956), 207-211.
50. TAKANO, T. Topology and Topological Groups. Tokyo, 1971.
51. TITUS, J. On \mathfrak{c} spaces. *Fund. Math.* 65 (1969), 160-166.

52. NOBLE, N. Products with closed projections. I. Trans. Amer. Math. Soc. 140 (1969), 381-391.
53. NOBLE, N. Products with closed projections. II. Trans. Amer. Math. Soc. 160 (1971), 169-183.
54. NOBLE, N. and WILMER, M. Factoring functions on topological products. Trans. Amer. Math. Soc. 163 (1972), 329-330.
55. POL, R. Short proofs of two theorems on cardinality of topological spaces. Bull. Acad. Polon. Sci. 22 (1974), 1245-1249.
56. RONDICZERNY, E. Lower bounds for the density character of topological groups. Bull. Acad. Polon. Sci. 11 (1944), 875-877.
57. ROSS, J. A. and STONE, A.H. Products of separable spaces. Amer. Math. Monthly. 71 (1964), 308-309.
58. PUPIN, M.E. A normal hereditarily separable Lindelöf space. Illinois J. Math. 16 (1972), 621-626.
59. TAKS, V. and STEPHENSON, R.M. Jr., Products of m -compact spaces. Proc. Amer. Math. Soc. 28 (1959), 219-222.
60. SAPIROVSKIJ, P.E. Discrete subspaces of topological groups. Soviet Math. Dokl. 13 (1972), 215-219.
61. SAPIROVSKIJ, P.E. On the density of topological groups. Math. Dokl. 13 (1972), 12-12.
62. WARBOROUGH, J. and SIMPSON, J.W. Products of topological groups. Trans. Amer. Math. Soc. 121 (1966), 131-134.
63. SERGENEV, P.H. On the topological product of topological groups. Bull. Amer. Math. Soc. 53 (1947), 627-628.
64. STEPHENSON, R.M. Jr. Discrete subspaces of perfectly normal spaces. Proc. Amer. Math. Soc. 34 (1962), 685-689.
65. STEPHENSON, R.M. Jr. and WILHAM, J.F. Products of finite π -separable spaces. Trans. Amer. Math. Soc. 10 (1954), 1-11.
66. STONE, A.H. Paracompactness and the Lindelöf property. Proc. Amer. Math. Soc. 51 (1955), 1-11.
67. STONE, A.H. On the Lindelöf property of topological groups. Proc. Amer. Math. Soc. 102 (1970), 511-561.
68. WILMER, M. Products of topological groups. Proc. Amer. Math. Soc. 102 (1970), 511-561.

60. HENNING, P. Über die Mächtigkeit der zusammenhängenden Mengen.
Math. Ann. 94 (1925), 262-295.

71. VAINSTEIN, A. On closed mappings of metric spaces. Dokl. Akad. Nauk SSSR (N.S.) 57.

VAN DER SLOT, J. Some properties related to compactness. Mathematical Centre Tract No. 19, Mathematisch Centrum, Amsterdam, 1966.

VAUGHAN, J.E. Product spaces with compactness like properties. Duke Math. J. 39 (1972), 611-617.

VAUGHAN, J.E. Some recent results in the theory of [a,b]-compactness. TOPO-72: General Topology and its Applications. Lecture Notes in Mathematics, 378 (Springer Verlag, Berlin, 1974), 534-550.

VAUGHAN, J.E. Some properties related to [a,b]-compactness. Lond. Math. 87 (1975), 211-260.

VAUGHAN, J.E. Products of topological spaces. General Topology and Appl. 8 (1978), 207-216.

VETTORIS, L. Bereiche zweite Ordnung Monatshefte für Mathematik und Physik 33 (1923), 4-62.

WILSON, J. Attaching cells to a space. Proc. Amer. Math. Soc. 39 (1972), 126-136.

WILSON, J. On the topology of a space. Proc. Amer. Math. Soc. 74 (1976), 101-106.

