

Simulating how animals learn: a new modelling framework applied to the process of optimal foraging

Peter R. Thompson^{1,*}

Mérodie Kunegel-Lion²

Mark A. Lewis^{1,3,4,5}

1. Department of Biological Sciences, University of Alberta, Edmonton, AB, Canada;
2. Canadian Forest Service, Northern Forestry Centre, Edmonton, AB, Canada;
3. Department of Mathematical and Statistical Sciences, University of Alberta, Edmonton, AB, Canada.
4. Department of Biology, University of Victoria, Victoria, BC, Canada.
5. Department of Mathematics and Statistics, University of Victoria, Victoria, BC, Canada.

* Corresponding author; e-mail: pt1@ualberta.ca.

Keywords: Bayesian MCMC, statistical decision theory, animal learning, optimal foraging, individual-based modelling

Manuscript type: Article.

Prepared using the suggested L^AT_EX template for *Am. Nat.*

Abstract

Animal learning has interested ecologists and psychologists for over a century. Mathematical
3 models that explain how animals store and recall information have gained attention recently.
Central to this work is statistical decision theory (SDT), which relates information uptake in
animals to Bayesian inference. SDT effectively explains many learning tasks in animals, but
6 extending this theory to predict how animals will learn in changing environments still poses a
challenge for ecologists. We addressed this shortcoming with a novel implementation of Bayesian
Markov Chain Monte Carlo (MCMC) sampling to simulate how animals sample environmental
9 information and learn as a result. We applied our framework to an individual-based model
simulating complex foraging tasks encountered by wild animals. Simulated “animals” learned
behavioral strategies that optimized foraging returns simply by following the principles of an
12 MCMC sampler. In these simulations, behavioral plasticity was most conducive to efficient for-
aging in unpredictable and uncertain environments. Our model suggests that animals prioritize
highly concentrated resources even when these resources are less available overall, in line with
15 existing knowledge on optimal foraging and ideal free distribution theory. Our innovative com-
putational modelling framework can be applied more widely to simulate the learning of many
other tasks in animals and humans.

Introduction

18

Animals do not know everything about the environments they live in (Fagan et al., 2013), and even if they did, the world is becoming more unpredictable every day (Masson-Delmotte et al., 21 2021). While evolutionary adaptations are typically too slow to match these changes (Bell and Collins, 2008; Chevin et al., 2010; Merilä and Hendry, 2014), many animals can exhibit multiple behavioral responses to a changing environment without modifying their genetic code in a phenomenon known as behavioral plasticity (DeWitt et al., 1998; Schmidt et al., 2010; Snell-Rood, 24 2013; Wong and Candolin, 2015). Examples of behavioral plasticity range from temporal adjustments in the phenology of frogs in the temperate forests of the eastern United States (Gibbs and 27 Breisch, 2001) to the settlement of urban areas by birds in Europe (Møller, 2009). The ability to incorporate external information into a revised behavioral strategy may confer a fitness benefit to animals living in uncertain environments (Donaldson-Matasci et al., 2008; Parrish, 2000), but the 30 conditions in which behavioral plasticity is adaptive are not well-understood (Wong and Candolin, 2015). Most forms of behavioral plasticity involve learning (Snell-Rood, 2013), which has a rich theoretical background (Pearce, 2008) that could provide important context to the problem.

33 Our understanding of how animals learn is largely derived from laboratory studies of simple tasks (Pearce, 2008), which elucidate important cognitive mechanisms for learning but do not particularly resemble the natural world. This rich field of study can be traced back to Pavlov's 36 work on conditioning and associative learning (Harris and Bouton, 2020; Pavlov, 1927), which spawned theoretical and experimental work assessing the formation and extinction of these associative relationships, along with an animal's ability to categorize stimuli into different groups 39 (Katz and Wright, 2006; Pearce, 1987; Rescorla and Wagner, 1972; Spence, 1936). As food is often used as a positive reinforcer for animals (Pavlov, 1927), it follows logically that "foraging" tasks can effectively display how animals learn to prioritize different food resources based on their 42 relative reward (Krebs et al., 1978; Lea et al., 2012). Many of these conclusions draw heavily from optimal foraging theory (Charnov, 1976), generating a connection between cognitive and

spatial ecology. When the proper data are available, animal movement and foraging processes
45 are among the easiest methods for characterizing memory and learning in wild animals (Fagan
et al., 2013; Lewis et al., 2021). The mechanistic clarity of laboratory experiments and the realism
of animal movement models are difficult to combine into one analysis, but individual-based sim-
48 ulation modelling may be an effective tool for generating realistic patterns with clear mechanistic
origins (DeAngelis and Diaz, 2019; Murphy et al., 2020; Tang and Bennett, 2010).

Cognitive psychologists and ecologists have identified a striking resemblance between learn-
51 ing and Bayesian inference. This is most clear when couched in terms of statistical decision
theory (SDT; Berger, 1985). Broadly speaking, SDT is a mathematical framework describing the
optimal way animals or humans should make decisions according to learned information (Dall
54 et al., 2005; Dayan and Daw, 2008; Schmidt et al., 2010). A key component of SDT is the use
of Bayes's theorem to represent how prior knowledge is updated through learning to produce
a refined, posterior distribution of belief (Berger, 1985). Bayes's theorem and its key principles
57 have been used to explain many learning processes (Griffiths et al., 2001; McNamara et al., 2006),
including Pavlovian conditioning (Courville et al., 2006), mate choice (Castellano et al., 2012;
Luttbeg, 1996), and optimal foraging (Green, 1980, 2006; Valone, 2006). The application of SDT
60 to optimal foraging problems has inspired the term "Bayesian foraging", which describes how
animals update their foraging preferences in a decision-theoretic manner (Green, 1980; Valone,
2006). Most of this work has focused on small-scale foraging tasks, but in reality, foraging is a
63 very complex process influenced by many cognitive cues (Fagan et al., 2013). Extending SDT to
a model that wholly encompasses animal movement and foraging will produce results that are
more realistic and applicable to vulnerable wildlife populations.

66 Bayesian Markov Chain Monte Carlo (MCMC) sampling is a simple algorithm that we can
use to simulate how animals learn. MCMC sampling uses a stochastic approach to calculate
the posterior distribution of a set of parameters based on prior distributions and data supplied
69 by the user (Raftery and Lewis, 1992). When applied to learning, these parameters represent
biological qualities of an animal, and the data represent information collected by animals through

empirical experience. The structure of the prior and posterior distributions reflects the relative
72 “belief” an animal possesses in a certain behavioral strategy (i.e., combination of parameters)
before and after incorporating “data”, respectively. The data enforce revised posterior belief in
certain behavioral strategies through an objective function, which depends on the parameters and
75 may also be stochastic. While the objective function in a MCMC sampling procedure is typically
a probability distribution function (or likelihood function) of some sort, it does not need to be
continuous nor does it need to integrate to 1 over the sampled domain. Instead of using MCMC
78 to find the global optimum of a likelihood function, we can use it to identify behavioral strategies
that result in globally optimal fitness. In this example, the objective function would represent
the net energetic yield afforded by a specific strategy. Under this framework, MCMC simulates
81 how “animals” sample information by executing the task and evaluating the energy afforded
by different behavioral strategies (i.e., parameter values). Behavioral strategies that consistently
produce less favorable objective function values are less likely to accumulate probability mass in
84 the posteriors.

One complete run of the MCMC algorithm, which we henceforth refer to as a “chain”, con-
sists of many iterations. In each iteration the sampler draws random parameter values and calls
87 the objective function at those values, either accepting or rejecting the parameters based on the
function value. The number of iterations in a chain has important mathematical and biological
interpretations. Chains with more iterations allow for more extensive modification of the
90 priors, which biologically represent a simulated animal’s relative belief in different behavioral
strategies. With that in mind, we suggest that the number of iterations in a chain represents the
amount of information the animal gathers in its environment. We can more effectively ensure
93 that the animal consistently develops the same posterior belief in identically parameterized, but
randomly independent, chains when these chains have more iterations (this is mathematically
akin to ensuring the algorithm converges; Cowles and Carlin, 1996; Raftery and Lewis, 1992).
96 Some MCMC algorithms leave iterations at the beginning of the chain out of the posterior dis-
tribution, classifying them as “burn-in” iterations. The burn-in period was designed to enhance

chain convergence (Cowles and Carlin, 1996) but by omitting the behavioral strategies employed
99 at the beginning of the simulation process, the posterior distributions no longer include infor-
mation the animal gathered during the unrealistically “naive” (given the structure of the priors)
stages of learning.

102 During the sampling process, MCMC allows for the acceptance of suboptimal objective func-
tion values (i.e., lower than previous values) to search the parameter space more completely and
avoid local optima. The rate at which these suboptimal values are accepted can be likened to the
105 range of behavioral strategies an animal may try in a given environment. Animals that accept a
wide variety of strategies, even when they may not be optimal, could be thought of as displaying
behavioral plasticity. Consistently following the optimal behavioral strategy could be thought
108 of as displaying environmental canalization, a term used to characterize a lack of phenotypic
variation in reaction to environmental change (Gaillard and Yoccoz, 2003; Gibson and Wagner,
2000; Liefting et al., 2009). The simplest way to enforce this in the model is to introduce an
111 exponent $k > 0$ which is applied to the objective function during sampling. We can think of k
as an index of canalization, implying that lower values of k correspond to high behavioral plas-
ticity. Animals that possess high plasticity frequently sample many behavioral strategies amid
114 environmental uncertainty in what is commonly referred to as bet-hedging (Donaldson-Matasci
et al., 2008; Nevoux et al., 2010).

We expanded on existing implementations of SDT by coupling an individual-based simula-
117 tion model for animal movement with memory (Avgar et al., 2013) to a Bayesian model simu-
lating how animals learn to forage optimally. Our algorithm incorporates an objective function
measuring the net energetic intake of a foraging bout, given a set of parameters controlling ani-
120 mal behavior. To this end, the posterior distribution of these parameters obtained after sampling
reflects what simulated animals learned about the relative optimality of different foraging tech-
niques. We tested how effectively animals adjusted to unexpected and abrupt changes in the
123 distribution and abundance of resources on the landscape. We found that animals with higher
behavioral plasticity performed more efficient foraging returns after these abrupt changes, but

were less efficient when the environment did not change. Our framework displays how SDT can
126 be extended to the simulation of realistic ecological processes that, if formulated correctly, can
make effective predictions when data are lacking.

Methods

129 *The learning model*

We used Bayesian Markov Chain Monte Carlo (MCMC) sampling to simulate how animals learn
to adjust their behavior based on indicators of success. The effectiveness with which an animal
132 executes a certain task was quantified by an objective function f . Animals “sample” different
parameter values (i.e., behavioral strategies) and evaluate their optimality by calculating f ; de-
pending on the value of f , the animal may be more or less likely to attempt similar strategies as
135 represented by the posterior distribution of behavioral strategies.

We parameterized the MCMC sampler in a way that produced consistent and biologically
realistic results. We used uniform priors for each of the behavioral parameters under the as-
138 sumption that animals were not initially biased towards any strategy. Using relatively uninfor-
mative priors necessitated that we added a burn-in period to our chains, and we chose to omit
the first $N_{burn} = 500$ iterations of each chain from the posterior distribution to this end. Choos-
141 ing the number of iterations per chain (including burn-in), N_{iter} , was a careful optimization of
the trade-off between computational expense and consistency. Chains with more iterations take
longer to simulate but they also more accurately represent what simulated animals have learned.
144 We analyzed chains of different sizes to evaluate the fewest iterations necessary to produce con-
sistent posterior distributions, finding that $N_{iter} = 2000$ optimized the trade-off (see Appendix
B for more detail). This produced posterior distributions with $N_{iter} - N_{burn} = 1500$ parameter
147 values. We also tested many different values for k , the exponent applied to f during sampling:
 $k = 5, 10, 20, 50, 100, 200, 500, 1000$. Parameter values used in this study are summarized in Table
1. We ran our algorithm in Julia 1.6.2 using the Turing library, which offers a number of different

Par	Description	Value
MCMC algorithm parameters		
N_{iter}	Number of MCMC iterations per chain	2000
N_{burn}	Number of iterations in burn-in period	500
k	Exponent of objective function f	Many values
Behavioral parameters		
β	Degree of reliance on memory	Not fixed
γ	Likelihood to make long navigations	Not fixed
q	Default expectation of habitat quality	Not fixed
h	Relative preference for resource Q_1	Not fixed
Movement parameters (see Appendix A)		
N_r	Number of potential points of interest simulated	1000
λ	Exponent of C values when choosing point of interest	10
ρ	Average step length on navigations	2
κ	von Mises angular correlation parameter for navigations	10
Objective function parameters		
T_{train}	Length of training portion of each track	1000
T_{test}	Length of test portion of each track	1000
v	Energetic loss per 1 cell length of movement	0.05
N_{avg}	Number of tracks incorporated into one f call	5
Landscape parameters (see Appendix A)		
\underline{Q}	Threshold for landscape patches	0.6 or 0.9
d_L	Rate of resource depletion per time step	1.0
r_L	Recovery rate of depleted resources per time step	0.025

Table 1: Description of model parameters. The four parameters under the section “Behavioral parameters” are incorporated into the objective function f , and sampled in the Bayesian MCMC algorithm.

150 MCMC samplers. We used a static sampler that did not require tuning or calculating a proposal
function, which typically requires a function gradient (Sengupta et al., 2015). Parameters with
infinite support were log-transformed and bounded on finite intervals determined by assessing
153 their biological meaning.

Application of the model to foraging

We tested our modelling framework with an optimal foraging task involving the individual-
156 based simulation of animal movement across a continuous-space landscape. Our individual-
based model (IBM) for movement is heavily inspired by Avgar et al. (2013) and contains four
parameters mediating the behavioral strategy of simulated animals. We provide a summary of
159 the model and parameters below, but see Appendix A for a more detailed explanation of the
process using the ODD (Overview, Design Concepts, and Details) protocol (Grimm et al., 2006).

Simulated animals move on a landscape characterized as a a 100 x 100 arbitrary length unit
162 (lu) square in two-dimensional continuous space. The landscape has two independently dis-
tributed “resources” that provide an energetic benefit to the animal. In the interest of producing
movements similar to empirically observed location data, animals take discrete-time “steps” ev-
165 ery 1 arbitrary time unit (tu). Animals perceive, remember, and recall the quality of previously
visited foraging patches to make informed movement decisions. We make four key assumptions
about how animals do this, listed below:

168 (A1) Animals exhibit a preference for one of the two resources on the landscape and bias their
movements accordingly.

(A2) Animals remember the resource density of areas they have previously visited, but the ani-
171 mals’ reliance on memory decreases over time.

(A3) All points that the animal has not visited are perceived by the animal as having equal value,
regardless of their spatial or temporal position.

174 (A4) Animals are more likely to navigate to nearby points, all else held equal.

The foraging quality of any point \mathbf{x} at any time t , which we denote $Q(\mathbf{x}, t)$, ranges from 0 to 1 and is composed of two independent foraging resources, $Q_1(\mathbf{x}, t)$ and $Q_2(\mathbf{x}, t)$. While $Q(\mathbf{x}, t) =$
177 $(Q_1(\mathbf{x}, t) + Q_2(\mathbf{x}, t))/2$ across the landscape, we allow animals to exhibit “habitat selection” for the different resources on the landscape (Assumption A1). The behavioral parameter h ranges from 0 to 1 and mediates the animal’s relative preference for Q_1 and Q_2 . Simulated animals
180 perceive Q_1 and Q_2 as independent entities, and when computing the animal’s perceived foraging quality for any point \mathbf{x} and time t , we use $\tilde{Q}(\mathbf{x}, t) = hQ_1(\mathbf{x}, t) + (1 - h)Q_2(\mathbf{x}, t)$ as opposed to $Q(\mathbf{x}, t)$ (Figure 1).

183 Simulated animals perceive new information about resources on the landscape and encode this information into spatial memory. Many different animals use memory to guide their foraging movements (Bracis et al., 2018; Clayton and Dickinson, 1998; Panakhova et al., 1984; Potts and
186 Lewis, 2016; Ranc et al., 2021; Schlägel and Lewis, 2014), but heavy reliance on spatial memory is accompanied by numerous energetic costs (Fagan et al., 2013). The behavioral parameter $\beta \geq 0$ quantifies the extent to which simulated animals rely on their memory of previous foraging
189 experiences. As β increases the animal relies less on its memory, potentially a strategy to adapt to temporally variable environments (Fagan et al., 2013). We note that unlike memory decay, a neurological process (Thomas and Riccio, 1979), the mechanism displayed here represents the
192 animal’s conscious choice not to rely on the memory of previous experiences.

Animals make a naive, uninformed “guess” about the resource quality of locations they have not visited, and per Assumption A3, this guess is constant across space and time. Specifically,
195 any location will be assigned the value $q \in [0, 1]$ as long as that location remains unvisited by the animal. Larger values of q suggest that the animal is more “optimistic” about the quality of unexplored areas (Avgar et al., 2013; Berger-Tal and Avgar, 2012), and will more frequently visit
198 these areas as a result.

Once the animal generates an expectation of resource quality across the landscape, it must choose a location to navigate to. Assumption A4 states that animals are more likely to navigate




Figure 1: Schematic describing the generation of $C(x, t)$, the animal's estimation of resource quality across the environment. The animal weights two independently distributed resources and incorporates newly perceived information into C based on the perception function $p(x, x_t)$. Note the incremental updating of C as the animal moves to a new location (x_t , pictured in blue on the bottom right).

201 towards nearby points than faraway points. This idea follows logically from the marginal value
theorem (Charnov, 1976), which considers the energetic cost of travel to other patches. We in-
cluded behavioral parameter $\gamma \geq 0$ to quantify this relationship. As γ increases, the probability
204 that the animal will navigate to a faraway point decreases; even if the animal believes there are
resources far away, it may opt for nearby resource patches instead, a tactic many animals adopt
as a risk avoidance mechanism (Gehr et al., 2020).

207 The animal’s perceived resource quality for any point \mathbf{x} and time t , denoted $C(\mathbf{x}, t)$, depends
on these four assumptions. This function consists of a weighted average of three quantities:
newly perceived information (weighted by perception function $p(\mathbf{x}, \mathbf{y})$), memorized information
210 (weighted by memory function $m(t)$), and the naive expectation q .

$$p(\mathbf{x}, \mathbf{y}) = \exp\left(-\frac{d(\mathbf{x}, \mathbf{y})}{\rho}\right), \quad (1)$$

$$m(t) = \exp(-\beta t), \quad (2)$$

$$C(\mathbf{x}, t) = \underbrace{p(\mathbf{x}, \mathbf{x}_t)\tilde{Q}(\mathbf{x}, t)}_{\text{perception}} + \underbrace{(1 - p(\mathbf{x}, \mathbf{x}_t))\left(m(1)C(\mathbf{x}, t - 1) + (1 - m(1))q\right)}_{\substack{\text{memory} \\ \text{expectation}}}. \quad (3)$$

The perception function relies on the assumption that animals perceive nearby information
more accurately than faraway information (Avagar et al., 2015; Fagan et al., 2017; Fletcher et al.,
213 2013), where $d(\mathbf{x}, \mathbf{y})$ is the distance between \mathbf{x} and \mathbf{y} and ρ is the animal’s average movement
speed in lu/tu. A positive association between movement capability and perceptual range has
been documented across many animal taxa (Kiltie, 2000; Møller and Erritzøe, 2010).

216 *Calculating the objective function*

We designed an objective function f measuring the energetic benefit afforded by a certain be-
havioral strategy. We divided these simulated foraging bouts into “training” and “test” sections

219 of durations T_{train} and T_{test} , respectively, and only measured f over the test section. Avgar et al. (2013) made a similar correction to allow animals to develop an initial memory of their simulated landscape, producing movement paths that resemble empirically collected animal location data. 222 We subtracted the animal’s total resource intake across the simulation by the energetic loss as a result of movement, calculated as the animal’s total distance traveled multiplied by a proportionality constant $v \geq 0$ (Table 1). Our function f consists of an average of N_{avg} independent 225 movement tracks so it effectively characterizes the expected value of any parameter combination. We define f_i , the net energetic intake from the i^{th} of these tracks, by summing the energetic gains collected at each location \mathbf{x}_t along the animal’s path:

$$f_i(\beta, \gamma, q, h|\mathbf{Q}) = \frac{\sum_{t=T_{train}+1}^{T_{train}+T_{test}} Q(\mathbf{x}_t, t) - v \sum_{t=T_{train}+1}^{T_{train}+T_{test}} d(\mathbf{x}_t, \mathbf{x}_{t-1})}{T_{test}}, \quad (4)$$

$$f(\beta, \gamma, q, h|\mathbf{Q}) = \frac{1}{N_{avg}} \sum_{i=1}^{N_{avg}} f_i(\beta, \gamma, q, h|\mathbf{Q}). \quad (5)$$

228 *Scenarios of environmental change*

We randomly generated spatially autocorrelated resource landscapes (see Appendix A for further detail) and used them to simulate abrupt landscape-level changes in the environment. Bayesian 231 inference allows for the iterative updating of prior expectations based on previous analyses (El-lison, 2004). The posterior distributions of our behavioral parameters represent knowledge accumulated by a simulated animal, which we can use as more “informative” priors for a second 234 MCMC chain. Each of our “scenarios” of environmental change contains two stages, where each stage has a unique Q_1 and Q_2 (Figure 2). The scenarios we generated incorporate two “types” of landscape, which can be visually compared in the first chain of Scenario A (Figure 2). Here, Q_1 is 237 much more abundant and widely distributed than Q_2 , but Q_2 is richer than Q_1 in the small area where it can be found. Scenario A serves as a “control” where the environment does not change; we would expect the animal to identify an optimal strategy and retain this strategy for both

240 chains. In Scenarios B and C, only one of the resources switches between being widely abundant
and locally available (the difference being the directionality of this change), and in Scenario D,
both resources swap.

243 We ran the MCMC algorithm with each of the four scenarios and a suite of k values (5, 10, 20,
50, 100, 200, 500, 1000) to evaluate how these quantities affected optimal foraging behavior. For
each value of k and scenario, we ran algorithm 12 independent times. We obtained posteriors
246 for the first and second chains of each run for the four parameters $\beta, \gamma, h,$ and q , along with a
posterior distribution of f_i values (1500 iterations after burn-in \times 5 f_i per f call \times 12 chains =
90000 total f_i calls) for each k and scenario.

249

Results

Posterior distribution of parameters

Under the same environmental circumstances, 12 independently simulated MCMC runs pro-
252 duced similar posterior distributions, suggesting that $N_{iter} = 2000$ and $N_{burn} = 500$ is sufficient
for convergence (a subset of these are displayed in Figure 3). In most circumstances, simulated
animals displayed a relatively “pessimistic” expectation of unvisited food patches, as suggested
255 by posterior distributions concentrated around low values of q . Posterior distributions of β were
relatively spread out across all values, suggesting that long-term reliance on memory only has
a minimal advantage over short-term reliance in these simulations. Simulated animals avoided
258 long-distance navigations, opting instead for values of γ close to 1 frequently (Figure 3). Most
notably, though, animals simulated in Scenario A (Figure 2) exhibited a strong preference for
resource Q_2 , which was much less abundant across the landscape than Q_1 . This is indicated by
261 posterior distributions for h concentrated around lower values.




Figure 2: Different scenarios of environmental change used in our simulations. Scenario A is a “control” where the environment, composed of two resources Q_1 and Q_2 , does not change at all. In Scenarios B and C, Q_1 stays the same, but Q_2 becomes more or less abundant than Q_1 , respectively. In Scenario D, the distributions of Q_1 and Q_2 “swap”.




Figure 3: Posterior density plots for one independent runs of the MCMC algorithm, taken from the first chain of Scenario A (see Figure 2) with $k = 10$. Greater probability mass at certain parameter values indicates higher belief in that value optimizing the net energetic gain function f . Note, in particular, the animal's preference for resource Q_2 , which in this case is much less widely available but provides a larger energetic benefit than Q_1 where it can be found (Figure 2).




Figure 4: Example violinplots detailing the distribution of objective function f_i , which represents the net energetic gain from a simulated animal foraging bout. These two violinplots are taken from the second chain of Scenario A (see Figure 2), with k taking on two different values.

Posterior distribution of objective function values

Both the scenario of environmental change and the MCMC parameter k affected the second chain's posterior distribution of f_i values. Typically, the spread of these distributions increased as k decreased, especially in Scenario A, where they appear similar to delta functions at $k = 500$ and $k = 1000$ (Figure 4). In scenarios where the environment changed dramatically (e.g., Scenario D; Figure 2), these distributions took on different shapes, sometimes becoming bimodal (Figure 5).

More specifically, the effect of MCMC parameter k on the distribution of objective function f_i values depended on the scenario of environmental change (Figure 6). In Scenario A (Figure 2), simulated animals performed more consistently and efficiently with large values of k than with small k . In Scenario B, k had a much smaller effect on foraging success than Scenario A, although the spread of f_i values was larger with smaller k (Figure 6). The posterior distributions of f_i from Scenario C resemble those from Scenario A at low k , but appear to take on a skewed, slightly bimodal shape at higher k . In Scenario D, intermediate values of k ($k = 100$ and $k = 200$)




Figure 5: Example violinplots detailing the distribution of objective function f_i , which represents the net energetic gain from a simulated animal foraging bout. These two violinplots are drawn from the second chain of Scenarios A and D (Figure 2), respectively, with $k = 200$ for each.

276 produced foraging bouts that were, on average, more efficient than at large values of k (Figure
 6). The distribution of f_i values was distinctly bimodal with large k , and as k increased, more
 probability mass was concentrated in the second, lower mode.

279 Discussion

Predicting how animals will adjust to environmental change is an important but complex ecolog-
 ical problem. We developed a Bayesian model that simulates how animals sample information
 282 about their environments to develop a posterior distribution of optimal foraging behavior. Our
 model builds on statistical decision theory, which has long been used to explain how animals
 learn from a Bayesian perspective (Berger, 1985; Dall et al., 2005; McNamara and Houston, 1980).
 285 We applied our learning model to a complex, continuous-space foraging task to be completed by
 simulated spatially informed foragers (Avgar et al., 2013). In the presence of two independently
 distributed resources with equal energetic return, animals simulated in our model prioritized
 288 resources that were concentrated within small, sparsely distributed patches. Animals that ex-




Figure 6: Effect of MCMC parameter k on foraging efficiency in simulated animals under four different scenarios of temporal environmental change (see Figure 2 for detail on each scenario). Each individual violinplot represents a sample of 90000 f_i values (12 independent runs of MCMC \times 1500 f values per run \times 5 f_i values per f call) representing the net energetic gain from a single simulated movement track. The red line represents the mean of all f_i values from each k value.

hibited canalized behaviour displayed consistently efficient foraging returns in temporally predictable environments, but environmental canalization became maladaptive when we introduced sudden, unpredictable changes to the landscape. Our results suggest that Bayesian MCMC can be used to simulate how animals, and potentially even humans, learn a wide variety of tasks in an ever-changing world.

When faced with the choice of two resources, simulated animals chose the resource that was available in smaller, but more heavily concentrated patches (Figure 3). We expected to observe a preference for the more abundant Q_1 , but our model showed that they instead opted for the less abundant but richer Q_2 . This finding suggests that simulated animals occupy areas with the highest possible habitat suitability, a key principle of ideal free distribution (IFD) theory (Cantrell et al., 2007; Fretwell and Lucas, 1969). Many patterns predicted by IFD theory can be seen in our results even though our IBM did not incorporate competition between individuals (this could be an interesting topic for future work). Specifically, IFD theory predicts that individuals residing in poor habitat will adjust for the lack of resource abundance by adopting larger home ranges (Haché et al., 2013). Simulated animals in our model centralized their movements around small plentiful resource patches, producing smaller home range sizes than individuals that foraged on less concentrated resources. Similarly, the resource dispersion hypothesis predicts that animals will occupy larger home ranges when resources are less spatially concentrated (Macdonald, 1983; Macdonald and Johnson, 2015). Increasing the speed or breadth of resource depletion or further decreasing the spatial availability of these concentrated resources could modify this relationship.

The wide variety of behavioral strategies adopted by simulated animals with high behavioral plasticity during sampling produced variable energetic outcomes. Behavioral plasticity allows animals to exhibit a variety of foraging strategies simply as a result of learning and adjusting to new environmental drivers (Parrish, 2000). Animals with highly canalized behavior (i.e., low plasticity) would be expected to perform one foraging strategy consistently (Gaillard and Yoccoz, 2003; Snell-Rood, 2013; Wong and Candolin, 2015), and we frequently saw that in our simulations. This is also unsurprising from an analytical perspective, since k (specificity) also resembles the

number of “clones” used in data cloning algorithms (Lele et al., 2007). This consistency also suggests that there is minimal stochastic variation in the value returned by our objective function f_i when our behavioral parameters were held constant.

Our simulations strongly suggest that behavioral plasticity is adaptive when the environment changes dramatically and unexpectedly. Animals simulated in temporally constant environments had unimodal distributions of energetic return, but those simulated in temporally unpredictable environments had a second mode centered around a lower energetic intake (Figure 5). The latter group of simulated animals foraged efficiently until the distribution of resources suddenly changed, rendering the original strategy suboptimal. Animals with high behavioral plasticity shifted their resource preferences depending on the environment, for better or for worse (Dunn et al., 2020; Parrish, 2000; van Baaren and Candolin, 2018). Animals with very low behavioral plasticity continued to forage according to their initial, now suboptimal, strategy, while animals with intermediate levels of behavioral plasticity adjusted their foraging strategies more effectively (Figure 6). Animals with very high behavioral plasticity performed a diverse array of foraging strategies, many of which were too inefficient to produce optimal foraging returns.

While behavioral plasticity is typically considered an adaptive trait, some animals suffer from it. Ecological traps are resources that appear beneficial to animals but, in reality, do not confer a fitness benefit (e.g., mayflies lay their eggs on asphalt because it reflects light similarly to water; Kriska et al., 1998). Ecological traps have become more frequent in the Anthropocene due to the proliferation of man-made novel objects in natural environments (Robertson and Chalfoun, 2016). A typical consequence of behavioral plasticity is an increased likelihood to explore unfamiliar stimuli (Mettke-Hofmann et al., 2009; Snell-Rood, 2013), which is believed to associate behavioral plasticity and vulnerability to ecological traps (Robertson and Chalfoun, 2016). The results from our simulation study corroborate empirical evidence that environmental canalization can be more effective than behavioral plasticity in some environments.

Translocated animals represent an effective way to test our model, displaying behavior similar to our simulations. Animal translocation and reintroduction protocols have many purposes,

ranging from the displacement of potentially dangerous animals (Milligan et al., 2018) to the restoration of populations and ecosystems (Polak and Saltz, 2011; Seddon et al., 2007). Translocated animals are abruptly brought to entirely new environments where they must learn to forage optimally or face heightened mortality risk. The nature of these protocols makes them an effective real-life test for our model, and many of the predictions offered by our model are verified from translocation studies. Translocated elk (*Cervus canadensis*) displayed different foraging behavior depending on the environmental conditions in their original home range and the environmental change they underwent (Falcón-Cortés et al., 2021). Specifically, elk translocated between two very different environments (resembling our Scenario D) were more exploratory and less reliant on memory than those translocated between similar environments, suggesting a shift in behavior from their original home ranges (Falcón-Cortés et al., 2021). As another case study, greater prairie-chickens (*Tympanuchus cupido*) typically sought out habitat similar to that of their natal ranges, suggesting a strong prior preference for resources found in their old environments (Kemink and Kesler, 2013). Here, canalization was detrimental to the birds' survival, adding support to the pattern observed in panel D of Figure 3. Translocations and reintroductions are frequently practiced across a wide array of animal taxa, but they are still risky and unpredictable (Berger-Tal and Saltz, 2014). The principles drawn from our analysis provide an improved forecast for the efficacy of these protocols in different ecological systems.

Conclusion

We developed a modelling framework that innovatively applies the principles of Bayesian statistics to animal foraging and learning. Much of what we currently know about animal learning comes from manipulative experiments conducted with captive animals (Pearce, 2008). Many of these studies have been critical for unearthing the mechanisms behind animal cognition, memory, and learning (Pavlov, 1927; Rescorla and Wagner, 1972), but they do not replicate the conditions wild animals experience. By incorporating the prevailing mathematical theory behind

animal learning, our modelling framework fills this gap. Our results with respect to continuous-
369 space foraging align with optimal foraging theory (Charnov, 1976), ideal free distribution theory
(Fretwell and Lucas, 1969), and prevailing knowledge on behavioral plasticity (Robertson et al.,
2013; Wong and Candolin, 2015). With that being said, our model for learning is general enough
372 that it need not be confined to optimal foraging. Specifically, any problem that can be char-
acterized in the form of an objective function and a set of parameters representing behavior is
tractable for our framework. This could include movement on different spatial or temporal scales,
375 social learning, or communication. Even more thought-provoking is the potential for our mod-
elling framework to predict how humans learn and make decisions. While the decisions made
by animals can affect the individual's fitness and survival, human decisions have the potential
378 to reverberate much more widely, which has become even more apparent during the COVID-19
pandemic (Bavel et al., 2020). Through these potential applications and more, our computational
modelling framework has the capacity to address challenging problems in cognitive science.

381

Author contributions

PRT, MKL, and MAL conceived the idea for the model. PRT wrote all computer programs and ran
all analyses. PRT wrote the first draft of the manuscript, upon which MKL and MAL provided
384 extensive feedback.

Appendix A: ODD Protocol

Purpose

387 We developed a model heavily influenced by Avgar et al. (2013) to simulate the movement of spatially informed foragers. The model includes four parameters that, when combined, quantify an animal's foraging strategy. These parameters are intended to measure behaviorally plastic 390 qualities of an animal as opposed to genetic or morphological traits. We assessed the adaptive value of different foraging strategies using a net energetic gain metric, which weighs the animal's resource intake against the energetic cost of movement. We do not specifically liken the model 393 to any animal taxon, but we note that many common behavioral processes (e.g., migration and sociality) are not included in the model.

State variables and scales

396 The model consists of one individual (henceforth referred to as an "animal") that moves throughout a bounded spatial landscape. The animal performs discrete-time, continuous-space movements at constant temporal intervals of 1 arbitrary time unit (tu). The landscape is a 100 x 100 399 arbitrary length unit (lu) square in two-dimensional space. Each spatial point on the landscape \mathbf{x} and time index t has a resource quality $Q(\mathbf{x}, t) \in [0, 1]$ representing the energetic value of resources at that point. For mathematical convenience, we formulated $Q(\mathbf{x}, t)$ as a piecewise 402 constant function; all \mathbf{x} in any 1x1 lu "grid cell" have the same value of $Q(\mathbf{x}, t)$ at any time t . To prevent animals from getting "trapped" in corners or boundaries of the landscape, we assume that landscapes have wrap-around boundaries (e.g., if the animal moves far enough to the left, it 405 will eventually end up on the right side of the grid).

The landscape has two unique resources that are added together to produce the total resource quality $Q(\mathbf{x}, t)$ for each point and time. We define $Q_1(\mathbf{x}, t)$ and $Q_2(\mathbf{x}, t)$ to be the quality values 408 for the first and second resources at point \mathbf{x} and time t , respectively. Both of these resource

functions can take on values between 0 and 1, so to ensure that $Q(\mathbf{x}, t)$ is defined properly, we set $Q(\mathbf{x}, t) = (Q_1(\mathbf{x}, t) + Q_2(\mathbf{x}, t)) * 0.5$ for every point \mathbf{x} and time t .

411 We incorporated depletion-recovery dynamics to the landscape to ensure animals would be incentivized to move. When the animal visits any point in a grid cell, it consumes and depletes that cell's resources. Specifically, we decrement $Q_1(\mathbf{x}, t)$ and $Q_2(\mathbf{x}, t)$ by resource depletion parameter d_L for every point \mathbf{x} in the cell the animal visits at time t . If d_L is greater than the resource value at that time, the cell is depleted entirely and is assigned a resource value of 0. Each depleted resource recovers by r_L units each time step until reaches its original, pre-depletion value.
414
417 We fixed d_L and r_L for all simulations (Table 1).

Process overview and scheduling

We tracked information storage in simulated animals using $C(\mathbf{x}, t)$, which represents the animal's estimation of resource quality for each point and time. As the animal perceives and remembers new information through movement, C is updated. The animal moves by choosing a "point of interest" to navigate to based on C . Points of interest may take more than 1 tu to reach, reflecting the numerous timescales at which animals make movement decisions (Blackwell et al., 2016; McClintock et al., 2014).
420
423

Design concepts

426 *Fitness:* Simulated animals perform the most basic version of "fitness-seeking" in that they search for points with a higher concentration of resources. Following Assumption A1, animals exhibit "habitat selection" for the different resources on the landscape. We introduce the parameter $h \in [0, 1]$ to quantify this relationship. When the animal visits a new location, it stores the value of that location as $\tilde{Q}(\mathbf{x}, t) = hQ_1(\mathbf{x}, t) + (1 - h)Q_2(\mathbf{x}, t)$ rather than $Q(\mathbf{x}, t)$ (Figure 1). Per Assumption A4, animals will be more likely to navigate to nearby points, as this minimizes locomotive cost as well as the opportunity cost of navigating through potentially resource-poor
429
432

habitat on the way to a faraway point of interest.

435 *Sensing:* Animals are not omniscient and must obtain information via perception. Typically, animals perceive nearby information more accurately (Avgar et al., 2015; Fagan et al., 2017). Mathematically, we formalize this using a perception function $p(\mathbf{x}, \mathbf{y})$. This function measures
438 how accurately (ranging from 0 to 1) an animal located at \mathbf{x} perceives information about \mathbf{y} . We chose an exponential decay function (similar to Avgar et al., 2013) to represent this relationship:

$$p(\mathbf{x}, \mathbf{y}) = \exp\left(-\frac{d(\mathbf{x}, \mathbf{y})}{\rho}\right), \quad (\text{A1})$$

where $d(\mathbf{x}, \mathbf{y})$ is the distance between \mathbf{x} and \mathbf{y} , accounting for wrap-around boundaries. We as-
441 sume that the animal's perceptual ability increases with ρ , the parameter governing the animal's locomotive capability.

Memory: Assumption A2 states that the animal's reliance on memory decreases as the time since
444 the formation of that memory increases. Mathematically, we used an exponential decay function to represent this (similarly to Avgar et al., 2013). The function $m(t)$ ranges between 0 and 1 and quantifies the animal's reliance on memory as a function of how long ago the memory was
447 formed. Simply put, $m(t) = \exp(-\beta t)$. If $\beta = 0$, the animal effectively has an infinite memory, and as β becomes infinitely large, the animal begins to neglect its memory entirely.

450 *Prediction:* The animal estimates the resource quality at any point on the landscape using perception and memory, but if it has never visited a location on the grid, it must still make a naive "guess" about the resource quality there (Avgar et al., 2013; Berger-Tal and Avgar, 2012). As-
453 sumption A3 states that this guess is constant across space and time; in other words, the animal will treat all unvisited points equally throughout the simulation. We can represent this guess with $q \in [0, 1]$. Larger values of q will result in more exploratory movement as animals assign
456 higher value to unvisited areas.




Figure A0: Flowchart describing the individual-based simulation model for animal movement. At each time step, animals update their perception of the environment C , occasionally using it to choose a point of interest (POI) to navigate to. This navigation can take any number of time steps, as the animal does not typically stop navigating until it reaches the point.

Stochasticity: Animal movement paths are stochastic, and as a result animals will not always visit
 459 the patch that confers the highest expected benefit (i.e., the highest value of C). That being said, points with higher values of C are still more likely to be chosen as points of interest. When the animal is not currently en route to a point of interest, a new point of interest is picked
 462 using a Monte Carlo sampling technique. This involves simulating $N_r = 1000$ possible points of interest $\mathbf{x}_{t,1}, \mathbf{x}_{t,2}, \dots, \mathbf{x}_{t,N_r}$ and randomly picking one (denoted \mathbf{x}_t^P) based on the value of C . More specifically,

$$P(\mathbf{x}_{t,i} = \mathbf{x}_t^P) = \frac{C(\mathbf{x}_{t,i}, t)^\lambda}{\sum_{j=1}^{N_r} C(\mathbf{x}_{t,j}, t)^\lambda}, \quad (\text{A2})$$

465 for any positive integer $i \leq N_r$. We include a fixed constant $\lambda \geq 0$ that controls the “determinism” of the animal’s movements: as λ increases, it is more likely to choose the point with the highest value of C .

468 We simulate the $\mathbf{x}_{t,i}$ as end points of a movement “step” beginning at \mathbf{x}_{t-1} , where the lengths of each step follow an exponential distribution. The shape of this distribution results in smaller

step lengths being more frequently sampled, following Assumption A4. We define $\gamma \geq 0$ as
471 the “rate” parameter of the exponential distribution, quantifying the strength of the relationship
between distance and point-of-interest selection. As γ approaches 0, every point on the grid has
an equal chance of being selected (assuming equal values of C). If γ is large, all $\mathbf{x}_{t,i}$ will be close
474 to the animal and it will not undertake long-distance navigations very often.

The animal navigates to points of interest by performing a biased random walk (Figure A0).
The lengths of each step along the navigation are simulated from a gamma distribution with
477 mean and variance ρ . This distribution has an entirely positive support and is roughly bell-
shaped for most values of ρ , including the value we used (Table 1). If simulated step lengths are
longer than the distance to the point of interest (i.e., the animal would “overshoot” its destina-
480 tion), the animal goes directly to the point of interest instead. Otherwise, it takes a step of the
simulated length towards the point of interest. The heading of this step is simulated from a von
Mises distribution where the mean heading is the heading required to reach the point of interest.
483 The concentration parameter for this distribution, $\kappa \geq 0$, is a fixed quantity in this model (Table
1). It is recommended that large values of κ , which cause more directed movement to the point
of interest, are used here. If one of the steps on the animal’s navigation ends on a point that has
486 better resources than the point of interest (i.e., $\tilde{Q}(\mathbf{x}_{t+1}, t+1) > C(\mathbf{x}_t^P, t+1)$), the animal “forgets”
about the point of interest and prioritizes foraging at the newfound location. The algorithm
restarts whenever the animal arrives at its point of interest.

489
Observation: We collected information about the animal’s movement as well as its cumulative
resource intake. We keep track of the animal’s location \mathbf{x}_t , as well as the value of $Q(\mathbf{x}_t, t)$, for
492 each time step t in the track. Note that while the animal exhibits relative preference for resources
using \tilde{Q} , it still takes in equal amounts of both resources when it visits a patch.

495 Our model does not implement interaction or collectives since animals are solitary on the land-
scape. While we assume that animals can “adapt” to environmental conditions by modifying

$\beta, \gamma, q,$ and h between simulations, we do not allow for adaptation within a single simulation. We
498 are not particularly focused on emergent properties such as home range formation.

Initialization

At the beginning of each simulation, we randomly generate a landscape and initialize the animal
501 at a random point on that landscape. Initially, $C(\mathbf{x}, 0) = q$ for every point \mathbf{x} , as the animal has no
prior experience on the grid.

Input

504 For each simulated animal movement path, we supplied two randomly generated landscapes
(for Q_1 and Q_2 respectively) as inputs. We simulated our landscapes as Gaussian random fields,
implying that each cell on the grid is a component of a multivariate Gaussian random variable
507 (Schlather, 2012). In this case, the covariance between any two cells depends on the wrap-around
distance between the two cells (closer cells have higher covariance). We then scaled the values
such they all fell between 0 and 1.

510 To more accurately capture the patchiness of many real-world habitats, we defined a cut-off
value \underline{Q} that could be used to make these landscapes more patchy. Under this rule, any grid cell
with a value of Q below \underline{Q} would be set to 0. Increasing \underline{Q} decreases the overall resource quality
513 of the landscape and is more likely to confine the animal to specific high-quality patches. Here,
we used landscapes with $\underline{Q} = 0.6$ and $\underline{Q} = 0.9$ (Figure 2).

Submodels

516 Our main submodel is the calculation of C , the animal's spatial map of perceived resource quality.
This calculation is composed of three mechanisms: perception ($p(\mathbf{x}, \mathbf{y})$), memory ($m(t)$), and
default expectation (q). Figure 1 displays how these quantities are combined and weighted to
519 produce C . This is mathematically formalized below:

$$\begin{aligned}
C(\mathbf{x}, t) = & \underbrace{p(\mathbf{x}, \mathbf{x}_t)\tilde{Q}(\mathbf{x}, t)}_{\text{perception}} + \\
& (1 - p(\mathbf{x}, \mathbf{x}_t)) \left(\underbrace{m(1)C(\mathbf{x}, t - 1)}_{\text{memory}} + \underbrace{(1 - m(1))q}_{\text{expectation}} \right).
\end{aligned} \tag{A3}$$

Note that $m(1) = \exp(-\beta)$, which resembles the model from Avgar et al. (2013).

Appendix B: Determining the appropriate number of MCMC

iterations

522

We determined an optimal number of iterations per MCMC chain by identifying when additional iterations did not substantially affect the posterior distribution of the four behavioral parameters.

525

If some value N were to be sufficient as the number of iterations per chain, we would expect that a chain simulated for N iterations would produce similar posteriors when we added additional iterations to the chain. If simulating more iterations produced negligibly different posteriors, it

528

is not computationally worthwhile to perform those iterations. To that end, we ran a chain of the MCMC sampling algorithm for our foraging task with 5000 iterations (what we deemed to be the largest computationally reasonable value). We then took the first N iterations of that chain

531

and compared the posterior distribution from that subset with a slightly larger subset, the first $N + 500$ iterations. We used a static MCMC sampler in our analysis so the individual iterations were independent of each other, rendering this process similar to comparing two separate chains.

534

We compared posterior distributions using the earth mover's distance, also known as the Wasserstein distance, a common tool for comparing multivariate distributions across many fields (Potts et al., 2014; Rubner et al., 2000; Vaserstein, 1969). The earth mover's distance approximates

537

the energy required to spatially transform one probability distribution such that it resembles another. As a result, lower values of this metric suggest higher distributional similarity, and an earth mover's distance of 0 is only achieved between two perfectly identical probability distri-

540

butions. Plotting the earth mover's distance against N , the proposed number of iterations, for many different values of N (ranging from 500 to 4500 by 100) led us to identify $N_{iter} = 2000$ as the appropriate number of iterations (Figure B0). We ran the process described above five

543

independent times to ensure that this relationship was similar with different random runs of the algorithm.




Figure B0: Relationship between the number of iterations in a MCMC chain used to simulate the foraging task and distributional similarity, measured using the earth mover's distance. We calculated the earth mover's distance between the first N iterations of the chain and the first $N + 500$ iterations to evaluate the difference that adding 500 iterations would make to the posterior distribution of animal behaviour. The coloured lines represent five individual runs of the process, and the thicker black line represents the mean earth mover's distance across these runs.

Literature Cited

- 546 Avgar, T., Baker, J. A., Brown, G. S., Hagens, J. S., Kittle, A. M., Mallon, E. E., McGreer, M. T.,
Mosser, A., Newmaster, S. G., Patterson, B. R., and et al. (2015). Space-use behaviour of wood-
land caribou based on a cognitive movement model. *Journal of Animal Ecology*, 84(4):1059–1070.
- 549 Avgar, T., Deardon, R., and Fryxell, J. M. (2013). An empirically parameterized individual based
model of animal movement, perception, and memory. *Ecological Modelling*, 251:158–172.
- Bavel, J. J. V., Baicker, K., Boggio, P. S., Capraro, V., Cichocka, A., Cikara, M., Crockett, M. J.,
552 Crum, A. J., Douglas, K. M., Druckman, J. N., Drury, J., Dube, O., Ellemers, N., Finkel, E. J.,
Fowler, J. H., Gelfand, M., Han, S., Haslam, S. A., Jetten, J., Kitayama, S., Mobbs, D., Napper,
L. E., Packer, D. J., Pennycook, G., Peters, E., Petty, R. E., Rand, D. G., Reicher, S. D., Schnall,
555 S., Shariff, A., Skitka, L. J., Smith, S. S., Sunstein, C. R., Tabri, N., Tucker, J. A., Linden, S.
v. d., Lange, P. v., Weeden, K. A., Wohl, M. J. A., Zaki, J., Zion, S. R., and Willer, R. (2020).
Using social and behavioural science to support covid-19 pandemic response. *Nature Human*
558 *Behaviour*, 4(5):460–471.
- Bell, G. and Collins, S. (2008). Adaptation, extinction and global change. *Evolutionary Applications*,
1(1):3–16.
- 561 Berger, J. O. (1985). *Statistical Decision Theory and Bayesian Analysis*. Springer Series in Statistics.
Springer New York.
- Berger-Tal, O. and Avgar, T. (2012). The glass is half-full: Overestimating the quality of a novel
564 environment is advantageous. *PLOS ONE*, 7(4):e34578.
- Berger-Tal, O. and Saltz, D. (2014). Using the movement patterns of reintroduced animals to
improve reintroduction success. *Current Zoology*, 60(4):515–526.
- 567 Bracis, C., Gurarie, E., Rutter, J. D., and Goodwin, R. A. (2018). Remembering the good and the

bad: memory-based mediation of the food–safety trade-off in dynamic landscapes. *Theoretical Ecology*, 11(3):305–319.

570 Cantrell, R. S., Cosner, C., Deangelis, D. L., and Padron, V. (2007). The ideal free distribution as an evolutionarily stable strategy. *Journal of Biological Dynamics*, 1(3):249–271.

573 Castellano, S., Cadeddu, G., and Cermelli, P. (2012). Computational mate choice: Theory and empirical evidence. *Behavioural Processes*, 90(2):261–277.

Charnov, E. L. (1976). Optimal foraging, the marginal value theorem. *Theoretical Population Biology*, 9(2):129–136.

576 Chevin, L.-M., Lande, R., and Mace, G. M. (2010). Adaptation, plasticity, and extinction in a changing environment: Towards a predictive theory. *PLOS Biology*, 8(4):e1000357.

579 Clayton, N. S. and Dickinson, A. (1998). Episodic-like memory during cache recovery by scrub jays. *Nature*, 395(6699):272–4.

Courville, A. C., Daw, N. D., and Touretzky, D. S. (2006). Bayesian theories of conditioning in a changing world. *Trends in Cognitive Sciences*, 10(7):294–300.

582 Cowles, M. K. and Carlin, B. P. (1996). Markov chain monte carlo convergence diagnostics: A comparative review. *Journal of the American Statistical Association*, 91(434):883–904.

585 Dall, S., Giraldeau, L., Olsson, O., Mcnamara, J., and Stephens, D. (2005). Information and its use by animals in evolutionary ecology. *Trends in Ecology & Evolution*, 20(4):187–193.

Dayan, P. and Daw, N. D. (2008). Decision theory, reinforcement learning, and the brain. *Cognitive, Affective, & Behavioral Neuroscience*, 8(4):429–453.

588 DeAngelis, D. L. and Diaz, S. G. (2019). Decision-making in agent-based modeling: A current review and future prospectus. *Frontiers in Ecology and Evolution*, 6:237.

- DeWitt, T. J., Sih, A., and Wilson, D. S. (1998). Costs and limits of phenotypic plasticity. *Trends in Ecology & Evolution*, 13(2):77–81.
- Donaldson-Matasci, M. C., Lachmann, . M., and Bergstrom, C. T. (2008). Phenotypic diversity as an adaptation to environmental uncertainty. *Evolutionary Ecology Research*, 10(4):493–515.
- Dunn, R. E., Wanless, S., Daunt, F., Harris, M. P., and Green, J. A. (2020). A year in the life of a north atlantic seabird: behavioural and energetic adjustments during the annual cycle. *Scientific Reports*, 10(11):5993.
- Ellison, A. M. (2004). Bayesian inference in ecology. *Ecology Letters*, 7(6):509–520.
- Fagan, W. F., Gurarie, E., Bewick, S., Howard, A., Cantrell, R. S., and Cosner, C. (2017). Perceptual ranges, information gathering, and foraging success in dynamic landscapes. *The American Naturalist*, 189(5):474–489.
- Fagan, W. F., Lewis, M. A., Auger-Méthé, M., Avgar, T., Benhamou, S., Breed, G., LaDage, L., Schlägel, U. E., Tang, W.-w., Papastamatiou, Y. P., and et al. (2013). Spatial memory and animal movement. *Ecology Letters*, 16(10):1316–1329.
- Falcón-Cortés, A., Boyer, D., Merrill, E., Frair, J. L., and Morales, J. M. (2021). Hierarchical, memory-based movement models for translocated elk (*cervus canadensis*). *Frontiers in Ecology and Evolution*, 9:497.
- Fletcher, R. J., Maxwell, C. W., Andrews, J. E., and Helmeý-Hartman, W. L. (2013). Signal detection theory clarifies the concept of perceptual range and its relevance to landscape connectivity. *Landscape Ecology*, 28(1):57–67.
- Fretwell, S. D. and Lucas, H. L. (1969). On territorial behavior and other factors influencing habitat distribution in birds. *Acta Biotheoretica*, 19(1):16–36.
- Gaillard, J.-M. and Yoccoz, N. G. (2003). Temporal variation in survival of mammals: A case of environmental canalization? *Ecology*, 84(12):3294–3306.

- Gehr, B., Bonnot, N. C., Heurich, M., Cagnacci, F., Ciuti, S., Hewison, A. J. M., Gaillard, J.-M.,
615 Ranc, N., Premier, J., Vogt, K., Hofer, E., Ryser, A., Vimercati, E., and Keller, L. (2020). Stay
home, stay safe—site familiarity reduces predation risk in a large herbivore in two contrasting
study sites. *Journal of Animal Ecology*, 89(6):1329–1339.
- 618 Gibbs, J. P. and Breisch, A. R. (2001). Climate warming and calling phenology of frogs near
Ithaca, New York, 1900–1999. *Conservation Biology*, 15(4):1175–1178.
- Gibson, G. and Wagner, G. (2000). Canalization in evolutionary genetics: a stabilizing theory?
621 *BioEssays*, 22(4):372–380.
- Green, R. F. (1980). Bayesian birds: A simple example of Oaten's stochastic model of optimal
foraging. *Theoretical Population Biology*, 18(2):244–256.
- 624 Green, R. F. (2006). A simpler, more general method of finding the optimal foraging strategy for
Bayesian birds. *Oikos*, 112(2):274–284.
- Griffiths, T. L., Kemp, C., and Tenenbaum, J. B. (2001). *Bayesian Models of Cognition*, page 59–100.
627 Cambridge University Press, 1 edition.
- Grimm, V., Berger, U., Bastiansen, F., Eliassen, S., Ginot, V., Giske, J., Goss-Custard, J., Grand, T.,
Heinz, S. K., Huse, G., and et al. (2006). A standard protocol for describing individual-based
630 and agent-based models. *Ecological Modelling*, 198(1):115–126.
- Haché, S., Villard, M.-A., and Bayne, E. M. (2013). Experimental evidence for an ideal free
distribution in a breeding population of a territorial songbird. *Ecology*, 94(4):861–869.
- 633 Harris, J. A. and Bouton, M. E. (2020). Pavlovian conditioning under partial reinforcement: The
effects of nonreinforced trials versus cumulative conditioned stimulus duration. *Journal of
Experimental Psychology: Animal Learning and Cognition*, 46(3):256–272.
- 636 Katz, J. S. and Wright, A. A. (2006). Same/different abstract-concept learning by pigeons. *Journal
of Experimental Psychology: Animal Behavior Processes*, 32(1):80–86.

- 639 Kemink, K. M. and Kesler, D. C. (2013). Using movement ecology to inform translocation efforts:
a case study with an endangered lekking bird species. *Animal Conservation*, 16(4):449–457.
- Kiltie, R. A. (2000). Scaling of visual acuity with body size in mammals and birds. *Functional Ecology*, 14(2):226–234.
- 642 Krebs, J. R., Kacelnik, A., and Taylor, P. (1978). Test of optimal sampling by foraging great tits.
Nature, 275(5675):27–31.
- Kriska, G., Horváth, G., and Andrikovics, S. (1998). Why do mayflies lay their eggs en
645 masse on dry asphalt roads? water-imitating polarized light reflected from asphalt attracts
ephemeroptera. *Journal of Experimental Biology*, 201(15):2273–2286.
- Lea, S. E., McLaren, I. P., Dow, S. M., and Graft, D. A. (2012). The cognitive mechanisms of
648 optimal sampling. *Behavioural Processes*, 89(2):77–85.
- Lele, S. R., Dennis, B., and Lutscher, F. (2007). Data cloning: easy maximum likelihood estimation
for complex ecological models using bayesian markov chain monte carlo methods. *Ecology*
651 *Letters*, 10(7):551–563.
- Lewis, M. A., Fagan, W. F., Auger-Méthé, M., Frair, J., Fryxell, J. M., Gros, C., Gurarie, E., Healy,
S. D., and Merkle, J. A. (2021). Learning and animal movement. *Frontiers in Ecology and*
654 *Evolution*, 9.
- Liefting, M., Hoffmann, A. A., and Ellers, J. (2009). Plasticity versus environmental canalization:
Population differences in thermal responses along a latitudinal gradient in *Drosophila serrata*.
657 *Evolution*, 63(8):1954–1963.
- Luttbeg, B. (1996). A comparative bayes tactic for mate assessment and choice. *Behavioral Ecology*,
7(4):451–460.
- 660 Macdonald, D. W. (1983). The ecology of carnivore social behaviour. *Nature*,
301(58995899):379–384.

- Macdonald, D. W. and Johnson, D. D. P. (2015). Patchwork planet: the resource dispersion
663 hypothesis, society, and the ecology of life. *Journal of Zoology*, 295(2):75–107.
- Masson-Delmotte, V., Zhai, P., Pirani, A., L., C. S., Péan, C., Berger, S., Caud, N., Chen, Y., Gold-
farb, L., Gomis, M. I., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J. B. R., Maycock, T. K.,
666 Waterfield, T., Yelekçi, O., Yu, R., and Zhou, B. (2021). Climate change 2021: The physical
science basis. contribution of working group i to the sixth assessment report of the intergov-
ernmental panel on climate change. Technical report, Intergovernmental Panel on Climate
669 Change.
- McNamara, J. and Houston, A. (1980). The application of statistical decision theory to animal
behaviour. *Journal of Theoretical Biology*, 85(4):673–690.
- 672 McNamara, J. M., Green, R. F., and Olsson, O. (2006). Bayes' theorem and its applications in
animal behaviour. *Oikos*, 112(2):243–251.
- Merilä, J. and Hendry, A. P. (2014). Climate change, adaptation, and phenotypic plasticity: the
675 problem and the evidence. *Evolutionary Applications*, 7(1):1–14.
- Mettke-Hofmann, C., Lorentzen, S., Schlicht, E., Schneider, J., and Werner, F. (2009). Spa-
tial neophilia and spatial neophobia in resident and migratory warblers (sylvia). *Ethology*,
678 115(5):482–492.
- Milligan, S., Brown, L., Hobson, D., Frame, P., and Stenhouse, G. (2018). Factors affecting the
success of grizzly bear translocations. *The Journal of Wildlife Management*, 82(3):519–530.
- 681 Murphy, K. J., Ciuti, S., and Kane, A. (2020). An introduction to agent-based models as an acces-
sible surrogate to field-based research and teaching. *Ecology and Evolution*, 10(22):12482–12498.
- Møller, A. P. (2009). Successful city dwellers: a comparative study of the ecological characteristics
684 of urban birds in the western palearctic. *Oecologia*, 159(4):849–858.
- Møller, A. P. and Erritzøe, J. (2010). Flight distance and eye size in birds. *Ethology*, 116(5):458–465.

- 687 Nevoux, M., Forcada, J., Barbraud, C., Croxall, J., and Weimerskirch, H. (2010). Bet-hedging response to environmental variability, an intraspecific comparison. *Ecology*, 91(8):2416–2427.
- Panakhova, E., Buresova, O., and Bures, J. (1984). Persistence of spatial memory in the morris water tank task. *International Journal of Psychophysiology*, 2(1):5–10.
- 690 Parrish, J. D. (2000). Behavioral, energetic, and conservation implications of foraging plasticity during migration. *Studies in Avian Biology*, 20:18.
- Pavlov, I. (1927). *Conditioned reflexes*. Oxford University Press.
- 693 Pearce, J. M. (1987). A model for stimulus generalization in pavlovian conditioning. *Psychological Review*, 94(1):61–73.
- Pearce, J. M. (2008). *Animal Learning and Cognition: An Introduction*. Psychology Press, 3 edition.
- 696 Polak, T. and Saltz, D. (2011). Reintroduction as an ecosystem restoration technique. *Conservation Biology*, 25(3):424–424.
- Potts, J. R., Auger-Méthé, M., Mokross, K., and Lewis, M. A. (2014). A generalized residual technique for analysing complex movement models using earth mover’s distance. *Methods in Ecology and Evolution*, 5(10):1012–1022.
- 702 Potts, J. R. and Lewis, M. A. (2016). How memory of direct animal interactions can lead to territorial pattern formation. *Journal of The Royal Society Interface*, 13(118):20160059.
- Raftery, A. E. and Lewis, S. (1992). How many iterations in the gibbs sampler? In *In Bayesian Statistics 4*, page 763–773. Oxford University Press.
- 705 Ranc, N., Moorcroft, P. R., Ossi, F., and Cagnacci, F. (2021). Experimental evidence of memory-based foraging decisions in a large wild mammal. *Proceedings of the National Academy of Sciences*, 118(15):e2014856118.

- 708 Rescorla, R. and Wagner, A. (1972). A theory of pavlovian conditioning: Variations in the effectiveness of reinforcement and nonreinforcement. In Black, A. and Prokasy, W., editors, *Classical conditioning II: Current research and theory*, pages 64–99. Appleton-Century-Crofts.
- 711 Robertson, B. A. and Chalfoun, A. D. (2016). Evolutionary traps as keys to understanding behavioral maladaptation. *Current Opinion in Behavioral Sciences*, 12:12–17.
- Robertson, B. A., Rehage, J. S., and Sih, A. (2013). Ecological novelty and the emergence of
714 evolutionary traps. *Trends in Ecology & Evolution*, 28(9):552–560.
- Schlather, M. (2012). Construction of covariance functions and unconditional simulation of random fields. In Porcu, E., Montero, J., and Schlather, M., editors, *Advances and Challenges in*
717 *Space-time Modelling of Natural Events*, Lecture Notes in Statistics, page 25–54. Springer.
- Schlägel, U. E. and Lewis, M. A. (2014). Detecting effects of spatial memory and dynamic information on animal movement decisions. *Methods in Ecology and Evolution*, 5(11):1236–1246.
- 720 Schmidt, K. A., Dall, S. R. X., and Gils, J. A. V. (2010). The ecology of information: an overview on the ecological significance of making informed decisions. *Oikos*, 119(2):304–316.
- Seddon, P. J., Armstrong, D. P., and Maloney, R. F. (2007). Developing the science of reintroduction biology. *Conservation Biology*, 21(2):303–312.
723
- Sengupta, B., Friston, K. J., and Penny, W. D. (2015). Gradient-free mcmc methods for dynamic causal modelling. *NeuroImage*, 112:375–381.
- 726 Snell-Rood, E. C. (2013). An overview of the evolutionary causes and consequences of behavioural plasticity. *Animal Behaviour*, 85(5):1004–1011.
- Spence, K. W. (1936). The nature of discrimination learning in animals. *Psychological Review*,
729 43(5):427–449.
- Tang, W. and Bennett, D. A. (2010). Agent-based modeling of animal movement: A review. *Geography Compass*, 4(7):682–700.

- 732 Thomas, D. A. and Riccio, D. C. (1979). Forgetting of a cs attribute in a conditioned suppression
paradigm. *Animal Learning & Behavior*, 7(2):191–195.
- Valone, T. J. (2006). Are animals capable of bayesian updating? an empirical review. *Oikos*,
735 112(2):252–259.
- van Baaren, J. and Candolin, U. (2018). Plasticity in a changing world: behavioural responses to
human perturbations. *Current Opinion in Insect Science*, 27:21–25.
- 738 Wong, B. B. and Candolin, U. (2015). Behavioral responses to changing environments. *Behavioral
Ecology*, 26(3):665–673.

References Cited Only in the Online Enhancements

- 741 Blackwell, P. G., Niu, M., Lambert, M. S., and LaPoint, S. D. (2016). Exact bayesian inference for
animal movement in continuous time. *Methods in Ecology and Evolution*, 7(2):184–195.
- McClintock, B. T., Johnson, D. S., Hooten, M. B., Hoef, J. M. V., and Morales, J. M. (2014). When to
744 be discrete: the importance of time formulation in understanding animal movement. *Movement
Ecology*, 2(21):14.
- Potts, J. R., Auger-Méthé, M., Mokross, K., and Lewis, M. A. (2014). A generalized residual
747 technique for analysing complex movement models using earth mover’s distance. *Methods in
Ecology and Evolution*, 5(10):1012–1022.
- Rubner, Y., Tomasi, C., and Guibas, L. J. (2000). The earth mover’s distance as a metric for image
750 retrieval. *International Journal of Computer Vision*, 40(2):99–121.
- Vaserstein, L. N. (1969). Markov processes over denumerable products of spaces, describing large
systems of automata. *Problemy Peredachi Informatsii*, 5(3):64–72.