

CANADIAN THESES ON MICROFICHE

I.S.B.N.

THESES CANADIENNES SUR MICROFICHE



National Library of Canada
Collections Development Branch

Canadian Theses on
Microfiche Service

Ottawa, Canada
K1A 0N4

Bibliothèque nationale du Canada
Direction du développement des collections

Service des thèses canadiennes
sur microfiche

NOTICE

The quality of this microfiche is heavily dependent upon the quality of the original thesis submitted for microfilming. Every effort has been made to ensure the highest quality of reproduction possible.

If pages are missing, contact the university which granted the degree.

Some pages may have indistinct print especially if the original pages were typed with a poor typewriter ribbon or if the university sent us a poor photocopy.

Previously copyrighted materials (journal articles, published tests, etc.) are not filmed.

Reproduction in full or in part of this film is governed by the Canadian Copyright Act, R.S.G. 1970, c. C-30. Please read the authorization forms which accompany this thesis.

THIS DISSERTATION
HAS BEEN MICROFILMED
EXACTLY AS RECEIVED

AVIS

La qualité de cette microfiche dépend grandement de la qualité de la thèse soumise au microfilmage. Nous avons tout fait pour assurer une qualité supérieure de reproduction.

S'il manque des pages, veuillez communiquer avec l'université qui a conféré le grade.

La qualité d'impression de certaines pages peut laisser à désirer, surtout si les pages originales ont été dactylographiées à l'aide d'un ruban usé ou si l'université nous a fait parvenir une photocopie de mauvaise qualité.

Les documents qui font déjà l'objet d'un droit d'auteur (articles de revue, examens publiés, etc.) ne sont pas microfilmés.

La reproduction, même partielle, de ce microfilm est soumise à la Loi canadienne sur le droit d'auteur, SRC 1970, c. C-30. Veuillez prendre connaissance des formules d'autorisation qui accompagnent cette thèse.

LA THÈSE A ÉTÉ
MICROFILMÉE TELLE QUE
NOUS L'AVONS REÇUE



National Library
of Canada

Bibliothèque nationale
du Canada

0-315-08911-3

Canadian Theses Division Division des thèses canadiennes

Ottawa, Canada
K1A 0N4

56828

PERMISSION TO MICROFILM — AUTORISATION DE MICROFILMER

• Please print or type — Ecrire en lettres moulees ou dactylographier

Full Name of Author — Nom complet de l'auteur

HENRIETTE ALBERTA KELKER

Date of Birth — Date de naissance

OCTOBER 11 1949

Country of Birth — Lieu de naissance

THE NETHERLANDS

Permanent Address — Résidence fixe

5410 MC ST. EDMONTON T6H 3E1

Title of Thesis — Titre de la thèse

THE EFFECTS OF INTERACTION BETWEEN GENOTYPES
ON SINGLE PLANT SELECTION OF WHEAT
(TRITICUM AESTIVUM)

University — Université

UNIVERSITY OF ALBERTA

Degree for which thesis was presented — Grade pour lequel cette thèse fut présentée

PH D

Year this degree conferred — Année d'obtention de ce grade

1982

Name of Supervisor — Nom du directeur de thèse

DR E A CLARK

Permission is hereby granted to the NATIONAL LIBRARY OF CANADA to microfilm this thesis and to lend or sell copies of the film.

The author reserves other publication rights, and neither the thesis nor extensive extracts from it may be printed or otherwise reproduced without the author's written permission.

L'autorisation est, par la présente, accordée à la BIBLIOTHÈQUE NATIONALE DU CANADA de microfilmer cette thèse et de prêter ou de vendre des exemplaires du film.

L'auteur se réserve les autres droits de publication, ni la thèse ni de longs extraits de celle-ci ne doivent être imprimés ou autrement reproduits sans l'autorisation écrite de l'auteur.

Date

APRIL 23 1982

Signature

H A Kelker

THE UNIVERSITY OF ALBERTA

THE EFFECTS OF INTERACTION BETWEEN GENOTYPES ON SINGLE PLANT
SELECTION OF WHEAT (*TRITICUM AESTIVUM*)

by

HENRIETTE A. KELKER

A THESIS

SUBMITTED TO THE FACULTY OF GRADUATE STUDIES AND RESEARCH
IN PARTIAL FULFILMENT OF THE REQUIREMENTS FOR THE DEGREE

OF DOCTOR OF PHILOSOPHY

IN

PLANT BREEDING

DEPARTMENT OF PLANT SCIENCE

EDMONTON, ALBERTA

1982

THE UNIVERSITY OF ALBERTA

RELEASE FORM

NAME OF AUTHOR HENRIETTE A. KELKER
TITLE OF THESIS THE EFFECTS OF INTERACTION BETWEEN
GENOTYPES ON SINGLE PLANT SELECTION OF
WHEAT (TRITICUM AESTIVUM)

DEGREE FOR WHICH THESIS WAS PRESENTED DOCTOR OF PHILOSOPHY
YEAR THIS DEGREE GRANTED 1982

Permission is hereby granted to THE UNIVERSITY OF
ALBERTA LIBRARY to reproduce single copies of this
thesis and to lend or sell such copies for private,
scholarly or scientific research purposes only.

The author reserves other publication rights, and
neither the thesis nor extensive extracts from it may
be printed or otherwise reproduced without the author's
written permission.

(SIGNED) *H.A. Kelker*.....

PERMANENT ADDRESS:

...5416...110...st.....
...Edmonton.....
...Alberta...T6H1Z1...

DATED ..April...23....1982

THE UNIVERSITY OF ALBERTA
FACULTY OF GRADUATE STUDIES AND RESEARCH

The undersigned certify that they have read, and recommend to the Faculty of Graduate Studies and Research, for acceptance, a thesis entitled THE EFFECTS OF INTERACTION BETWEEN GENOTYPES ON SINGLE PLANT SELECTION OF WHEAT (*TRITICUM AESTIVUM*) submitted by HENRIETTE A. KELKER in partial fulfilment of the requirements for the degree of DOCTOR OF PHILOSOPHY in PLANT BREEDING.

..... *E. Ann Clark*

Supervisor

..... *Arthur D. Strickland*

..... *John H. ...*

External Examiner

Date..... *September 30, 1981*

Abstract

In the first year of the study, the response of wheat plants to genotype interaction was studied on monocultures and mixtures of seven wheat genotypes, at seven different interplant spacings. The genotypes differed significantly in their response to genotype interaction.

In the second year, mixtures and monocultures of the same genotypes were grown at two different interplant spacings. The mean variation of the number of tillers, plant weight and yield of the genotypes grown in mixtures at the closer spacing, and of kernel weight at the wider spacing, differed significantly from the mean variation in monoculture. For the plants grown in mixtures, an increase in spacing was accompanied by an increase in the plant to plant variation of the number of tillers, the number of heads, plant weight, yield, kernels per plant, kernel weight and harvest index. Similar effects were observed for plants grown in monocultures.

With the exception of height of the flag leaf, plant height and harvest index, the effect of genotype interaction on plant characters differed significantly among genotypes. Correlations between pairs of characters, measured on the same genotypes and their associates, revealed that the characters which affect (or respond to) genotype interaction are not the same at different spacings and in different genotypes.

In the third year, the effect of initial seed size on plant interactions in five monocultures was examined. Number

of tillers, number of heads, and kernel weight were affected by initial seed size in plots from uniform seed. Plants from large seeds produced more tillers than did plants from small seeds, but plants from small seed produced larger kernels than did plants from large seeds. In plots from mixed seed sizes, however, neighbor interaction caused plants from small seeds to produce shorter plants, bearing lighter seeds, than did plants from large seed.

The data demonstrated that the distribution of many characters, in monoculture as well as in mixture, is not normal. It was shown that, if one selects the highest values for a character, characters which have a negatively skewed distribution are selected with less error than are characters for which the distribution is normal or positively skewed.

Yield trials of mixtures and monocultures were repeated for three years. Only in 1977 did the mixtures yield significantly more than the mean of the monocultures of the components. Interactions between the genotypes differed from year to year.

It was concluded, that the efficiency of selection of single plants from a segregating population is affected by variation due to different initial seed sizes, variation due to interaction between plants from different size seeds, and variation due to interaction between different genotypes, as well as by the frequency distribution of the observed characters.

It was recommended that, in order to evaluate the suitability of a character for selection in early generations, the distribution of this character in a multi-component mixture, as well as its sensitivity to genotype interaction, should be evaluated in conjunction with the estimation of its heritability.

Acknowledgement

I wish to thank my advisor, Dr. K. G. Briggs, for encouraging me to undertake this project, and for his interest and advice during the following years of study.

I am indebted to Dr. E. A. Clark, for accepting the position of substitute advisor, for her critical reading of the manuscript, and for her helpful comments.

I thankfully acknowledge the help which I received from Mr. K. Kutchera and the staff at Parkland farm, with all field operations and the collection of data.

The financial assistance of the Canadian Wheat Board is greatly appreciated.

Lastly I would like to thank my husband, Douglas, for his support, patience and knowledgable advice, and my children for providing me with much needed diversion.

Table of Contents

Chapter	Page
1. Introduction	1
2. Literature Review	3
2.1 Terminology	3
2.2 Mixtures vs. monocultures	5
2.3 Types of interaction and interaction effects	9
2.4 Effects of plant interaction on plant growth	14
2.4.1 Plasticity of plant characters	14
2.4.2 Competition for water and nutrients	18
2.4.3 Competition for light	20
2.4.4 Effects of seed size on plant interactions	23
2.4.5 Effects of density on plant interactions ..	28
2.4.6 Changes in the distribution of characters during growth	33
2.4.7 Effects of genotype interaction on grain filling and maturity	35
2.5 Effects of interaction between genotypes on single plant selection	38
3. Short Description of the Experiments	44
4. Materials and Methods - General Information	46
4.1 Characteristics of test sites	46
4.2 Plant material used	47
5. Interaction Between Genotypes at Different Interplant Spacings	49
5.1 Materials and methods	49
5.1.1 Field plot design	49
5.1.2 Seed preparation and data collection	50

5.2	Analysis of the data	52
5.3	Results	52
5.4	Discussion	57
	Relationships between Plant Characters Measured on Mixed Genotypes	58
6.1	Materials and methods	58
6.2	Analysis of the data	60
6.3	Results	63
6.3.1	Effects of genotype interaction on plant-to-plant variation	64
6.3.2	Effects of genotype interaction on the expression of various plant characters ...	68
6.3.3	Relative performance of genotypes in mixture	69
6.3.4	Relationships between characters of interacting genotypes	76
6.3.5	Interaction between genotypes during the early stages of growth	79
6.4	Discussion	81
6.4.1	Effects of genotype interaction on plant-to-plant variation	81
6.4.2	Effects of genotype interaction on mean plant performance	84
7.	Effects of Seed Size on Plant Interaction	88
7.1	Materials and methods	88
7.2	Analysis of the data	89
7.3	Results	93
7.3.1	Effects of seed size on mean plant performance and sample distribution of characters	93
7.3.2	Effects of seed size on interaction between neighbours	102
7.4	Discussion	106

8.	The Relationship Between Mixture Yield and Monoculture Yield of the Components.	110
8.1	Materials and methods	110
8.2	Analysis of the data	111
8.3	Results	118
8.3.1	Effects of genotype interaction on plot yields	120
8.3.2	Mixture efficiency and its correlation with yield	123
8.3.3	Diallel analysis of variance	126
8.3.4	Mixture efficiency	128
8.3.5	Regression of W on V	129
8.4	Discussion	132
9.	Discussion of the Results with Respect to Single Plant Selection in Early Generations	135
9.1	Effects of non-normal character distribution on single plant selection	136
9.2	Effects of genotype interaction on single plant selection	139
9.3	Effects of interplant spacing on single plant selection	140
9.4	Effects of seed size on single plant selection	144
9.5	Suitability of various characters for single plant selection	145
10.	References	149
11.	List of Appendices	167

List of Tables

Table	Page
1. The effect of genotype, genotype mixing and interplant spacing on single plant characters, as determined by analyses of variance.....	55
2. Comparison of the means of characters of single plants, grown in monoculture and in mixture, at various plant densities, by means of Students t-test.....	56
3. The effects of genotype, genotype mixing and interplant spacing, on the variance of single plant characters, as determined by analyses of variance.....	65
4. The effects of genotype mixing and interplant spacing on the average variation among plants, of single plant characters, as determined by a variance ratio F-test.....	66
5. The effect of genotype, genotype mixing and interplant spacing on single plant characters, as determined by analyses of variance.....	70
6. Comparison of single plant characters of four	

genotypes in monoculture and in mixture, at two interplant spacings, by means of 'Relative Yield' values.....	71
7. Correlations of single plant characters of five genotypes, with characters of associated mixture components, at two interplant spacings.....	77
8. Correlations of single plant characters of five genotypes, with characters of associated mixture components, in machine seeded plots, at two growth stages.....	80
9. Plant characters which were measured or derived at four stages of growth in 1979.....	90
10. The effect of genotype and seed treatment on the variance of single plant characters, at different stages of growth, as determined by analyses of variance.....	94
11. The effect of growth stage, genotype, and seed treatment on the variance and skewness of single plant characters, as determined by analyses of variance.....	95
12a. The effect of genotype and seed treatment	

on single plant characters, measured at the 3-5 leaf stage, as determined by analyses of variance.....	97
12b. The effect of genotype and seed treatment on single plant characters, measured at the jointing stage, as determined by analyses of variance.....	98
12c. The effect of genotype and seed treatment on single plant characters, measured at the heading stage, as determined by analyses of variance.....	99
12d. The effect of genotype and seed treatment on single plant characters, measured at maturity, as determined by analyses of variance.....	100
13. Correlated characters of neighbouring plants, at different stages of growth, in rows of plants from large seeds, small seeds or alternated large and small seeds.....	104
14. Frequency distribution of mixture yield, relative to the yield of the components grown in monoculture.....	119
15. The effect of year, genotype monoculture and genotype mixing on plot yield, as determined by an analysis of variance.....	121

16. The effect of genotype monoculture and genotype mixing on plot yield in each of three test years, as determined by analyses of variance.....122
17. Relationships between mixture yield and Specific Mixture Efficiency, within and between years, as expressed by Spearmans rank-order correlation coefficient.....124
18. Rank-orders of the monoculture yields of eight genotypes, and the mean yields of their mixtures, as observed at the Edmonton and Ellerslie locations..125
19. The average array and specific effects of genotype on mixture yield, as determined by diallel analyses of variance for each of the three test years.....127
20. Simple correlations between monoculture yield and Average Mixture Efficiency, in 1977, 1978, and 1979.....130
- 21a. Estimates of heritability of single plant characters, grown at a 3 cm interplant spacing, as computed from monoculture and mixture data.....141
- 21b. Estimates of heritability of single plant characters, grown at a 9 cm interplant spacing,

as computed from monoculture and mixture data.....142

List of Figures

Figure	Page
1a-d. Graphic illustration of four possible types of genotype interaction effects on genotype and mixture yield of two genotypes, A and B, grown in binary mixtures of varying proportions ('Replacement Series').....	12
2a-e. Graphic illustration of the genotype interaction effects on genotype and mixture yield in five genotype mixtures, each grown at two different interplant spacings in 1978.....	75
3a-c. V-W regression lines, illustrating the relative ability of genotypes to dominate in a mixture, in the years 1977, 1978 and 1979.....	131
4. The effect of skewness and kurtosis of the distribution of a plant character on single plant selection.....	138

1. Introduction

In the history of wheat cultivation, the use of homogeneous cultivars is relatively new. Cultivated wheat populations used to be, and in some countries still are, composed of a large number of genotypes. For centuries, selection was practised by farmers for the plant types best suited for their needs. This resulted in plant types with large free threshing seeds, and increased yields, with better flour quality and with adaptation to a wide range of climatic conditions (Feldman, 1974). Although the principles of selection have been practiced in a more methodical way since the second half of the 18th century (Darwin, 1859), plant breeding has been recognised as a science only since the last part of the 19th century (Allard, 1960).

Almost all current selection methods for small cereals are based on line selection and its modifications (Sneep et al., 1979). These involve the selection of single plants or progeny rows in the early segregating generations.

It is widely recognised that single plant selection for quantitative characters, such as seed yield, in the segregating generations of a breeding program, is largely ineffective. Several studies have attempted to associate various plant characters in the F₂ or F₃ generations, with the expression of characters in more advanced generations (McGinnis and Shebeski, 1968; Hamblin and Donald, 1974). Interaction between genotypes is considered to be an important cause of the phenotypic variation which hinders

single plant selection (Kiesselbach, 1923; Christian and Gray, 1941). Yet, the effects of specific characters of one plant on the phenotype of its associates, and the implications of these effects for single plant selection, are unclear.

The present study focuses on the interactions between genotypes of common spring wheat (*Triticum aestivum* L.). The study was undertaken to determine the responsiveness of specific plant characters to the presence of other genotypes, and to evaluate the factors associated with the ability to perform well in a mixture situation. The results are discussed in relation to current plant breeding practices.

2. Literature Review

2.1 Terminology

Plant interactions have been studied by scientists from various branches of science, and each discipline has its own definitions and nomenclature. In 'The Origin of Species', Darwin (1859) discussed the competition which takes place between individuals within both the plant and the animal kingdoms. He preferred to use the term 'struggle for existence'. The term 'competition' was used by Odum (1971), who stated that competition, in the broadest sense, refers to the interaction of two organisms striving for the same thing. Harper (1961) described the different approaches of agronomists, ecologists and geneticists to plant competition. From an agronomist's point of view, competition is 'the response of plants to density-induced shortages'. The ecologist considers competition as 'all those forces by which one organism succeeds at the expense of another', while the geneticist deals with the 'effect of interaction operating between individuals of different genotypes within a population'.

A frequently quoted definition of competition was given by Clements et al. (1929), and reads as follows:

'Competition arises from the reaction of one plant upon the physical factors about it and the effect of the modified factors upon its competitors. In the exact sense, two plants

do not compete with each other as long as the water content, the nutrient material, the light and the heat are in excess of the needs of both. When the immediate supply of a single necessary factor falls below the combined demands of the plants competition begins.'

Because the word competition is commonly associated with negative effects, its use has been avoided by several authors. Harper (1961) preferred the term 'interference', while Trenbath (1974, 1975) and Trenbath and Harper (1973) used the term 'neighbour effects'. Mather (1961) distinguished 'competition' from 'cooperation', and pointed out that individuals which are competing in one respect may be cooperating in another.

Hall (1974) attributed the responses of plants to their neighbours to both competitive and non-competitive interference. Competitive interference, as defined by Hall, is the same as competition for essential growth factors, as defined by Clements et al. (1929). Non-competitive interference occurs when an individual responds to changes in its environment, which are caused by other plants and which do not fall into the first category. As an example of non-competitive interference, Hall named the positive effect which a legume can have on the growth of a non-legume. This happens when nitrogen, fixed by the legume, becomes available to the non-legume through mineralisation (Henzell and Vallis, 1977).

The word 'interference' implies a one-way process. The mutual influences of plants upon each other can therefore perhaps be best described by the word 'interaction'. The effects of interaction between genotypes are measured as the phenotypic deviations of plants grown in mixed culture from those grown in pure culture, under otherwise similar growing conditions.

2.2 Mixtures vs. monocultures

The literature on mixtures of crop cultivars goes back at least as far as the last part of the 19th century, and the value and practicality of mixtures as a crop has been a much disputed issue. Von Rümker (1892) reviewed several of the older European studies and advocated the use of mixtures in an effort to increase the yield per unit area, especially where it concerned grains used for animal feed.

Montgomery (1912) concluded that, on the average, both oats and winter wheat had higher grain yields when grown in cultivar mixtures than when grown alone. Unfortunately he examined only two cultivars of each species in this study. Similar observations were made on wheat by Engelke (1935), who also noted that total tiller density was always higher in mixtures than in monocultures.

Data on a mixture diallel of four wheat cultivars (Nuding, 1936) were reanalysed by Simmonds (1962). He subsequently showed that the mean yield of the components in

monoculture was negatively correlated with their performance in mixture. The yields of the six mixtures all exceeded the mean yield of their component monocultures, and the highest yields were obtained from mixtures of genotypes which were known to have a wide range of adaptation.

In contrast to the above reports, Heuser (1938) concluded from the trials of five wheat cultivars and their binary mixtures, that monocultures yield at least as well as, and sometimes better than, mixtures do.

Three genotypes of wheat, which were similar in morphological characters, development and yield, produced yields in mixture which were similar to their yield in monoculture (Frankel, 1939). The similarity of the genotypes in this study might explain the lack of effect from genotype interaction.

A review of the more recent literature on the performance of mixtures of genotypes and mixtures of species was made by Trenbath (1974). The reviewed studies examined the biomass production of several forage, cereal and oil seed species. In 344 two-component mixtures, 60% of the mixtures - which is significantly ($\alpha \leq 0.01$) more than half - yielded more than the mean of the component monocultures, while 24% yielded more than the higher yielding monoculture. This suggests that, on the average, biomass production per plant is better in mixtures than in monocultures.

After evaluating a number of different types of rice (*Oryza* sp. L.) mixtures, mixing techniques and cultural

practices, Roy (1960) concluded that the chances for gaining or for losing yield potential through the formation of mixtures are about equal and that most advantages of mixtures are obtained when the soil conditions are poor. An advantage of mixtures over monocultures under extreme environmental conditions was also observed for barley (*Hordeum vulgare* L.) grain yield (Clay and Allard, 1969), in a study involving twentythree mixtures, evaluated over two years and five locations.

• The performance of a mixture of two wheat cultivars (Pitic 62 and Neepawa), sown at various seeding rates differed in the two years of the study (Baker, 1977). This was ascribed to the fact that Pitic 62 was better able to compensate for poor survival after early drought conditions, when grown in pure stand, than when grown in mixture. The reason for this different ability to recover was not discussed, however. Further study of this observation could clarify the process of interaction which takes place between these two wheat genotypes.

Simmonds (1962) wrote an extensive review on the variation in crop plants, and indicated that mixtures, besides being often slightly superior in yield, are usually also more reliable in their performance, especially under conditions of stress.

It thus appears that in the majority of cases, stress conditions affect mixtures less than they do monocultures. With this in mind, it is interesting to note that in a study

by Shorter and Frey (1979), which was conducted over two years and at two locations, the genotype x year, the genotype x location, and the genotype x year x location interactions were insignificant most of the time (12 out of 16) for mixtures, and were always significant for monocultures.

For lima beans (*Phaseolus lunatus* L.), it was observed that the stability of performance is related to the degree of genetic variation in the population. Allard (1961) studied the performance of ten lima bean populations: three monocultures, three two-component mixtures, one three-component mixture and the advanced generations (F7 and F9) from crosses between these three genotypes. He found that, although the monocultures performed most often either very poorly or very well, the mixtures consistently performed at a mediocre level, while the bulk populations performed well most of the time. He concluded that the genetic diversity of the mixed and bulk populations apparently made them more stable.

The yield of five two- and three-way mixtures of barley genotypes appeared to be more stable over years and locations than were the yields of the component genotypes grown in monoculture (Early and Qualset, 1971). At the same time, a tendency was observed for the plant-to-plant variation within a genotype to be greater in mixtures than in monocultures. This is perhaps not surprising, since the micro-environment within the crop stand also is more

variable in a mixture of plant types than in a monoculture.

The reviewed studies suggest that mixtures often perform in a more stable manner than do monocultures. In order to evaluate the relative performance of mixtures and monocultures, it is therefore advisable to repeat any study of competition effects for a number of years. Close observation of single plant characters under both mixture and monoculture conditions during these years could reveal the mechanism underlying the stability of mixtures, and could possibly explain why certain genotypes are better combiners in a mixture

2.3 Types of interaction and interaction effects

Interactions between plants occur in monoculture as well as in mixed populations. Since developmental patterns of genotypes within a species can differ considerably, interplant relations within a mixture of genotypes will differ from those in monoculture. Interaction between plants of two different species or two genotypes of the same species can be detrimental, neutral or beneficial to either or both associates. Depending on the degree of interaction and the proportion in which each component occurs in the mixture, the mixture yields can be less than, equal to, or more than what can be expected from the monocultures of each component.

Since interactions vary with the growth stage of each of the plants involved, variation in the relationships between the individuals of a population can be brought about by varying the relative growth stages of the interacting plants. This can be achieved through variations in the time of emergence (Ross and Harper, 1972), variations in the seeding depth (Black, 1956), or variations in seed size of the population members (Montgomery, 1912; Kaufmann and McFadden, 1960). Interactions between components of a mixture can also be varied through variations in density (Kira et al., 1953; Chebib et al., 1973) or variations in the proportions of the mixture components (de Wit, 1960; Hill, 1974).

The effect on mixture yield of varying proportions of mixture components at a constant overall density has been studied by de Wit (1960) and by Hill (1974). De Wit introduced the term 'replacement series' for experiments, in which the performance of monocultures and mixtures containing a range of different proportions of two types of plants are assessed.

In de Wit's experiments, population pressure was maintained at a constant level by using optimal seeding rates appropriate for each of the components. Thus, in an experiment involving various proportions of oats and peas, oats were given 31 cm²/kernel and peas were given 139 cm²/kernel in all plots (de Wit, 1961).

De Wit's model assumes that the genotypes in a mixture exclude each other and crowd for the same 'space'. Space is assumed to be uniformly distributed and encompasses all essential growth factors. De Wit considered it undesirable to define these factors, since: 'such a description is not necessary, always inaccurate and therefore inadvisable.' (de Wit, 1960). Hall (1974) explained, however, that identification of a particular limiting factor could give valuable insight into the interaction process.

A replacement diagram consists of curves which represent the measurements on yield or other characters for each mixture component, obtained from a series of mixtures of different proportions, as well as a curve representing the sum of the component yields (total mixture yield) for each mixture. These diagrams can take on any of the basic shapes given in the figures 1a to 1d, and can be explained as follows, using plot yield as an example:

1. Interaction between genotypes has no effect on the yield of either one of the mixture components (Figure 1a).
2. Interaction between genotypes decreases the yield of genotype A, while it increases the yield of B (Figure 1b). The combined effects, however, cause no change in the total yield of the mixture. One can speak in this case of 'complementary competition' (Shuts et al., 1968).
3. Interaction between genotypes decreases the yields of both A and B (Figure 1c). This can result in a total




Fig. 1a.d Graphic illustration of four possible types of genotype interaction effects on genotype and mixture yield of two genotypes, A and B, grown in binary mixtures of varying proportions ('Replacement Series').

%A and %B in the mixture are marked along the X-axis, reading from left to right for genotype A, and from right to left for genotype B.

mixture yield intermediate between the yields of A and B grown in monoculture (section 'p') or in a total mixture yield which is less than the monoculture yield of the lower yielding genotype (section 'q'). In the latter case one can speak of 'underyielding' of the mixture (Trenbath, 1974).

4. Interaction between genotypes increases the yields of both A and B (Figure 1d), resulting in a total mixture yield which is lower than the yield of the better monoculture, but higher than the yield of the poorer monoculture (section 'p'), or in the mixture yield being higher than the yield of the better monoculture (section 'q'). In the latter case one can speak of 'overyielding' of the mixture (Trenbath, 1974).

An understanding of the mechanisms leading to overyielding could be exploited to increase yield. In general, overyielding occurs in mixtures of species which exhibit marked differences in their pattern of growth. These differences provide the potential to utilise the available resources more efficiently (de Wit, 1961; Trenbath, 1974; Clark, 1980). In herbage plants, Hill (1974) demonstrated that a 50:50 mixture is not usually the optimal composition to maximise forage yield. This conclusion agrees with the diagram in Figure 1d, which shows that overyielding is more likely to occur in mixtures containing a higher proportion of the better yielding genotype.

Differences between genotypes of cultivated cereals are probably more subtle than are those between the components of a mixture of species, because plants of the same species are more similar in growth habit and nutrient requirements. Nevertheless, the presence of genetic differences implies the existence of differences in the various aspects of plant growth. Thus, the potential exists for complicated interplant relationships, which are different from those occurring in a monoculture.

A great number of factors play a role in plant interactions. There is therefore little hope that a generalised set of rules can be designed from which the outcome of plant interactions can be accurately predicted. Hill (1974) recognised this when he presented a theoretical model for the prediction of the value of binary mixtures of herbage species. He emphasised that plant interaction is a dynamic phenomenon, and that relationships can be transient.

2.4 Effects of plant interaction on plant growth

2.4.1 Plasticity of plant characters

The ability of plant parts and processes to respond to variations in the environment has been extensively discussed by Bradshaw (1964). He presented evidence that the plasticity of a certain character

- a. is specific for that character,

- b. is specific in relation to particular environmental influences,
- c. is specific in direction,
- d. is under genetic control, not necessarily related to heterozygosity, and
- e. can be radically altered by selection.

His theories on the role of plasticity of plant characters in evolution and adaptation apply directly to the interpretation of interactions between plants of different genotype. Bradshaw proposed that the plasticity of a character is related to the duration of meristematic activity which pertains to its development. Stemplength, for example, is controlled by long periods of meristematic activity and will therefore be more susceptible to environmental influences than are characters which are formed rapidly, such as reproductive structures. However, within this framework, there exists great variation among species as well as among genotypes within species.

Bradshaw explained that as a result of natural selection, a character whose stability is important for survival will likely show greater stability than will a character for which some plasticity is not a disadvantage. Harper (1961) pointed out that seed size is often surprisingly stable, thus ensuring an equal starting capital for each seedling. Significant variations in seed size due to interaction between genotypes, however, have been reported for wheat

(Christian and Gray, 1944; Chapman et al., 1969).

For some characters, plasticity can play a great role in adaptation. Height of many species appears to be a very plastic character. Trenbath and Harper (1973) described the different abilities of several oat species (*Avena* sp. L.) to adapt to different competitive situations through changes in height. *Avena sativa* was found to be capable of a stem extension of 10 cm above its monoculture value when grown in association with the taller *A. ludoviciana*. This resulted, for *A. sativa*, in a 20% increase in kernel weight over its monoculture value. They discussed the biological significance of this 'extension response' and concluded that such a mechanism must lead to a more even sharing of radiation. In natural communities this would conserve genotypes and thus diversity.

In corn the rapid elongation rate of one plant has an accelerating effect on the elongation rate of its neighbours (Hozumi et al., 1955). This effect appeared to be transient, however, resulting in an oscillating pattern of elongation rate in each plant.

Because environmental conditions in the field vary a great deal, stability of certain characters necessarily implies plasticity in others. Plant height, for example, is in a monoculture usually very uniform. Plasticity must therefore exist in some characters which determine plant height, such as elongation rate at various stages of development. Plasticity of the components of a certain

character can result in either stability or plasticity of the resultant character. Plant weight and grain yield of cereals are not considered as stable characters, nor are most of their components. However, harvest index, which is the ratio of grain yield and plant weight, is surprisingly stable over a range of environments (Donald and Hamblin, 1976).

If the degree of plasticity of various characters is genotype specific, as suggested by Bradshaw, then this could have important implications for a genotypes ability to perform in a mixture. Sakai (1955) concluded from a study of twelve barley cultivars that this 'competitive ability' was not associated with any of the characters he measured: plant height, maturity, seed size, habit of growth (erect vs. prostrate), heading habit (spring or winter) or grain yield. He concluded that competitive ability is a biological character in its own right and is under genetic control.

This conclusion has been criticised by several authors (Donald, 1963; Harper, 1965). Harper (1965) pointed out the incompleteness of Sakai's list of characters and stressed the importance of further investigations involving characters such as depth and extent of the root system, leaf area and height of the flag leaf. Sakai's theory on competitive ability as a genetic character is in agreement, however, with Bradshaw's views (1964) on the genetic control of the plasticity of each individual character.

2.4.2 Competition for water and nutrients

Suneson (1949) described experiments involving a mixture of four similarly adapted barley cultivars. Over sixteen years, two of the four components virtually disappeared from the population, even though one of them, Vaughn, had a significantly better yield potential and resistance to leaf diseases than any of the others when grown in pure stand. Atlas, the cultivar which eventually dominated the mixture, had the poorest leaf disease record and a mean yield below the median of the component cultivars.

Subsequent attempts to analyse the relationships between Atlas and Vaughn revealed that competition for light was not a decisive factor (Edwards and Allard, 1963), but that competition was primarily for moisture and nutrients (Hartmann and Allard, 1964). Lee (1960) observed that the response to competition between the two cultivars appeared abruptly at the jointing stage, when Atlas developed a dense mass of crown roots. Since Vaughn did not develop as large a root mass, Lee suggested that Atlas would become more efficient in gathering nutrients and would have the advantage when both varieties have to draw from the same soil mass.

Competition between roots usually starts long before the shoots are sufficiently developed to cause significant mutual shading (Milthorpe, 1961). But, competition for nutrients often leads to competition for light, since an

unevenly distributed nutrient supply will result in a similar variation in shoot growth. The study of nutrient competition without consideration of competition for light is therefore possible only under conditions which artificially prevent intermingling of the leaves of neighbouring plants.

Such studies have been performed by Donald (1958) on a mixture of grasses and by Aspinall and Milthorpe (1959) on barley and white persicaria (*Polygonum lepathifolium* L.) Both studies involved mixtures of species grown in pots.

Aspinall (1960) concluded that root competition, presumably for nutrients, since water was supplied in adequate amounts, reduced growth of the less successful component more than did shoot competition. Donald emphasised the interaction which occurs between competition for light, competition for nutrients, and factors indirectly affected by the two.

Both authors mentioned differences in embryo size and in the size of the root systems of species as the main causes of competitive interference in the early stages. These same factors should then play a role in pure stands if random variation occurs in initial seed size and rate of germination.

2.4.3 Competition for light

For most crop species, photosynthetic rate in an individual leaf is saturated at 150,000 - 200,000 lux (Donald, 1961). This is well below the flux density experienced by the top layer of a canopy on a sunny day. In a closed crop stand a large part of the foliage is shaded and will receive light intensities which are below the level of light saturation, except perhaps during the midday hours on a very bright day.

Horizontal as well as vertical distribution of the leaves and leaf angle will affect the amount of light which can be intercepted and utilised by a crop. Erect leaves will allow more light to penetrate into the canopy, enabling a larger leaf area to take part in photosynthesis.

Competition for light within a canopy will not occur until the photosynthetically active radiation which reaches the surface of the leaf falls below the saturation level. With higher light intensities, competition will occur deeper inside the canopy.

Donald (1967) described the ideal wheat plant for cultivation at high densities, such as are customary for most cereals, and suggested that it should have small erect leaves. This would maximise the area which is illuminated. Nichiporovich (1967) also discussed this subject and stressed that a different leaf angle will be optimal at different latitudes or climates. According to the latter, some cereals have an arrangement of leaves in which the top

leaves are approximately vertical, the middle leaves are inclined at intermediate angles and the lower leaves are nearly horizontal. Nichiporovich considered this the ideal arrangement for plants to be grown at high densities. In a mixture of genotypes, this type of plant would also be less susceptible to shading by taller plants, than would types with more horizontal leaves.

Donald (1961) stated that competition for light takes place between leaves rather than between plants or species. This is well illustrated by the studies of Brougham (1958) on competition for light in a sward of white clover. It was shown that, following defoliation, mature petiole length of successive leaves became longer, suggesting an attempt to reach daylight through the increasingly dense canopy.

Because the growth patterns of genotypes differ from each other, the start of plant interactions and their effects on subsequent processes will be different in mixtures than in monocultures. Trenbath and Harper (1973) found that, under optimal moisture and nutrient conditions, reactions of *Avena* species to genotype interaction could be explained in terms of the shading experienced at different stages of development. Early shading appeared to have a depressive effect on tiller number, while shading of the flag leaf during grain filling had a pronounced effect on seed weight.

The effects of shading on wheat at different points in development were studied by Fischer (1975). He found that

the period of rapid spike growth was the most sensitive to shading, resulting in fewer grains per spikelet and low tiller survival.

A negative correlation between shoot elongation rate and shoot length tended to equalise the height of corn plants during the first few weeks of growth (Hozumi et al., 1955). This effect caused temporary variations in the shoot elongation rate of individual plants. Later in the season, when competition became more severe, this correlation changed into a positive one, suggesting suppression of small plants by taller neighbors.

Growth rate of the high yielding dwarf wheat cultivar Yecora was found to be sensitive to low shading intensities, of three to four weeks duration, at various times during the growing season (Fischer, 1975). Shading at any time reduced crop growth rate. The strongest reductions in yield, however, occurred when the plants were shaded during the month before anthesis. Although the shading treatments in these experiments involved complete plots, one can expect shading by taller neighbours to have similar effects.

Yoda et al. (1957) found that the growth of rose mallow (*Hibiscus moschentos* L.) seedlings, during the latter part of the growing season, was more closely related to the relative size rather than to the absolute size of neighbouring plants. They discussed Hozumi's findings as well as their own, and concluded that the change from a negative to a positive correlation between plant size and

growth rate marked the point where the growth reductions due to reduced photosynthesis surpassed the etiolating effects of competition for light. The change from a negative to a positive correlation occurred earlier in dense plantings than at wider interplant spacings, supporting this theory.

An increase in the mean plant height of a mixed wheat population was observed by Khalifa and Qualset (1975), who planted the progeny of a cross between short statured and standard height wheats, without artificial selection, for six generations. They found that the frequency of short statured plants in the population decreased, suggesting that tall plants were more successful in setting seed, in spite of a significant negative correlation between height and yield.

In a similar study Busch and Luizzi (1979) found no evidence of such a directional shift. They ascribed this lack of change to either a lesser genetic range in height in their material or to the lesser expression of height differences in the more limiting dryland environment of North Dakota, compared to that of California.

2.4.4 Effects of seed size on plant interactions .

The place that an individual occupies within the hierarchy of a plant population seems to be largely determined in the very early stages of plant development (Harper, 1977). Seed size is one of the first factors affecting this early development, and it was shown by

Austenson and Walton (1970) that a two or three fold difference in weight can exist among kernels of wheat in the same head.

The effect of seed size on plant development has been quantitatively studied by various authors. Kiesselbach (1924) reviewed a number of investigations made on cereals around the turn of the century, and added to this many of his own findings. He concluded that:

1. When small and large seeds of various cereal crops were space planted, small seeds developed into considerably smaller plants than did large seeds, giving correspondingly lower yields (on the average 19% less). This result was ascribed to the differences in energy reserves of the seeds, and hence, in initial seedling vigour.
2. When equal numbers of small and large seeds were planted at a rate optimal for the large seeds, the reduction in yield of the plants from small seeds was less than the reduction at wide spacings, although the small seed plants still yielded on the average 11% less than did the plants from large seed.
3. When equal weights of the two types of seed were planted, the plants from small seeds yielded on the average 3% less than did the plants from large seeds. The greater number of small seeds planted compensated almost completely for the smaller size. When equal weights of large seeds and unselected seeds were

planted, the plants from unselected seeds consistently yielded slightly more than the plants from large seeds. Averaged over 131 test years, the advantage of unsorted seed over large seed was 2.1%. The author did not speculate on the cause of this more efficient performance of a mixture of seed sizes.

Interaction between plants from different seed size within a genotype, and between genotypes differing in growth rate, was studied by Christian and Gray (1941). In both late and early cultivars of wheat, seed size affected the number of tillers and the number of heads per plant. Differences between the genotypes in their performance in mixture affected the the number of heads per plant but not the number of tillers per plant. Thousand kernel weight of the early genotype was significantly reduced by competition from the late genotype, but the latter was not affected.

A similar study was conducted by Kaufmann and McFadden (1960). They studied the competitive interactions between barley plants grown from large and small seeds, and concluded that plants from small seed are more negatively affected by increased plant density than are plants from large seed. The main character responsible for the yield reduction in response to increased density was the number of heads per plant. They reported a ratio of yield from large and small plants of 1:0.78 at a 10 cm equidistant spacing and a ratio of 1:0.54 at a 5 cm equidistant spacing. Comparison of yield from plots seeded to equal numbers of

either uniformly large seeds or uniformly small seeds revealed, as did the studies of Kiesselbach (1924) and Christian and Gray (1941), a higher yield for the large-seed plots.

The effects of seed size on components of yield were studied in more detail by Austenson and Walton (1970). They found that, within each of three cultivars of wheat, total yield, grain yield, straw yield, the number of heads per plant and number of seeds per plant were significantly correlated with initial seed weight. Variation in initial seed weight accounted for from 2.5 to 4.5% of the variation in these characters at maturity. Thousand kernel weight and number of seeds per head appeared to be unrelated to initial seed size.

Black (1958) demonstrated that, in two swards of subterranean clover, derived from equal numbers of seedlings but from different seed sizes, all plants in both swards eventually reached the same size. When large and small seeds were mixed, however, the plants from large seeds progressively suppressed the plants from small seeds and eventually made up 90% of the dry weight and leaf area of the sward.

Black noted that the relative growth rate of both the swards from large seeds and small seeds declined once a critical leaf area was reached. The plants from large seed reached this point faster than did the plants from small seed.

Kaufmann and McFadden (1960) reported that the vigorous growth of barley plants from large seeds resulted in an earlier development of second and subsequent leaves and of tillers than was the case for plants from small seed, but no difference was reported in the onset of flowering.

The importance of seed size and seedling growth rate to interplant relationships was demonstrated by Aspinall and Milthorpe (1959) and Aspinall (1960), who studied the interaction between barley and white persicaria. They concluded that, although in pure culture white persicaria has a faster relative growth rate than does barley, the latter has a larger embryo and initial seedling size, enabling it to establish a larger root system more rapidly and thus to gain an advantage when grown in a mixture. Restricted nutrient supply appeared to depress the growth of white persicaria more than did shading by the barley plants.

At the onset of flowering of the barley plants, when barley root growth and leaf expansion decrease, the persicaria plants immediately increased their growth rate, demonstrating the changes in interplant relations which can occur as plant development progresses.

Litav and Isti (1974a,b) compared the growth of seedlings of large and small seeded spinach (*Spinacia oleracea* L.) strains. Both strains gave the same individual plant yield in pure culture. The authors concluded from their experiments, which were grown at two levels of fertility, that a larger embryonic capital is only an

advantage when nutrients are in short supply. An experiment involving two seed sizes and two fertility levels, showed that an increase in soil fertility enables seedlings derived from smaller seeds to maintain an advantage when seeded earlier than the larger seeds.

A study of a more qualitative nature was made by { McDaniel (1969), who investigated the relationship of seed weight, seedling vigour and mitochondrial metabolism in barley. He found that seedling fresh weight, seedling mitochondrial protein and mitochondrial biochemical activity were positively correlated with seed weight. Although the number of mitochondria per unit fresh weight was the same for all classes of seed size, relatively more mitochondrial protein was present in seedlings from large seeds, resulting in a higher respiratory activity. The greater energy production in seedlings from large seeds thus allows a higher growth rate of these seedlings than of the seedlings from smaller seeds of the same genotype (McDaniel, 1969).

Seed size, thus, affects early growth rate of seedlings and is therefore an important factor in determining the competitive advantage or disadvantage of a plant at the time that a shortage of growth factors occurs.

2.4.5 Effects of density on plant interactions

Interaction between plants will start earlier in dense stands than in more widely spaced plantings. However, there is no direct effect of plant density upon plant growth. All

responses to density are indirectly measured as responses to limits on growth factors. Using wheat plants, grown at four different densities, Clements et al. (1929) observed that the effects of density resembled the effects of a water shortage. Pot cultures with different levels of soil moisture, fertiliser and light intensity, confirmed their theories and they concluded that, under their growing conditions, with increasing density, competition for water is most severe, with competition for nutrients and for light playing important, but secondary, roles.

Harper and Clatworthy (1963) studied changes in light absorption in swards of *Trifolium repens* L. and *T. fragiferum* L. grown at two densities. For both species, initial differences between the two densities in light extinction and leaf area had almost disappeared after fifteen weeks of growth. Apparently light was the limiting factor at both densities and an LAI of about five for *T. repens* and of six for *T. fragiferum* was the maximum that could be maintained, under the provided light regime.

Increasing population density, therefore, may be regarded as an increase in competition for light, nutrients and water among associated genotypes. Donald (1958) emphasised that, even if competition occurs for only a single factor, there will be interaction between direct and indirect effects. Competition for nutrients, for example, may affect shoot growth, and this may modify competition for light.

Harper (1961) distinguished two ways in which plants may react to density. Firstly, an increased density may result in a reduced chance for survival. He described the results of studies concerning several species of *Papaver*. It appeared that most of these species regulate their numbers in response to increases in density of their own kind, independent of the density of other species. This forms an important mechanism in nature, which prevents elimination through crowding by neighbouring species.

Secondly, there may be a plastic reaction to density during the development of a plant. This response was illustrated by Harper with the following example, describing the growth of *Agrostemma githago* L. under various conditions of density and in different mixture situations. When grown in pure stand, the number of plants per unit area, which were produced from different amounts of seed, was fairly constant. All plants bore an approximately equal number of capsules and gave equal seed yields. In association with different species, however, the surviving plants bore an equal number of capsules but gave a different seed yield. Unmixed, the *Agrostemma* plants in the study gave about 30,000 seeds/sq.yd. In association with wheat, the same the number of plants produced about 12,000 seeds/sq.yd., and in association with sugar beets, about 22,000 seeds/sq.yd.

This type of response, when it occurs in a mixture of genotypes such as encountered in the early generations of a breeding program, could have important effects on the

outcome of selection. The occurrence of such interactions is suggested by observations made by Phung and Rathjen (1976, 1977), who reported that grain yield of wheat plants was affected by the frequency of plants of the same genotype, amidst a population of plants of a different genotype.

Genotypes in segregating populations of corn responded to changes in density with changes in seed yield (CIMMYT, 1972). They noted that density affected the selection of corn, and that different genotypes were selected at different plant densities.

A study of the effects of variation in spacing, seed size and genotype on plant-to-plant variation of wheat, showed that increased interplant spacing was by far the most effective in increasing plant-to-plant variation for characters, such as plant dry weight and yield components (Chebib et al., 1973). However, interaction between genotypes and variation in seed size also consistently, but insignificantly, increased variation among plants.

Contrary to the findings of Chebib et al., Kelker and Briggs (1978) observed a decrease in variance with increased spacing for several plant characters measured on seven cultivars of wheat. They concluded that the tendency for variance to respond to changes in spacing is cultivar specific, and that not all plant characters respond to the same degree.

Hozumi et al. (1955) used uniform seed of yellow dent corn to observe the growth of individual seedlings and their

response to neighbouring plants. The seeds were grown in boxes outdoors. They found that large plants in a row tended to suppress their neighbours, and vice versa, resulting in an alternation of large and small plants. This was clearly shown by auto-correlations of plant weights within a row. Correlations with first, third and fifth neighbours were negative, while correlations with second and fourth neighbours were positive. Plant weight in their study was estimated from stem diameter and plant height. The importance of plant shape to plant interactions was clearly illustrated by the observation that correlation coefficients of actual plant weight showed a similar pattern, but were much smaller. At wider spacing, the neighbour effects appeared to be restricted to first and second neighbours only (Yoda et al., 1957).

In an earlier paper by Kira et al. (1953), the authors found quite different results for the growth of soybean plants. In this study, an hexagonal planting arrangement was used, with equal distances between all plants. The correlation between the weight of an individual and the mean weight of the six nearest neighbours appeared to be positive rather than negative. The authors did not speculate on the reason for the observed behaviour. It can be noted, though, that while the corn experiment was performed in boxes, the soybean experiment was grown in the field where soil heterogeneity might have played a larger role. This would tend to create a positive correlation between neighbours

which were similarly affected by varying soil conditions. It is also possible that the different planting arrangement was responsible for the different results, because in rows, the effects of one plant are exerted onto two neighbours, while six neighbours are involved in the interactions between plants grown in hexagons. One can expect a proportional dilution of neighbour effects in this case, decreasing the strength of neighbour correlations.

Thus, because changes in plant density tend to change the time at which neighbour interactions start, one can expect neighbour interactions which are density specific. In a mixture of genotypes or species, the density of plants from the same genotype, as well as the density of plants from different genotypes may affect plant growth independently.

2.4.6 Changes in the distribution of characters during growth

The shape of the frequency distributions of various characters, assessed at different times during the growing season, is illustrative of the dynamic nature of the population. Mean and variance give only a partial description of the distribution curve. A more precise picture is obtained when skewness and kurtosis are included.

Skewness is a measure of asymmetry of the curve. A positively skewed distribution has a mean to the left of the median, and a 'heavy' positive tail (L-shaped). A negatively

skewed distribution has approximately a J-shaped form (Sokal and Rolf, 1969).

Kurtosis further specifies the shape of the curve. Generally, when no skewness is present, a positive kurtosis means that the character has more values around the mean and in the tails and less in the intermediate regions, than does a character with a normal distribution does. The curve is thus sharply peaked with flat long tails. A negative kurtosis, in the absence of skewness, indicates more values in the intermediate regions (Sokal and Rolf, 1969). However, many different shapes of curves may have similar values of kurtosis, and this parameter is not necessarily a measure of peakedness of the curve (Kendall and Stuart, 1977). Strongly skewed distributions tend to have a positive kurtosis, due to the heavy tail on one side of the curve.

Koyama and Kira (1956) pointed out that the frequency distribution of plant weight of many species was normal shortly after emergence, but became more positively skewed (L-shaped) with time. This effect was stronger at higher plant densities. On the other hand, plant height most often developed a distribution which was negatively skewed (J-shaped). The authors observed that higher mortality occurred in those stands which developed a skewed weight distribution during early growth, while in populations which kept a normal distribution, mortality was low. The two species for which height distributions were reported, corn and ragweed, both have a determinate growth habit. In these

plants, genetic factors impose an upper limit to plant height, while no lower limit is present. This could explain the J-shape of the distribution of height. No mention was made of a possible different shaped distribution for plants of an indeterminate growth habit.

Non-normal distributions were also observed on characters of barley, grown in simulated segregating rows at Beaverlodge, Alberta (H. A. Kelker, unpublished). The measured characters (the number of tillers, height, kernels per head and others) were those which are normally used as criteria for plant selection in a breeding program, and non-normality of their distribution could affect the accuracy with which superior genotypes are selected.

2.4.7 Effects of genotype interaction on grain filling and maturity

Although plant interactions can evoke responses in a great number of plant characters, the plant breeder is, ultimately, only interested in the final effect of these interactions on grain yield, and in many regions, on maturity.

Final crop yield of a high yielding dwarf wheat cultivar, Yecora, appeared to be well buffered against small reductions in crop photosynthesis at various times during the growing season (Fischer, 1975). Crop growth rate, on the other hand, responded to shading at any time. There are various phenomena which can explain the stability of crop

yield. For example, Fischer (1975) also observed that this cultivar adapts to variations in light intensity or plant spacing with variations in tiller survival. In this case morphological plasticity of single plants is related to stability of crop yield.

Another mechanism aiding the stability of crop yield was described by Evans and Wardlaw (1976). They reported that, of the dry matter which accumulates in grain on unstressed plants, 90 to 95% comes from photosynthates produced after anthesis. The remaining quantity comes from reserves stored in vegetative plant parts. Plants under stress from shading or defoliation, however, will draw upon reserves to a much greater extent.

Although actual grain filling does not start until several days after anthesis, shading immediately after anthesis had a significant effect on kernel weight of wheat (Ford and Thorne, 1975). Shading during this period affected the subsequent capacity of grains to accumulate carbohydrates, possibly through limiting the number of endosperm cells, or through a mechanism which limits the amount of carbohydrates that can be translocated. Shading during this period, however, did not affect the amount of nitrogen in the mature grains, thus resulting in grains with a relatively higher grain protein percentage (Jenner, 1979).

The negative correlation which exists between the number of kernels per plant and kernel weight (Fisher, 1975; Jenner, 1979), once more illustrates the mechanism through

which yield can be buffered against variations in individual yield components. Variations in the number of kernels per head were brought about by various degrees of shading in the period just prior to anthesis (Fischer, 1975). Adjustments in subsequent kernel size, however, could not completely compensate for the reduced number of kernels, and reductions in yield were observed as a result from pre-anthesis shading.

Effects of plant interaction on maturity have generally been overlooked in studies on mixtures of genotypes, possibly because visual assessment of maturity of single plants is generally inaccurate. More reliable measurements can be obtained from the moisture content of the grain as an indirect indication of maturity, although this is a very time consuming technique (Somerville, 1977).

One could, however, anticipate that genotype interactions may affect maturity of mixture components. For example, variations in the onset of flowering, related to varying levels of nutrients in the leaves, have been reported for several plant species (Aitken, 1974). Such a situation could arise from the unequal sharing of available nutrients by the components of a mixture. Variation in the onset of flowering may affect the timing of subsequent developmental stages. Somerville (1977) reported a high correlation between date of flowering and maturity for both wheat and barley.

2.5 Effects of interaction between genotypes on single plant selection

The need for early identification of genotypes which have good agronomic qualities and high yield was clearly demonstrated by Shebeski (1967). He showed with a theoretical example, that with advancing generations, the proportion of plants which can be expected to possess the best combination of genetic characters decreases very rapidly. Thus, selection should ideally take place in the F₂ generation.

It is possible to select for a number of simply inherited traits, such as disease reaction or height, at an early stage of a breeding program (Sneep et al., 1979). However, characters which are influenced by many different internal as well as external factors, such as yield, cannot be correctly assessed on a single plant basis. McGinnis and Shebeski (1968) illustrated the ineffectiveness of visual selection for yield in an F₂ generation of wheat. They compared the results of visual selection of single plants by three plant breeders, with the results of random selection. They found that there was no significant difference between the mean yield of the F₃-progeny plots derived from randomly selected lines and derived from lines selected by plant breeders.

Because visual selection of single plants for yield *per se* is ineffective, several studies have dealt with the question whether selection for other characters would be

effective in increasing yield.

Rasmusson and Cannell (1970) found selection for the number of heads in two populations of barley to be as ineffective as selection for yield itself. Yield was reduced when selection was practiced for high number of kernels per head. Selection for high kernel weight was effective in increasing yield in one of the populations.

An increase in grain yield, associated with mass selection for large kernel size in wheat, was found by Derera and Bhatt (1972). This increase appeared not to be correlated in any way with quality characters, such as test weight, milling extract, wheat protein, flour protein and kernel hardness (Bhatt and Derera, 1973). Röbbelen (1979), however, explained that in the past, increased grain yields of many crops have led to a relative reduction in seed protein, which, in cereals, is mainly due to the increase of the endosperm relative to the embryo.

Morphologic and agronomic characters of 22 cultivars of wheat were studied by Nass (1973). He found that the yield per ear, the number of heads per plant, harvest index, and kernels per head were all positively associated with plot yield. Yield per head and heads per plot had the strongest associations, but because the yield per head and heads per plant are negatively correlated with each other, it would be difficult to select for both traits at the same time. Nass suggested that selection for moderate expression of these two characters could lead to increased yield.

Hamblin and Donald (1974) found no consistent correlations between yield per head and ears per plant in the F3 and the F5 generations of a barley cross. But they did find that tall F3 plants with long leaves tended to give low yields in the F5 generation. They suggested that the greater amount of interaction between genotypes which occurs in the F3 generation caused suppression of the higher yielding types, resulting in the observed negative correlation.

It was concluded from studies on barley that interactions between genotypes caused significant shifts in the expression of plant yield, heads per unit area, and the number of kernels per head, causing genotypes with a high yield potential to yield poorly (Wiebe et al., 1961). If yield in monoculture is the criterion for selection, the poorest rather than the best plants should thus be selected from the early generations.

In the past, yield improvements have been paralleled by a steady increase in harvest index, a character which appears to be relatively insensitive to variations in environmental conditions (Donald and Hamblin, 1976). However, this was an unplanned side effect, since no selection for harvest index was practiced. The question can be asked whether the effects would be reversible: would selection for high harvest index be accompanied by an increase in yield? From studies with F3 and F5 generation plants of barley, it was concluded that harvest index is not

predictive for yield from one environment (F3) to another (F5) (Hamblin and Donald, 1974). However, comparison of single plants of wheat, grown in pots in the greenhouse and in plots in the field, showed that single plant harvest index is a good predictor of crop yield in the field (Syme, 1972).

Selection for harvest index in the F2 generation of two wheat crosses doubled the grain yield per plant in the F3 generation (Bhatt, 1977). In addition, the positive correlation of yield with harvest index was enhanced. While selection for low harvest index resulted in a correspondingly low harvest index in the F3 generation, selection for high harvest index resulted in a population segregating for high and medium harvest index, thus allowing further selection for this character.

Although Donald and Hamblin (1976) reported that harvest index is relatively little affected by variations in density, resulting in a low genotype x density interaction, Nass (1980) found that indirect selection for grain yield through harvest index gave better results when practiced at commercial density than when practiced at low density. He ascribed this to the smaller plant-to-plant variation for harvest index which exists at high density.


The value of selection for harvest index was studied by Rosielle and Frey (1975a,b), using 1200 F9-derived lines of oats, grown in hill plots. They concluded that selection for harvest index alone was 43% as effective as was selection

for grain yield. If selection for height and maturity were combined with selection for harvest index, this percentage increased to 70%, indicating a good relationship between yield and harvest index within the limits of acceptable height and maturity, in homozygous lines.

Harvest index of main shoots appeared to be an even better indicator of crop yield. (Fischer and Kertesz, 1976). Using 30 genotypes of bread wheat, correlation coefficients were computed between the yield of plots at commercial crop spacing with harvest index of main shoots, harvest index of whole plants and grain yield of spaced plants, respectively. They obtained values 0.57 (sign., $\alpha = 0.01$), 0.49 (sign., $\alpha = 0.01$) and 0.20 (n.s.) respectively.

Because interaction between genotypes evokes reactions which are specific to each mixture situation, it is not surprising that Hamblin and Rosielle (1978) found great differences between genetic parameters of crosses when estimated in mixture and when estimated in monoculture. They demonstrated, using published data from different authors, that interaction between genotypes can increase or decrease the observed additive and dominance effects by a considerable amount. They showed cases where these parameters, measured in a monoculture and measured in a mixture, differed by more than 100%. In some cases, the values were about equal in magnitude, but opposite in sign.

In addition to an effect on the mean values of characters, positive and negative effects of plant



interaction on the variances were reported (Hamblin and Rosielle, 1978). As a result, estimates of heritability which are based on the estimation of variance components in a mixture of genotypes, would be either under- or over-estimated. 'This may lead to excessive effort being put into crosses and parents where competitive effects have increased heritability estimates and to the rejection of crosses and parents where they have reduced heritability estimates.' (Hamblin and Rosielle, 1978). The authors then suggested that, if early generation selection is to be effective, techniques should be developed which accurately assess the confounding effects of competition on the estimation of genetic parameters, and which could be used to correct these parameters for plant interaction effects.

In order to be able to assess the confounding effects of interaction between genotypes on the expression of genetic characters, it is necessary to know which plant characters are affected by genotype interaction, how interaction affects these characters and which characters of neighbouring genotypes are associated with these effects. The present study was conducted in an attempt to contribute fundamental knowledge on these subjects, focusing on the implications of genotype interaction for breeding of spring wheat.

3. Short Description of the Experiments

The experiments for this study were conducted during the growing seasons of 1977, 1978, and 1979, and were designed to investigate different aspects of the interaction between wheat genotypes.

During the first year of the study, eight genotypes and their binary mixtures were planted accurately at eight different interplant spacings. Morphologic characters were measured on single plants at maturity. It was hoped to learn from this test how the effects of competition were expressed at various spacings in each of the studied genotypes.

For the second year of the study, five of these genotypes were grown at three different spacings in monocultures and binary mixtures, and a detailed study was made of the relationships between plant characters of competing genotypes. Effects of interaction on mean plant performance as well as on plant to plant variation were determined.

During the last year of the study, the role of seed size and subsequent plant development on the development of adjacent plants in a row was investigated. This experiment involved only monocultures of the five genotypes used in the second year.

In addition to the single plant experiments, replicated yield trials, containing eight genotypes in monoculture and all possible binary mixtures, were grown in each of the three years. These tests were conducted to illustrate how

each of the genotypes affected mixture yields, and how the performance of mixtures varied over the three years.

4. Materials and Methods - General Information

The specific procedures followed for each of the experiments in this study are discussed in the sections preceding the presentation of the results for each test. General information pertinent to all the experiments is given in the following sections.

4.1 Characteristics of test sites

Experiments were conducted at the Edmonton Research Station in 1977 and in 1979, and at the Ellerslie Research Station in 1978. Both stations are operated by the University of Alberta, Edmonton, Alberta, Canada.

The sites are located about 10 km apart, at a latitude of 53° 24' N and at an altitude of 694 m above sea level. Appendix 1 shows the mean monthly temperatures and precipitation during the three growing seasons in which the experiments were conducted.

Both sites are located in the thin black soil zone and the soil is classified as silty clay loam.

Fertilizer was applied in each spring prior to seeding, at rates recommended for wheat, based on soil analyses conducted by the Soil and Feed Testing Laboratory of Alberta Agriculture (Appendix 2). During the fall of 1977 and 1978, Avadex was applied to the test sites for control of wild oats. In 1977 and 1978, herbicides were applied during the early stages of growth, for control of the common broadleaf

weeds, stinkweed, green smartweed, pigweed and hempnettle (Appendix 2). In addition, all plots were hand weeded when necessary.

4.2 Plant material used

Eight wheat genotypes were used in the test: Pitic 62, Glenlea, Park, Neepawa, 70M110001, 70M009002, Norquay and NB701. Park and Neepawa are Canada Western Red Spring wheats, while the other genotypes are Utility type wheats. The genotypes were chosen to represent a wide range of agronomic characters such as height, maturity, tiller number and seed size (Appendix 3). Seven of these genotypes had been previously studied, providing extra information on their performance at the Edmonton and Ellerslie test sites (Attinaw, 1977; Somerville, 1977).

As a seed source for Pitic 62, Glenlea, Park, Neepawa and Norquay, foundation seed or registered seed was used. For the genotypes 70M110001, 70M009002 and NB701, clean, sound seed was obtained from yield trials conducted as part of The University of Alberta wheat breeding program.

In each year, kernel weights and germination rate of all genotypes were determined on the seed lots to be planted (Appendix 4). Germination tests were conducted in petri dishes containing two rounds of filterpaper (Wattman, no. 40) and 4 ml of distilled water. The dishes, containing 100 seeds each, were placed in a Seedburo incubator, model 2100,

at 36°C, and germination was determined at 24, 48, 72 and 96 hrs.

5. Interaction Between Genotypes at Different Interplant Spacings

5.1 Materials and methods

5.1.1 Field plot design

During the first year of the study, monocultures and binary mixtures of the eight genotypes were tested over a wide range of spacings, to observe genotypic differences in response to variations in the environment.

Each of these mixtures and monocultures was grown at eight different spacings, ranging from 7 to 60 cm between plants. To make the most efficient use of space, a 'wheelplot' design was used, in which each wheelplot represented one mixture or monoculture grown at seven spacings. Seeds were placed on a grid made up of concentric circles and 'spokes', and seeds of the competing genotypes were alternated on the spokes and circles, such that each plant was surrounded by four plants of the associated genotype. In proceeding outward from the centre of the circle, the distance between plants increased as the spokes diverged, and the distance between concentric circles increased.

Plant density was calculated as the average of the distance from one plant to its four immediate neighbours, which were situated on either the same spoke or on the same

circle. The resulting spacings were 7, 9, 12, 15, 19, 24 and 31 cm between plants or 204, 124, 69, 44, 28, 17 and 10 plants per m² respectively. Commercial seeding rates in the Edmonton area range from 56 to 101 kg/ha (Alberta Farm Guide, 1976), which converts to approximately 140-250 seeds/m².

Each plot had the shape of a three quarter circle with a radius of 2 m, containing twenty four spokes and nine concentric circles. The plants on the innermost and outermost circles, as well as on the first and the twenty fourth spoke, were discarded as guard plants.

In addition to the seven spacings in the wheelplots, each genotype was grown in a rectangular plot, containing 10 plants each, planted at an equidistant spacing of 60 cm. At this spacing plant interactions were assumed to be absent.

5.1.2 Seed preparation and data collection

To reduce seeding time, all seeds were glued onto cotton string (10/6 S), made by Dominion Textile (Texmade), using LePage's Multi Use Bondfast. This method had previously been used in the field for several years with good success (Dr. V. Burrows, Chief, Cereal Section, Research Branch, Ottawa, Agriculture Canada, 1976; personal communication). To keep the 24 strings (spokes) of each plot separated, they were labelled and tied, in the order in which they had to appear in the field, onto a stake which at seeding time could be placed in the centre of the plot. A

strip of paper was subsequently placed on top of the strings and strings and paper together were rolled around a cardboard tube.

To facilitate placement of the strings in the field, a square metal frame was constructed, containing a bar which could rotate around a central axis. Furrows were drawn with a metal shoe which slid along this bar. After placement of the strings, the furrows were closed and packed using a board.

Before harvest, the location of each plant was recorded on plot maps and all plants were labelled. All plants which had all four neighbours present were harvested when 75% of the tillers appeared to be ripe. Measurements were taken on plant height, height of the flag leaf blade, the number of tillers, the number of fertile heads (heads with a minimum of five seeds), the number of kernels per plant, grain yield per plant, and above ground plant dry weight. From these measurements, the following characters were calculated: extrusion length (calculated as plant height - height of the flagleaf blade), the number of kernels per head, weight per 1000 kernels, the yield per head and harvest index (calculated as plant grain yield/ plant dry weight).

Because the present study approached the problem of genotype interaction as it is encountered by the plant breeder when he makes selections in the field, measurements of height and related characters were taken on two randomly chosen tillers of each plant. In this way it was hoped to

obtain a more realistic estimate of the variation which is observed in the field, than would be obtained from measurements on the main stem only.

5.2 Analysis of the data

The wheelplots were arranged according to a randomised block design with two replications which each occupied two banks of plots. The 60 x 60 cm spaced plants were replicated four times and grown at the end of each bank.

Because the spacing treatments were arranged systematically within the wheelplots, the requirements of randomisation were not met. The spacing effects indicated by the analysis of variance are thus confounded with effects due to treatment arrangement. The results cannot be considered as more than an indication of possible effects of spacing and will have to be further tested in subsequent experiments.

5.3 Results

Pitic 62 failed to germinate sufficiently and all treatments involving this genotype were dropped from the 1977 tests. Additional germination and or growth problems were encountered in the remaining plots, resulting in a very low number of usable plants overall. More than half of the plots (31 out of 56) had a survival rate of less than 75%. Recorded survival rate in some plots was as low as 43%.

while a few plots (including all 60 x 60 cm spaced plants) were abandoned completely.

In search for an explanation for the poor germination, soil tests were performed in the fall of 1977. The results indicated levels of available P ranging from 39 to 47 lb/2M on the east side of the field to 14 lb/2M along the west side of the field. Although no specific symptoms of P-deficiency were noted during the growing season, all but four of the sparsely populated plots were located in the western halves of the banks, suggesting a direct or indirect relationship between the phosphorus content of the soil and survival rate.

Seeding was interrupted by two days of heavy rain, causing some of the seeds to become uncovered, which probably added to the low germination rates in some plots.

In an effort to get as much information from the data as possible, the test was treated as a completely randomised design. Plants from both replicates were pooled, and only those treatments were included for which data had been recorded on a minimum of five plants. For none of the genotypes were data available from all possible genotype combinations (Appendix 5). Means and variances of the characters measured at each treatment illustrate the observed range of performance (Appendix 6). Since data about monoculture performance were obtained for only four genotypes (Glenlea, Park, 70M110001 and Norquay), tests for the effects due to genotype interaction had to be limited to

these four genotypes.

Spacing effects and genotype effects were significant ($\alpha \leq 0.01$) for all characters (Table 1). Mixing significantly affected height of the flag leaf, plant height, 1000 kernel weight and harvest index. However, interactions between effects of genotype interaction and spacing were significant for plant weight, yield and the number of kernels per plant, indicating that neighbouring plants affected these characters at some of the spacings tested. Interactions between the effects of genotype interaction and genotype were significant for all characters except the number of heads, the number of tillers, and kernels per plant, suggesting that the genotypes differed in their response to genotype interaction. Three way interactions were significant for the number of heads, the number of tillers, weight, yield, and the number of kernels per plant.

Comparison of mixture and monoculture performance of each of the four genotypes at each of the spacings tested, indicated the specific densities at which the various characters were affected (Table 2). While interaction - indicated by significant t-values - was evident for Glenlea at a density of 10 plants/m², Park was only affected at 124 and 69 plants/m² and 70M110001 only at 124 plants/m². None of the genotypes showed significant effects of interaction at 44 and at 204 plants/m². It is likely that at the highest plant density, the general effects of density on plant growth were so strong that the more subtle effects of

Table 1. The effect of genotype, genotype-mixing, and interplant spacing on single plant characters, as determined by analyses of variance.¹

All plants were grown in wheelplots in 1977, and measured at maturity. The data are only from plants which had all four neighbours present.

Source of variation	df	Mean squares of characters ²											
		T	H	Wt	Y	FL	Ht	ExL	K/P	K/H	Y/H	Kwt	HI
Genotypes (G)	3	160.3**	96.2**	6645**	1162**	3510**	32504**	23481**	280477**	2665**	16.35**	2299**	0.191**
Mixing (M)	1	0.0	13.8	1135	109	505**	8859**	17	1264	360	1.16	444**	0.037*
Spacings (S)	6	1915.1**	907.2**	31510**	7103**	83**	346**	278**	3988200**	11672**	15.93**	153**	0.082**
G × M	3	7.1	19.0	1008*	237*	986**	3614**	201**	77860	1668**	4.46**	916**	0.118**
G × S	18	49.5**	46.7**	1035**	243**	22	50	37	74868*	341	0.67	18	0.015*
M × S	6	42.7	28.0	1062**	252**	22	41	19	98303*	610	0.56	59	0.009
G × M × S	9	40.3	41.0**	1183**	198*	34	83	40	83213*	265	0.96*	40	0.013
Residual	574	22.5	16.3	361	94	29	80	55	42857	325	0.46	42	0.009

1. * significant, $\alpha \leq 0.05$; ** significant, $\alpha \leq 0.01$.

2. Abbreviation of characters:

T = Number of tillers per plant

H = Number of fertile heads per plant

Wt = Plant dry weight

Y = Grain yield per plant

FL = Height of the flag leaf blade

Ht = Shoot length

ExL = Extrusion length

K/P = Number of kernels per plant

K/H = Number of kernels per head

Y/H = Grain yield per head

Kwt = 1000 kernel weight

HI = Harvest Index

Table 2. Comparison of the means of characters of single plants, grown in monoculture and in mixture, at various plant densities, by means of Students t-test.¹

All plants were grown in wheelplots in 1977, and measured at maturity. The data are only from plants which had all four neighbours present.

Genotype	Density ² (plants/m ²)	df	t-values for measured characters ²									
			lit	H	T	Wt	Y	K/P	K/H	Y/H	Kwt	HI
Glenlea	204	22	0.54	0.21	0.31	0.71	0.00	0.19	0.48	0.91	0.24	2.24
"	124	33	1.28	0.06	1.27	0.58	1.07	1.14	2.37*	2.23*	0.52	1.76
"	69	33	1.01	0.63	1.66	0.90	1.88	1.63	2.46*	2.79**	1.89	1.85
"	44	27	1.44	0.71	0.94	0.29	0.37	0.24	1.16	0.98	0.09	0.92
"	28	18	0.51	2.23*	3.46**	2.47*	3.03**	2.86**	0.31	0.45	0.43	2.66*
"	17	18	0.71	1.04	0.06	0.13	0.44	0.55	2.81**	2.15*	1.09	0.80
"	10	17	0.57	0.29	1.37	0.73	2.39*	2.72**	2.23*	2.00	2.38*	2.15*
Park	204	23	1.39	0.13	0.17	0.40	0.52	0.33	0.71	0.97	0.89	0.66
"	124	39	1.59	2.01*	2.45*	1.62	1.77	1.86	1.16	1.22	0.67	1.35
"	69	30	3.54**	1.20	0.82	1.31	1.18	1.79	0.33	0.60	0.41	1.41
"	44	26	1.86	0.06	0.27	0.16	0.66	0.06	0.18	1.05	1.83	1.46
"	28	27	0.29	0.91	0.31	0.35	1.16	0.83	0.11	0.74	0.66	1.79
"	17	24	0.34	0.26	0.45	0.47	1.26	0.57	0.05	0.45	0.71	1.59
"	10	24	0.50	1.78	0.56	0.90	0.89	0.74	1.15	0.94	1.38	1.68
79M110001	204	23	0.20	0.28	0.22	0.25	0.73	0.54	1.72	1.98	0.40	1.82
"	124	20	0.93	1.37	2.09*	2.26*	2.31*	2.28*	1.43	1.78	1.01	0.88
"	69	21	1.86	0.60	1.66	1.51	0.20	0.59	0.28	0.85	1.63	1.15
"	44	25	1.19	0.28	0.40	1.11	1.64	1.54	0.95	0.27	0.25	0.44
"	10	17	0.78	0.35	0.77	0.90	0.92	0.91	1.55	1.59	0.00	0.45
Merquay	124	10	0.69	0.28	1.06	0.13	0.90	0.98	2.30*	2.29*	0.72	0.42
"	10	15	0.41	0.28	0.13	0.14	0.92	0.68	1.21	1.46	1.73	1.35

1. * significant, $\alpha \leq 0.05$, ** significant, $\alpha \leq 0.01$

2. character abbreviations defined in Table 1

interaction between different genotypes could not be distinguished with the experimental techniques used.

5.4 Discussion

Because the data used for the analysis of this test come from a small number of observations only, further experiments are necessary to verify the conclusions about the effects of plant interactions on the genotypes involved over the range of spacings tested. Nevertheless, a number of comments can be made regarding the observed effects.

The fact that the genotypes differed in their response to genotype interaction, and that each did so at different spacings, indicates that interpretation of mixing experiments performed with a limited number of genotypes and environments could lead to variable conclusions.

It appeared that Glenlea, which is the tallest, latest and most vigorous of the four genotypes, was the only genotype to show effects of genotype interaction at a density as low as 10 plants/m². Possible methods of interaction are restricted to light interception by neighbouring plants, and interaction between root systems. Since shading by neighbours only occurred during part of each day, interaction between root systems is probably the most influential factor at this interplant spacing.

6. Relationships between Plant Characters Measured on Mixed Genotypes

6.1 Materials and methods

In 1978, a test was conducted to investigate, in more detail than in 1977, the phenotypic responses of genotypes to genotype interaction at different spacings. To keep the experiment manageable, the number of genotypes had to be restricted to five: Pitic 62, Glenlea, Park, 70M009002 and Norquay. They were chosen, based on the 1977 yield trials, to represent a wide range of Average Mixture Efficiency (see Section 8.2, p.92), and for their recognisability in a mixture. They were grown in rows 30 cm apart, at plant spacings within rows of 1, 3 and 9 cm. This resulted in densities of 333, 111 and 37 plants/m². Seeds were glued onto strings as described in Section 5.1.2, and placed in furrows which were drawn with a pointed stick along a board.

A split plot design was used with three replicates. The main plot treatments were mixtures and monocultures, and the subplot treatments were spacings. The mainplots, which were 6 m long, were divided into three subplots of equal length. Each plot consisted of three rows ~~of which~~ only the middle one was sampled. Three plants at each end of each subplot were discarded as guards.

From both the 3 cm spacing and the 9 cm spacing, a random sample of six plants per genotype was taken at

maturity, using only plants which had both neighbour plants present. Measurements were taken on the height of the the flag leaf blade, the extrusion length (height above the flag leaf blade), the head length, the yield per plant, the number of kernels per plant, the number of tillers per plant, the number of fertile heads per plant, and plant dry weight. From these measurements were computed the number of kernels per head, the yield per head, the 1000 kernel weight and the harvest index. As in 1977, measurements on height and related characters were taken on a random sample of two tillers per plant.

It was the intention to examine the root distributions of the genotypes in mixture, and therefore the complete bulk yield test was replicated several times, so that enough extra plots were available for destructive soil core sampling. Unseasonably high soil moisture content, however, made it impossible to obtain good soil cores, and this objective was abandoned. Thus, extra plots were available, and tiller samples were taken from these at flowering and at the mid-dough stage. Measurements were taken on the flag leaf area, the height of the flag leaf blade, the tiller length, the extrusion length and the head length. Each sample consisted of a 1 m section of row, chosen at random from one of the middle rows of the plots. The samples taken at the two developmental stages were not replicated, so that differences between mixing treatments may be confounded with differences among plots. At each stage, sampling was

completed in one day, so that differences among the genotypes will reflect genotypic differences in rate of development.

6.2 Analysis of the data

Because it was not possible to accurately separate the plants which were grown at the 1 cm spacing, only data obtained from the 3 cm and the 9 cm spaced plants were included. The data were analysed to detect effects of genotype interaction on within genotype variation, as well as on mean plant performance.

Variance, skewness and kurtosis of the measured characters were computed for each sample, as well as for the combined mixture data for each genotype. Because significant deviations from normality occurred, Bartlett's test for homogeneity of variances could not be used, since this test is extremely sensitive to non-normality (Scheffé, 1959). Scheffé described an approximate test for the comparison of variances, based on the natural logarithms of the sample variances. This analysis was used to test the effect of genotype, genotype interaction and spacing on the variances of all characters.

Overall differences in variation between plants grown in monoculture and plants grown in mixture, were determined by means of a variance ratio F-test, using the mean variances of all genotypes grown in mixture and grown in

monoculture. For these comparisons the natural logarithms of the data were used (Lewontin, 1966).

Although the test was designed as a split plot, it was decided that an error term with more degrees of freedom was desirable, which would make it possible to detect smaller effects of genotype interaction than would be possible with a split plot analysis of variance. Therefore a randomised block analysis of variance was used to test the effects of genotype, spacing and genotype interaction.

Mixture efficiency was expressed for components of yield and plant weight, using the technique described by de Wit (1960), as the 'Relative Yield Total' (RYT):

where Y_{AB} and Y_{BA} are, respectively, the mean plant yields, or values of the character concerned, of the genotypes A and B when grown in association with each other, and Y_{AA} and Y_{BB} are the monoculture yields of the two genotypes. The expressions Y_{AB}/Y_{AA} and Y_{BA}/Y_{BB} are called the 'Relative Yield' (RY) of genotypes A and B, respectively, expressing the ratio of the yield of a genotype in mixture to its yield in monoculture. A value of 1 for the RY of a mixture component indicates that the effects of interaction between genotypes are not different from those which plants of the same genotype have on each other. A RY greater than 1 suggests that the associated genotype makes demands on the

environment which are either different from, or less than, those of the genotype under investigation. A RY of less than 1 indicates that interaction between the genotypes negatively affects the growth of the genotype in question, compared to its growth in monoculture.

A RYT of 1 would indicate effects of a compensating nature, or the absence of any effect on either of the components. A RYT greater than 1 suggests that the genotypes are, at least partially, occupying different niches, and thus are not sharing the same supply of resources. A value less than 1 indicates an antagonistic effect of one or both genotypes on their associate.

Both RY and RYT values were computed for mixtures and their components.

Since no simple test is available to determine whether RY or RYT values differ significantly from 1, they were evaluated with a t-test, comparing monoculture and mixture values. RY values were called significant if the difference between mixture and monoculture yield differed significantly from 0. Significance of the RYT values was determined using a t-test comparing expected yields as can be computed from monoculture values, with observed performance in mixture.

Within each individual plant, many characters are correlated. When investigating the relationships between competing plants, these within plant correlations can play a confounding role. Unfortunately, the number of genotype combinations in the experiment yielded an insufficient

number of degrees of freedom to investigate partial correlation coefficients between all characters of associated genotypes. Only correlations between character pairs have therefore been investigated. For this, multiple regression analyses were used, using indicator variables to correct for replicate effects (Steel and Torrie, 1980).

6.3 Results

During the growing season, two large bare spots, in which no plants grew, occurred in the plot area. Soil tests, performed after harvest to determine levels of available soil nutrients, soil pH and physical soil characteristics, did not reveal any abnormalities which could be related to the bare spots. All monoculture plots of Pitic 62 happened to be planted in these areas, so that no comparisons could be made between the performance of this genotype in monoculture and in mixture. Additional mishaps at harvest time resulted in the loss of several more treatments. For each genotype, mean values of the measured characters, when grown in monoculture and in mixture at 3 cm and at 9 cm spacings, are given in Appendix 7.

Although one can expect genotype interactions to affect flowering time and subsequent maturity of the mixture components (Aitken, 1974; Somerville, 1977), measurements of maturity on single plants were not made, because the methods available for rapid screening of a large number of samples

are generally inaccurate (Somerville, 1977). Many more replications would have been needed than were available. Visual assessment of genotype maturity in mixture, however, did not indicate any effects of genotype interaction, and no data on this character have been reported.

6.3.1 Effects of genotype interaction on plant-to-plant variation

An analysis of variance using the natural logarithm of the sample variances, showed no effect of genotype interaction on plant-to-plant variation in any of the characters (Table 3). Significant differences between genotypes occurred for variances of all characters except the number of tillers, the number of heads, kernel weight and harvest index, and changes in spacing affected the number of tillers, the number of heads, plant weight, yield, kernels per plant and extrusion length. When, for each genotype, the results of all mixtures were combined, however, variation was, on the average, greater in mixture than in monoculture for the number of tillers, plant weight and yield at the 3 cm spacing, and for kernel weight at the 9 cm spacing (Table 4). Harvest index, height of the flag leaf, extrusion length and height were less variable in mixture than in monoculture at the 3 cm spacing, while kernel weight, height, and extrusion length showed a lesser variability in mixture than in monoculture at the 9 cm spacing.

Table 3. Effects of genotype, genotype mixing and interplant spacing, on the variance¹ of single plant characters, as determined by analysis of variance².

All plants were grown in four-row plots in 1978. The data are from randomly selected plants, which had both neighbours present, and which were measured at maturity.

Source of variation	df	Mean squares of characters ³												
		T	H	Wt	Y	FL	Kwt	Ext	HI	FL	Ht	HL	ExL	Y/H
Genotypes (G)	3	0.66	1.13	4.13**	4.58**	2.29**	1.49	4.02**	0.16	2.33**	5.47**	2.78**	3.14**	5.40**
Mixing (M)	1	0.83	0.13	0.02	0.22	0.20	0.91	0.50	2.13	0.47	0.40	0.70	0.09	2.67
Spacings (S)	1	51.37**	54.88**	67.18**	66.95**	68.17**	0.85	1.19	0.03		0.17	0.44	2.74*	0.04
G × M	3	0.24	0.17	0.28	0.12	0.09	4.45*	0.22	0		1.60*	0.93	0.47	0.17
G × S	3	0.97	0.38	0.62	0.57	0.75	0.92	0.17	0.00	1.11	2.04	0.20	0.30	0.47
M × S	1	0.95	0.08	0.01	0.04	0.01	0.01	0.13	0.49	0.18	0.02	3.16	0.72	0.01
G × M × S	3	0.59	0.33	0.34	0.36	0.20	0.48	0.15	1.92	0.36	0.18	0.10	0.40	0.22
Residual	66	0.57	0.48	0.44	0.43	0.46	1.46	0.95	1.21	0.59	0.83	0.70	0.67	0.87

1. The natural logarithm of the variances were used.

2. * significant, $\alpha \leq 0.05$; ** significant, $\alpha \leq 0.01$.

3. Character abbreviations defined in Table 1. HL = Head length.

Table 4. The effects of genotype mixing and interplant spacing on the average variation among plants, of single plant characters, as determined by a variance ratio F-test¹.

All plants were grown in four-row plots in 1978. The data are from randomly selected plants, which had both neighbours present, and which were harvested at maturity.

Character ²	Inter-plant spacing (cm)	Coefficient of variation ³		F-values	
		Monoculture	Mixture	F ₁ ⁴	F ₂ ⁵
T	3	25.1	36.0	2.06**	2.01**
H	3	28.2	31.7	1.26	1.82*
Wt	3	29.5	37.6	1.63*	1.68*
Y	3	33.1	39.0	1.39	1.48
K/P	3	31.2	36.6	1.38	1.68*
Kwt	3	7.6	8.9	1.37	2.04** ⁶
K/H	3	18.5	19.5	1.11	1.01
HI	3	12.1	8.4	2.08**	1.89**
Fl	3	14.7	13.3	1.22	1.28
Ht	3	12.2	9.5	1.64*	1.75**
Exl	3	17.5	14.8	2.08**	2.32**
HL	3	9.4	13.9	1.38	1.22

Character ²	Interplant spacing (cm)	Coefficient of variation ³		F-values	
		Monoculture	Mixture	F_3^7	
T	9	35.6	38.7	1.18	1.16
H	9	38.0	38.9	1.05	1.51**
Wt	9	38.2	40.3	1.11	1.15
Y	9	40.3	42.3	1.10	1.18
K/P	9	40.4	41.6	1.06	1.29*
Kwt	9	5.3	10.0	3.56**	1.26
K/H	9	18.6	18.0	<i>1.06</i>	<i>1.18</i>
HI	9	8.8	9.0	1.05	1.15
F1	9	13.0	13.4	1.06	1.01
Ht	9	9.2	9.0	<i>1.03</i>	<i>1.10</i>
Exl	9	11.5	11.4	<i>1.02</i>	<i>1.69**</i>
HL	9	9.5	12.7	1.08	<i>1.04</i>

1. * significant, $\alpha \leq 0.05$; ** significant, $\alpha \leq 0.01$
2. Character abbreviations defined in Table 1. HL = Head length.
3. Averages of the coefficients of variation of four genotypes.
4. $F_1 = s^2(\ln x, \text{mixture}) / s^2(\ln x, \text{monoculture})$, df(180,60)
5. $F_2 = s^2(\ln x, 9 \text{ cm, monoculture}) / s^2(\ln x, 3 \text{ cm, monoculture})$, df(60,60).
6. F-values given in italics are the reciprocal of the values defined in footnotes 4, 5 and 7.
7. $F_3 = s^2(\ln x, 9 \text{ cm, mixture}) / s^2(\ln x, 3 \text{ cm, mixture})$, df(175,180).

Values of skewness and kurtosis averaged over the four genotypes grown in monoculture and in mixture, are given in Appendix 8. For some characters, in particular kernel weight and harvest index at the 3 cm spacing and number of heads and kernel weight at the 9 cm spacing, skewness was stronger in mixture than in monoculture. Other characters, such as kernels per head and height of the flag leaf at the 3 cm spacing and height at the 9 cm spacing, showed a skewed distribution in monoculture and a near normal distribution in mixture. In most cases, the kurtosis was less in mixture than in monoculture, indicating that, in mixture, more values occurred in the intermediate regions, with a lower concentration of values around the mean and in the tails. An analysis of variance, performed on the values of skewness and kurtosis, did not detect any significant effects of genotype interaction. However, the values were based on samples of only six plants per replicate. An analysis based on larger numbers of plants would be necessary to draw more meaningful conclusions with respect to the effect of genotype interaction on skewness or kurtosis.

6.3.2 Effects of genotype interaction on the expression of various plant characters

Although it was demonstrated in the previous section that the distributions of most characters deviated significantly from normal, no suitable transformations were found to correct this. Analyses were thus performed on

untransformed data.

As expected, the genotypes differed significantly from each other for most characters (Table 5). Mixing of plants with individuals of a different genotype had a significant effect only on head length, on kernels per head and on yield per head. However, genotype x mixing interaction effects were significant for all characters except height of the flag leaf, plant height, kernels per head and harvest index, suggesting differences between the genotypes in their reaction to genotype interaction.

All characters except height and harvest index were affected by an increase in spacing, and replicate effects were significant for extrusion length, 1000 kernel weight and harvest index. Genotypes differed for most characters in their response to spacing, as indicated by the significant genotype x spacing interactions. Mixing x spacing interactions were only significant for weight and harvest index.

6.3.3 Relative performance of genotypes in mixture

To evaluate the effect of genotype interaction on each character in each genotype, RY values were calculated (Table 6), and the effects of genotype interaction were tested for significance with a t-test. It can be seen that Glenlea, in most cases, yielded better in mixture than in monoculture. This effect was associated with an increased number of heads, and to a lesser extent with an increased

Table 5. The effect of genotype, genotype mixing, and interplant spacing on single plant characters, as determined by analyses of variance¹.

All plants were grown in four-row plots in 1978. The data are from randomly selected plants, which had both neighbours present and which were measured at maturity.

Source of variation	df	Mean squares of characters ²												
		T	H	Wt	Y	PL	HC	ExL	HL	K/P($\times 10^{-3}$)	K/H	Y/H($\times 10^2$)	KWt	WI($\times 10^3$)
Genotypes (G)	3	5	24**	1794**	399**	9857**	16497**	1318**	320**	76**	4098**	2496**	5641**	58**
Mixing (M)	1	5	1	164	18	74	131	59	23**	15	528**	121**	36	2
Spacings (S)	1	1340**	953**	22031**	4338**	934**	19	1057**	40**	220**	3338**	932**	314**	1
Replications	2	2	1	2	5	29	41	119*	2	5	9	1	51*	8**
G \times M	3	20	16**	444**	91**	11	77	202**	6**	36**	102	39*	67**	3
G \times S	3	11	6	600**	106**	167*	42	89*	5*	23*	25	6	51**	6**
M \times S	1	1	3	3	7	39	1	58	0	2	2	0	0	9*
Residual	467	5	4	73	15	48	30	33	2	8	46	12	14	2

1. * significant, $\alpha \leq 0.05$; ** significant, $\alpha \leq 0.01$.

2. Character abbreviations defined in Table 1. HL = Head length.

Table 6 (continued)

Mixture	HI		Y/H		FL		HT		ExL		HL	
	3 cm	9 cm	3 cm	9 cm	3 cm	9 cm	3 cm	9 cm	3 cm	9 cm	3 cm	9 cm
Glenlea + Pitic 62	1.00	1.08*	1.08	1.01	0.96	1.03	1.01	1.04	1.13	1.03	1.08	0.98
+ Park	0.94	1.07	1.07	1.00	0.94	0.07	1.01	0.97	1.15	0.97	1.07	1.00
+ 70M009002	0.99	1.02	1.10	0.98	0.95	1.03	1.02	1.03	1.17*	1.02	1.09	0.99
+ Norquay	0.97	1.11**	1.14	0.86	0.96	0.89	1.04	0.93	1.20*	0.98	1.05	0.96
Park + Pitic 62	1.05	1.09	1.17	0.92	0.98	1.00	1.04	0.96	1.11	0.90	0.97	0.87
+ Glenlea	1.00	0.98	0.91	0.70**	0.97	0.96	1.02	0.93	1.08	0.89	1.01	0.84*
+ 70M009002	0.98	1.03	0.95	0.88	0.92	0.95	0.98	0.94	1.08	0.92	1.01	0.92
+ Norquay	1.01	0.99	1.24	0.96	0.96	1.04	1.02	0.99	1.11	0.96	1.12	0.94
70M009002 + Glenlea	1.01	1.06	0.78**	0.83*	0.99	0.98	0.88	0.98	0.84*	1.01	0.88	1.08
+ Park	1.00	1.03	0.89	0.95	1.02	0.97	0.95	0.99	0.90	1.05	0.99	1.11
Norquay + Pitic 62	--	0.99	--	0.93	--	1.00	--	1.00	--	0.99	--	0.98
+ Glenlea	0.96	0.99	0.60**	0.94	0.90	1.01	0.88	0.98	0.87	1.04	0.88	1.00
+ Park	0.96	0.99	0.86	0.94	1.04	0.97	1.00	0.98	0.95	0.99	0.92	0.98

1. RY = character in mixture
character in monoculture, see Section 6.2.

2. * significant, $\alpha < 0.05$, ** significant, $\alpha < 0.01$.

3. Character abbreviations defined in Table 1. HL = Head length.

4. No data available.

kernel weight. At the 9 cm spacing, harvest index of Glenlea in two mixtures was significantly higher than in monoculture.

With one exception, Park yielded less in mixture than in monoculture. Reduction in yield was associated with a lesser number of heads per plant, and in one mixture with fewer kernels per head.

Data for 70M009002 were only obtained for two mixture combinations. Yield in combination with Glenlea was less than in monoculture at both spacings and was associated with a reduced number of kernels per head. The number of heads per plant did not show any change from the number in monoculture. Park did not significantly reduce the yield of 70M009002.

Data for Norquay in association with Pitic 62 were only available for the 9 cm spacing. Although yield was less than in monoculture, none of the RY values were significantly less than 1. At the 3 cm spacing, the performance of Norquay in the presence of Glenlea was poor, and the number of kernels per head and 1000 kernel weight were both significantly ($\alpha \leq 0.01$) reduced. RY for the number of heads per plant was, in spite of its low value (0.72), not significantly less than 1, due to the small number of observations for this treatment. Association with Park reduced the yield of Norquay at the 3 cm spacing and increased yield at the 9 cm spacing, but neither value (0.87 and 1.12) was significant.

Overall, Glenlea tended to produce RY values greater than 1 for plant weight, yield and components of yield, while values less than 1 were more frequent among the other genotypes. This suggests that Glenlea plants grown in mixture were able to utilise a greater amount of resources than when grown in monoculture. The other genotypes, in particular Park, were more often negatively affected by associated genotypes.

Relative yield totals, derived from the RY values, appeared to be close to 1 for most cases (Appendix 9). A graphic representation of the performance of plants grown in monoculture and in mixture further illustrates the effect of mixture components on each other. Figures 2a through 2e illustrate the relationship between the yields of the genotypes grown in binary mixtures, and their yields in monoculture. In these graphs the monocultures and mixtures are seen as a 'replacement series' (de Wit, 1960), with the proportions of the genotypes in mixture being 1:0, 0.5:0.5 and 0:1, respectively.

At the 9 cm spacing, the RYT of the mixtures Glenlea+Pitic 62 and Glenlea+Norquay was greater than 1, and in both cases, the mixture yield surpassed the yield of the better component grown in monoculture.

At the 3 cm spacing, none of the RYT values were greater than 1, suggesting a detrimental effect of genotype interaction on plant yield at this spacing.




Fig. 2a-e. Graphic illustration of the genotype interaction effects on genotype and mixture yield in five genotype mixtures, each grown at two different interplant spacings in 1978. Along the X-axis, the populations containing 100% of the first genotype, 50% of each genotype, and 100% of the second genotype, are marked with the letters A, B and C, respectively.

6.3.4 Relationships between characters of interacting genotypes

One of the objectives of this study was to investigate the relationship between the measured values of characters on neighbouring genotypes. Correlations were therefore computed between the plant characters of each genotype in the test, and a number of characters of the associated genotypes (Table 7).

For most genotypes, the characters which played a role in plant interaction at the 3 cm and at the 9 cm spacings were not the same. Only Pitic 62 showed negative relationships of several characters with height of its associates at both spacings. In addition, the neighbour characters by which a genotype was affected differed from genotype to genotype.

Most characters of Pitic 62 were negatively correlated with the height of its associates. Since Pitic 62 was the latest maturing genotype in the test, and could throughout the growing season be identified in the field through its slow development, it is possible that this genotype showed more clearly the effects of interaction with taller neighbours, affecting plant weight, as well as grain yield and its components.

At the 3 cm spacing, most characters of Glenlea were positively associated with the weight and/or yield and the harvest index of its associates. Only tiller number was negatively related to the height of the associated

Table 7. Correlations¹ of single plant characters of five genotypes, with characters of associated mixture components, at two interplant spacings.

All plants were grown in four-row plots in 1978. The data pairs were averages of randomly selected plants, which had both neighbours present, and which were measured at maturity.

Char- acter ²		Characters of associated mixture components with which correlations occurred			
Inter- plant spacing (cm)	Plot 62	Glenn	Part	704009992	Horquay
T	3	FL ⁰⁰	T ⁰ , H ⁰	FL ⁰⁰ , H ⁰	HI ⁰⁰
H	3	FL ⁰⁰ , H ⁰	T ⁰ , H ⁰	FL ⁰⁰ , H ⁰	HI ⁰⁰
Wt	3	FL ⁰⁰ , H ⁰	T ⁰ , H ⁰	FL ⁰⁰ , H ⁰	FL ⁰⁰ , H ⁰
Y	3	FL ⁰⁰ , H ⁰	T ⁰ , H ⁰	FL ⁰⁰ , H ⁰	FL ⁰⁰ , H ⁰
K/P	3	FL ⁰⁰ , H ⁰	T ⁰	FL ⁰⁰ , H ⁰	FL ⁰⁰ , H ⁰
Ewt	3	FL ⁰⁰	Wt ⁰⁰ , Y ⁰⁰	Wt ⁰⁰ , Y ⁰⁰	FL ⁰⁰ , H ⁰
K/H	3	FL ⁰⁰	Wt ⁰⁰ , Y ⁰⁰	Wt ⁰⁰ , Y ⁰⁰	FL ⁰⁰ , H ⁰
HI	3	FL ⁰⁰	Wt ⁰⁰ , Y ⁰⁰	Wt ⁰⁰ , Y ⁰⁰	FL ⁰⁰ , H ⁰
PL	3	Y ⁰⁰ , Wt ⁰⁰	Wt ⁰⁰ , Y ⁰⁰	Wt ⁰⁰ , Y ⁰⁰	FL ⁰⁰ , H ⁰
Ht	3	FL ⁰⁰	Wt ⁰⁰ , Y ⁰⁰	Wt ⁰⁰ , Y ⁰⁰	FL ⁰⁰ , H ⁰
ILL	3	Y ⁰⁰ , Wt ⁰⁰	Wt ⁰⁰ , Y ⁰⁰	Wt ⁰⁰ , Y ⁰⁰	FL ⁰⁰ , H ⁰
ExL	3	FL ⁰⁰ , H ⁰	Wt ⁰⁰ , Y ⁰⁰	Wt ⁰⁰ , Y ⁰⁰	FL ⁰⁰ , H ⁰
Y/H	3	FL ⁰⁰ , H ⁰	Wt ⁰⁰ , Y ⁰⁰	Wt ⁰⁰ , Y ⁰⁰	FL ⁰⁰ , H ⁰
T	9	H ⁰	Wt ⁰⁰ , Y ⁰⁰	Wt ⁰⁰ , Y ⁰⁰	FL ⁰⁰ , H ⁰
H	9	FL ⁰⁰ , H ⁰	Wt ⁰⁰ , Y ⁰⁰	Wt ⁰⁰ , Y ⁰⁰	FL ⁰⁰ , H ⁰
Wt	9	FL ⁰⁰ , H ⁰	Wt ⁰⁰ , Y ⁰⁰	Wt ⁰⁰ , Y ⁰⁰	FL ⁰⁰ , H ⁰
Y	9	FL ⁰⁰ , H ⁰	Wt ⁰⁰ , Y ⁰⁰	Wt ⁰⁰ , Y ⁰⁰	FL ⁰⁰ , H ⁰
K/P	9	FL ⁰⁰ , H ⁰	Wt ⁰⁰ , Y ⁰⁰	Wt ⁰⁰ , Y ⁰⁰	FL ⁰⁰ , H ⁰
Ewt	9	FL ⁰⁰	Wt ⁰⁰ , Y ⁰⁰	Wt ⁰⁰ , Y ⁰⁰	FL ⁰⁰ , H ⁰
K/H	9	FL ⁰⁰	Wt ⁰⁰ , Y ⁰⁰	Wt ⁰⁰ , Y ⁰⁰	FL ⁰⁰ , H ⁰
HI	9	FL ⁰⁰	Wt ⁰⁰ , Y ⁰⁰	Wt ⁰⁰ , Y ⁰⁰	FL ⁰⁰ , H ⁰
FL	9	FL ⁰⁰	Wt ⁰⁰ , Y ⁰⁰	Wt ⁰⁰ , Y ⁰⁰	FL ⁰⁰ , H ⁰
Ht	9	FL ⁰⁰	Wt ⁰⁰ , Y ⁰⁰	Wt ⁰⁰ , Y ⁰⁰	FL ⁰⁰ , H ⁰
ILL	9	FL ⁰⁰	Wt ⁰⁰ , Y ⁰⁰	Wt ⁰⁰ , Y ⁰⁰	FL ⁰⁰ , H ⁰
ExL	9	FL ⁰⁰	Wt ⁰⁰ , Y ⁰⁰	Wt ⁰⁰ , Y ⁰⁰	FL ⁰⁰ , H ⁰
Y/H	9	FL ⁰⁰ , H ⁰	Wt ⁰⁰ , Y ⁰⁰	Wt ⁰⁰ , Y ⁰⁰	FL ⁰⁰ , H ⁰

1. * significant, $\alpha \leq 0.05$; ** significant, $\alpha \leq 0.01$.

2. Character abbreviations defined in Table 1. HL = Head length.

3. - negative correlations, + positive correlations.

genotypes. At the 9 cm spacing, however, weight, 1000 kernel weight, height and extrusion length of Glenlea were negatively correlated with yield, weight and harvest index of associated genotypes. It was earlier shown (Table 6), that Glenlea yielded better in most mixture situations than in monoculture. This could occur either if the demands of associates were less than those of Glenlea, leaving the latter a larger share of the available resources, or through 'hogging' of resources by Glenlea, resulting in suppression of the associated genotypes, accompanied by reductions in yield. RY values of less than 1, for the genotypes which were grown in combination with Glenlea (Table 6), support the latter hypothesis.

Park showed very few significant correlations at either the 3 cm or the 9 cm spacing. This could mean that either Park was not much affected by genotype interaction, or that Park was affected by its associates in an inconsistent manner. Since performance of Park in mixture was typically less than its performance in monoculture (Table 6), it can be concluded that Park responded differently to the various associated genotypes.

At the 3 cm spacing, many characters of both 70M009002 and Norquay were negatively correlated with the height of their associates, as was earlier described for Pitic 62. Unlike Pitic 62, however, these relationships did not exist at the 9 cm spacing. Pitic 62, 70M009002 and Norquay all have a medium height, but Pitic 62 tends to produce more

tillers (Appendix 7). It is possible that the tillers of Pitic 62 intermingle more with the tillers of neighbouring plants than do the tillers of the two other genotypes. This could result in effects of competition for light at both interplant spacings in mixtures containing Pitic 62, while for 70M009002 and Norquay, the increase in spacing from 3 cm to 9 cm was sufficient to reduce competition for light to an insignificant level.

Overall, it appears that competition for light plays an important role in the relationships between interacting genotypes, in particular for the shorter types and at the higher plant density. However, the negative correlations between the yield components of associated genotypes which occurred at the 9 cm spacing, indicate that competition for other growth factors did occur as well.

6.3.5 Interaction between genotypes during the early stages of growth

Mean values and standard deviations for characters measured on plants, grown in monoculture, at the flowering stage and at the mid-dough stage are reported in Appendices 10 and 11.

Again, correlations between characters of associated genotypes were calculated (Table 8). Positive correlations appeared to be more common than negative ones.

At the flowering stage, however, the flag leaf area of most genotypes appeared to be negatively correlated with

Table 8. Correlations¹ of single plant characters of five genotypes, with characters of associated mixture components, in machine seeded plots, at two growth stages.

All plants were grown in four-row plots in 1978. The data pairs were averages of randomly selected tillers.

Char- acter ²	Growth stage ³	Characters of associated mixture components with which correlations occurred			
		Pittic 62 4	Glenlea	Rark	Norquay
FL	F		ExL ^{oo} , A ^{oo}		ExL ^o
Ht	F				Ht ^{oo} , A ^{oo}
A	F	ExL ^{oo} , Ht ^o	A ^o	A ^o	Ht ^{oo} , A ^o
ExL	F		ExL ^{oo}	ExL ^{oo} , Ht ^{oo}	FL ^o
HL	F	FL ^o , Ht ^{oo}	ExL ^o , A ^o	ExL ^o , Ht ^o	ExL ^{oo}
FL	M				
Ht	M	ExL ^o		ExL ^o	ExL ^o
A	M	ExL ^o			FL ^o
ExL	M				
HL	M	ExL ^o		ExL ^o	

1. ^o significant, $\alpha \leq 0.05$; ^{oo} significant, $\alpha \leq 0.01$.

2. Character abbreviations defined in Table 1. HL = head length, A = Area of the flag leaf blade.

3. F = Flowering stage, M = Mid-dough stage.

4. - negative correlations, + positive correlations.

height components in associated genotypes. Variations in photosynthesis due to competition for light, or other factors correlated with height, could have affected leaf growth. This effect was not observed at the mid-dough stage, suggesting that leaf senescence, which sets in earlier in some genotypes, and in some plants within genotypes, than in others, caused a greater variation in the measurements of this character at flowering.

Plant height and height of the flag leaf of Norquay were negatively correlated with height of its associates at the flowering stage, but not at the mid-dough stage.

Overall, it appears that the effects of genotype interaction on the expression of the measured characters at both stages of growth were small. This might be partially caused by the greater amount of plant-to-plant variation which occurs in machine seeded plots than in plots with accurately spaced plants. Interaction associated with differences in height, however, appeared to occur at both stages of growth.

6.4 Discussion

6.4.1 Effects of genotype interaction on plant-to-plant variation

The variances of measurements of plants grown in monocultures and plants grown in mixtures were not

significantly different, possibly because a systematic planting pattern was used. A genotype, which is grown in a mixed stand in which two genotypes are alternated and accurately spaced within the row, will always have the same type of neighbours, even though these are different from neighbouring plants in a monoculture. This consistency in neighbour relationships may explain the observed lack in variation.

Sakai (1955) observed that a genotype which tends to dominate in a mixture shows greater variation when grown in mixture with a less dominant type, than when grown alone, or in combination with another dominating type. He did not discuss any changes in variation which could occur in a mixture of genotypes with an equal ability to dominate. One could, however, expect effects on within genotype variation which are related to the relative amount of random variation present in the two genotypes. A variable genotype, in combination with a uniform genotype, could conceivably show a reduction in variation, because the environment it experiences in such a mixture is more uniform than its environment in monoculture. Similarly, the other mixture component could show an increase in variation.

Any of these situations would result in a significant genotype x mixture interaction effect on plant-to-plant variation. In the present study, however, this interaction was only significant for 1000 kernel weight (Table 3).

When observing for each genotype the combined data from several binary mixtures, however, an increase in plant-to-plant variation was found for the number of tillers, weight and yield at the 3 cm spacing, and for kernel weight at the 9 cm spacing (Table 4). These effects may well be the result of the greater amount of environmental variation which is caused by the range of genotypes in the mixtures.

At both the 3 cm and the 9 cm spacing, the distribution of several characters, in particular kernel weight and harvest index, deviated significantly from normal. Although the analysis of variance is insensitive to small deviations from normality, other conclusions based on estimates of variation might be biased to a greater extent. The effect of non-normality on the expected gain from selection in a breeding program will be discussed in Chapter 9.

Chebib et al. (1973) found that in a mixture of segregating F3 lines of wheat, the effect of interaction between genotypes on plant-to-plant variation was insignificant for all measured characters. His results might be partially explained though, by the fact that he obtained his segregating population from a cross between the genotypes Pembina and Manitou, which have a similar genetic back ground', and therefore would produce a progeny which is

 'Pembina=Thatcher x R.L.2564(R.L.2564=McMurchy-Exchange x Redman')
 Manitou=(Thatcher'-Frontana x Thatcher'-Kenya Farmer) x Thatcher'-P.I.170925

genetically and morphologically much more uniform than the progeny from a more diverse cross would be.

Q Kelker and Briggs (1978) reported both increases and decreases in variation as a response to interaction of several genotypes. In their experiment, seven genotypes of wheat, including the ones used in the present study, were mixed within rows, simulating a segregating population. It was concluded that the tendency to increase or decrease in variation is genotype-specific.

The number of genotypes in the present study was not large enough to be able to verify Kelker and Briggs' conclusion, although the results from the present experiment do not disagree with their results. In Kelker and Briggs' study, the variation of the genotypes Glenlea, Park and 70M009002 each increased in variation due to genotype interaction at a density of 87 plants/ m², in particular for the characters heads per plant, yield and plant weight. In addition, variation for height of Pitic 62, Glenlea, and Norquay, and harvest index of Pitic 62, Glenlea, Park and Norquay decreased. Similar effects were observed in the present study (Appendix 7).

6.4.2 Effects of genotype interaction on mean plant performance

Significant changes in the mean values of several characters, related to either total plant weight or to plant yield, were observed, as a response to genotype interaction.

Harvest index, which is the ratio of plant weight and plant yield, was very little affected, illustrating the stability which a complex character can maintain in spite of the plasticity of its components.

For some genotypes, the response to genotype interaction was quite different at 3 cm and at 9 cm spacing. This was demonstrated most clearly by Glenlea, the characters of which showed positive correlations with most characters of neighbours at the 3 cm spacing, while negative correlations were more prevalent at the 9 cm spacing. Thus, a change in interplant relationships occurred with increasing plant spacing. A possible explanation is that at the 3 cm spacing, the available growth factors were limiting to both genotypes in the mixtures, thus preventing the expression of genotypic differences. At the 9 cm spacing, resources were less limiting and plants would have a greater chance to develop according to their genetic potential. This could have resulted in negative relationships between the measured characters of Glenlea and its associates at the 9 cm but not at the 3 cm spacing. This hypothesis is supported by the observation in 1977, that effects of genotype interaction were insignificant at the highest plant density, where all plants were severely reduced in growth.

Competitive interaction between plants frequently leads to negative correlations between characters of the genotypes combined in a mixture. However, positive correlations are observed as well. Interaction between plants of different

oat species, resulting in positive correlations between the height of mixture components, has been described by Trenbath and Harper (1973).

Factors which cause differences between replicates, such as soil heterogeneity or a difference in elevation, could also be the cause of a positive correlation between characters of genotypes which are combined in a mixture. In the present study, however, it was not possible to distinguish between positive correlations due to plant interaction and those due to other factors.

There is some evidence in the literature (Troughton and Whittington, 1968), that maturity is positively correlated with root mass, because late flowering plants can invest more time in the development of an extensive root system. If maturity is positively correlated with root mass, then Pitic and Glenlea, which were the latest maturing genotypes in this study, could have had a competitive advantage through a larger root system. This could cause negative correlations of other characters specific to these genotypes, such as the number of tillers or height, with the yield components of their associates. Whether this was the case can only be confirmed through studies involving measurements on the root systems during the growing season. This hypothesis could also explain why Park, which was the earliest genotype in the study, generally performed poorly in mixture.

At the 9 cm spacing, negative correlations existed between kernel weight and the number of tillers and the

number of heads on neighbours. These relationships suggest a different type of interaction than observed at the 3 cm spacing. Although interactions related to the greater nutrient requirements of plants with more tillers are probably of importance, one has also to consider the effect which variation in the number of tillers could have on the competition for light between neighbouring plants. It is possible that at the lower plant density, plants with many tillers interfere more with the light reception of their neighbors than do plants with less tillers, resulting in the observed correlations.

Interaction between genotypes in a mixture occurs throughout most of the growing season. In addition to measurements at maturity, a detailed study of the growth and development of each mixture component could therefore provide information on the causal basis for the observed results of interaction. Several studies have indicated that differences in the developmental pattern of neighbouring plants greatly influence their interaction (Christian and Gray, 1941; Trenbath and Harper, 1973; Trenbath, 1975; Harper, 1977; Clark, 1980). The present study was not designed to perform such a detailed growth analysis. However, data collected at different times during the summer suggest that interactions occurring before flowering, and specifically those affecting growth of the flag leaf, may ultimately affect grain filling through an altered photosynthetic area (Hsu and Walton, 1971).

7. Effects of Seed Size on Plant Interaction

7.1 Materials and methods

In 1979, seed of the genotypes Pitic 62, Glenlea, Park, Norquay, and 70M009002 was sorted, through sieving and hand picking, into large, medium, and small size seeds (Appendix 12). The medium size seeds were discarded, and the rest seeded in plots, containing only large seeds, only small seeds, or small and large seeds alternated within the rows.

All seeds were glued onto strings (see Section 5.1.2) with 3 cm between seeds. The plots each consisted of three 2.5 m long rows, 30 cm apart, and all treatments were replicated three times. Eighteen adjacent plants (54 cm) from the middle rows were sampled at the 3-5 leaf stage, at the jointing stage, at the heading stage and at maturity. Since sampling at each time was completed in one day, the five genotypes were not at exactly the same stage of development.

The locations of the samples were randomised within each row, and two plants at the end of each sample were allowed as guards. At the time of sampling, all plants in the section of row to be harvested were numbered, and measurements were taken on each plant. Empty places in the row were recorded and plants with one or both first neighbours missing were discarded.

The characters which were measured varied with the time of sampling, as indicated in Table 9, and were chosen, for each developmental stage, as features which could easily be distinguished.

7.2 Analysis of the data

The data were analysed to investigate the effects of seed size and uniformity on plant performance, on within genotype variation, and on the interaction between neighbouring plants.

Variance, skewness and kurtosis of the measured characters were computed for each sample. The effects of genotype, seed size and uniformity on these parameters were examined for each sample date with an analysis of variance. Because several of the distributions of measured characters deviated significantly from normality, a natural log transformation of the variances was used (see Section 6.2)

Because plants were missing at the end of some rows, it was very difficult to identify the exact sequence number of each plant. This especially created a problem in the plots with mixed seed. It was, therefore, impossible to compare the performance of plants from large and small seeds in mixture, with the performance of the same type of plants in plots from uniform seed.

Differences between the genotypes, between the replicates, and between the performance of plants from

Table 9. Plant characters which were measured or derived at four stages of growth in 1979.

Measured Character	Abbreviation	Growth Stage ¹			
		1	2	3	4
Height (to tip of longest leaf) (cm)	Ht ₁	x			
Height (of top most node) (cm)	Ht ₂		x		
Height (to tip of head, including awns) (cm)	Ht ₃			x	
Number of leaves	L _m	x			
Number of nodes	N		x		
Dry weight (g)	Wt	x	x	x	x
Number of tillers	T	x	x	x	x
Number of heads	H				x
Height of flag leaf (cm)	FL				x
Number of spikelets/head	Sp/H				x
Yield/plant (g)	Y				x
Number of kernels/plant	K/P _{max}				x

Derived Character	Growth Stage ¹			
	Abbreviation	1	2	3 4
Harvest Index = Yield (g)/Dry Weight (g)	HI			x
1000 kernel weight = Yield (g)/Number of kernels x 1000 (g)	Kwt			x
Number of kernels/head = Number of kernels/Number of heads	K/H			x
Extrusion length = Height - height of flag leaf (cm)	ExL			x

1. 3-5 leaf stage

2. jointing stage

3. heading stage

4. maturity

large, small, and mixed seed were examined, for each sample date, with an analysis of variance.

To investigate the interactions between adjacent plants from small, large, or mixed seed, simple correlation coefficients were computed for all character combinations on pairs of neighbouring plants (1st with 2nd, 2nd with 3rd, etc.).

It was intended to compute these correlations for second neighbours as well as for first neighbours. Missing plants, however, severely reduced the number of pairs for which correlations could be computed. For every plant which was missing, four of the first neighbour pairs and five of the second neighbour pairs were invalidated, or 23.5% and 31.3% respectively of the data. Since an average of 1.6 plants per plot were missing (10%), correlations on second neighbours have not been included.

In 1978, it was noticed that considerable variation in height existed among plants of the line 70M009002. At the end of the 1979 growing season, it became apparent that this genotype was not a pure line. There was a marked variation in height and awn length, and it was decided to eliminate this line from all tests involving single plant data. No meaningful comparisons concerning plant interactions can be made between treatments if there are already genetic differences in plant form, and possibly growth pattern, within the line under investigation. It is now believed that this line possessed a degree of outcrossing, perhaps as high

as 5%, which would explain the increase in heterogeneity as seed was increased during the three years of this study (Dr. K. G. Briggs, Professor of Plant Breeding, University of Alberta, Edmonton, Alberta, 1980, personal communication).

7.3 Results

7.3.1 Effects of seed size on mean plant performance and sample distribution of characters

Mean values of the plants from small seeds, from large seeds, and from mixed seeds, at each of the measured growth stages are shown in Appendices 13-16. Values of skewness and kurtosis of the samples are given in Appendix 17.

At the 3-5 leaf stage the plants from mixed seed had a significantly lower variance ($\alpha \leq 0.05$) for number of leaves than the plants from the uniform seed (Table 10). At maturity, only the variance of head length was significantly ($\alpha \leq 0.05$) affected by seed treatment. The head length of plants from small seed was much more variable than was the head length of plants from large or mixed seed.

The skewness and kurtosis of the characters within sample dates were not significantly affected by the treatments. However, distributions did differ significantly among growth stages for those characters which could be compared: the number of tillers, weight and height (sample 3, 4 only, Table 11). Kurtosis of all three characters was

Table 10. The effect of genotype and seed treatment¹ on the variance² of single plant characters, at different stages of growth, as determined by analyses of variance³.

All plants were grown in three-row plots in 1979. The data are from consecutive plants in a row, excluding plants with missing neighbours.

Source of variation	df	Mean squares of characters ⁴	
		Number of leaves at the 3-5 leaf stage	Head length at maturity
Genotype (G)	3	9591	13978
Seed treatment (S)	2	15672*	24183*
Replicate (R)	2	667	14899
G × S	6	4458	6429
G × R	6	11539	7085
S × R	4	5445	7661
Residual	12	3729	6127

1. Seeds were sorted into large and small seeds (see Appendix 10) and seeded in rows with uniformly large seeds (treatment 1), uniformly small seeds (treatment 2), or large and small seeds alternated (treatment 3).
2. The natural logarithm of the variances was used.
3. * significant, $\alpha \leq 0.05$.
4. Characters which showed no significant differences have been excluded.

Table 11. The effect of growth stage, genotype, and seed treatment¹ on the variance² and skewness of single plant characters, as determined by analyses of variance³.

All plants were grown in three-row plots in 1979. The data are from consecutive plants in a row, excluding plants with missing neighbours.

Source of variation	df	Mean squares of distribution parameters				
		Variance			Skewness	
		Ht ⁴	T	Wt	Ht	T
Growth stage (GS)	3	12.48**	25.68**	554.14**	3.26	5.14*
Genotype (G)	3	2.03**	4.80**	1.23	0.36	2.16*
Seed treatment (S)	2	0.11	0.68	1.87	1.08	0.29
Replication (R)	2	0.26	0.72	0.82	0.10	1.04
GS × G	9	2.51**	1.07	3.13	0.33	0.66
GS × S	6	0.76	0.53	1.82	1.55	1.05
GS × R	6	0.26	0.29	0.59	2.60	0.72
G × S	6	0.57	0.44	1.49	1.62	0.24
G × R	6	1.07	0.40	1.74	0.35	1.93
S × R	4	0.51	0.56	4.39*	1.68	0.76
Residual	12	0.47	0.57	1.71	0.85	0.79

1. As defined in Table 10.

2. The natural logarithm of the variances was used.

3. * significant, $\alpha \leq 0.05$; ** significant, $\alpha \leq 0.01$.

4. Character abbreviations defined in Table 1.

unaffected by growth stage, as was skewness of the weight distribution. However, the skewness of the distribution of tiller number changed significantly ($\alpha \leq 0.01$) from an, on the average, negative value at the first three sample dates, to a positive one at maturity (Appendix 17).

The skewness of the height distribution increased from an average value of -0.34 at the heading stage, to a value of -0.69 during the last part of the growing season (Appendix 8). Although the difference has only a significance level of $\alpha = 0.057$ it does illustrate the increase in skewness which was earlier observed by Koyama and Kira (1956).

At each growth stage, differences among genotypes were significant for most characters (Table 12a through d). At the first three growth stages, the genotype x treatment interactions were more often significant than were the treatment effects. This suggests that genotypes were affected differently by the various seed treatments throughout the growing season. At maturity, the seed treatment main effects and the genotype x seed treatment interactions were predominantly insignificant, as were the seed treatment effects. Height and dry weight, which had shown significant seed treatment effects at the jointing and at the heading stage, did not show these effects any longer. Because maximum height was not obtained by most plants until after heading, this disappearance of seed treatment effects illustrates the tendency of plants to grow out to a uniform

Table 12a. The effect of genotype and seed treatment¹ on single plant characters, measured at the 3-5 leaf stage, as determined by analyses of variance.²

All plants were grown in three-row plots in 1979. The data are from consecutive plants in a row, excluding plants with missing neighbours.

Source of variation	df	Mean squares of characters ³			
		Ht ₁	L	T	Wt (x 10 ³)
Genotypes (G)	3	322**	0.28	8.31**	5**
Seed treatment (S)	2	10	0.82	0.97	3
Replicates (R)	2	47**	4.46**	0.68	2
G × S	6	23	2.16**	2.77**	3*
G × R	6	58	1.39**	1.45	4**
S × R	4	1	6.87**	1.47	1
G × S × R	12	31	2.02**	1.74**	3*
Residual	392	12	0.32	0.70	1

1. As defined in Table 10.

2. * significant, $\alpha \leq 0.05$; ** significant, $\alpha \leq 0.01$.

3. Character abbreviations defined in Table 9.

Table 12b. The effect of genotype and seed treatment¹ on single plant characters, measured at the jointing stage, as determined by analysis of variance.²

All plants were grown in three-row plots in 1979. The data are from consecutive plants in a row, excluding plants with missing neighbours.

Source of variation	df	Mean squares of characters ³			
		Ht ₂	N	T	Wt
Genotypes (G)	3	147**	11.21**	12.29	3.24**
Seed treatment (S)	2	61*	0.26	6.99	0.58
Replicates (R)	2	78**	12.03**	3.39	1.74**
G × S	6	51**	2.03**	4.56	0.62*
G × R	6	36*	0.84	7.42	0.44
S × R	4	116**	2.05**	5.55	0.38
G × S × R	11	63**	2.12**	3.77	0.46
Residual	315	17	0.45	4.90	0.28

1. As defined in Table 10.

2. * significant, $\alpha \leq 0.05$; ** significant, $\alpha \leq 0.01$.

3. Character abbreviations defined in Table 9.

Table 12c. The effect of genotype and seed treatment¹ on single plant characters, measured at the heading stage, as determined by analyses of variance.²

All plants were grown in three-row plots in 1979. The data are from consecutive plants in a row, excluding plants with missing neighbours.

Source of variation	df	Mean squares of characters ³		
		Ht ₃	T	Wt
Genotypes	3	18462**	68.7**	1.69
Seed treatment	2	200**	0.7	9.63*
Replicates	2	187**	12.7*	2.08
G × S	6	270**	13.0**	9.77**
G × R	6	45	7.5	3.34
S × R	4	283**	6.1	7.07*
G × S × R	12	134**	7.5*	3.57
Residuals	383	28	4.0	2.67

1. As defined in Table 10.

2. * significant, $\alpha \leq 0.05$; ** significant, $\alpha \leq 0.01$.

3. Character abbreviations defined in Table 9.

Table 12d. The effect of genotype and seed treatment¹ on single plant characters, measured at maturity, as determined by analysis of variance².

All plants were grown in three-row plots in 1979. The data are from consecutive plants in a row, excluding plants with missing neighbours.

Source of variation	df	Mean squares of characters ³					
		Ht ₃	T	H	Wt	Y	K/P(x 10 ⁻²) HI(x 10 ³)
Genotypes	(G) 3	15437**	73.9**	55.1**	1297**	339.6**	2156** 73**
Seed treatment (S)	2	34	11.7*	9.4*	48	10.7	145 3
Replicates	(R) 2	51	14.1*	14.6**	58	10.4	145 1
G x S	6	66	5.0	4.6	85	20.6*	153* 3
G x R	6	17	5.5	5.2	84	12.9	77 7**
S x R	4	42	11.4**	4.6	28	4.8	55 4
G x S x R	12	69	10.4**	7.2**	100*	10.3	84 7**
Residual	404	59	3.4	2.9	50	40.6	58 2

Source of variation	df	Mean squares of characters					
		Kwt	K/H	Fl	ExL	Sp/H	HL
Genotypes	3	4557.6**	3582**	8409**	5606**	11.2**	508.1**
Seed treatment (S)	2	42.9**	43	30	10	0.4	0.6
Replicates (R)	2	115.2**	184*	38	25	11.6**	0.7
G x S	6	7.1	94	94	15	1.1	3.7
G x R	6	19.5*	44	46	11	4.4*	16.9**
S x R	4	13.0	55	6	15	3.1	7.3*
G x S x R	12	15.6*	103*	71	4	2.1	8.4**
Residual	404	8.9	56	47	17	1.7	3.0

1 As defined in Table 10.

2 * significant, $\alpha \leq 0.05$; ** significant, $\alpha \leq 0.01$.

3 Character abbreviations defined in Table 9.

height, which is characteristic of their genotype.

The mean tiller number per plant dropped from jointing to maturity, for all genotypes except Park (Appendices 14 through 16). Tiller number was not significantly affected by seed treatment at the jointing and the heading stages, a significant genotype x seed treatment interaction ($\alpha \leq 0.01$) occurred at the heading stage, and at maturity, a significant treatment effect ($\alpha \leq 0.05$) existed for the number of tillers and the number of heads per plant (Table 12b through d).

At maturity, the plants from small seeds had fewer tillers and heads than did the plants from large seeds, for all genotypes except Glenlea (Appendix 16). Since seed treatment effects earlier in the season were not significant, this suggests that plants from small seed on the average lost more tillers than did plants from large seeds. Thousand kernel weight of the plants from large seeds was significantly smaller ($\alpha \leq 0.01$) than that of the plants from small seeds (Table 12d).

7.3.2 Effects of seed size on interaction between neighbours

The correlations of all measured characters of each plant with those of their first neighbours gave an unwieldy total of 1572 correlation coefficients. It would be of little value to look at the magnitude of each of these coefficients individually. More important is to see if any patterns in character relations exist. If no correlation

exists between two characters of neighbouring plants, the probability that the coefficients for all four genotypes in a treatment will be of the same sign just by chance is $1 \times 1/2 \times 1/2 \times 1/2 = 0.125$. Therefore, the value of the coefficients does not have to be very high to be meaningful, if they all have the same sign. In order to accept the hypothesis that a correlation exists between characters measured on neighbouring plants, in each of the four genotypes given the same treatment, with $\alpha \leq 0.01$, each of the coefficients has to have the same sign and a significance level less than or equal to 0.47, since $2 \times (0.47)^4 = 0.01$. Using this value, all correlation coefficients were examined for each of the three treatments (Table 13).

Throughout development, positive correlations were more common than negative ones. As was explained in Section 6.4.2, plant interactions as well as environmental conditions can be responsible for this. Conditions which affect a short section of row, such as variation in depth of sowing, wheel compaction or irregular nutrient distribution, could result in higher or lower values for all plants in that section, compared with plants in the rest of the plot, and, thus, result in positive correlations between neighbours.

At the jointing stage, the only negative correlation was the one between tiller number and the number of nodes of neighbouring plants, derived from mixed seed sizes.

Table 13. Correlated¹ of neighbouring plants, at different stages of growth in rows of plants from large seeds, small seeds or alternated large and small seeds.

All plants were grown in three-row plots in 1979. The data are from overlapping pairs of neighbouring plants in a row: 1st+2nd, 2nd+3rd, 3rd+4th, etc.

Growth ¹ stage	Character ³	Correlated (+, -) characters of neighbouring plants ⁴		
3-SL	Ht ₁	+Ht ₁ 2,	+Ht ₁ 3	
3-SL	L	+Ht ₁ 3,	+L1,	+L3
3-SL	Wt	+Ht ₁ 3		
J	Ht ₂	+Ht ₂ 1,	+Ht ₂ 2	
J	N	+Ht ₂ 1,	+N1,	+N3
J	T	+N1,	-N3,	+T1
J	Wt	+Ht ₂ 2,	+N2	
H	Ht ₃	+Ht ₃ 1,	+Ht ₃ 2	+Ht ₃ 3
H	T	+T2,	-T3	
H	Wt	+Ht ₃ 1,	+Ht ₃ 2,	+Wt2
M	Ht ₃	-KWt ₃		
M	T	-HI1,	-KWt1	
M	H	-HI1,	-KWt1	
M	K/H	-KWt1		
M	HI	+HI1		
M	KWt	+KWt1		

1. All correlations are significant with $\alpha \leq 0.01$.
2. 3-SL = 3-5 Leaf stage, J = Jointing stage, H = Heading Stage,
M = Maturity.
3. Character abbreviations defined in Table 9.
4. Last number indicates seed treatment as defined in Table 10.

At the heading stage, the tiller numbers of neighbouring plants, derived from mixed seed, were negatively correlated. However, at this stage, tiller number of some genotypes was affected by seed treatment also (Table 12c), and the observed correlation, therefore, does not necessarily imply the occurrence of neighbour interactions. At maturity, a significant negative correlation existed between the number of tillers and harvest index and between number of heads and harvest index of plants from large seeds, and between height and 1000 kernel weight, and between stem length and 1000 kernel weight, of plants from mixed seed. Since seed size effects on tiller number, head number and kernel weight were observed in plots seeded to uniformly sized seed, one would expect negative correlations between these characters, in plots from mixed seed sizes. Their absence indicates possible effects of neighbour interaction. For plants derived from small seeds, significant correlations were all positive.

7.4 Discussion

The drop in tiller number, which was observed after heading for plants from all seed treatments, was more severe for plants from small seed than for those from mixed or large seed. This resulted in significantly fewer tillers and fewer heads on plants from small seeds at maturity.

For most genotypes, however, yield per plant was not significantly affected by seed treatment. The larger 1000 kernel weight, observed for plants derived from small seeds, demonstrates compensation for the reduction in head number, through an increase in seed weight. A number of factors could have contributed to this result. Firstly, some of the assimilates which were acquired before anthesis, and which had been stored in the lost tillers may have been relocated and later been used in grain filling on the remaining tillers (Evans and Wardlaw, 1976). Secondly, since competition for light takes place between leaves rather than between plants (Donald, 1961), leaves of plants with few tillers could experience less competitive interference from leaves on other tillers than would leaves on plants with many tillers, given the same interplant spacing. A higher rate of photosynthesis per leaf could therefore occur in plants with few tillers, resulting in more assimilates per tiller and, thus, the potential for larger kernels.

The data from this test give only partial support to the findings of Hozumi et al. (1955), who found negative correlations between dry weights and between shoot lengths of neighbouring corn plants derived from uniform seed. Negative correlations did occur in the present experiment, but not very frequently. The negative correlations which were observed among plants from large seed, all occurred between tillers or heads per plant and those characters which pertain to grain filling (harvest index, 1000 kernel

weight). This suggests that variation in tiller number is, directly or indirectly, related to the process of grainfilling in adjacent plants. No negative correlations showed up in the plots from small seed. This could possibly be the result of smaller root systems, which would reduce interaction between neighbouring plants.

Although alternating height differences between neighbours existed throughout the growing season in all three treatments, at maturity only the mixed seed treatment showed significant effects related to differences in height. It was not possible to determine with certainty whether small seeds gave rise to short or tall plants. If one assumes the former, than seed size in the mixed seed treatment had an effect on kernel weight opposite to the effect observed in the uniform seed treatments. The negative correlation, which occurred in the mixed seed treatment between height and kernel weight, suggests that plants from small seed with tall neighbours yielded seeds with low kernel weight and vice versa. Thus, interaction between the tall and short plants affected plant growth in such a way that tall plants were able to acquire a relatively greater amount of one or more of the factors limiting to grain filling. It thus appears that interactions between neighbours derived from different size seeds, were different than those which occurred between plants from uniform seeds, affecting in particular the character kernel weight.

The results of the present experiment indicate that differences in seed size can be a source of variation among phenotypes, both through effects on plant development as well as through effects on neighbour relationships. Although these conclusions are based on observations in monocultures, similar effects can be expected in a mixture of genotypes. Effects of interaction between genotypes in a segregating population might thus be confounded with the effects of seed size which occurred among the seeds of the propagated lines.

8. The Relationship Between Mixture Yield and Monoculture Yield of the Components.

8.1 Materials and methods

Yield trials were grown in 1977 and 1979 in Edmonton, and in 1978 in Ellerslie. The trials involved eight genotypes (see Section 4.2), grown in monoculture and in all possible binary combinations, arranged in a randomised block design with three replications.

Plots consisted of four 6 m long rows, spaced 23 cm apart, in banks which were separated by 2 m wide pathways. The rows were seeded in all three years in a N-S direction, guard plots were seeded at the end of each bank, and at harvest, a 30 cm strip at both ends of each plot was removed. Only the center two rows of each plot were harvested with a small plot harvester¹. Using a visual assesment of maturity, each plot was harvested when more than 50% of the plants appeared to be ripe.

To obtain an equal number of plants of each genotype per plot, percent germination and 1000 kernel weight of each genotype were used to calculate the weight of seed necessary to give an average density of 257 seeds/m², or an average of 1.7 cm between seeds within rows. For an average weight of 40 g/1000 kernels, this seeding rate is equivalent to 100 kg viable seed per ha. These are similar seeding rates and

¹Designed and produced by J. Fitzsimmons and T. Snider, Dept. of Genetics, University of Alberta, Edmonton, Alberta.

plant spacings to those normally used for standard yield trials of wheat at both stations.

The plots were seeded with a small-plot disk seeder, designed in 1970 by the Canadian Department of Agriculture, Swift Current, Saskatchewan. A cone-shaped seed divider was attached to partition the seed equally into the four drill rows.

Harvesting conditions in 1978 were very wet, causing some loss of yield and considerable lodging in one corner of the field. Because the amount of loss was hard to estimate, the data have not been adjusted. None of the genotypes was infested with significant levels of disease or insect pests in any of the three years.

8.2 Analysis of the data

It was the objective of this study, to observe the stability of mixture yield over years, and to investigate the relationship between mixture yield and the yield of their components grown in monoculture.

By means of a randomised block analysis of variance, the variation among years, among monocultures, among mixtures and between mixtures and monocultures was examined.

From the yields of the monocultures in each block (replicate), the expected yield of each mixture in the same block was calculated as:

$$Y(AB) = Y(A) + Y(B) / 2$$

where $Y(AB)$, $Y(A)$ and $Y(B)$ are the plot yields of the mixture of genotypes A and B, and their monocultures, respectively. The ratio of the observed yield and the expected yield of a mixture of two genotypes, hereafter referred to as Specific Mixture Efficiency (SME), was calculated as:

using the same notation as above. If, for example, a mixture of the genotypes A and B has a plot yield of 1400 g, and the monoculture plot yields of A and B are 1000 g and 1500 g, respectively, then the SME of the mixture AB would be $2 \times 1400 / (1000 + 1500) = 1.12$.

The SME resembles the RYT which was used in Chapter 6. The RYT, however, is calculated from the RY of each of the components, and hence, will have values which are numerically slightly different from the SME values. The meaning of values less than one, greater than one, and equal to 1 is the same for both, however.

Significance of the SME values was tested with a t-test analogous to the one described for RYT (see Section 6.2).

Spearman's rank order correlation coefficients were computed for the correlations between grain yield of the mixtures and SME in each of the three years, as well as for the correlations between yields in 1977 and 1978, yields in 1977 and 1979, and yields in 1978 and 1979, respectively. The same correlations were computed for the SME values.

Mixtures of two genotypes can be seen as analogues of the F1 generation of a genetic cross, and hence, techniques similar to those which are used to determine genetic parameters, can be used to analyse mixture properties. An analysis of variance for diallel crosses was developed by Hayman (1954). In his method, the total variation is partitioned into a component due to 'arrays' (columns of the diallel table), a set of three orthogonal interaction components, and an error term. Morley Jones (1965) described the use of this technique to analyse data from a half diallel.

Hayman's technique of diallel analysis has been modified and used by various authors to analyse competition experiments (Williams, 1962; Eberhart et al., 1964). In most of those cases, the competing individuals could be readily identified, allowing data to be collected from each of the components of the mixtures. Hay (1974) described the use of Hayman's analysis of variance for experiments in which the competing individuals cannot be separated.

The components of variation in the Hayman analysis are:

1. The array sum of squares, with $n-1$ degrees of freedom for an $n \times n$ diallel. Each column of the full diallel table comprises an array. The examination of the arrays of a half diallel thus requires that each data point is used twice in the computation. Morley Jones (1965) computed the SS for arrays as follows:

where Y_r is the sum of the r th array and Y_{rr} is the value of the r th monoculture (diagonal values of the diallel table). Significant differences among the arrays would indicate that differences exist between the general ability of the genotypes to perform in a mixture.

2. An interaction term, b , which can be partitioned into three orthogonal components:
 - a. b_1 tests the effect of genotype interaction and allows examination of the difference between mixtures and monocultures, with 1 degree of freedom.
 - b. b_2 tests the differences between genotypes in their interaction with all other genotypes. The sums of squares are computed using for each array the differences between twice the mixture values and the monoculture values:

Using the same notation as above, b_2 has $n-1$ degrees of freedom. If no interactions between genotypes occur, this component measures strictly the differences in yielding ability among genotypes.

- c. b_3 tests the effect due to specific combinations of genotypes and is computed by subtraction of the SS's for arrays for b_1 and b_2 from the total SS.

The formula which was used by Morley Jones for array SS, was used by Griffing (1956) to test differences in General Combining Ability (GCA). He used the remaining variation as a whole, to determine effects of specific combining ability (SCA). The use of Griffing's technique for the analysis of mixture experiments was suggested by Jensen and Federer (1965). Hayman's partitioning of the interaction SS, however, allows a closer examination of the types of interaction which take place and of their relative magnitude.

GCA is a parameter which was originally introduced to assess the value of a genotype for breeding purposes. An extensive review of this subject was given by Sprague and Tatum (1942). Because high yield is an objective of most breeding programs, an indicator of both yielding ability of a genotype and its value as a parent in crosses, is useful for the selection of breeding material. In experiments which study the mechanisms underlying mixture efficiency, however, the yield of a mixture is only of interest in relation to the monoculture yields of its components. Average Mixture Efficiency (AME) can be expressed as the ratio of the observed mixture yields to the mixture yields which can be expected based on the monoculture yields of the mixture components:

using the same notation as before. If, for example, binary

mixtures are formed from the genotypes A, B, and C, and the monoculture plot yields are 900 g, 950 g, and 1000 g, respectively, and the mixture plot yields are 1000 g each, the AME for genotype A would be

$$2 \times 3000 / (3 \times 900 + 900 + 950 + 1000) = 1.09.$$

The AME provides a measure of the average effect of a genotype on mixture yield, using its monoculture yield as a point of reference. The GCA gives an indication of a genotype's ability to perform in mixture using the experiment mean as a point of reference.

Jinks (1954) suggested the use of regression of the covariances (W) of offspring and parents of a diallel cross on the variances (V) of the offspring, to investigate dominance relationships. Harper (1965) and Hill and Shimamoto (1973) used a similar technique to determine the genotypic differences in 'competitive ability' of plants grown in mixtures. For the analysis of mixture experiments, the positions of the array variances along the X-axis of the resulting graph indicate the ability of the genotypes to dominate in the mixture. Dominating types will influence the performance of the mixtures in which they grow to a larger extent than will less dominating types, resulting in more uniform array components. Their arrays are, therefore, expected to have smaller variances and covariances. As a result these values will lie near the origin of the graph. Without any specific interactions between genotypes, a regression line can be fitted with a slope of approximately

1. Deviations from 1 will result from mixtures which perform much better or much worse than expected, causing an increase in variance and a decrease in covariance.

If a linear relationship exists between V and W, the sum ($W + V$) can be correlated with the monoculture yields (or any other character that might play a role in genotype interaction) to determine whether there is an association of monoculture yield with the ability to dominate in a mixture.

For the analysis of variance of the mixture yield trials from 1977, 1978 and 1979, Hay's directions (1974) were followed. Regression lines of W on V were fitted according to Jink's method (Jinks, 1954). SME and AME values were computed, and for comparison, Griffing's (1956) formula was used to compute the effects of General and Specific Combining Ability (GCA and SCA) for each genotype or genotype combination. In order to assess the significance of the AME values, a t-test was used to compare the observed mixture yield with the expected yield based on monoculture values. Significance of the GCA values was tested with an F-test, using MS values obtained from the analysis of variance. Correlations between monoculture yields and AME were computed, to investigate the relationship between AME and yielding ability.

8.3 Results

In spite of satisfactory performance in the 1977 germination test (84.7%), Pitic 62 failed to germinate in the field that year, and in 1977, all plots involving this genotype had to be discarded.

Monoculture and mixture yields varied substantially in the three years (Appendix 18a through c). Besides the different locations involved, the weather patterns in the three years were quite dissimilar (Appendix 1), which probably contributed to these yield differences.

The yield levels of the mixtures, in relation to those of the component monocultures, were divided into the following four categories (Table 14):

1. below the monoculture yield of the lower yielding component in that year (C1),
2. between the monoculture yield of C1 and the mean of the component monoculture yields (C2),
3. between C2 and the monoculture yield of the higher yielding component (C3), and
4. above C3.

Significantly ($\alpha \leq 0.05$) more than half of the mixtures yielded more than the mid-component yield, suggesting, on the average, a greater efficiency of resource utilisation by mixtures than by monocultures in these three years. However, only 4 of the mixtures (14.3%) yielded more than the mid component yield in all three years. None yielded consistently less.

Table 14. Frequency distribution of mixture yield relative to the yield of the components grown in monoculture.

All plots were machine seeded, and harvested at maturity, in 1977, 1978 and 1979.

Mixture	Frequency distribution of mixture yield ¹			
	$Y_M < C_1$	$C_1 < Y_M < C_2$	$C_2 < Y_M < C_3$	$Y_M > C_3$
Pitic 62 + Glenlea				2
Pitic 62 + Park			1	1
Pitic 62 + Neepawa			1	1
Pitic 62 + 70M110001				2
Pitic 62 + 70MD09002				2
Pitic 62 + Norquay	1			1
Pitic 62 + NB 701				2
Glenlea + Park			1	2
Glenlea + Neepawa			2	1
Glenlea + 70M110001		2		1
Glenlea + 70MD09002		1		2
Glenlea + Norquay			2	1
Glenlea + NB 701	2			1
Park + Neepawa	1		1	1
Park + 70M110001	1	1	1	
Park + 70MD09002	1		2	
Park + Norquay		1	1	1
Park + NB 701	1		1	1
Neepawa + 70M110001	1	1	1	
Neepawa + 70MD09002		1	2	
Neepawa + Norquay	1	1		1
Neepawa + NB 701		1		2
70M110001 + 70MD09002	1	1	1	
70M110001 + Norquay		2		1
70M110001 + NB 701	1		2	
70MD09002 + Norquay		2		1
70MD09002 + NB 701		2	1	
Norquay + NB 701			1	2
	11	16	21	29
Percent of total	(14.3)	(20.8)	(27.3)	(37.7)

1. Number of seasons in which the mixture yield (Y_M) fell in the specified categories; where

C_1 = Lower yielding component

C_2 = Mid-component yield

C_3 = Higher yielding component

2. Data for Pitic 62 were only obtained in two years.

8.3.1 Effects of genotype interaction on plot yields

An analysis of variance of the combined data of the three test years showed that year effects as well as effects among mixtures and monocultures were significant ($\alpha \leq 0.01$) (Table 15). Partitioning of the treatment SS into components containing variation due to differences among monocultures, due to differences among mixtures and due to differences between mixtures and monocultures, revealed that mixture and monoculture yields did not differ significantly when observed for a period of three years. Significant interaction effects existed between years and mixtures and between years and monocultures, suggesting a year effect on the performance of individual genotypes when compared in mixture and in monoculture.

Analysis of each year individually (Table 16) revealed the absence of significant differences between monocultures in 1978. This fact is probably largely responsible for the year effect reported in Table 16. The wet fall weather in 1978 caused considerable lodging in one corner of the field, which resulted in a significant ($\alpha \leq 0.01$) replication effect, and probably obscured true genotype effects. The differences among mixture yields were significant in each of the three years ($\alpha \leq 0.05$, $\alpha \leq 0.01$, $\alpha \leq 0.01$). Only in 1977 did mixture and monoculture yields differ significantly ($\alpha \leq 0.05$).

Table 15. The effect of year, genotype monoculture and genotype mixing on plot yield, as determined by analysis of variance.¹

All plots were machine seeded, and harvested at maturity, in 1977, 1978 and 1979.

Source of variation	df	Mean square
Years (Y)	2	2913881**
Treatments (T)	27 ²	26432**
Genotype		
Monocultures (Mono)	6	51901**
Mixtures (Mix)	20	18963*
Mix vs Mono	1	22990
Y × T	54	33183**
Y × Mono	12	45503**
Y × Mix	40	30069**
Y × Mix vs Mono	2	21547
Residual	168	11487

1. * significant, $\alpha \leq 0.05$; ** significant, $\alpha \leq 0.01$.

2. Data for Pitic 62 have been excluded for all years.

Table 16. The effect of genotype monoculture and genotype mixing on plot yield in each of three test years, as determined by analyses of variance.¹

All plots were machine seeded, and harvested at maturity, in 1977, 1978 and 1979.

Year	Source of variation	df	Mean square
1977	Treatments:		
	Genotype		
	Monocultures (Mono)	6	68725**
	Mixtures (Mix)	20	21543*
	Mono vs Mix	1	60896*
	Replications	2	22021
	Residual	53	10846
1978	Treatments:		
	Genotype		
	Monocultures (Mono)	7	22197
	Mixtures (Mix)	27	29860**
	Mono vs Mix	1	6513
	Replications	2	138430**
	Residual	107	12336
1979	Treatments:		
	Genotype		
	Monocultures (Mono)	7	48808**
	Mixtures (Mix)	27	43720**
	Mono vs Mix	1	14597
	Replications	2	3242
	Residual	107	8682

1. * significant, $\alpha \leq 0.05$, ** significant, $\alpha \leq 0.01$.

8.3.2 Mixture efficiency and its correlation with yield

Both the mean mixture yields and SME values differed from year to year. For example, the Park/Neepawa mixture yielded well above the expected yield in 1977, while in 1979, the yield of the same combination failed to reach even the level of the lower yielding component (Appendix 18a through c).

Spearman's rank-order correlation coefficients were computed for the following correlations (Table 17):

1. mixture yield and SME in 1977, in 1978 and in 1979,
2. mixture yields in 1977 and 1978, in 1977 and 1979, and in 1978 and 1979, and
3. SME's in 1977 and 1978, in 1977 and 1979, and in 1978 and 1979.

No consistent relationship seemed to exist between the mixture yields and the SME values in any of the three years. The relative mixture yields in 1978 appeared to be quite different from those in 1977 and in 1979, and either climatic conditions or the different location could have influenced relative yields.

Attinaw (1977) grew seven of the eight genotypes which were involved in this study, at three different locations, including Edmonton and Ellerslie. Table 18 shows the rank order for yield, in each year, for the genotypes grown in his study and in this study at the two locations. Some genotypes appeared quite consistent in their relative position (e.g. Glenlea and Park), while others were more

Table 17. Relationship between mixture yield and Specific Mixture Efficiency¹, within and between years, as expressed by Spearmans rank order correlation coefficient².

Correlated Parameters ³	Correlation coefficient
$Y_{77} - Y_{78}$	-0.37*
$Y_{77} - Y_{79}$	0.58**
$Y_{78} - Y_{79}$	-0.29
$Y_{77} - \text{SME}_{77}$	0.19
$Y_{78} - \text{SME}_{78}$	0.26
$Y_{79} - \text{SME}_{79}$	0.74**
$\text{SME}_{77} - \text{SME}_{78}$	-0.21
$\text{SME}_{77} - \text{SME}_{79}$	0.29
$\text{SME}_{78} - \text{SME}_{79}$	0.06

1. As defined in Section 8.2.

2. * significant, $\alpha \leq 0.05$; ** significant, $\alpha \leq 0.01$.

3. Y_{77} , Y_{78} , Y_{79} = mixture yield in 1977, 1978 and 1979, respectively.

SME_{77} , SME_{78} , SME_{79} = Specific Mixture Efficiency in 1977, 1978 and 1979, respectively.

Table 18. Rank orders of the monoculture yields of eight genotypes, and the mean yields of their mixtures, as observed at the Edmonton and Ellerslie locations.

Genotype	Rank orders									
	Monocultures					Mixtures				
	1975 ¹ Edmonton	1977 ¹ Edmonton	1979 ² Edmonton	1975 ¹ Ellerslie	1978 ² Ellerslie	1977 ² Edmonton	1978 ² Ellerslie	1979 ² Edmonton		
Pitic 62	1	--	1	1	5	--	4	1		
Glenlea	4	4	3	3	4	5	2	3/4		
Park	7	7	8	7	7	6	7	7		
Neepawa	5	6	6	5	1	7	1	8		
70M110001	2	3	5	2	2	3	3	5		
70M09002	3	1	4	6	8	1	8	2		
Norquay	6	2	7	4	3	2	5	6		
NB 701	--	5	2	--	6	4	6	3/4		

1. Data from Attinaw (1977)

2. Data from present study

3. Genotype not represented in test of that year

unstable (e.g. 70M009002). In 1978, however, the genotypes Neepawa and Pitic 62 showed quite a different relative yield level than in any of the other years. This erratic behaviour also determined the performance of the mixtures containing these two genotypes, as illustrated by the ranking of the array means. The variable yields obtained from both monocultures and mixtures over the three years thus appear to relate to factors other than the change in location.

8.3.3 Diallel analysis of variance

As was discussed in Section 8.2, analysis of the data as a mixture diallel allows a more detailed interpretation of the results.

Array means differed significantly in each of the three years ($\alpha \leq 0.01$) (Table 19). In the absence of interaction effects between the genotypes, this is a reflection of genotype differences in yielding ability. In the previous section it was shown, however, that in 1978, the differences among genotypes in monoculture were insignificant. A significant array effect in this year, thus, suggests the occurrence of a significant interaction between genotypes in mixture in that year.

The b1 effect, which contrasts mixture and monoculture yields, was significant in 1977 only ($\alpha \leq 0.05$), indicating that interaction between genotypes was present in that year also. Only in 1979 was the b2 effect significant ($\alpha \leq 0.01$), indicating that the effects of genotype interaction on yield

Table 1: The average array and specific effects of genotype on mixture yield, as determined by diallel analyses of variance¹ for each of the three test years.

All plots were machine seeded, and harvested at maturity.

Year	Source of variation ²	df	Mean square
1977	Replications	2	22021
	Arrays	6	100804**
	Mixtures vs monocultures	1	60896*
	Average genotype interaction	6	7519
	Specific genotype interaction	14	13805
	Residual	53	10846
1978	Replications	2	138430**
	Arrays	7	176335**
	Mixtures vs monocultures	1	6513
	Average genotype interactions	7	6901
	Specific genotype interaction	20	18877
	Residual	70	12336
1979	Replications	2	3242
	Arrays	7	168180**
	Mixtures vs monocultures	1	14597
	Average genotype interaction	7	27834**
	Specific genotype interaction	20	7501
	Residual	70	8682

1. * significant, $\alpha \leq 0.05$; ** significant, $\alpha \leq 0.01$.

2. Sources of variation were partitioned according to the technique described by Hayman (1954); see Section 8.2.

differed among genotypes.

Hay (1974) explained that, when the additions made by the interactions to the array values (mixtures) are correlated with the diagonal values (monocultures), '...the differences between the arrays will be inflated and the array mean square will be more likely to reveal interactions than the b2 mean square.' It can be noted, though, that this is only the case if the correlation is positive. In the present study, the correlation between monoculture yield and Specific Mixture Efficiency was insignificant in two of the three years, and significantly positive ($\alpha \leq 0.01$) in the third year (Table 17). Simmonds (1962), however, investigated the relationship between the difference between mixture yield and mean monoculture yield, and the mean monoculture yield of the components, and reported a negative correlation. Thus, a positive correlation cannot be considered to be a general rule.

The b3 effect, which measures variation due to specific mixtures, was not significant in any of the three years, suggesting that no specific combination of genotypes performed exceptionally well or exceptionally poorly in any of the three years.

8.3.4 Mixture efficiency

The AME values for each of the genotypes in each of the three years show the ability of each genotype to perform in a mixture situation (Appendix 19). The values for GCA, which

merely reflect the relative yield level of the mixtures in the test, were computed for comparison. Slight discrepancies between the rank order for array mean yield and for GCA are due to the fact that monoculture yields are used twice in the computation of GCA and once in the computation of the array mean yield.

Great variation existed from year to year among AME values of each genotype. In 1977, all genotypes had AME values greater than 1, suggesting a contribution to the mixture yield which is greater than expected from monoculture values. In 1978, however, less than half the genotypes seemed to perform better than expected in mixture.

Correlation coefficients of AME with monoculture yield for the three years were insignificant (Table 20), and it can be concluded that no consistent relationship existed between monoculture yield and the ability to perform in a mixture.

8.3.5 Regression of W on V


The regression of the covariances (W) of the array values and monoculture values (diagonal values of the diallel table) on the array variances (V), provides information on the relative ability of the genotypes to dominate in mixture (Figure 3). Because the genotype Pitic 62 was eliminated from the 1977 tests, the values for that year are not directly comparable with the other years.

Table 20. Simple correlations (r) between monoculture yield and Average Mixture Efficiency¹, in 1977, 1978 and 1979.

Year ²	r
1977	-0.48
1978	-0.05
1979	0.57

1. See Section 8.2

2. Seven genotypes were used in 1977, eight in each of the years 1978 and 1979.



- 1 Pitic 62
- 2 Glenlea
- 3 Park
- 4 Neepawa
- 5 70M110001
- 6 70M009002
- 7 Norquay
- 8 NB 701





Fig. 3a-c. V-W regression lines, illustrating the relative ability of genotypes to dominate in a mixture, in the years 1977, 1978 and 1979. V = variance of the array (mixture) values, W = covariance of the array values and the diagonal (monoculture) values.

As was already shown by the previous analyses, in 1978 the behaviour of the mixtures was erratic. Although the slopes of the regression lines in 1977 and 1979 were both close to one, the ranking of the genotypes according to their behaviour in mixture was quite different in each of the years. Glenlea, 70M009002, Neepawa and Park generally maintained their positions, while 70M110001, Norquay and NB701 showed quite different behaviours in mixture in the two years.

The V-W regression line shows along the x-axis the genotypes in order of their dominance in the mixtures. Correlation of the rank-order of dominance in mixture of the genotypes with their monoculture yields, resulted in a Spearman's rank-order correlation coefficient of 0.25 (n.s.) for 1977 and of 0.99 (sign $\alpha \leq 0.001$) for 1979 (Figure 3).

It can thus be concluded that the ability to dominate in a mixture was highly variable from year to year and not consistently related to yield in monoculture.

8.4 Discussion

It has been suggested that the ability to perform well in a mixture is negatively correlated to yield in monoculture (Wiebe et al., 1961; Simmonds, 1962). The present study, however, did not support this theory. Significant differences existed in every year in the relative ability of the genotypes to dominate in mixture. No

relationship was found between the ability to dominate in mixture and monoculture yield in the same year.

Baker (1977) reported variable results for Neepawa, Pitic 62, and a mixture of the two, and ascribed this to the fact that Pitic 62 was better able to compensate for poor survival after early drought conditions, when grown in pure culture than when grown in combination with Neepawa.

DePauw et al. (1981) included the same genotypes in a study comparing yield at five locations over four years. They also found great variation in the performance of the genotype Pitic 62. They ascribed this to frost damage in the post-anthesis stage. In the present study, however, neither post-anthesis frost nor early drought occurred in any of the three years.

It is probable that the different responses of the genotypes to the variable climatic conditions were responsible for the year to year variation. Identification of genotype-specific responses to the environment, however, would require a qualitative study of genotype-environment interactions.

The results of this experiment do support the findings of those investigators who concluded that mixtures, on the average, yield more than the expected mean of the component monocultures (Table 14). This suggests that a positive interaction can occur between the components of a mixture, resulting in a more efficient use of available resources by one or both of the mixture components. Variation within and

between years, however, was so large that the yield advantage of mixtures over monocultures was statistically insignificant (Table 15). No specific binary mixture was identified which consistently gave a high yield advantage.

9. Discussion of the Results with Respect to Single Plant Selection in Early Generations

Inefficiency of single plant selection in early generations can be ascribed to two major sources of error:

1. The amount of additive genetic variation present in the population is low, which makes it difficult to identify those genotypes which harbour the superior genetic material one wishes to select.
2. Effects of genotype x environment interaction on plant growth can cause undesirable phenotypic variation.

The relative magnitude of these factors is expressed by the heritability (H^2) of a character, which is computed as follows:

$$H^2 = \text{Genetic variance} / \text{Total variance}$$

Estimates of heritability can be obtained in a variety of ways, and the outcome will vary with the experimental design used and the method of computation employed (Allard, 1960; Falconer, 1960). Experiments with a large number of replications will reduce the environmental variance and, thus, give higher estimates for H^2 . However, single plant selections from early generations have to be made from unreplicated plots, and the improvement of the selected population over the parental population depends entirely on how accurately superior genotypes can be identified within those plots.

9.1 Effects of non-normal character distribution on single plant selection

The expected gain from selection can be expressed as

where k is the selection differential, which takes into account the mean and the variance of the parental population, the mean of the selected fraction, and the selection intensity; σ is the phenotypic standard deviation; and H^2 is the heritability of the observed character (Allard, 1960). This expression assumes a normal distribution of the units to be selected, whether they are mean values of characters of plants grown in rows or observations on single plants. Reports on non-normal distributions of several characters of various plant species were given by Koyama and Kira (1956) and by Harper (1977).

The present study showed that the genotypes grown in binary mixtures with a range of associated genotypes at the 3 cm spacing, showed significant deviations from normality for all characters, except the number of kernels per head and height of the flag leaf (Appendix 8). At the 9 cm spacing, deviations from normality occurred for the number of tillers, the number of heads, kernels per plant, 1000 kernel weight and harvest index.

Deviations from normality will affect the accuracy with which genotypes can be distinguished, and will lead to gains from selection which are different than those expected based on the above formula.

The effect which a positively skewed distribution would have on single plant selection is graphically illustrated in Figure 4a. If genotypes with a high value of a certain character are to be selected, a positive skewness would increase the probability for less desirable genotypes to be present in the selected fraction. Similarly, a negative skewness (or selection of plants with the lowest values for a certain character from a population with a positively skewed distribution) would result in a selected fraction with more desirable genotypes than would have been the case if the character had been normally distributed.

To quantify this effect, the distribution of chi-square, which is a typical skewed distribution, can be conveniently used (Appendix 20). It has been shown, that the effect of skewness on single plant selection became much larger when the frequency of superior genotypes in the population was low. (Dr. D. H. Kelker, Associate Professor of Statistics, University of Alberta, Edmonton, Alberta, 1981, Personal communication). A positive skewness would, thus, result in gains from selection which are much lower than expected, based on estimates of heritability. Similarly, a negative skewness would result in a higher than expected gain from selection, if the highest values from the population were selected.

Considering these facts, Appendix 8 leads to the conclusion that response to selection at the 3 cm spacing would be less than expected for the number of tillers, the




Fig. 4.

The effect of skewness and kurtosis of the distribution of a plant character on single plant selection. $G_1 - G_4$ represent genotypes in a mixture, with means at equal distances apart. a_1 , a_2 and a_3 are the ranges of genotypes selected in populations with a normal distribution, a positive skewness, and a positive kurtosis, respectively. $a_3 < a_1 < a_2$.

number of heads, weight, yield and kernels per plant, and greater than expected for 1000 kernel weight, harvest index, height and extrusion length, provided that the tallest individuals were selected.

As was mentioned earlier, little can be said about the effect of a significant kurtosis in the presence of skewness. In the absence of skewness, however, a positive kurtosis is likely to have an adverse effect on selection (Figure 4b), because the selected fraction would be contaminated with less desirable genotypes to a greater extent than would be the case if the character had been normally distributed. Similarly, better results can be expected with a negative kurtosis.

At the 9 cm spacing, selection for a large number of tillers or a large number of kernels per head could result in smaller than expected gains, while selection of plants with a high harvest index or large kernels could result in a higher than expected gain.

9.2 Effects of genotype interaction on single plant selection

The results from the experiment conducted in 1978 suggest that within-genotype variation of several characters would be significantly increased by genotype interaction if the plants were grown in a multi-component mixture. At the 3 cm spacing, the number of tillers, weight and yield

were affected by genotype interaction, while at the 9 cm spacing, 1000 kernel weight showed an increase in variation (Table 4).

The effect of genotype interaction on estimates of heritability is shown in Tables 21a and 21b for the data from the experiment grown in 1978. Analyses of variance for each interplant spacing, using a random model (Steel and Torrie, 1980), provided the variance components which were used to estimate the values of H^2 (Briggs et al, 1978). Differences between the two estimates suggest that genotype interaction can lead to higher estimates of heritability, in particular for the characters weight, yield and kernels per plant. This observation supports the idea that H^2 is not always a reliable predictor for the ease with which a character can be improved through selection.

9.3 Effects of interplant spacing on single plant selection

The within-genotype variation for several characters of plants grown in monoculture increased significantly with an increase in spacing (Table 4). The mean variation of the four genotypes, when grown in mixture, increased significantly with the increase in spacing for all characters, except kernels per head and components of height.

This suggests that identification of superior genotypes would be more successful at the 3 cm than at the 9 cm

Table 21a. Estimates of heritability of single plant characters, grown at a 3 cm interplant spacing, as computed from monoculture and mixture data of four genotypes.

All plants were grown in four-row plots in 1978. The data are from randomly selected plants, which had both neighbours present, and which were harvested at maturity.

Character ²	Heritability	
	Computed from monoculture data	Computed from mixture data
T	0.57	0.57
H	0.66	0.70
Wt	0.16	0.50
Y	0.35	0.63
K/P	0.04	0.35
Kwt	0.91	0.94
K/H	0.70	0.83
HI	0.63	0.63
FL	0.88	0.89
Ht	0.83	0.83
ExL	0.01	0.48
HL	0.87	0.90

1. Heritability = $\frac{\text{Genetic variation}}{\text{Total variation}}$

2. Character abbreviations defined in Table 1. HL = Head length.

Table 21b. Estimates of heritability of single plant characters, grown at a 9 cm interplant spacing as computed from monoculture and mixture data of four genotypes.

All plants were grown in four row plots in 1978. The data are from randomly selected plants, which had both neighbours present, and which were harvested at maturity.

Character ²	Heritability	
	Computed from monoculture data	Computed from mixture data
T	0.0	0.0
H	0.0	0.32
Wt	0.48	0.67
Y	0.50	0.71
K/P	0.19	0.45
Kwt	0.97	0.97
K/H	0.71	0.77
HI	0.71	0.71
FL	0.91	0.91
Ht	0.74	0.74
ExL	0.78	0.87
HL	0.87	0.94

1. Heritability = $\frac{\text{Genetic variation}}{\text{Total variation}}$

2. Character abbreviations defined in Table 1. HL = Head length.

spacing, because the genetic variance at the 3 cm spacing and at the the 9 cm spacing is approximately the same, differing only by an amount representing genotype x spacing interaction.

Other studies led to different conclusions, however. In an earlier study concerning the same genotypes as were used in the present experiment, a decrease rather than an increase in variation with an increase in spacing was observed for most characters (Kelker and Briggs, 1978). The different results may be partially explained by the fact that in the present study, the difference between the two seeding rates was much larger than in the earlier study. In addition, the distance between rows was larger in the present experiment than in the previous study, resulting in a different spatial arrangement of the plants, which would cause different neighbour interactions (Harper, 1961; Donald, 1963). It is unlikely, though, that these factors can account for a complete reversal of the effect of an increase in spacing on plant-to-plant variation.

Nass (1980) observed that, while biological yield and grain yield of plants from two spring wheat crosses responded each in a different manner to a ten-fold increase in plant spacing, harvest index of plants from both crosses was less variable at a higher (commercial) plant density. Nass recommended growing plants at commercial crop densities, if plants with high harvest index were to be selected. The commercial seeding rate for the humid Prince

Edward Island environment, however, is more than double the rate used in Alberta.

The fact that different seeding rates as well as different planting patterns were used in the various studies, makes it difficult to compare their results. It can be concluded, though, that an increase in plant spacing does not always result in an increase in plant-to-plant variation. The contrasting results from the two Alberta studies, concerning the same genotypes, suggest that location effects and year effects may play an important role.

The present study showed that mean values of most characters were significantly affected by spacing effects as well as by genotype x spacing interaction effects (Table 5). This could lead to different results from selection at different spacings (CIMMYT, 1972; Nass, 1980). For this reason, as well as to avoid a possible increase in variation, selection should be practiced at a density which approximates the commercial seeding rate of the region to which the selected genotypes have to be adapted.

9.4 Effects of seed size on single plant selection

The experiments performed to evaluate the effect of variation in initial seed size on plant-to-plant variation, suggested that variation in seed size can be an additional source of error in the identification of superior genotypes.

The number of heads, the number of tillers and 1000 kernel weight appeared to vary with initial seed size, while harvest index and 1000 kernel weight were affected by variations in seed size among neighbouring plants. Thus, the use of uniform seed would reduce the plant-to-plant variation observed in the selection plots.

This has earlier been suggested by Christian and Gray (1941) and by Chebib et al. (1973). However, the sorting of seed should aim for seed of medium size, to avoid selection for large kernelweight. The latter could be an undesirable side effect, because the number of kernels per head and 1000 kernel weight are negatively correlated (Fischer, 1975 ;Jenner, 1979). Reports of studies investigating the effects of selection for increased kernel size on yield are contradictory (Bhatt and Derera, 1973; Rasmusson and Cannell, 1970), while the percent protein in the kernels tends to decline with an increase in kernel size (Röbbelen, 1979). The merits of seeding uniform seed would, therefore, have to be evaluated for each selection program in light of its breeding objectives.

9.5 Suitability of various characters for single plant selection

The formula for the calculation of heritability shows that the highest values will be obtained for characters which:

- a show a large amount of additive genetic variation, and
 - b are little affected by random environmental variation.
- The previous sections of this chapter explained that, in addition, the best chances for genetic improvement through selection exist for those characters which, in a segregating population grown at a commercial crop density:
- c are little affected by variations in initial seed size,
 - d are little affected by neighbour interactions arising from variations in seed size,
 - e are little affected by genotype interactions, and
 - f have a distribution which is negatively skewed (if the lowest values are to be selected the skewness should be positive).

For most characters, there are one or more factors by which selection is hindered. Estimates of heritability would encompass the factors mentioned in the points a, b, c and d. To what extent H^2 is biased through genotype interaction effects would depend on the method of calculation (Hamblin and Rosielle, 1978). The effects of non-normality are not accounted for.

Because only homozygous genotypes were used in this study, there is only additive genetic variation, and a hypothetical mixture of these genotypes would be unrepresentative of a segregating population. But the characters measured in this study can be evaluated with respect to the remaining points which determine the efficiency of selection. It then appears that the characters

1000 kernel weight and harvest index did fullfill these requirements best (Tables 4, 5, 12d), and gains from selection could be higher than suggested by an estimate of heritability. Based on the data from this study, the number of tillers, the number of heads and yield can be expected to respond poorly to single plant selection, resulting in less genetic improvement than expected. Of the remaining characters, none had a distribution which would interfere with predictions of gain from selection. Genotype interaction, however, biased estimates of heritability for plant weight, yield and kernels per plant.

It was explained that kernel weight is not always a reliable indicator of yield and may affect percent protein of the selected genotypes. Selection for high harvest index, as an indirect method for selection for yield, however, has shown promising results (Donald and Hamblin, 1976; Fischer and Kertesz, 1976; Bhatt, 1977). The present study gives support to the observation that selection for high harvest index can result in high gains from selection (Bhatt, 1977).

The results from the present study suggest, that, in order to evaluate the suitability of a character for selection in early generations, the distribution of this character in a multi-component mixture, as well as its sensitivity to genotype interaction, should be evaluated at a commercial plant density, in conjunction with the estimation of its heritability. The combined information

could then lead to a more realistic expectation of the progress which can be made through selection.

10. References

- Aitken, Y. (1974). Flowering time, climate and genotype. Melbourne Univ. Press, Carlton, Victoria.
- Alberta Farm Guide. (1976). Alberta Agriculture, Edmonton.
- Allard, R. W. (1960). Principles of plant breeding. Wiley and Sons, Inc., New York.
- Allard, R. W. (1961). Relationship between genetic diversity and consistency of performance in different environments. *Crop Sci.* 1:127-33.
- Aspinall, D. (1960). An analysis of competition between barley and white persicaria. II. Factors determining the course of competition. *Ann. appl. Biol.* 48:637-54.
- Aspinall, D. and F. L. Milthorpe. (1959). An analysis of competition between barley and white persicaria. I. The effects on growth. *Ann. appl. Biol.* 47:156-72.
- Attinaw, A. (1977). Effects of seeding rates and dates on wheat. M.Sc. Thesis, The Univ. of Alberta, Edmonton, Alberta.
- Austenson, H. M. and P. D. Walton. (1970). Relationships

between initial seed weight and mature plant characters in spring wheat. *Can. J. Plant Sci.* 50:53-8.

Baker, R. J. (1977). Yield of pure and mixed stands of two spring wheat cultivars sown at five rates of seeding. *Can. J. Plant Sci.* 57:1005-7.

Bhatt, G. M. (1977). Response to two-way selection for harvest index in two wheat crosses. *Aust. J. agric. Res.* 28:29-36.

Bhatt, G. M. and N. F. Derera. (1973). Associated changes in the attributes of wheat populations mass-selected for seed size. *Aust. J. agric. Res.* 24:179-86.

Black, J. N. (1956). The influence of seed size and depth of sowing on pre-emergence and early vegetative growth of subterranean clover. *Aust. J. agric. Res.* 7:98-109.

Black, J. N. (1958). Competition between plants of different initial seed sizes in swards of subterranean clover, with particular reference to leaf area and the light microclimate. *Aust. J. agric. Res.* 9:299-318.

Bradshaw, A. D. (1964). Evolutionary significance of phenotypic plasticity in plants. *Adv. Genet.* 13:115-55.

Briggs, K. G., D. G. Paris and H. A. Kelker. (1978).

Effectiveness of selection for plant characters of barley in simulated segregating rows. *Euphytica* 27:157-66.

Brougham, R. W. (1958). Leaf development in swards of white clover (*Trifolium repens*). *New Zeal. J. of agric. Res.* 1:707-18.

Busch, R. H. and D. Luizzi. (1979). Effects of intergenotypic competition on plant height, days to heading and grain yield of F2 through F5 bulks of spring wheat. *Crop Sci.* 19:815-9.

Chapman, S. R., R. W. Allard and J. Adams. (1969). Effect of planting rate and genotypic frequency on yield and seed size in mixtures of two wheat varieties. *Crop Sci.* 9:575-576.

Chebib, F. S., S. B. Helgason and P. J. Kaltsikes. (1973). Effect of variation in plant spacing, seed size and genotype on plant-to-plant variability in wheat. *Z. Pflanzenzuchtg.* 69:301-32.

Christian, C. S. and S. G. Gray. (1941). Interplant competition in mixed wheat populations and its relation to single plant selection. *J. Council Sci. Ind. Res.* 14:59-68.

CIMMYT (1972). Report on progress toward increasing yields of maize and wheat. pp. 1967-68. Intern. Maize and Wheat Improvement Center, El Batan, Mexico.

Clark, E. A. (1980). Physiological basis for transgressive yielding in bean-maize intercrops. Ph.D. Thesis, Iowa State Univ., Ames, Iowa.

Clay, R. E. and R. W. Allard. (1969). A comparison of the performance of homogeneous and heterogeneous barley populations. *Crop Sci.* 9:407-12.

Clements, F. E., J. E. Weaver and H. C. Hanson. (1929). Competition in cultivated crops. Carnegie Inst. Wash.. Publ. No. 398.

Darwin, C. (1859). On the origin of species. A facsimile of the first edition. Harvard Univ. Press, Cambridge, Mass.

DePauw, R.M., D. G. Faris and C. J. Williams. (1981). Genotype-environment interaction of yield in cereal crops in northwestern Canada. *Can. J. Plant Sci.* 61:255-63.

Derera, N. F. and G. M. Bhatt. (1972). Effectiveness of mechanical mass selection in wheat. *Aust. J. agric. Res.* 23:761-8.

- Donald, C. M. (1958). The interaction of competition for light and for nutrients. *Aust. J. agric. Sci.* 9:421-35.
- Donald, C. M. (1961). Competition for light in crops and pastures. In *Mechanisms in biological competition*. F. L. Milthorpe (ed.). S.E.B. Symp. XV, pp. 287-313. Cambridge Univ. Press.
- Donald, C. M. (1963). Competition among crop and pasture plants. *Adv. Agron.* 15:1-118.
- Donald, C. M. (1967). The breeding of crop ideotypes. *Euphytica* 17:385-403.
- Donald, C. M. and J. Hamblin. (1976). The biological yield and harvest index of cereals as agronomic and plant breeding criteria. *Adv. Agron.* 28:361-405.
- Early, H. L. and C. O. Qualset. (1971). Complementary competition in cultivated barley. *Euphytica* 20:400-409.
- Eberhart, S. A., L. H. Penny and G. F. Sprague. (1964) Intraplot competition among maize single crosses. *Crop Sci.* 4:467-71.
- Edwards, K. J. R. and R. W. Allard. (1963). The influence of light intensity on competitive ability. *Amer. Nat.*

97:243-248.

Engelke, H. (1935). Versuche mit Weizenmischsaaten. *J. Landw.* 83:63-86.

Evans, L. T. and I. F. Wardlaw. (1976). Aspects of the comparative physiology of grain yield in cereals. *Adv. Agron.* 24:301-59.

Falconer, D.S. (1960). Introduction to quantitative genetics. Ronald Press Co., New York.

Feldman, M. (1974). Wheats. In Evolution of crop plants. pp.120-28. N. W. Simmonds (ed.). Longman, New York.

Fischer, R. A. (1975). Yield potential in a dwarf spring wheat and the effect of shading. *Crop Sci* 15:607-13.

Fischer, R. A. and Z. Kertesz. (1976). Harvest index in spaced populations and grain weight in microplots as indicators of yielding ability in spring wheat. *Crop. Sci.* 16:55-9.

Ford, M. A. and G. N. Thorne. (1975). Effect of variation in temperature and light intensity at different times on growth and yield of spring wheat. *Ann. appl. Biol.* 80:283-99.

- Frankel, O. H. (1939). Analytical yield investigations on New Zealand wheat. *J. agric. Sci.* 29:249-61.
- Griffing, B. (1956). Concept of general and specific combining ability in relation to diallel crossing systems. *Aust. J. biol. Sci.* 9:463-93.
- Hall, R. L. (1974). Analysis of the nature of interference between plants of different species. I. Concepts and extension of the de Wit analysis to examine effects. *Aust. J. agric. Res.* 25:739-47.
- Hamblin, J. and C. M. Donald. (1974). The relationship between plant form, competitive ability and grain yield in a barley cross. *Euphytica* 23:535-42.
- Hamblin, J. and A. A. Rosielle. (1978). Effect of interaction between genotypes on genetic parameter estimation. *Crop Sci.* 18:51-54.
- Harper, J. L. (1961). Approaches to the study of plant competition. In *Mechanisms in biological competition*. F. L. Milthorpe (ed.). S.E.B. Symp. XV, pp. 1-39.
- Harper, J. L. (1965). The nature and consequences of interference amongst plants. In *Genetics today*. Proc. 11th Intern. Cong. Genet., The Hague, pp. 465-82.

Pergamon Press, Oxford.

Harper, J. L. (1977). Population biology of plants. Acad. Press, San Francisco.

Harper, J. L. and J. N. Clatworthy. (1963). The comparative biology of closely related species. VI. Analysis of the growth of *Trifolium repens* and *T. fragiferum* in pure and mixed populations. *J. exp. Bot.* 14:172-90.

Hartmann, R. W. and R. W. Allard. (1964). Effect of nutrient and moisture levels on competitive ability in barley. *Crop Sci.* 4:424-6.

Hay, D. A. (1974). A quantitative method for analysing competition where individuals cannot be identified. *Aust. J. biol. Sci.* 27:133-196.

Hayman, B. I. (1954). The analysis of variance of diallel tables. *Biometrics* 10:235-44.

Henzell, F. L. and I. Vallis. (1977). Transfer of nitrogen between legumes and other crops. In Biological nitrogen fixation in farming systems of the tropics. A. Ayanaba and P. J. Dart (eds). Wiley and Sons. New York.

Heuser, W. (1938). Versuche mit Sortenmischungen von

Winterweizen. *Mitt. Landw.* 8:5-6.

Hill, J. (1974). Methods of analysing competition with special reference to herbage plants. III. Monocultures vs binary mixtures. *J. agric. Sci., Camb.* 83:57-65.

Hill, J. and Y. Shimamoto. (1973). Methods of analysing competition with special reference to herbage plants. I. Establishment. *J. agric. Sci., Camb.* 81:77-89.

Hozumi, K., H. Koyama and T. Kira. (1955). Intraspecific competition among higher plants. IV. A preliminary account on the interaction between adjacent individuals. *J. Inst. Polytechn. Osaka City Univ.*, D6:121-30.

Hsu, P. and P. D. Walton. (1971). Relationships between yield and its components and structures above the flag leaf node in spring wheat. *Crop Sci.* 11:190-193.

Jenner, C. F. (1979). Grain-filling in wheat plants shaded for brief periods after anthesis. *Aust. J. Plant Physiol.* 6:629-41.

Jensen, N. F. and W. T. Federer. (1965). Competing ability in wheat. *Crop Sci.* 5:449-52.

Jinks, J. L. (1954). The analysis of continuous variation in

a diallel cross of *Nicotiana rustica* varieties. *Genetics* 39:767-88.

Kaufmann, M. L. and A. D. McFadden. (1960). The competitive interaction between barley plants grown from large and small seeds. *Can. J. Pl. Sci.* 40:623-9.

Kelker, H. A. and K. G. Briggs. (1978). The effects of interaction between genotypes and interplant spacings in simulated segregating rows of wheat. *Proc. 5th Intern. Wheat Genet. Symp.*, New Delhi, pp. 778-86.

Kendall, M. and A. Stuart. (1977). The advanced theory of statistics. Vol. I. McMillan, New York.

Khalifa, M. A. and C. O. Qualset. (1975). Intergenotypic competition between tall and dwarf wheats. II. In hybrid bulks. *Crop Sci.* 15:640-4.

Kiesselbach, T. A. (1923). Competition as a source of error in comparative corn yields. *J. Amer. Soc. Agron.* 15:199-215.

Kiesselbach, T. A. (1924). Relation of seed size to the yield of small grain crops. *J. Amer. Soc. Agron.* 16:670-82.

Kira, T., H. Ogawa and N. Sakazaki. (1953). Intraspecific competition among higher plants. I.

Competition-yield-density interrelationship in regularly dispersed populations. *J. Inst. Polytechn. Osaka City Univ.*, D4:1-16.

Koyama, M. and T. Kira. (1956). Intraspecific competition among higher plants. VIII. Frequency distribution of individual plant weight as affected by the interaction between plants. *J. Inst. Polytech., Osaka City Univ.*, 7:73-94.

Lee, J. A. (1960). A study of plant competition in relation to development. *Evolution* 14:18-28.

Lewontin, R. C. (1966). On the measurements of relative variability. *Syst. Zool.* 14:141-2

Litav, M. and D. Isti. (1974a). Root competition between two strains of *Spinacia oleracea*. I. Pattern and processes. *J. appl. Ecol.* 11:1007-16.

Litav, M. and D. Isti. (1974b). Root competition between two strains of *Spinacia oleracea*. II. Effects of nutrient supply and of non-simultaneous emergence. *J. appl. Ecol.* 11:1017-25.

- Mather, K. (1961). Competition and cooperation. In
Mechanisms in biological competition. F. L. Milthorpe
(ed.). S.E.B. Symp. XV. pp. 164-81.
- McDaniel, R. G. (1969). Relationships of seed weight,
seedling vigour and mitochondrial metabolism in barley.
Crop Sci. 9:823-27.
- McGinnis, R. C. and L. H. Shebeski. (1968). The reliability
of single plant selection for yield in F₂. Proc. 3rd
Intern. Wheat Genet. Symp., Canberra. pp. 410-415.
- Milthorpe, F. L. (1961). The nature and analysis of
competition between plants of different species. In
Mechanisms in biological competition. F. L. Milthorpe
(ed.). S.E.B. Symp. XV, pp. 330-355.
- Montgomery, E. G. (1912). Competition in cereals. Nebraska
Agric. Exp. Stn. Bull. 127.
- Morley Jones, R. (1965). Analysis of variance of the half
diallel table. *Heredity* 20:117-21.
- Nass, H. G. (1973). Determination of characters for yield
selection in spring wheat. *Can. J. Plant Sci.* 53:755-62.
- Nass, H. G. (1980). Harvest index as a criterion for grain

yield in two spring wheat crosses grown at two population densities. *Can. J. Plant Sci.* 60:1141-6.

Nichiporovich, A. A. (1967). Aims of research on the photosynthesis of plants as a factor in productivity. In *Photosynthesis of productive systems*, pp. 3-36. A. A. Nichiporovich (ed.), Israel Prog. Sci. Trans., Jerusalem.

Nuding, J. (1936). Leistung und Ertragsstruktur von Winterweizensorten in Reinsaat und Mischung in verschiedenen deutschen Anbaugebieten. *Pflanzenbau* 12:382-447.

Odum, E. P. (1971). *Fundamentals of ecology*. Saunders, Philadelphia.

Phung, T. K. and A. J. Rathjen. (1976). Frequency-dependent advantage in wheat. *Theor. appl. Genet.* 48:289-97.

Phung, T. K. and A. J. Rathjen. (1977). Mechanisms of frequency-dependent advantage in wheat. *Aust. J. agric. Res.* 28:187-202.

Rasmusson, D. C. and R. Q. Cannell. (1970). Selection for grain yield and components of yield in barley. *Crop Sci.* 10:51-4.

Röbbelen, G. (1979). The challenge of breeding for improved protein crops. In Seed protein improvement in cereals and grain legumes. Vol. I, pp. 27-42. Intern. Atomic Energy Agency, Vienna.

Rosielle, A. A. and K. J. Frey. (1975a). Estimates of selection parameters associated with harvest index in oat lines derived from a bulk population. *Euphytica* 24:121-31.

Rosielle, A. A. and K. J. Frey. (1975b). Application of restricted selection for grain yield improvement in oats. *Crop Sci.* 15:544-7.

Ross, M. A. and J. L. Harper. (1972). Occupation of biological space during seedling establishment. *J. Ecol.* 60:77-88.

Roy, S. K. (1960). Interaction between rice varieties. *J. Genet.* 5:137-52.

Rümker, K. von (1892). Über Mengsaat verschiedener Weizensorten. *Fühlings Landw. Z.* 41:46-51, 92-7.

Sakai, K. I. (1955). Competition in plants in relation to selection. Cold Spring Harbour Symp. Quant. Biol. 20:137-57.

- Scheffé, H. (1959). The analysis of variance. Wiley Publications in Statistics, Shewhart and Wflks. (eds.), Wiley and Sons, New York.
- Shebeski, L. H. (1967). Wheat and breeding. Can. Centennial Wheat Symp. pp. 249-72. K. F. Nielsen (Ed). Modern Press, Saskatoon..
- Shorter, R. and K. J. Frey. (1979). Relative yields of mixtures and monocultures of oat genotypes. *Crop Sci.* 19:548-53.
- Shut, W. M., C. A. Brim and S. A. Usanis. (1968). Intergenotypic competition in plant populations. I. Feedback systems with stable equilibria in populations of auto~~ous~~mous homozygous lines. *Crop Sci.* 8:61-6.
- Simmonds, N. W. (1962). Variability in crop plants, its use and conservation. *Biol. Rev.* 37:442-65.
- Sneep, J., B. R. Murty and H. F. Utz. (1979). Current breeding methods. In Plant breeding perspectives. J. Sneep and A. J. T. Hendriksen (eds.), Pudoc, Wageningen.
- Sokal, R. R. and F. J. Rohlf. (1969). Biometry. Freeman and Co., San Francisco.

- Somerville, S. C. (1977). Yield and maturity estimates in cereals. M.Sc. Thesis, The Univ. of Alberta, Edmonton, Alberta.
- Sprague, G. F. and L. A. Tatum (1942). General vs. specific combining ability in single crosses of corn. *J. Amer. Soc. Agron.* 34:923-32.
- Steel, R. G. D. and J. H. Torrie. (1980). Principles and procedures of statistics. McGraw-Hill, Toronto.
- Suneson, C. A. (1949). Survival of four barley varieties in a mixture. *Agron. J.* 41:459-61.
- Syme, J. R. (1972). Single-plant characters as a measure of field plot performance of wheat cultivars. *Aust. J. agric. Res.* 23:753-60.
- Trenbath, B. R. (1974). Biomass productivity of mixtures. *Adv. Agron.* 26:177-210.
- Trenbath, B. R. (1975). Neighbour effects in the genus *Avena*. III. A diallel approach. *J. appl. Ecol.* 12:189-200.
- Trenbath, B. R. and J. L. Harper. (1973). Neighbour effects in the genus *Avena*. I. Comparison of crop species. *J. appl. Ecol.* 10:379-99.

- Troughton, A., W. J. Whittington. (1968). The significance of genetic variation in root systems. *In* Root growth, Proc. 45th Easter School in Agric. Sci., Univ. of Nottingham, pp. 296-313, W. J. Whittington (ed.).
- Wiebe, G. A., F. C. Petr and H. Stevens. (1961). Interplant competition between barley genotypes. *In* Statistical genetics and plant breeding, pp. 546-53. W. D. Hanson and H. F. Robinson (eds.). Publ. 982, Natl. Acad. Sci., Natl. Res. Council., Wash. D.C.
- Williams, E. J. (1962). The analysis of competition experiments. *Aust. J. biol. Sci.* 15:509-525.
- Wit, C. T. de (1960). On competition. *Versl. landbouwk. Onderz. Ned.* 66:1-82.
- Wit, C. T. de (1961). Space relationships within populations of one or more species. Mechanisms in biological competition. F. L. Milthorpe (ed.). S.E.B. Symp. XV, pp. 314-329.
- Yates, F. and W. G. Cochran. (1938). The analysis of groups of experiments. *J. agric. Sci.* 28:556-80.
- Yoda, K., T. Kira and K. Hozumi. (1957). Intra specific competition among higher plants. IX. Further analysis of

-
- ~ the competitive interaction between adjacent individuals. *J. Inst. Polytech.*, Osaka City Univ., D8:161-78.

11. List of Appendices

Appendix	Page
1. Mean monthly temperatures and precipitation near the test sites, for the growing seasons of 1977, 1978 and 1979, and the long term averages...	171
2. Details regarding fertiliser and herbicide application at the 1977, 1978 and 1979 locations.....	172
3. Agronomic characters of wheat genotypes in the study.....	173
4. Thousand kernel weight and percent germination of the seed used for the experiments in 1977, 1978 and 1979.....	175
5. Number of plants available, per treatment, from each density, for analysis of the wheelplot experiment grown in 1977.....	176
6. Descriptive statistics of characters of plants, grown in monoculture and in mixture at various interplant spacings, measured at maturity in 1977....	177
7. Descriptive statistics of characters of plants, grown in monoculture and in mixture at two	

interplant spacings, measured at maturity in 1978....	214
8. Distribution parameters of characters of single plants, grown in monoculture and in mixture, at two interplant spacings.....	229
9. Relative Yield Totals for several characters of plants grown in mixtures, at two interplant spacings in 1978.....	230
10. Descriptive statistics of characters of plants, grown in monoculture and in mixture, in machine seeded plots, measured at the flowering stage, in 1978	231
11. Descriptive statistics of characters of plants, grown in monoculture and in mixture, in machine seeded plots, measured at the middough stage, in 1978	235
12. Variation in thousand kernel weights among the seed classes used in 1979.....	239
13. Descriptive statistics of characters of plants from large, small and mixed size seeds,	

measured at the 3-5 leaf stage, in 1979.....	240
14. Descriptive statistics of characters of plants from large, small and mixed size seeds, measured at the jointing stage, in 1979.....	244
15. Descriptive statistics of characters of plants from large, small and mixed size seeds, measured at the heading stage, in 1979.....	247
16. Descriptive statistics of characters of plants from large, small and mixed size seeds, measured at maturity, in 1979.....	250
17. Distribution parameters of characters of single plants, measured at different stages of growth, derived from large seeds, small seeds, and alternated large and small seeds.....	258
18a. Yield and Specific Mixture Efficiency of mixtures grown in 1977.....	259
18b. Yield and Specific Mixture Efficiency of mixtures grown in 1978.....	260
18c. Yield and Specific Mixture Efficiency of mixtures grown in 1979.....	261

-
19. Performance of mixtures in each of the three years, containing specific genotypes, as expressed by the Array mean yield, the average Mixture Efficiency, and the General Combining Ability.....262
20. Calculation of the effect of skewness on single plant selection.....263

Appendix 1. Mean monthly temperatures and precipitation near the test sites, for the growing seasons of 1977, 1978 and 1979, and the long term averages.

A. Edmonton

Month	Temperature (°C)			Rainfall (mm)		
	1977	1979	Long term average ¹	1977	1979	Long term average
May	11.8	8.7	10.9	134	38	37
June	15.6	14.9	14.7	11	63	75
July	15.6	17.8	17.5	107	122	83
August	13.4	16.5	15.9	91	39	72
September	9.6	12.9	10.9	40	22	36

B. Ellerslie

Month	Temperature (°C)		Rainfall (mm)	
	1978	Long term average ²	1978	Long term average
May	10.5	10.5	59	47
June	16.0	14.0	38	84
July	16.5	16.0	114	85
August	14.5	15.0	94	68
September	12.0	10.0	34	36

1 Average of 99 years. Information obtained from the Annual Meteorological Summary for the Edmonton Municipal Airport, issued by Environment Canada, 1980.

2 Average of 17 years. Information obtained from the Department of Geography, Division of Meteorology, 1980.

Appendix 2. Details regarding fertiliser and herbicide application to the 1977, 1978 and 1979 locations.

Location	Fertiliser ¹ (kg/ha)			Herbicide		
	N	P	K	Trade name ²	Growth stage	Rate of application
Edmonton 1977	14	34	0	Buctril M	3-5 leaf	237 ml/ha
Ellerslie 1978	14	34	0	Avadex	N. A. ³	13.6 kg/ha ⁴
				MCPA-K	3-5 leaf	237 ml/ha
				Banvel	3-5 leaf	44 ml/ha
Edmonton 1979	5.5	14	0	Avadex	N.A.	14.2 kg/ha

1. Actual amounts N, P and K applied, derived from granular mono ammonium phosphate ($\text{NH}_4\text{K}_2\text{PO}_4$) in a broadcast application; applied two weeks before seeding and incorporated by disking.

2. Active ingredients:

Buctril M: 3,5-dibromo-4-hydroxybenzotrile
 Avadex: S-(2,3,3-trichloroallyl) disopropyl thiocarbamate
 MCPA-K: K-salt of 2-Methyl, 4-chlorophenoxyacetic acid
 Banvel: 2 methoxy-3,6-dichlorobenzoic acid

3. N. A. = not applicable.

Appendix 3. Agronomic characters of wheat genotypes in the study.

Genotype	Maturity ¹	Height (cm) ²	Tiller number per plant ³	1000 kernel weight (g) ⁴
Pitic 62	+6	85	4.5	36.7
Glenlea	+8	108	2.6	44.6
	-1	101	4.5	33.4
Neepawa	0	102	No data	32.9
70M110001 ⁵	+3	91	No data	44.2
70M09002 ⁶	+4	86	3.3	42.8
Norquay	+4	82	3.4	36.6
NB 701 ⁷	+1	108	No data	43.0

1. Number of days earlier or later than Neepawa (Attinaw, 1977).

2. Attinaw, 1977

3. Average of 1978, 1979 tests, plants 3 cm apart in rows 30 cm apart

4. Average of 1977, 1978, 1979 seedlots used.

5. Pedigree of 70M110001 = CIANO S × ((CIANO × SON-KI. REND)8156)

6. Pedigree of 70M00900 = CIANO S × ((SON 64-Y 50ES × G10)INIA'S')
7. Data obtained from 1976 Co-op NonBread Wheat Test, Edmonton.
Pedigree of NB701 = ((SON 64-Y 50ES × G10)INIA) × INIA'S' - NAPO

Appendix 4. Thousand kernel weight (1000 kwt) and percent germination (% germ) of the seed used for the experiments in 1977, 1978 and 1979.

Genotype	1977		1978		1979	
	1000 kwt	% germ	1000 kwt	% germ	1000 kwt	% germ
Pitic 62	34.3	84.0	39.1	99.5	No data	98.5
Glenlea	42.5	95.0	41.8	93.5	44.5	97.5
Park	33.5	96.0	31.8	99.0	34.9	98.5
Neepawa	32.1	99.0	33.2	99.0	33.5	82.5
70M110001	35.9	98.0	47.9	94.5	48.9	99.0
70M009002	33.6	92.0	48.9	100.0	45.9	99.5
Norquay	35.8	88.5	36.6	98.5	37.3	87.0
NB 701	38.9	98.0	43.1	85.0	47.1	91.0

Appendix 5. Number of plants available per treatment, from each density, for analysis of the wheelplot experiment, grown in 1977.

Genotype	Associate Genotype	Plants per treatment						
		204 ¹	124	69	44	28	17	10
Glenlea	Glenlea	11	14	13	13	15	13	8
	Neepawa	7	7	7	6	5	7	5
	70M110001	6	7	8	5	2	-	6
	70M009002	9	-	10	5	5	-	-
	NB 701	-	7	7	5	-	-	-
Park	Park	6	14	12	11	15	13	13
	Neepawa	-	5	6	-	-	-	-
	70M110001	7	8	-	-	-	-	-
	70M009002	6	7	7	8	7	6	7
	NB 701	6	7	7	9	7	7	6
Neepawa	Glenlea	7	10	7	8	6	7	6
	Park	-	7	7	-	-	-	-
	70M110001	6	-	-	-	-	-	-
	Norquay	-	-	-	-	6	7	-
	NB 701	7	6	6	9	7	6	6
70M110001	70M110001	16	10	10	11	12	8	9
	Glenlea	6	6	6	7	-	-	5
	Park	8	6	-	-	-	-	-
	Neepawa	5	-	-	-	-	-	-
	NB 701	-	-	7	9	-	-	5
70M009002	Glenlea	5	7	6	5	5	6	5
	Park	7	6	6	-	6	8	7
	NB 701	6	9	6	7	7	8	9
Norquay	Norquay	-	6	6	7	-	-	6
	Neepawa	-	-	-	-	8	7	6
	NB 701	9	6	-	-	-	-	8
NB 701	Glenlea	-	6	6	5	-	-	-
	Park	8	6	-	5	6	7	9
	Neepawa	7	7	7	7	7	7	8
	70M110001	-	-	5	6	-	-	5
	70M009002	5	5	6	8	5	5	7
	Norquay	5	8	-	-	-	5	-

1. Plant density (plants/m²)

2. Less than five plants available

Appendix 6. Descriptive statistics of characters of plants, grown in monoculture and in mixture at various interplant spacings, measured at maturity in 1977.

APPENDIX 6

03/16/82

PAGE 2

FILE W077 (CREATION DATE = 03/16/82)

DESCRIPTION OF SUBPOPULATIONS
 CRITERION VARIABLE BY CUL PLANT HEIGHT CM
 BROKEN DOWN BY CUL CULTIVAR
 BY SPAC INTER-PLANT SPACING
 BY COMP TREATMENT

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
FOR ENTIRE POPULATION			88417 0000	93 4'98	15 8889	282 8870	1049
CUL	2	6LENLEA	20231 0000	100 4341	8 8044	77 8197	182
SPAC	1	204 PLANTS M	2895 0000	102 2917	10 0748	122 8504	24
COMP	1	MONOCULTURE	258 0000	102 6364	6 1820	68 4845	11
COMP	2	MIXTURE	1448 0000	101 5332	13 2908	178 8810	13
SPAC	2	3 CM SPACING	3882 0000	103 7714	9 8825	99 8521	35
COMP	1	MONOCULTURE	1886 0000	102 1429	15 0018	225 0849	14
COMP	2	MIXTURE	2426 0000	105 8238	3 8810	18 0819	21
SPAC	3	5 CM SPACING	3897 0000	101 3429	9 2417	85 4084	35
COMP	1	MONOCULTURE	1474 0000	103 3848	3 4288	11 7584	13
COMP	2	MIXTURE	2423 0000	100 1384	11 2897	127 8887	22
SPAC	4	44 PLANTS M	2232 0000	101 4483	8 7447	78 8784	28
COMP	1	MONOCULTURE	1482 0000	104 0000	2 8820	8 8887	13
COMP	2	MIXTURE	1750 0000	108 3750	11 2778	127 1833	16
SPAC	5	28 PLANTS M	2223 0000	101 1500	4 2212	17 8184	20
COMP	1	MONOCULTURE	1863 0000	110 8467	3 7007	13 8982	15
COMP	2	MIXTURE	580 0000	102 0000	6 9582	35 8080	5
SPAC	6	17 PLANTS M	2202 0000	100 1000	5 8830	32 4188	20
COMP	1	MONOCULTURE	1440 0000	110 7892	5 8898	31 3860	13
COMP	2	MIXTURE	762 0000	106 8571	6 0845	37 1428	10
SPAC	7	10 PLANTS M	2080 0000	107 8947	6 6210	74 3218	19
COMP	1	MONOCULTURE	874 0000	109 2500	5 2265	28 8000	8

APPENDIX 5

02/16/82

PAGE 3

CRITERION VARIABLE HT

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
COMP	2	MIXTURE	1176 0000	106 9091	10 6499	111 3900	111
CUL	2	PARK	18818 0000	85 7285	7 3821	54 0527	207
SPAC	1	204 PLANTS H	2442 0000	87 7280	7 0977	50 3787	281
COMP	1	MONOCULTURE	887 0000	101 1887	8 7417	22 8687	88
COMP	2	MIXTURE	1526 0000	86 8318	7 2984	52 8012	181
SPAC	2	3 CM SPACING	2884 0000	86 4390	9 0229	81 4024	411
COMP	1	MONOCULTURE	1283 0000	88 8000	10 8484	113 2482	141
COMP	2	MIXTURE	2561 0000	84 8519	7 8039	60 9002	271
SPAC	3	6 CM SPACING	3188 0000	88 0000	9 0737	26 7419	321
COMP	1	MONOCULTURE	1220 0000	102 5000	4 2989	18 4648	121
COMP	2	MIXTURE	1926 0000	86 8000	4 2518	18 0388	201
SPAC	4	44 PLANTS H	2886 0000	86 8286	8 2324	67 7726	281
COMP	1	MONOCULTURE	1082 0000	89 2828	4 0810	16 6548	111
COMP	2	MIXTURE	1882 0000	82 7089	6 5182	60 5588	171
SPAC	5	28 PLANTS H	2618 0000	87 0880	2 7410	12 8881	261
COMP	1	MONOCULTURE	1489 0000	87 2887	2 1278	8 7810	181
COMP	2	MIXTURE	1288 0000	86 8671	4 8177	15 5188	161
SPAC	6	17 PLANTS H	2288 0000	82 1828	6 8217	40 8384	221
COMP	1	MONOCULTURE	1182 0000	81 8822	7 8418	58 2674	121
COMP	2	MIXTURE	1204 0000	82 8184	6 1717	28 0887	121
SPAC	7	10 PLANTS H	2284 0000	80 8288	6 1289	27 8288	281
COMP	1	MONOCULTURE	91 1828	8 1080	6 1080	27 2677	121
COMP	2	MIXTURE	1189 0000	89 8221	6 2308	40 0789	121
CUL	4	YONGE	12488 0000	86 4048	6 8824	78 2688	1211
SPAC	1	204 PLANTS H	1908 0000	86 4000	6 2288	58 8888	201
COMP	2	MIXTURE	1808 0000	86 4000	6 2288	58 8888	201
SPAC	2	3 CM SPACING	2206 0000	100 2174	10 2848	108 2887	221
COMP	2	MIXTURE	2206 0000	100 2174	10 2848	108 2887	221

APPENDIX 5

02/16/82

PAGE 4

CRITERION VARIABLE HT

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
SPAC	3	6 CM SPACING	1828 0000	88 8000	8 0518	64 8218	201
COMP	2	MIXTURE	1828 0000	88 8000	8 0518	64 8218	201
SPAC	4	44 PLANTS H	1818 0000	85 1768	10 9872	120 2784	171
COMP	2	MIXTURE	1818 0000	85 1768	10 9872	120 2784	171
SPAC	5	28 PLANTS H	1829 0000	88 2832	4 8288	22 2188	181
COMP	2	MIXTURE	1829 0000	88 2832	4 8288	22 2188	181
SPAC	6	17 PLANTS H	1828 0000	81 8000	6 2627	38 2211	201
COMP	2	MIXTURE	1828 0000	81 8000	6 2627	38 2211	201
SPAC	7	10 PLANTS H	1084 0000	88 8887	6 7220	45 2222	121
COMP	2	MIXTURE	1084 0000	88 8887	6 7220	45 2222	121
CUL	5	MONQUAY	10888 0000	78 1301	10 8872	117 8788	1481
SPAC	1	204 PLANTS H	2828 0000	78 4000	10 8878	117 8881	281
COMP	1	MONOCULTURE	1212 0000	78 8128	6 7180	22 2828	181
COMP	2	MIXTURE	1626 0000	76 0528	13 8802	184 8082	191
SPAC	2	3 CM SPACING	1878 0000	76 2182	16 0187	226 8884	221
COMP	1	MONOCULTURE	788 0000	78 8000	9 2888	87 8080	101
COMP	2	MIXTURE	882 0000	72 8822	18 4880	340 8824	121
SPAC	3	6 CM SPACING	1780 0000	78 0870	12 2816	182 8182	221
COMP	1	MONOCULTURE	708 0000	70 8000	7 4304	55 2111	101
COMP	2	MIXTURE	1041 0000	80 0789	14 1088	198 0789	121
SPAC	4	44 PLANTS H	2088 0000	78 8288	10 8817	111 8488	271
COMP	1	MONOCULTURE	811 0000	72 7272	8 7481	32 0182	111
COMP	2	MIXTURE	1266 0000	78 8288	12 8882	160 8187	181
SPAC	5	28 PLANTS H	844 0000	70 2322	4 2822	18 4242	121
COMP	1	MONOCULTURE	844 0000	70 2322	4 2822	18 4242	121

APPENDIX 5

02/16/82

PAGE 5

CRITERION VARIABLE HT

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
SPAC	8	17 PLANTS M	885 0000	78 5260	4 2408	17 8821	81
COMP	1	MONOCULTURE	885 0000	78 5260	4 2408	17 8821	81
SPAC	7	10 PLANTS M	1423 0000	74 8947	8 1711	68 7881	181
COMP	1	MONOCULTURE	889 0000	73 3333	2 8600	8 3000	81
COMP	2	MIXTURE	783 0000	78 3000	17 1981	123 2444	101
CUL	8	YOM000003	10184 0000	77 8188	8 2538	68 7818	181
SPAC	1	304 PLANTS M	1488 0000	82 8888	11 0887	123 3028	181
COMP	2	MIXTURE	1488 0000	82 8888	11 0887	123 3028	181
SPAC	2	3 CM SPACING	1728 0000	78 5488	7 0287	48 4028	221
COMP	2	MIXTURE	1728 0000	78 5488	7 0287	48 4028	221
SPAC	3	6 CM SPACING	1447 0000	80 3889	7 2772	52 8878	181
COMP	2	MIXTURE	1447 0000	80 3889	7 2772	52 8878	181
SPAC	4	44 PLANTS M	888 0000	82 4187	8 8188	77 7187	121
COMP	2	MIXTURE	888 0000	82 4187	8 8188	77 7187	121
SPAC	5	28 PLANTS M	1314 0000	73 0000	8 4772	30 0000	181
COMP	2	MIXTURE	1314 0000	73 0000	8 4772	30 0000	181
SPAC	8	17 PLANTS M	1888 0000	78 4848	8 7884	48 1848	221
COMP	2	MIXTURE	1888 0000	78 4848	8 7884	48 1848	221
SPAC	7	10 PLANTS M	1870 0000	74 7618	7 2488	52 4808	211
COMP	2	MIXTURE	1870 0000	74 7618	7 2488	52 4808	211
CUL	7	BORQUAY	6371 0000	77 8408	8 8188	41 1884	881
SPAC	1	304 PLANTS M	733 0000	81 4444	2 7888	7 7778	81
COMP	2	MIXTURE	733 0000	81 4444	2 7888	7 7778	81
SPAC	2	3 CM SPACING	883 0000	78 4187	3 3828	11 1782	121
COMP	1	MONOCULTURE	473 0000	78 8333	3 3808	18 8887	81

APPENDIX 6

02/16/82

PAGE 6

CRITERION VARIABLE HT

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
COMP	2	MIXTURE	480 0000	80 0000	3 8332	13 2000	81
SPAC	3	6 CM SPACING	440 0000	73 3333	8 2408	27 4887	81
COMP	1	MONOCULTURE	440 0000	73 3333	8 2408	27 4887	81
SPAC	4	44 PLANTS M	881 0000	78 7143	7 7881	81 2381	71
COMP	1	MONOCULTURE	881 0000	78 7143	7 7881	81 2381	71
SPAC	5	28 PLANTS M	888 0000	83 2800	7 8778	63 8438	81
COMP	2	MIXTURE	888 0000	83 2800	7 8778	63 8438	81
SPAC	6	17 PLANTS M	1141 0000	78 0887	8 2438	27 4882	181
COMP	2	MIXTURE	1141 0000	78 0887	8 2438	27 4882	181
SPAC	7	10 PLANTS M	887 0000	73 8167	7 3788	84 4670	121
COMP	1	MONOCULTURE	448 0000	74 8333	3 8880	13 8887	81
COMP	2	MIXTURE	438 0000	73 0000	10 2178	104 4000	81
CUL	8	NETO1	18284 0000	108 3888	8 8188	88 8887	1831
SPAC	1	304 PLANTS M	2782 0000	110 4800	7 4728	88 8433	281
COMP	2	MIXTURE	2782 0000	110 4800	7 4728	88 8433	281
SPAC	2	3 CM SPACING	3830 0000	110 3125	8 8830	91 8347	321
COMP	2	MIXTURE	3830 0000	110 3125	8 8830	91 8347	321
SPAC	2	6 CM SPACING	2832 0000	108 2800	8 2780	38 4130	281
COMP	2	MIXTURE	2832 0000	108 2800	8 2780	38 4130	281
SPAC	4	44 PLANTS M	3288 0000	108 0000	7 7888	60 2000	311
COMP	2	MIXTURE	3288 0000	108 0000	7 7888	60 2000	311
SPAC	5	28 PLANTS M	1888 0000	108 3333	8 9452	48 2383	181
COMP	2	MIXTURE	1888 0000	108 3333	8 9452	48 2383	181
SPAC	8	17 PLANTS M	2381 0000	88 2883	10 8848	113 7373	241
COMP	2	MIXTURE	2381 0000	88 2883	10 8848	113 7373	241

PAGE 3

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
SPAC	1	10 PLANTS IN MIXTURE	2811 0000	88 8310	8 0224	30 2802	2811
COMP	2	MIXTURE	2811 0000	88 8310	8 0224	30 2802	2811
TOTAL CASES =	1000						
MISSING CASES =	31 00	2 8 PCY					

PAGE 6

DESCRIPTION OF SUBPOPULATIONS

CRITERION VARIABLE		DESCRIPTION OF SUBPOPULATIONS
FL	BROKEN DOWN BY	HEIGHT OF THE FLAG LEAF BLADE CM
BY	SPAC	CULTIVAR
BY	CPMC	INTER-PLANT SPACING
BY		TRIFOLIUM

VARIABLE	COUNT	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
FOR ENTIRE POPULATION			37001 0000	38 3083	7 8172	81 1088	1000
CUL	2	ELENLEA	7181 0000	38 3482	6 4161	41 1868	182
SPAC	1	204 PLANTS M	833 0000	38 8780	4 0358	16 2880	24
COMP	1	MONOCULTURE	433 0000	38 2626	3 7785	14 2648	11
COMP	2	MIXTURE	800 0000	38 4615	4 2818	18 8389	13
SPAC	2	3 CM SPACING	1408 0000	40 0000	4 4721	20 0000	35
COMP	1	MONOCULTURE	840 0000	38 8714	5 0340	25 3407	14
COMP	2	MIXTURE	880 0000	40 0824	5 8920	18 1876	21
SPAC	3	9 CM SPACING	1383 0000	38 8428	9 0844	82 5261	26
COMP	1	MONOCULTURE	831 0000	40 8482	12 8486	194 4744	12
COMP	2	MIXTURE	832 0000	37 8182	4 3486	18 8177	22
SPAC	4	44 PLANTS M	1148 0000	38 8882	7 1840	81 8227	28
COMP	1	MONOCULTURE	818 0000	39 8184	8 3886	88 0887	18
COMP	2	MIXTURE	633 0000	38 8626	6 0324	36 3382	16
SPAC	6	88 PLANTS M	887 0000	40 3880	7 8486	83 1888	20
COMP	1	MONOCULTURE	887 0000	39 8800	8 8206	74 3182	18
COMP	2	MIXTURE	210 0000	42 0000	6 8682	38 0000	8
SPAC	8	17 PLANTS M	778 0000	38 8880	8 2408	37 4832	20
COMP	1	MONOCULTURE	801 0000	38 8285	8 2481	38 4026	19
COMP	2	MIXTURE	277 0000	38 8714	8 3807	28 0824	7
SPAC	7	10 PLANTS M	732 0000	38 8283	4 8138	20 3743	19
COMP	1	MONOCULTURE	307 0000	38 3780	5 2827	27 8884	8

APPENDIX 8

02/10/82

PAGE 8

CRITERION VARIABLE PL

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
COMP	2	MIXTURE	426 0000	20 3000	4 1830	17 2646	111
CUL	3	204 PLANTS M	890 0000	41 1000	6 2030	27 0700	207
SPAC	1	MONOCULTURE	900 0000	30 0000	6 3343	40 1233	201
COMP	1	MIXTURE	200 0000	42 0000	3 0000	12 0000	61
COMP	2	MIXTURE	742 0000	30 1000	6 0467	40 0773	101
SPAC	2	3 CM SPACING	1007 0000	61 3000	6 3140	27 1030	411
COMP	1	MONOCULTURE	804 0000	42 1420	4 0000	24 0000	141
COMP	2	MIXTURE	1003 0000	40 1010	6 1000	27 0200	271
SPAC	3	6 CM SPACING	1210 0000	41 1000	6 4070	30 1120	331
COMP	1	MONOCULTURE	820 0000	43 3333	2 2000	5 1010	121
COMP	2	MIXTURE	700 0000	30 0000	6 4204	41 2211	201
SPAC	4	44 PLANTS M	1177 0000	42 0307	6 1703	34 1030	201
COMP	1	MONOCULTURE	400 0000	44 3636	3 0000	10 0000	21
COMP	2	MIXTURE	800 0000	40 0000	6 0000	40 0000	171
SPAC	5	20 PLANTS M	1201 0000	43 4000	3 0240	10 0010	201
COMP	1	MONOCULTURE	900 0000	43 3333	3 0000	10 0000	101
COMP	2	MIXTURE	811 0000	43 0420	4 1207	17 0100	141
SPAC	6	17 PLANTS M	1001 0000	40 0231	3 0211	10 0030	201
COMP	1	MONOCULTURE	822 0000	40 0231	4 2017	10 0000	121
COMP	2	MIXTURE	810 0000	30 0231	3 0000	12 0000	121
SPAC	7	10 PLANTS M	1000 0000	30 0000	3 0000	10 0000	201
COMP	1	MONOCULTURE	810 0000	30 0000	4 0000	20 0000	121
COMP	2	MIXTURE	400 0000	40 0000	2 0000	5 0000	121
CUL	4	700000000	4471 0000	34 1000	6 0000	31 0000	1311
SPAC	1	204 PLANTS M	890 0000	34 7000	6 0000	34 0000	201
COMP	2	MIXTURE	800 0000	34 7000	6 0000	34 0000	201
SPAC	2	3 CM SPACING	742 0000	34 0000	4 0000	22 0000	231
COMP	2	MIXTURE	702 0000	34 0000	4 0000	22 0000	231

APPENDIX 8

02/10/82

PAGE 10

CRITERION VARIABLE PL

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
SPAC	3	9 CM SPACING	710 0000	30 0000	6 3010	40 0737	201
COMP	2	MIXTURE	710 0000	30 0000	6 3010	40 0737	201
SPAC	4	44 PLANTS M	800 0000	32 0000	6 0733	47 2020	171
COMP	2	MIXTURE	800 0000	32 0000	6 0733	47 2020	171
SPAC	5	20 PLANTS M	843 0000	33 0000	4 0000	10 0000	101
COMP	2	MIXTURE	843 0000	33 0000	4 0000	10 0000	101
SPAC	6	17 PLANTS M	800 0000	33 0000	3 0000	12 0000	201
COMP	2	MIXTURE	800 0000	33 0000	3 0000	12 0000	201
SPAC	7	10 PLANTS M	421 0000	30 0000	3 0000	10 0000	121
COMP	2	MIXTURE	421 0000	30 0000	3 0000	10 0000	121
CUL	5	NORWAY	4400 0000	30 0000	6 0000	31 0000	1401
SPAC	1	204 PLANTS M	1000 0000	31 0000	4 0000	10 0000	201
COMP	1	MONOCULTURE	800 0000	31 0000	3 0000	10 0000	101
COMP	2	MIXTURE	870 0000	30 0000	6 0000	20 0000	101
SPAC	2	3 CM SPACING	840 0000	28 0000	6 0000	40 0000	221
COMP	1	MONOCULTURE	300 0000	30 0000	3 0000	10 0000	101
COMP	2	MIXTURE	240 0000	20 0000	6 0000	40 0000	121
SPAC	3	9 CM SPACING	700 0000	31 0000	4 0000	10 0000	231
COMP	1	MONOCULTURE	310 0000	31 0000	4 0000	10 0000	101
COMP	2	MIXTURE	410 0000	31 0000	4 0000	10 0000	121
SPAC	4	44 PLANTS M	800 0000	31 0000	7 0000	50 0000	201
COMP	1	MONOCULTURE	310 0000	31 0000	3 0000	10 0000	101
COMP	2	MIXTURE	400 0000	30 0000	6 0000	40 0000	101
SPAC	5	20 PLANTS M	300 0000	20 0000	4 0000	21 0000	121
COMP	1	MONOCULTURE	300 0000	20 0000	4 0000	21 0000	121

APPENDIX 8

03/18/82

PAGE 11

CRITERION VARIABLE PL

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
SPAC	8	17 PLANTS M	259 0000	32 3750	4 4701	18 8821	81
COMP	1	MONOCULTURE	259 0000	32 3750	4 4701	18 8821	81
SPAC	7	10 PLANTS M	302 0000	31 8842	3 4489	11 8847	181
COMP	1	MONOCULTURE	285 0000	31 8887	1 8028	3 2500	81
COMP	2	MIXTURE	317 0000	31 7000	4 8717	20 8800	181
CUL	8	70M000002	3409 0000	26 0228	8 3128	28 2228	1211
SPAC	1	204 PLANTS M	811 0000	28 3888	8 1488	28 4888	181
COMP	2	MIXTURE	811 0000	28 3888	8 1488	28 4888	181
SPAC	2	3 CM SPACING	883 0000	28 8848	4 8883	22 8888	221
COMP	2	MIXTURE	883 0000	28 8848	4 8883	22 8888	221
SPAC	3	8 CM SPACING	488 0000	27 1111	7 8823	83 3887	181
COMP	2	MIXTURE	488 0000	27 1111	7 8823	83 3887	181
SPAC	4	44 PLANTS M	304 0000	26 3333	6 3880	40 4242	121
COMP	2	MIXTURE	304 0000	26 3333	6 3880	40 4242	121
SPAC	5	28 PLANTS M	448 0000	24 7222	3 3821	11 8888	181
COMP	2	MIXTURE	448 0000	24 7222	3 3821	11 8888	181
SPAC	6	17 PLANTS M	888 0000	28 7273	4 8822	18 8173	221
COMP	2	MIXTURE	888 0000	28 7273	4 8822	18 8173	221
SPAC	7	10 PLANTS M	802 0000	23 9048	4 8280	21 3888	211
COMP	2	MIXTURE	802 0000	23 9048	4 8280	21 3888	211
CUL	7	BORQUAY	1883 0000	28 3843	4 8880	21 8878	881
SPAC	1	204 PLANTS M	271 0000	28 1111	8 4887	41 8111	81
COMP	2	MIXTURE	271 0000	28 1111	8 4887	41 8111	81
SPAC	2	3 CM SPACING	241 0000	28 4187	5 1778	28 8188	121
COMP	1	MONOCULTURE	182 0000	27 8888	2 4488	8 8888	81

APPENDIX 8

03/18/82

PAGE 12

CRITERION VARIABLE PL

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
COMP	2	MIXTURE	178 0000	28 8333	8 8482	48 1887	81
SPAC	3	8 CM SPACING	182 0000	28 3333	3 8287	14 8887	81
COMP	1	MONOCULTURE	182 0000	28 3333	3 8287	14 8887	81
SPAC	4	44 PLANTS M	182 0000	27 4288	2 1482	4 8180	71
COMP	1	MONOCULTURE	182 0000	27 4288	2 1482	4 8180	71
SPAC	5	28 PLANTS M	288 0000	31 8750	4 8881	18 8884	81
COMP	2	MIXTURE	288 0000	31 8750	4 8881	18 8884	81
SPAC	6	17 PLANTS M	408 0000	27 2000	3 7284	13 8887	181
COMP	2	MIXTURE	408 0000	27 2000	3 7284	13 8887	181
SPAC	7	10 PLANTS M	334 0000	27 8333	4 8838	18 8182	121
COMP	1	MONOCULTURE	171 0000	28 8000	3 8341	14 7000	81
COMP	2	MIXTURE	163 0000	27 1887	4 8388	20 8887	81
CUL	8	HS701	7812 0000	28 3188	8 8721	47 2387	1821
SPAC	1	204 PLANTS M	813 0000	28 8200	8 8784	28 7888	281
COMP	2	MIXTURE	813 0000	28 8200	8 8784	28 7888	281
SPAC	2	3 CM SPACING	1287 0000	27 7188	5 8178	28 1788	321
COMP	2	MIXTURE	1287 0000	27 7188	5 8178	28 1788	321
SPAC	3	8 CM SPACING	881 0000	40 8417	4 2781	18 3828	241
COMP	2	MIXTURE	881 0000	40 8417	4 2781	18 3828	241
SPAC	4	44 PLANTS M	1222 0000	38 4184	12 1378	147 3183	311
COMP	2	MIXTURE	1222 0000	38 4184	12 1378	147 3183	311
SPAC	5	28 PLANTS M	712 0000	28 8888	5 4788	28 8281	181
COMP	2	MIXTURE	712 0000	28 8888	5 4788	28 8281	181
SPAC	6	17 PLANTS M	800 0000	27 8000	8 8888	47 8438	241
COMP	2	MIXTURE	800 0000	27 8000	8 8888	47 8438	241

184

APPENDIX 6 03/16/82 PAGE 13

CRITERION VARIABLE PL

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
SPAC	7	16 PLANTS M	1097 0000	27 6278	4 3951	18 3621	201
COMP	2	MIXTURE	1097 0000	27 6278	4 3951	18 3621	201
TOTAL CASES *	1000						
MISSING CASES *	22 94	2 9 PET					

APPENDIX 6 03/16/82 PAGE 14

FILE W977 (CREATION DATE: 03/16/82)

CRITERION VARIABLE N OF SUB POPULATIONS

CRITERION VARIABLE	N	DESCRIPTION OF SUB POPULATIONS	SUM	MEAN	STD DEV	VARIANCE	N
BROKEN DOWN BY		CUL					
BY		SPAC					
BY		COMP					
VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
FOR ENTIRE POPULATION			7821 0000	7 2888	5 0084	25 0848	10641
CUL	2	ELENLEA	1800 0000	8 5385	7 7422	45 4675	1811
SPAC	1	204 PLANTS M	84 0000	3 6167	7917	3 2101	241
COMP	1	MONOCULTURE	44 0000	4 0000	7888	3 2000	111
COMP	2	MIXTURE	59 0000	3 5452	8640	3 4744	131
SPAC	2	3 CM SPACING	151 0000	4 4412	3988	1 5610	341
COMP	1	MONOCULTURE	55 0000	4 4615	3914	1 5259	121
COMP	2	MIXTURE	83 0000	4 4288	4343	2 0571	211
SPAC	3	6 CM SPACING	102 0000	5 4657	8210	3 3169	351
COMP	1	MONOCULTURE	68 0000	5 2304	8892	2 8268	121
COMP	2	MIXTURE	124 0000	5 6254	8851	3 8615	221
SPAC	4	44 PLANTS M	225 0000	8 2089	2 3580	5 0985	251
COMP	1	MONOCULTURE	111 0000	8 8265	1 7818	3 1928	121
COMP	2	MIXTURE	197 0000	7 9378	2 6186	6 8675	181
SPAC	5	28 PLANTS M	240 0000	12 0000	5 6068	31 3684	291
COMP	1	MONOCULTURE	155 0000	10 5333	4 3493	18 6381	151
COMP	2	MIXTURE	85 0000	16 4000	7 1274	50 6000	91
SPAC	6	17 PLANTS M	280 0000	14 5000	8 1000	37 2105	301
COMP	1	MONOCULTURE	202 0000	15 5285	5 0589	25 6025	121
COMP	2	MIXTURE	88 0000	12 5714	7 7428	59 9624	71
SPAC	7	10 PLANTS M	285 0000	20 7895	8 4553	71 5028	181
COMP	1	MONOCULTURE	151 0000	20 1250	8 1113	26 1250	81

APPENDIX 8		03/10/82				PAGE 18	
CRITERION VARIABLE M							
VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
COMP	2	MIXTURE	234 0000	21 2727	10 4704	108 8183	111
CUL	3	PARE	1237 0000	8 4803	2 3888	11 4708	208
SPAC	1	204 PLANTS M	102 0000	4 0400	1 7080	2 9100	204
COMP	1	MONOCULTURE	24 0000	4 0000	1 8574	3 0000	81
COMP	2	MIXTURE	78 0000	4 1083	1 8982	2 8772	107
SPAC	2	3 CM SPACING	188 0000	4 8500	2 3883	8 8807	407
COMP	1	MONOCULTURE	74 0000	8 0000	1 8247	1 5000	121
COMP	2	MIXTURE	120 0000	4 4444	2 8908	7 0288	271
SPAC	3	6 CM SPACING	178 0000	8 8838	2 1077	4 4428	821
COMP	1	MONOCULTURE	74 0000	8 1087	1 8880	3 8081	121
COMP	2	MIXTURE	108 0000	8 2880	2 1878	4 8288	201
SPAC	4	44 PLANTS M	188 0000	8 7887	2 8884	7 2118	281
COMP	1	MONOCULTURE	78 0000	8 8182	2 8882	8 7888	111
COMP	2	MIXTURE	118 0000	8 7887	2 8887	8 8812	171
SPAC	5	28 PLANTS M	244 0000	8 4128	2 1122	4 4888	201
COMP	1	MONOCULTURE	121 0000	8 0887	2 4831	8 0887	181
COMP	2	MIXTURE	123 0000	8 7887	1 8723	2 7887	141
SPAC	6	18 PLANTS M	222 0000	8 8788	8 1708	28 7328	281
COMP	1	MONOCULTURE	108 0000	8 3077	2 3283	11 8041	121
COMP	2	MIXTURE	118 0000	8 8482	8 8888	44 4744	121
SPAC	7	10 PLANTS M	201 0000	7 7308	4 0288	18 2048	201
COMP	1	MONOCULTURE	82 0000	8 2848	4 1841	17 2844	121
COMP	2	MIXTURE	118 0000	8 0788	3 8484	12 8788	121
CUL	4	700000002	808 0000	8 8213	4 0004	18 0000	1211
SPAC	1	204 PLANTS M	88 0000	4 2880	2 1481	4 8184	201
COMP	2	MIXTURE	88 0000	4 2880	2 1481	4 8184	201
SPAC	2	3 CM SPACING	88 0000	4 2880	1 8838	3 4742	221
COMP	2	MIXTURE	88 0000	4 2880	1 8838	3 4742	221

APPENDIX 8		03/10/82		PAGE 18			
CRITERION VARIABLE M							
VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
SPAC	3	3 CM SPACING	128 0000	8 4000	2 4288	8 9388	201
COMP	2	MIXTURE	128 0000	8 4000	2 4288	8 9388	201
SPAC	4	44 PLANTS M	101 0000	8 8412	1 8834	3 8328	171
COMP	2	MIXTURE	101 0000	8 8412	1 8834	3 8328	171
SPAC	5	28 PLANTS M	182 0000	8 8282	2 4700	12 0400	181
COMP	2	MIXTURE	182 0000	8 8282	2 4700	12 0400	181
SPAC	6	17 PLANTS M	188 0000	8 2880	2 8884	12 8342	201
COMP	2	MIXTURE	188 0000	8 2880	2 8884	12 8342	201
SPAC	7	10 PLANTS M	148 0000	12 4187	8 2888	38 1742	121
COMP	2	MIXTURE	148 0000	12 4187	8 2888	38 1742	121
CUL	5	MOROCCO	821 0000	8 1817	4 8742	20 8240	1481
SPAC	1	204 PLANTS M	124 0000	8 8288	2 0781	4 2227	281
COMP	1	MONOCULTURE	82 0000	7 8378	1 1818	1 3888	181
COMP	2	MIXTURE	71 0000	7 7888	2 8424	8 8828	101
SPAC	2	3 CM SPACING	108 0000	8 0478	1 8021	3 2478	211
COMP	1	MONOCULTURE	88 0000	8 8000	2 0111	4 0444	101
COMP	2	MIXTURE	80 0000	4 8488	2 2778	2 8772	111
SPAC	3	6 CM SPACING	108 0000	4 8817	1 8888	2 3818	821
COMP	1	MONOCULTURE	42 0000	4 2000	1 8888	3 8887	101
COMP	2	MIXTURE	81 0000	4 8822	1 2888	1 8841	121
SPAC	4	44 PLANTS M	200 0000	7 4074	2 8880	8 1728	271
COMP	1	MONOCULTURE	88 0000	8 8888	2 8888	8 8888	111
COMP	2	MIXTURE	112 0000	7 0000	2 8048	7 8887	181
SPAC	5	28 PLANTS M	80 0000	8 8887	3 8780	12 7878	121
COMP	1	MONOCULTURE	80 0000	8 8887	3 8780	12 7878	121

APPENDIX 6

02/10/82

PAGE 17

CRITERION VARIABLE M

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
SPAC	1	17 PLANTS M	88 0000	8 8250	8 8070	82 1260	81
COMP	1	MONOCULTURE	88 0000	8 8250	8 8070	82 1260	81
SPAC	7	10 PLANTS M	228 0000	12 0000	7 7680	80 0000	101
COMP	1	MONOCULTURE	102 0000	11 3232	6 0000	36 0000	81
COMP	2	MIXTURE	126 0000	12 6000	8 3267	87 1956	101
CUL	8	700000002	888 0000	8 7834	4 8019	24 0282	121
SPAC	1	204 PLANTS M	81 0000	3 3884	1 2887	1 8834	101
COMP	2	MIXTURE	81 0000	3 3884	1 2887	1 8834	101
SPAC	2	3 CM SPACING	83 0000	3 7727	1 4778	2 1840	121
COMP	2	MIXTURE	83 0000	3 7727	1 4778	2 1840	121
SPAC	3	8 CM SPACING	84 0000	4 8887	2 0884	4 3828	101
COMP	2	MIXTURE	84 0000	4 8887	2 0884	4 3828	101
SPAC	4	44 PLANTS M	80 0000	6 8887	2 3484	8 5182	121
COMP	2	MIXTURE	80 0000	6 8887	2 3484	8 5182	121
SPAC	5	28 PLANTS M	118 0000	8 8111	3 3238	11 0782	101
COMP	2	MIXTURE	118 0000	8 8111	3 3238	11 0782	101
SPAC	6	17 PLANTS M	208 0000	8 4848	4 8327	20 8488	121
COMP	2	MIXTURE	208 0000	8 4848	4 8327	20 8488	121
SPAC	7	10 PLANTS M	251 0000	11 8824	7 3788	84 4478	121
COMP	2	MIXTURE	251 0000	11 8824	7 3788	84 4478	121
CUL	7	880000	880 0000	8 8887	8 7842	22 4888	81
SPAC	1	204 PLANTS M	48 0000	4 4444	0 8818	0 7778	81
COMP	2	MIXTURE	40 0000	4 4444	0 8818	0 7778	81
SPAC	2	3 CM SPACING	82 0000	8 1887	1 8824	3 8887	121
COMP	1	MONOCULTURE	30 0000	8 0000	2 8288	8 4000	81

APPENDIX 6

02/10/82

PAGE 18

CRITERION VARIABLE M

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
COMP	2	MIXTURE	32 0000	8 3332	1 5088	2 2867	81
SPAC	3	8 CM SPACING	31 0000	8 1887	2 8384	8 8887	81
COMP	1	MONOCULTURE	31 0000	8 1887	2 8384	8 8887	81
SPAC	4	44 PLANTS M	84 0000	7 7142	2 8804	7 2381	71
COMP	1	MONOCULTURE	84 0000	7 7142	2 8804	7 2381	71
SPAC	5	28 PLANTS M	82 0000	11 8250	3 4280	11 8884	81
COMP	2	MIXTURE	82 0000	11 8250	3 4280	11 8884	81
SPAC	6	17 PLANTS M	174 0000	11 8000	7 0880	48 9714	181
COMP	2	MIXTURE	174 0000	11 8000	7 0880	48 9714	181
SPAC	7	10 PLANTS M	128 0000	11 3332	7 8488	82 1818	121
COMP	1	MONOCULTURE	72 0000	12 0000	8 8128	82 4000	81
COMP	2	MIXTURE	84 0000	10 8887	8 7428	48 4887	81
CUL	8	88001	1278 0000	7 8188	4 8882	28 8788	181
SPAC	1	204 PLANTS M	104 0000	4 1888	1 2477	1 8887	28
COMP	2	MIXTURE	104 0000	4 1888	1 2477	1 8887	28
SPAC	2	3 CM SPACING	132 0000	4 2883	1 8772	2 8128	311
COMP	2	MIXTURE	132 0000	4 2883	1 8772	2 8128	311
SPAC	3	8 CM SPACING	131 0000	8 4882	2 0212	4 0881	241
COMP	2	MIXTURE	131 0000	8 4882	2 0212	4 0881	241
SPAC	4	44 PLANTS M	208 0000	8 7887	2 7712	7 8788	311
COMP	2	MIXTURE	208 0000	8 7887	2 7712	7 8788	311
SPAC	5	28 PLANTS M	188 0000	11 8888	2 8888	8 8887	181
COMP	2	MIXTURE	188 0000	11 8888	2 8888	8 8887	181
SPAC	6	17 PLANTS M	288 0000	11 8870	8 4344	28 8378	231
COMP	2	MIXTURE	288 0000	11 8870	8 4344	28 8378	231

APPENDIX B

03/16/82

PAGE 19

CRITERION VARIABLE H

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
SPAC	7	10 PLANTS M	345 0000	12 0345	5 4834	142 0345	281
COMP	2	MIXTURE	345 0000	12 0345	5 4834	142 0345	281
TOTAL CASES =	1000						
MISSING CASES =	25 OR	3 3 PCT					

APPENDIX B

03/16/82

PAGE 20

FILE WP77 (CREATION DATE = 03/16/82)

DESCRIPTION OF SUB POPULATIONS

CRITERION VARIABLE BROKEN DOWN BY	T BY BY	CUL SPAC COMP	TILLERS PER PLANT CULTIVAR INTER-PLANT SPACING TREATMENT	SUM	MEAN	STD DEV	VARIANCE	N
FOR ENTIRE POPULATION				10255 0000	5 7766	5 2632	26 8774	10481
CUL	2	6LENLEA		2073 0000	11 3848	5 3191	27 8520	182
SPAC	1	204 PLANTS M		123 0000	5 1250	3 1112	9 6783	24
COMP	1	MONOCULTURE		54 0000	4 8081	2 8784	8 2808	11
COMP	2	MIXTURE		69 0000	5 3077	3 4006	11 6841	13
SPAC	2	3 CM SPACING		186 0000	5 3143	2 0289	4 1042	35
COMP	1	MONOCULTURE		87 0000	4 7857	1 8472	3 4121	14
COMP	2	MIXTURE		119 0000	5 8857	3 1055	4 4323	21
SPAC	3	6 CM SPACING		249 0000	7 1143	2 6933	7 2807	35
COMP	1	MONOCULTURE		80 0000	5 1928	1 8788	3 5077	12
COMP	2	MIXTURE		169 0000	7 8818	3 0458	9 2748	22
SPAC	4	24 PLANTS M		320 0000	11 0345	3 3433	11 1773	20
COMP	1	MONOCULTURE		135 0000	10 3848	1 8807	3 5331	13
COMP	2	MIXTURE		185 0000	11 0808	4 1507	17 0825	16
SPAC	5	28 PLANTS M		289 0000	14 6500	5 2448	26 8874	20
COMP	1	MONOCULTURE		101 0000	12 7323	4 8788	23 7810	18
COMP	2	MIXTURE		188 0000	21 8000	5 3195	26 3000	5
SPAC	6	17 PLANTS M		378 0000	18 6500	5 0523	25 5395	20
COMP	1	MONOCULTURE		247 0000	18 0000	4 3589	19 0000	13
COMP	2	MIXTURE		132 0000	18 8571	5 6189	32 5095	7
SPAC	7	10 PLANTS M		516 0000	27 1879	7 7283	59 8859	18
COMP	1	MONOCULTURE		195 0000	24 3750	5 3457	40 2878	8

APPENDIX 8

03/10/82

PAGE 21

CRITERION VARIABLE Y

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
COMP	3	MIXTURE	221 0000	29 1818	4 2603	84 8638	111
CUL	3	PARK	1048 0000	8 8130	4 8788	20 8833	207
SPAC	1	204 PLANTS M	134 0000	8 2800	2 3238	4 8000	28
COMP	1	MONOCULTURE	23 0000	8 8000	1 8708	3 8000	8
COMP	2	MIXTURE	101 0000	8 2188	2 3817	8 8738	18
SPAC	2	3 CM SPACING	288 0000	8 8810	3 1847	8 8834	41
COMP	1	MONOCULTURE	114 0000	8 1428	3 0881	8 8848	14
COMP	2	MIXTURE	188 0000	8 7407	2 8880	8 7378	27
SPAC	3	9 CM SPACING	281 0000	7 8438	3 9278	8 1862	32
COMP	1	MONOCULTURE	101 0000	8 4187	2 8888	8 3881	12
COMP	2	MIXTURE	180 0000	7 8888	3 9348	8 2188	20
SPAC	4	44 PLANTS M	248 0000	8 7888	3 8848	14 8282	28
COMP	1	MONOCULTURE	88 0000	8 8888	4 3288	18 8888	11
COMP	2	MIXTURE	168 0000	8 8882	3 8882	12 3824	17
SPAC	5	28 PLANTS M	304 0000	10 8828	2 8871	8 8818	28
COMP	1	MONOCULTURE	188 0000	10 3232	3 1881	8 8887	18
COMP	2	MIXTURE	118 0000	10 8428	2 8878	4 4811	14
SPAC	6	17 PLANTS M	217 0000	12 1823	6 8881	21 3818	28
COMP	1	MONOCULTURE	188 0000	12 8823	6 4218	28 3874	12
COMP	2	MIXTURE	182 0000	11 8823	8 8488	28 3874	12
SPAC	7	18 PLANTS M	228 0000	12 8888	8 8888	24 8288	28
COMP	1	MONOCULTURE	184 0000	11 8482	8 1488	27 8877	12
COMP	2	MIXTURE	171 0000	12 1828	8 8144	22 8877	12
CUL	4	700000002	1288 0000	8 8187	8 8832	28 8388	121
SPAC	1	204 PLANTS M	118 0000	8 7888	3 4888	11 8888	20
COMP	2	MIXTURE	118 0000	8 7888	3 4888	11 8888	20
SPAC	3	3 CM SPACING	182 0000	8 1728	2 8788	8 8778	22
COMP	2	MIXTURE	182 0000	8 1728	2 8788	8 8778	22

APPENDIX 8

03/10/82

PAGE 22

CRITERION VARIABLE Y

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
SPAC	3	9 CM SPACING	188 0000	8 4888	3 8217	14 8818	28
COMP	2	MIXTURE	188 0000	8 4888	3 8217	14 8818	28
SPAC	4	44 PLANTS M	148 0000	8 8882	2 8888	8 7874	17
COMP	2	MIXTURE	148 0000	8 8882	2 8888	8 7874	17
SPAC	5	28 PLANTS M	228 0000	12 8828	3 7828	14 1827	18
COMP	2	MIXTURE	228 0000	12 8828	3 7828	14 1827	18
SPAC	6	17 PLANTS M	288 0000	12 8888	4 8878	18 8832	28
COMP	2	MIXTURE	288 0000	12 8888	4 8878	18 8832	28
SPAC	7	18 PLANTS M	182 0000	18 1887	8 8888	88 8882	12
COMP	2	MIXTURE	182 0000	18 1887	8 8888	88 8882	12
CUL	8	NORWAY	181 0000	8 8888	8 8782	28 7888	148
SPAC	1	204 PLANTS M	182 0000	4 2828	2 3288	8 8888	28
COMP	1	MONOCULTURE	88 0000	4 2888	1 8827	1 4888	18
COMP	2	MIXTURE	84 0000	4 4211	3 8888	9 8881	18
SPAC	2	3 CM SPACING	128 0000	8 8828	3 1887	10 8281	28
COMP	1	MONOCULTURE	73 0000	7 3888	3 8888	18 8887	18
COMP	2	MIXTURE	88 0000	4 8887	1 3233	2 8888	12
SPAC	3	9 CM SPACING	184 0000	8 2888	2 8288	8 3824	22
COMP	1	MONOCULTURE	83 0000	8 3888	2 3888	8 8888	10
COMP	2	MIXTURE	91 0000	7 8888	2 4823	8 1888	12
SPAC	4	44 PLANTS M	283 0000	8 3784	3 8248	14 8288	27
COMP	1	MONOCULTURE	187 0000	8 7272	4 1888	17 4182	11
COMP	2	MIXTURE	188 0000	8 1288	3 8888	13 8822	18
SPAC	5	28 PLANTS M	188 0000	8 7888	4 8282	20 8882	12
COMP	1	MONOCULTURE	188 0000	8 7888	4 8282	20 8882	12

APPENDIX 8
CRITERION VARIABLE Y

02/18/82

PAGE 23

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
SPAC	8	17 PLANTS M	88 0000	10 7800	7 3241	53 8428	81
COMP	1	MONOCULTURE	88 0000	10 7800	7 3241	53 8428	81
SPAC	7	10 PLANTS M	312 0000	18 4211	8 2333	68 2360	181
COMP	1	MONOCULTURE	132 0000	14 8887	8 7034	75 7800	81
COMP	2	MIXTURE	180 0000	18 0000	10 0233	100 8687	181
CUL	8	700000002	1248 0000	8 8118	8 1138	67 3768	131
SPAC	1	204 PLANTS M	78 0000	4 2322	3 1629	4 4163	181
COMP	2	MIXTURE	78 0000	4 2322	3 1629	4 4163	181
SPAC	3	3 CM SPACING	108 0000	4 7727	1 7877	3 2318	231
COMP	2	MIXTURE	108 0000	4 7727	1 7877	3 2318	231
SPAC	3	8 CM SPACING	112 0000	8 2778	3 8887	8 8830	181
COMP	2	MIXTURE	112 0000	8 2778	3 8887	8 8830	181
SPAC	4	44 PLANTS M	122 0000	10 2800	3 4148	11 8881	121
COMP	2	MIXTURE	122 0000	10 2800	3 4148	11 8881	121
SPAC	8	28 PLANTS M	178 0000	8 7778	4 2778	18 3007	181
COMP	2	MIXTURE	178 0000	8 7778	4 2778	18 3007	181
SPAC	8	17 PLANTS M	308 0000	14 0488	8 2088	38 8218	221
COMP	2	MIXTURE	308 0000	14 0488	8 2088	38 8218	221
SPAC	7	10 PLANTS M	344 0000	18 3810	8 8438	34 1478	211
COMP	2	MIXTURE	344 0000	18 3810	8 8438	34 1478	211
CUL	7	800000002	840 0000	12 1738	7 1413	60 8887	81
SPAC	1	204 PLANTS M	78 0000	4 4444	0 8818	0 7778	81
COMP	2	MIXTURE	78 0000	4 4444	0 8818	0 7778	81
SPAC	3	3 CM SPACING	88 0000	8 8887	3 2848	10 7878	181
COMP	1	MONOCULTURE	88 0000	7 8887	4 2888	18 8887	81

APPENDIX 8
CRITERION VARIABLE Y

02/18/82

PAGE 24

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
COMP	2	MIXTURE	34 0000	8 8887	1 8088	2 2887	81
SPAC	3	8 CM SPACING	47 0000	7 8333	3 8707	15 7887	81
COMP	1	MONOCULTURE	47 0000	7 8333	3 8707	15 7887	81
SPAC	4	44 PLANTS M	90 0000	12 8871	2 8838	8 1428	71
COMP	1	MONOCULTURE	90 0000	12 8871	2 8838	8 1428	71
SPAC	8	28 PLANTS M	132 0000	16 8000	3 8848	13 8288	81
COMP	2	MIXTURE	132 0000	16 8000	3 8848	13 8288	81
SPAC	8	17 PLANTS M	223 0000	14 8887	8 8888	38 8381	181
COMP	2	MIXTURE	223 0000	14 8887	8 8888	38 8381	181
SPAC	7	10 PLANTS M	228 0000	18 0000	8 4423	71 2727	121
COMP	1	MONOCULTURE	112 0000	18 8887	8 8388	78 8887	81
COMP	2	MIXTURE	118 0000	18 3333	8 7880	78 8887	81
CUL	8	80701	1812 0000	8 8018	8 7322	32 8884	1831
SPAC	1	204 PLANTS M	140 0000	8 8000	1 8028	3 2800	281
COMP	2	MIXTURE	140 0000	8 8000	1 8028	3 2800	281
SPAC	3	3 CM SPACING	174 0000	8 4378	2 2882	8 2883	231
COMP	2	MIXTURE	174 0000	8 4378	2 2882	8 2883	231
SPAC	3	8 CM SPACING	188 0000	8 8187	2 8180	8 8148	241
COMP	2	MIXTURE	188 0000	8 8187	2 8180	8 8148	241
SPAC	4	44 PLANTS M	274 0000	8 8387	3 8818	13 4088	211
COMP	2	MIXTURE	274 0000	8 8387	3 8818	13 4088	211
SPAC	8	28 PLANTS M	228 0000	12 7222	3 3220	11 8388	181
COMP	2	MIXTURE	228 0000	12 7222	3 3220	11 8388	181
SPAC	8	17 PLANTS M	384 0000	18 1887	8 8882	27 1884	241
COMP	2	MIXTURE	384 0000	18 1887	8 8882	27 1884	241

APPENDIX 6
CRITERION VARIABLE 7
02/16/82
PAGE 26

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
SPAC	7	10 PLANTS M	455 0000	10 0345	5 8084	20 3202	20
COMP	2	MIXTURE	455 0000	10 0345	5 8084	20 3202	20
TOTAL CASES = 1000							
MISSING CASES = 31 OR 3.1 PCT							

APPENDIX 6
FILE WP77 (CREATION DATE = 02/16/82)
02/16/82
PAGE 26

CRITERION VARIABLE 7
ORDERED DOWN BY
BY
BY

DESCRIPTION OF SUBPOPULATIONS
WT
CUL
SPAC
COMP
CULTIVAR
INTER-PLANT SPACING
TREATMENT

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
FOR ENTIRE POPULATION			48123 3225	48 0510	28 8858	823 4380	1000
CUL	2	GLENLEA	10824 2487	58 1480	28 9136	1288 7476	181
SPAC	1	204 PLANTS M	744 4889	31 0204	14 9801	223 5083	24
COMP	1	MONOCULTURE	357 2889	33 3882	16 5810	275 2816	11
COMP	2	MIXTURE	377 2100	28 0189	13 7464	189 5149	12
SPAC	2	3 CM SPACING	1120 3888	32 2888	7 8875	61 7404	25
COMP	1	MONOCULTURE	428 8489	31 3484	8 0405	64 8802	14
COMP	2	MIXTURE	691 5399	32 8308	7 8865	61 8822	21
SPAC	3	8 CM SPACING	1238 4887	38 1831	10 2451	104 8828	25
COMP	1	MONOCULTURE	489 8889	38 1831	8 8787	75 3187	12
COMP	2	MIXTURE	688 4188	38 3827	11 0834	122 8410	22
SPAC	4	44 PLANTS M	1828 3888	54 8211	12 2460	148 8387	28
COMP	1	MONOCULTURE	700 8188	53 8881	8 8872	47 1882	12
COMP	2	MIXTURE	888 8880	58 2880	16 7420	247 8120	18
SPAC	5	28 PLANTS M	1443 8384	72 1870	22 4488	503 8388	20
COMP	1	MONOCULTURE	957 7288	65 8487	19 4882	380 1800	18
COMP	2	MIXTURE	486 2100	61 2420	21 4128	458 8480	8
SPAC	6	17 PLANTS M	1870 4889	84 8228	31 4381	988 1881	20
COMP	1	MONOCULTURE	1288 8288	88 2023	22 8887	854 8488	12
COMP	2	MIXTURE	880 8400	87 2828	64 8071	2016 8423	7
SPAC	7	10 PLANTS M	2388 1888	124 8926	28 7888	717 8381	18
COMP	1	MONOCULTURE	958 1088	119 3887	18 8841	288 8827	8

APPENDIX 6

03/16/82

PAGE 27

CRITERION VARIABLE WT

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
COMP	2	MIXTURE	114 0500	128 5500	21 2071	073 8818	(11)
CUL	2	PARK	552 2288	35 7088	13 2816	178 5160	(200)
SPAC	1	204 PLANTS M	527 2488	25 0940	5 4200	28 4881	(25)
COMP	1	MONOCULTURE	145 8400	24 3067	5 0088	25 0878	(8)
COMP	2	MIXTURE	451 5088	25 3438	5 5830	32 0700	(10)
SPAC	2	3 CM SPACING	1131 8088	27 8002	7 2853	54 8806	(41)
COMP	1	MONOCULTURE	422 1100	30 1514	5 8427	42 8073	(14)
COMP	2	MIXTURE	708 4888	25 2774	7 5772	57 4135	(27)
SPAC	3	8 CM SPACING	1023 0788	32 2837	5 3310	40 0813	(32)
COMP	1	MONOCULTURE	408 8088	34 1504	5 1031	37 2474	(12)
COMP	2	MIXTURE	623 2688	31 1835	5 3488	40 3088	(20)
SPAC	4	44 PLANTS M	1002 2288	37 1587	5 5810	50 8565	(27)
COMP	1	MONOCULTURE	412 4088	37 4818	5 7404	54 5835	(11)
COMP	2	MIXTURE	590 8188	35 8282	5 7418	78 4135	(16)
SPAC	5	28 PLANTS M	1354 2288	45 7010	5 1757	64 2480	(29)
COMP	1	MONOCULTURE	551 8288	47 1227	10 9438	119 7874	(15)
COMP	2	MIXTURE	802 4088	47 3207	7 1857	51 5773	(14)
SPAC	6	15 PLANTS M	1224 0388	47 0785	17 2348	300 4888	(25)
COMP	1	MONOCULTURE	822 8088	45 8823	15 8784	242 8252	(12)
COMP	2	MIXTURE	551 1888	45 4745	15 4354	377 7742	(12)
SPAC	7	10 PLANTS M	1188 5888	45 7150	14 5351	211 2685	(25)
COMP	1	MONOCULTURE	550 7288	43 1321	15 9084	275 4217	(12)
COMP	2	MIXTURE	638 8688	45 2889	12 2682	150 4888	(12)
CUL	4	700000002	4583 7381	35 7527	15 8020	248 7017	(121)
SPAC	1	204 PLANTS M	445 8088	22 4405	5 2118	27 1627	(20)
COMP	2	MIXTURE	445 8088	22 4405	5 2118	27 1627	(20)
SPAC	2	3 CM SPACING	577 3488	25 1022	5 4300	41 3443	(23)
COMP	2	MIXTURE	577 3488	25 1022	5 4300	41 3443	(23)

APPENDIX 6

03/16/82

PAGE 28

CRITERION VARIABLE WT

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
SPAC	3	8 CM SPACING	540 2888	32 0150	5 7804	55 2854	(20)
COMP	2	MIXTURE	540 2888	32 0150	5 7804	55 2854	(20)
SPAC	4	44 PLANTS M	554 8288	32 8371	5 0514	54 8251	(17)
COMP	2	MIXTURE	554 8288	32 8371	5 0514	54 8251	(17)
SPAC	5	28 PLANTS M	555 3288	45 0175	11 7882	138 3033	(19)
COMP	2	MIXTURE	555 3288	45 0175	11 7882	138 3033	(19)
SPAC	6	17 PLANTS M	539 2888	45 8700	14 8408	220 2480	(20)
COMP	2	MIXTURE	539 2888	45 8700	14 8408	220 2480	(20)
SPAC	7	10 PLANTS M	555 8088	55 5758	23 4897	552 2382	(12)
COMP	2	MIXTURE	555 8088	55 5758	23 4897	552 2382	(12)
CUL	5	NORQUAY	5570 4880	35 1528	22 5808	555 0454	(145)
SPAC	1	204 PLANTS M	547 8888	24 2180	7 1483	51 0984	(25)
COMP	1	MONOCULTURE	252 1288	23 8831	4 7380	22 4488	(15)
COMP	2	MIXTURE	455 4288	24 4553	5 5108	77 6302	(19)
SPAC	2	3 CM SPACING	534 2488	25 8295	5 1088	52 8324	(22)
COMP	1	MONOCULTURE	322 2188	33 2220	10 8547	117 8240	(10)
COMP	2	MIXTURE	202 0300	25 1882	5 4552	25 7700	(12)
SPAC	3	8 CM SPACING	554 4888	25 4885	7 4171	55 0121	(25)
COMP	1	MONOCULTURE	255 7288	25 8730	5 1800	55 0082	(10)
COMP	2	MIXTURE	295 7688	30 4438	5 4680	41 8088	(12)
SPAC	4	44 PLANTS M	1053 0888	40 4848	10 5544	112 9978	(27)
COMP	1	MONOCULTURE	475 1888	43 2988	12 0682	144 1688	(11)
COMP	2	MIXTURE	575 8888	34 5555	10 1055	102 1210	(18)
SPAC	5	28 PLANTS M	457 3488	35 9454	15 2885	255 3488	(12)
COMP	1	MONOCULTURE	457 3488	35 9454	15 2885	255 3488	(12)

APPENDIX 8			03/16/82				PAGE 29	
CRITERION VARIABLE WT								
VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N	
SPAC	8	17 PLANTS M	481 8400	57 7050	28 8408	878 8783	8	
COMP	1	MONOCULTURE	481 8400	57 7050	28 8408	878 8783	8	
SPAC	7	10 PLANTS M	1412 0788	74 3200	37 1387	1378 3188	10	
COMP	1	MONOCULTURE	888 1888	88 8411	28 8838	788 2814	8	
COMP	2	MIXTURE	818 8888	81 8810	43 8428	1831 8848	10	
CUL	8	700000002	8782 2880	81 8408	42 0288	1788 4887	120	
SPAC	1	204 PLANTS M	828 1188	34 4811	41 7818	1744 8488	10	
COMP	2	MIXTURE	828 1188	34 4811	41 7818	1744 8488	10	
SPAC	2	3 CM SPACING	1888 8887	88 8888	78 8848	8712 8373	20	
COMP	2	MIXTURE	1888 8887	88 8888	78 8848	8712 8373	20	
SPAC	3	8 CM SPACING	888 8888	38 8822	8 1282	37 4888	18	
COMP	2	MIXTURE	888 8888	38 8822	8 1282	37 4888	18	
SPAC	4	44 PLANTS M	878 8288	47 8182	28 1818	487 8871	12	
COMP	2	MIXTURE	878 8288	47 8182	28 1818	487 8871	12	
SPAC	5	28 PLANTS M	788 3188	44 4887	13 4887	178 7788	18	
COMP	2	MIXTURE	788 3188	44 4887	13 4887	178 7788	18	
SPAC	6	17 PLANTS M	1482 8288	88 4788	23 8883	1143 8828	22	
COMP	2	MIXTURE	1482 8288	88 4788	23 8883	1143 8828	22	
SPAC	7	10 PLANTS M	1884 8888	88 8882	27 1788	738 8878	20	
COMP	2	MIXTURE	1884 8888	88 8882	27 1788	738 8878	20	
CUL	7	888888888	3488 8888	88 8884	28 8874	737 8431	88	
SPAC	1	204 PLANTS M	231 1888	28 8888	4 8848	28 7431	8	
COMP	2	MIXTURE	231 1888	28 8888	4 8848	28 7431	8	
SPAC	2	3 CM SPACING	241 2888	28 4342	8 8138	74 1881	12	
COMP	1	MONOCULTURE	172 8888	28 7787	11 8218	128 7478	8	

APPENDIX 8		03/16/82					PAGE 30	
CRITERION VARIABLE WT								
VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE		
COMP	2	MIXTURE	188 8488	28 8817	8 8178	23 2888	81	
SPAC	3	8 CM SPACING	182 4888	38 4188	12 8848	181 1841	81	
COMP	1	MONOCULTURE	182 4888	38 4188	12 8848	181 1841	81	
SPAC	4	44 PLANTS M	313 7888	43 8287	8 8777	88 8884	71	
COMP	1	MONOCULTURE	313 7888	43 8287	8 8777	88 8884	71	
SPAC	8	28 PLANTS M	884 8888	78 8812	8 7318	48 3178	81	
COMP	2	MIXTURE	884 8888	78 8812	8 7318	48 3178	81	
SPAC	8	17 PLANTS M	848 8888	83 8887	21 7388	472 8881	18	
COMP	2	MIXTURE	848 8888	83 8887	21 7388	472 8881	18	
SPAC	7	10 PLANTS M	887 8888	78 8417	24 4317	1188 8417	12	
COMP	1	MONOCULTURE	482 8400	77 1887	28 8888	1888 8887	8	
COMP	2	MIXTURE	448 8888	74 1787	21 8348	1812 4388	8	
CUL	8	88781	8842 4288	82 4384	28 7188	881 4824	182	
SPAC	1	204 PLANTS M	788 7788	31 4712	4 3832	18 8881	28	
COMP	2	MIXTURE	788 7788	31 4712	4 3832	18 8881	28	
SPAC	2	3 CM SPACING	888 8888	38 8838	8 7348	48 3848	31	
COMP	2	MIXTURE	888 8888	38 8838	8 7348	48 3848	31	
SPAC	3	8 CM SPACING	818 8188	38 3888	8 8883	88 8888	88	
COMP	2	MIXTURE	318 8188	38 3288	8 8233	88 8888	24	
SPAC	4	44 PLANTS M	1438 8387	48 4887	11 2888	128 8831	31	
COMP	2	MIXTURE	1438 8387	48 4887	11 2888	128 8831	31	
SPAC	8	28 PLANTS M	1228 8888	87 8888	12 3888	183 4348	18	
COMP	2	MIXTURE	1228 8888	87 8888	12 3888	183 4348	18	
SPAC	8	17 PLANTS M	1844 8888	78 8828	28 8881	788 7731	24	
COMP	2	MIXTURE	1844 8888	78 8828	28 8881	788 7731	24	

APPENDIX 6 03/10/82 PAGE 21

CRITERION VARIABLE WT

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
SPAC	7	10 PLANTS M	2371 0200	81 7000	24 3021	884 8743	(20)
COMP	2	MIXTURE	2371 0300	81 7000	24 3021	884 8743	(20)
TOTAL CASES =	1000						
MISSING CASES =	35 OR	3 2 PET					

APPENDIX 6 03/10/82 PAGE 22

FILE WP77 (CREATION DATE = 03/10/82)

DESCRIPTION OF SUBPOPULATIONS

CRITERION VARIABLE	BY	DESCRIPTION	SEED YIELD PER PLANT g
BROKEN DOWN	BY	CUL	CULTIVAR
	BY	SPAC	INTER-PLANT SPACING
	BY	COMP	TREATMENT

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
FOR ENTIRE POPULATION			14804 4174	18 3026	12 8480	185 0650	(1025)
CUL	2	BLENDLEA	2820 4484	20 1638	18 2221	332 0818	(180)
SPAC	1	204 PLANTS M	177 2400	7 7081	8 8273	33 8878	(22)
COMP	1	MONOCULTURE	84 8600	7 7126	7 2169	62 8886	(11)
COMP	2	MIXTURE	92 3800	7 6992	4 6385	20 8798	(12)
SPAC	2	3 CM SPACING	271 7100	7 9916	3 1188	9 7288	(24)
COMP	1	MONOCULTURE	94 4800	7 2888	3 2433	10 8188	(12)
COMP	2	MIXTURE	177 2300	8 4390	3 0311	9 1878	(21)
SPAC	3	6 CM SPACING	361 2100	10 3203	4 1891	17 3817	(26)
COMP	1	MONOCULTURE	112 5800	8 8888	3 8728	14 8988	(13)
COMP	2	MIXTURE	248 6300	11 3023	4 1082	18 8823	(22)
SPAC	4	44 PLANTS M	486 0900	17 1638	8 3800	28 7300	(28)
COMP	1	MONOCULTURE	217 0400	16 8884	3 2172	10 3803	(13)
COMP	2	MIXTURE	278 8600	17 4266	8 7147	48 8870	(16)
SPAC	5	28 PLANTS M	810 1300	28 5070	9 8384	62 8482	(20)
COMP	1	MONOCULTURE	336 2000	22 3823	8 3744	70 1300	(16)
COMP	2	MIXTURE	174 8400	34 8880	8 8081	48 3801	(8)
SPAC	6	17 PLANTS M	781 3488	37 8878	18 4182	237 8888	(20)
COMP	1	MONOCULTURE	473 8888	38 4300	12 3887	181 8878	(13)
COMP	2	MIXTURE	277 7600	39 8880	20 8888	440 3888	(7)
SPAC	7	10 PLANTS M	1061 7600	55 8827	30 4837	418 8802	(19)
COMP	1	MONOCULTURE	382 3088	44 1637	18 3778	238 4884	(8)

APPENDIX B

03/10/83

PAGE 33

CRITERION VARIABLE Y

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
COMP	2	MIXTURE	706 4788	64 4073	20 6183	400 8516	11
CUL	3	PARK	1874 4887	9 8364	6 8680	33 3307	307
SPAC	1	204 PLANTS M	114 7000	4 8680	2 4793	6 1467	28
COMP	1	MONOCULTURE	24 7400	4 1233	2 2404	5 0193	8
COMP	2	MIXTURE	80 8800	4 7347	2 8887	8 7088	10
SPAC	2	3 CM SPACING	219 2000	8 3466	3 1306	9 8261	61
COMP	1	MONOCULTURE	81 2700	6 5193	2 7446	7 8223	14
COMP	2	MIXTURE	128 0300	4 7416	2 1883	10 2288	27
SPAC	3	8 CM SPACING	241 1400	7 8386	2 8738	8 6440	33
COMP	1	MONOCULTURE	100 0100	8 3343	2 8821	8 7188	12
COMP	2	MIXTURE	141 1300	7 0688	3 1487	9 8988	30
SPAC	4	44 PLANTS M	288 3400	10 2078	4 0380	16 2248	28
COMP	1	MONOCULTURE	104 3500	8 6683	3 8644	14 8880	11
COMP	2	MIXTURE	181 8900	10 7083	4 2008	17 6451	19
SPAC	8	28 PLANTS M	407 3088	14 0482	4 7748	22 7893	38
COMP	1	MONOCULTURE	198 8200	13 0847	6 4028	28 2088	18
COMP	2	MIXTURE	211 4800	15 1084	3 8134	16 2146	14
SPAC	6	17 PLANTS M	261 8288	13 8084	6 1801	37 8494	28
COMP	1	MONOCULTURE	200 2400	15 4083	8 7373	48 2884	13
COMP	2	MIXTURE	181 2000	12 4048	8 2777	28 8188	13
SPAC	7	10 PLANTS M	342 1288	13 1888	8 0888	36 7183	28
COMP	1	MONOCULTURE	187 2300	12 0848	7 0121	48 1883	13
COMP	2	MIXTURE	184 0800	14 3331	4 8888	24 8883	13
CUL	4	700000002	1318 4488	18 2084	7 2788	82 8784	188
SPAC	1	204 PLANTS M	84 8700	4 2288	1 8883	3 8384	20
COMP	2	MIXTURE	84 8700	4 2288	1 8883	3 8384	20
SPAC	2	3 CM SPACING	118 8800	8 4348	2 3782	8 8887	22
COMP	2	MIXTURE	118 8800	8 4348	2 3782	8 8887	22

APPENDIX B

03/10/83

PAGE 34

CRITERION VARIABLE Y

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
SPAC	3	8 CM SPACING	184 2000	8 2100	4 1278	17 0270	20
COMP	2	MIXTURE	784 2000	8 2100	4 1278	17 0270	20
SPAC	4	44 PLANTS M	181 7400	8 8288	3 3288	11 0808	17
COMP	2	MIXTURE	181 7400	8 8288	3 3288	11 0808	17
SPAC	8	28 PLANTS M	283 4400	13 8883	4 8844	20 8337	18
COMP	2	MIXTURE	283 4400	13 8883	4 8844	20 8337	18
SPAC	6	17 PLANTS M	318 8888	16 8888	7 3138	82 4877	20
COMP	2	MIXTURE	318 8888	16 8888	7 3138	82 4877	20
SPAC	7	10 PLANTS M	212 3700	19 3084	12 3884	183 7848	11
COMP	2	MIXTURE	212 3700	19 3084	12 3884	183 7848	11
CUL	8	NORWAY	1748 2887	12 0288	11 8010	124 8840	148
SPAC	1	204 PLANTS M	188 3400	8 3240	11 1880	11 1880	28
COMP	1	MONOCULTURE	82 3700	8 7731	2 3704	5 1646	18
COMP	2	MIXTURE	93 8700	4 8488	4 0878	16 4837	18
SPAC	2	3 CM SPACING	147 0100	7 0005	3 8880	16 8878	21
COMP	1	MONOCULTURE	81 8800	8 1888	4 8888	21 7888	8
COMP	2	MIXTURE	66 0300	8 4182	2 8188	8 8188	12
SPAC	3	8 CM SPACING	188 3800	7 1804	3 8883	7 8784	22
COMP	1	MONOCULTURE	70 8400	7 0840	3 1883	10 8888	18
COMP	2	MIXTURE	84 8400	7 2854	3 8880	8 7382	12
SPAC	4	44 PLANTS M	348 5188	12 8482	6 8284	31 8880	27
COMP	1	MONOCULTURE	188 1888	18 0186	6 1811	37 8388	11
COMP	2	MIXTURE	184 3800	11 8228	4 9228	24 2212	16
SPAC	8	28 PLANTS M	184 8100	12 7082	7 4881	88 0718	12
COMP	1	MONOCULTURE	184 8100	12 7082	7 4881	88 0718	12

APPENDIX 6

03/16/82

PAGE 38

CRITERION VARIABLE Y

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
SPAC	6	17 PLANTS M	163 6000	26 4750	16 6206	276 2466	61
COMP	1	MONOCULTURE	163 6000	26 4750	16 6206	276 2466	61
SPAC	7	10 PLANTS M	566 7200	26 8327	17 6116	320 6261	101
COMP	1	MONOCULTURE	223 2700	26 8169	13 3281	177 6206	61
COMP	2	MIXTURE	223 4600	23 6470	21 2800	462 6000	101
CUL	6	700000002	1923 0800	16 4028	14 6346	223 0407	1201
SPAC	1	204 PLANTS M	66 6600	4 9260	2 0799	4 3261	161
COMP	2	MIXTURE	66 6600	4 9260	2 0799	4 3261	161
SPAC	2	3 CM SPACING	121 0600	6 5023	2 4644	6 2221	221
COMP	2	MIXTURE	121 0600	6 5023	2 4644	6 2221	221
SPAC	3	6 CM SPACING	111 1200	6 6460	3 1230	6 7630	161
COMP	2	MIXTURE	111 1200	6 6460	3 1230	6 7630	161
SPAC	4	44 PLANTS M	176 3000	14 6417	6 6516	60 1366	121
COMP	2	MIXTURE	176 3000	14 6417	6 6516	60 1366	121
SPAC	6	26 PLANTS M	264 4200	14 1360	7 1222	60 7264	161
COMP	2	MIXTURE	264 4200	14 1360	7 1222	60 7264	161
SPAC	6	17 PLANTS M	666 6700	26 2716	16 4032	326 6700	221
COMP	2	MIXTURE	666 6700	26 2716	16 4032	326 6700	221
SPAC	7	10 PLANTS M	672 6600	23 6260	16 2646	264 6244	201
COMP	2	MIXTURE	672 6600	23 6260	16 2646	264 6244	201
CUL	7	BORQUAY	1066 7600	16 6040	12 3633	163 3673	661
SPAC	1	204 PLANTS M	41 6400	6 2426	1 6062	3 6414	61
COMP	2	MIXTURE	41 6400	6 2426	1 6062	3 6414	61
SPAC	2	3 CM SPACING	66 6300	7 6664	4 3166	16 6262	111
COMP	1	MONOCULTURE	66 7000	6 6600	6 4240	26 6200	61

APPENDIX 6

03/16/82

PAGE 39

CRITERION VARIABLE Y

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
COMP	2	MIXTURE	32 6200	6 6660	2 4042	6 7602	61
SPAC	3	6 CM SPACING	62 7000	10 4600	6 2206	36 6114	61
COMP	1	MONOCULTURE	62 7000	10 4600	6 2206	36 6114	61
SPAC	4	44 PLANTS M	62 6100	13 6017	6 6607	32 2706	61
COMP	1	MONOCULTURE	62 6100	13 6017	6 6607	32 2706	61
SPAC	6	26 PLANTS M	176 6000	26 6667	4 6736	24 7260	71
COMP	2	MIXTURE	176 6000	26 6667	4 6736	24 7260	71
SPAC	6	17 PLANTS M	307 6100	20 6060	6 3263	66 6606	161
COMP	2	MIXTURE	307 6100	20 6060	6 3263	66 6606	161
SPAC	7	10 PLANTS M	341 6500	26 4633	17 2737	301 6446	121
COMP	1	MONOCULTURE	166 7600	23 1317	20 3637	414 6602	61
COMP	2	MIXTURE	142 7700	23 7860	14 0263	197 6742	61
CUL	6	86701	3067 6100	16 6624	11 7613	136 6213	1611
SPAC	1	204 PLANTS M	167 6700	7 6146	2 2240	4 6462	261
COMP	2	MIXTURE	167 6700	7 6146	2 2240	4 6462	261
SPAC	2	3 CM SPACING	232 6000	7 2716	3 1641	6 6367	321
COMP	2	MIXTURE	232 6000	7 2716	3 1641	6 6367	321
SPAC	3	6 CM SPACING	261 6000	10 6760	4 1663	17 3630	241
COMP	2	MIXTURE	261 6000	10 6760	4 1663	17 3630	241
SPAC	4	44 PLANTS M	422 6200	14 4160	6 4060	26 2466	301
COMP	2	MIXTURE	422 6200	14 4160	6 4060	26 2466	301
SPAC	6	26 PLANTS M	440 6600	24 4026	6 6766	43 2662	161
COMP	2	MIXTURE	440 6600	24 4026	6 6766	43 2662	161
SPAC	6	17 PLANTS M	666 6600	26 6646	12 6021	166 4626	221
COMP	2	MIXTURE	666 6600	26 6646	12 6021	166 4626	221

APPENDIX B
CRITERION VARIABLE Y

03/16/82

PAGE 27

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
SPAC	7	10 PLANTS M	843 4888	28 0888	11 7048	127 0888	281
COMP	2	MIXTURE	843 4888	28 0888	11 7048	127 0888	281
TOTAL CASES =			1080				
MISSING CASES =			48 OR	4 Z PET			

APPENDIX B

03/16/82

PAGE 30

FILE WPT7 (CREATION DATE = 03/16/82)

CRITERION VARIABLE Y

DESCRPTION OF SUBPOPULATIONS
BROKEN DOWN BY
BY CUL
BY SPAC
BY COMP
NUMBER OF KERNELS PER PLANT
CULTIVAR
INTER-PLANT SPACING
TREATMENT

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
FOR ENTIRE POPULATION			351302 0000	380 0888	212 3884	87666 6161	10221
CUL	2	BLENDA	72328 0000	406 3820	377 8888	142883 8888	1781
SPAC	1	20 PLANTS M	3887 0000	184 8822	134 2100	18012 2281	231
COMP	1	MONOCULTURE	1783 0000	180 2727	174 4768	30441 8182	111
COMP	2	MIXTURE	1784 0000	148 5000	81 6478	6288 7273	121
SPAC	2	3 CM SPACING	5288 0000	195 8888	84 8710	3888 8207	341
COMP	1	MONOCULTURE	1848 0000	142 0000	87 8188	3242 8887	131
COMP	2	MIXTURE	3443 0000	183 8824	82 2747	2732 8476	211
SPAC	3	5 CM SPACING	7680 0000	202 5714	78 2817	5688 8403	381
COMP	1	MONOCULTURE	2281 0000	178 2368	73 3134	5374 8890	131
COMP	2	MIXTURE	4788 0000	218 1364	72 8441	5423 4687	221
SPAC	4	44 PLANTS M	8383 0000	334 0387	88 4888	8303 7384	281
COMP	1	MONOCULTURE	4488 0000	328 8882	88 3188	4887 1616	121
COMP	2	MIXTURE	4888 0000	328 8887	117 8888	13882 4888	181
SPAC	5	28 PLANTS M	10884 0000	834 2000	208 1432	42848 7474	201
COMP	1	MONOCULTURE	7048 0000	488 8887	178 0828	32878 8824	181
COMP	2	MIXTURE	3838 0000	727 8000	188 8828	28178 7000	81
SPAC	6	17 PLANTS M	14888 0000	787 8847	303 3721	82834 8888	191
COMP	1	MONOCULTURE	8888 0000	727 8187	242 8182	58881 1742	121
COMP	2	MIXTURE	5738 0000	818 2887	402 8834	16318 8848	71
SPAC	7	10 PLANTS M	21773 0000	1148 8474	428 1611	18188 1637	191
COMP	1	MONOCULTURE	8884 0000	870 8000	288 1518	87708 4288	81

APPENDIX 6

02/10/82

PAGE 30

CRITERION VARIABLE NR

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
COMP	2	MIXTURE	14800 0000	1240 2727	423 4040	170230 2162	111
CUL	3	PARK	50013 0000	270 5042	171 7000	29000 3200	207
SPAC	1	204 PLANTS M	2000 0000	110 0000	50 7324	2440 5000	20
COMP	1	MONOCULTURE	577 0000	112 5232	50 5700	2540 3007	51
COMP	2	MIXTURE	2210 0000	122 0000	50 5342	2602 1111	101
SPAC	2	3 CM SPACING	8930 0000	144 8241	60 2772	6400 4070	411
COMP	1	MONOCULTURE	2400 0000	170 1420	72 0232	5167 3020	141
COMP	2	MIXTURE	2400 0000	120 2000	60 4310	5022 7000	271
SPAC	3	6 CM SPACING	6700 0000	212 3700	82 1771	6702 0000	221
COMP	1	MONOCULTURE	2027 0000	244 7000	50 0000	5110 7000	121
COMP	2	MIXTURE	2000 0000	102 0000	72 5340	5200 6200	201
SPAC	4	44 PLANTS M	7000 0000	200 1700	124 0200	15007 0220	201
COMP	1	MONOCULTURE	2100 0000	200 0100	131 0410	17270 0020	111
COMP	2	MIXTURE	4000 0000	200 1170	124 0000	15020 7200	171
SPAC	5	20 PLANTS M	11070 0000	300 1724	127 1740	16172 2102	201
COMP	1	MONOCULTURE	5701 0000	300 0007	122 4372	17020 0201	101
COMP	2	MIXTURE	5070 0000	410 0420	122 7700	15072 2242	141
SPAC	6	17 PLANTS M	10047 0000	400 5000	200 2022	40100 0000	201
COMP	1	MONOCULTURE	5017 0000	422 0700	180 5120	20700 4102	121
COMP	2	MIXTURE	5030 0000	300 0231	221 0070	52702 7420	121
SPAC	7	10 PLANTS M	10000 0000	307 0007	107 0000	30007 3204	201
COMP	1	MONOCULTURE	4007 0000	300 0000	200 2207	42017 0007	121
COMP	2	MIXTURE	5017 0000	410 0022	100 1200	30700 7200	121
CUL		700000000	20001 0000	200 1702	104 2000	27707 2010	1001
SPAC		204 PLANTS M	2000 0000	117 0000	52 0400	2010 4737	201
COMP		MIXTURE	2200 0000	117 0000	52 0400	2010 4737	201
SPAC	2	3 CM SPACING	2172 0000	144 2272	61 0271	3707 0002	221
COMP	2	MIXTURE	2172 0000	144 2272	61 0271	3707 0002	221

APPENDIX 6

02/10/82

PAGE 40

CRITERION VARIABLE NR

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
SPAC	3	1 CM SPACING	4000 0000	227 0000	122 2400	14044 2737	201
COMP	2	MIXTURE	4000 0000	227 0000	122 2400	14044 2737	201
SPAC	4	44 PLANTS M	4430 0000	200 5002	117 3001	13772 2024	171
COMP	2	MIXTURE	4430 0000	200 5002	117 3001	13772 2024	171
SPAC	5	20 PLANTS M	7100 0000	372 0474	110 1070	12200 1027	101
COMP	2	MIXTURE	7100 0000	372 0474	110 1070	12200 1027	101
SPAC	6	17 PLANTS M	8702 0000	430 1000	174 4100	30422 2002	201
COMP	2	MIXTURE	8702 0000	430 1000	174 4100	30422 2002	201
SPAC	7	10 PLANTS M	6000 0000	300 0001	204 0122	02072 0000	111
COMP	2	MIXTURE	6000 0000	300 0001	204 0122	02072 0000	111
CUL	5	BROUQUAY	40440 0000	270 0370	201 7000	70300 7202	1401
SPAC	1	204 PLANTS M	4002 0000	110 0102	70 7020	5002 2100	201
COMP	1	MONOCULTURE	1000 0000	124 0070	40 7000	2377 0020	101
COMP	2	MIXTURE	2007 0000	110 2000	60 0207	0022 2007	101
SPAC	2	3 CM SPACING	3212 0000	102 0024	60 3200	7001 0470	211
COMP	1	MONOCULTURE	1702 0000	100 1111	107 0700	11000 2011	011
COMP	2	MIXTURE	1420 0000	110 2222	62 0744	2700 0070	121
SPAC	3	6 CM SPACING	3727 0000	102 0020	60 0020	3720 1344	221
COMP	1	MONOCULTURE	1024 0000	100 0000	67 0000	4000 0000	101
COMP	2	MIXTURE	2703 0000	100 0022	67 3020	2202 7200	121
SPAC	4	44 PLANTS M	5017 0000	211 7007	100 2070	27022 1000	271
COMP	1	MONOCULTURE	4002 0000	200 2727	200 0000	42000 0102	111
COMP	2	MIXTURE	4200 0000	272 1070	110 0072	14201 0000	101
SPAC	5	20 PLANTS M	3700 0000	210 0007	174 0020	30200 2424	121
COMP	1	MONOCULTURE	3700 0000	210 0007	174 0020	30200 2424	121

APPENDIX 6

03/10/82

PAGE 41

CRITERION VARIABLE NW

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
SPAC	6	17 PLANTS M	3730 0000	466 2500	400 7022	167680 0200	81
COMP	1	MONOCULTURE	3730 0000	466 2500	400 7022	167680 0200	81
SPAC	7	10 PLANTS M	13400 0000	700 4737	431 8183	186404 4884	191
COMP	1	MONOCULTURE	6525 0000	612 8689	339 8988	115306 1111	81
COMP	2	MIXTURE	7005 0000	700 5000	602 8874	362006 2776	101
CUL	6	700000002	47021 0000	370 2441	288 0327	128004 4884	127
SPAC	1	204 PLANTS M	2010 0000	111 8887	48 8480	2404 7080	181
COMP	2	MIXTURE	2010 0000	111 8887	48 8480	2404 7080	181
SPAC	3	3 CM SPACING	2000 0000	121 8182	88 8094	3001 2887	221
COMP	2	MIXTURE	2000 0000	121 8182	88 8094	3001 2887	221
SPAC	3	3 CM SPACING	2000 0000	121 8182	88 8094	3001 2887	221
COMP	2	MIXTURE	2000 0000	121 8182	88 8094	3001 2887	221
SPAC	4	44 PLANTS M	4007 0000	340 8833	170 7302	29148 8108	121
COMP	2	MIXTURE	4007 0000	340 8833	170 7302	29148 8108	121
SPAC	6	26 PLANTS M	6407 0000	360 8444	187 8260	35018 8261	181
COMP	2	MIXTURE	6407 0000	360 8444	187 8260	35018 8261	181
SPAC	6	17 PLANTS M	13111 0000	624 3233	428 7468	183823 0233	211
COMP	2	MIXTURE	13111 0000	624 3233	428 7468	183823 0233	211
SPAC	7	10 PLANTS M	18001 0000	604 0500	398 0818	158680 8831	201
COMP	2	MIXTURE	18001 0000	604 0500	398 0818	158680 8831	201
CUL	7	800000000	20028 0000	450 8482	318 8428	101861 3822	881
SPAC	1	204 PLANTS M	1110 0000	120 8000	88 8200	4428 7142	91
COMP	2	MIXTURE	1110 0000	120 8000	88 8200	4428 7142	91
SPAC	3	3 CM SPACING	2334 0000	212 1818	128 1738	16018 8838	111
COMP	1	MONOCULTURE	1677 0000	246 1887	167 8881	24082 8887	81

APPENDIX 6

03/10/82

PAGE 42

CRITERION VARIABLE NW

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
COMP	2	MIXTURE	867 0000	171 4000	88 2883	4706 3000	51
SPAC	3	3 CM SPACING	1731 0000	268 5000	182 7682	33337 8000	81
COMP	1	MONOCULTURE	1731 0000	268 5000	182 7682	33337 8000	81
SPAC	4	44 PLANTS M	2310 0000	365 0000	188 8770	24810 4000	81
COMP	1	MONOCULTURE	2310 0000	365 0000	188 8770	24810 4000	81
SPAC	6	26 PLANTS M	4872 0000	607 4286	140 1814	19882 8100	71
COMP	2	MIXTURE	4872 0000	607 4286	140 1814	19882 8100	71
SPAC	6	17 PLANTS M	8481 0000	683 8000	232 8428	84122 8428	181
COMP	2	MIXTURE	8481 0000	683 8000	232 8428	84122 8428	181
SPAC	7	10 PLANTS M	9211 0000	767 8833	418 2190	172408 8108	121
COMP	1	MONOCULTURE	6107 0000	661 1887	473 4178	224124 1887	121
COMP	2	MIXTURE	6104 0000	664 0000	372 0268	138404 0000	81
CUL	6	80701	70200 0000	437 8000	371 4780	137886 8874	181
SPAC	1	204 PLANTS M	4447 0000	177 8800	81 8888	2702 8422	281
COMP	2	MIXTURE	4447 0000	177 8800	81 8888	2702 8422	281
SPAC	2	3 CM SPACING	5420 0000	188 8882	71 8888	8183 7812	321
COMP	2	MIXTURE	5420 0000	188 8882	71 8888	8183 7812	321
SPAC	3	3 CM SPACING	6082 0000	282 8260	100 8884	10123 7228	241
COMP	2	MIXTURE	6082 0000	282 8260	100 8884	10123 7228	241
SPAC	4	44 PLANTS M	10837 0000	381 3233	122 2888	17787 8402	201
COMP	2	MIXTURE	10837 0000	381 3233	122 2888	17787 8402	201
SPAC	6	26 PLANTS M	11140 0000	619 3888	217 8887	47383 8488	181
COMP	2	MIXTURE	11140 0000	619 3888	217 8887	47383 8488	181
SPAC	6	17 PLANTS M	17181 0000	747 4248	378 1888	143006 3478	221
COMP	2	MIXTURE	17181 0000	747 4248	378 1888	143006 3478	221

APPENDIX 6

02/10/82

PAGE 43

CRITERION VARIABLE ON

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
SPAC	7	10 PLANTS M	24444 0000	842 0000	611 3027	201431 4832	201
COMP	2	MIXTURE	24444 0000	842 0000	611 3027	201431 4832	201
TOTAL CASES *	1000						
MISSING CASES *	40 00	4 4 PCT					

APPENDIX 6

02/10/82

PAGE 44

FILE WPTT (CREATION DATE = 02/10/82)

CRITERION VARIABLE		DESCRIPTION OF SUBPOPULATIONS					
BROKEN DOWN BY		BY		EXTENSION LENGTH CM			
				CULTIVAR			
				INTER-PLANT SPACING			
				TREATMENT			
VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
FOR ENTIRE POPULATION			61344 0000	66 5344	12 5076	155 4055	1000
CUL	2	6LX18A	13120 0000	72 0670	9 1930	83 7022	182
SPAC	1	204 PLANTS M	1782 0000	73 4167	9 0025	81 0707	24
COMP	1	MONOCULTURE	817 0000	74 2727	7 0132	62 5182	11
COMP	2	MIXTURE	965 0000	72 0022	11 6002	134 5841	12
SPAC	2	3 CM SPACING	2682 0000	73 7714	8 0040	64 0620	35
COMP	1	MONOCULTURE	1016 0000	72 8714	11 8127	134 8781	14
COMP	2	MIXTURE	1666 0000	75 8714	4 6223	19 5671	21
SPAC	3	6 CM SPACING	2634 0000	72 4000	10 0442	110 7785	35
COMP	1	MONOCULTURE	943 0000	72 5355	12 8114	165 2682	12
COMP	2	MIXTURE	1691 0000	72 3152	9 3828	86 0368	23
SPAC	4	44 PLANTS M	2064 0000	71 0021	10 2120	100 2974	20
COMP	1	MONOCULTURE	857 0000	74 2444	10 5794	111 0231	12
COMP	2	MIXTURE	1207 0000	69 0185	9 0600	80 0050	18
SPAC	5	28 PLANTS M	1418 0000	70 0000	7 0016	55 2727	20
COMP	1	MONOCULTURE	1005 0000	71 0000	7 2851	52 7010	15
COMP	2	MIXTURE	413 0000	70 0000	9 0277	81 0000	5
SPAC	6	17 PLANTS M	1424 0000	71 2000	6 7715	45 8526	20
COMP	1	MONOCULTURE	920 0000	72 2200	6 7842	46 0265	12
COMP	2	MIXTURE	504 0000	69 2557	6 6242	44 5714	7
SPAC	7	10 PLANTS M	1218 0000	69 3664	6 6394	44 1245	18
COMP	1	MONOCULTURE	857 0000	70 8750	6 0612	36 1200	8

APPENDIX 8

02/10/82

PAGE 48

CRITERION VARIABLE EL

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
COMP	2	MIXTURE	781 0000	66 2727	8 0232	61 4182	11
CUL	3	PARK	11207 0000	64 8232	6 8867	47 0618	207
SPAC	1	204 PLANTS M	1444 0000	67 7600	4 8801	23 8232	20
COMP	1	MONOCULTURE	381 0000	68 8000	3 7283	13 8000	8
COMP	2	MIXTURE	1063 0000	67 8283	6 2214	37 2832	10
SPAC	3	3 CM SPACING	2267 0000	66 0483	7 6880	68 7678	41
COMP	1	MONOCULTURE	748 0000	66 3871	7 8218	66 0824	14
COMP	2	MIXTURE	1488 0000	64 3704	7 7487	66 0114	27
SPAC	3	8 CM SPACING	1882 0000	67 6786	6 8088	48 3710	32
COMP	1	MONOCULTURE	710 0000	68 1887	5 0422	28 4242	12
COMP	2	MIXTURE	1142 0000	67 1000	7 8876	68 2828	20
SPAC	4	44 PLANTS M	1808 0000	63 8838	4 7848	22 8881	20
COMP	1	MONOCULTURE	808 0000	68 8000	5 8784	28 8000	11
COMP	2	MIXTURE	984 0000	63 1788	4 8130	21 2784	17
SPAC	6	28 PLANTS M	1884 0000	63 8882	5 2812	27 6788	20
COMP	1	MONOCULTURE	888 0000	63 8332	5 4872	28 7810	18
COMP	2	MIXTURE	748 0000	63 2142	6 2208	27 2882	14
SPAC	8	17 PLANTS M	1248 0000	61 7388	6 2828	68 8048	20
COMP	1	MONOCULTURE	888 0000	68 7882	6 4474	71 2888	12
COMP	2	MIXTURE	888 0000	62 8822	6 2408	68 8041	12
SPAC	7	10 PLANTS M	1248 0000	61 7882	6 8877	48 2448	20
COMP	1	MONOCULTURE	872 0000	61 7882	6 3882	68 8888	12
COMP	2	MIXTURE	872 0000	61 7882	6 1888	28 8822	12
CUL	4	7000000000	8827 0000	61 2748	6 0838	62 1847	181
SPAC	1	204 PLANTS M	1212 0000	68 8888	6 2888	67 8878	20
COMP	2	MIXTURE	1212 0000	68 8888	6 2888	67 8878	20
SPAC	3	3 CM SPACING	1822 0000	68 2174	6 0481	64 7222	22
COMP	2	MIXTURE	1822 0000	68 2174	6 0481	64 7222	22

APPENDIX 8

02/10/82

PAGE 48

CRITERION VARIABLE EL

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
SPAC	3	8 CM SPACING	1224 0000	61 4000	11 8117	138 8188	20
COMP	2	MIXTURE	1224 0000	61 4000	11 8117	138 8188	20
SPAC	4	44 PLANTS M	1082 0000	62 8284	11 0120	121 2847	17
COMP	2	MIXTURE	1082 0000	62 8284	11 0120	121 2847	17
SPAC	6	28 PLANTS M	1188 0000	62 4211	5 8378	38 2873	18
COMP	2	MIXTURE	1188 0000	62 4211	5 8378	38 2873	18
SPAC	6	17 PLANTS M	1171 0000	68 8888	6 8812	32 8888	20
COMP	2	MIXTURE	1171 0000	68 8888	6 8812	32 8888	20
SPAC	7	10 PLANTS M	842 0000	63 8832	6 0174	28 1742	12
COMP	2	MIXTURE	842 0000	63 8832	6 0174	28 1742	12
CUL	6	MONOCULTURE	8411 0000	44 2138	6 8878	68 8888	181
SPAC	1	204 PLANTS M	1884 0000	44 4000	7 8788	63 8888	20
COMP	1	MONOCULTURE	784 0000	44 8000	4 7748	22 8000	18
COMP	2	MIXTURE	488 0000	44 7388	10 8488	100 8828	18
SPAC	3	3 CM SPACING	1831 0000	46 8638	11 0882	123 1710	22
COMP	1	MONOCULTURE	488 0000	48 8000	10 8824	118 8888	18
COMP	2	MIXTURE	888 0000	48 2888	11 8788	128 7888	12
SPAC	3	8 CM SPACING	1822 0000	44 4248	11 2182	128 8072	22
COMP	1	MONOCULTURE	288 0000	38 8000	7 8277	48 2888	18
COMP	2	MIXTURE	888 0000	48 2388	12 8878	187 8822	12
SPAC	4	44 PLANTS M	1181 0000	48 8877	8 2187	68 8818	28
COMP	1	MONOCULTURE	422 0000	42 2000	5 4732	28 8888	18
COMP	2	MIXTURE	788 0000	48 8828	10 8111	112 8888	18
SPAC	6	28 PLANTS M	488 0000	48 8000	4 7801	22 8888	12
COMP	1	MONOCULTURE	488 0000	48 8000	4 7801	22 8888	12

APPENDIX B

02/16/82

PAGE 47

CRITERION VARIABLE EL

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
SPAC	5	17 PLANTS H	306 0000	36 3600	6 5192	42 0000	81
COMP	1	MONOCULTURE	306 0000	36 3600	6 5192	42 0000	81
SPAC	7	10 PLANTS H	821 0000	43 3105	6 3085	40 0022	191
COMP	1	MONOCULTURE	276 0000	41 0007	2 9500	8 7000	81
COMP	2	MIXTURE	446 0000	44 6000	6 3427	40 0000	191
CUL	6	700000002	8765 0000	61 7020	6 3288	38 7867	1311
SPAC	1	204 PLANTS H	876 0000	64 1857	6 2194	67 0000	161
COMP	2	MIXTURE	876 0000	64 1857	6 2194	67 0000	161
SPAC	2	3 CM SPACING	1126 0000	61 6600	7 1767	51 4012	221
COMP	2	MIXTURE	1126 0000	61 6600	7 1767	51 4012	221
SPAC	3	6 CM SPACING	960 0000	63 2774	4 7370	22 4477	161
COMP	2	MIXTURE	960 0000	63 2774	4 7370	22 4477	161
SPAC	4	44 PLANTS H	686 0000	67 0833	4 2310	17 0016	121
COMP	2	MIXTURE	686 0000	67 0833	4 2310	17 0016	121
SPAC	5	26 PLANTS H	600 0000	46 2774	4 1641	17 0000	161
COMP	2	MIXTURE	600 0000	46 2774	4 1641	17 0000	161
SPAC	6	17 PLANTS H	1004 0000	49 7272	6 6321	31 7218	221
COMP	2	MIXTURE	1004 0000	49 7272	6 6321	31 7218	221
SPAC	7	10 PLANTS H	1000 0000	60 6571	4 7664	22 0200	211
COMP	2	MIXTURE	1000 0000	60 6571	4 7664	22 0200	211
CUL	7	RD004	3016 0000	46 6362	6 7612	22 0780	601
SPAC	1	204 PLANTS H	482 0000	61 3232	6 0416	36 0000	81
COMP	2	MIXTURE	482 0000	61 3232	6 0416	36 0000	81
SPAC	2	3 CM SPACING	612 0000	61 0000	4 4721	20 0000	121
COMP	1	MONOCULTURE	311 0000	61 6222	3 6200	16 3607	81

APPENDIX B

02/16/82

PAGE 48

CRITERION VARIABLE EL

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
COMP	2	MIXTURE	301 0000	60 1667	6 1020	26 0007	81
SPAC	3	6 CM SPACING	266 0000	46 0000	3 6322	13 2000	61
COMP	1	MONOCULTURE	266 0000	46 0000	3 6322	13 2000	61
SPAC	4	44 PLANTS H	360 0000	61 2857	7 1348	50 0000	71
COMP	1	MONOCULTURE	360 0000	61 2857	7 1348	50 0000	71
SPAC	5	26 PLANTS H	611 0000	61 3750	6 0126	26 1260	81
COMP	2	MIXTURE	611 0000	61 3750	6 0126	26 1260	81
SPAC	6	17 PLANTS H	722 0000	48 8667	6 6416	24 1224	161
COMP	2	MIXTURE	722 0000	48 8667	6 6416	24 1224	161
SPAC	7	10 PLANTS H	662 0000	46 0833	6 3666	40 4470	121
COMP	1	MONOCULTURE	276 0000	46 3232	6 7726	46 0007	81
COMP	2	MIXTURE	276 0000	46 3232	6 6666	42 0007	81
CUL	8	RD701	12276 0000	67 0820	6 6670	62 6669	1621
SPAC	1	204 PLANTS H	1640 0000	73 6600	7 6164	66 0400	261
COMP	2	MIXTURE	1640 0000	73 6600	7 6164	66 0400	261
SPAC	2	3 CM SPACING	2222 0000	72 6626	6 6766	72 6626	321
COMP	2	MIXTURE	2222 0000	72 6626	6 6766	72 6626	321
SPAC	3	6 CM SPACING	1661 0000	66 2000	6 2666	66 1721	241
COMP	2	MIXTURE	1661 0000	66 2000	6 2666	66 1721	241
SPAC	4	44 PLANTS H	2064 0000	66 6600	12 6300	164 7600	311
COMP	2	MIXTURE	2064 0000	66 6600	12 6300	164 7600	311
SPAC	5	26 PLANTS H	1164 0000	66 7774	6 6666	26 6201	161
COMP	2	MIXTURE	1164 0000	66 7774	6 6666	26 6201	161
SPAC	6	17 PLANTS H	1461 0000	61 7063	6 6717	24 4764	261
COMP	2	MIXTURE	1461 0000	61 7063	6 6717	24 4764	261

APPENDIX 8

03/16/82

PAGE 48

CRITERION VARIABLE EL

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
SPAC	7	30 PLANTS M	1716 0000	60 1034	4 6386	23 3818	28
COMP	2	MIXTURE	1716 0000	60 1034	4 6386	23 3818	28
TOTAL CASES = 1080							
MISSING CASES = 32 OR 3.0 PCT							

APPENDIX 8

03/16/82

PAGE 50

FILE WPT7

(CREATION DATE = 03/16/82)

DESCRIPTION OF SUBPOPULATIONS

CRITERION VARIABLE BROKEN DOWN BY	KN CUL BY SPAC BY COMP	DESCRIPTION CULTIVAR INTER-PLANT SPACING TREATMENT	SUM	MEAN	STD DEV	VARIANCE	N
FOR ENTIRE POPULATION			48886 7801	47 2272	38 0484	802 7881	1080
CUL	2	SLHLEA	7881 8181	43 1848	17 4313	303 8818	178
SPAC	1	304 PLANTS M	821 8777	38 7338	10 8788	120 8381	33
COMP	1	MONOCULTURE	380 3187	34 8742	14 8880	218 8784	11
COMP	2	MIXTURE	441 8611	36 7888	8 8124	62 8118	12
SPAC	2	3 CM SPACING	1200 7281	36 3188	7 8348	61 3888	34
COMP	1	MONOCULTURE	488 7738	31 8211	7 8838	61 8781	13
COMP	2	MIXTURE	788 8523	37 8844	7 8888	68 1184	21
SPAC	3	8 CM SPACING	1301 4780	37 1880	8 2888	64 7888	38
COMP	1	MONOCULTURE	423 8904	32 8484	8 8388	38 2482	13
COMP	2	MIXTURE	878 3888	39 8788	8 7888	68 8334	22
SPAC	4	44 PLANTS M	1173 8888	41 8888	8 8482	78 1888	38
COMP	1	MONOCULTURE	818 1222	38 8888	8 8788	8 8788	12
COMP	2	MIXTURE	888 8888	43 7218	11 8888	138 1888	18
SPAC	8	28 PLANTS M	818 8882	48 7888	8 3218	88 8818	38
COMP	1	MONOCULTURE	881 8882	48 8872	8 8784	78 7734	18
COMP	2	MIXTURE	234 8880	48 8880	7 1788	61 8224	8
SPAC	8	17 PLANTS M	1078 4884	68 7828	18 8882	348 3873	18
COMP	1	MONOCULTURE	887 3882	48 8480	12 1387	147 3880	12
COMP	2	MIXTURE	481 1071	70 1582	21 8888	483 1220	7
SPAC	7	10 PLANTS M	1188 8837	62 8838	34 1888	1188 8233	18
COMP	1	MONOCULTURE	382 3312	44 8414	14 3882	208 1881	8

APPENDIX B

02/10/82

PAGE 81

CRITERION VARIABLE KM

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
COMP	2	MIXTURE	830 7826	76 0882	28 8478	1488 8088	11
CUL	3	PARK	8871 1043	41 8873	18 4878	237 3888	208
SPAC	1	204 PLANTS M	723 2727	28 2848	8 8888	73 2878	28
COMP	1	MONOCULTURE	183 8187	27 1828	8 7888	23 8887	8
COMP	2	MIXTURE	878 8878	38 8821	8 2882	28 8188	18
SPAC	3	3 CM SPACING	1188 7234	28 7432	8 8882	73 1888	48
COMP	1	MONOCULTURE	418 8187	31 8821	7 2812	84 8827	12
COMP	2	MIXTURE	773 8227	28 8881	8 8888	88 7823	27
SPAC	3	8 CM SPACING	1288 8828	48 2178	12 2882	181 2888	22
COMP	1	MONOCULTURE	471 2182	38 2888	8 2828	38 8871	12
COMP	2	MIXTURE	818 7888	48 7873	18 8428	223 2828	28
SPAC	4	44 PLANTS M	1181 1223	42 8888	7 8888	81 2887	28
COMP	1	MONOCULTURE	484 2273	42 2824	8 1781	38 1481	12
COMP	2	MIXTURE	728 8881	42 7888	8 8188	78 8788	12
SPAC	5	28 PLANTS M	1288 8783	47 8188	18 8878	112 8281	22
COMP	1	MONOCULTURE	718 8824	47 2888	18 8827	188 2888	18
COMP	2	MIXTURE	878 8188	47 8882	11 1214	123 8878	18
SPAC	6	17 PLANTS M	1282 8821	52 4888	18 8888	288 1788	28
COMP	1	MONOCULTURE	878 8418	52 2724	18 8888	118 8888	12
COMP	2	MIXTURE	882 8882	52 8288	28 8728	423 2722	12
SPAC	7	18 PLANTS M	1428 4828	54 8847	18 8828	288 1887	28
COMP	1	MONOCULTURE	771 4822	58 2188	18 8728	128 4828	12
COMP	2	MIXTURE	888 2288	58 4888	28 8887	888 2888	12
CUL	4	700000002	8188 8818	48 2748	18 7844	248 2818	128
SPAC	1	204 PLANTS M	848 1842	27 4882	8 4147	88 8288	28
COMP	2	MIXTURE	848 1842	27 4882	8 4147	88 8288	28
SPAC	3	3 CM SPACING	783 1878	38 8844	8 8288	88 8188	22
COMP	2	MIXTURE	783 1878	38 8844	8 8288	88 8188	22

APPENDIX B

02/10/82

PAGE 82

CRITERION VARIABLE KM

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
SPAC	2	8 CM SPACING	728 7881	38 2278	12 8288	184 8824	28
COMP	2	MIXTURE	728 7881	38 2278	12 8288	184 8824	28
SPAC	4	44 PLANTS M	778 2882	48 8884	21 8878	488 1722	17
COMP	2	MIXTURE	778 2882	48 8884	21 8878	488 1722	17
SPAC	6	28 PLANTS M	888 2887	48 8428	18 8828	117 7887	18
COMP	2	MIXTURE	888 2887	48 8428	18 8828	117 7887	18
SPAC	8	17 PLANTS M	1082 8282	58 1818	18 8888	288 2482	28
COMP	2	MIXTURE	1082 8282	58 1818	18 8888	288 2482	28
SPAC	7	18 PLANTS M	488 8888	41 8888	8 8781	73 8228	11
COMP	2	MIXTURE	488 8888	41 8888	8 8781	73 8228	11
CUL	5	WORMHAY	8882 2841	48 8788	18 8821	288 2821	144
SPAC	1	204 PLANTS M	887 8888	38 4877	8 2878	78 8184	28
COMP	1	MONOCULTURE	487 2888	31 8782	8 8848	44 4182	18
COMP	2	MIXTURE	488 8823	28 2884	8 2288	88 1812	18
SPAC	2	3 CM SPACING	882 3823	33 1181	11 4121	138 2282	28
COMP	1	MONOCULTURE	322 4788	37 8822	11 1281	123 8781	8
COMP	2	MIXTURE	328 8238	28 8821	11 8888	122 8884	11
SPAC	3	8 CM SPACING	828 2818	38 8178	8 2718	78 8888	22
COMP	1	MONOCULTURE	388 2288	38 8828	7 8882	88 2188	18
COMP	2	MIXTURE	482 8888	38 8828	8 1448	83 8128	12
SPAC	4	44 PLANTS M	1088 7248	48 8184	14 8788	187 8888	27
COMP	1	MONOCULTURE	481 1217	43 7282	14 8272	223 1211	11
COMP	2	MIXTURE	818 8823	28 4781	12 8884	182 2872	18
SPAC	5	28 PLANTS M	888 1882	48 8888	12 8782	171 8842	12
COMP	1	MONOCULTURE	888 1882	48 8888	12 8782	171 8842	12

204

APPENDIX 8 03/16/82 PAGE 53

CRITERION VARIABLE KH

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
SPAC	6	17 PLANTS M	413 3508	51 8868	13 4008	178 8738	81
COMP	1	MONOCULTURE	413 3508	51 8868	13 4008	178 8738	81
SPAC	7	10 PLANTS M	1288 3213	87 8480	33 6088	1128 8200	181
COMP	1	MONOCULTURE	488 8718	55 8080	10 3888	107 8864	81
COMP	2	MIXTURE	788 7884	78 8788	43 3870	1888 3874	181
CUL	6	700000003	8878 4120	54 1807	27 8887	1813 8882	1271
SPAC	1	204 PLANTS M	887 2833	23 1824	18 1888	103 1848	181
COMP	2	MIXTURE	887 2833	23 1824	18 1888	103 1848	181
SPAC	2	3 CM SPACING	788 2288	31 8288	12 2888	188 8188	221
COMP	2	MIXTURE	788 2288	31 8288	12 2888	188 8188	221
SPAC	3	8 CM SPACING	848 1880	40 3283	18 8801	284 8387	181
COMP	2	MIXTURE	848 1880	40 3283	18 8801	284 8387	181
SPAC	4	44 PLANTS M	888 8874	60 8488	10 8821	114 3218	121
COMP	2	MIXTURE	888 8874	60 8488	10 8821	114 3218	121
SPAC	8	28 PLANTS M	1871 8321	58 8818	28 4828	781 3281	181
COMP	2	MIXTURE	1871 8321	58 8818	28 4828	781 3281	181
SPAC	8	17 PLANTS M	1888 1882	58 8812	44 2788	1888 8712	211
COMP	2	MIXTURE	1888 1882	58 8812	44 2788	1888 8712	211
SPAC	7	10 PLANTS M	1787 8873	88 2888	87 8211	3384 8888	281
COMP	2	MIXTURE	1787 8873	88 2888	87 8211	3384 8888	281
CUL	7	NORWAY	2788 2880	87 8188	48 8828	1884 2871	881
SPAC	1	204 PLANTS M	248 8833	38 7884	8 3488	88 7181	81
COMP	2	MIXTURE	248 8833	38 7884	8 3488	88 7181	81
SPAC	2	3 CM SPACING	438 8373	38 8843	12 1827	147 7128	111
COMP	1	MONOCULTURE	278 7883	48 1181	11 8880	188 8178	81

APPENDIX 8 03/16/82 PAGE 54

CRITERION VARIABLE KH

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
COMP	2	MIXTURE	188 8180	31 8838	7 3428	53 8133	81
SPAC	3	8 CM SPACING	338 8278	58 8880	18 8888	243 3880	81
COMP	1	MONOCULTURE	338 8278	58 8880	18 8888	243 3880	81
SPAC	4	44 PLANTS M	328 4288	54 7378	18 2788	223 4748	81
COMP	1	MONOCULTURE	328 4288	54 7378	18 2788	223 4748	81
SPAC	8	28 PLANTS M	388 8833	58 8878	17 3478	308 8488	71
COMP	2	MIXTURE	388 8833	58 8878	17 3478	308 8488	71
SPAC	8	17 PLANTS M	1888 8884	70 8888	88 8880	4287 4813	181
COMP	2	MIXTURE	1888 8884	70 8888	88 8880	4287 4813	181
SPAC	7	10 PLANTS M	888 8888	78 8878	48 8732	1878 8888	121
COMP	1	MONOCULTURE	888 8888	83 8280	48 8734	2288 4871	81
COMP	2	MIXTURE	388 3888	68 8800	31 8837	1018 2882	81
CUL	8	88781	18888 8182	88 2701	48 8441	2203 7482	1881
SPAC	1	204 PLANTS M	1888 8888	43 8840	7 1817	51 2801	281
COMP	2	MIXTURE	1888 8888	43 8840	7 1817	51 2801	281
SPAC	2	3 CM SPACING	1281 4118	41 8884	8 8718	73 4787	311
COMP	2	MIXTURE	1281 4118	41 8884	8 8718	73 4787	311
SPAC	3	8 CM SPACING	1188 2287	48 8812	7 8878	82 3788	281
COMP	2	MIXTURE	1188 2287	48 8812	7 8878	82 3788	281
SPAC	4	44 PLANTS M	1883 7380	62 1248	17 8118	308 8880	281
COMP	2	MIXTURE	1883 7380	62 1248	17 8118	308 8880	281
SPAC	8	28 PLANTS M	1828 4320	87 8240	14 1838	208 8888	181
COMP	2	MIXTURE	1828 4320	87 8240	14 1838	208 8888	181
SPAC	8	17 PLANTS M	1788 7707	73 8488	28 8888	1288 8888	221
COMP	2	MIXTURE	1788 7707	73 8488	28 8888	1288 8888	221

APPENDIX 8 02/16/82 PAGE 59

CRITERION VARIABLE KM

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
SPAC	7	10 PLANTS M	2881 9261	88 8678	88 8687	8821 2822	(28)
COMP	2	MIXTURE	2881 9261	88 8678	88 8687	8821 2822	(28)
TOTAL CASES =	1080						
MISSING CASES =	81 BR	4.7 PCT					

APPENDIX 8 02/16/82 PAGE 60

FILE WP77 (CREATION DATE = 02/16/82)

DESCRIPTION OF SUBPOPULATIONS

CRITERION VARIABLE	VM	SEED YIELD PER HEAD
BROKEN DOWN BY	CUL	CULTIVAR
BY	SPAC	INTER-PLANT SPACING
BY	COMP	TREATMENT

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
FOR ENTIRE POPULATION			1868 0893	1 8071	1 0328	1 0662	(1822)
CU	2	81ERLEA	388 0632	2 1392	0 8822	0 7782	(180)
SPAC	1	208 PLANTS M	41 8094	1 8178	0 4814	0 2317	(23)
COMP	1	MONOCULTURE	18 8426	1 7221	0 5848	0 3420	(11)
COMP	2	MIXTURE	22 8669	1 9058	0 3873	0 1349	(12)
SPAC	2	3 CM SPACING	58 8771	1 7808	0 4431	0 1963	(34)
COMP	1	MONOCULTURE	40 8386	1 5874	0 4884	0 2178	(13)
COMP	2	MIXTURE	40 8415	1 8183	0 3872	0 1498	(21)
SPAC	3	9 CM SPACING	65 8227	1 8748	0 5290	0 2789	(35)
COMP	1	MONOCULTURE	20 8213	1 5786	0 4082	0 1688	(13)
COMP	2	MIXTURE	45 1013	2 0801	0 5210	0 2714	(22)
SPAC	4	44 PLANTS M	80 3123	2 0787	0 5411	0 2928	(29)
COMP	1	MONOCULTURE	26 8183	1 8705	0 2055	0 0422	(13)
COMP	2	MIXTURE	34 8880	2 1625	0 7028	0 4940	(18)
SPAC	5	22 PLANTS M	43 8273	2 1914	0 4073	0 1689	(20)
COMP	1	MONOCULTURE	22 8069	2 1871	0 4180	0 1758	(18)
COMP	2	MIXTURE	11 3214	2 2843	0 4088	0 1648	(8)
SPAC	6	17 PLANTS M	54 8802	2 7280	0 8471	0 8871	(28)
COMP	1	MONOCULTURE	31 8894	2 4223	0 6262	0 3921	(13)
COMP	2	MIXTURE	23 8898	3 2887	1 3188	1 4783	(7)
SPAC	7	10 PLANTS M	55 0442	3 0680	1 8421	2 8884	(18)
COMP	1	MONOCULTURE	17 8888	2 2381	0 7644	0 5881	(8)

APPENDIX 6

02/16/82

PAGE 87

CRITERION VARIABLE YH

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
COMP	2	MIXTURE	40 1855	3 6908	1 8784	3 5206	11
CUL	3	PARK	304 1484	1 4768	0 5195	0 2699	201
SPAC	1	204 PLANTS M	27 4426	1 0977	0 3842	0 1328	25
COMP	1	MONOCULTURE	5 8222	0 9722	0 2282	0 0858	5
COMP	2	MIXTURE	21 6092	1 1272	0 3032	0 1840	18
SPAC	2	3 CM SPACING	43 8908	1 0908	0 2884	0 1238	40
COMP	1	MONOCULTURE	15 4820	1 1816	0 3582	0 1161	12
COMP	2	MIXTURE	28 1378	1 0421	0 3727	0 1398	27
SPAC	3	5 CM SPACING	45 4788	1 4211	0 4247	0 1800	22
COMP	1	MONOCULTURE	18 2201	1 3806	0 2222	0 0484	12
COMP	2	MIXTURE	28 1486	1 4872	0 5264	0 2761	20
SPAC	4	44 PLANTS M	43 8841	1 6791	0 2626	0 1242	28
COMP	1	MONOCULTURE	18 2184	1 4822	0 3848	0 1461	11
COMP	2	MIXTURE	27 8678	1 8202	0 3287	0 1087	17
SPAC	5	28 PLANTS M	40 4452	1 6795	0 3088	0 1887	28
COMP	1	MONOCULTURE	24 2828	1 6168	0 4481	0 2008	18
COMP	2	MIXTURE	24 1614	1 7280	0 3478	0 1208	14
SPAC	6	17 PLANTS M	48 7881	1 7982	0 8087	0 2887	28
COMP	1	MONOCULTURE	22 8780	1 8442	0 2792	0 1422	12
COMP	2	MIXTURE	22 8041	1 7842	0 5212	0 3880	12
SPAC	7	10 PLANTS M	44 4108	1 6618	0 5184	0 3728	28
COMP	1	MONOCULTURE	28 8712	1 8747	0 3184	0 1020	12
COMP	2	MIXTURE	22 7284	1 7482	0 8041	0 4485	12
CUL	4	70000002	184 8818	1 4222	0 8448	0 2888	120
SPAC	1	204 PLANTS M	18 2881	0 8822	0 2424	0 1178	20
COMP	2	MIXTURE	18 2881	0 8822	0 2424	0 1178	20
SPAC	2	3 CM SPACING	28 1887	1 3288	0 2877	0 1882	22
COMP	2	MIXTURE	28 1887	1 3288	0 2877	0 1882	22

APPENDIX 6

02/16/82

PAGE 88

CRITERION VARIABLE YH

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
SPAC	3	8 CM SPACING	28 1172	1 3088	0 4482	0 2018	28
COMP	2	MIXTURE	28 1172	1 3088	0 4482	0 2018	28
SPAC	4	44 PLANTS M	28 3228	1 4892	0 3240	0 1080	17
COMP	2	MIXTURE	28 3228	1 4892	0 3240	0 1080	17
SPAC	5	28 PLANTS M	32 8414	1 7180	0 3888	0 1488	19
COMP	2	MIXTURE	32 8414	1 7180	0 3888	0 1488	18
SPAC	6	17 PLANTS M	38 2441	1 8172	0 8081	0 8482	28
COMP	2	MIXTURE	38 2441	1 8172	0 8081	0 8482	28
SPAC	7	10 PLANTS M	18 8042	1 4888	0 5110	0 2611	11
COMP	2	MIXTURE	18 8042	1 4888	0 5110	0 2611	11
CUL	5	NORQUAY	285 0081	1 7709	0 8028	0 8487	144
SPAC	1	204 PLANTS M	45 4230	1 2881	0 3948	0 1587	28
COMP	1	MONOCULTURE	22 8812	1 4362	0 3122	0 0981	18
COMP	2	MIXTURE	22 4817	1 1817	0 4288	0 1811	18
SPAC	2	3 CM SPACING	30 0314	1 5016	0 4878	0 2378	20
COMP	1	MONOCULTURE	18 3427	1 7048	0 8888	0 2288	9
COMP	2	MIXTURE	14 8877	1 2282	0 4418	0 1888	11
SPAC	3	8 CM SPACING	38 8288	1 8820	0 2877	0 1280	22
COMP	1	MONOCULTURE	18 8887	1 8881	0 3222	0 1028	18
COMP	2	MIXTURE	18 8782	1 8288	0 2888	0 1488	12
SPAC	4	44 PLANTS M	45 0082	1 7840	0 5187	0 2880	27
COMP	1	MONOCULTURE	18 8847	1 8888	0 4112	0 1881	11
COMP	2	MIXTURE	28 1145	1 8222	0 5821	0 2401	16
SPAC	5	28 PLANTS M	24 3208	2 0287	0 4821	0 2422	12
COMP	7	MONOCULTURE	24 3208	2 0287	0 4821	0 2422	12

APPENDIX 8

03/10/82

PAGE 58

CRITERION VARIABLE YH

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
SPAC	0	17 PLANTS M	18 0000	2 250	0 7305	0 5400	81
COMP	1	MONOCULTURE	18 0000	2 250	0 7305	0 5400	81
SPAC	7	10 PLANTS M	53 725	2 527	1 2188	1 7418	181
COMP	1	MONOCULTURE	71 070	2 342	0 4837	0 2437	81
COMP	2	MIXTURE	32 550	2 555	1 0765	2 5175	101
CUL	0	700000002	285 7245	2 3635	1 4789	2 1613	1281
SPAC	1	204 PLANTS M	25 3003	1 4665	0 3718	0 1381	161
COMP	2	MIXTURE	25 3003	1 4665	0 3718	0 1381	161
SPAC	2	3 CM SPACING	31 4594	1 4300	0 5217	0 2723	221
COMP	2	MIXTURE	31 4594	1 4300	0 5217	0 2723	221
SPAC	3	5 CM SPACING	25 8285	1 5017	0 5002	0 5403	181
COMP	2	MIXTURE	25 8285	1 5017	0 5002	0 5403	181
SPAC	4	64 PLANTS M	25 5712	2 1555	0 4555	0 2357	121
COMP	2	MIXTURE	25 5712	2 1555	0 4555	0 2357	121
SPAC	5	25 PLANTS M	43 0823	2 3035	1 1272	1 2706	181
COMP	2	MIXTURE	43 0823	2 3035	1 1272	1 2706	181
SPAC	6	17 PLANTS M	51 0155	2 7725	1 7383	3 0215	221
COMP	2	MIXTURE	51 0155	2 7725	1 7383	3 0215	221
SPAC	7	10 PLANTS M	73 0051	2 5544	2 2555	5 0575	201
COMP	2	MIXTURE	73 0051	2 5544	2 2555	5 0575	201
CUL	7	HORQUAY	124 9320	2 0755	1 2033	1 0727	501
SPAC	1	204 PLANTS M	5 1733	0 1834	0 0355	0 0055	81
COMP	2	MIXTURE	5 1733	0 1834	0 0355	0 0055	81
SPAC	2	3 CM SPACING	15 4570	1 5570	0 4155	0 1747	111
COMP	1	MONOCULTURE	15 4570	1 5570	0 4155	0 1747	81

APPENDIX 8

03/10/82

PAGE 59

CRITERION VARIABLE YH

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
COMP	2	MIXTURE	5 1624	1 2325	0 2504	0 0627	51
SPAC	3	5 CM SPACING	11 3371	1 5555	0 5051	0 2573	51
COMP	7	MONOCULTURE	11 3371	1 5555	0 5051	0 2573	51
SPAC	4	64 PLANTS M	11 7355	1 5551	0 4555	0 2171	51
COMP	1	MONOCULTURE	11 7355	1 5551	0 4555	0 2171	51
SPAC	5	25 PLANTS M	15 0875	2 1554	0 5554	0 4347	71
COMP	2	MIXTURE	15 0875	2 1554	0 5554	0 4347	71
SPAC	6	17 PLANTS M	25 5513	2 4434	1 5441	3 7755	151
COMP	2	MIXTURE	25 5513	2 4434	1 5441	3 7755	151
SPAC	7	10 PLANTS M	34 2555	2 5555	1 5554	2 4315	121
COMP	1	MONOCULTURE	20 5554	2 4514	1 5751	2 5552	51
COMP	2	MIXTURE	13 2762	2 2397	1 2550	1 5503	51
CUL	5	H9701	414 3354	2 3515	1 2355	1 5257	1501
SPAC	1	204 PLANTS M	45 5512	1 4344	0 2335	0 0547	251
COMP	2	MIXTURE	45 5512	1 4344	0 2335	0 0547	251
SPAC	2	3 CM SPACING	55 3051	1 7542	0 3751	0 1437	311
COMP	2	MIXTURE	55 3051	1 7542	0 3751	0 1437	311
SPAC	3	5 CM SPACING	57 0575	1 5555	0 3151	0 1015	501
COMP	2	MIXTURE	47 5572	1 5555	0 3151	0 1015	241
SPAC	4	64 PLANTS M	55 4555	2 1522	0 5551	0 4552	301
COMP	2	MIXTURE	55 4555	2 1522	0 5551	0 4552	301
SPAC	5	25 PLANTS M	4 5555	2 3142	0 5151	0 2753	151
COMP	2	MIXTURE	4 5555	2 3142	0 5151	0 2753	151
SPAC	5	17 PLANTS M	54 4351	2 5515	1 1555	1 3552	231
COMP	2	MIXTURE	54 4351	2 5515	1 1555	1 3552	231

APPENDIX 8

03/16/82

PAGE 51

CRITERION VARIABLE VM

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
SPAC	7	10 PLANTS M	85 8433	3 2201	2 4646	6 0200	(20)
COMP	2	MIXTURE	83 8433	3 2201	2 4646	6 0200	(20)
TOTAL CASES =	1000						
MISSING CASES =	48 OR	4 4 PCT					

APPENDIX 8

03/16/82

PAGE 52

FILE WP77 (CREATION DATE = 03/16/82)

CRITERION VARIABLE BWT CUL WEIGHT PER 1000 KERNELS G

BROKEN DOWN BY CUL CULTIVAR

BY SPAC INTER-PLANT SPACING

BY COMP TREATMENT

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
FOR ENTIRE POPULATION			42688 7874	41 2488	7 2888	53 2888	(1022)
CUL	2	OLENLEA	8851 0077	48 7348	5 0682	25 4880	(178)
SPAC	1	204 PLANTS M	1178 1888	51 2882	3 1304	8 7888	(23)
COMP	1	MONOCULTURE	884 3181	50 7889	3 7874	14 4201	(11)
COMP	2	MIXTURE	820 8834	51 7378	2 4688	5 8888	(12)
SPAC	2	3 CM SPACING	1717 3338	50 8088	4 4677	18 7834	(34)
COMP	1	MONOCULTURE	880 8488	50 0031	5 8828	30 8448	(13)
COMP	2	MIXTURE	1087 7829	59 8238	2 7184	13 8841	(21)
SPAC	3	8 CM SPACING	1785 1987	50 1488	5 7838	33 8848	(38)
COMP	1	MONOCULTURE	821 7048	47 8234	5 8231	43 8888	(13)
COMP	2	MIXTURE	1123 4938	51 8238	4 8841	23 8822	(22)
SPAC	4	44 PLANTS M	1378 1382	48 3888	6 8889	42 8180	(20)
COMP	1	MONOCULTURE	841 8880	48 3788	2 2778	5 8828	(13)
COMP	2	MIXTURE	757 3388	48 1488	8 7829	77 3182	(18)
SPAC	5	28 PLANTS M	988 7877	47 8388	2 1878	4 7884	(30)
COMP	1	MONOCULTURE	718 7428	47 7182	2 2208	4 8228	(28)
COMP	2	MIXTURE	241 8848	48 2110	2 2888	5 2418	(8)
SPAC	6	17 PLANTS M	938 2883	48 8184	7 0882	50 2818	(18)
COMP	1	MONOCULTURE	803 1088	50 2888	8 7784	48 8888	(12)
COMP	2	MIXTURE	328 1824	48 8889	7 8217	58 8784	(7)
SPAC	7	10 PLANTS M	834 0728	48 1817	3 1848	8 8884	(18)
COMP	1	MONOCULTURE	497 8888	50 8882	3 2888	10 7881	(8)

APPENDIX 6

03/16/82

PAGE 63

CRITERION VARIABLE KWT

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
COMP	2	MIXTURE	526 4671	47 8662	2 4648	6 7828	11
CUL	3	PARK	7397 8488	36 7344	6 1882	37 0004	207
SPAC	1	304 PLANTS M	828 8888	27 1708	4 0428	24 4264	30
COMP	1	MONOCULTURE	212 8884	26 8897	3 2898	11 4867	8
COMP	2	MIXTURE	716 8881	27 6782	6 2184	28 2860	10
SPAC	2	3 CM SPACING	1488 1211	36 7344	6 5768	31 1204	41
COMP	1	MONOCULTURE	488 8221	34 8221	7 8822	87 2042	14
COMP	2	MIXTURE	878 2888	36 1888	4 2288	18 7368	27
SPAC	3	6 CM SPACING	1627 2288	36 8222	4 8782	24 7828	32
COMP	1	MONOCULTURE	428 7828	36 8828	6 2410	26 8288	12
COMP	2	MIXTURE	718 8722	36 8228	4 8888	22 7871	30
SPAC	4	44 PLANTS M	1622 8188	36 8112	4 8888	22 8878	28
COMP	1	MONOCULTURE	362 8488	34 8881	6 8828	26 7881	11
COMP	2	MIXTURE	888 8888	36 2188	3 8478	12 2888	17
SPAC	5	28 PLANTS M	1621 8888	36 8787	6 8842	43 8424	28
COMP	1	MONOCULTURE	821 7787	34 7882	6 7848	78 8488	18
COMP	2	MIXTURE	888 8172	36 8227	3 8112	8 8182	14
SPAC	6	17 PLANTS M	888 8888	34 7877	6 8828	26 8424	30
COMP	1	MONOCULTURE	481 8728	36 8888	4 8248	18 1887	12
COMP	2	MIXTURE	482 8888	34 8778	6 8112	26 1241	12
SPAC	7	10 PLANTS M	888 2888	34 8724	3 8744	26 8181	30
COMP	1	MONOCULTURE	428 1244	32 8488	2 8878	8 2242	12
COMP	2	MIXTURE	488 1881	36 2888	2 8818	18 8888	12
CUL	4	YONGGONGGONG	4822 2841	36 8217	6 1288	26 4181	120
SPAC	1	304 PLANTS M	782 2222	36 1818	6 1888	27 8247	30
COMP	2	MIXTURE	782 2222	36 1818	6 1888	27 8247	30
SPAC	2	3 CM SPACING	814 8632	27 8287	4 8818	26 7118	22
COMP	2	MIXTURE	814 8632	27 8287	4 8818	26 7118	22

APPENDIX 6

03/16/82

PAGE 64

CRITERION VARIABLE KWT

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
SPAC	2	6 CM SPACING	728 8208	36 4210	3 8812	12 8822	20
COMP	2	MIXTURE	728 8208	36 4210	3 8812	12 8822	20
SPAC	4	44 PLANTS M	888 8887	36 8880	7 2110	82 4808	17
COMP	2	MIXTURE	888 8887	36 8880	7 2110	82 4808	17
SPAC	5	28 PLANTS M	781 8188	36 8428	2 7442	7 8208	18
COMP	2	MIXTURE	781 8188	36 8428	2 7442	7 8208	18
SPAC	6	17 PLANTS M	711 8827	36 8841	4 8884	22 8428	30
COMP	2	MIXTURE	711 8827	36 8841	4 8884	22 8428	30
SPAC	7	10 PLANTS M	388 7828	32 2448	7 8288	61 2881	12
COMP	2	MIXTURE	388 7828	32 2448	7 8288	61 2881	12
CUL	5	BORQUAY	8288 2222	46 1284	4 8888	24 8182	188
SPAC	1	304 PLANTS M	1888 8881	46 4172	3 8848	12 8882	30
COMP	1	MONOCULTURE	788 4888	46 2187	2 8888	4 2878	18
COMP	2	MIXTURE	888 1882	46 7424	4 4248	18 8778	18
SPAC	2	3 CM SPACING	882 2787	46 8782	6 1888	26 8844	21
COMP	1	MONOCULTURE	418 8284	46 1817	2 2828	8 2884	8
COMP	2	MIXTURE	828 7442	43 8884	6 4888	42 8882	12
SPAC	3	6 CM SPACING	1022 1218	46 4848	3 2887	18 8772	32
COMP	1	MONOCULTURE	487 1788	46 7128	2 8488	4 1848	18
COMP	2	MIXTURE	888 8824	43 8248	2 8178	14 8721	12
SPAC	4	44 PLANTS M	1147 2888	42 4682	6 2487	28 8888	27
COMP	1	MONOCULTURE	471 4881	42 8888	6 8882	48 4442	11
COMP	2	MIXTURE	878 7824	42 2248	6 2247	28 1888	18
SPAC	5	28 PLANTS M	828 2288	46 1817	4 8128	18 1882	12
COMP	1	MONOCULTURE	828 2288	46 1817	4 8128	18 1882	12

APPENDIX 5

02/10/82

PAGE 06

CRITERION VARIABLE KWT

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
SPAC	8	17 PLANTS M	361 8848	45 1081	5 7887	48 0488	81
COMP	1	MONOCULTURE	361 8848	45 1081	5 7887	48 0488	81
SPAC	7	10 PLANTS M	608 1910	42 2788	5 2884	28 1218	101
COMP	1	MONOCULTURE	361 2061	42 2888	5 2178	26 2787	81
COMP	2	MIXTURE	422 8838	42 2884	5 7837	32 1048	101
CUL	8	700000002	6808 8812	42 2827	5 8812	32 2882	1271
SPAC	1	204 PLANTS M	817 8190	48 2888	5 4228	41 2848	181
COMP	2	MIXTURE	817 8190	48 2888	5 4228	41 2848	181
SPAC	2	3 CM SPACING	1068 1300	48 8241	4 7884	22 8000	221
COMP	2	MIXTURE	1068 1300	48 8241	4 7884	22 8000	221
SPAC	3	8 CM SPACING	714 8218	44 8278	5 8788	32 2482	181
COMP	2	MIXTURE	714 8218	44 8278	5 8788	32 2482	181
SPAC	4	44 PLANTS M	814 8788	42 8188	5 2428	28 8828	121
COMP	2	MIXTURE	814 8788	42 8188	5 2428	28 8828	121
SPAC	5	28 PLANTS M	727 8284	40 2888	5 8078	34 8828	181
COMP	2	MIXTURE	727 8284	40 2888	5 8078	34 8828	181
SPAC	6	17 PLANTS M	888 2888	42 2878	4 8888	21 8742	211
COMP	2	MIXTURE	888 2888	42 2878	4 8888	21 8742	211
SPAC	7	10 PLANTS M	828 8488	41 8024	5 8828	32 2884	201
COMP	2	MIXTURE	828 8488	41 8024	5 8828	32 2884	201
CUL	7	8888888	2274 8722	38 8224	5 8728	28 7421	881
SPAC	1	204 PLANTS M	312 4172	38 8821	4 1828	17 2282	81
COMP	2	MIXTURE	312 4172	38 8821	4 1828	17 2282	81
SPAC	2	3 CM SPACING	418 2878	38 1281	2 8218	8 8284	111
COMP	1	MONOCULTURE	228 1788	37 8288	2 2888	18 2884	81

APPENDIX 5

02/10/82

PAGE 06

CRITERION VARIABLE KWT

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
COMP	2	MIXTURE	184 2188	38 8821	2 7888	7 2271	81
SPAC	2	8 CM SPACING	282 8722	32 8788	7 1818	51 1284	81
COMP	1	MONOCULTURE	282 8722	32 8788	7 1818	51 1284	81
SPAC	4	44 PLANTS M	218 8888	38 8488	6 8888	44 8787	81
COMP	1	MONOCULTURE	218 8888	38 8488	6 8888	44 8787	81
SPAC	5	28 PLANTS M	284 8277	37 8488	3 8287	12 8888	71
COMP	2	MIXTURE	284 8277	37 8488	3 8287	12 8888	71
SPAC	8	17 PLANTS M	827 8884	38 8837	4 8428	21 8882	181
COMP	2	MIXTURE	827 8884	38 8837	4 8428	21 8882	181
SPAC	7	10 PLANTS M	417 1427	34 7818	5 1872	37 2888	121
COMP	1	MONOCULTURE	228 4288	37 8728	3 7287	12 4882	81
COMP	2	MIXTURE	181 7872	31 8812	7 8188	68 2874	81
CUL	8	88701	7418 8822	48 8727	5 8888	34 8487	1811
SPAC	1	204 PLANTS M	1881 8888	42 4884	3 7828	14 2171	281
COMP	2	MIXTURE	1881 8888	42 4884	3 7828	14 2171	281
SPAC	2	3 CM SPACING	1288 1828	42 4887	5 1428	28 4884	321
COMP	2	MIXTURE	1288 1828	42 4887	5 1428	28 4884	321
SPAC	3	8 CM SPACING	1847 8287	42 8474	4 2214	18 7812	241
COMP	2	MIXTURE	1847 8287	42 8474	4 2214	18 7812	241
SPAC	4	44 PLANTS M	1227 8241	41 2848	4 2821	17 8872	301
COMP	2	MIXTURE	1227 8241	41 2848	4 2821	17 8872	301
SPAC	8	28 PLANTS M	722 8242	48 7228	6 8787	38 8288	181
COMP	2	MIXTURE	722 8242	48 7228	6 8787	38 8288	181
SPAC	8	17 PLANTS M	888 2284	38 8712	5 8878	32 8874	221
COMP	2	MIXTURE	888 2284	38 8712	5 8878	32 8874	221

APPENDIX B 03/16/82 PAGE 87

CRITERION VARIABLE	BY	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
SPAC	1	10 PLANTS M	1000 8703	27.2714	8.3023	70.4478	(20)
COMP	2	MIXTURE	1000 8703	27.2714	8.3023	70.4478	(20)
TOTAL CASES *	1000						
MISSING CASES *	48 00	4.4 PCT					

APPENDIX B 03/16/82 PAGE 88

FILE WPT7 (CREATION DATE = 03/16/82)

CRITERION VARIABLE	BY	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
FOR ENTIRE POPULATION			7000 3000	8.5771	47.6782	2273.0102	(1021)
CUL	2	ELENLEA	665 8181	3.8314	2.0483	4.1985	(170)
SPAC	1	204 PLANTS M	104 8285	4.8325	1.3247	1.7615	(23)
COMP	1	MONOCULTURE	57 8437	5.2312	1.4688	2.1505	(13)
COMP	2	MIXTURE	49 0029	4.0838	0.9307	0.8663	(12)
SPAC	2	3 CM SPACING	158 8189	4.8652	2.1088	4.4380	(34)
COMP	1	MONOCULTURE	70 8283	5.4489	2.0526	4.2204	(13)
COMP	2	MIXTURE	87 7815	4.1801	1.0420	1.0854	(21)
SPAC	3	6 CM SPACING	150 3801	4.2950	2.1012	4.4152	(38)
COMP	1	MONOCULTURE	54 8678	5.1321	2.7633	7.7477	(13)
COMP	2	MIXTURE	82 7722	3.8076	1.4280	2.0395	(22)
SPAC	4	44 PLANTS M	104 2077	3.8946	2.1384	4.5633	(28)
COMP	1	MONOCULTURE	42 8187	3.2782	0.2555	0.1284	(13)
COMP	2	MIXTURE	62 8911	4.3727	4.2788	18.2008	(16)
SPAC	5	26 PLANTS M	55 8322	3.8480	0.2997	0.1827	(20)
COMP	1	MONOCULTURE	45 8528	3.0635	0.2604	0.1447	(18)
COMP	2	MIXTURE	12 8795	2.8950	0.1188	0.0138	(5)
SPAC	6	17 PLANTS M	55 8078	2.7804	0.4580	0.2078	(20)
COMP	1	MONOCULTURE	38 8760	2.8404	0.4082	0.1665	(12)
COMP	2	MIXTURE	18 8828	2.8890	0.5470	0.2991	(7)
SPAC	7	10 PLANTS M	47 8447	2.5024	1.0785	1.1620	(19)
COMP	1	MONOCULTURE	24 8729	3.0718	1.4293	2.0430	(9)

APPENDIX 8

02/10/82

PAGE 88

CRITERION VARIABLE N1

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
COMP	2	MIXTURE	22 8718	2 0883	0 4881	0 2379	11
CUL	3	PARK	2458 8834	11 8412	58 5800	3718 0083	208
SPAC	1	204 PLANTS M	198 4007	7 8888	8 1831	37 8882	28
COMP	1	HOMOECULTURE	88 8882	8 2288	8 8871	88 7888	8
COMP	2	MIXTURE	148 8884	7 2818	8 2884	27 8888	18
SPAC	3	3 CM SPACING	1704 2478	41 8704	228 8888	48848 8888	41
COMP	1	HOMOECULTURE	1478 7284	108 8884	277 8888	142888 8128	14
COMP	2	MIXTURE	224 8824	8 2288	8 1181	27 2841	27
SPAC	3	8 CM SPACING	181 8844	4 7381	1 2128	1 7241	22
COMP	1	HOMOECULTURE	81 8888	4 2218	0 8112	0 8882	12
COMP	2	MIXTURE	88 8888	4 8844	1 8827	2 2882	20
SPAC	4	44 PLANTS M	108 2888	3 8888	0 8888	0 2244	27
COMP	1	HOMOECULTURE	44 8788	4 8881	0 8818	0 4244	11
COMP	2	MIXTURE	68 2188	3 7884	0 4847	0 2348	18
SPAC	5	28 PLANTS M	104 8818	3 8188	1 278	1 2781	28
COMP	1	HOMOECULTURE	88 4884	3 8887	1 4882	2 8228	18
COMP	2	MIXTURE	48 3888	3 2428	0 8881	0 2882	18
SPAC	6	17 PLANTS M	84 2217	3 8278	0 4887	0 7088	28
COMP	1	HOMOECULTURE	43 8778	3 2782	0 4877	0 4887	13
COMP	2	MIXTURE	88 4488	3 8882	0 8888	0 8128	12
SPAC	7	18 PLANTS M	183 8472	3 8834	1 2888	1 8482	28
COMP	1	HOMOECULTURE	87 1218	0 8848	1 7828	2 1774	12
COMP	2	MIXTURE	48 8184	3 8328	0 8214	0 2718	12
CUL	4	700000000	888 1428	8 2888	8 4881	42 8881	128
SPAC	1	204 PLANTS M	218 7882	18 8488	12 8878	187 2482	28
COMP	2	MIXTURE	218 7882	18 8488	12 8878	187 2482	28
SPAC	2	3 CM SPACING	148 2887	8 8884	8 8888	28 2488	22
COMP	2	MIXTURE	148 2887	8 8884	8 8888	28 2488	22

APPENDIX 8

02/10/82

PAGE 70

CRITERION VARIABLE N1

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
SPAC	3	8 CM SPACING	88 8887	4 2228	1 2824	1 8478	28
COMP	2	MIXTURE	88 8887	4 2228	1 2824	1 8478	28
SPAC	4	44 PLANTS M	78 2888	4 4828	3 2888	10 8288	17
COMP	2	MIXTURE	78 2888	4 4828	3 2888	10 8288	17
SPAC	5	28 PLANTS M	83 4871	3 2428	0 8788	0 2284	18
COMP	2	MIXTURE	83 4871	3 2428	0 8788	0 2284	18
SPAC	6	17 PLANTS M	88 7448	3 2872	1 0388	1 8828	28
COMP	2	MIXTURE	88 7448	3 2872	1 0388	1 8828	28
SPAC	7	18 PLANTS M	41 8828	3 8887	2 2818	8 8778	11
COMP	2	MIXTURE	41 8828	3 8887	2 2818	8 8778	11
CUL	5	HOMOECULTURE	1288 8884	8 2141	28 2418	1888 8288	148
SPAC	1	204 PLANTS M	282 2487	8 2888	11 4482	121 8884	28
COMP	1	HOMOECULTURE	74 8887	4 8884	1 4881	2 1888	18
COMP	2	MIXTURE	218 1888	11 4822	14 8282	222 1884	18
SPAC	2	3 CM SPACING	884 8882	28 4222	108 4278	10888 7818	21
COMP	1	HOMOECULTURE	28 7184	3 8878	1 8881	1 1218	8
COMP	2	MIXTURE	818 1878	42 8881	128 7818	17818 8788	12
SPAC	3	8 CM SPACING	108 8288	4 2481	1 8278	2 2248	22
COMP	1	HOMOECULTURE	28 2488	3 241	0 8222	0 8718	18
COMP	2	MIXTURE	88 8878	4 8882	1 8882	2 2718	12
SPAC	4	44 PLANTS M	141 2881	4 1248	3 2122	10 8788	27
COMP	1	HOMOECULTURE	41 8184	3 7828	2 8888	8 4228	11
COMP	2	MIXTURE	88 7887	4 2888	3 2888	12 8888	18
SPAC	5	28 PLANTS M	41 8124	3 4827	2 0224	4 1248	12
COMP	1	HOMOECULTURE	41 8124	3 4827	2 0224	4 1248	12

APPENDIX 6			02/10/82				PAGE 71
CRITERION VARIABLE #1							
VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
SPAC COMP	0	17 PLANTS M	83 8378	0 8022	11 8838	138 7288	81
	1	MONOCULTURE	83 8378	0 8022	11 8838	138 7288	81
SPAC COMP	7	10 PLANTS M	81 8782	2 7147	0 8077	0 3883	181
	1	MONOCULTURE	28 8388	2 7618	0 7818	0 8888	81
COMP	2	MIXTURE	28 8422	2 8842	0 8821	0 2144	181
CUL SPAC	0	700000002	822 4882	0 4782	14 8778	188 1818	1271
	1	204 PLANTS M	188 4744	0 8897	12 8388	187 1837	181
COMP	2	MIXTURE	188 4744	0 8897	12 8388	187 1837	181
SPAC COMP	3	3 CM SPACING	334 8328	18 8248	38 8862	882 2322	211
	8	MIXTURE	334 8328	18 8248	38 8862	882 2322	211
SPAC COMP	3	8 CM SPACING	78 1888	4 8888	1 7788	3 1818	181
	2	MIXTURE	78 1888	4 8888	1 7788	3 1818	181
SPAC COMP	4	44 PLANTS M	41 3822	2 4418	0 7288	0 8271	121
	2	MIXTURE	41 3822	2 4418	0 7288	0 8271	121
SPAC COMP	8	28 PLANTS M	88 8322	2 8882	1 4847	2 1888	181
	2	MIXTURE	88 8322	2 8882	1 4847	2 1888	181
SPAC COMP	8	17 PLANTS M	88 7884	4 1288	4 8888	21 8834	221
	2	MIXTURE	88 7884	4 1288	4 8888	21 8834	221
SPAC COMP	7	10 PLANTS M	81 8848	2 8882	0 4827	0 2328	201
	2	MIXTURE	81 8848	2 8882	0 4827	0 2328	201
CUL SPAC	7	NORWAY	288 7882	2 7888	1 8812	3 3884	881
	1	204 PLANTS M	41 1818	0 1277	0 8211	0 8742	81
COMP	2	MIXTURE	41 1818	0 1277	0 8211	0 8742	81
SPAC COMP	2	3 CM SPACING	87 8888	4 2142	1 8812	2 7287	111
	1	MONOCULTURE	24 8878	4 1148	2 2287	4 8388	81

APPENDIX 6		02/10/82					PAGE 72
CRITERION VARIABLE #1							
VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
COMP	2	MIXTURE	22 7888	4 8838	0 7228	0 8228	81
SPAC	3	8 CM SPACING	22 2224	3 8784	2 2778	8 8882	81
COMP	1	MONOCULTURE	22 2224	3 8784	2 2778	8 8882	81
SPAC	4	44 PLANTS M	20 7428	3 4872	1 8788	1 1888	81
COMP	1	MONOCULTURE	20 7428	3 4872	1 8788	1 1888	81
SPAC	8	28 PLANTS M	20 1848	2 8848	0 3228	0 1882	71
COMP	2	MIXTURE	20 1848	2 8848	0 3228	0 1882	71
SPAC	8	17 PLANTS M	48 8148	3 2878	0 7888	0 4888	181
COMP	2	MIXTURE	48 8148	3 2878	0 7888	0 4888	181
SPAC	7	10 PLANTS M	44 2782	3 8888	3 2881	10 8378	121
COMP	1	MONOCULTURE	18 8818	2 4778	0 4811	0 1887	81
COMP	2	MIXTURE	28 8142	4 9181	4 4881	18 4888	81
CUL	8	NORWAY	878 7888	8 4428	22 7887	818 8841	1881
SPAC	1	204 PLANTS M	118 2888	4 4148	0 8288	0 8818	281
COMP	2	MIXTURE	118 2888	4 4148	0 8288	0 8818	281
SPAC	3	3 CM SPACING	448 8888	14 4784	84 8447	2888 8378	311
COMP	2	MIXTURE	448 8888	14 4784	84 8447	2888 8378	311
SPAC	3	8 CM SPACING	84 8182	3 8171	1 2228	1 7481	261
COMP	8	MIXTURE	84 8182	3 8171	1 2228	1 7481	261
SPAC	4	44 PLANTS M	117 8122	2 8384	2 8888	7 8881	301
COMP	2	MIXTURE	117 8122	2 8384	2 8888	7 8881	301
SPAC	8	28 PLANTS M	81 4884	2 8888	0 4271	0 1828	181
COMP	2	MIXTURE	81 4884	2 8888	0 4271	0 1828	181
SPAC	8	17 PLANTS M	70 8128	3 8781	1 8811	1 1888	231
COMP	2	MIXTURE	70 8128	3 8781	1 8811	1 1888	231
SPAC	7	10 PLANTS M	88 8382	2 8841	0 8817	0 8881	881
COMP	2	MIXTURE	88 8382	2 8841	0 8817	0 8881	881
TOTAL CASES :			1888				
MISSING CASES :			48 OR	4 8 PCT			

POOR COPY
COPIE DE QUALITEE INFERIEURE

214

Appendix 7. Descriptive statistics of characters of plants, grown in monoculture and in mixture at two interplant spacings, measured at maturity in 1978.

POOR COPY
COPIE DE QUALITEE INFERIEURE

215

APPENDIX 7

02/10/82

PAGE 2

FILE SPHRC CREATION DATE = 10/20/81

CRITERION VARIABLE		DESCRIPTION		OF SUBPOPULATIONS			
BROKEN DOWN BY		CUL		TILLERS PER PLANT			
BY		SPAC		CULTIVAR			
BY		MYR		INTER-PLANT SPACING			
BY		TREATMENT					
VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
FOR ENTIRE POPULATION			2017 0000	6 0083	3 1373	9 7700	669
CUL	1	PITIC 62	670 0000	6 0512	4 3600	18 0088	62
	2	3 CM SPACING	217 0000	4 6208	1 3930	1 0570	42
SPAC	2	MIXTURE	217 0000	4 6208	1 2889	1 0570	42
SPAC	3	9 CM SPACING	363 0000	10 3824	4 8117	23 1824	36
	2	MIXTURE	363 0000	10 3824	4 8117	23 1824	36
CUL	2	BLEND	988 0000	9 6363	3 0840	9 5113	170
	2	3 CM SPACING	309 0000	3 8786	7 1633	1 3833	86
SPAC	1	MIXTURE	62 0000	3 4444	0 9218	0 8497	16
MYR	2	MIXTURE	247 0000	3 7424	1 2193	1 4668	60
SPAC	3	9 CM SPACING	648 0000	7 6666	3 1788	10 1088	60
	1	MIXTURE	113 0000	6 2774	2 4444	6 0771	18
MYR	2	MIXTURE	636 0000	7 6624	3 2808	10 7621	60
CUL	3	PARK	788 0000	9 8888	2 3888	5 6574	127
	2	3 CM SPACING	204 0000	4 6788	1 3931	1 0400	60
SPAC	1	MIXTURE	58 0000	5 3636	0 8000	0 6400	11
MYR	2	MIXTURE	246 0000	4 5370	1 4808	2 1024	54
SPAC	3	9 CM SPACING	454 0000	7 3226	2 4888	6 0910	62
	1	MIXTURE	113 0000	8 4107	2 2344	4 9924	12
MYR	2	MIXTURE	341 0000	6 6200	2 2860	5 1302	60
CUL	4	TOMORROW	850 0000	9 8511	2 8958	7 2879	84
	2	3 CM SPACING	189 0000	4 1087	1 1201	1 2544	48
SPAC	1	MIXTURE	77 0000	4 2778	1 0741	1 1528	76
MYR							

APPENDIX 7

02/10/82

PAGE 3

CRITERION VARIABLE Y

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
MYR	2	MIXTURE	112 0000	4 0000	1 1647	1 3333	(88)
SPAC	2	0 CM SPACING	201 0000	7 0304	2 7122	7 2612	(68)
MYR	1	MONOCULTURE	144 0000	8 0000	2 0000	8 7680	(18)
MYR	2	MIXTURE	217 0000	7 2222	2 0000	8 0000	(30)
CUL	5	MONOCULTURE	521 0000	8 0221	2 1020	8 4288	(88)
SPAC	2	3 CM SPACING	143 0000	2 8722	1 8010	2 2776	(28)
MYR	1	MONOCULTURE	70 0000	2 8222	1 2722	1 6240	(18)
MYR	2	MIXTURE	73 0000	4 0000	2 0422	4 1732	(18)
SPAC	2	0 CM SPACING	428 0000	7 2000	2 0000	8 0024	(88)
MYR	1	MONOCULTURE	128 0000	7 0000	2 1804	10 1176	(18)
MYR	2	MIXTURE	200 0000	7 1428	2 0000	8 0000	(82)
TOTAL CASES =	870						
MISSING CASES =	1 00	0.2 PCT.					

APPENDIX 7

02/10/82

PAGE 4

FILE SPHERE (CREATION DATE = 12/30/81)

CRITERION VARIABLE Y

CRITERION VARIABLE	BY	DESCRIPTION	SUM	MEAN	STD DEV	VARIANCE	N
SPAC	BY	HEADS PER PLANT					
MYR	BY	CULTIVAR					
MYR	BY	INTER-PLANT SPACING					
MYR	BY	TREATMENT					
VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
FOR ENTIRE POPULATION			3043 0000	8 2480	2 8281	8 0007	(688)
CUL	1	PITIC 82	484 0000	8 8888	2 1040	8 4288	(82)
SPAC	2	3 CM SPACING	187 0000	2 8888	1 4178	2 0102	(48)
MYR	2	MIXTURE	187 0000	2 8888	1 4178	2 0102	(48)
SPAC	2	0 CM SPACING	277 0000	8 1471	2 1443	8 4288	(34)
MYR	2	MIXTURE	277 0000	8 1471	2 1443	8 4288	(34)
CUL	2	GLENNLEA	608 0000	4 7412	2 4430	8 4288	(170)
SPAC	2	3 CM SPACING	272 0000	2 2221	1 0014	1 0020	(64)
MYR	1	MONOCULTURE	88 0000	2 0000	1 0000	1 1144	(18)
MYR	2	MIXTURE	217 0000	2 2270	0 8888	0 8776	(88)
SPAC	2	0 CM SPACING	924 0000	8 2093	2 8443	8 4732	(88)
MYR	1	MONOCULTURE	88 0000	4 7778	1 7878	3 1243	(18)
MYR	2	MIXTURE	448 0000	8 8888	2 8820	8 2204	(88)
CUL	3	PARK	708 0000	8 8827	2 1883	4 8224	(127)
SPAC	2	3 CM SPACING	291 0000	4 4769	1 3122	1 7221	(88)
MYR	1	MONOCULTURE	88 0000	8 2727	0 8048	0 8182	(11)
MYR	2	MIXTURE	223 0000	4 3148	1 2282	1 7080	(88)
SPAC	2	0 CM SPACING	418 0000	8 7419	2 3488	8 0001	(82)
MYR	1	MONOCULTURE	88 0000	8 2500	2 2404	8 4772	(12)
MYR	2	MIXTURE	218 0000	8 2500	2 2212	4 9243	(80)
CUL	4	700000002	830 0000	8 8283	2 8788	7 1708	(84)
SPAC	2	3 CM SPACING	183 0000	2 8782	1 1830	1 3888	(48)
MYR	1	MONOCULTURE	74 0000	4 1111	1 1218	1 2810	(18)

APPENDIX 7

03/16/82

PAGE 6

CRITERION VARIABLE N

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
MTR	2	MIXTURE	100 0000	2 0000	1 0274	1 0556	(20)
SPAC	3	0 CM SPACING	247 0000	7 2282	2 7646	7 6447	(48)
MTR	1	MONOCULTURE	123 0000	7 3680	3 0706	9 4261	(18)
MTR	2	MIXTURE	214 0000	7 1333	2 0082	0 7402	(20)
CUL	5	MONOCULTURE	534 0000	5 0026	2 0076	0 3081	(00)
SPAC	2	3 CM SPACING	133 0000	3 0044	1 2380	1 5326	(20)
MTR	1	MONOCULTURE	70 0000	2 0000	1 2783	1 6340	(18)
MTR	2	MIXTURE	63 0000	3 0000	1 3006	1 6412	(18)
SPAC	3	0 CM SPACING	401 0000	6 0022	3 0237	9 2021	(00)
MTR	1	MONOCULTURE	124 0000	6 0000	3 1039	9 6340	(18)
MTR	2	MIXTURE	277 0000	6 0000	3 0206	9 2294	(42)
TOTAL CASES *	STD						
MISSING CASES *	1 OR	0 2 PCT					

APPENDIX 7

03/16/82

PAGE 6

FILE SPHERE (CREATION DATE = 12/30/81)

DESCRIPTION OF SUBPOPULATIONS

CRITERION VARIABLE		WT	DRY WEIGHT PER PLANT	POPULATIONS			
BROKEN DOWN BY		CUL	CULTIVAR				
BY	BY	BY	INTER-PLANT SPACING				
BY	BY	BY	TREATMENT				
		SPAC					
		MTR					
VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
FOR ENTIRE POPULATION:			11137 6862	10 7127	12 0896	146 8761	(606)
CUL	1	PITIC 82	1796 1194	21 0478	14 0063	210 4038	(08)
SPAC	2	3 CM SPACING	635 3307	13 2360	4 0013	24 0230	(48)
MTR	2	MIXTURE	635 3307	13 2360	4 0013	24 0230	(48)
SPAC	3	0 CM SPACING	1139 7887	33 0222	16 2873	234 0088	(34)
MTR	2	MIXTURE	1139 7887	33 0222	16 2873	234 0088	(34)
CUL	2	GLENLEA	3867 6486	22 0622	13 7837	189 1842	(168)
SPAC	2	3 CM SPACING	1147 4094	13 0262	4 0787	21 6898	(63)
MTR	1	MONOCULTURE	207 3399	12 1066	4 0706	21 6146	(17)
MTR	2	MIXTURE	940 1696	16 2448	4 0230	21 3728	(60)
SPAC	3	0 CM SPACING	2710 1481	31 0041	13 0446	191 8706	(66)
MTR	1	MONOCULTURE	478 2798	26 0090	10 1277	102 7728	(18)
MTR	2	MIXTURE	2232 8694	33 2412	16 3077	267 2088	(67)
CUL	3	PARK	1814 8990	14 2880	6 0702	46 0631	(127)
SPAC	2	3 CM SPACING	870 3196	10 2128	3 7853	14 4042	(66)
MTR	1	MONOCULTURE	181 4388	12 0200	2 6186	6 3443	(18)
MTR	2	MIXTURE	616 8797	8 7902	3 0874	14 6782	(62)
SPAC	3	0 CM SPACING	1144 3784	16 0677	7 1223	60 7277	(62)
MTR	1	MONOCULTURE	309 6199	26 0017	7 2732	62 8899	(12)
MTR	2	MIXTURE	834 7585	16 0062	6 0074	36 0973	(60)
CUL	4	700000002	1826 4692	18 0620	10 7884	116 3472	(92)
SPAC	2	3 CM SPACING	648 0886	12 4080	3 0876	13 6730	(44)
MTR	1	MONOCULTURE	220 1089	14 3818	3 7247	13 6731	(18)

APPENDIX 7

03/16/82

PAGE 7

CRITERION VARIABLE WT

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
MTR	2	MIXTURE	316.8706	11.2614	3.3218	10.4446	(28)
SPAC	3	9 CM SPACING	1288.4798	38.9787	10.8488	113.2386	(48)
MTR	1	MONOCULTURE	826.2388	28.2487	12.3821	153.2742	(18)
MTR	2	MIXTURE	783.2397	28.0747	9.2383	87.1477	(30)
CUL	5	NORWAY	1883.7882	18.4142	10.7882	118.8884	(88)
SPAC	2	3 CM SPACING	427.7888	11.8823	5.8817	35.8232	(38)
MTR	1	MONOCULTURE	248.8388	13.8133	4.8871	20.8887	(18)
MTR	2	MIXTURE	182.7888	10.1833	8.8880	28.8888	(18)
SPAC	3	9 CM SPACING	1428.8884	23.8327	10.7821	118.8876	(88)
MTR	1	MONOCULTURE	482.8888	28.1884	11.1242	123.7482	(18)
MTR	2	MIXTURE	883.8886	23.4888	10.8821	114.1076	(42)

TOTAL CASES = 870
MISSING CASES = 0 OR 0.0 PCT

APPENDIX 7

03/18/82

PAGE 8

FILE SPHERC (CREATION DATE = 12/30/81)

CRITERION VARIABLE WT

DESCRIPTION OF SUBPOPULATIONS

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
FOR ENTIRE POPULATION			4901.1880	8.5888	8.4084	28.2182	(870)
CUL	1	PITIC 82	777.8887	8.4828	8.3218	39.8821	(82)
SPAC	2	3 CM SPACING	278.8888	8.8333	2.4888	6.2342	(48)
MTR	2	MIXTURE	278.8888	8.8333	2.4888	6.2342	(48)
SPAC	3	9 CM SPACING	487.8888	12.8383	8.8180	42.4847	(34)
MTR	2	MIXTURE	487.8888	14.8383	8.8180	42.4847	(34)
CUL	2	BLEND 82	1878.1882	8.8800	8.8187	38.8428	(170)
SPAC	2	3 CM SPACING	804.8888	8.8071	2.8848	4.3482	(84)
MTR	1	MONOCULTURE	84.8000	8.2722	2.3887	5.8880	(18)
MTR	2	MIXTURE	488.8888	8.2078	2.8888	4.8278	(88)
SPAC	3	9 CM SPACING	1171.8884	12.8232	8.8388	38.4318	(88)
MTR	1	MONOCULTURE	184.8888	10.8187	4.3017	18.8880	(18)
MTR	2	MIXTURE	878.8888	14.8288	8.2328	38.8478	(88)
CUL	2	PARK	747.8887	8.8414	3.8272	8.1643	(128)
SPAC	2	3 CM SPACING	271.8888	4.1187	1.8287	2.8482	(88)
MTR	1	MONOCULTURE	80.2000	8.8187	1.2848	1.8781	(12)
MTR	2	MIXTURE	211.8888	3.8204	1.8388	2.8748	(84)
SPAC	3	9 CM SPACING	478.7888	7.8742	3.1048	8.8287	(82)
MTR	1	MONOCULTURE	127.2000	10.8883	3.2843	10.8884	(12)
MTR	2	MIXTURE	348.4888	8.8700	2.8444	8.8828	(80)
CUL	4	700000002	880.8888	8.1888	8.1824	28.8471	(88)
SPAC	2	3 CM SPACING	280.8888	8.8874	1.7823	3.1228	(88)
MTR	1	MONOCULTURE	117.8000	8.8444	1.8871	3.8881	(18)

APPENDIX 7

02/16/82

PAGE 9

CRITERION VARIABLE Y

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
MTR	2	MIXTURE	142 8898	5 1036	1 4886	2 2488	281
SPAC	3	9 CM SPACING	800 1884	12 5042	5 1082	26 0826	481
MTR	1	MONOCULTURE	241 7889	13 4332	5 7384	32 9088	181
MTR	2	MIXTURE	256 3899	11 8487	4 7668	22 1425	301
CUL	5	BERGUA	638 7887	8 7378	5 1240	28 2882	881
SPAC	2	3 CM SPACING	187 3889	5 2088	2 2877	5 2327	381
MTR	1	MONOCULTURE	186 9089	8 0680	2 0823	4 2121	181
MTR	2	MIXTURE	78 8080	4 2811	2 2478	5 0821	181
SPAC	3	9 CM SPACING	881 3888	10 8887	5 1884	28 8828	881
MTR	1	MONOCULTURE	287 8889	11 8332	5 4888	28 8888	181
MTR	2	MIXTURE	443 7889	10 8887	5 1221	26 2482	421

TOTAL CASES = 870

APPENDIX 7

02/16/82

PAGE 10

FILE SPHERE (CREATION DATE = 12/30/81)

CRITERION VARIABLE DESCRIPTION OF SUBPOPULATIONS

CRITERION VARIABLE		DESCRIPTION	NUMBER OF KERNELS PER PLANT				
BROKEN DOWN BY		CUL	CULTIVAR				
BY		SPAC	INTER-PLANT SPACING				
BY		MTR	TREATMENT				
VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
FOR ENTIRE POPULATION			118888 0000	210 8226	126 8218	18023 1111	(870)
CUL	1	PITIC 82	21420 0000	261 2186	173 2322	30069 3832	(821)
SPAC	2	3 CM SPACING	7884 0000	183 8332	85 1088	4228 7378	(481)
MTR	2	MIXTURE	7884 0000	183 8332	85 1088	4228 7378	(481)
SPAC	3	9 CM SPACING	13888 0000	398 7088	185 3432	34382 0827	(341)
MTR	2	MIXTURE	13888 0000	398 7088	185 3432	34382 0827	(341)
CUL	2	GLENLEA	33728 0000	188 4000	112 2882	12803 8077	(1701)
SPAC	2	3 CM SPACING	10471 0000	124 8848	40 8728	1870 8881	(841)
MTR	1	MONOCULTURE	2087 0000	114 2778	48 7487	2082 0389	(181)
MTR	2	MIXTURE	8414 0000	127 4848	38 3424	1847 8228	(841)
SPAC	3	9 CM SPACING	23287 0000	270 4302	113 1838	12803 7088	(881)
MTR	1	MONOCULTURE	3802 0000	218 7778	88 8882	7328 2888	(181)
MTR	2	MIXTURE	18388 0000	284 8324	118 7878	13808 7722	(881)
CUL	3	PARK	21287 0000	188 1484	81 2888	8809 4802	(1281)
SPAC	2	3 CM SPACING	7881 0000	118 1081	44 2878	1881 2888	(881)
MTR	1	MONOCULTURE	1878 0000	138 8332	38 8212	1380 3322	(121)
MTR	2	MIXTURE	8183 0000	114 8000	44 8308	2018 7830	(841)
SPAC	3	9 CM SPACING	13408 0000	218 2288	82 0308	8727 4238	(821)
MTR	1	MONOCULTURE	3817 0000	283 8822	84 8881	7070 8108	(121)
MTR	2	MIXTURE	9889 0000	187 7800	70 8888	4883 8078	(801)
CUL	4	YOMODI	22203 0000	228 2021	128 8310	16888 8148	(841)
SPAC	2	3 CM SPACING	7123 0000	188 8882	44 1701	1880 8887	(881)
MTR	1	MONOCULTURE	3121 0000	173 8444	44 1741	1881 2487	(181)

APPENDIX 7

03/18/83

PAGE 11

CRITERION VARIABLE WK

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
MTR	2	MIXTURE	4002 0000	142 9288	40 4002	1032 0814	(20)
SPAC	3	8 CM SPACING	18070 0000	212 8882	120 8771	18084 0408	(48)
MTR	1	MONOCULTURE	5812 0000	328 8444	148 4108	21148 8487	(18)
MTR	2	MIXTURE	9188 0000	308 3887	121 8531	14778 1878	(30)
CUL	8	NORWAY	21380 0000	222 7083	133 1228	17724 3772	(88)
SPAC	2	3 CM SPACING	4738 0000	131 8278	88 3428	3062 8278	(38)
MTR	1	MONOCULTURE	2730 0000	181 1111	80 7884	2880 8782	(18)
MTR	2	MIXTURE	2018 0000	111 8444	83 8732	2812 1144	(18)
SPAC	3	8 CM SPACING	18848 0000	277 4187	138 4027	18808 7048	(88)
MTR	1	MONOCULTURE	5322 0000	288 8887	148 2877	21182 7888	(18)
MTR	2	MIXTURE	11322 0000	288 8882	133 4738	17818 2712	(42)
TOTAL CASES =			870				

APPENDIX 7

03/18/83

PAGE 12

FILE SPHERG (CREATION DATE = 12/30/81)

DESCRIPTION OF SUB POPULATIONS

CRITERION VARIABLE BROKEN DOWN BY	BY	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
FOR ENTIRE POPULATION				23118 3103	40 8884	8 8782	48 8882	(870)
CUL	1	PITIC 82	2888 8823	28 4217	4 4188	18 8342		82
SPAC	2	3 CM SPACING	1718 8881	35 8122	4 8272	24 2778		88
MTR	2	MIXTURE	1718 8881	35 8122	4 8272	24 2778		48
SPAC	3	8 CM SPACING	1287 8872	37 2823	3 4738	12 8871		34
MTR	2	MIXTURE	1287 8872	37 2823	3 4738	12 8871		34
CUL	2	GLENLBA	8322 3281	48 8848	4 7342	22 4138		170
SPAC	2	3 CM SPACING	4038 4008	48 8824	4 8280	24 2880		84
MTR	1	MONOCULTURE	838 7813	48 8323	7 8028	88 2882		18
MTR	2	MIXTURE	3208 8184	48 8308	3 8213	14 8788		88
SPAC	3	8 CM SPACING	4388 8244	48 8383	4 3804	18 2788		88
MTR	1	MONOCULTURE	888 8471	48 8887	3 4884	11 8238		18
MTR	2	MIXTURE	3383 3773	48 8028	4 8288	21 4878		88
CUL	3	PARK	4441 8873	34 7028	3 8232	12 4137		128
SPAC	2	3 CM SPACING	2282 8280	34 2807	2 8128	7 8121		88
MTR	1	MONOCULTURE	430 8100	38 8788	1 8824	2 8304		12
MTR	2	MIXTURE	1832 0180	33 8283	2 8003	8 4120		58
SPAC	3	8 CM SPACING	2178 3894	38 1818	4 1281	17 0188		82
MTR	1	MONOCULTURE	432 1847	38 0128	1 1282	1 2888		12
MTR	2	MIXTURE	1747 2447	34 8448	4 8482	20 8877		80
CUL	4	TOMCO8002	3887 3830	38 1828	3 8438	18 8711		84
SPAC	2	3 CM SPACING	1871 2830	38 3328	3 8308	18 2700		48
MTR	1	MONOCULTURE	872 7784	37 3788	3 2713	10 7018		18

APPENDIX 7

03/10/82

PAGE 13

CRITERION VARIABLE KWT

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
MTR	2	MIXTURE	888 5135	38 8812	4 2853	17 8848	23
SPAC	3	8 CM SPACING	1918 8100	38 8169	3 2843	10 2874	48
MTR	1	MONOCULTURE	740 1288	41 1168	2 8193	8 8137	18
MTR	2	MIXTURE	1178 8714	38 1087	3 3483	11 2113	30
CUL	5	MONOCULTURE	2780 1728	38 3768	2 8838	8 7785	88
SPAC	2	3 CM SPACING	1418 8813	38 3873	2 8848	8 8814	38
MTR	1	MONOCULTURE	723 7888	40 2888	2 8188	4 8787	18
MTR	2	MIXTURE	881 2882	38 4888	2 8228	7 8878	18
SPAC	3	8 CM SPACING	2388 1113	38 4188	3 1881	10 1838	88
MTR	1	MONOCULTURE	718 8788	38 4817	1 8884	3 4888	18
MTR	2	MIXTURE	1884 4488	38 3814	8 8288	13 1788	48

TOTAL CASES = 878

APPENDIX 7

03/10/82

PAGE 14

FILE SPHERE (CREATION DATE = 12/30/81)

DESCRIPTION OF SUBPOPULATIONS

CRITERION VARIABLE
BROKEN DOWN BY
BY SPAC
BY MTR

NUMBER OF KERNELS PER HEAD
CULTIVAR
INTER-PLANT SPACING
TREATMENT

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
FOR ENTIRE POPULATION			32084 1702	38 8288	9 8444	84 8813	888
CUL	1	PITIC 82	3721 8482	48 3788	12 2144	148 1827	82
SPAC	2	3 CM SPACING	2888 7488	43 8873	13 2478	178 8883	48
MTR	2	MIXTURE	2888 7488	43 8873	13 2478	178 8883	48
SPAC	3	8 CM SPACING	1884 2888	48 8888	8 8887	87 3228	38
MTR	2	MIXTURE	1884 2888	48 8888	8 8887	87 3228	38
CUL	2	8LENLEA	7828 8818	41 2474	7 2883	82 8833	178
SPAC	2	3 CM SPACING	3844 8888	38 8774	8 1283	37 8418	88
MTR	1	MONOCULTURE	878 7488	37 2838	8 1714	38 8888	18
MTR	2	MIXTURE	2878 1488	38 8828	8 1188	37 4887	88
SPAC	3	8 CM SPACING	3788 1831	43 8883	7 3827	84 8828	88
MTR	1	MONOCULTURE	818 7842	48 4888	8 1887	87 1883	18
MTR	2	MIXTURE	2881 2878	43 8488	7 1237	88 7488	88
CUL	3	PARK	3883 8888	38 8828	8 8483	44 1887	127
SPAC	2	3 CM SPACING	1782 8184	38 1841	8 2883	28 8288	88
MTR	1	MONOCULTURE	282 7878	38 7888	8 8788	31 8878	118
MTR	2	MIXTURE	1418 8188	38 2828	8 2748	28 8888	84
SPAC	3	8 CM SPACING	1881 8422	32 1138	8 8438	42 8223	82
MTR	1	MONOCULTURE	431 7888	38 8888	8 8228	38 1883	18
MTR	2	MIXTURE	1888 2788	31 1888	8 3827	48 8888	88
CUL	4	YONK88882	3814 8847	41 8442	7 7843	88 1288	84
SPAC	2	3 CM SPACING	1824 7322	38 8881	7 4411	88 3883	48
MTR	2	MONOCULTURE	777 2888	43 1833	7 4788	88 8284	18

APPENDIX 7

02/10/82

PAGE 15

CRITERION VARIABLE KH

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
MTR	2	MIXTURE	1047 4833	37 4883	6 8078	43 8281	281
SPAC	3	8 CM SPACING	2088 8218	43 8378	7 8473	88 4811	481
MTR	1	MONOCULTURE	802 2407	44 8748	7 3881	84 8878	181
MTR	2	MIXTURE	1287 4888	43 8188	7 8828	81 8848	381
CUL	8	ROSNAY	3738 8847	38 8182	8 3471	88 8734	981
SPAC	2	3 CM SPACING	1278 8781	38 8282	8 1184	88 8888	281
MTR	1	MONOCULTURE	788 7888	39 4283	7 1817	81 1487	181
MTR	2	MIXTURE	888 1888	31 8284	7 2231	82 1734	181
SPAC	3	8 CM SPACING	2088 8787	48 8888	7 8871	81 8818	881
MTR	1	MONOCULTURE	784 7811	42 4834	8 8178	88 3484	181
MTR	2	MIXTURE	1882 8778	48 3823	8 8418	48 8182	421

TOTAL CASES = 878
 MISSING CASES = 1 OR 0.2 PCT

APPENDIX 7

02/10/82

PAGE 16

FILE SPHERE (CREATION DATE = 12/30/81)

DESCRIPTION OF SUBPOPULATIONS
 CRITERION VARIABLE
 BROKEN DOWN BY
 BY CUL HARVEST INDEX
 BY SPAC CULTIVAR
 BY MTR INTER-PLANT SPACING
 BY TREATMENT

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
FOR ENTIRE POPULATION			248 2882	0 4281	0 0882	0 0023	8881
CUL	1	PITIC 82	38 8718	0 4388	0 0884	0 0044	821
SPAC	2	3 CM SPACING	21 0048	0 4378	0 0733	0 0084	481
MTR	2	MIXTURE	21 0048	0 4378	0 0733	0 0084	481
SPAC	3	8 CM SPACING	18 8874	0 4433	0 0883	0 0032	341
MTR	2	MIXTURE	18 8874	0 4432	0 0883	0 0032	341
CUL	2	GLENSA	73 1100	0 4282	0 0482	0 0018	1881
SPAC	2	3 CM SPACING	38 8848	0 4408	0 0412	0 0017	821
MTR	1	MONOCULTURE	7 8878	0 4610	0 0880	0 0038	171
MTR	2	MIXTURE	28 8188	0 4281	0 0383	0 0012	881
SPAC	3	8 CM SPACING	38 8288	0 4287	0 0387	0 0018	881
MTR	1	MONOCULTURE	7 3348	0 4878	0 0821	0 0087	181
MTR	2	MIXTURE	28 1887	0 4387	0 0322	0 0010	871
CUL	3	PARK	81 4721	0 4883	0 0431	0 0018	1271
SPAC	2	3 CM SPACING	28 8283	0 3888	0 0387	0 0018	881
MTR	1	MONOCULTURE	4 7884	0 3873	0 0840	0 0041	121
MTR	2	MIXTURE	21 1888	0 3883	0 0327	0 0011	821
SPAC	3	8 CM SPACING	38 8887	0 4130	0 0487	0 0021	821
MTR	1	MONOCULTURE	4 8881	0 4873	0 0278	0 0008	121
MTR	2	MIXTURE	20 8877	0 4132	0 0482	0 0024	801
CUL	4	YONGGONG	42 2242	0 4880	0 0388	0 0013	821
SPAC	2	3 CM SPACING	18 8848	0 4818	0 0380	0 0012	481
MTR	1	MONOCULTURE	7 1888	0 4488	0 0844	0 0020	181

APPENDIX 7

02/10/82

PAGE 17

CRITERION VARIABLE M1

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
MTR	2	MIXTURE	12 8838	0 4520	0 0200	0 0008	281
SPAC	3	9 CM SPACING	22 3894	0 4668	0 0282	0 0012	481
MTR	1	MONOCULTURE	8 1497	0 4623	0 0218	0 0010	181
MTR	2	MIXTURE	14 2167	0 4740	0 0281	0 0012	301
CUL	8	MONOCULTURE	42 4071	0 4417	0 0418	0 0017	881
SPAC	2	3 CM SPACING	18 7277	0 4272	0 0403	0 0021	381
MTR	1	MONOCULTURE	8 0308	0 4482	0 0428	0 0018	181
MTR	2	MIXTURE	7 7088	0 4282	0 0474	0 0022	181
SPAC	3	9 CM SPACING	28 8894	0 4448	0 0382	0 0015	801
MTR	1	MONOCULTURE	8 0827	0 4481	0 0387	0 0012	181
MTR	2	MIXTURE	18 8867	0 4428	0 0407	0 0017	421

TOTAL CASES = 870
MISSING CASES = 0 OR 0.0 PCT

APPENDIX 7

02/10/82

PAGE 18

FILE SPHERE (CREATION DATE = 12/30/81)

CRITERION VARIABLE M1

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
FOR ENTIRE POPULATION			30120 0000	82 8421	10 2888	108 4021	8701
CUL	1	PITIC 82	4328 0000	82 8024	7 2872	52 2486	821
SPAC	2	3 CM SPACING	2881 0000	83 8782	7 4418	55 3828	481
MTR	2	MIXTURE	2881 0000	83 8782	7 4418	55 3828	481
SPAC	3	9 CM SPACING	1747 0000	81 3824	8 8108	47 7888	341
MTR	2	MIXTURE	1747 0000	81 3824	8 8108	47 7888	341
CUL	2	GLORIA	10881 0000	82 8828	7 7247	88 8288	1701
SPAC	2	3 CM SPACING	5488 0000	88 2482	8 8703	43 1888	181
MTR	1	MONOCULTURE	1240 0000	88 8888	8 8878	44 8878	181
MTR	2	MIXTURE	4248 0000	84 2788	8 2808	38 8887	881
SPAC	4	8 CM SPACING	8182 0000	80 8222	7 8142	82 8847	881
MTR	1	MONOCULTURE	1877 0000	88 8322	8 8888	88 8888	181
MTR	2	MIXTURE	4088 0000	88 8728	7 8888	82 8184	881
CUL	3	PARK	8718 0000	82 4888	8 2814	88 2810	1281
SPAC	2	3 CM SPACING	2887 0000	83 8828	7 8828	82 4183	881
MTR	1	MONOCULTURE	881 0000	84 2808	10 1388	102 7888	121
MTR	2	MIXTURE	2886 0000	83 8148	7 8112	88 4178	841
SPAC	3	9 CM SPACING	2188 0000	80 8818	8 2888	88 8812	821
MTR	1	MONOCULTURE	818 0000	81 8222	7 3841	84 8822	121
MTR	2	MIXTURE	2840 0000	80 8080	8 8820	74 8871	801
CUL	4	YONGE	4288 0000	88 8887	8 8778	34 8848	881
SPAC	2	3 CM SPACING	2117 0000	88 8217	8 8883	44 2228	481
MTR	1	MONOCULTURE	823 0000	88 2778	8 8182	42 4477	181

APPENDIX 7

02/10/82

PAGE 10

CRITERION VARIABLE 8

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
MTR	2	MIXTURE	1284 0000	45 8571	5 8822	47 0698	281
SPAC	3	9 CM SPACING	2109 0000	45 1875	5 0557	25 5558	461
MTR	1	MONOCULTURE	814 0000	45 2222	4 1045	17 5946	181
MTR	2	MIXTURE	1285 0000	45 1887	5 8775	31 1092	291
CUL	5	MOROMAY	4128 0000	42 8104	5 8878	32 3473	981
SPAC	2	3 CM SPACING	1850 0000	42 0555	5 5594	42 0254	281
MTR	1	MONOCULTURE	777 0000	42 1887	7 1145	50 8175	181
MTR	2	MIXTURE	773 0000	42 0444	5 1594	27 5279	181
SPAC	3	9 CM SPACING	2578 0000	42 8623	5 1825	28 5800	981
MTR	1	MONOCULTURE	784 0000	42 5555	5 4464	41 5555	181
MTR	2	MIXTURE	1795 0000	42 7381	4 8589	28 7824	421
TOTAL CASES =			570				

APPENDIX 7

02/10/82

PAGE 20

FILE SPHERS (CREATION DATE = 12/30/81)

CRITERION VARIABLE 8

CRITERION VARIABLE		DESCRIPTION OF SUBPOPULATIONS						
VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N	
FOR ENTIRE POPULATION			47694 0000	83 1649	12 0110	169 2873	5701	
CUL	1	PTIC 82	8472 0000	78 8288	8 9770	80 5872	421	
	2	3 CM SPACING	3768 0000	78 9375	8 3349	87 1227	481	
	MTR	2	MIXTURE	3769 0000	78 9375	8 3349	87 1227	481
SPAC	3	9 CM SPACING	2883 0000	78 9118	8 5880	73 7193	341	
	MTR	3	MIXTURE	2883 0000	78 9118	8 5880	73 7193	341
CUL	2	ELENLEA	18004 0000	84 1412	8 9153	78 4119	1701	
	2	3 CM SPACING	7865 0000	85 0595	7 4742	55 8638	881	
	MTR	1	MONOCULTURE	1891 0000	83 8444	7 7850	50 7814	181
MTR	2	MIXTURE	6284 0000	85 3628	7 4188	55 0042	651	
SPAC	3	9 CM SPACING	8019 0000	83 2442	10 0643	101 5936	851	
	MTR	1	MONOCULTURE	1880 0000	82 3233	10 5830	112 0000	181
	MTR	2	MIXTURE	6238 0000	83 2208	10 0298	100 5924	851
CUL	3	PARK	11278 0000	88 8084	11 2841	128 8784	1281	
	3	3 CM SPACING	8871 0000	88 9545	12 5181	158 8267	881	
	MTR	1	MONOCULTURE	1055 0000	87 5187	20 3850	410 2852	121
MTR	2	MIXTURE	4816 0000	89 1852	10 3258	108 0443	541	
SPAC	3	9 CM SPACING	8898 0000	86 8267	9 8517	97 2622	821	
	MTR	1	MONOCULTURE	1118 0000	83 1887	7 8067	62 5182	121
	MTR	2	MIXTURE	4380 0000	87 8000	10 0871	101 3489	801
CUL	4	10M000002	8835 0000	70 8857	8 9249	48 0029	941	
	3	3 CM SPACING	3217 0000	82 3348	8 2292	67 8845	481	
	MTR	1	MONOCULTURE	1313 0000	72 8444	7 0989	50 4055	181

APPENDIX 7

02/16/82

PAGE 21

CRITERION VARIABLE 9

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
MTR	2	MIXTURE	1004 0000	88 0000	8 4847	71 4818	281
SPAC	3	9 CM SPACING	2410 0000	71 2202	8 4187	20 2203	481
MTR	1	MONOCULTURE	1201 0000	72 2778	8 2006	28 0048	181
MTR	2	MIXTURE	2118 0000	70 8000	8 8747	28 0728	201
CUL	8	NORWAY	8813 0000	72 8104	8 4800	28 8200	881
SPAC	3	3 CM SPACING	2894 0000	72 0888	8 0884	28 0884	281
MTR	1	MONOCULTURE	1222 0000	72 4444	8 8488	21 0088	181
MTR	2	MIXTURE	1272 0000	70 8887	8 1482	27 7087	181
SPAC	3	9 CM SPACING	4210 0000	71 8633	8 1732	28 7034	881
MTR	1	MONOCULTURE	1205 0000	71 8444	8 7004	48 7028	181
MTR	2	MIXTURE	3024 0000	72 0000	8 4222	18 8810	421
TOTAL CASES =			870				

APPENDIX 7

02/16/82

PAGE 22

FILE SPHERE (CREATION DATE = 12/30/81)

CRITERION VARIABLE 9 DESCRIPTION OF SUBPOPULATIONS

CRITERION VARIABLE	HL	HEADLENGTH CM					
ORDERED DOWN BY	CUL	CULTIVAR					
BY	SPAC	INTER-PLANT SPACING					
BY	MTR	TREATMENT					
VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
FOR ENTIRE POPULATION			8081 0000	10 8884	2 0884	4 2763	8701
CUL	1	PITIC 82	1021 0000	12 4812	1 8180	2 6210	821
SPAC	2	3 CM SPACING	884 0000	12 1887	1 8880	2 8248	481
MTR	3	MIXTURE	884 0000	12 1887	1 8880	2 8248	481
SPAC	3	9 CM SPACING	437 0000	12 8828	1 8880	2 8838	241
MTR	8	MIXTURE	437 0000	12 8828	1 8880	2 8838	241
CUL	2	GLORIA	1889 0000	11 1708	1 4088	1 9788	1701
SPAC	2	3 CM SPACING	804 0000	10 8088	1 2032	1 8882	801
MTR	1	MONOCULTURE	182 0000	10 8887	1 1278	1 2841	181
MTR	4	MIXTURE	718 0000	10 8488	1 2801	1 8228	881
SPAC	3	9 CM SPACING	881 0000	11 8233	1 4202	2 0171	881
MTR	1	MONOCULTURE	210 0000	11 8887	1 8718	2 4708	181
MTR	8	MIXTURE	781 0000	11 4088	1 2878	1 8888	801
CUL	3	PARK	1018 0000	7 8287	1 2048	1 4817	1281
SPAC	2	3 CM SPACING	488 0000	7 4848	1 0708	1 1488	881
MTR	1	MONOCULTURE	81 0000	7 8833	0 8148	0 2882	121
MTR	2	MIXTURE	483 0000	7 4830	1 1888	1 2477	841
SPAC	3	9 CM SPACING	821 0000	8 4032	1 1888	1 3883	821
MTR	1	MONOCULTURE	110 0000	8 1887	0 7177	0 8182	121
MTR	2	MIXTURE	411 0000	8 2200	1 1830	1 3888	801
CUL	4	1000000002	1071 0000	11 2828	1 8212	2 8883	881
SPAC	2	3 CM SPACING	828 0000	11 4248	1 7884	2 8887	481
MTR	1	MONOCULTURE	218 0000	12 0000	1 7180	2 8412	181

APPENDIX 7

02/10/82

PAGE 23

CRITERION VARIABLE NL

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
MTR	2	MIXTURE	310 0000	11 0714	1 7108	2 9577	28
SPAC	3	9 CM SPACING	545 0000	11 3042	1 4048	2 2339	48
MTR	1	MONOCULTURE	218 0000	12 1111	1 4807	2 1048	18
MTR	2	MIXTURE	327 0000	10 9000	1 3481	1 8172	30
CUL	5	BOROMAY	1078 0000	11 1078	1 1203	1 2852	98
SPAC	2	3 CM SPACING	394 0000	10 8444	1 0940	1 1888	38
MTR	1	MONOCULTURE	267 0000	11 8000	0 7871	0 6000	18
MTR	2	MIXTURE	187 0000	10 3685	1 1448	1 3108	18
SPAC	3	9 CM SPACING	881 0000	11 3500	1 1172	1 2482	88
MTR	1	MONOCULTURE	205 0000	11 2889	1 1080	1 2281	18
MTR	2	MIXTURE	478 0000	11 2323	1 0888	1 2032	42
TOTAL CASES =			570				

APPENDIX 7

02/10/82

PAGE 24

FILE SPHERE (CREATION DATE = 12/30/81)

DESCRIPTION OF SUBPOPULATIONS

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
FOR ENTIRE POPULATION			23385 0000	40 8634	5 8208	48 0228	570
CUL	1	PITIC 82	3155 0000	36 4786	7 1081	51 3388	32
SPAC	2	3 CM SPACING	1782 0000	37 1250	6 1178	38 8088	48
MTR	2	MIXTURE	1782 0000	37 1250	6 1178	38 8088	48
SPAC	3	9 CM SPACING	1373 0000	40 3624	5 0782	25 7888	38
MTR	2	MIXTURE	1373 0000	40 3624	5 0782	25 7888	38
CUL	2	CLENLEA	7282 0000	42 8688	6 4782	41 8421	170
SPAC	2	3 CM SPACING	3404 0000	40 8228	5 8389	44 0788	84
MTR	1	MONOCULTURE	843 0000	38 7222	5 0604	25 6068	18
MTR	2	MIXTURE	2781 0000	41 8222	5 4388	41 4322	88
SPAC	3	9 CM SPACING	3888 0000	48 7442	5 8113	31 4887	88
MTR	1	MONOCULTURE	813 0000	48 1887	5 3818	40 8328	18
MTR	2	MIXTURE	3035 0000	48 8224	5 4322	38 9188	88
CUL	3	PARK	8878 0000	44 3884	7 8082	57 8888	128
SPAC	2	3 CM SPACING	2806 0000	42 8488	6 4274	71 0210	88
MTR	1	MONOCULTURE	485 0000	41 2800	12 2271	182 2648	12
MTR	2	MIXTURE	2313 0000	42 8222	7 4184	55 4072	84
SPAC	3	9 CM SPACING	2670 0000	48 3902	5 1228	27 4881	82
MTR	1	MONOCULTURE	809 0000	60 7500	5 0205	25 2088	12
MTR	2	MIXTURE	2781 0000	46 2200	5 8118	34 8808	80
CUL	4	TOMORRO2	2421 0000	38 3838	5 2632	27 8881	84
SPAC	2	3 CM SPACING	1828 0000	38 2478	6 8118	38 1430	88
MTR	1	MONOCULTURE	688 0000	38 8887	5 0381	36 4708	18

227

APPENDIX 7

02/10/82

PAGE 26

CRITERION VARIABLE EL

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
MTR	2	MIXTURE	930 0000	33 2142	5 0211	25 2118	281
SPAC	2	0 CM SPACING	1705 0000	37 3050	4 2212	17 0038	401
MTR	1	MONOCULTURE	705 0000	38 1867	4 2737	18 3047	181
MTR	2	MIXTURE	1020 0000	35 3232	3 8881	15 1004	301
CUL	5	BERGWAY	2850 0000	45 1875	4 1018	17 5708	601
SPAC	2	3 CM SPACING	1435 0000	29 8444	4 5184	21 3111	301
MTR	1	MONOCULTURE	752 0000	41 7778	4 2803	18 0854	181
MTR	2	MIXTURE	835 0000	35 1111	4 2225	15 0825	181
SPAC	2	0 CM SPACING	2421 0000	40 3500	3 8481	15 0672	601
MTR	1	MONOCULTURE	715 0000	35 7778	4 2731	18 1242	181
MTR	2	MIXTURE	1705 0000	40 8952	3 7510	14 2908	421
TOTAL CASES =			570				

APPENDIX 7

02/10/82

PAGE 26

FILE SPHERE CREATION DATE = 12/30/81

CRITERION VARIABLE EL

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
FOR ENTIRE POPULATION			507 8268	1 8920	0 5385	0 2901	5701
CUL	1	PYIC 82	134 9181	1 8482	0 4802	0 2307	821
SPAC	2	3 CM SPACING	73 8814	1 8269	0 4930	0 2430	481
MTR	2	MIXTURE	73 8814	1 8350	0 4930	0 2430	481
SPAC	2	0 CM SPACING	51 2387	1 8011	0 2287	0 1067	341
MTR	2	MIXTURE	51 2387	1 8011	0 2287	0 1067	341
CUL	2	GLEULEA	345 9283	2 9204	0 4819	0 2324	1701
SPAC	2	3 CM SPACING	158 9286	1 8862	0 4130	0 1897	801
MTR	1	MONOCULTURE	31 3233	1 7407	0 8045	0 3855	181
MTR	2	MIXTURE	125 6023	1 8020	0 2287	0 1104	801
SPAC	2	0 CM SPACING	158 2888	2 2008	0 4508	0 2032	801
MTR	1	MONOCULTURE	40 7887	2 2850	0 4825	0 2320	181
MTR	2	MIXTURE	145 4441	2 1836	0 4442	0 1973	801
CUL	3	PARE	20 2770	1 8105	0 2882	0 0878	1201
SPAC	2	3 CM SPACING	10 8807	0 8807	0 2418	0 0585	801
MTR	1	MONOCULTURE	10 1884	0 8800	0 2388	0 1148	121
MTR	2	MIXTURE	48 4555	0 8873	0 2181	0 0475	501
SPAC	2	0 CM SPACING	70 7214	1 1407	0 2951	0 0871	821
MTR	1	MONOCULTURE	15 5585	1 2005	0 2215	0 0481	121
MTR	2	MIXTURE	55 1818	1 1022	0 3090	0 0900	801
CUL	4	YONGGONGGONG	150 1081	1 5869	0 3895	0 1300	801
SPAC	2	3 CM SPACING	55 7416	1 4902	0 3401	0 1188	481
MTR	1	MONOCULTURE	29 1243	1 5188	0 3251	0 1057	181

FOOR COPY
COPIE DE QUALITEE INFERIEURE

228

APPENDIX Y

3/16/82

PAGE 37

CRITERION VARIABLE YH

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
MYR	2	MIXTURE	37 6923	1 3420	0 3067	0 0953	281
SPAC	3	0 CM SPACING	83 3886	1 7969	0 3250	0 1056	461
MYR	1	MONOCULTURE	32 8880	1 8216	0 3200	0 1024	181
MYR	2	MIXTURE	60 3864	1 8768	0 3197	0 1022	301
CUL	6	CONWAY	147 3272	1 8348	0 3549	0 1259	901
SPAC	2	3 CM SPACING	50 6418	1 4067	0 3897	0 1518	301
MYR	1	MONOCULTURE	28 8902	1 5553	0 3197	0 0953	181
MYR	2	MIXTURE	22 9617	1 2281	0 3301	0 1090	181
SPAC	3	0 CM SPACING	98 6982	1 8116	0 3277	0 1074	901
MYR	1	MONOCULTURE	20 1402	1 5765	0 3438	0 1181	181
MYR	2	MIXTURE	88 6981	1 8648	0 3564	0 0878	421
TOTAL CASES =			870				

Appendix 8. Distribution parameters of characters of single plants, grown in monoculture and in mixture, at two interplant spacings.

All plants were grown in four-row plots in 1978. The data are from randomly selected plants, which had both neighbours present, and which were harvested at maturity.

Character ¹	Spacing (cm)	Skewness ^{2,3}		Kurtosis ³	
		Monoculture	Mixture	Monoculture	Mixture
T	3	0.47	0.88**	1.21	1.66**
H	3	0.53	0.40*	1.02	0.60
Wt	3	0.72*	0.38*	1.44*	0.78*
Y	3	0.91**	0.43*	0.71	0.42
K/P	3	0.59	0.47*	0.97	0.40
Kwt	3	0.54	-0.88**	3.41**	2.18**
K/H	3	0.90**	-0.16	1.53	0.52
HI	3	-0.09	-0.64**	5.99**	1.51**
FL	3	-0.82**	-0.29	0.87	0.57
Ht	3	-1.15**	-0.86**	1.57*	1.29**
ExL	3	0.40	-0.54**	1.51*	-0.13
HL	3	0.33	-0.40	0.65	0.93
T	9	0.20	0.63**	-0.48	0.49
H	9	-0.09	0.72**	-0.75	0.93*
Wt	9	0.12	0.29	-0.13	0.03
Y	9	-0.15	0.32	-0.46	0.12
K/P	9	-0.03	0.47*	-0.51	0.31
Kwt	9	-0.16	-1.36**	-0.02	3.40**
K/H	9	-0.34	-0.09	1.86**	0.04
HI	9	-1.46**	-0.89**	3.10**	2.64**
FL	9	0.33	0.31	-0.23	-0.16
Ht	9	-0.93**	0.07	1.24*	-0.08
ExL	9	-0.53	-0.37	0.52	-0.12
HL	9	-0.28	-0.41	0.33	0.58

1. Character abbreviations defined in Table 1, HL = Head Length.
2. Values of skewness and kurtosis are averages of four genotypes and three replicates.
3. * significant, $\alpha \leq 0.05$; ** significant, $\alpha \leq 0.01$.

Appendix 9. Relative Yield Totals¹ for several characters of plants grown in mixtures at two interplant spacings in 1978.

Mixture	Character ²															
	Y		T		H		Kvt		HI		Mc		K/H		Y/H	
	3 cm	9 cm	3 cm	9 cm	3 cm	9 cm	3 cm	9 cm	3 cm	9 cm	3 cm	9 cm	3 cm	9 cm	3 cm	9 cm
Glenlea + Park	0.94	1.13	1.03	1.15	0.97	1.25*	1.00	0.97	0.97	1.03	0.96	1.08	0.97	0.87	0.99	0.85**
Glenlea + 70M09002	0.98	1.01	1.02	1.08	1.04	1.11	0.99	0.99	1.00	1.04	0.97	0.98	0.95	0.92	0.94	0.91
Glenlea + Norquay	0.88	1.12	0.99	1.12	0.95	1.23*	0.99	0.98	0.97	1.05	1.14	1.08	0.87	0.91	1.00	0.90
Park + 70M09002	0.78*	0.86	0.91	0.83	0.88	0.88	0.96	1.00	0.99	1.03	0.79**	0.84	0.92	0.98	0.92	0.92
Park + Norquay	1.00	0.93	1.11	0.95	1.00	0.97	0.99	1.01	0.99	0.99	0.97	0.98	1.02	0.95	1.05	0.95

Section 6.2.

* significant, $\alpha \leq 0.05$; ** significant, $\alpha \leq 0.01$.

2. Character abbreviations defined in Table 1.

POOR COPY
COPIE DE QUALITEE INFERIEURE

231

Appendix. 10. Descriptive statistics of characters of plants, grown
in monoculture and in mixture, in machine seeded plots,
measured at the flowering stage, in 1978.

POOR COPY
COPIE DE QUALITEE INFERIEURE

232

APPENDIX 10

03/16/82

PAGE 2

FILE FLAGMERS (CREATION DATE = 03/16/82)

CRITERION VARIABLE 0 DESCRIPTION OF SUBPOPULATIONS
BROKEN DOWN BY CUL HEIGHT OF THE FLAG LEAF BLADE CM
BY COMP CULTIVAR TREATMENT

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
FOR ENTIRE POPULATION			72272 0000	66 1666	12 2666	161 2632	1310
CUL	1	PITIC 62	16639 0000	61 6722	6 3692	70 6472	324
COMP	1	MONOCULTURE	4088 0000	61 2266	7 3329	63 7716	80
COMP	2	MIXTURE	12741 0000	62 2172	6 7206	76 0473	244
EUL	2	GLENLEA	16376 0000	67 8155	6 4666	72 0647	271
COMP	1	MONOCULTURE	4114 0000	66 6667	7 6426	66 6636	80
COMP	2	MIXTURE	12262 0000	67 8019	6 7436	76 4603	211
CUL	3	PARK	16663 0000	66 3112	13 0639	170 6662	266
COMP	1	MONOCULTURE	3232 0000	63 6667	6 4663	41 6766	80
COMP	2	MIXTURE	13731 0000	60 7666	13 6726	166 2627	226
CUL	4	YONGG8002	9696 0000	67 6666	7 7677	60 2366	216
COMP	1	MONOCULTURE	2776 0000	67 6136	7 6676	66 6616	86
COMP	2	MIXTURE	7217 0000	67 4663	7 6310	61 2241	162
EUL	5	BERGWAY	10086 0000	66 1006	7 6616	66 1662	216
COMP	1	MONOCULTURE	2666 0000	66 0000	7 6613	66 6621	86
COMP	2	MIXTURE	7441 0000	66 6667	7 6626	62 3166	160

TOTAL CASES = 1310

APPENDIX 10

03/10/82

PAGE 3

FILE PLANNERS (CREATION DATE = 03/10/82)

CRITERION VARIABLE
BROKEN DOWN BY
BY CUL
COMP

DESCRIPTION OF SUBPOPULATIONS
PLANT HEIGHT CM
CULTIVAR
TREATMENT

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
FOR ENTIRE POPULATION			92380 0000	73 7271	16 4848	271 0836	1253
CUL	1	PITIC 82	28816 0000	84 2489	11 8887	134 2880	324
COMP	1	MONOCULTURE	8142 0000	84 2780	10 8440	118 7718	80
COMP	2	MIXTURE	18674 0000	84 2277	11 8141	138 8728	244
CUL	2	GLENLEA	23130 0000	88 3808	11 8828	141 4881	271
COMP	1	MONOCULTURE	4877 0000	82 8800	12 2037	148 8297	80
COMP	2	MIXTURE	18183 0000	88 0332	11 7438	137 8178	211
CUL	3	PARK	21832 0000	88 0323	15 2384	232 2084	248
COMP	1	MONOCULTURE	8178 0000	88 3000	15 0607	228 8227	80
COMP	2	MIXTURE	18854 0000	88 8881	15 2821	233 8804	188
CUL	4	70M000002	13872 0000	84 8332	11 8278	132 1127	210
COMP	1	MONOCULTURE	3789 0000	82 8482	10 1107	102 2253	80
COMP	2	MIXTURE	8884 0000	84 8847	12 0884	148 4081	182
CUL	5	NORQUAY	13088 0000	88 1488	10 8877	120 8487	200
COMP	1	MONOCULTURE	2888 0000	88 4088	11 7724	128 8902	80
COMP	2	MIXTURE	8170 0000	88 0388	10 8887	118 4830	181
TOTAL CASES =	1210						
MISSING CASES =	87 00	4 4 PCT					

APPENDIX 10

03/10/82

PAGE 4

FILE PLANNERS (CREATION DATE = 03/10/82)

CRITERION VARIABLE
BROKEN DOWN BY
BY CUL
COMP

DESCRIPTION OF SUBPOPULATIONS
HEADLENGTH CM
CULTIVAR
TREATMENT

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
FOR ENTIRE POPULATION			14088 0000	11 2524	2 7777	7 7188	1252
CUL	1	PITIC 82	4188 0000	12 8323	1 8828	2 8048	224
COMP	1	MONOCULTURE	1031 0000	12 8878	1 4488	2 1011	80
COMP	2	MIXTURE	3127 0000	12 8188	1 8283	2 8448	244
CUL	2	GLENLEA	3027 0000	11 2088	2 0788	4 3201	271
COMP	1	MONOCULTURE	888 0000	11 1323	1 8848	3 4288	80
COMP	2	MIXTURE	2389 0000	11 2278	2 1418	4 8881	211
CUL	3	PARK	2028 0000	8 2810	3 8888	12 8838	248
COMP	1	MONOCULTURE	481 0000	8 0187	1 1122	1 2370	80
COMP	2	MIXTURE	1887 0000	8 3282	4 0740	16 8873	188
CUL	4	70M000002	2828 0000	12 0428	1 8278	3 7841	210
COMP	1	MONOCULTURE	782 0000	12 1024	1 8228	3 3224	80
COMP	2	MIXTURE	1827 0000	12 0187	1 8848	3 8400	182
CUL	5	NORQUAY	2328 0000	11 8200	1 8380	2 8788	200
COMP	1	MONOCULTURE	704 0000	11 8222	1 8828	3 8818	80
COMP	2	MIXTURE	1822 0000	11 8028	1 8083	2 2881	181
TOTAL CASES =	1210						
MISSING CASES =	87 00	4 4 PCT					

APPENDIX 10

03/16/82

PAGE 8

FILE PLASMERS (CREATION DATE = 03/16/82)

----- D E S C R I P T I O N O F S U B P O P U L A T I O N S -----
CRITERION VARIABLE EL EXTRUSION LENGTH CM
BROKEN DOWN BY CUL CULTIVAR
BY COMP TREATMENT

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
FOR ENTIRE POPULATION			20106 0000	16 2612	21 7046	476 0001	(1200)
CUL	1	PITIC 82	3977 0000	12 3127	6 1662	26 8174	(323)
COMP	1	MONOCULTURE	1044 0000	13 0800	6 3289	28 8038	(80)
COMP	2	MIXTURE	2933 0000	12 0760	6 1286	26 2716	(243)
CUL	2	GLENLEA	4762 0000	17 8261	6 8612	87 2422	(271)
COMP	1	MONOCULTURE	882 0000	16 2822	7 6262	58 2421	(80)
COMP	2	MIXTURE	3880 0000	18 8212	10 3484	104 8893	(211)
CUL	3	PARK	4889 0000	17 9246	40 8848	1647 8483	(288)
COMP	1	MONOCULTURE	1846 0000	22 4322	11 6309	122 8616	(80)
COMP	2	MIXTURE	3043 0000	12 8226	44 4118	1972 4069	(226)
CUL	4	TOMCOCOCOC	3677 0000	17 8322	6 3246	40 0037	(210)
COMP	1	MONOCULTURE	820 0000	16 8346	6 2894	38 8826	(80)
COMP	2	MIXTURE	2857 0000	17 4146	6 3128	28 8892	(182)
CUL	5	MORQUAY	2833 0000	13 2827	21 8140	487 1882	(218)
COMP	1	MONOCULTURE	1264 0000	20 4084	6 4704	41 8882	(80)
COMP	2	MIXTURE	1729 0000	10 8882	24 8882	600 2678	(180)

TOTAL CASES = 1210
MISSING CASES = 1 OR 0 1 PCT

APPENDIX 10

03/16/82

PAGE 8

FILE PLASMERS (CREATION DATE = 03/16/82)

----- D E S C R I P T I O N O F S U B P O P U L A T I O N S -----
CRITERION VARIABLE AREA FLAGLEAF BLADE AREA CM SQ
BROKEN DOWN BY CUL CULTIVAR
BY COMP TREATMENT

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
FOR ENTIRE POPULATION			31216 8827	24 1698	6 0616	64 8916	(1200)
CUL	1	PITIC 82	8736 8882	27 0462	6 7224	46 1812	(323)
COMP	1	MONOCULTURE	2170 8886	27 4772	6 1688	38 2882	(78)
COMP	2	MIXTURE	6566 8887	26 8062	6 6927	47 8086	(244)
CUL	2	GLENLEA	7846 8886	27 8886	7 2886	82 2877	(270)
COMP	1	MONOCULTURE	1846 8887	27 4480	7 4641	86 7123	(80)
COMP	2	MIXTURE	6001 3088	26 1018	7 2827	82 7616	(210)
CUL	3	PARK	4821 8880	17 7878	6 1316	66 1207	(277)
COMP	1	MONOCULTURE	1006 2886	16 7717	6 6489	42 8492	(80)
COMP	2	MIXTURE	3815 2882	18 0429	6 5107	72 4316	(217)
CUL	4	TOMCOCOCOC	4640 1880	22 0862	6 6444	44 1476	(210)
COMP	1	MONOCULTURE	1246 8887	23 2666	7 2016	51 8621	(80)
COMP	2	MIXTURE	3394 1883	21 8826	6 3886	40 8111	(182)
CUL	5	MORQUAY	8473 8886	24 8812	6 4462	41 8624	(218)
COMP	1	MONOCULTURE	1488 7887	26 1820	6 4011	29 1721	(80)
COMP	2	MIXTURE	3985 2882	24 8206	6 6042	46 2978	(180)

TOTAL CASES = 1210
MISSING CASES = 11 OR 0 8 PCT

POOR COPY
COPIE DE QUALITEE INFERIEURE

235

Appendix 11. Descriptive statistics of characters of plants, grown in
monoculture and in mixture, in machine seeded plots,
measured at the middough stage, in 1978.

APPENDIX 11

03/18/82

PAGE 2

FILE PLASMERS (CREATION DATE = 03/18/82)

CRITERION VARIABLE BROKEN DOWN BY		DESCRIPTION OF SUBPOPULATIONS					
VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
FOR ENTIRE POPULATION			66030 0000	64 3972	11 8228	132 7712	(1260)
CUL	1	PITIC 82	15088 0000	82 2882	8 4822	71 8480	(288)
COMP	1	MONOCULTURE	3128 0000	82 3187	9 2782	85 0064	(80)
COMP	2	MIXTURE	11920 0000	82 2607	8 2628	68 9983	(228)
CUL	2	GLENLEA	20114 0000	86 8028	10 8882	111 4881	(302)
COMP	1	MONOCULTURE	3548 0000	81 7787	12 2428	142 2472	(88)
COMP	2	MIXTURE	16488 0000	87 7727	8 7828	88 1180	(242)
CUL	3	PARK	14827 0000	88 0844	8 0814	62 4711	(284)
COMP	1	MONOCULTURE	2048 0000	81 2280	8 0208	64 3227	(40)
COMP	2	MIXTURE	12488 0000	88 7800	8 1048	62 8880	(224)
CUL	4	TOMORROW	10558 0000	47 7847	7 1822	51 0282	(221)
COMP	1	MONOCULTURE	2821 0000	47 1822	8 8287	48 1184	(80)
COMP	2	MIXTURE	7728 0000	47 8814	7 2280	62 2424	(161)
CUL	5	HORQUAY	8784 0000	48 1782	6 8821	64 8824	(184)
COMP	1	MONOCULTURE	2848 0000	42 8782	7 7881	60 8884	(88)
COMP	2	MIXTURE	5218 0000	48 7278	6 0887	37 2088	(128)
TOTAL CASES =			1260				

APPENDIX 11

03/18/82

PAGE 2

FILE PLASMERS (CREATION DATE = 03/18/82)

CRITERION VARIABLE BROKEN DOWN BY		DESCRIPTION OF SUBPOPULATIONS					
VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
FOR ENTIRE POPULATION			108208 0000	88 2887	14 7788	218 2180	(1260)
CUL	1	PITIC 82	23222 0000	80 9782	10 8888	118 7881	(288)
COMP	1	MONOCULTURE	4872 0000	81 2880	11 2282	128 0880	(80)
COMP	2	MIXTURE	18480 0000	80 8211	10 7182	114 8388	(228)
CUL	2	GLENLEA	28888 0000	98 3078	13 0818	171 2885	(302)
COMP	1	MONOCULTURE	5888 0000	94 2202	13 8148	182 6220	(88)
COMP	2	MIXTURE	23000 0000	98 3004	12 7187	161 7182	(242)
CUL	3	PARK	24248 0000	81 8822	12 7222	162 1028	(284)
COMP	1	MONOCULTURE	3838 0000	88 4000	14 1282	188 3388	(40)
COMP	2	MIXTURE	20410 0000	82 4888	12 3881	142 7120	(224)
CUL	4	TOMORROW	18484 0000	74 8228	8 8288	72 7080	(221)
COMP	1	MONOCULTURE	4482 0000	74 8822	9 4027	88 4088	(80)
COMP	2	MIXTURE	12001 0000	74 8484	8 2088	67 2274	(161)
CUL	5	HORQUAY	14482 0000	74 4848	8 8181	88 8180	(184)
COMP	1	MONOCULTURE	4328 0000	74 8287	11 0882	122 2388	(88)
COMP	2	MIXTURE	10124 0000	74 4812	6 8274	77 8224	(128)
TOTAL CASES =			1260				

APPENDIX 11

03/10/82

PAGE 4

FILE PLASMERS (CREATION DATE = 03/10/82)

DESCRIPTION OF SUBPOPULATIONS

CRITERION VARIABLE BROKEN DOWN BY	EL CUL COMP	DESCRIPTION CULTIVAR TREATMENT	SUM	MEAN	STD DEV	VARIANCE	N
VARIABLE	CODE	VALUE LABEL					
FOR ENTIRE POPULATION			14306 0000	11 2784	2 8288	6 3761	(1288)
CUL	1	PITIC 82	3226 0000	12 2647	2 2812	5 0685	(288)
COMP	1	MONOCULTURE	802 0000	12 2667	1 6260	2 3276	(80)
COMP	2	MIXTURE	3024 0000	12 2632	2 4060	5 7863	(228)
CUL	2	GLEHLEA	3226 0000	11 0430	1 7005	2 8916	(302)
COMP	1	MONOCULTURE	800 0000	8 9881	2 1612	4 7878	(80)
COMP	2	MIXTURE	2747 0000	11 2045	1 4609	2 1062	(242)
CUL	3	PARK	2147 0000	8 1328	1 0828	1 2029	(284)
COMP	1	MONOCULTURE	207 0000	7 8760	1 1851	1 4048	(40)
COMP	2	MIXTURE	1840 0000	8 2143	1 0824	1 1286	(224)
CUL	4	700000002	2746 0000	12 4344	1 9280	3 7186	(221)
COMP	1	MONOCULTURE	777 0000	12 9809	1 9851	3 9806	(80)
COMP	2	MIXTURE	1871 0000	12 2422	1 8724	3 5097	(181)
CUL	5	NOROUAY	2252 0000	11 8134	1 8400	2 7047	(184)
COMP	1	MONOCULTURE	885 0000	11 4855	1 8806	2 8848	(88)
COMP	2	MIXTURE	1866 0000	11 8768	1 8224	2 8363	(126)
TOTAL CASES = 1288							

APPENDIX 11

03/10/82

PAGE 5

FILE PLASMERS (CREATION DATE = 03/10/82)

DESCRIPTION OF SUBPOPULATIONS

CRITERION VARIABLE BROKEN DOWN BY	EL CUL COMP	DESCRIPTION CULTIVAR TREATMENT	SUM	MEAN	STD DEV	VARIANCE	N
VARIABLE	CODE	VALUE LABEL					
FOR ENTIRE POPULATION			30176 0000	30 8203	7 8248	68 1298	(1267)
CUL	1	PITIC 82	8263 0000	28 8910	5 5854	31 2084	(288)
COMP	1	MONOCULTURE	1733 0000	28 8633	4 9290	24 2985	(80)
COMP	2	MIXTURE	6530 0000	28 6404	5 7889	33 2794	(228)
CUL	2	GLEHLEA	8575 0000	31 8106	7 6403	58 2740	(302)
COMP	1	MONOCULTURE	1814 0000	32 4407	8 2321	67 7680	(80)
COMP	2	MIXTURE	7861 0000	31 8670	7 4890	56 2348	(242)
CUL	3	PARK	8712 0000	38 7879	8 8672	78 4481	(284)
COMP	1	MONOCULTURE	1487 0000	37 1760	8 9811	80 6809	(40)
COMP	2	MIXTURE	8225 0000	38 7188	8 8624	78 3824	(224)
CUL	4	700000002	8928 0000	38 8888	5 2608	27 8872	(221)
COMP	1	MONOCULTURE	1882 0000	27 7000	5 4898	30 2476	(80)
COMP	2	MIXTURE	4376 0000	38 9899	5 1692	26 8206	(181)
CUL	5	NOROUAY	8686 0000	39 4716	5 2785	28 8963	(182)
COMP	1	MONOCULTURE	1783 0000	38 7414	8 2263	68 7818	(88)
COMP	2	MIXTURE	3905 0000	38 8269	4 8693	23 8940	(126)
TOTAL CASES = 1268							
MISSING CASES = 2 OR 0.2 PCT							

POOR COPY
COPIE DE QUALITEE INFERIEURE

238

APPENDIX 11

03/10/82

PAGE 8

FILE PLASMERS (CREATION DATE = 03/10/82)

CRITERION VARIABLE	DESCRIPTION	FLAGSLAP	SLADE AREA CM SQ	POPULATION		
BROKEN DOWN BY	CUL	CULTIVAR	TREATMENT			
	BY	COMP				
VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE
FOR ENTIRE POPULATION			34578 1030.	27 4000	9 1000	62 9360
CUL	1	PITIC 62	8878 1063	30 2411	7 7867	60 3811
COMP	1	MONOCULTURE	1708 3000	20 8723	6 6212	43 8400
COMP	2	MIXTURE	8860 7987	30 2110	6 6810	44 8280
CUL	2	SLERLEA	8811 8883	33 8387	8 8873	67 8886
COMP	1	MONOCULTURE	1720 1007	20 1000	11 8810	120 7888
COMP	2	MIXTURE	8191 8886	33 8888	8 1886	64 8116
CUL	3	PARK	4888 7888	18 3877	5 8834	38 8846
COMP	1	MONOCULTURE	881 3888	18 6388	5 4182	28 3873
COMP	2	MIXTURE	4364 3888	18 8276	5 8282	38 8483
CUL	4	708888882	5784 8887	28 8488	7 8881	58 8814
COMP	1	MONOCULTURE	1721 2888	28 8883	7 8438	48 8017
COMP	2	MIXTURE	4023 8881	28 8884	7 4288	58 1687
CUL	5	NORWAY	8288 3888	27 1884	6 2823	38 8830
COMP	1	MONOCULTURE	1828 8887	28 3882	6 8118	47 7731
COMP	2	MIXTURE	3630 4882	28 8848	6 6787	38 7654
TOTAL CASES =	1288					
MISSING CASES =	11 OR	0.8 PCT				

Appendix 12. Variation in thousand kernel weights (g) among the
seed classes used in 1979.

Genotype	Seed class		
	Small	Large	Unsorted ¹
Pitic 62	29.8	40.8	--- ²
Glenlea	32.0	45.6	44.5
Park	25.8	36.0	34.9
70M009002	25.6	46.2	45.9
Norquay	25.5	38.0	37.3

1 Only sorted small and large seeds were used in the
seed size experiment.

2 No data available.

POOR COPY
COPIE DE QUALITEE INFERIEURE

240

Appendix 13. Descriptive statistics of characters of plants from
large, small and mixed size seeds, measured at the 3-5
leaf stage, in 1979.

POOR COPY
COPIE DE QUALITEE INFERIEURE

241

APPENDIX 13

02/10/83

PAGE 2

FILE SOL1 (CREATION DATE = 12/30/81)

CRITERION VARIABLE BROKEN DOWN BY		DESCRIPTION OF CULTIVAR TREATMENT		SUM POPULATION			
VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
FOR ENTIRE POPULATION							
			8802 5000	17 8478	3 7881	14 1008	482
CUL	0		1780 5000	17 1202	3 0173	9 1063	104
TR	1	LARGE SEED	749 0000	17 4323	3 1803	10 0447	42
TR	2	SMALL SEED	828 0000	17 3611	2 7376	7 4844	36
TR	3	MIXED SEED	405 5000	18 8346	2 7030	7 3112	36
CUL	1	PITIC 62	1480 5000	18 8239	2 8636	8 1432	106
TR	1	LARGE SEED	689 0000	18 1316	1 9408	3 7686	36
TR	2	SMALL SEED	575 0000	18 9118	3 0632	9 3877	34
TR	3	MIXED SEED	566 5000	17 7727	3 3728	11 3784	32
CUL	2	CLENLEA	2128 0000	19 1712	3 1856	10 1677	117
TR	1	LARGE SEED	748 0000	19 8318	3 8742	15 0092	36
TR	2	SMALL SEED	748 5000	18 8447	3 6732	9 4447	36
TR	3	MIXED SEED	632 5000	18 1671	2 1817	4 8731	36
CUL	3	PARK	1878 5000	19 1788	4 4888	19 8108	88
TR	1	LARGE SEED	815 0000	18 6384	4 8743	21 8480	33
TR	2	SMALL SEED	806 5000	19 6323	5 4180	29 2586	31
TR	3	MIXED SEED	857 0000	19 2624	3 1468	9 8946	36
CUL	4	70M000002	884 0000	18 0626	4 0637	16 4326	64
TR	1	LARGE SEED	281 0000	18 2609	3 8986	15 9870	23
TR	2	SMALL SEED	228 0000	18 0626	3 7864	14 3282	16
TR	3	MIXED SEED	375 0000	18 8200	4 3120	18 8033	25

TOTAL CASES = 482

APPENDIX 13

02/16/62

PAGE 3

FILE SOL1 (CREATION DATE = 12/30/61)

DESCRIPTION OF SUBPOPULATIONS							
CRITERION VARIABLE BROKEN DOWN BY	L1 CUL TR	NUMBER OF CULTIVAR TREATMENT					
VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
FOR ENTIRE POPULATION			1762 0000	3 8888	0 8748	0 4640	(482)
CUL	0		307 0000	3 8192	0 4976	0 2476	(104)
TR	1	LARGE SEED	180 0000	3 7142	0 5982	0 3564	(42)
TR	2	SMALL SEED	140 0000	3 8888	0 3884	0 1507	(30)
TR	3	MIXED SEED	101 0000	3 8846	0 4318	0 1862	(26)
CUL	1	PITIC 62	300 0000	3 8871	0 8176	0 3812	(100)
TR	1	LARGE SEED	120 0000	3 3188	0 4711	0 2219	(30)
TR	2	SMALL SEED	120 0000	3 7847	0 6989	0 4884	(30)
TR	3	MIXED SEED	130 0000	3 9394	0 4982	0 2482	(33)
CUL	2	ELLENLEA	401 0000	3 8126	0 8204	0 3640	(111)
TR	1	LARGE SEED	140 0000	3 8842	0 8819	0 4381	(38)
TR	2	SMALL SEED	120 0000	3 3844	0 6334	0 4011	(30)
TR	3	MIXED SEED	130 0000	3 8000	0 4726	0 2228	(36)
CUL	3	PARK	348 0000	3 8810	0 7818	0 8788	(88)
TR	1	LARGE SEED	130 0000	3 8264	0 7424	0 5511	(32)
TR	2	SMALL SEED	111 0000	3 8808	0 8808	0 7808	(21)
TR	3	MIXED SEED	117 0000	3 4412	0 8808	0 4388	(34)
CUL	4	700000002	232 0000	3 8280	0 8887	0 8086	(64)
TR	1	LARGE SEED	88 0000	3 8281	1 1140	1 2411	(22)
TR	2	SMALL SEED	55 0000	3 8000	0 8226	0 4000	(16)
TR	3	MIXED SEED	88 0000	3 8200	0 8226	0 8787	(26)
TOTAL CASES =			482				

APPENDIX 13

02/16/62

PAGE 4

FILE SOL1 (CREATION DATE = 12/30/61)

DESCRIPTION OF SUBPOPULATIONS							
CRITERION VARIABLE BROKEN DOWN BY	L1 CUL TR	TILLAGE PER PLANT CULTIVAR TREATMENT					
VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
FOR ENTIRE POPULATION			821 0000	1 8106	0 8948	0 8008	(482)
CUL	0		223 0000	2 1442	0 8178	0 6882	(104)
TR	1	LARGE SEED	100 0000	2 2810	0 8084	0 6269	(42)
TR	2	SMALL SEED	70 0000	1 8444	0 7808	0 6254	(30)
TR	3	MIXED SEED	53 0000	2 0288	0 8847	0 3888	(26)
CUL	1	PITIC 62	223 0000	2 1238	0 8888	0 6018	(105)
TR	1	LARGE SEED	88 0000	2 2832	0 7847	0 6318	(38)
TR	2	SMALL SEED	88 0000	2 0000	0 8211	0 8488	(34)
TR	3	MIXED SEED	89 0000	2 0000	0 8788	0 8807	(33)
CUL	2	ELLENLEA	178 0000	1 8888	0 7188	0 8178	(111)
TR	1	LARGE SEED	87 0000	1 8000	0 7260	0 5270	(38)
TR	2	SMALL SEED	82 0000	1 8878	0 8148	0 8838	(30)
TR	3	MIXED SEED	89 0000	1 8000	0 8038	0 3847	(36)
CUL	3	PARK	188 0000	1 8828	0 8787	0 8820	(88)
TR	1	LARGE SEED	88 0000	1 8887	0 8881	0 3842	(32)
TR	2	SMALL SEED	80 0000	1 8128	0 8427	0 7118	(21)
TR	3	MIXED SEED	81 0000	1 7841	0 8818	0 3802	(34)
CUL	4	700000002	123 0000	2 0781	1 2782	1 8287	(64)
TR	1	LARGE SEED	81 0000	2 8522	1 8844	1 8844	(22)
TR	2	SMALL SEED	28 0000	1 8828	0 8828	0 8282	(16)
TR	3	MIXED SEED	47 0000	1 8880	1 1882	1 3880	(26)
TOTAL CASES =			482				

APPENDIX 13

02/16/82

PAGE 8

FILE S01 (CREATION DATE 4/12/80/81)

CRITERION VARIABLE		DESCRIPTION		SUB POPULATIONS			
BROKEN DOWN BY		BY		CULTIVAR TREATMENT			
VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
FOR ENTIRE POPULATION			44 3170	0 0018	0 0401	0 0018	(402)
CUL	0		10 0200	0 1000	0 0400	0 0018	1000
TR	1	LARGE SEED	4 0030	0 1100	0 0000	0 0000	400
TR	2	SMALL SEED	3 0030	0 0007	0 0000	0 0000	300
TR	3	MIXED SEED	3 1000	0 0000	0 0000	0 0000	300
CUL	1	PITIC SZ	0 3420	0 0000	0 0000	0 0000	1000
TR	1	LARGE SEED	2 0070	0 0000	0 0000	0 0000	300
TR	2	SMALL SEED	2 0000	0 0000	0 0000	0 0000	300
TR	3	MIXED SEED	2 0700	0 0000	0 0000	0 0000	300
CUL	2	GLENEA	10 0070	0 0001	0 0001	0 0000	1100
TR	1	LARGE SEED	2 0000	0 1000	0 0400	0 0000	300
TR	2	SMALL SEED	2 0000	0 1000	0 0000	0 0000	300
TR	3	MIXED SEED	2 0000	0 0001	0 0000	0 0000	300
CUL	3	PARK	0 0000	0 0001	0 0000	0 0000	900
TR	1	LARGE SEED	2 0000	0 0000	0 0000	0 0000	300
TR	2	SMALL SEED	2 0000	0 0000	0 0000	0 0000	300
TR	3	MIXED SEED	2 0000	0 0001	0 0000	0 0000	300
CUL	4	TOMORROW	0 3000	0 0000	0 0000	0 0000	000
TR	1	LARGE SEED	2 0000	0 0000	0 0000	0 0000	300
TR	2	SMALL SEED	0 0710	0 0000	0 0000	0 0000	100
TR	3	MIXED SEED	2 1000	0 0000	0 0000	0 0000	300
TOTAL CASES =			402				

POOR COPY
COPIE DE QUALITEE INFERIEURE

244

Appendix 14. Descriptive statistics of characters of plants from large, small and mixed size seeds, measured at the jointing stage, in 1979.

APPENDIX 14

02/10/82

PAGE 2

FILE SOL2 (CREATION DATE = 12/30/81)

CRITERION VARIABLE		DESCRIPTION OF SUBPOPULATIONS					
BROKEN DOWN BY		CULTIVAR					
BY		TREATMENT					
VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
FOR ENTIRE POPULATION			3220 5000	9 8244	4 8401	23 4204	(407)
CUL	0		874 0000	8 0042	4 0884	24 8438	(88)
TR	1	LARGE SEED	343 0000	8 0077	4 0470	24 4818	(35)
TR	2	SMALL SEED	277 0000	8 1471	4 2475	18 2504	(34)
TR	3	MIXED SEED	205 0000	8 8348	6 0818	25 7708	(23)
CUL	1	PITIC 82	308 0000	7 8827	3 8283	15 4317	(52)
TR	1	LARGE SEED	228 0000	8 0714	3 8926	15 1814	(28)
TR	2	SMALL SEED	88 0000	8 1428	3 8186	15 3828	(14)
TR	3	MIXED SEED	87 0000	8 7500	3 7878	14 3472	(10)
CUL	2	GLENLEA	467 0000	8 2288	8 2888	27 8812	(84)
TR	1	LARGE SEED	261 0000	7 8182	4 9822	24 8321	(27)
TR	2	SMALL SEED	247 0000	8 1820	8 8790	31 1247	(27)
TR	3	MIXED SEED	233 0000	11 2867	4 8528	23 8471	(30)
CUL	3	PARK	838 0000	10 8128	4 8288	21 3882	(88)
TR	1	LARGE SEED	248 0000	12 4843	8 2888	29 0387	(28)
TR	2	SMALL SEED	244 0000	10 1428	4 2771	18 2827	(28)
TR	3	MIXED SEED	306 0000	10 1823	3 8888	15 1888	(30)
CUL	4	70M000002	849 3000	10 7808	4 2227	17 8388	(78)
TR	1	LARGE SEED	228 8000	11 7428	4 0818	18 4144	(28)
TR	2	SMALL SEED	181 0000	8 0800	4 0888	18 4711	(28)
TR	3	MIXED SEED	239 5000	10 8818	4 2788	18 2882	(31)
TOTAL CASES =			407				

APPENDIX 14

02/10/82

PAGE 3

FILE SOL2 (CREATION DATE = 12/30/81)

CRITERION VARIABLE		DESCRIPTION OF SUBPOPULATIONS					
BROKEN DOWN BY		TILLERS PER PLANT					
BY		CULTIVAR					
		TREATMENT					
VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
FOR ENTIRE POPULATION			2241 0000	8 5081	2 0843	4 3442	(407)
CUL	0		504 0000	8 2800	1 8723	2 8000	(88)
TR	1	LARGE SEED	208 0000	8 2844	1 3711	1 8798	(29)
TR	2	SMALL SEED	171 0000	8 0284	1 7838	2 1808	(34)
TR	3	MIXED SEED	128 0000	8 5852	1 9731	2 8822	(23)
CUL	1	PITIC 82	281 0000	8 8882	3 2088	10 2488	(52)
TR	1	LARGE SEED	148 0000	8 2887	3 3874	11 8488	(28)
TR	2	SMALL SEED	83 0000	8 8428	2 0222	4 0824	(14)
TR	3	MIXED SEED	80 0000	8 0000	3 8872	18 1111	(10)
CUL	2	GLENLEA	483 0000	8 1382	1 7888	3 2287	(84)
TR	1	LARGE SEED	174 0000	4 7827	1 8844	2 7782	(27)
TR	2	SMALL SEED	168 0000	8 7774	2 1842	4 8410	(27)
TR	3	MIXED SEED	153 0000	8 1888	1 8784	2 1827	(30)
CUL	3	PARK	489 0000	8 8880	1 8182	3 2287	(88)
TR	1	LARGE SEED	194 0000	8 8088	1 8887	2 7778	(28)
TR	2	SMALL SEED	180 0000	8 8788	2 1882	4 8188	(28)
TR	3	MIXED SEED	175 0000	8 8887	1 8888	2 4884	(30)
CUL	4	70M000002	874 0000	8 0000	2 1284	4 8288	(78)
TR	1	LARGE SEED	178 0000	8 3871	2 0224	4 0888	(28)
TR	2	SMALL SEED	118 0000	8 0800	1 8187	3 8727	(28)
TR	3	MIXED SEED	178 0000	8 7418	2 3880	5 8878	(31)
TOTAL CASES =			407				

APPENDIX 14

02/10/82

PAGE 4

FILE S0L2 (CREATION DATE = 12/30/81)

DESCRIPTION OF SUBPOPULATIONS							
CRITERION VARIABLE BROKEN DOWN BY	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
FOR ENTIRE POPULATION			918 0000	2 2508	0 8281	0 6858	407
CUL	0		211 0000	2 1978	0 8287	0 6867	98
TR	1	LARGE SEED	81 0000	2 2322	0 8883	0 8070	38
TR	2	SMALL SEED	70 0000	2 0588	0 8143	0 6631	34
TR	3	MIXED SEED	60 0000	2 1728	0 7188	0 5168	23
CUL	1	PITIC 82	90 0000	1 7308	0 8593	0 7381	52
TR	1	LARGE SEED	81 0000	1 8214	0 7228	0 5228	28
TR	2	SMALL SEED	23 0000	1 8428	0 7448	0 5548	14
TR	3	MIXED SEED	16 0000	1 8000	0 5184	0 2687	10
CUL	2	CLNLEA	203 0000	2 1888	0 8238	0 6787	84
TR	1	LARGE SEED	87 0000	1 8108	0 8078	0 6523	37
TR	2	SMALL SEED	58 0000	2 1882	0 7287	0 5312	27
TR	3	MIXED SEED	77 0000	2 8887	0 8281	0 6858	30
CUL	3	PARK	218 0000	2 4618	0 7812	0 6100	88
TR	1	LARGE SEED	77 0000	2 7800	0 7818	0 6122	28
TR	2	SMALL SEED	89 0000	2 4842	0 8222	0 6768	28
TR	3	MIXED SEED	54 0000	2 1222	0 8712	0 7588	30
CUL	4	YOM008002	202 0000	2 8578	0 7827	0 6122	79
TR	1	LARGE SEED	78 0000	2 8788	0 7724	0 5968	28
TR	2	SMALL SEED	80 0000	2 8000	0 7888	0 6222	20
TR	3	MIXED SEED	77 0000	2 4828	0 7888	0 6222	31
TOTAL CASES =			407				

APPENDIX 14

02/10/82

PAGE 5

FILE S0L2 (CREATION DATE = 12/30/81)

DESCRIPTION OF SUBPOPULATIONS							
CRITERION VARIABLE BROKEN DOWN BY	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
FOR ENTIRE POPULATION			548 8888	1 3412	0 8488	0 7200	407
CUL	0		122 3000	1 3888	0 4728	0 2248	98
TR	1	LARGE SEED	55 8800	1 4348	0 3838	0 1474	38
TR	2	SMALL SEED	48 8000	1 3878	0 5184	0 2687	34
TR	3	MIXED SEED	30 8500	1 3412	0 8887	0 7888	23
CUL	1	PITIC 82	78 1700	1 4488	0 8280	0 6858	52
TR	1	LARGE SEED	28 2400	1 3887	0 8822	0 7888	28
TR	2	SMALL SEED	20 4200	1 4888	0 3388	0 1128	14
TR	3	MIXED SEED	18 8100	1 8810	0 8287	0 6858	10
CUL	2	CLNLEA	128 8700	1 4422	0 8820	0 7888	84
TR	1	LARGE SEED	80 8800	1 3888	0 8811	0 7888	37
TR	2	SMALL SEED	38 2200	1 4888	0 8822	0 7888	27
TR	3	MIXED SEED	48 8800	1 8830	0 4278	0 1831	30
CUL	3	PARK	88 1400	1 1178	0 4822	0 2328	88
TR	1	LARGE SEED	30 8200	1 1007	0 8224	0 6734	28
TR	2	SMALL SEED	32 1200	1 1471	0 8281	0 6858	28
TR	3	MIXED SEED	23 2000	1 1087	0 4080	0 1684	30
CUL	4	YOM008002	108 8800	1 3388	0 8282	0 7200	79
TR	1	LARGE SEED	42 2800	1 8088	0 8280	0 6858	28
TR	2	SMALL SEED	18 1800	0 8878	0 3888	0 1488	20
TR	3	MIXED SEED	48 1800	1 4282	0 8027	0 6427	31
TOTAL CASES =			407				

POOR COPY
COPIE DE QUALITEE INFERIEURE

247

Appendix 15. Descriptive study of characters of plants from large, small and mixed size seeds, measured at the heading stage, in 1979.

POOR COPY
COPIE DE QUALITEE INFERIEURE

248

APPENDIX 15

02/18/82

PAGE 2

FILE 00L3 (CREATION DATE = 12/30/81)

CRITERION VARIABLE		DESCRIPTION		SUBPOPULATIONS			
BROKEN DOWN BY		HT1	PLANT HEIGHT CM				
		CUL	CULTIVAR				
		BY	TREATMENT				
VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
FOR ENTIRE POPULATION			13481 0000	28 4881	12 8408	159 7908	487
CUL	0		2359 0000	28 8487	9 2787	86 0579	92
TR	1	LARGE SEED	844 0000	24 5694	7 8887	61 9165	28
TR	2	SMALL SEED	800 0000	28 0188	9 2884	86 4594	32
TR	3	MIXED SEED	674 0000	28 1042	11 0029	121 0648	24
CUL	1	PITIC 82	1707 0000	18 1084	3 5036	12 2781	108
TR	1	LARGE SEED	634 0000	18 1870	3 0284	9 8459	25
TR	2	SMALL SEED	608 0000	18 4385	3 9788	16 8290	31
TR	3	MIXED SEED	663 0000	18 8000	3 4858	12 2180	42
CUL	2	GLENELE	3121 0000	31 0000	5 5837	31 2900	101
TR	1	LARGE SEED	1171 0000	31 8448	2 6401	8 9703	37
TR	2	SMALL SEED	1004 0000	33 4867	5 9484	35 3854	30
TR	3	MIXED SEED	956 0000	28 1176	6 4458	41 5483	34
CUL	3	PARK	4883 0000	47 8429	8 8834	47 3814	88
TR	1	LARGE SEED	1888 0000	48 8874	5 8488	34 2100	38
TR	2	SMALL SEED	1382 0000	48 4888	7 7882	60 8098	30
TR	3	MIXED SEED	1417 0000	47 2333	6 8880	43 5471	30
CUL	4	70M008002	1814 0000	28 8000	5 8345	48 7182	80
TR	1	LARGE SEED	788 0000	30 7800	5 3484	28 8088	28
TR	2	SMALL SEED	314 0000	20 8333	6 0847	36 7810	16
TR	3	MIXED SEED	600 0000	28 3421	5 7780	33 3626	19
TOTAL CASES =			487				

APPENDIX 16

02/16/82

PAGE 3

FILE SBL3 (CREATION DATE = 12/30/81)

DESCRIPTION OF SUBPOPULATIONS							
CRITERION VARIABLE	BY	BY	TILLERS PER PLANT				
BROKEN DOWN BY	CUL	TR	CULTIVAR				
			TREATMENT				
VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
FOR ENTIRE POPULATION			2387 0000	5 1704	2 1002	4 4107	(487)
CUL	0		488 0000	5 3182	1 8452	3 4080	(82)
TR	1	LARGE SEED	208 0000	5 8844	1 8177	3 3040	(38)
TR	2	SMALL SEED	188 0000	5 2813	1 7841	3 1718	(32)
TR	3	MIXED SEED	118 0000	4 7817	1 8232	3 7273	(24)
CUL	1	PITIC 82	810 0000	5 7847	2 8070	5 7884	(108)
TR	1	LARGE SEED	201 0000	5 8909	2 8184	5 5227	(33)
TR	2	SMALL SEED	184 0000	4 8877	2 8092	5 2888	(31)
TR	3	MIXED SEED	268 0000	5 0714	2 3416	5 4824	(42)
CUL	2	GLENLEA	398 0000	3 8408	1 2478	1 5584	(101)
TR	1	LARGE SEED	128 0000	3 8488	1 7884	0 8221	(27)
TR	2	SMALL SEED	128 0000	4 8223	1 1388	1 2820	(30)
TR	3	MIXED SEED	127 0000	3 7383	1 8820	2 4430	(34)
CUL	3	PARK	824 0000	5 3488	1 7840	3 2188	(88)
TR	1	LARGE SEED	213 0000	5 8082	1 4248	2 0202	(38)
TR	2	SMALL SEED	184 0000	5 4887	2 1813	4 8713	(30)
TR	3	MIXED SEED	147 0000	4 9000	1 7878	3 1888	(30)
CUL	4	7000000002	348 0000	5 7887	2 2727	5 1880	(80)
TR	1	LARGE SEED	188 0000	5 8788	2 3482	5 8138	(28)
TR	2	SMALL SEED	81 0000	5 4000	2 8873	7 1142	(18)
TR	3	MIXED SEED	107 0000	5 8218	1 8822	3 4878	(18)
TOTAL CASES =			487				

APPENDIX 16

02/16/82

PAGE 4

FILE SBL3 (CREATION DATE = 12/30/81)

DESCRIPTION OF SUBPOPULATIONS							
CRITERION VARIABLE	BY	BY	DRY WEIGHT PER PLANT				
BROKEN DOWN BY	CUL	TR	CULTIVAR				
			TREATMENT				
VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
FOR ENTIRE POPULATION			1881 8488	4 8741	1 8768	2 4821	(487)
CUL	0		370 8800	4 0281	1 8888	3 4818	(82)
TR	1	LARGE SEED	182 4700	4 2382	1 8781	2 4808	(38)
TR	2	SMALL SEED	118 8300	3 7188	1 2028	1 4488	(32)
TR	3	MIXED SEED	88 2800	4 1387	1 8482	3 7088	(24)
CUL	1	PITIC 82	428 8800	4 0882	1 8784	2 8288	(108)
TR	1	LARGE SEED	127 8800	4 1884	1 8788	2 8218	(33)
TR	2	SMALL SEED	128 8800	4 0887	1 7704	2 1342	(31)
TR	3	MIXED SEED	188 8200	3 8488	1 8488	2 7088	(42)
CUL	2	GLENLEA	407 7400	4 0370	1 4447	2 0871	(101)
TR	1	LARGE SEED	142 2700	3 8481	1 0871	1 1778	(27)
TR	2	SMALL SEED	142 1300	4 7377	1 8214	2 3148	(30)
TR	3	MIXED SEED	123 2400	3 8278	1 8828	2 4111	(34)
CUL	3	PARK	383 3400	4 0137	1 4128	1 9883	(88)
TR	1	LARGE SEED	170 8700	4 4882	1 3277	1 7827	(38)
TR	2	SMALL SEED	111 7000	3 7223	1 4728	2 1888	(30)
TR	3	MIXED SEED	110 8700	3 8888	1 3842	1 7828	(30)
CUL	4	7000000002	280 1300	4 2388	1 8848	3 4744	(80)
TR	1	LARGE SEED	121 0300	5 0288	1 8888	3 4873	(28)
TR	2	SMALL SEED	82 8400	3 8882	1 8887	3 4478	(18)
TR	3	MIXED SEED	78 2800	3 8811	1 8882	2 8771	(18)
TOTAL CASES =			487				

POOR COPY
COPIE DE QUALITEE INFERIEURE

250

Appendix 16. Descriptive statistics of characters of plants from large,
small and mixed size seeds, measured at maturity, in 1979.

POOR COPY
COPIE DE QUALITEE INFERIEURE

251

APPENDIX 10

03/16/83

PAGE 2

FILE SBL4 CREATION DATE = 12/30/81
SUBFILE P401 P402 P403 P404 P405 P406 P407 P408 P409 P410 P411 P412
P413 P414 P415 P416 P417 P418 P419 P420 P421 P422 P423 P424
P425 P426 P427 P428 P429 P430 P431 P432 P433 P434 P435 P436
P437 P438 P439 P440 P441 P442 P443 P444 P445 P446 P447 P448
P449 P450 P451 P452 P453 P454 P455 P456 P457 P458 P459 P460
P461 P462 P463 P464 P465 P466 P467 P468 P469 P470 P471 P472
P473 P474 P475 P476 P477 P478 P479 P480 P481 P482 P483 P484
P485 P486 P487 P488 P489 P490 P491 P492 P493 P494 P495 P496
P497 P498 P499 P500

CRITERION VARIABLE Y1 CUL YR
BROKEN DOWN BY CUL YR
DESCRIPTION OF SUB POPULATIONS
TILLERS PER PLANT
CULTIVAR
TREATMENT

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
FOR ENTIRE POPULATION			2210 0000	4 8775	2 0842	4 3507	444
CUL	1	PITIC SEED	757 0000	5 4480	2 0857	4 3503	139
TR	1	SMALL SEED	271 0000	5 1591	2 0903	4 3695	44
TR	2	LARGE SEED	238 0000	5 4884	2 4041	5 7788	43
TR	3	MIXED SEED	250 0000	4 8077	1 8723	2 4721	52
CUL	2	GLENEA	408 0000	3 7850	1 4144	2 0005	107
TR	1	SMALL SEED	128 0000	3 7850	1 4041	1 9715	34
TR	2	LARGE SEED	154 0000	3 8500	1 4587	2 1304	40
TR	3	MIXED SEED	126 0000	3 7879	1 4088	1 9846	33
CUL	3	PARK	850 0000	5 8585	2 1312	4 5421	117
TR	1	SMALL SEED	289 0000	5 8474	2 3587	5 5541	48
TR	2	LARGE SEED	135 0000	5 0000	1 7541	3 0789	27
TR	3	MIXED SEED	246 0000	5 8503	2 0722	4 2939	64
CUL	4	YONGGONG002	384 0000	4 8135	2 1617	4 6289	81
TR	1	SMALL SEED	129 0000	5 1500	2 1547	4 6400	30
TR	2	LARGE SEED	76 0000	4 2889	1 7187	2 9575	18
TR	3	MIXED SEED	180 0000	5 0000	2 3365	5 4595	38
TOTAL CASES =	444						

APPENDIX 10

03/16/83

PAGE 4

FILE	SOLA	(CREATION DATE = 12/30/81)	P401	P402	P403	P404	P405	P406	P407	P408	P409	P410	P411	P412
SUBFILE	P413	P414	P415	P416	P417	P418	P419	P420	P421	P422	P423	P424	P425	P426
	P427	P428	P429	P430	P431	P432	P433	P434	P435	P436	P437	P438	P439	P440
	P441	P442	P443	P444	P445	P446	P447	P448	P449	P450	P451	P452	P453	P454
	P455	P456	P457	P458	P459	P460	P461	P462	P463	P464	P465	P466	P467	P468
	P469	P470	P471	P472	P473	P474	P475	P476	P477	P478	P479	P480	P481	P482
	P483	P484	P485	P486	P487	P488	P489	P490	P491	P492	P493	P494	P495	P496
	P497	P498	P499	P500	P501	P502	P503	P504	P505	P506	P507	P508	P509	P510
	P511	P512	P513	P514	P515	P516	P517	P518	P519	P520	P521	P522	P523	P524
	P525	P526	P527	P528	P529	P530	P531	P532	P533	P534	P535	P536	P537	P538
	P539	P540	P541	P542	P543	P544	P545	P546	P547	P548	P549	P550	P551	P552
	P553	P554	P555	P556	P557	P558	P559	P560	P561	P562	P563	P564	P565	P566
	P567	P568	P569	P570	P571	P572	P573	P574	P575	P576	P577	P578	P579	P580
	P581	P582	P583	P584	P585	P586	P587	P588	P589	P590	P591	P592	P593	P594
	P595	P596	P597	P598	P599	P600	P601	P602	P603	P604	P605	P606	P607	P608
	P609	P610	P611	P612	P613	P614	P615	P616	P617	P618	P619	P620	P621	P622
	P623	P624	P625	P626	P627	P628	P629	P630	P631	P632	P633	P634	P635	P636
	P637	P638	P639	P640	P641	P642	P643	P644	P645	P646	P647	P648	P649	P650
	P651	P652	P653	P654	P655	P656	P657	P658	P659	P660	P661	P662	P663	P664
	P665	P666	P667	P668	P669	P670	P671	P672	P673	P674	P675	P676	P677	P678
	P679	P680	P681	P682	P683	P684	P685	P686	P687	P688	P689	P690	P691	P692
	P693	P694	P695	P696	P697	P698	P699	P700	P701	P702	P703	P704	P705	P706
	P707	P708	P709	P710	P711	P712	P713	P714	P715	P716	P717	P718	P719	P720
	P721	P722	P723	P724	P725	P726	P727	P728	P729	P730	P731	P732	P733	P734
	P735	P736	P737	P738	P739	P740	P741	P742	P743	P744	P745	P746	P747	P748
	P749	P750	P751	P752	P753	P754	P755	P756	P757	P758	P759	P760	P761	P762
	P763	P764	P765	P766	P767	P768	P769	P770	P771	P772	P773	P774	P775	P776
	P777	P778	P779	P780	P781	P782	P783	P784	P785	P786	P787	P788	P789	P790
	P791	P792	P793	P794	P795	P796	P797	P798	P799	P800	P801	P802	P803	P804
	P805	P806	P807	P808	P809	P810	P811	P812	P813	P814	P815	P816	P817	P818
	P819	P820	P821	P822	P823	P824	P825	P826	P827	P828	P829	P830	P831	P832
	P833	P834	P835	P836	P837	P838	P839	P840	P841	P842	P843	P844	P845	P846
	P847	P848	P849	P850	P851	P852	P853	P854	P855	P856	P857	P858	P859	P860
	P861	P862	P863	P864	P865	P866	P867	P868	P869	P870	P871	P872	P873	P874
	P875	P876	P877	P878	P879	P880	P881	P882	P883	P884	P885	P886	P887	P888
	P889	P890	P891	P892	P893	P894	P895	P896	P897	P898	P899	P900	P901	P902
	P903	P904	P905	P906	P907	P908	P909	P910	P911	P912	P913	P914	P915	P916
	P917	P918	P919	P920	P921	P922	P923	P924	P925	P926	P927	P928	P929	P930
	P931	P932	P933	P934	P935	P936	P937	P938	P939	P940	P941	P942	P943	P944
	P945	P946	P947	P948	P949	P950	P951	P952	P953	P954	P955	P956	P957	P958
	P959	P960	P961	P962	P963	P964	P965	P966	P967	P968	P969	P970	P971	P972
	P973	P974	P975	P976	P977	P978	P979	P980	P981	P982	P983	P984	P985	P986
	P987	P988	P989	P990	P991	P992	P993	P994	P995	P996	P997	P998	P999	P1000
	P1001	P1002	P1003	P1004	P1005	P1006	P1007	P1008	P1009	P1010	P1011	P1012	P1013	P1014
	P1015	P1016	P1017	P1018	P1019	P1020	P1021	P1022	P1023	P1024	P1025	P1026	P1027	P1028
	P1029	P1030	P1031	P1032	P1033	P1034	P1035	P1036	P1037	P1038	P1039	P1040	P1041	P1042
	P1043	P1044	P1045	P1046	P1047	P1048	P1049	P1050	P1051	P1052	P1053	P1054	P1055	P1056
	P1057	P1058	P1059	P1060	P1061	P1062	P1063	P1064	P1065	P1066	P1067	P1068	P1069	P1070
	P1071	P1072	P1073	P1074	P1075	P1076	P1077	P1078	P1079	P1080	P1081	P1082	P1083	P1084
	P1085	P1086	P1087	P1088	P1089	P1090	P1091	P1092	P1093	P1094	P1095	P1096	P1097	P1098
	P1099	P1100	P1101	P1102	P1103	P1104	P1105	P1106	P1107	P1108	P1109	P1110	P1111	P1112
	P1113	P1114	P1115	P1116	P1117	P1118	P1119	P1120	P1121	P1122	P1123	P1124	P1125	P1126
	P1127	P1128	P1129	P1130	P1131	P1132	P1133	P1134	P1135	P1136	P1137	P1138	P1139	P1140
	P1141	P1142	P1143	P1144	P1145	P1146	P1147	P1148	P1149	P1150	P1151	P1152	P1153	P1154
	P1155	P1156	P1157	P1158	P1159	P1160	P1161	P1162	P1163	P1164	P1165	P1166	P1167	P1168
	P1169	P1170	P1171	P1172	P1173	P1174	P1175	P1176	P1177	P1178	P1179	P1180	P1181	P1182
	P1183	P1184	P1185	P1186	P1187	P1188	P1189	P1190	P1191	P1192	P1193	P1194	P1195	P1196
	P1197	P1198	P1199	P1200	P1201	P1202	P1203	P1204	P1205	P1206	P1207	P1208	P1209	P1210
	P1211	P1212	P1213	P1214	P1215	P1216	P1217	P1218	P1219	P1220	P1221	P1222	P1223	P1224
	P1225	P1226	P1227	P1228	P1229	P1230	P1231	P1232	P1233	P1234	P1235	P1236	P1237	P1238
	P1239	P1240	P1241	P1242	P1243	P1244	P1245	P1246	P1247	P1248	P1249	P1250	P1251	P1252
	P1253	P1254	P1255	P1256	P1257	P1258	P1259	P1260	P1261	P1262	P1263	P1264	P1265	P1266
	P1267	P1268	P1269	P1270	P1271	P1272	P1273	P1274	P1275	P1276	P1277	P1278	P1279	P1280
	P1281	P1282	P1283	P1284	P1285	P1286	P1287	P1288	P1289	P1290	P1291	P1292	P1293	P1294
	P1295	P1296	P1297	P1298	P1299	P1300	P1301	P1302	P1303	P1304	P1305	P1306	P1307	P1308
	P1309	P1310	P1311	P1312	P1313	P1314	P1315	P1316	P1317	P1318	P1319	P1320	P1321	P1322
	P1323	P1324	P1325	P1326	P1327	P1328	P1329	P1330	P1331	P1332	P1333	P1334	P1335	P1336
	P1337	P1338	P1339	P1340	P1341	P1342	P1343	P1344	P1345	P1346	P1347	P1348	P1349	P1350
	P1351	P1352	P1353	P1354	P1355	P1356	P1357	P1358	P1359	P1360	P1361	P1362	P1363	P1364
	P1365	P1366	P1367	P1368	P1369	P1370	P1371	P1372	P1373	P1374	P1375	P1376	P1377	P1378
	P1379	P1380	P1381	P1382	P1383	P1384	P1385	P1386	P1387	P1388	P1389	P1390	P1391	P1392
	P1393	P1394	P1395	P1396	P1397	P1398	P1399	P1400	P1401	P1402	P1403	P1404	P1405	P1406
	P1407	P1408	P1409	P1410	P1411	P1412	P1413	P1414	P1415	P1416	P1417	P1418	P1419	P1420
	P1421	P1422	P1423	P1424	P1425	P1426	P1427	P1428	P1429	P1430	P1431	P1432	P1433	P1434
	P1435	P1436	P1437	P1438	P1439	P1440	P1441	P1442	P1443	P1444	P1445	P1446	P1447	P1448
	P1449	P1450	P1451	P1452	P1453	P1454	P1455	P1456	P1457	P1458	P1459	P1460	P1461	P1462
	P1463	P1464	P1465	P1466	P1467	P1468	P1469	P1470	P1471	P1472	P1473	P1474	P1475	P1476
	P1477	P1478	P1479	P1480	P1481	P1482	P1483	P1484	P1485	P1486	P1487	P1488	P1489	P1490
	P1491	P1492	P1493	P1494	P1495	P1496	P1497	P1498	P1499	P1500	P1501	P1502	P1503	P1504
	P1505	P1506	P1507	P1508	P1509	P1510	P1511	P1512	P1513	P1514	P1515	P1516	P1517	P1518
	P1519	P1520	P1521	P1522	P1523	P1524	P1525	P1526	P1527	P1528	P1529	P1530	P1531	P1532
	P1533	P1534	P1535	P1536	P1537	P1538	P1539	P1540	P1541	P1542	P1543	P1544	P1545	P1546
	P1547	P1548	P1549	P1550	P1551	P1552	P1553	P1554	P1555	P1556	P1557	P1558	P1559	P1560
	P1561	P1562	P1563	P1564	P1565	P1566	P1567	P1568	P1569	P1570	P1571	P1572	P1573	P1574
	P1575	P1576	P1577	P1578	P1579	P1580	P1581	P1582	P1583	P1584	P1585	P1586	P1587	P1588
	P1589	P1590	P1591	P1592	P1593	P1594	P1595	P1596	P1597	P1598	P1599	P1600	P1601	P1602
	P1603	P1604	P1605	P1606	P1607	P1608	P1609	P1610	P1611	P1612	P1613	P1614	P1615	P1616
	P1617	P1618	P1619	P1620	P1621	P1622	P1623	P1624	P1625	P1626	P1627	P1628	P1629	P1630
	P1631	P1632	P1633	P1634	P1635	P1636	P1637	P1638	P1639	P1640	P1641	P1642	P1643	P1644
	P1645	P1646	P1647	P1648	P1649	P1650	P1651	P1652	P1653	P1654	P1655	P1656	P1657	P1658
	P1659	P1660	P1661	P1662	P1663	P1664	P1665	P1666	P1667	P1668	P1669	P1670	P1671	P1672
	P1673	P1674	P1675	P1676	P1677	P1678	P1679	P1680	P1681	P1682	P1683	P1684	P1685	P1686
	P1687	P1688	P1689	P1690	P1691	P1692	P1693	P1694	P1695	P1696	P1697	P1698	P1699	P1700
	P1701	P1702	P1703	P1704	P1705	P1706	P1707							

03/10/02

PAGE 8

SUSPENSE	P401	P402	P403	P404	P405	P406	P407	P408	P409	P410	P411	P412
P414	P415	P416	P417	P418	P419	P420	P421	P422	P423	P424	P425	P426
P511	P512	P513	P514	P515	P516	P517	P518	P519	P520	P521	P522	P523
P600	P610	P611	P612	P613	P614	P615	P616	P617	P618	P619	P620	P621

CRITERION VARIABLE	DESCRIPTION OF SUBPOPULATIONS
BROKEN DOWN BY	Y1 SEED YIELD PER PLANT
	CUL CULTIVAR
BY	TR TREATMENT

VARIABLE	CASE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
FOR ENTIRE POPULATION			3317 0300	7 5067	3 5063	12 2023	(443)
CUL	1	PITIC SEED	1304 3000	0 3043	3 7873	14 1020	(130)
TR	1	SMALL SEED	403 3000	10 0070	3 4031	11 0000	(40)
TR	2	LARGE SEED	300 7000	0 2000	4 0003	16 0070	(43)
TR	3	MIXED SEED	426 3000	0 3713	3 1223	0 6110	(52)
CUL	2	ELERLEA	023 4000	7 7430	3 1332	0 8171	(107)
TR	1	SMALL SEED	253 0000	7 4002	3 0070	0 0474	(30)
TR	2	LARGE SEED	322 0000	0 0072	3 3004	11 4270	(40)
TR	3	MIXED SEED	201 0100	7 0330	3 0002	0 0371	(23)
CUL	3	PARK	020 0000	0 0000	2 3073	0 0003	(117)
TR	1	SMALL SEED	201 7700	0 4723	3 7720	7 0003	(40)
TR	2	LARGE SEED	130 4300	0 1307	3 4003	0 7000	(27)
TR	3	MIXED SEED	248 4700	0 0470	1 0033	2 4710	(44)
CUL	4	YOMOOO0002	040 4300	0 0100	3 1174	0 7100	(70)
TR	1	SMALL SEED	103 3300	0 0332	3 0007	0 0000	(20)
TR	2	LARGE SEED	114 0500	0 3000	1 0030	3 0100	(10)
TR	3	MIXED SEED	200 0300	7 0017	2 0000	14 7011	(20)
TOTAL CASES =			444				
MISSING CASES =			2 OR	0 0 PCT			

03/10/82

PAGE 7

SUSPENSE	P401	P402	P403	P404	P405	P406	P407	P408	P409	P410	P411	P412
	P414	P415	P401	P402	P403	P404	P405	P406	P407	P408	P409	P410
	P411	P412	P414	P415	P401	P402	P403	P404	P405	P406	P407	P408
	P409	P410	P411	P412	P413	P414	P415	P416	P417	P418	P419	P420

CRITERION VARIABLE		DESCRIPTION OF SUBPOPULATIONS	
BROKEN DOWN BY	FI	HEIGHT OF THE FLAG LEAF BLADE CM	
BY	CUL	CULTIVAR	
	TR	TREATMENT	

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
FOR ENTIRE POPULATION			27735 0000	62 4862	10 2334	100 7326	444
CUL	1	PITIC SEED	8784 0000	63 2882	7 7877	60 1823	139
TR	1	SMALL SEED	2722 0000	64 1364	5 5187	32 5888	44
TR	2	LARGE SEED	2877 0000	61 0930	6 8741	47 1500	43
TR	3	MIXED SEED	3345 0000	64 3285	6 9433	48 3224	42
CUL	2	GLNLEA	7816 0000	73 8813	8 8384	78 7722	107
TR	1	SMALL SEED	2801 0000	73 8888	5 1476	26 4884	34
TR	2	LARGE SEED	3020 0000	76 8000	7 4827	55 8223	40
TR	3	MIXED SEED	2395 0000	72 6768	7 3810	54 8889	33
CUL	3	PARE	6832 0000	68 2032	6 4888	42 2002	109
TR	1	SMALL SEED	2718 0000	69 1087	6 8105	46 8880	40
TR	2	LARGE SEED	1888 0000	67 2288	6 7148	45 0883	27
TR	3	MIXED SEED	2287 0000	68 1138	6 7280	46 2668	44
CUL	4	7000000002	4193 0000	61 7864	6 8887	31 3888	81
TR	1	SMALL SEED	1273 0000	60 8200	4 8638	20 8267	28
TR	2	LARGE SEED	902 0000	60 1111	4 7288	22 4878	18
TR	3	MIXED SEED	2018 0000	63 1083	6 3483	40 3129	28

APPENDIX 18		03/18/82										PAGE 8	
FILE	SOL#	CREATION DATE = 12/30/81											
SUBFILE	P001	P002	P003	P004	P005	P006	P007	P008	P009	P010	P011	P012	
	P010	P015	P001	P002	P003	P004	P005	P006	P007	P008	P009	P010	
	P011	P012	P010	P015	P001	P002	P003	P004	P005	P006	P007	P008	
	P009	P010	P011	P012	P013								
CRITERION VARIABLE													
BROKEN DOWN BY													
BY													
CUL													
TREATMENT													
SUB POPULATIONS													
VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N						
FOR ENTIRE POPULATION			43266 0000	97 4610	12 7777	162 2700	(444)						
CUL	1	PITTS 82	12002 0000	84 1842	7 8628	62 2446	(120)						
TR	1	SMALL SEED	4167 0000	84 4773	6 8728	75 2320	(44)						
TR	2	LARGE SEED	3876 0000	82 4410	8 8088	74 6630	(43)						
TR	3	MIXED SEED	4959 0000	86 4036	6 6237	42 8662	(82)						
CUL	2	GLENSDA	11088 0000	111 8316	6 7851	76 6095	(107)						
TR	1	SMALL SEED	3806 0000	111 8412	6 5119	32 0870	(34)						
TR	2	LARGE SEED	4504 0000	112 8000	11 1336	123 8265	(40)						
TR	3	MIXED SEED	2686 0000	110 7879	6 2483	37 5946	(33)						
CUL	3	PARR	11620 0000	98 8281	7 1091	50 8296	(117)						
TR	1	SMALL SEED	4878 0000	98 4783	7 1701	51 4106	(40)						
TR	2	LARGE SEED	3702 0000	100 0741	7 1572	51 3251	(27)						
TR	3	MIXED SEED	4440 0000	100 0465	7 1657	51 3067	(40)						
CUL	4	TOMM000002	8547 0000	80 8372	8 0684	37 0448	(81)						
TR	1	SMALL SEED	2998 0000	80 3300	4 8664	22 0600	(26)						
TR	2	LARGE SEED	1418 0000	78 8697	4 8683	20 1176	(16)						
TR	3	MIXED SEED	3133 0000	82 1842	7 2265	52 2064	(38)						
TOTAL CASES =			444										

APPENDIX 18			03/10/82										PAGE 8	
FILE	SBL4	(CREATION DATE = 12/30/81)												
SUBFILE	P401	P402	P403	P404	P405	P406	P407	P408	P409	P410	P411	P412		
	P410	P411	P412	P413	P414	P415	P416	P417	P418	P419	P420	P421		
	P421	P422	P423	P424	P425	P426	P427	P428	P429	P430	P431	P432		
	P433	P434	P435	P436	P437	P438	P439	P440	P441	P442	P443	P444		
CRITERION VARIABLE														
BROKEN DOWN BY														
BY														
DESCRIPTION OF SUSPENSION														
CULTIVAR														
TREATMENT														
PER HEAD														
VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N							
FOR ENTIRE POPULATION			3787 0000	8 4242	1 3728	1 8846	4461							
CUL	1	PIYIC SE	1207 0000	8 8836	1 2742	1 6237	1207							
TR	1	SMALL SEED	382 0000	8 5818	0 8680	0 7537	44							
TR	2	LARGE SEED	358 0000	8 4884	1 8018	2 2564	43							
TR	3	MIXED SEED	466 0000	8 5482	1 3044	1 7014	52							
CUL	2	GLEHLEA	820 0000	8 8881	1 4628	2 1108	1071							
TR	1	SMALL SEED	287 0000	8 7283	1 8261	2 6248	34							
TR	2	LARGE SEED	346 0000	8 5800	1 1447	1 3103	40							
TR	3	MIXED SEED	277 0000	8 3838	1 8180	2 6812	29							
CUL	3	PARK	942 0000	8 0512	1 4072	1 8501	1171							
TR	1	SMALL SEED	383 0000	7 8012	1 4020	1 8657	44							
TR	2	LARGE SEED	216 0000	8 0741	1 8171	2 3920	27							
TR	3	MIXED SEED	343 0000	8 2048	1 1118	1 2363	44							
CUL	4	TOMMOOSES	888 0000	8 6172	1 2606	1 6622	811							
TR	1	SMALL SEED	211 0000	8 4400	1 8287	2 3400	25							
TR	2	LARGE SEED	181 0000	8 8444	1 2113	1 4873	18							
TR	3	MIXED SEED	326 0000	8 5788	1 0813	1 1803	38							
TOTAL CASES =			446											

APPENDIX 10

02/18/82

PAGE 10

FILE SBL4 (CREATION DATE = 12/30/81)

SUSFILE P401 P402 P403 P404 P405 P406 P407 P408 P409 P410 P411 P412
P413 P414 P415 P416 P417 P418 P419 P420 P421 P422 P423 P424
P425 P426 P427 P428 P429 P430 P431 P432 P433 P434 P435 P436

CRITERION VARIABLE HL1 HEADLENGTH CM
BROKEN DOWN BY CUL CULTIVAR
BY TR TREATMENT

DESCRIPTION OF SUB POPULATIONS

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
FOR ENTIRE POPULATION			5132 0000	11 8721	2 6173	6 8803	(444)
CUL	1	PITIC 82	1847 0000	14 0072	2 0126	4 0507	(128)
TR	1	SMALL SEED	810 0000	13 8838	1 9982	3 9919	(44)
TR	2	LARGE SEED	895 0000	14 0888	2 1646	4 6855	(43)
TR	3	MIXED SEED	732 0000	14 0788	1 9288	3 7198	(82)
CUL	2	SLERLEA	1278 0000	11 9188	2 3818	5 7183	(107)
TR	1	SMALL SEED	422 0000	12 4412	2 0837	7 0418	(34)
TR	2	LARGE SEED	478 0000	11 8788	2 2218	4 9327	(40)
TR	3	MIXED SEED	377 0000	11 4242	2 2843	5 1288	(32)
CUL	3	PARK	1106 0000	9 4781	1 2888	1 6788	(117)
TR	1	SMALL SEED	418 0000	9 1827	1 1888	1 4224	(48)
TR	2	LARGE SEED	384 0000	9 7778	1 3888	1 9288	(27)
TR	3	MIXED SEED	425 0000	9 6881	1 2888	1 6788	(44)
CUL	4	700000002	888 0000	9 8783	1 2847	1 6744	(81)
TR	1	SMALL SEED	243 0000	9 7288	1 8144	2 2833	(28)
TR	2	LARGE SEED	178 0000	9 8444	1 0888	1 1844	(18)
TR	3	MIXED SEED	368 0000	10 1878	1 1514	1 2887	(38)
TOTAL CASES =	444						

APPENDIX 10

02/18/82

PAGE 11

FILE SBL4 (CREATION DATE = 12/30/81)

SUSFILE P401 P402 P403 P404 P405 P406 P407 P408 P409 P410 P411 P412
P413 P414 P415 P416 P417 P418 P419 P420 P421 P422 P423 P424
P425 P426 P427 P428 P429 P430 P431 P432 P433 P434 P435 P436

CRITERION VARIABLE HL1 HEADLENGTH CM
BROKEN DOWN BY CUL CULTIVAR
BY TR TREATMENT

DESCRIPTION OF SUB POPULATIONS

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
FOR ENTIRE POPULATION			80080 0000	181 8420	87 0410	7878 1287	(441)
CUL	1	PITIC 82	32885 0000	238 8827	101 1188	10224 3888	(138)
TR	1	SMALL SEED	12100 0000	278 0000	84 8888	9022 4372	(44)
TR	2	LARGE SEED	18888 0000	234 8372	114 4284	13088 1388	(43)
TR	3	MIXED SEED	19887 0000	205 8182	82 7474	7013 8271	(82)
CUL	2	SLERLEA	18812 0000	181 0888	88 3188	3837 4883	(106)
TR	1	SMALL SEED	4878 0000	148 4118	58 8720	3420 6738	(34)
TR	2	LARGE SEED	8278 0000	188 7888	88 1823	4248 8888	(40)
TR	3	MIXED SEED	4764 0000	148 4788	87 8387	3282 8000	(32)
CUL	3	PARK	17872 0000	181 0427	83 0884	3878 2818	(117)
TR	1	SMALL SEED	7883 0000	183 8783	78 4878	8883 7881	(48)
TR	2	LARGE SEED	3818 0000	141 2883	84 8878	4212 8888	(27)
TR	3	MIXED SEED	6774 0000	183 8848	48 2877	2148 8888	(44)
CUL	4	700000002	13881 0000	178 7723	78 2887	8884 8218	(78)
TR	1	SMALL SEED	4882 0000	182 8888	71 8827	8182 8188	(28)
TR	2	LARGE SEED	2817 0000	188 8888	48 8848	2488 8788	(18)
TR	3	MIXED SEED	6882 0000	182 8323	87 1888	7881 4888	(38)
TOTAL CASES =	444						
MISSING CASES =	2 88	0.7 PCT					

APPENDIX 10

03/16/82

PAGE 13

FILE	SUBFILE	P401	P402	P403	P404	P405	P406	P407	P408	P409	P410	P411	P412
		P414	P415	P416	P417	P418	P419	P420	P421	P422	P423	P424	P425
		P426	P427	P428	P429	P430	P431	P432	P433	P434	P435	P436	P437
		P438	P439	P440	P441	P442	P443	P444	P445	P446	P447	P448	P449
		P450	P451	P452	P453	P454	P455	P456	P457	P458	P459	P460	P461
		P462	P463	P464	P465	P466	P467	P468	P469	P470	P471	P472	P473
		P474	P475	P476	P477	P478	P479	P480	P481	P482	P483	P484	P485
		P486	P487	P488	P489	P490	P491	P492	P493	P494	P495	P496	P497
		P498	P499	P500	P501	P502	P503	P504	P505	P506	P507	P508	P509
		P510	P511	P512	P513	P514	P515	P516	P517	P518	P519	P520	P521
		P522	P523	P524	P525	P526	P527	P528	P529	P530	P531	P532	P533
		P534	P535	P536	P537	P538	P539	P540	P541	P542	P543	P544	P545
		P546	P547	P548	P549	P550	P551	P552	P553	P554	P555	P556	P557
		P558	P559	P560	P561	P562	P563	P564	P565	P566	P567	P568	P569
		P570	P571	P572	P573	P574	P575	P576	P577	P578	P579	P580	P581
		P582	P583	P584	P585	P586	P587	P588	P589	P590	P591	P592	P593
		P594	P595	P596	P597	P598	P599	P600	P601	P602	P603	P604	P605
		P606	P607	P608	P609	P610	P611	P612	P613	P614	P615	P616	P617
		P618	P619	P620	P621	P622	P623	P624	P625	P626	P627	P628	P629
		P630	P631	P632	P633	P634	P635	P636	P637	P638	P639	P640	P641
		P642	P643	P644	P645	P646	P647	P648	P649	P650	P651	P652	P653
		P654	P655	P656	P657	P658	P659	P660	P661	P662	P663	P664	P665
		P666	P667	P668	P669	P670	P671	P672	P673	P674	P675	P676	P677
		P678	P679	P680	P681	P682	P683	P684	P685	P686	P687	P688	P689
		P690	P691	P692	P693	P694	P695	P696	P697	P698	P699	P700	P701
		P702	P703	P704	P705	P706	P707	P708	P709	P710	P711	P712	P713
		P714	P715	P716	P717	P718	P719	P720	P721	P722	P723	P724	P725
		P726	P727	P728	P729	P730	P731	P732	P733	P734	P735	P736	P737
		P738	P739	P740	P741	P742	P743	P744	P745	P746	P747	P748	P749
		P750	P751	P752	P753	P754	P755	P756	P757	P758	P759	P760	P761
		P762	P763	P764	P765	P766	P767	P768	P769	P770	P771	P772	P773
		P774	P775	P776	P777	P778	P779	P780	P781	P782	P783	P784	P785
		P786	P787	P788	P789	P790	P791	P792	P793	P794	P795	P796	P797
		P798	P799	P800	P801	P802	P803	P804	P805	P806	P807	P808	P809
		P810	P811	P812	P813	P814	P815	P816	P817	P818	P819	P820	P821
		P822	P823	P824	P825	P826	P827	P828	P829	P830	P831	P832	P833
		P834	P835	P836	P837	P838	P839	P840	P841	P842	P843	P844	P845
		P846	P847	P848	P849	P850	P851	P852	P853	P854	P855	P856	P857
		P858	P859	P860	P861	P862	P863	P864	P865	P866	P867	P868	P869
		P870	P871	P872	P873	P874	P875	P876	P877	P878	P879	P880	P881
		P882	P883	P884	P885	P886	P887	P888	P889	P890	P891	P892	P893
		P894	P895	P896	P897	P898	P899	P900	P901	P902	P903	P904	P905
		P906	P907	P908	P909	P910	P911	P912	P913	P914	P915	P916	P917
		P918	P919	P920	P921	P922	P923	P924	P925	P926	P927	P928	P929
		P930	P931	P932	P933	P934	P935	P936	P937	P938	P939	P940	P941
		P942	P943	P944	P945	P946	P947	P948	P949	P950	P951	P952	P953
		P954	P955	P956	P957	P958	P959	P960	P961	P962	P963	P964	P965
		P966	P967	P968	P969	P970	P971	P972	P973	P974	P975	P976	P977
		P978	P979	P980	P981	P982	P983	P984	P985	P986	P987	P988	P989
		P990	P991	P992	P993	P994	P995	P996	P997	P998	P999	P1000	P1001
		P1002	P1003	P1004	P1005	P1006	P1007	P1008	P1009	P1010	P1011	P1012	P1013
		P1014	P1015	P1016	P1017	P1018	P1019	P1020	P1021	P1022	P1023	P1024	P1025
		P1026	P1027	P1028	P1029	P1030	P1031	P1032	P1033	P1034	P1035	P1036	P1037
		P1038	P1039	P1040	P1041	P1042	P1043	P1044	P1045	P1046	P1047	P1048	P1049
		P1050	P1051	P1052	P1053	P1054	P1055	P1056	P1057	P1058	P1059	P1060	P1061
		P1062	P1063	P1064	P1065	P1066	P1067	P1068	P1069	P1070	P1071	P1072	P1073
		P1074	P1075	P1076	P1077	P1078	P1079	P1080	P1081	P1082	P1083	P1084	P1085
		P1086	P1087	P1088	P1089	P1090	P1091	P1092	P1093	P1094	P1095	P1096	P1097
		P1098	P1099	P1100	P1101	P1102	P1103	P1104	P1105	P1106	P1107	P1108	P1109
		P1110	P1111	P1112	P1113	P1114	P1115	P1116	P1117	P1118	P1119	P1120	P1121
		P1122	P1123	P1124	P1125	P1126	P1127	P1128	P1129	P1130	P1131	P1132	P1133
		P1134	P1135	P1136	P1137	P1138	P1139	P1140	P1141	P1142	P1143	P1144	P1145
		P1146	P1147	P1148	P1149	P1150	P1151	P1152	P1153	P1154	P1155	P1156	P1157
		P1158	P1159	P1160	P1161	P1162	P1163	P1164	P1165	P1166	P1167	P1168	P1169
		P1170	P1171	P1172	P1173	P1174	P1175	P1176	P1177	P1178	P1179	P1180	P1181
		P1182	P1183	P1184	P1185	P1186	P1187	P1188	P1189	P1190	P1191	P1192	P1193
		P1194	P1195	P1196	P1197	P1198	P1199	P1200	P1201	P1202	P1203	P1204	P1205
		P1206	P1207	P1208	P1209	P1210	P1211	P1212	P1213	P1214	P1215	P1216	P1217
		P1218	P1219	P1220	P1221	P1222	P1223	P1224	P1225	P1226	P1227	P1228	P1229
		P1230	P1231	P1232	P1233	P1234	P1235	P1236	P1237	P1238	P1239	P1240	P1241
		P1242	P1243	P1244	P1245	P1246	P1247	P1248	P1249	P1250	P1251	P1252	P1253
		P1254	P1255	P1256	P1257	P1258	P1259	P1260	P1261	P1262	P1263	P1264	P1265
		P1266	P1267	P1268	P1269	P1270	P1271	P1272	P1273	P1274	P1275	P1276	P1277
		P1278	P1279	P1280	P1281	P1282	P1283	P1284	P1285	P1286	P1287	P1288	P1289
		P1290	P1291	P1292	P1293	P1294	P1295	P1296	P1297	P1298	P1299	P1300	P1301
		P1302	P1303	P1304	P1305	P1306	P1307	P1308	P1309	P1310	P1311	P1312	P1313
		P1314	P1315	P1316	P1317	P1318	P1319	P1320	P1321	P1322	P1323	P1324	P1325
		P1326	P1327	P1328	P1329	P1330	P1331	P1332	P1333	P1334	P1335	P1336	P1337
		P1338	P1339	P1340	P1341	P1342	P1343	P1344	P1345	P1346	P1347	P1348	P1349
		P1350	P1351	P1352	P1353	P1354	P1355	P1356	P1357	P1358	P1359	P1360	P1361
		P1362	P1363	P1364	P1365	P1366	P1367	P1368	P1369	P1370	P1371	P1372	P1373
		P1374	P1375	P1376	P1377	P1378	P1379	P1380	P1381	P1382	P1383	P1384	P1385
		P1386	P1387	P1388	P1389	P1390	P1391	P1392	P1393	P1394	P1395	P1396	P1397
		P1398	P1399	P1400	P1401	P1402	P1403	P1404	P1405	P1406	P1407	P1408	P1409
		P1410	P1411	P1412	P1413	P1414	P1415	P1416	P1417	P1418	P1419	P1420	P1421
		P1422	P1423	P1424	P1425	P1426	P1427	P1428	P1429	P1430	P1431	P1432	P1433
		P1434	P1435	P1436	P1437	P1438	P1439	P1440	P1441	P1442	P1443	P1444	P1445
		P1446	P1447	P1448	P1449	P1450	P1451	P1452	P1453	P1454	P1455	P1456	P1457
		P1458	P1459	P1460	P1461	P1462	P1463	P1464	P1465	P1466	P1467	P1468	P1469
		P1470	P1471	P1472	P1473	P1474	P1475	P1476	P1477	P1478	P1479	P1480	P1481
		P1482	P1483	P1484	P1485	P1486	P1487	P1488	P1489	P1490	P1491	P1492	P1493
		P1494	P1495	P1496	P1497	P1498	P1499	P1500	P1501	P1502	P1503	P1504	P1505
		P1506	P1507	P1508	P1509	P1510	P1511	P1512	P1513	P1514	P1515	P1516	P1517
		P1518	P1519	P1520	P1521	P1522	P1523	P1524	P1525	P1526	P1527	P1528	P1529
		P1530	P1531	P1532	P1533	P1534	P1535	P1536	P1537	P1538	P1539	P1540	P1541
		P1542	P1543	P1544	P1545	P1546	P1547	P1548	P1549	P1550	P1551	P1552	P1553
		P1554	P1555	P1556	P1557	P1558	P1559	P1560	P1561	P1562	P1563	P1564	P1565
		P1566	P1567	P1568	P1569	P1570	P1571	P1572	P1573	P1574	P1575	P1576	P1577
		P1578	P1579	P1580	P1581	P1582	P1583	P1584	P1585	P1586	P1587	P1588	P1589
		P1590	P1591	P1592	P1593	P1594	P1595	P1596	P1597	P1598	P1599	P1600	P1601
		P1602	P1603	P1604	P1605	P1606	P1607	P1608	P1609	P1610	P1611	P1612	P1613
		P1614	P1615	P1616	P1617	P1618	P1619	P1620	P1621	P1622	P1623	P1624	P1625
		P1626	P1627	P1628	P1629	P1630	P1631	P1632	P1633	P1634	P1635	P1636	P1637
		P1638	P1639	P1640	P1641	P1642	P1643	P1644	P1645	P1646	P1647	P1648	P1649
		P1650	P1651	P16									

PAGE 14

MSBFILE	P401	P402	P403	P404	P405	P406	P407	P408	P409	P410	P411	P412
	P415	P501	P502	P503	P504	P505	P506	P507	P508	P509	P510	P511
	P511	P512	P514	P515	P501	P502	P504	P505	P507	P508	P509	P510
	P505	P510	P511	P512	P513	P515						

CRITERION VARIABLE		DESCRIPTION OF SUSPOPULATION
BROKEN DOWN BY	EN1	NUMBER OF KERNELS PER HEAD
BY	CUL	CULTIVAR
	TR	TREATMENT

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
FOR ENTIRE POPULATION			17384 6882	40 2083	8 0875	82 7648	(431)
CUL	1	PITIC SEED	8214 8001	48 8002	8 8245	74 8887	(128)
TR	1	SMALL SEED	2090 5812	47 5124	8 4025	41 0055	(44)
TR	2	LARGE SEED	1705 2556	43 7882	10 5903	111 8200	(41)
TR	3	MIXED SEED	2324 7322	45 8614	8 4250	70 2288	(51)
CUL	2	BLUENLA	4288 7070	41 8279	7 1147	50 8194	(102)
TR	1	SMALL SEED	1307 8886	42 3825	7 8809	82 8840	(32)
TR	2	LARGE SEED	1808 4288	41 2316	8 5986	43 8647	(30)
TR	3	MIXED SEED	1282 7023	41 3775	8 9279	48 1263	(31)
CUL	3	PARK	3782 8521	32 8874	5 7050	44 8880	(118)
TR	1	SMALL SEED	1424 4880	38 8873	5 8156	43 7882	(48)
TR	2	LARGE SEED	878 8301	33 8812	8 4828	71 8577	(38)
TR	3	MIXED SEED	1478 7272	34 3880	5 0457	28 4668	(42)
CUL	4	TOMMOOOOOO	3089 1144	28 8888	7 8109	62 8610	(77)
TR	1	SMALL SEED	861 8915	38 4877	7 8108	51 0081	(28)
TR	2	LARGE SEED	727 4208	40 4124	8 2489	58 7184	(18)
TR	3	MIXED SEED	1270 9910	40 5880	7 8825	61 8810	(34)
TOTAL CASES =			446				
MISSING CASES =			13 OR	2 9 PCT			

PAGE 18

SUBFILE	P401	P402	P403	P404	P405	P406	P408	P410	P411	P412
	P416	P415	P501	P502	P504	P504	P507	P507	P510	
	P511	P512	P516	P515	P601	P603	P604	P606	P607	P608
	P609	P610	P611	P612	P613	P615				

DESCRIPTION OF SUBPOPULATIONS		
CRITERION VARIABLE	SLI	EXTRUSION LENGTH CM
BROKEN DOWN BY	CUL	CULTIVAR
BY	TR	TREATMENT

VARIABLE	CODE	VALUE LABEL	SUM	MEAN	STD DEV	VARIANCE	N
FOR ENTIRE POPULATION			10413 0000	23 4927	7 3226	53 7086	444
CUL	1	PITIC SEED	2352 0000	16 9209	4 4544	19 8415	139
TR	1	SMALL SEED	735 0000	16 4773	5 2226	28 7304	46
TR	2	LARGE SEED	743 0000	17 2791	3 9661	15 7298	43
TR	3	MIXED SEED	884 0000	17 0060	2 7838	7 8038	52
CUL	2	GLENNLEA	2778 0000	25 9368	4 5831	21 0051	107
TR	1	SMALL SEED	862 0000	25 9412	4 2895	18 4813	50
TR	2	LARGE SEED	1008 0000	25 2350	4 2448	18 4661	60
TR	3	MIXED SEED	884 0000	25 7879	4 3246	18 7346	33
CUL	3	PARK	3740 0000	31 9554	3 3606	11 4299	117
TR	1	SMALL SEED	1428 0000	31 2609	3 2279	10 4193	48
TR	2	LARGE SEED	862 0000	33 9887	6 2807	16 1534	27
TR	3	MIXED SEED	1428 0000	33 2727	2 8314	8 0186	44
CUL	4	7000000002	1845 0000	18 0884	3 3226	11 1648	81
TR	1	SMALL SEED	482 0000	18 0890	3 2265	10 7923	28
TR	2	LARGE SEED	338 0000	18 5111	3 5170	12 3893	18
TR	3	MIXED SEED	719 0000	18 8211	3 7155	13 8064	26

TOTAL CASES - 444

Appendix 17. Distribution parameters of characters of single plants, measured at different stages of growth, derived from large seeds, small seeds and alternated large and small seeds.

All plants were grown in three row plots in 1979. Plants with missing neighbours are excluded.

Character	Skewness ^{1,2}			Kurtosis ²		
	Large seed	Small seed	Mixed seed	Large seed	Small seed	Mixed seed
3-5 Leaf Stage:						
Ht ₁	-0.78*	-0.85*	-0.38	1.59*	0.92	0.85
L	-0.80*	-0.34	-0.59	2.60**	1.62*	2.18**
T	-0.59	0.03	-0.59	0.91	-0.01	0.81
Wt	0.29	0.07	0.06	1.24	0.39	0.53
Jointing Stage:						
Ht ₂	0.11	0.45	0.12	1.04	-0.21	-0.35
M	-0.00	-0.19	-0.21	0.98	0.58	-0.61
T	0.07	-0.44	-0.10	0.36	0.31	0.02
Wt	0.42	0.18	0.30	1.42*	-0.29	0.08
Heading Stage:						
Ht ₃	-0.23	-0.39	-0.39	1.01	1.29	1.19
T	-0.57	-0.54	-0.10	0.55	0.32	-0.03
Wt	0.19	0.22	0.15	0.05	-0.59	0.55
Maturity:						
Ht ₃	-0.37	-1.22**	-0.49	0.76	2.45**	0.53
T	0.60	0.14	0.39	0.75	0.61	-0.10
Wt	0.33	0.01	0.41	1.02	0.85	0.38
M	0.27	0.18	0.34	0.57	0.64	0.24
FL	0.13	-0.47	-0.12	1.31	0.49	0.24
ExL	0.16	-0.27	-0.06	0.39	3.06**	0.36
HL	0.06	-1.06*	-0.59	1.31	2.91**	0.94
Sp/H	-0.56	-0.59	-0.50	0.85	0.64	0.10
Y/P	0.34	-0.09	0.35	1.15	0.77	0.31
K/P	0.41	-0.06	0.35	1.58*	0.78	0.22
HI	-0.80*	-1.28**	-0.15	1.95**	2.23**	2.38**
Krt	-0.50	-0.14	-0.51	0.57*	-0.25	0.82
K/H	-0.33	-0.75*	-0.23	1.84**	0.76	1.03

1. Values of skewness and kurtosis are averages of the values of three replicates and four genotypes.

2. * significant, $\alpha \leq 0.05$; ** significant, $\alpha \leq 0.01$.

Appendix 18a. Plot yield and Specific Mixture Efficiency¹ of mixtures grown in 1977.

Mixture	Plot yield (g)		Specific Mixture Efficiency	
	Mean	Std. Dev.	Mean	Std. Dev.
For Entire Population	1345	138	1.05	0.10
Glenlea	1249	154		
Glenlea + Park	1278	64	1.10	0.09
Glenlea + Neepawa	1361	179	1.12	0.06
Glenlea + 70M110001	1277	80	0.98	0.05
Glenlea + 70M009002	1346	27	0.98	0.07
Glenlea + Norquay	1352	150	1.02	0.16
Glenlea + NB 701	1353	110	1.09	0.05
Park	1096	105		
Park + Neepawa	1227	130	1.08	0.06
Park + 70M110001	1275	97	1.05	0.16
Park + 70M009002	1401	56	1.07	0.01
Park + Norquay	1364	8	1.04	0.03
Park + NB 701	1347	145	1.16	0.13
Neepawa	1170	86		
Neepawa + 70M110001	1302	153	1.03	0.14
Neepawa + 70M009002	1344	65	1.00	0.01
Neepawa + Norquay	1239	103	0.96	0.04
Neepawa + NB 701	1302	126	1.08	0.03
70M110001	1358	253		
70M110001 + 70M009002	1498	123	1.05	0.06
70M110001 + Norquay	1482	92	1.07	0.03
70M110001 + NB 701	1303	70	1.03	0.19
70M009002	1514	64		
70M009002 + Norquay	1451	114	0.99	0.06
70M009002 + NB 701	1508	88	1.10	0.05
Norquay	1418	108		
Norquay + NB 701	1421	36	1.07	0.07
NB 701	1240	203		

1. See Section 8.2.

Appendix 18b. Plot yield and Specific Mixture Efficiency of mixtures grown in 1978.

Mixture	Plot yield (g)		Specific Mixture Efficiency	
	Mean	Std. Dev.	Mean	Std. Dev.
For Entire Population	1012	138	1.02	0.07
Pitic 62	966	251		
Pitic 62 + Glenlea	1111	116	1.11	0.07
Pitic 62 + Park	1014	47	1.08	0.11
Pitic 62 + Neepawa	1112	61	1.08	0.13
Pitic 62 + 70M110001	1105	92	1.09	0.08
Pitic 62 + 70M009002	1006	47	1.11	0.05
Pitic 62 + Norquay	912	79	0.92	0.13
Pitic 62 + NB 701	993	229	1.04	0.03
Glenlea	1043	138		
Glenlea + Park	1085	62	1.10	0.09
Glenlea + Neepawa	1087	115	1.02	0.20
Glenlea + 70M110001	1197	27	1.13	0.02
Glenlea + 70M009002	1067	48	1.13	0.10
Glenlea + Norquay	1063	57	1.02	0.02
Glenlea + NB 701	912	160	0.92	0.10
Park	933	57		
Park + Neepawa	1023	77	1.00	0.06
Park + 70M110001	911	40	0.91	0.00
Park + 70M009002	815	84	0.91	0.09
Park + Norquay	992	118	1.00	0.07
Park + NB 701	880	186	0.93	0.10
Neepawa	1112	92		
Neepawa + 70M110001	1007	125	0.92	0.10
Neepawa + 70M009002	1000	92	1.02	0.13
Neepawa + Norquay	1141	103	1.06	0.12
Neepawa + NB 701	1127	105	1.10	0.11
70M110001	1078	113		
70M110001 + 70M009002	873	124	0.92	0.11
70M110001 + Norquay	1054	66	0.99	0.04
70M110001 + NB 701	1050	168	1.03	0.08
70M009002	859	159		
70M009002 + Norquay	873	2	0.92	0.05
70M009002 + NB 701	894	175	0.99	0.16
Norquay	1049	38		
Norquay + NB 701	1106	31	1.12	0.08
NB 701	941	179		

Appendix 18c. Plot yield and Specific Mixture Efficiency of mixtures
grown in 1979.

Mixture	Plot yield (g)		Specific Mixture Efficiency	
	Mean	Std. Dev.	Mean	Std. Dev.
For Entire Population	1248	140	1.03	
Pitic 62	1342	80		
Pitic 62 + Glenlea	1436	36	1.09	0.10
Pitic 62 + Park	1313	11	1.11	0.06
Pitic 62 + Neepawa	1339	59	1.10	0.05
Pitic 62 + 70M110001	1379	29	1.05	0.09
Pitic 62 + 70M009002	1511	80	1.14	0.02
Pitic 62 + Norquay	1437	99	1.19	0.13
Pitic 62 + NB 701	1401	123	1.05	0.12
Glenlea	1312	166		
Glenlea + Park	1238	99	1.06	0.03
Glenlea + Neepawa	1199	75	1.00	0.13
Glenlea + 70M110001	1175	81	0.91	0.11
Glenlea + 70M009002	1313	112	1.01	0.14
Glenlea + Norquay	1271	98	1.08	0.20
Glenlea + NB 701	1272	63	0.97	0.08
Park	1033	104		
Park + Neepawa	1002	38	0.94	0.04
Park + 70M110001	1153	34	1.00	0.04
Park + 70M009002	1212	40	1.04	0.01
Park + Norquay	1082	73	1.02	0.03
Park + NB 701	1324	134	1.12	0.08
Neepawa	1102	125		
Neepawa + 70M110001	1122	44	0.94	0.05
Neepawa + 70M009002	1178	36	0.98	0.04
Neepawa + Norquay	1067	58	0.98	0.08
Neepawa + NB 701	1181	30	0.97	0.02
70M110001	1290	130		
70M110001 + 70M009002	1285	67	1.00	0.08
70M110001 + Norquay	1141	22	0.96	0.07
70M110001 + NB 701	1175	167	0.90	0.14
70M009002	1299	43		
70M009002 + Norquay	1301	62	1.10	0.09
70M009002 + NB 701	1307	45	1.00	0.02
Norquay	1082	122		
Norquay + NB 701	1226	80	1.01	0.07
NB 701	1327	73		

Appendix 19. Performance of mixtures in each of the three years, containing specific genotypes, as expressed by the Array mean yield (Array \bar{Y}), the Average Mixture Efficiency (AME)^{1,2}, and the General Combining Ability (GCA)².

Year	Genotype	Array \bar{Y}	Std. Dev.	AME	GCA
1977	Pitic 62	--	--	--	--
	Glenlea	1317	118	1.04**	-28.0
	Park	1280	134	1.07**	-71.8*
	Neepawa	1278	134	1.04**	-69.4*
	70M110001	1357	152	1.02*	17.3
	70M009002	1438	104	1.03*	100.9*
	Norquay	1391	121	1.03*	51.0*
	NB 701	1354	138	1.07**	-0.0
1978	Pitic 62	1028	142	1.05**	7.6
	Glenlea	1071	119	1.05**	54.3
	Park	957	121	0.99	-52.2
	Neepawa	1076	106	1.02*	65.6*
	70M110001	1035	138	1.00	24.5
	70M009002	924	128	1.00	-86.1*
	Norquay	1024	110	1.01	13.0
	NB 701	988	177	1.02*	-26.6
1979	Pitic 62	1395	92	1.09**	129.4*
	Glenlea	1277	116	1.01	32.1
	Park	1170	137	1.04**	-81.8*
	Neepawa	1149	112	0.99	-91.5*
	70M110001	1215	117	0.97**	-19.6
	70M009002	1301	112	1.03**	49.8*
	Norquay	1201	145	1.04**	-51.8*
	NB 701	1277	119	1.00	33.4

1. See Section 8.2.

2. * significant, $\alpha \leq 0.05$; ** significant, $\alpha \leq 0.01$

Appendix 20.

Calculation of the effect of skewness on single plant selection.

Assume a 50:50 mixture of two genotypes G1 and G2. The mean value for a character X of G1 is 0.3589 or less than the mean of this character for G2. (The value of 0.3589 had to be chosen due to the restricted number of values listed in chi-square tables). Assume further that X has a chi-square distribution with 30 degrees of freedom, and thus a variance of 2×30 , a skewness of 0.516, and a kurtosis of 0.400. If the 0.75% of the population having the highest values for X were selected, the selected fractions would consist of $1/3$ G1 and $2/3$ G2 (Figure 5). The probability of selecting G1 rather than G2 is $P(\text{Error})=0.333$.

If the character, X, had had a standard normal distribution, it can be found, through trial and error, that:

$$P(z > 2.29) = 0.011$$


$$P(z > 2.29 + 0.3589) = P(z > 2.6489) = 0.00401$$

$$\text{Total selected fraction} = (0.011 + 0.00401) / 2 = 0.0075$$

The probability to select G1 rather than G2 is

$$P(\text{Error}) = 0.00401 / 0.01501 = 0.267.$$

The probability to select G1 rather than G2 is thus $0.333 / 0.267 = 1.25$ times larger in the case where the distribution of X is skewed, than in the case where the



d.f. = 30

$\sigma^2 = 2 \times 30 = 60$

Skewness = 0.516

Kurtosis = 0.400

$$\mu_{G_1} - \mu_{G_2} = \frac{53.672 - 50.892}{\sqrt{60}} = 0.3589$$

Fig. 5. Graphic illustration of selection in a 50:50 mixture of two genotypes, G_1 and G_2 , for a character x , which has a positively skewed distribution. The selected fraction would, in this case, consist of $1/3 G_1$ and $2/3 G_2$.

distribution of X is normal.

P(Error) will increase, however when the frequency of G2 in the population decreases.

Assume a mixture of G1 and G2, with means for the character X which are 1.096 σ apart, and in which the frequency of G2 is 1/10 the frequency of G1. The distribution of X is the same as in the previous example. If this time 0.11% of the population is selected, then the probability of selecting G1 rather than G2 is

$$P(\text{Error}) = \frac{10(0.0005)}{1.0 + 10(0.0005)} = \frac{0.005}{0.01} = 0.5$$

If X had had a standard normal distribution, then it can be found, by trial and error, that

$$P(Z > 2.33) = 0.01$$

$$P(Z > 2.33 + 1.096) = P(Z > 3.426) = 0.0002$$

$$P(\text{Error}) = 10(0.0002)/(1 \times 0.02 + 0.0002) = 0.1$$

The probability to select G1 is thus 5/2=2.5 times larger in the case where X is skewed than in the case of a normal distribution of X.

On the other hand, if the distribution of X is negatively skewed, and has a curve which is the mirror image of the curve used in the above examples, the probability to select the superior genotype, G2, is greater than would be the case if X were normally distributed.

For a 50:50 mixture of G1 and G2, with the means 0.151 s apart, it can be calculated, in the same manner as before, that the probability to select G2 is 1.19 times greater than would be the case for a normal distribution, if 0.75 % of the population is selected. If the ratio of G1:G2 is 10:1, and the means are 0.3851 s apart, the probability to select G2 rather than G1 is 1.49 times greater than would be the case if X were normally distributed and 0.11% were selected.