

INFORMATION TO USERS

This manuscript has been reproduced from the microfilm master. UMI films the text directly from the original or copy submitted. Thus, some thesis and dissertation copies are in typewriter face, while others may be from any type of computer printer.

The quality of this reproduction is dependent upon the quality of the copy submitted. Broken or indistinct print, colored or poor quality illustrations and photographs, print bleedthrough, substandard margins, and improper alignment can adversely affect reproduction.

In the unlikely event that the author did not send UMI a complete manuscript and there are missing pages, these will be noted. Also, if unauthorized copyright material had to be removed, a note will indicate the deletion.

Oversize materials (e.g., maps, drawings, charts) are reproduced by sectioning the original, beginning at the upper left-hand corner and continuing from left to right in equal sections with small overlaps.

Photographs included in the original manuscript have been reproduced xerographically in this copy. Higher quality 6" x 9" black and white photographic prints are available for any photographs or illustrations appearing in this copy for an additional charge. Contact UMI directly to order.

**ProQuest Information and Learning
300 North Zeeb Road, Ann Arbor, MI 48106-1346 USA
800-521-0600**

UMI[®]

UNIVERSITY OF ALBERTA

**POSSIBLE ROLES FOR THE VACUOLAR H⁺-ATPASE AND
F₁F₀-ATPASE IN ALUMINUM RESISTANCE**

BY

CHRISTIE ANNE HAMILTON



A THESIS

SUBMITTED TO THE FACULTY OF GRADUATE STUDIES AND RESEARCH

IN PARTIAL FULFILLMENT OF THE REQUIREMENTS FOR

THE DEGREE OF DOCTOR OF PHILOSOPHY

IN

MOLECULAR BIOLOGY AND GENETICS

DEPARTMENT OF BIOLOGICAL SCIENCES

EDMONTON, ALBERTA

SPRING 2002



**National Library
of Canada**

**Acquisitions and
Bibliographic Services**

**395 Wellington Street
Ottawa ON K1A 0N4
Canada**

**Bibliothèque nationale
du Canada**

**Acquisitions et
services bibliographiques**

**395, rue Wellington
Ottawa ON K1A 0N4
Canada**

Your file Votre référence

Our file Notre référence

The author has granted a non-exclusive licence allowing the National Library of Canada to reproduce, loan, distribute or sell copies of this thesis in microform, paper or electronic formats.

The author retains ownership of the copyright in this thesis. Neither the thesis nor substantial extracts from it may be printed or otherwise reproduced without the author's permission.

L'auteur a accordé une licence non exclusive permettant à la Bibliothèque nationale du Canada de reproduire, prêter, distribuer ou vendre des copies de cette thèse sous la forme de microfiche/film, de reproduction sur papier ou sur format électronique.

L'auteur conserve la propriété du droit d'auteur qui protège cette thèse. Ni la thèse ni des extraits substantiels de celle-ci ne doivent être imprimés ou autrement reproduits sans son autorisation.

0-612-68580-2

Canada

**UNIVERSITY OF ALBERTA
LIBRARY RELEASE FORM**

Name of Author: Christie Anne Hamilton
Title of Thesis: Possible Roles for the Vacuolar H⁺-ATPase and
Mitochondrial F₁F₀-ATPase in Aluminum Resistance
Degree: Doctor of Philosophy
Year this Degree Granted: 2002

Permission is hereby granted to the University of Alberta Library to reproduce single copies of this thesis and to lend or sell such copies for private, scholarly or scientific research purposes only.

The author reserves other publication rights, and neither the thesis nor extensive extracts from it may be printed or otherwise reproduced without the author's written permission.


.....Christie Anne Hamilton.....

Christie Anne Hamilton
105 9th St NW
Calgary AB T2N 1T1

Date: January 22, 2002

FACULTY OF GRADUATE STUDIES AND RESEARCH


The undersigned certify that they have read, and recommend to the Faculty of Graduate Studies and Research for acceptance, a thesis entitled **POSSIBLE ROLES FOR THE VACUOLAR H⁺-ATPASE AND F₁F₀-ATPASE IN ALUMINUM RESISTANCE** submitted by **CHRISTIE ANNE HAMILTON** in partial fulfillment of the requirements for the degree of **DOCTOR OF PHILOSOPHY** in Molecular Biology and Genetics.


.....
Gregory Taylor, Co-Supervisor


.....
Allen Good, Co-Supervisor


.....
Peter Constabel


.....
Bernard Lemire


.....
Jocelyn Ozga


.....
Donald R. Ort, External Examiner

Date: Jan 17, 2002

ABSTRACT

Crop yields are significantly reduced under acidic soil conditions, where the speciation of aluminum (Al) compounds changes so that Al becomes soluble and/or toxic. Recent research has focused on developing crop cultivars resistant to soluble Al in the soil. Previous research in our laboratory identified two 51 kD, Al-induced proteins from an Al-resistant cultivar of wheat (*Triticum aestivum* L.).

This thesis describes the purification and identification of the 51 kD, aluminum-induced proteins from wheat. Peptide microsequence analysis revealed that the purified peptides were homologous to subunits of the vacuolar H⁺-ATPase (V-ATPase) and the mitochondrial ATP synthase (F₁F₀-ATPase). Both V-ATPase and F₁F₀-ATPase activities were induced by Al in the Al-resistant cultivar, PT741, and not in other cultivars tested. Based on these results, I hypothesized that increased V-ATPase activity could contribute to Al resistance by counteracting cytoplasmic acidosis caused by Al-inhibition of the plasma membrane ATPase. Vacuolar ATPase activity could maintain cytoplasmic pH homeostasis by transporting protons into the vacuole. Alternatively, it may be required to allow tonoplast transmembrane transport of other molecules, such as organic acids. The ATP required to fuel this process could be provided by the mitochondrial ATP synthase.

This hypothesis was tested in the yeast and *Arabidopsis* model systems. Yeast V-ATPase mutant strains were hypersensitive to Al, while F₁F₀-ATPase mutant strains exhibited wild-type growth. Since the tonoplast membrane potential generated by V-ATPase activity is also thought to be required for NaCl tolerance in yeast, the scope of my research was expanded to include analysis of the yeast response to NaCl.

As was observed with Al, V-ATPase mutant strains were hypersensitive to NaCl while F_1F_0 -ATPase mutant strains exhibited wild-type growth. Transgenic *Arabidopsis* expressing a cDNA encoding the B subunit of the V-ATPase in the antisense orientation were also hypersensitive to Al and, to a lesser extent, NaCl.

In conclusion, this research describes the development and testing of a novel hypothesis that describes potential roles for the vacuolar H^+ -ATPase and the F_1F_0 -ATPase in Al and NaCl tolerance. Determination of the precise mechanism by which increased V-ATPase activity mediates Al and NaCl tolerance awaits future investigation.

ACKNOWLEDGEMENTS

I would like to thank my co-supervisors, Dr Gregory Taylor and Dr Allen Good, for their patience and encouragement during the course of this work. With their support I felt free to pursue my own path. I would also like to acknowledge the valuable advice of my committee members, Dr Peter Constabel, Dr Jocelyn Ozga and Dr Bernard Lemire. My external examiner, Dr Donald Ort, started my defense by stating that he would never let a Ph.D. student feel let down by the ease of his or her defense. I would like to thank him for maintaining his record. Financial support for this work was provided by the Natural Sciences and Engineering Research Council of Canada (research grants to Dr Gregory Taylor and Dr Allen Good), the Canadian Wheat Board Graduate Fellowship Program and by the University of Alberta Department of Biological Sciences.

This achievement would not have been possible without the encouragement and unconditional support of my parents. I would also like to thank my lab colleagues - in particular Ms Mary dePauw, Dr John Vidmar, Ms Julie Stephens, Dr Urmila Basu and Dr Jan Slaski - for many helpful and enjoyable discussions over the years, and for their friendship. I have also enjoyed the support of my friends Tanya Martin, Dr Quynh Tran, Jonathan Burke, Leah Hrab, BJ Daniels, Jay White and Bill Klimke. Finally, I would like to thank my fiance, Dr Richard Fekete. I look forward to a future of learning and love during our long-term collaboration.

TABLE OF CONTENTS

CHAPTER	PAGE
1. Introduction	1
1.1 Agricultural Significance of Aluminum	1
1.2 Effects of Aluminum on Plant Cellular Metabolism	1
1.3 The Search for a Primary Toxic Lesion	4
1.4 Genetic Variation in Aluminum Tolerance	5
1.5 Mechanisms of Aluminum Resistance	7
1.5.1 Al-induced Release of Organic Anions	7
1.5.2 Plant-Induced Elevation of Rhizosphere pH	8
1.5.3 Exudation of Al-binding Polypeptides	9
1.5.4 Support for External Resistance from Inhibitor Studies	10
1.5.5 Modulation of Intracellular Enzyme Activity	10
1.5.6 Sequestration of Al in the Vacuole	11
1.6 Structure of the Vacuolar H ⁺ -ATPase	14
1.7 Function of the Vacuolar H ⁺ -ATPase	21
1.8 Control of ATPase Expression	21
1.9 Objectives	24
1.10 Literature cited	28
2. Induction of Vacuolar ATPase and Mitochondrial ATP Synthase by Aluminum in an Aluminum-Resistant Cultivar of <i>Triticum aestivum</i> L.	44

2.1	Introduction	44
2.2	Methods	47
2.3	Results	53
	2.3.1 Purification of RMP51	53
	2.3.2 Peptide Microsequence Analysis	56
	2.3.3 V-ATPase and F ₁ F ₀ -ATPase are Al-induced	64
	2.3.4 Cultivar Screen for ATPase Induction	67
2.4	Discussion	73
2.5	Literature cited	79
3.	Vacuolar H⁺-ATPase, but not Mitochondrial ATP Synthase, is Required for Aluminum Resistance in <i>Saccharomyces cerevisiae</i>	88
3.1	Introduction	88
3.2	Methods	92
3.3	Results	97
	3.3.1 Response of yeast ATPases to aluminum	97
	3.3.2 V-ATPase mutants, but not F ₁ F ₀ -ATPase mutants, are hypersensitive to Al	102
3.4	Discussion	103
3.5	Literature cited	108

4.	Vacuolar H⁺-ATPase, but not Mitochondrial ATP Synthase, is Required for NaCl Tolerance in <i>Saccharomyces cerevisiae</i>	113
4.1	Introduction	113
4.2	Methods	115
4.3	Results	118
4.3.1	Response of yeast ATPases to NaCl	118
4.3.2	V-ATPase mutants, but not F ₁ F ₀ -ATPase mutants, are hypersensitive to NaCl	124
4.4	Discussion	130
4.5	Literature cited	133
5.	Involvement of the Vacuolar H⁺-ATPase in Al and NaCl Tolerance in <i>Arabidopsis thaliana</i>	137
5.1	Introduction	137
5.2	Methods	140
5.3	Results	143
5.3.1	The Effect of Al on Root Elongation in Wild-type <i>Arabidopsis</i>	143
5.3.2	The Effect of NaCl on Root Elongation in Wild-type	

<i>Arabidopsis</i>	144
5.3.3 Production of Transgenic <i>Arabidopsis</i> Expressing the V-ATPase B Subunit in the Antisense Orientation	149
5.3.4 Phenotype Testing of Transgenic <i>Arabidopsis</i>	157
5.4 Discussion	169
5.5 Literature cited	173
6. Concluding Discussion	178
6.1 Literature cited	185

LIST OF TABLES

Table	Title	Page
1-1	Classification of Al-induced genes	13
1-2	Subunit composition of the vacuolar ATPase (V-ATPase) and functional homologies to subunits of the mitochondrial ATP synthase (F ₁ F ₀ -ATPase)	16
2-1	Enrichment of marker enzymes in endomembrane-enriched membrane fractions isolated from root tips of <i>Triticum aestivum</i> L. cv PT741	55
3-1	Yeast strains used in this study	94
5-1	Determination of copy number in transgenic <i>Arabidopsis thaliana</i> as measured by segregation of hygromycin resistance in T ₂ progeny	156
5-2	Three factor analysis of variance of root elongation measurements of transgenic and parent (C24) lines in the presence of dexamethasone and/or (A) Al (B) NaCl.	168

LIST OF FIGURES

Figure	Title	Page
1-1	Global distribution of acidic soils	3
1-2	Structural comparison of the vacuolar ATPase (V-ATPase) and mitochondrial ATP synthase (F_1F_0 -ATPase)	18
2-1	Fractionation of endomembrane-associated proteins by continuous elution electrophoresis	58
2-2	Two-dimensional gel analysis of crude endomembrane proteins and purified RMP51	60
2-3	Identification of RMP51 by immunoblotting	63
2-4	The effect of Al on activity of the V-ATPase, F_1F_0 -ATPase and P-ATPase	66
2-5	The effect of Al on transcript levels of V-ATPase and F_1F_0 -ATPase Subunits	69
2-6	The effect of Al on ATPase specific activity in total microsomal membranes prepared from 1 cm root tips of PT741, Atlas-66 and Maringa (Al-resistant), and from Scout-66 and Katepwa (Al-sensitive)	71
3-1	Model representing the energy balance hypothesis for V-ATPase and F_1F_0 -ATPase-mediated Al resistance in wheat	91
3-2	The effect of Al on cellular ATPase activities (F_1F_0 -ATPase,	

	V-ATPase, P-ATPase) in wild type yeast	99
3-3	The effect of Al on expression of V-ATPase and F ₁ F ₀ -ATPase	101
3-4	The effect of ATPase mutations on Al resistance in yeast	105
4-1	The effect of NaCl on cellular ATPase activities (F ₁ F ₀ -ATPase, P-ATPase, V-ATPase) in wild type yeast	121
4-2	The effect of NaCl on expression of F ₁ F ₀ -ATPase and V-ATPase	123
4-3	The effect of ATPase mutations on NaCl tolerance in yeast	126
4-4	Halotolerance of V-ATPase and F ₁ F ₀ -ATPase mutant strains	129
5-1	The effect of Al on root elongation in wild-type <i>Arabidopsis</i> (ecotype C24)	146
5-2	The effect of NaCl on root elongation in wild-type <i>Arabidopsis</i> (ecotype C24)	148
5-3	The T-DNA construct designed for glucocorticoid-inducible expression of the B subunit of the vacuolar ATPase (At57) in the antisense orientation	151
5-4	Analysis of transgenic lines of <i>Arabidopsis</i> by PCR amplification of a 650 bp fragment of hygromycin phosphotransferase and a 598 bp fragment specific to the transgene	153
5-5	The effect of transgene expression on Al resistance in <i>Arabidopsis</i> , measured by root elongation in the presence and absence of 30 μM dexamethasone (DEX)	160
5-6	The effect of transgene expression on NaCl resistance in <i>Arabidopsis</i> , measured by root elongation in the presence and	

	absence of 30 μ M dexamethasone (DEX)	162
5-7	The effect of dexamethasone application on Al and NaCl resistance in wild-type <i>Arabidopsis thaliana</i> (ecotype C24) in the presence and absence of 30 μ M dexamethasone (DEX)	164
5-8	The effect of transgene expression on Al and NaCl resistance in <i>Arabidopsis thaliana</i> (ecotype C24, transgenic line 34) in the presence and absence of 30 μ M dexamethasone (DEX)	166

1. Introduction


1.1 Agricultural Significance of Aluminum

Crop yields are significantly reduced under acidic soil conditions, where the speciation of aluminum (Al) naturally present in soil changes such that Al becomes soluble and toxic. Acidic soils comprise approximately 40% of the world's arable land, and this area (Fig 1-1) is likely expanding due to acidic precipitation, use of nitrogenous fertilizers and poor land management practices in newly cleared tropical forests (von Euxkull and Mutert, 1995). The poor fertility of acidic soils is aggravated by physical factors such as low water-holding capacity, susceptibility to crusting, erosion and compaction (von Euxkul! and Mutert, 1995). Historically, Al toxicity has been ameliorated by liming soil. This increases soil pH so that Al becomes insoluble and/or nontoxic. Since this method is expensive and fails to correct subsoil acidity, recent research has focused on developing crop cultivars resistant to soluble Al in soil.

1.2 Effects of Aluminum on Plant Cellular Metabolism

Aluminum disturbs cellular metabolism and inhibits growth of Al-sensitive plants. Decreased growth correlates with inhibition of cell division (Morimura *et al.*, 1978) and elongation (Wallace and Anderson, 1984; Taylor and Foy, 1985), reduced mineral uptake by roots (Baligar *et al.*, 1987), decreased activity of calmodulin-

Figure 1-1. Global distribution of acidic soils (indicated in red). For the purpose of this figure, topsoils of pH 5.5 or less are considered significantly acidic. (Adapted from van Wambeke, 1976; Wright *et al.* (eds), 1990)



regulated enzymes (Putterill and Gardner, 1988; Siegel and Haug, 1983, Suhayda and Haug, 1985; Slaski, 1989), reduced DNA synthesis (Matsumoto and Morimura, 1980; Wallace and Anderson, 1984) and disruption of the Golgi apparatus in root cap cells (Bennet *et al.*, 1987).

1.3 The Search for a Primary Toxic Lesion

Recent research has attempted to identify the primary toxic lesion caused by Al. However, the multitude of symptoms exhibited by plants grown in the presence of Al has made it difficult to determine a single, primary cause for growth inhibition. One mechanism that might account for the range of symptoms observed is interaction of Al with calcium (Taylor, 1988; Rengel, 1992; Kochian, 1995). Aluminum interferes with transport of calcium (Ca^{2+}) across the plasma membrane by blocking Ca^{2+} channels (Ding *et al.*, 1993; Pineros and Tester, 1995). However, this is unlikely to be directly responsible for Al toxicity, since root growth can be inhibited at low Al activities without affecting Ca^{2+} uptake (Ryan *et al.*, 1997). Aluminum also interacts with calcium by disrupting Ca^{2+} homeostasis in the cytoplasm, either by affecting its activity (Nichol *et al.*, 1993) or by interfering with its role as a secondary messenger, perhaps through the inositol 1,4,5-triphosphate pathway (Jones and Kochian, 1995). In *Arabidopsis*, Al exposure results in prolonged elevation in cytoplasmic Ca^{2+} , which occurs after growth inhibition (Jones *et al.*, 1998a). This suggests that alteration of Ca^{2+} homeostasis is not a required event for Al toxicity. In contrast, exposure to toxic concentrations of Al causes a rapid reduction in cytoplasmic Ca^{2+} in cultured tobacco

cells, which correlates with growth inhibition (Jones *et al.*, 1998b). To address this contradictory data, Plieth *et al.* (1999) constructed transgenic *Arabidopsis* expressing the calcium-activated photoprotein, aequorin, to allow measurement of Ca^{2+} in intact whole roots exposed to Al. Their results suggest that by blocking Ca^{2+} channels, Al prevents Ca^{2+} influx and makes plants vulnerable to low pH (Plieth *et al.*, 1999). However, their model cannot account for the ability of other cations to ameliorate Al toxicity (Ryan *et al.*, 1997). Aluminum could also exert its toxic effect by displacing Ca^{2+} from critical sites in the apoplasm, where Ca^{2+} normally binds to negatively charged proteins and pectins in the cell wall (Hanson, 1984; Kinraide and Parker, 1987; Kinraide *et al.*, 1994). However, this hypothesis is not supported by the observation that other cations can alleviate Al stress, some even more effectively than added Ca^{2+} (Ryan *et al.*, 1997). A number of other lesions, such as peroxidation of membrane lipids, have also been proposed as possible primary toxic lesions (Yoshino *et al.*, 1999). However, it is perhaps unlikely that a single primary toxic lesion exists due to the extreme reactivity of Al with cellular components.

1.4 Genetic Variation in Aluminum Tolerance

Many species exhibit genetic variability in their responses to Al stress. Although Al-resistant cultivars of wheat and other crop plants have been developed and studied (reviewed by Carver and Ownby, 1995), it is difficult to separate the effects of different resistance mechanisms due to their possible interactions and compensatory effects. In barley (*Hordeum vulgare*), evaluation of 37 genotypes of

diverse origin suggested that Al tolerance is controlled by a single locus with multiple alleles (Minella and Sorrels, 1992). Studies using other genotypes suggested that at least two dominant, independent loci are involved in the Al tolerance phenotype (Gallego and Benito, 1997). Although originally thought to be controlled by a single gene (*Alm1*, Moon *et al.*, 1997), Al tolerance in maize (*Zea mays*), was later shown to be controlled by two loci, one located on chromosome 6 and the other on chromosome 10 (Sibov *et al.*, 1999). In rye (*Secale cereale* L.), Al tolerance is controlled by at least two independent and dominant loci (*Alt1* and *Alt3*) located on chromosomes 6RS and 4R (Gallego *et al.*, 1998). In some wheat genotypes (for example Alikat) aluminum tolerance behaves as a single gene trait in crosses (Basu *et al.*, 1997). However, many different genes have been implicated in other populations, suggesting that Al tolerance in wheat is a dominant trait controlled by several genes (Aniol, 1991; Berzonsky, 1992). The genetic basis of Al resistance in dicots has not been extensively studied, but in soybean (*Glycine max*), it is associated with five quantitative trait loci (Bianchi-Hall *et al.*, 2000). The genetic complexity associated with Al tolerance makes it difficult to understand the contributions of different components. However, this very complexity is a potentially valuable resource for breeding programs to increase the Al tolerance of locally adapted commercial cultivars. Furthermore, molecular techniques have provided new tools to separate different components of Al resistance so that they can be studied in isolation.

1.5 Mechanisms of Aluminum Resistance

Aluminum resistance may be mediated by both internal and external mechanisms. External mechanisms detoxify Al in the apoplast or prevent it from crossing the plasma membrane to reach sensitive cytoplasmic sites, while internal mechanisms detoxify Al once it has crossed the plasma membrane, either by immobilization or sequestration (Taylor, 1991).

1.5.1. Al-induced Release of Organic Anions

One example of an external Al resistance mechanism is Al-induced release of organic anions, which chelate Al in the apoplast and limit its entry into the symplast. Increased exudation of organic anions such as citrate and malate correlates with increased Al resistance in snapbean (Miyasaka *et al.*, 1991), wheat (Delhaize *et al.*, 1993; Ryan *et al.*, 1994; Basu *et al.*, 1994c) and maize (Pellet *et al.*, 1994). Addition of malate to the growth medium restores root elongation to control levels in an Al-sensitive cultivar of wheat (Basu *et al.*, 1994c), suggesting that malate plays a role in Al resistance. The role of organic anions in Al tolerance has been investigated directly using transgenic plants. Overexpression of citrate synthase in transgenic tobacco (*Nicotiana tabacum*) and papaya (*Carica papaya*) caused increased citrate efflux, which correlated with Al tolerance (de la Fuente *et al.*, 1997). This interesting result was not reproducible, however, since subsequent studies using independent lines of transgenic tobacco found that overexpression of citrate synthase had no effect on citrate efflux levels or Al tolerance (Delhaize *et al.*, 2001). Since

organic acids exist as organic acid anions in the cytoplasm, involvement of Al-activated anion channels in Al resistance has been hypothesized (Delhaize and Ryan, 1994). This is supported by evidence showing that anion channel antagonists block Al-induced efflux of malate in an Al-tolerant cultivar of wheat (Ryan *et al.*, 1994). Protein kinase inhibitors can also inhibit malate efflux, suggesting that activation of an Al-activated plasma membrane anion channel may be controlled by protein phosphorylation (Osawa and Matsumoto, 2001). Aluminum-activated, malate-permeable channels have been characterized using the patch-clamp technique in maize and wheat (Pineros and Kochian, 2001; Zhang *et al.*, 2001). Continuing research in this area has focused on identification of these channels.

1.5.2 Plant-Induced Elevation of Rhizosphere pH

A second mechanism of external Al resistance is plant-induced elevation of rhizosphere pH. This is an attractive hypothesis because the chemistry of Al favours formation of insoluble and/or nontoxic Al species at near neutral pH. Early studies using 20 winter and 20 spring wheat cultivars observed a negative correlation between cultivar tolerance to Al and pH changes of nutrient solutions (Taylor and Foy, 1985 a,b). These pH differences were caused by differences in ammonium and nitrate nutrition (Taylor and Foy, 1985c), but subsequent studies found that the relative tolerance of Atlas-66 (Al-resistant) and Scout-66 (Al-sensitive) was unaffected by the ratio of ammonium:nitrate and by solution pH (Taylor, 1987). There were also no discernable differences in the rhizosphere pH of Atlas-66 and

Scout-66, as measured with pH microelectrodes (Miyasaka *et al.*, 1989). However, recent experiments suggest that plant-induced elevation of rhizosphere pH may occur in *Arabidopsis*. An Al-resistant mutant exhibits an Al-induced increase in rhizosphere pH mediated by increased net proton influx into the cytoplasm (Degenhardt *et al.*, 1998).

1.5.3 Exudation of Al-binding Polypeptides

A third mechanism of external Al resistance is the exudation of polypeptides which may bind Al or counteract its growth-inhibitory effects in the apoplasm. Aluminum-resistant and Al-sensitive cultivars of wheat show differences in polypeptide exudation by roots under conditions of Al stress (Delhaize *et al.*, 1990). In Al-resistant cultivars, specific polypeptides accumulate in exudates and have greater association with Al than those exuded by sensitive cultivars. This implies that polypeptides specifically produced by Al-resistant cultivars may facilitate external Al resistance (Basu *et al.*, 1994b, 1997), although the physiological significance of most of these proteins has not yet been ascertained. A 23 kD root exudate protein from an Al-resistant isolate of wheat (Alikat = Katepwa*3/Maringa) is induced by Al treatment, binds Al and segregates with Al resistance in the progeny of a cross between doubled-haploid lines of Alikat (Al-resistant) and Katepwa (Al-sensitive). This suggests that this 23 kD protein is involved in mediating Al resistance (Basu *et al.*, 1997). The purified protein has been identified as an apoplastic form of

superoxide dismutase, and may be involved in preventing oxidative damage caused by Al (Basu *et al.*, 2001).

1.5.4 Support for External Resistance from Inhibitor Studies

The involvement of external resistance mechanisms, such as organic anion and polypeptide export, is supported by the effects of biological inhibitors on Al uptake. Excised roots from Al-resistant wheat cultivars show increased net Al uptake when exposed to dinitrophenol (DNP), an uncoupler of oxidative phosphorylation. A similar effect has been observed in intact single cells of the giant algae, *Chara corallina*, where treatment with DNP causes increased net Al uptake into the protoplasm (Taylor *et al.*, 2000). Based on these observations, it appears that an energy-dependent exclusion mechanism is involved in Al-resistance, perhaps mediated by exudation of chelator ligands such as malate and/or polypeptides to the rhizosphere (Zhang and Taylor, 1991). Cycloheximide, an inhibitor of cytoplasmic protein synthesis, decreases Al uptake by Al-sensitive cultivars, but not by resistant cultivars, suggesting that proteins encoded by nuclear genes might be involved in regulating uptake of Al across the plasma membrane and may play a role in Al susceptibility (Zhang *et al.*, 1995).

1.5.5 Modulation of Intracellular Enzyme Activity

Observations of changing intracellular enzyme activity support the involvement of internal mechanisms in Al resistance. Activities of intracellular

enzymes such as glucose-6-phosphate dehydrogenase, 6-phosphogluconate dehydrogenase (Slaski *et al.*, 1996) and vacuolar H⁺-ATPase (Kasai *et al.*, 1992, 1993; Hamilton *et al.*, 2001) are stimulated by Al. Transcripts encoding other intracellular proteins such as glutathione S-transferase (Richards *et al.*, 1998), and a fimbrin-like protein (Cruz-Ortega *et al.*, 1997) from wheat, and transcripts encoding aldolase, peroxidase, glutathione S-transferase, blue-copper-binding protein, superoxide dismutase and a reticuline:oxygen oxidoreductase homolog from *Arabidopsis* are also induced by Al (Richards *et al.*, 1998). Most of these Al-induced gene products (Table 1-1) appear to be involved in general stress response, since they are also induced by oxidative stress (Richards *et al.*, 1998), other toxic metals, low calcium levels and wounding (Snowden *et al.*, 1995; Hamel *et al.*, 1998). The involvement of blue-copper-binding protein, glutathione S-transferase, peroxidase and GDP-dissociation inhibitor in Al resistance has been confirmed in transgenic *Arabidopsis*, where their overexpression confers increased Al resistance (Ezaki *et al.*, 2000).

1.5.6 Sequestration of Al in the Vacuole

Perhaps the most compelling example of a possible internal Al resistance mechanism is sequestration of Al in the vacuole, where it would likely be chelated by organic acids or phosphates. Although measurement of intracellular Al is technically difficult, Al uptake across the plasma membrane has been arguably demonstrated in

Table 1-1. Classification of AI-induced genes.

Gene	Identity	Species	Reference
Pathogenesis-Related			
<i>Wali5</i>	Bowman-Birk protease inhibitor	<i>T. aestivum</i>	Snowden and Gardner, 1993
<i>AtBP1</i>	Bowman-Birk protease inhibitor	<i>A. thaliana</i>	Richards <i>et al.</i> , 1997
Oxidative Stress-Related			
<i>AtBCB</i>	Blue copper binding protein	<i>A. thaliana</i>	Richards <i>et al.</i> , 1997
<i>AtPOX</i>	Peroxidase	<i>A. thaliana</i>	Richards <i>et al.</i> , 1997
<i>AtSOD</i>	Superoxide dismutase	<i>A. thaliana</i>	Richards <i>et al.</i> , 1997
<i>TaSOD</i>	Superoxide dismutase	<i>T. aestivum</i>	Basu <i>et al.</i> , 2001
<i>NtPOX</i>	Reticuline:oxygen oxidoreductase	<i>A. thaliana</i>	Richards <i>et al.</i> , 1997
<i>ParB</i>	Peroxidase	<i>N. tabacum</i>	Ezaki <i>et al.</i> , 1995
	Glutathione S-transferase	<i>N. tabacum</i>	Ezaki <i>et al.</i> , 1995
General Stress-Related			
<i>HSP150</i>	Heat shock protein	<i>S. cerevisiae</i>	Ezaki <i>et al.</i> , 1998
Unknown Function			
<i>SED1</i>	Putative membrane protein	<i>S. cerevisiae</i>	Ezaki <i>et al.</i> , 1998
<i>Para</i>	No homologs	<i>N. tabacum</i>	Ezaki <i>et al.</i> , 1995
<i>Wali1</i>	Homology to metallothioneins	<i>T. aestivum</i>	Snowden and Gardner, 1993
<i>Wali2</i>	No Homologs	<i>T. aestivum</i>	Snowden and Gardner, 1993
<i>Wali3</i>	No Homologs	<i>T. aestivum</i>	Snowden and Gardner, 1993
<i>Wali4</i>	Phenylalanine ammonia lyase	<i>T. aestivum</i>	Snowden and Gardner, 1993
<i>Wali6</i>		<i>T. aestivum</i>	Snowden and Gardner, 1994
<i>Wali7</i>	Fimbrin-like protein	<i>T. aestivum</i>	Snowden and Gardner, 1994
	Aldolase	<i>T. aestivum</i>	Cruz-Ortega <i>et al.</i> , 1997
		<i>A. thaliana</i>	Richards <i>et al.</i> , 1998

Saccharomyces cerevisiae (Ezaki *et al.*, 1999; Jagannatha Rao *et al.*, 1997), *Triticum aestivum* (Zhang and Taylor, 1989), *Phaseolus vulgaris* (McDonald-Stephens and Taylor, 1995), *Glycine max* (Lazof *et al.*, 1994) and the algae *Chara corallina* (Taylor *et al.*, 2000). It is generally agreed that the majority of Al is associated with the cell wall, but it is also taken up into the vacuole (Taylor *et al.*, 2000). Based on observations using the Al-binding fluorescent dye, morin, Ezaki *et al.* (1999) suggested that intracellular Al does not associate with the yeast vacuole. However, low levels such as those measured in *Chara* (Taylor *et al.*, 2000) would not be detected using this method. The mechanism by which sequestration in the vacuole may occur is unknown, but Al may enter the vacuole nonspecifically via transporters of essential metals or as an unidentified conjugate. In either case, vacuolar ion uptake utilizes proton motive force generated primarily by the vacuolar H⁺-ATPase (V-ATPase).

1.6 Structure of the Vacuolar H⁺-ATPase


The vacuolar H⁺-ATPase (V-ATPase) and the closely related mitochondrial F₁F₀-ATPase are complex, multisubunit enzymes (Fig 1-2, Table 1-2). They share both a common structure and a common mechanism of action. Their structures consist of catalytic (V₁ and F₁, respectively) and membrane sectors (V₀ and F₀,

Table 1-2. Subunit composition of the vacuolar ATPase (V-ATPase) and functional homologies to subunits of the mitochondrial ATP synthase (F_1F_0 -ATPase).

V-ATPase Subunit	Functional Homolog in F ₁ F ₀ -ATPase	Function	Molecular Mass (kD)
V₁ Constituents		ATP Binding and Catalysis	500-600
A	β	ATP hydrolysis (synthesis in F ₁)	67-73
B	α	ATP binding, activity regulation*	55-60
C	unknown	Assembly*	44
D	γ	Spindle*	32
E	γ	Spindle*	31
F	ε	V ₁ -V ₀ contact, regulation*	14
G	F ₀ -b	Stabilization	12-16
H	unknown	V ₁ stabilization	50-54
V₀ Constituents		Proton Translocation	250-300
a	b	Activity, assembly, targeting	97-115
d	unknown	Assembly, stabilization	32-38
c	c	Proton translocation	16
c'	c	Proton translocation	17
c''	c	Proton translocation	17

*These functions have been tentatively assigned based on similarity to F₁F₀-ATPases

Figure 1-2. Structural comparison of the vacuolar ATPase (V-ATPase) and mitochondrial ATP synthase (F_1F_0 -ATPase). A generalized V-ATPase is shown on the left and a generalized F_1F_0 -ATPase is shown on the right. The functions (if known) of each subunit are described in Table 1-2 and in the text. Adapted in part from Sze *et al.*, 1999.



respectively). The catalytic sectors are responsible for energy conversion, while the membrane sectors are responsible for proton transport through the membrane. The V-ATPase is localized to most membranes of the secretory pathway and is particularly abundant in the tonoplast of the plant vacuole. It functions in energizing these membranes to allow secondary active transport, and in acidifying internal compartments. The F_1F_0 -ATPase is localized to the mitochondria and chloroplasts, where it functions predominantly in ATP synthesis. Interestingly, the V-ATPase is essential in all organisms tested, with the exception of yeast, while many types of cells can survive without a functional F_1F_0 -ATPase (King and Attardi, 1989; Dow *et al.*, 1997; Bowman *et al.*, 2000). The characteristic phenotype of yeast mutants lacking V-ATPase activity is sensitivity to calcium in the growth medium, inability to grow in medium buffered to neutral pH and inability to utilize nonfermentable carbon sources (Graham *et al.*, 2000).

The catalytic sector of the V-ATPase consists of three copies each of the A and B subunits, which are approximately 25% identical at the amino acid level to the β and α subunits of the F_1F_0 -ATPase, respectively. This implies evolution from common ancestral genes (Holzenburg *et al.*, 1993; Davies *et al.*, 1996; Bowman *et al.*, 1988; Manolson *et al.*, 1988; Nelson *et al.*, 1989). Subunit A (67-73 kD) contains the ATP hydrolytic domain (Arai *et al.*, 1987), while subunit B (55-60 kD) is responsible for ATP binding, and may be involved in regulation of V-ATPase activity (Manolson *et al.*, 1985; Zhang *et al.*, 1995; Peng, 1995). The function of Subunit C (44 kD) is not known, but it may be involved in assembly of V_1 (Ho *et al.*, 1993).

Subunits D (32 kD) and E (31 kD) show regions of similarity to the γ subunit of F_1 , and may functionally associate to form a homolog of the F_1 spindle (Tomashek *et al.*, 1996). Subunit F (14 kD), analogous to the ϵ subunit of F_1 , may have a role in stabilizing contacts between V_1 and V_0 (Graham *et al.*, 1994) or it may have a regulatory role (Nelson *et al.*, 1994). Subunit G (16 kD) shows regions of similarity to the stabilizing b subunit of F_0 , although in the *Manduca sexta* midgut, it is present in three copies rather than the two copies of subunit b found in the F_1F_0 -ATPase (Supekova *et al.*, 1995).

The membrane sectors of the V-ATPase and F_1F_0 -ATPase (V_0 and F_0 , respectively) have similar functions in proton translocation across the membrane, but V_0 has some unique properties. The proteolipid subunits (subunit c in both enzymes) are homologous, but the V_0 proteolipid is 16 kD while the F_0 proteolipid is only 8 kD (Mandel *et al.*, 1988). This difference may have arisen by gene duplication and fusion of an ancestral gene homologous to that encoding the F_0 proteolipid (Nelson, 1992). In yeast, the V_0 sector contains several copies of subunit c and one copy each of homologous proteolipids, c' and c'' , all of which are required for activity (Powell *et al.*, 2000). In contrast, the F_0 sector consists of variable numbers of subunit c (Hirata *et al.*, 1997, Powell *et al.*, 2000). These differences are thought to affect protein packing within the proton channel. The channel in V_0 is wider than that in F_0 , allowing for proton slip when the proton motive force (pmf, $\Delta\mu_{H^+}$) becomes too great to allow further proton pumping (Perzov *et al.*, 2001).

1.7 Function of the Vacuolar H⁺-ATPase

In a typical plant cell, activity of the V-ATPase, using ATP produced by the F₁F₀-ATPase and in conjunction with the vacuolar H⁺-pyrophosphatase, is responsible for maintaining a proton electrochemical gradient across the membranes of endomembrane compartments. These include the endoplasmic reticulum, golgi, coated vesicles, provacuoles and vacuoles (Herman *et al.*, 1994; Robinson *et al.*, 1996). The electrochemical gradient allows secondary active transport of cationic solutes into and anionic solutes out of the vacuolar lumen. Energetically favourable transport of cations out of and anions into the lumen is carried out by electrogenic channels. Due to its wide distribution within the cell, the V-ATPase is thought to be involved in a number of cellular activities in addition to ion transport. For example, vesicle acidification is required for ligand-receptor dissociation, protein-protein interaction (Stevens and Forgac, 1997), protein trafficking (Morano and Klionsky, 1994) and membrane fusion (Ungermann *et al.*, 1998). These processes are required for processes such as vacuole biogenesis, cell division and secretion.

1.8 Control of ATPase Expression

Activities of cellular proton pumps, which use ATP generated by the F₁F₀-ATPase, are coordinated in response to growth, development and changes in the environment. For example, transport activities of the V-ATPase and plasma membrane H⁺-ATPase (P-ATPase) regulate turgor pressure during growth.

Coordinated activity of these enzymes is also required to maintain pH homeostasis during pH changes in the environment (Young *et al.*, 1998). This is achieved by sequestration of protons in the vacuole by increased V-ATPase activity combined with extrusion of protons from the cytoplasm by increased P-ATPase activity. A similar mechanism is used by NaCl tolerant plants, which sequester Na⁺ in the vacuole and also actively transport Na⁺ out of the cell across the plasma membrane. This phenomenon has been extensively studied in the facultative halophyte, *Mesembryanthemum crystallinum* (iceplant), and in the halotolerant sugar beet, *Beta vulgaris*. In both species, V-ATPase expression is stimulated by NaCl treatment (Ratajczak *et al.*, 1994; Kirsch *et al.*, 1996). Increased activity of V-ATPase allows accumulation of Na⁺ in the vacuole via the Na⁺/H⁺ antiport protein, NHX1. The importance of vacuolar H⁺/Na⁺ antiport has been confirmed by overexpressing *NHX1* in transgenic *Arabidopsis*, where it causes enhanced NaCl tolerance (Apse *et al.*, 1999). Toxic Na⁺ ions are also removed from the cytoplasm by active transport through the plasma membrane Na⁺ transporter, ENA1, which is energized by the plasma membrane H⁺-ATPase (Garcia-deblas *et al.*, 1993).

The mechanisms by which vacuolar ATPase activity is modulated by developmental stage and the environment are not clear. Short-term adjustments in activity may occur via reversible dissociation of the peripheral V₁ subunits from the V₀ sector, as has been observed during cold stress in *Vigna radiata* (Matsuura-Endo *et al.*, 1992) and during glucose deprivation in *Saccharomyces cerevisiae* (Kane, 1995). Vacuolar ATPase activity is also affected by Cl⁻ levels, pH and the ratio of ATP to

ADP (Dietz *et al.*, 2001), although there is no evidence for a direct regulatory role of these factors. The lipid environment also has an effect on V-ATPase activity. V-ATPase activity is stimulated by phospholipids (Yamanishi and Kasamo, 1993), while it is inhibited by glycolipids (Kasamo *et al.*, 2000). This could be caused by changes in membrane fluidity or by conformational changes in the V-ATPase (Kasamo *et al.*, 2000). The presence of additional subunits can also affect V-ATPase activity. This has been best characterized in the salt response of *M. crystallinum*, where a change in V-ATPase subunit structure (induced subunits E_i and D_i) during salt exposure correlates with increased enzyme activity (Ratajczak *et al.*, 2000). Several small proteins have also been identified as inhibitors (Zhang K *et al.*, 1992) and activators (Xie *et al.*, 1993; Zhang *et al.*, 1992) of V-ATPase activity. Interestingly, the V-ATPase also responds to the redox status of the cell. The V-ATPase A and E subunits contain cysteine residues which form intramolecular disulphide bonds under oxidizing conditions (Feng and Forgac, 1994; Dietz *et al.*, 2001) when the cytoplasm is not acidified. In the event of cytoplasmic acidosis, this inhibition cannot occur, further illustrating the importance of the V-ATPase in stress responses (Dietz *et al.*, 2001).

Long-term adjustments in V-ATPase activity are thought to be controlled at the level of gene expression, although this phenomenon has not been studied in detail. Transcript levels of the V-ATPase A subunit increase in response to salt stress in *Nicotiana tabacum* (Narasimhan *et al.*, 1991) and *Lycopersicon esculentum* (Binzel and Dunlap, 1994). In *M. crystallinum*, levels of transcripts encoding the V-ATPase

A, B, E and c subunits increase in response to salt stress (Low *et al.*, 1996; Tsiantis *et al.*, 1996; Golldack and Dietz, 2001). The effect of salt or other stresses on levels of transcripts encoding the other V-ATPase subunits has not been examined.

Interestingly, levels of the transcript encoding the V-ATPase c subunit also show diurnal regulation in leaves (Low *et al.*, 1996), although under normal growth conditions levels of transcripts encoding the V-ATPase A and B subunits remain constant. It is unknown whether the above effects on V-ATPase gene expression in response to developmental and environmental changes are due to transcriptional regulation or posttranscriptional events.

1.9 Objectives

Two 51 kD microsomal proteins are induced in roots of the Al-resistant cultivar of wheat, PT741, under conditions of Al stress (Basu *et al.*, 1994a). These proteins show dose-dependent accumulation in the presence of Al and disappear within 24 h upon removal of Al stress (Basu *et al.*, 1994a). The proteins are localized at the tonoplast of root tip cells, and segregate with the resistance phenotype (Taylor *et al.*, 1997). This is an exciting result because it suggests that at least one of these proteins may be involved in Al resistance. Localization to the tonoplast also suggests that they may be involved in a sequestration mechanism of Al resistance (Taylor *et al.*, 1997).

Differential expression of these 51 kD proteins in an Al-resistant cultivar of wheat raises several questions. Does expression of the proteins contribute to the

resistance phenotype? How is expression of the proteins regulated? What would be the effect of expressing the proteins in a heterologous, Al-sensitive genotype? To address these questions, my objectives were to purify and identify the 51 kD, Al-induced proteins from a tonoplast-enriched membrane fraction of Al-resistant wheat, to evaluate expression of the proteins, and to determine the effect of altering expression of the genes encoding these proteins in a heterologous system. My hypothesis is that eliminating expression of one or both 51 kD proteins will confer Al hypersensitivity to a heterologous genotype. Understanding the function of these Al-induced proteins will add to our knowledge of Al stress physiology, with potentially significant agricultural and economic impact. If we can alter expression of the 51 kD proteins, it may be possible to produce transgenic lines of common crop plants such as canola (*Brassica napus*) and wheat showing enhanced resistance to Al. This could increase crop yield in the presence of soluble Al, and contribute to improving agriculture in areas of soil acidity.

In Chapter Two, I describe purification of the 51 kD, Al-induced proteins from an endomembrane-enriched membrane fraction isolated from an Al-resistant cultivar of *Triticum aestivum* L. (cv PT741) grown in the presence of Al. The identity of these proteins has been determined using peptide microsequence data. Purified peptides were homologous to the B subunit of the vacuolar H⁺-ATPase (V-ATPase) and to the α and β subunits of F₁F₀-ATPase (mitochondrial ATP synthase). Both V-ATPase and F₁F₀-ATPase activities were induced by Al treatment in PT741 and not

in other cultivars tested. These results suggest that upregulation of ATPase activity in PT741 may be an adaptive response involved in Al resistance.

Yeast has previously been used as a model system to examine Al stress (MacDiarmid and Gardner 1996, 1998; Ezaki *et al.*, 1998, 1999). Use of yeast as a model is uniquely appropriate here, because yeast is the only eukaryote tested thus far that does not require a functional V-ATPase for growth (Dow *et al.*, 1997; Bowman *et al.*, 2000). In Chapter Three, I describe the use of the yeast model system to simply and directly test my hypothesis that upregulation of V-ATPase and F₁F₀-ATPase activity in PT741 may be an adaptive response involved in Al resistance. Increased activity of the V-ATPase, using ATP provided by the F₁F₀-ATPase, could be required to energize the tonoplast membrane, allowing transport of Al-complexing compounds such as organic acids out of the vacuole. It is also possible that increased V-ATPase activity is required to allow sequestration of Al within the vacuole. Although V-ATPase transcript and protein levels, as well as enzyme activity, are not affected by increasing Al concentrations in yeast as they are in plants, the observation that V-ATPase mutants are hypersensitive to Al supports my hypothesis. In wild-type yeast, F₁F₀-ATPase transcript, protein and activity levels all increase in response to Al, while F₁F₀-ATPase mutants do not show an Al hypersensitive phenotype.

To further examine the importance of the V-ATPase and the vacuole in metal tolerance, I investigated the effect of vacuolar ATPase mutations on salt (NaCl) tolerance in Chapter Four. Salt tolerance in yeast is a complex trait, involving regulation of membrane polarization, Na⁺ efflux and sequestration of Na⁺ in the

vacuole. Since transmembrane transport energized by H⁺-ATPases is common to all of these tolerance mechanisms, the responses of the plasma membrane H⁺-ATPase, vacuolar H⁺-ATPase and mitochondrial F₁F₀-ATPase to NaCl stress were characterized. Since the vacuolar ATPase is responsible for generating the proton motive force required for import of cations (such as Na⁺) into the vacuole, strains lacking this activity should be hypersensitive to NaCl. We found that strains lacking vacuolar ATPase activity were in fact hypersensitive to NaCl, while strains lacking ATP synthase were not.

My observations in wheat and yeast suggest that V-ATPase activity is required for Al and NaCl tolerance. Examining the effect of V-ATPase mutations in plants, however, would provide the most convincing test possible of our hypothesis. Therefore, in Chapter Five, I describe the construction and phenotypic analysis of transgenic *Arabidopsis thaliana* expressing a cDNA encoding the V-ATPase B subunit in the antisense orientation. Decreased V-ATPase activity caused transgenic plants to be hypersensitive to both Al and NaCl, as measured by differences in root elongation. This approach allowed me to directly test my hypothesis and further characterize the involvement of the V-ATPase in mediating Al and NaCl tolerance.

1.10 Literature cited

Aniol A (1991) Genetics of acid tolerant plant. In Wright RJ, Baligar VC and Murrman RP (eds). Plant-soil interactions at low pH. Proceedings of the Second International Symposium on Plant-Soil Interactions at Low pH. pp 1007-1017

Apse MP, Aharon GS, Snedden WA and Blumwald E (1999) Salt tolerance conferred by overexpression of a vacuolar Na⁺/H⁺ antiport in Arabidopsis. Science **285**: 1256-1258

Arai H, Berne M and Forgac M (1987) Inhibition of the coated vesicle proton pump and labeling of a 17,000-dalton polypeptide by N,N'-dicyclohexylcarbodiimide. J Biol Chem **262**: 11006-11011

Baligar VC, Wright RJ, Kinraide TB, Foy CD and Elgin JH Jr. (1987) Aluminum effects on growth, mineral uptake and efficiency ratios in red clover cultivars. Agron J **79**:1038-1044

Basu U, Basu A and Taylor GJ (1994a) Induction of microsomal membrane proteins in roots of an aluminum-resistant cultivar of *Triticum aestivum* L. under conditions of aluminum stress. Plant Physiol **104**: 1007-1013

Basu U, Basu A and Taylor GJ (1994b) Differential exudation of polypeptides by roots of aluminum-resistant and aluminum-sensitive cultivars of *Triticum aestivum* L. in response to aluminum stress. Plant Physiol **106**: 151-158

- Basu U, Godbold D and Taylor GJ (1994c)** Aluminum resistance in *Triticum aestivum* associated with enhanced exudation of malate. *J Plant Physiol* **144**: 747-753
- Basu U, Good AG and Taylor GJ (2001)** Transgenic *Brassica napus* plants overexpressing aluminium-induced mitochondrial manganese superoxide dismutase cDNA are resistant to aluminium. *Plant Cell Environ* (In Press)
- Bennett RJ, Breen CM and Fey MV (1987)** The effects of aluminum on root cap function and root development in *Zea mays* L. *Environ Exp Bot* **27**: 91-104
- Berzonsky WA (1992)** The genomic inheritance of aluminum tolerance in 'Atlas 66' wheat. *Genome* **35**: 689-693
- Bianchi-Hall CM, Carter TE Jr, Bailey MA, Mian MAR, Rufty TW, Ashley DA, Boerma HR, Arellano C, Hussey RS and Parrott WA (2000)** Aluminum tolerance associated with quantitative trait loci derived from soybean PI 416937 in hydroponics. *Crop Sci* **40**: 538-545
- Bowman BJ, Allen R, Wechsler MA and Bowman EJ (1988)** Isolation of genes encoding the *Neurospora* vacuolar ATPase. Analysis of *vma-2* encoding the 57-kDa polypeptide and comparison to *vma-1*. *J Biol Chem* **263**: 14002-14007
- Bowman BJ, Dschida WJ, Harris T and Bowman EJ (1988)** Bafilomycins: a class of inhibitors of membrane ATPases from microorganisms, animal cells and plant cells. *Proc Natl Acad Sci USA*. **85**: 7972-7976

- Bowman EJ, Kendle R and Bowman BJ** (2000) Disruption of *vma-1*, the gene encoding the catalytic subunit of the vacuolar H(+)-ATPase, causes severe morphological changes in *Neurospora crassa*. *J Biol Chem* **275**: 167-176
- Carver BF and Ownby JD** (1995) Acid soil tolerance in wheat. *Adv Agron* **54**: 117-173
- Cruz-Ortega R, Cushman JC and Ownby JD** (1997) cDNA clones encoding 1,3-beta-glucanase and a fimbrin-like cytoskeletal protein are induced by Al toxicity in wheat roots. *Plant Physiol* **114**: 1453-1460
- Davies SA, Kelly DC, Goodwin SF, Wang S-Z, Kaiser K and Dow JAT** (1996) Analysis and inactivation of *vha55*, the gene encoding the vacuolar ATPase B-subunit in *Drosophila melanogaster* reveals a larval lethal phenotype. *J Biol Chem* **271**: 30677-30684
- Delhaize E, Hebb DM and Ryan PR** (2001) Expression of a *Pseudomonas aeruginosa* citrate synthase gene in tobacco is not associated with either enhanced citrate accumulation or efflux. *Plant Physiol* **125**: 2059-67
- Delhaize E, Higgins TJV and Randall PJ** (1991) Aluminum tolerance in wheat: analysis of polypeptides in the root apices of tolerant and sensitive genotypes. *In* Wright, R.J., Baligar, V.C., and Murrman, R.P., eds. *Plant-soil interactions at low pH. Proceedings of the second International Symposium on Plant-Soil Interactions at Low pH, 24-29 June 1990, Beckley, West Virginia, USA.* pp 1071-1079. Kluwer Academic Publishers, Dordrecht, The Netherlands.

- Degenhardt J, Larsen PB, Howell SH, and Kochian LV** (1998) Aluminum resistance in the *Arabidopsis* mutant *alr-104* is caused by an aluminum-induced increase in rhizosphere pH. *Plant Physiol* **117**: 19-27
- Dietz KJ, Tavakoli N, Kluge C, Mimura T, Sharma SS, Harris GC, Chardonnens AN and Golldack D** (2001) Significance of the V-type ATPase for the adaptation to stressful growth conditions and its regulation on the molecular and biochemical level. *J Exp Bot* **52**: 1969-1980
- Ding JP, Badot P-M and Pickard BG** (1993) Aluminum and hydrogen ions inhibit a mechansosensory calcium-selective cation channel. *Aust J Plant Physiol* **20**: 771-778
- Dow JAT, Davies SA, Guo Y, Graham S, Finbow M and Kaiser K** (1997) Molecular genetic analysis of V-ATPase function in *Drosophila melanogaster*. *J Exp Biol* **200**: 237-245
- Ezaki B, Gardner RC, Ezaki Y, Kondo H and Matsumoto H** (1998) Protective roles of two aluminum (Al)-induced genes, *HSP150* and *SEDI* of *Saccharomyces cerevisiae*, in Al and oxidative stresses. *FEMS Microbiol Lett* **159**: 99-105
- Ezaki B, Sivaguru M, Ezaki Y, Matsumoto H and Gardner RC** (1999) Acquisition of aluminum tolerance in *Saccharomyces cerevisiae* by expression of the *BCB* or *NtGDII* gene derived from plants. *FEMS Microbiol Lett* **171**: 81-87

- Ezaki B, Gardner RC, Ezaki Y and Matsumoto H** (2000) Expression of aluminum-induced genes in transgenic *Arabidopsis* plants can ameliorate aluminum stress and/or oxidative stress. *Plant Physiol* **122**: 657-665
- de la Fuente JM, Ramirez-Rodriguez V, Cabrera-Ponce JL and Herrera-Estrella L** (1997) Aluminum tolerance in transgenic plants by alteration of citrate synthesis. *Science* **276**: 1566-1568
- Gallego FJ and Benito C** (1997) Genetic control of aluminum tolerance in rye (*Secale cereale* L.). *Theor Appl Genet* **95**: 393-399
- Gallego FJ, Calles B and Benito C** (1998) Molecular markers linked to the aluminium tolerance gene Alt1 in rye (*Secale cereale* L.). *Theor Appl Genet* **97**: 1104-1109
- Graham LA, Hill KJ and Stevens TH** (1994) VMA7 encodes a novel 14-kDa subunit of the *Saccharomyces cerevisiae* vacuolar H(+)-ATPase complex. *J Biol Chem* **269**: 25974-25977
- Graham LA, Powell B and Stevens TH** (2000) Composition and assembly of the yeast vacuolar H⁺-ATPase complex. *J Exp Biol* **203**: 61-70
- Hamel F, Breton C and Houde M** (1998) Isolation and characterization of wheat aluminum-regulated genes: Possible involvement of aluminum as a pathogenesis response regulator. *Planta* **205**: 531-538
- Hanson JB** (1984) The functions of calcium in plant nutrition. In **PB Tinker, A Lauchli**, eds, *Advances in Plant Nutrition*, Vol 1. Praeger Scientific, New York, pp 149-208

Herman EM, Li X, Su RT, Larsen P, Hsu H-T and Sze H (1994) Vacuolar-type H⁺-ATPases are associated with the endoplasmic reticulum and provacuoles of root tip cells. *Plant Physiol* **106**: 1313-1324

Hirata R, Graham LA, Takatsuki A, Stevens TH and Anraku Y (1997) VMA11 and VMA16 encode second and third proteolipid subunits of the *Saccharomyces cerevisiae* vacuolar membrane H⁺-ATPase. *J Biol Chem* **272**: 4795-4803

Ho MN, Hill KJ, Lindorfer MA and Stevens TH (1993) The *Saccharomyces cerevisiae* VMA6 gene encodes the 36-kDa subunit of the vacuolar H(+)-ATPase membrane sector. *J Biol Chem* **268**: 221-227

Holzenburg A, Jones PC, Franklin T, Pali T, Heimborg T, Marsh D, Findlay JBC and Finbow ME (1993) Evidence for a common structure for a class of membrane channels. *Eur J Biochem* **213**: 21-30

Jagannatha Rao KS and Easwaran KKK (1997) ²⁷Al-NMR studies of aluminum transport across yeast cell membranes. *Mol Cell Biochem* **175**: 59-63

Jones DL, Gilroy S, Larsen PB, Howell SH and Kochian LV (1998a) Effect of aluminum on cytoplasmic Ca²⁺ homeostasis in root hairs of *Arabidopsis thaliana* (L.). *Planta* **206**: 378-387

Jones DL, Kochian LV and Gilroy S (1998b) Aluminum induces a decrease in cytosolic calcium concentration in BY-2 tobacco cell cultures. *Plant Physiol* **116**: 81-89

Jones DL and Kochian LV (1995) Aluminum inhibition of the inositol 1,4,5-triphosphate signal transduction pathway in wheat roots: a role in aluminum toxicity? *Plant Cell* **7**: 1913-1922

Kane PM (1995) Disassembly and reassembly of the yeast vacuolar H⁺-ATPase in vivo. *J Biol Chem* **270**: 17025-17032

Kasai M, Sasaki M, Tanakamaru S, Yamamoto Y and Matsumoto H (1993) Possible involvement of abscisic acid in increases in activities of two vacuolar H⁺-pumps in barley roots under aluminum stress. *Plant Cell Physiol* **34**: 1335-1338

Kasai M, Sasaki M, Yamamoto Y and Matsumoto H (1992) Aluminum stress increases K⁺ efflux and activities of ATP- and P_i-dependent H⁺ pumps of tonoplast-enriched membrane vesicles from barley roots. *Plant Cell Physiol* **33**: 1035-1039

Kasamo K, Yamaguchi M and Nakamura Y (2000) Mechanism of the chilling-induced decrease in proton pumping across the tonoplast of rice cells. *Plant Cell Physiol* **41**: 840-849

King MP and Attardi G (1989) Human cells lacking mtDNA: Repopulation with exogenous mitochondria by complementation. *Science* **246**: 500-503

Kinraide TB and Parker DR (1987) Cation amelioration of aluminum toxicity in wheat. *Plant Physiol* **83**: 546-551

- Kinraide TB, Ryan PR and Kochian LV** (1994) Al^{3+} - Ca^{2+} interactions in aluminum rhizotoxicity. II. Evaluating the Ca^{2+} -displacement hypothesis. *Planta* **192**: 104-109
- Kirsch M, An Z, Viereck R, Low R and Rausch T** (1996) Salt stress induces an increased expression of V-type H^{+} -ATPase in mature sugar beet leaves. *Plant Mol Biol* **32**: 543-7
- Kochian LV** (1995) Cellular mechanisms of aluminum toxicity and tolerance in plants. *Annu Rev Plant Physiol Plant Mol Biol* **46**: 237-260
- Lazof DB, Goldsmith JG, Rufty GW and Linton RW** (1994) Rapid uptake of aluminum into cells of intact soybean root tips. A microanalytical study using secondary ion mass spectrometry. *Plant Physiol* **106**: 1107-1114
- Mandel M, Moriyama Y, Hulmes JD, Pan Y-CE, Nelson H and Nelson N** (1988) cDNA sequence encoding the 16-kDa proteolipid of chromaffin granules implies gene duplication in the evolution of H^{+} -ATPases. *PNAS* **85**: 5521-5524
- MacDiarmid CW and Gardner RC** (1996) Al toxicity in yeast: A role for Mg? *Plant Physiol* **112**: 1101-1109
- MacDiarmid CW and Gardner RC** (1998) Overexpression of the *Saccharomyces cerevisiae* magnesium transport system confers resistance to aluminum ion. *J Biol Chem* **273**: 1727-1732
- Manolson MF, Rea PA and Poole RJ** (1985) Identification of 3-O-(4-benzoyl)benzoyl adenosine 5'-triphosphate- and N,N'-

dicyclohexylcarbodiimide-binding subunits of a higher plant H⁺-translocating tonoplast ATPase. *J Biol Chem* **260**: 12273-12279

Manolson MF, Ouellette BFF, Filion M and Poole RJ (1988) cDNA sequence and homologies of the "57-kDa" nucleotide-binding subunit of the vacuolar ATPase from *Arabidopsis*. *J Biol Chem* **263**: 17987-17994

Matsumoto H and Morimura S (1980) Repressed template activity of chromatin of pea roots treated by aluminum. *Plant Cell Physiol* **21**: 951-959

Matsuura-Endo C, Maeshima M and Yoshida S (1992) Subunit composition of vacuolar membrane H⁺-ATPase from mung bean. *Eur J Biochem* **187**: 745-51

McDonald-Stephens JL and Taylor GJ (1995) Kinetics of aluminum uptake by cell suspensions of *Phaseolus vulgaris* L. *J Plant Physiol* **145**: 327-334

Minella E and Sorrells ME (1992) Aluminum tolerance in barley: Genetic relationships among genotypes of diverse origin. *Crop Sci* **32**: 593-598

Moon DH, Ottoboni LMM, Souza AP, Sibov ST, Gaspar M and Arruda P (1997) Somaclonal-variation-induced aluminum sensitive mutant from and aluminum-inbred maize tolerant line. *Plant Cell Rep* **16**: 686-691

Morano KA and Klionsky DJ (1994) Differential effects of compartment deacidification on the targeting of membranes and soluble proteins to the vacuole in yeast. *J Cell Sci* **107**: 2813-2824

Morimura S, Takahashi E and Matsumoto H (1978) Association of aluminum with nuclei and inhibition of cell division in onion (*Allium cepa*) roots. *Z Pflanzenphysiol Bd* **88**: 395-401

- Nelson N** (1992) Evolution of organellar proton-ATPases. *Biochim Biophys Acta* **1100**: 109-124
- Nelson H, Mandiyan S and Nelson N** (1989) A conserved gene encoding the 57-kDa subunit of the yeast vacuolar H⁺-ATPase. *J Biol Chem* **264**: 1775-1778
- Nelson H, Mandiyan S and Nelson N** (1994) The *Saccharomyces cerevisiae* *VMA7* gene encodes a 14-kDa subunit of the vacuolar H⁺-ATPase catalytic sector. *J Biol Chem* **269**: 24150-24155
- Nichol BE, Oliveira LA, Glass ADM and Siddiqi MY** (1993) The effects of aluminum on the influx of calcium, potassium, ammonium, nitrate and phosphate in an aluminum-sensitive cultivar of barley (*Hordeum vulgare* L.). *Plant Physiol.* **101**: 1263-1266
- Osawa H and Matsumoto H** (2001) Possible involvement of protein phosphorylation in aluminum-responsive malate efflux from wheat root apex. *Plant Physiol* **126**: 411-420
- Peng SB** (1995) Nucleotide labeling and reconstitution of the recombinant 58-kDa subunit of the vacuolar proton-translocating ATPase. *J Biol Chem* **270**: 16926-16931
- Perzov N, Padler-Karavani V, Nelson H and Nelson N** (2001) Features of V-ATPases that distinguish them from F-ATPases. *FEBS Lett* **504**: 223-228
- Pineros MA and Kochian LV** (2001) A patch-clamp study on the physiology of aluminum toxicity and aluminum tolerance in maize. Identification and characterization of Al³⁺-induced anion channels. *Plant Physiol* **125**: 292-305

- Pineros M and Tester M** (1995) Characterization of a voltage-dependent Ca^{2+} -selective channel from wheat roots. *Planta* **195**: 478-488
- Plieth C, Sattelmacher B, Hansen U-P and Knight MR** (1999) Low-pH-mediated elevations in cytosolic calcium are inhibited by aluminium: A potential mechanism for aluminium toxicity. *Plant J* **18**: 643-650
- Powell B, Graham LA and Stevens TH** (2000) Molecular characterization of the yeast vacuolar H^{+} -ATPase proton pore. *J Biol Chem* **275**: 23654-23660
- Putterill JJ and Gardner RC** (1988) Proteins with the potential to protect plants from Al^{3+} toxicity. *Biochim Biophys Acta* **964**: 137-145
- Rengel Z** (1992) Disturbance of cell Ca^{2+} homeostasis as a primary trigger of Al toxicity syndrome. *Plant Cell Environ* **15**: 931-938
- Richards KD, Schott EJ, Sharma YK, Davis KR and Gardner RC** (1998) Aluminum induces oxidative stress genes in *Arabidopsis thaliana*. *Plant Physiol* **116**: 409-418
- Robinson DG, Haschke H-P, Hinz G, Hoh B, Maeshima M and Marty F** (1996) Immunological detection of tonoplast polypeptides in the plasma membranes of pea cotyledons. *Planta* **198**: 95-103
- Ryan PR, Reid RJ and Smith FA** (1997) Direct evaluation of the Ca^{2+} -displacement hypothesis for Al toxicity. *Plant Physiol* **113**: 1351-1357
- Sibov ST, Gaspar M, Silva MJ, Ottoboni LMM, Arruda P and Souza AP** (1999) Two genes control aluminum tolerance in maize: Genetic and molecular mapping analyses. *Genome* **42**: 475-482

- Siegel N and Haug A** (1983) Calmodulin-dependent formation of membrane potential in barley root plasma membrane vesicles: A biochemical model of aluminum toxicity in plants. *Physiol Plant* **59**: 285-291
- Slaski JJ** (1989) Effect of aluminum on calmodulin-dependent and calmodulin-independent NAD kinase activity in wheat (*Triticum aestivum* L.) root tips. *J Plant Physiol* **133**: 696-701
- Slaski JJ** (1996) Aluminum resistance in wheat (*Triticum aestivum*) is associated with rapid, Al-induced changes in activities of glucose-6-phosphate dehydrogenase and 6-phosphoglucanate dehydrogenase in root apices. *Physiol Plant* **98**: 477-484
- Snowden KC and Gardner RC** (1993) Five genes induced by aluminum in wheat (*Triticum aestivum* L.) roots. *Plant Physiol* **103**: 855-861
- Stevens TH and Forgac M** (1997) Structure, function and regulation of the vacuolar H⁺-ATPase. *Annu Rev Cell Dev Biol* **13**: 779-808
- Suhayda CG and Haug A** (1985) Citrate chelation as a potential mechanism against aluminum toxicity in cells: The role of calmodulin. *Can J Biochem Cell Biol* **63**: 1167-1175
- Supekova L, Supek F and Nelson N** (1995) The *Saccharomyces cerevisiae* *VMA10* is an intron-containing gene encoding a novel 13-kDa subunit of vacuolar H⁺-ATPase. *J Biol Chem* **270**: 13726-13732
- Sze H, Li X and Palmgren MG** (1999) Energization of plant cell membranes by H⁺-pumping ATPases: Regulation and biosynthesis. *Plant Cell* **11**: 677-689

- Taylor GJ and Foy CD (1985a)** Mechanisms of aluminum tolerance in *Triticum aestivum* L. (wheat). I. Differential pH induced by winter cultivars in nutrient solutions. *Amer J Bot* **72**: 695-701
- Taylor GJ and Foy CD (1985b)** Mechanisms of aluminum tolerance in *Triticum aestivum* L. (wheat). II. Differential pH induced by spring cultivars in nutrient solutions. *Amer J Bot* **72**: 702-706
- Taylor GJ and Foy CD (1985c)** Mechanisms of aluminum tolerance in *Triticum aestivum* L. (wheat). IV. The role of ammonium and nitrate nutrition. *Can J Bot* **63**: 2181-2186
- Taylor GJ (1987)** Mechanisms of aluminum tolerance in *Triticum aestivum* L. (wheat). V. Nitrogen nutrition, plant-induced pH and tolerance to aluminum: correlation without causality? *Can J Bot* **66**: 694-699
- Taylor GJ (1988)** The physiology of aluminum phytotoxicity. *In* H Sigel, A Sigel eds, *Metal Ions in Biological Systems*, Vol 24. Marcel Dekker, New York, pp 123-163
- Taylor GJ (1991)** Current views of the aluminum stress response; the physiological basis of resistance. *In* D.D. Randall, D.G. Blevins, C.D. Miles, eds. *Current topics in plant biochemistry and physiology 1991. Ultraviolet-B radiation stress, aluminum stress, toxicity and resistance, boron requirements, stress and toxicity*. Vol 10. *Interdisciplinary Plant Biochemistry and Physiology Program*, University of Missouri-Columbia. pp 57-93

- Taylor GJ and Foy CD (1985)** Mechanisms of aluminum resistance in *Triticum aestivum* L. (wheat) I: Differential pH induced by winter cultivars in nutrient solutions. *Amer J Bot* **72**: 695-701
- Taylor GJ, Basu A, Basu U, Slaski JJ, Zhang G and Good A (1997)** Aluminum-induced, 51 kD, membrane-bound proteins are associated with resistance to Al in a segregating population of wheat. *Plant Physiol* **114**: 363-372
- Taylor GJ, McDonald-Stephens JL, Hunter DB, Bertsch PM, Elmore D, Rengel Z and Reid RJ (2000)** Direct measurement of aluminum uptake and distribution in single cells of *Chara corallina*. *Plant Physiol* **123**: 987-996
- Tomashek JJ, Sonnenburg JL, Artimovich JM and Klionsky DJ (1996)** Resolution of subunit interactions and cytoplasmic subcomplexes of the yeast vacuolar proton-translocating ATPase. *J Biol Chem* **271**: 10397-10404
- Ungermann C, Sato K and Wickner W (1998)** Defining the functions of trans-SNARE pairs. *Nature* **396**: 543-548
- Von Euxkull HR and Mutert E (1995)** Global extent, development and economic impact of acid soils. *Plant Soil* **171**: 1-15
- Van Wambeke A (1976)** Formation, distribution and consequences of acid soils in agricultural development *In* MJ Wright, ed., *Plant Adaptation to Mineral Stress in Problem Soils*. Proceedings of workshop on plant adaptation to mineral stress in problem soils, 22-23 Nov, Beltsville, MD, USA. pp 15-30

- Wright RJ, Baligar VC and Murrman RP**, eds. (1995) Plant Soil Interactions at Low pH. Proceeding of the Second International Symposium on Plant-Soil Interactions at Low pH, 24-29 June, Beckley, West Virginia, USA.
- Wallace SU and Anderson IC** (1984) Aluminum toxicity and DNA synthesis in wheat roots. *Agronomy J* **76**: 5-8
- Yoshino M, Ito M, Haneda M, Tsubouchi R and Murakami K** (1999) Prooxidant action of aluminum ion--stimulation of iron-mediated lipid peroxidation by aluminum. *Biometals* **12**: 237-240
- Young JC, DeWitt ND and Sussman MR** (1998) A transgene encoding a plasma membrane H⁺-ATPase that confers acid resistance in *Arabidopsis thaliana* seedlings. *Genetics* **149**: 501-507
- Zhang G and Taylor GJ** (1989) Kinetics of aluminum uptake by excised roots of aluminum-tolerant and aluminum-sensitive cultivars of *Triticum aestivum* L. *Plant Physiol* **91**: 1094-1099
- Zhang G and Taylor GJ** (1991) Effects of biological inhibitors on kinetics of aluminum uptake by excised roots and purified cell wall material of aluminum-resistant and aluminum-sensitive cultivars of *Triticum aestivum* L. *J Plant Physiol* **138**: 533-539
- Zhang G, Hoddinott J and Taylor GJ** (1994) Characterization of 1,3-β-D-glucan (callose) synthesis in roots of *Triticum aestivum* in response to aluminum toxicity. *J Plant Physiol* **144**: 229-234

Zhang G, Archambault DJ, Slaski JJ and Taylor GJ (1995) Effects of protein synthesis inhibitors on uptake of aluminum in aluminum-resistant and aluminum-sensitive cultivars of wheat. *J Plant Physiol* **147**: 457-462

Zhang J, Vasilyeva E, Feng Y and Forgac M (1995) Inhibition and labeling of the coated vesicle V-ATPase by 2-azido-[³²P]ATP. *J Biol Chem* **270**: 15494-15500

Zhang W-H, Ryan PR and Tyerman SD (2001) Malate-permeable channels and cation channels activated by aluminum in the apical cells of wheat roots. *Plant Physiol* **125**: 1459-1472

2. Induction of Vacuolar ATPase and Mitochondrial ATP Synthase by Aluminum in an Aluminum-Resistant Cultivar of *Triticum aestivum* L.¹

2.1 Introduction

A wide range of proteins are induced by aluminum stress in wheat and other plant species. Aluminum-induced proteins include membrane-bound (Basu *et al.*, 1994a; Taylor *et al.*, 1997; Cruz-Ortega *et al.*, 1997), cytosolic (Richards *et al.*, 1998), cytoskeletal (Cruz-Ortega *et al.*, 1997) and exudate (Basu *et al.*, 1994b, 1997) proteins. Many of these have been implicated as general stress response proteins (Snowden *et al.*, 1995; Hamel *et al.*, 1998). Others have been associated with oxidative and other stresses (Richards *et al.*, 1998; Hamel *et al.*, 1998). The potential roles of others are still unclear. However, few have been functionally characterized as to their possible roles in Al tolerance.

The majority of Al-induced gene products have been identified at the RNA level. These include the *wali* (wheat aluminum-induced) and *war* (wheat aluminum regulated) gene products (Snowden and Gardner, 1993; Richards *et al.*, 1994; Hamel

¹ A version of this chapter has been published. Hamilton CA, Good AG and Taylor GJ (2001) Induction of Vacuolar ATPase and Mitochondrial ATP Synthase by Aluminum in an Aluminum-Resistant Cultivar of *Triticum aestivum* L. *Plant Physiol* **125**: 2068-2077

et al., 1998), β -1,3-glucanase (Cruz-Ortega *et al.*, 1997), glutathione S-transferase (Richards *et al.*, 1998), a fimbrin-like protein (Cruz-Ortega *et al.*, 1997) and a mitochondrial Mn-superoxide dismutase (Basu *et al.*, 2001) from wheat. In *Arabidopsis*, transcripts encoding aldolase, peroxidase, glutathione S-transferase, blue-copper-binding protein, superoxide dismutase and a reticuline:oxygen oxidoreductase homolog are all induced by Al (Richards *et al.*, 1998). These gene products appear to be involved in general stress response, since they are also induced by oxidative stress (Richards *et al.*, 1998), other toxic metals, low calcium levels and wounding (Snowden *et al.*, 1995, Hamel *et al.*, 1998). Recently, expression of genes encoding the *Arabidopsis* blue-copper-binding protein, and tobacco glutathione S-transferase, peroxidase and GDP-dissociation inhibitor have been shown to confer resistance to Al in transgenic *Arabidopsis* (Ezaki *et al.*, 2000). Expression of a wheat Mn-superoxide dismutase also conferred increased Al resistance in transgenic canola (Basu *et al.*, 2001).

Several gene products have been shown to be Al-induced by analysis of their activity. These include glucose-6-phosphate dehydrogenase, 6-phosphogluconate dehydrogenase (Slaski *et al.*, 1996) and vacuolar H⁺-ATPase (Kasai *et al.*, 1992, 1993). In contrast, the plasma membrane H⁺-ATPase (Matsumoto, 1988; Sasaki *et al.*, 1995; Widell *et al.*, 1994) is inhibited by Al. Glucan synthase II is inhibited by Al *in vitro* (Widell *et al.*, 1994), but synthesis of its end-product, callose, is stimulated by Al stress *in vivo* (Zhang *et al.*, 1994).

Another group of gene products have been shown to be Al-induced by comparing protein profiles of Al-sensitive and Al-resistant cultivars during Al stress (Ownby and Hruschka, 1991; Picton *et al.*, 1991; Delhaize *et al.*, 1991; Cruz-Ortega and Ownby, 1993; Somers *et al.*, 1996). Only three Al-induced proteins have been shown to co-segregate with the Al-resistance phenotype in wheat. One of these is a 23 kD root exudate protein (Basu *et al.*, 1997), while the other two are 51 kD, tonoplast associated proteins (RMP51), which are the focus of this study. These 51 kD proteins are specifically induced by Al in root tips of an Al-resistant cultivar of wheat (PT741). They are newly synthesized once Al stress begins, accumulate in a dose- and time-dependent manner, and then decline to control levels with the removal of Al stress (Basu *et al.*, 1994). The identity of these proteins remained to be determined.

The aim of the present study was to identify and characterize the RMP51 proteins in order to better understand their role in the Al stress response. These proteins have been purified from an endomembrane-enriched membrane fraction isolated from root tips of an Al-resistant cultivar of *Triticum aestivum* L. (cv PT741) grown in the presence of Al. Their identity has also been determined using peptide microsequence data. Purified peptides were homologous to the B subunit of the V-ATPase and to the α and β subunits of F_1F_0 -ATPase (mitochondrial ATP synthase). Both V-ATPase and F_1F_0 -ATPase activities were induced by Al treatment in PT741 and not in other cultivars tested. These results suggest that upregulation of ATPase activity in PT741 may be an adaptive response involved in Al resistance.

2.2 Methods

2.2.1 Plant Material and Growth Conditions

Seeds of Al-resistant (PT741, Atlas-66, Maringa) and Al-sensitive (Scout-66, Katepwa) cultivars of *Triticum aestivum* L. (wheat) were surface sterilized in 0.5% (v/v) sodium hypochlorite for 15-20 min and germinated overnight in a 0.005 g L⁻¹ solution of the antifungal agent Vitavax (Uniroyal Chemical Ltd, Calgary, AB, Canada) in double distilled water to limit fungal growth. Seedlings were grown for 5 d on nylon mesh floating over 15 L of an aerated mineral nutrient solution (pH 4.30) containing (μ M): 2900 NO₃, 300 NH₄, 100 PO₄, 800 K, 1000 Ca, 300 Mg, 101 SO₄, 34 Cl, 60 Na, 10 Fe, 6 B, 2 Mn, 0.15 Cu, 0.5 Zn, 0.1 Mo and 10 EDTA. For Al exposure, 5 d old seedlings were transferred to solutions containing (μ M): 1000 Ca, 300 Mg, 300 NH₄, 2900 NO₃ and 0-100 AlCl₃ at pH 4.30 (Taylor *et al.*, 1997). The seedlings were grown in a growth chamber (16 h light, 20°C, 68% RH and 8 h darkness, 16°C, 85% RH) for 2 d. Taylor *et al.* (1997) demonstrated that 100 μ M AlCl₃ is optimal for induction of the proteins of interest in cv. PT741. After 2 d of Al exposure, 1 cm root tips were harvested for isolation of endomembranes and subsequent purification of the 51 kD proteins (RMP51).

2.2.2 Isolation of Endomembrane-Enriched Membranes

All steps involved in endomembrane preparation were carried out at 4°C. Root tips (1 cm) were finely chopped and immediately homogenized in a Proctor-

Silex blender in homogenization buffer (0.25 M sucrose, 50 mM MOPS-Tris pH 7.5, 5 mM EDTA and 5 mM ascorbic acid), 1 mL per g root tissue. The homogenate was then filtered through miracloth (Calbiochem) and centrifuged at 20,000g for 15 min. The supernatant was collected and centrifuged at 100,000g for 1 h. The microsomal membrane pellet was resuspended in gradient buffer (0.25 M sorbitol, 5 mM HEPES-BTP, pH 7.0) and loaded onto a two-step gradient of 5 mL each of 2% and 10% (w/w) Dextran T-70 prepared in gradient buffer (Kasai *et al.*, 1992). The gradient was centrifuged at 70,000g for 2 h, and the interface containing endomembranes was collected, diluted with gradient buffer, and centrifuged at 120,000g for 1 h. The endomembrane-enriched membrane pellet was resuspended in 10 mM Tris-Acetate, pH 7.9, 10% glycerol and either frozen at -80°C for marker enzyme analysis, or used immediately for further purification.

2.2.3 Membrane Marker Assays

To ensure that the isolated membrane fraction was enriched for endomembranes, marker enzyme analysis was carried out according to Briskin *et al.* (1987). Activities of the NO₃⁻- and bafilomycin-sensitive ATPases, vanadate-sensitive ATPase and glucan synthase II (GSII), and cytochrome c oxidase were used as markers for tonoplast, plasma membrane and mitochondria, respectively. Adenosine triphosphatase activity was assayed in a reaction mixture containing 30 mM Tris-MES (pH 8.0), 3 mM MgSO₄, 0.2% Triton X-100 (v/v), 50 mM KCl, and 3 mM ATP-Tris, in the presence or absence of 250 μM Na₃VO₄ (P-Type ATPase activity), 1 mM NaN₃ (F₁F₀-Type ATPase activity), 100 nM bafilomycin A₁ or 50 mM KNO₃ (V-

Type ATPase activity). Phosphatase activity was measured as described by Ames (1966).

2.2.4 Purification of RMP51

Separation of the 51 kD proteins (RMP51) from other endomembrane-associated proteins was achieved by continuous elution electrophoresis (Miniprep Cell: Bio-Rad Laboratories). Total endomembrane protein (up to 500 µg per run), was prepared for electrophoresis by adding an equivalent volume of SDS-PAGE loading buffer (0.125 M Tris-HCl pH 6.8, 4% SDS, 20% glycerol, 10% β-mercaptoethanol, 0.002% bromophenol blue) and heating at 95°C for 5 min. Separating conditions for SDS-PAGE were as follows: 1.5 cm (4%) stacking gel, 10 cm (10%) separating gel. Electrophoresis/elution was carried out at 4°C at a constant current of 5 mA. Fractions (200 µL) were collected after the dye front had run off the gel and analyzed for the presence of RMP51 by SDS-PAGE (Mini Protean II: Bio-Rad Laboratories). Fractions containing RMP51 from each run were pooled and desalted using Sephadex G-50 gel filtration chromatography (Nick Column: Amersham Pharmacia Biotech). Aliquots of desalted samples were then analyzed for purity using two-dimensional electrophoresis (Mini Protean II 2D Cell: Bio-Rad Laboratories) according to the manufacturer's directions. After confirming their purity by two-dimensional electrophoresis, the desalted fractions from each run were pooled for microsequence analysis.

2.2.5 SDS-Polyacrylamide Gel Electrophoresis (SDS-PAGE)

Samples were prepared for SDS-PAGE by adding an equivalent volume of SDS-PAGE loading buffer and heating at 95°C for 5 min. Running conditions for SDS-PAGE (Laemmli, 1970) were: 4% stacking gel (at 12.5 mA) and 10% resolving gel (at 25 mA).

2.2.6 Two-Dimensional Electrophoresis

First dimension isoelectric focusing gels [4% acrylamide, 9.2 M urea, 2.0% CHAPS, 1.6% 5/7 ampholyte (BioRad Laboratories), and 0.4% 3/10 ampholyte (BioRad Laboratories)] were prefocused at 200V/10 min, 300V/15 min, and 400V/15 min. Microsomal membrane samples were extracted and precipitated as described by Hurkman and Tanaka (1986). Approximately 250 ng purified protein or 20 µg crude protein was then mixed with IEF sample buffer [9.5 M urea, 2.0% CHAPS, 5% β-mercaptoethanol, 1.6% 5/7 ampholyte (BioRad Laboratories), 0.4% 3/10 ampholyte (BioRad Laboratories)] and heated at 70°C for 20 min. After changing electrolytes and sample loading, isoelectric focusing was run for 500V/10 min and 750V/3.5h using the Mini Protean II 2D Cell (Bio-Rad Laboratories). Second dimension electrophoresis was carried out as described above.

2.2.7 Protein Quantitation and Visualization

Quantitation of protein present in membrane samples was performed using either the Bradford assay (Bradford, 1970) or comparing the band density of samples and known quantities of a standard protein (chicken egg ovalbumin: Sigma A5503)

of similar molecular weight (45 kD) on denaturing polyacrylamide gels. Proteins present in polyacrylamide gels were visualized using either a modified Morrissey silver stain procedure (Morrissey, 1981; Merril *et al.*, 1981) or staining with Coomassie Brilliant Blue R-250 (CBB R-250 Staining Kit, Bio-Rad Laboratories).

2.2.8 Immunoblotting

Microsomal proteins and RMP51 were separated by SDS-PAGE and electroblotted onto nitrocellulose (0.45 μ m, BioRad Laboratories) membranes using the Mini Trans-Blot Cell (Bio-Rad Laboratories) with a transfer buffer of 25 mM Tris, 192 mM glycine (pH 8.3) and 20% methanol. Transfer was performed at 100V at 4⁰C for 1h. Membranes were blocked overnight in TBST (20 mM Tris, pH 7.5, 140 mM NaCl, 0.1% v/v TWEEN-20) and 5% w/v skim milk powder. Membranes were incubated with primary antibodies diluted in TBST at 37⁰C for 1h, followed by three 15 min washes in PBST (80 mM Na₂HPO₄, 20 mM NaH₂PO₄, 100 mM NaCl, 0.1% TWEEN-20). Membranes labeled with primary antibody were incubated with horseradish peroxidase-conjugated secondary antibodies (Sigma A6154) diluted 1:15,000 in PBST at 4⁰C for 1h, followed by three 15 min washes. Chemiluminescent detection (Kirkegaard and Perry Laboratories, 54-61-00) was carried out according to the manufacturer's directions.

2.2.9 Peptide Microsequence Analysis

A 10 μ g sample of purified RMP51 was electrophoresed and excised from a denaturing SDS-polyacrylamide gel and sent to the Harvard Microchemistry Facility

(Cambridge, MA) for tryptic digestion, peptide separation and microsequence analysis. With technical advice from Harvard Microchemistry, the most abundant tryptic peptides were selected for microsequence analysis. Sequence comparison of the selected peptides was performed using the BLAST algorithm (Altschul *et al.*, 1997).

2.2.10 RNA Isolation, Northern Hybridization and Analysis

RNA was isolated from 1 cm root tips using the RNeasy Plant Mini Kit (Qiagen) according to the manufacturer's directions. Each 100 mg tissue sample yielded approximately 50 µg of RNA. RNA concentration was estimated by measuring A_{260} and checked by running aliquots on non-denaturing 1% agarose gels. Size Standards (Gibco-BRL RNA ladder) were included on all gels. Northern transfers were carried out using GeneScreen Plus (DuPont) membranes according to the manufacturer's directions. Probes for Northern blots were prepared from cDNA fragments isolated from pHTB1 (Berkelman *et al.*, 1994) and pAH (Chaumont *et al.*, 1988) plasmids (V-ATPase B subunit from *Hordeum vulgare* and F_1F_0 -ATPase α subunit from *Nicotiana plumbaginifolia*, respectively). Plasmids were digested with appropriate restriction enzymes (Amersham Pharmacia Biotech) and the released cDNA inserts were isolated from agarose gels using the QiaQuick gel extraction kit (Qiagen). Probes were then prepared by random priming (Sambrook *et al.*, 1989) using 100 ng of each DNA template. Unincorporated nucleotides (including [32 P]dCTP) were removed using Sephadex G-50 size exclusion chromatography (Nick Column: Amersham Pharmacia Biotech). Membranes were prehybridized at

42⁰C with 100 µg ml⁻¹ denatured, sheared, herring sperm DNA (Sigma) for 4h. Hybridization was carried out overnight at 42⁰C and washed twice with 2X SSPE (1X SSPE: 0.15M NaCl, 0.01 M NaH₂PO₄-H₂O, 0.001 M EDTA-Na₂, pH 7.4) for 15 min at room temperature, twice with 2X SSPE, 2% SDS for 45 min at 65⁰C, and twice with 0.1X SSPE for 15 min at room temperature. Membranes were exposed to Kodak X-OMAT x-ray film at -80⁰C. Shorter exposures of autoradiograms were performed to confirm that the observed results were not due to film saturation. Transcript levels were measured by densitometry of autoradiographs using an Alphaimager 2000 Documentation and Analysis System (Version 5.1 Alpha Innotech Corp). For Fig preparation, representative lanes of triplicate samples were selected from the same blot and compiled using Adobe Photoshop^R version 5.5.

2.3 Results

2.3.1 Purification of RMP51

The first step in the purification of RMP51 was preparation of endomembrane-enriched membranes from Al-treated seedlings of *Triticum aestivum* L. cv PT741. Membrane enrichment was confirmed using marker enzyme analysis (Briskin *et al.*, 1987). Positive markers for the tonoplast (Bafilomycin A₁- and nitrate-sensitive ATPase activities) were enriched by 1.43x and 3.35x, respectively, while negative markers (vanadate-sensitive ATPase, glucan synthase II and cytochrome c oxidase activities) were reduced to 0.45x, 0.69x and 0.35x (Table 2-1). Other endomembrane-associated proteins were then separated from RMP51 by

Table 2-1. Enrichment of marker enzymes in endomembrane-enriched membrane fractions isolated from root tips of *Triticum aestivum* L. cv PT741. Endomembrane-enriched membranes were isolated from microsomal membranes using a two-step gradient of Dextran T70 (2% and 10%), from which the interface was collected. Data are representative of 3 independent membrane preparations. Values shown are means +/- SE.

Marker Enzyme	Cellular Location	Enrichment in Endomembrane Fraction*	
		Mean	SE
Bafilomycin A ₁ -Sensitive ATPase	Tonoplast	1.43	0.24
Nitrate-Sensitive ATPase	Tonoplast	3.35	1.15
Vanadate-Sensitive ATPase	Plasma Membrane	0.45	0.19
Glucan Synthase II	Plasma Membrane	0.69	0.21
Cytochrome C Oxidase	Mitochondria	0.35	0.02

*Enrichment in endomembrane fraction is calculated as endomembrane specific activity/microsomal specific activity

continuous elution electrophoresis using the Miniprep Cell (BioRad). Separating conditions were optimized to purify the RMP51 band in a single step (Fig 2-1). The optimized procedure was repeated five times, with the RMP51 proteins isolated as a single band each time.

To determine whether or not both RMP51 proteins were present in the purified 51 kD band, aliquots of the pooled fractions from each purification run were analyzed by two-dimensional electrophoresis. Although two RMP51 spots were visible in the crude endomembrane fraction (Fig 2-2A), only one spot was visible in the purified sample (Fig 2-2B). This may have been caused by the presence of residual SDS (even after detergent removal and addition of nonionic CHAPS) or by loss of one of the RMP51 proteins during purification. Only one protein spot was observed from every preparation. Since the results from each purification run were identical, all of the fractions containing RMP51 were pooled. Total protein yield from five independent preparations was approximately 15 μ g, sufficient for subsequent analysis by tryptic digestion and peptide microsequencing.

2.3.2 Peptide Microsequence Analysis

Tryptic digestion and fragment purification by HPLC were carried out at the Harvard Microchemistry Facility (Cambridge, MA). Expert advice from Harvard helped to select several peptides which were abundant and well-separated from other

Fig 2-1. Fractionation of endomembrane-associated proteins by continuous elution electrophoresis. Total endomembrane proteins were isolated from root tips of *Triticum aestivum* L. cv PT741 after 48h treatment with 100 μ M AlCl₃. Selected fractions (lanes 2-8) were analyzed for the presence of RMP51 (arrow) by SDS-PAGE and silver staining. Molecular weights of protein standards are given in kilodaltons. Fractions pooled for further analysis are indicated by asterisks (*).







Fig 2-2. Two-dimensional gel analysis of crude endomembrane proteins (A) and purified RMP51 (B). Molecular weights of protein standards (kD) are indicated on the left and the pH scale is shown on the top. RMP51 is indicated by solid arrows. Data shown are representative of 5 independent trials.



peptides by HPLC. Four of these peptides were selected, sequenced and subsequently identified using the BLAST sequence algorithm (Altschul *et al.*, 1997). Peptide 1 (QIYPPINVLPSLSR) was identical to amino acids 365-378 of the V-ATPase B subunit from barley (*Hordeum vulgare*, accession Q40079). Peptide 2 (FVAQGAYDTR) also showed 100% identity to the barley V-ATPase B subunit (accession Q40079, amino acids 440-449). Peptide 3 (FTAQANSEVSALLGR) showed 100% identity to amino acids 336-349 of the F₁F₀-ATPase β subunit from wheat (accession P20858), but no significant homology to the V-ATPase B subunit. Peptide 4 (TGSIVDVPAGK) showed 100% identity to amino acids 93-103 of the F₁F₀-ATPase α subunit from wheat (accession P12862).

Results from the BLAST sequence comparison suggest that the four tryptic fragments were derived from the ATP-binding and catalytic subunits of related ATPases (V-ATPase and F₁F₀-ATPase). The possibility that they are all derived from a single protein was checked by aligning the four RMP51 peptide sequences with the B subunit of the V-ATPase from barley and the α and β subunits of the F₁F₀-ATPase from wheat. The 3 ATPase subunits share only approximately 25% identity, and the peptide sequences are not derived from regions with significant sequence homology between them. Therefore, it is unlikely that any single protein could contain all four peptides. The fact that more than one protein is present in the RMP51 band was confirmed by Western analysis (Fig 2-3). Antibodies raised against the V-ATPase B subunit from *Vigna radiata* and the F₁F₀-ATPase α and β subunits from

Fig 2-3. Identification of RMP51 by immunoblotting. RMP51 and crude microsomal proteins isolated from AI-treated root tips of *Triticum aestivum* L. cv PT741 were separated by SDS-PAGE, immunoblotted and probed with polyclonal antibodies specific to the V-ATPase B subunit (from *Vigna radiata*) and the F₁F₀-ATPase α and β subunits (from *Saccharomyces cerevisiae*). Antibody labeling was detected using horseradish peroxidase-conjugated secondary antibodies and chemiluminescent detection.




Saccharomyces cerevisiae all cross-react with RMP51, suggesting that all three proteins are present.

2.3.3 V-ATPase and F_1F_0 -ATPase are Al-induced

Since RMP51 protein levels increase in a dose-dependent manner with Al treatment (Taylor *et al.*, 1997), levels of V-ATPase and F_1F_0 -ATPase activity in Al-treated and control seedlings were compared. Activities were measured in total microsomal membrane fractions isolated from 5d-old seedlings of *Triticum aestivum* L. cv PT741 exposed to 0-150 μM AlCl_3 for 2d (Fig 2-4). V-ATPase and F_1F_0 -ATPase activities were both induced during Al stress, although to different extents. Vacuolar ATPase activity increased to 181% of control (from 0.127 to 0.232 $\text{nmol Pi mg}^{-1} \text{min}^{-1}$) as concentrations of AlCl_3 increased from 0-100 μM , followed by a decline to 125% of the control level (0.159 $\text{nmol Pi mg}^{-1} \text{min}^{-1}$) at 150 μM . Mitochondrial F_1F_0 -ATPase activity increased to 899% of the control level (from 0.025 to 0.223 $\text{nmol Pi mg}^{-1} \text{min}^{-1}$) as concentrations of AlCl_3 increased from 0-100 μM , followed by a decline to 742% of the control level (0.184 $\text{nmol Pi mg}^{-1} \text{min}^{-1}$) at 150 μM AlCl_3 . Plasma membrane ATPase activity decreased to 57% of the control level (from 0.136 to 0.077 $\text{nmol Pi mg}^{-1} \text{min}^{-1}$) at 25 μM AlCl_3 and to 18% of the control level (0.025 $\text{nmol Pi mg}^{-1} \text{min}^{-1}$) at 100 μM . Since both F_1F_0 -ATPase and V-ATPase activities were induced by Al in a dose-dependent manner, it remains possible that the Al-induced RMP51 band initially identified by Basu *et al.* (1994a) consisted of both vacuolar and mitochondrial components.

Fig 2-4. The effect of Al on activity of the V-ATPase, F_1F_0 -ATPase and P-ATPase. ATPase specific activity was measured in total microsomal membranes prepared from 1 cm root tips of the Al-resistant cultivar, PT741. Five-d-old seedlings were treated with different concentrations of Al (0, 25, 50, 75, 100, 150 μ M) for 48h. Values are means \pm SE of three biological replicates and are representative of three independent trials.




In order to determine whether or not there is a transcriptional component to the induction of these ATPases, Northern analysis was performed on RNA isolated from the same tissues used for ATPase assays (Fig 2-5). Levels of the transcript encoding the B subunit of the V-ATPase showed an increase to 1.9x control from 0-100 μM AlCl_3 . This increase, although relatively small, was consistent in 3 independent experiments, and showed a pattern similar to that observed for V-ATPase activity (Fig 2-4). This suggests that induction of V-ATPase activity by Al may be transcriptionally mediated, although participation of translational and post-translational mechanisms cannot be excluded. In contrast, levels of the transcript encoding the α subunit of the F_1F_0 -ATPase remained constant over the entire range of Al concentrations tested. Since F_1F_0 -ATPase activity levels increased 899% over this range, it appears that this increase may be due to a translational or post-translational mechanism.

2.3.4 *Cultivar Screen for ATPase Induction*

If induction of V-ATPase and F_1F_0 -ATPase activities are general responses to Al stress, then induction should be observed in Al-resistant and Al-sensitive cultivars of wheat under stress conditions. To test this hypothesis, V-ATPase, F_1F_0 -ATPase and P-ATPase activities were measured in 3 Al-resistant and 2 Al-sensitive cultivars of wheat in control (0 μM AlCl_3) conditions, conditions stressful to sensitive cultivars (20 μM AlCl_3), and conditions stressful to resistant cultivars (100 μM AlCl_3) (Taylor

Fig 2-5. The effect of AI on transcript levels of V-ATPase and F₁F₀-ATPase subunits. RNA was isolated from the same tissues used for ATPase activity measurements in Fig 4 and Northern blotted. (A) Autoradiographs of Northern blots probed with ³²-P labeled cDNAs encoding the V-ATPase B subunit from *Hordeum vulgare* and the F₁F₀-ATPase α subunit from *Nicotiana plumbaginifolia*. To ensure equal RNA loading and transfer, membranes were also probed with a cDNA clone encoding the 25s rRNA from *Glycine max*. Results shown are representative of three independent trials. (B) Quantification of transcript levels shown in (A). Transcript levels are expressed as the density of each band relative to 25S rRNA on the autoradiograph. Values are means +/- SE of three independent replicates.



B)





Fig 2-6. The effect of Al on ATPase specific activity in total microsomal membranes prepared from 1 cm root tips of PT741, Atlas-66 and Maringa (Al-resistant), and from Scout-66 and Katepwa (Al-sensitive). Five-d-old seedlings of each cultivar were exposed to 0, 20 or 100 μ M Al for 48h. Values represent means \pm SE of three independent replicates.



et al., 1997). In cultivar PT741, V-ATPase activity increased by 315% (from 0.113 to 0.356 nmol Pi mg⁻¹ min⁻¹), even at 20 μM AlCl₃ (Fig 2-6A). Vacuolar ATPase activities in Atlas-66 and Maringa (Al-resistant) were unaffected at 20 μM AlCl₃, but reduced to 30% and 33% of control (from 0.174 to 0.050 nmol Pi mg⁻¹ min⁻¹ and from 0.097 to 0.032 nmol Pi mg⁻¹ min⁻¹, respectively) at 100 μM AlCl₃. In Scout-66 and Katepwa (Al-sensitive), a similar pattern was observed, with reductions to 62% and 54% of control (from 0.129 to 0.080 nmol Pi mg⁻¹ min⁻¹ and from 0.097 to 0.032 nmol Pi mg⁻¹ min⁻¹, respectively) observed at 100 μM AlCl₃.

Cultivar PT741 also showed induction of F₁F₀-ATPase activity in this experiment (Fig 2-6B). Control levels of F₁F₀-ATPase activity were slightly greater in PT741 than in the other cultivars tested, and activity increased to 180% of control (from 0.220 to 0.390 nmol Pi mg⁻¹ min⁻¹) at 20 μM AlCl₃ and to 214% of control (0.469 nmol Pi mg⁻¹ min⁻¹) at 100 μM AlCl₃. In Atlas-66, F₁F₀-ATPase activity increased to 142% of control (from 0.125 to 0.178 nmol Pi mg⁻¹ min⁻¹) at 20 μM AlCl₃, but activity declined to control levels at 100 μM AlCl₃. All other cultivars tested showed reduced F₁F₀-ATPase activities (Maringa, Scout-66 and Katepwa reduced to 59%, 40% and 47% of control, respectively) at 20 μM AlCl₃. In contrast to the above observations, decreased activity of the plasma membrane ATPase was a general effect of Al treatment in all cultivars tested (Fig 2-6C). Activities decreased to near zero in all cultivars, even at 20 μM AlCl₃. In summary, V-ATPase and F₁F₀-

ATPase activities were specifically induced in PT741, while P-ATPase activity declined in all cultivars tested.

2.4 Discussion

Differential screening of mRNA and protein profiles between Al-treated and untreated tissues or between Al-resistant and Al-sensitive wheat cultivars has revealed that expression of many genes is induced during Al stress (Snowden and Gardner, 1993; Snowden *et al.*, 1995; Richards *et al.*, 1994; Cruz-Ortega *et al.*, 1997; Hamel *et al.*, 1998; Basu *et al.*, 2001). However, only three Al-induced proteins (Basu *et al.*, 1994b; Taylor *et al.*, 1997) have been shown to co-segregate with the Al-resistance phenotype. In order to better understand the role of these proteins, their identities must be determined.

Aluminum stress causes a general decline in protein synthesis (^{35}S -methionine incorporation) in Al-sensitive cultivars of *Triticum aestivum*, while there is little to no effect in Al-resistant cultivars (Ownby and Hruschka, 1991; Rincon and Gonzales, 1991; Basu *et al.*, 1994a). Similar results have been observed in *Medicago sativa* (Campbell *et al.*, 1994). Since pretreatment of Al-resistant cultivars of wheat (Atlas-66, Grana) improves the resistance of plants to subsequent exposures, Aniol (1984) hypothesized that induction of new protein synthesis was involved in Al resistance. This hypothesis was supported by the subsequent observation that Al pretreatment induced Al tolerance in an Al-resistant cultivar of *Phaseolus vulgaris* L. (Cumming *et*

al., 1992). Basu *et al.* (1994a) previously reported that two tonoplast-associated proteins are induced by Al stress in root tips of an Al-resistant cultivar of *Triticum aestivum*, PT741. These proteins segregate with the Al-resistance phenotype in a cross between PT741 and Katepwa (Taylor *et al.*, 1997). In this work, I used continuous elution electrophoresis to purify the 51 kD band containing these proteins from Al-treated seedlings of *Triticum aestivum* L. cv PT741 (Fig 2-1). Sequence analysis of purified peptides was then used to identify RMP51. Of four peptides sequenced, two were identical to fragments of the V-ATPase B subunit from barley, one was identical to a fragment of the F₁F₀-ATPase α subunit from wheat and one was identical to a fragment of the F₁F₀-ATPase β subunit from wheat. Although RMP51 was originally identified as two proteins, it was nonetheless surprising that both mitochondrial and vacuolar proteins were present in the 51 kD band, which had been purified from an endomembrane-enriched fraction (Table 2-1). However, antibodies specific to the B subunit of the V-ATPase and the α and β subunits of the F₁F₀-ATPase all cross-reacted with the RMP51 band (Fig 2-3), suggesting that there were in fact at least three proteins present in the purified band.

To determine whether the V-ATPase and the F₁F₀-ATPase are Al-induced, ATPase activities were measured at various Al concentrations. Basu *et al.* (1994a) showed previously that RMP51 protein levels increased in an Al dose-dependent manner. If RMP51 is in fact a subunit of the V-ATPase or F₁F₀-ATPase, then activities of these enzymes should show a similar response to Al, assuming that all of the subunits are similarly induced. My data showed that V-ATPase and F₁F₀-ATPase

activities increased to 183% and 899% of control (respectively) with Al treatment, while P-ATPase activity decreased to 18% of control levels (Fig 2-4). Induction of V-ATPase (and H⁺-PP_iase) by Al has been previously demonstrated in tonoplast-enriched membrane vesicles from *Hordeum vulgare* roots by Kasai *et al.* (1992, 1993). These authors observed a 40-53% increase in V-ATPase activity upon Al exposure, depending upon the external Ca²⁺ concentration. These results were confirmed by Zhang *et al.* (1998). Inhibition of P-ATPase activity has also been previously reported. Matsumoto (1988) observed a 50% decrease in P-ATPase activity from *Hordeum vulgare* roots treated with 100 μM AlCl₃ at pH 6.5, and a 45% decrease after treatment with 1 mM AlCl₃ at pH 5.5 (Matsumoto *et al.*, 1992). In subsequent experiments, Sasaki *et al.* (1995) showed that P-ATPase activity in *Triticum aestivum* declined by 13-19% after treatment with 50 μM AlCl₃ at pH 4.5. Widell *et al.* (1994) observed a similar effect in *Picea abies* and *Triticum aestivum*.

The effect of Al treatment on F₁F₀-ATPase activity has not previously been investigated. I was initially surprised by the induction of F₁F₀-ATPase activity, since localization experiments showed no enrichment of the RMP51 band in mitochondrial fractions (Taylor *et al.*, 1997). However, it is possible that protein levels of the mitochondrial F₁F₀-ATPase subunits remain constant during Al stress, and that F₁F₀-ATPase activity is being modulated at the post-translational level (Stevens and Forgac, 1997). This is consistent with our observation that steady-state transcript levels remained constant while activity increased by 7.3x (Figs 2-4, 2-5). Since Al induces F₁F₀-ATPase activity, increased ATP synthase activity may be required to

support V-ATPase induction and other energy-dependent processes involved in Al resistance. Although ATP synthase activity was not directly measured, the F_1F_0 -ATPase acts almost exclusively as an ATP synthase under physiological conditions. Therefore, the increased F_1F_0 -ATPase activity observed here has been interpreted as increased ATP synthase activity *in vivo*.

If induction of V-ATPase and F_1F_0 -ATPase activities is a general effect of Al stress, then a similar response should be observed in all cultivars, whether Al-resistant or Al-sensitive. If induction is involved in mediating a general Al-resistance mechanism, induction should occur only in Al-resistant cultivars. Alternatively, induction may be a cultivar-specific resistance mechanism in PT741. To differentiate among these hypotheses, ATPase activities were measured in two additional Al-resistant cultivars (Atlas-66, Maringa) and two Al-sensitive cultivars (Scout-66, Katepwa). Our data show that Al-induction of V-ATPase and F_1F_0 -ATPase is unique to PT741 (Fig 2-6). In all other cultivars, V-ATPase and F_1F_0 -ATPase activities remained constant or declined with Al treatment.

It is interesting that this phenomenon was observed only in PT741 and not in the other two Al-resistant cultivars tested, neither of which showed increased levels of RMP51 protein (data not shown). The lack of an effect in Maringa is perhaps not surprising since research on this cultivar has correlated exudation of a 23 kD, Al-binding polypeptide with Al resistance, and Al-resistance is controlled by a single dominant gene (Basu *et al.*, 1997). In Atlas 66, Al resistance has also been correlated with enhanced exudation of malate during Al stress (Basu *et al.*, 1994c).

Several observations suggest that induction of V-ATPase and F_1F_0 -ATPase could be involved in mediating Al resistance in *Triticum aestivum* L. cv PT741. First, RMP51 protein (consisting of the V-ATPase B subunit and the F_1F_0 -ATPase α and β subunits) is newly synthesized upon Al treatment and accumulates in an Al dose-dependent manner (Basu *et al.*, 1994a). Second, accumulation of RMP51 segregates with the Al-resistance phenotype (Taylor *et al.*, 1997). Third, V-ATPase and F_1F_0 -ATPase activities increase in an Al dose-dependent manner only in the Al-resistant cultivar, PT741.

Induction of V-ATPase activity has been linked to salt tolerance in several species (Ballesteros *et al.*, 1996; Kirsch *et al.*, 1996). Induction of V-ATPase activity is thought to be a homeostatic mechanism required to provide energy for Na^+/H^+ antiport, which delivers Na^+ to the vacuole (Matsumoto and Chung, 1988; Reuveni *et al.*, 1990; Nakamura *et al.*, 1992; Colombo and Cerana, 1993). Kasai *et al.* (1992) suggested that a similar mechanism could be responsible for Al resistance in wheat, with an Al^{3+}/H^+ antiport system driving sequestration of Al in the vacuole. This suggestion should be taken with caution, however, since there is no direct evidence for an Al^{3+}/H^+ co-transporter. Increased V-ATPase activity could also be required as a homeostatic mechanism to maintain the cytoplasmic pH near neutrality. I have shown that Al exposure decreases plasma membrane ATPase activity, in agreement with results reported previously (Matsumoto, 1988; Matsumoto *et al.*, 1992; Sasaki *et al.*, 1995; Widell *et al.*, 1994). This could cause a decrease in cytoplasmic pH, with adverse physiological effects. Increased activity of V-ATPase, with energy balance

maintained by increased ATP synthase activity, could counteract these changes by transporting protons into the vacuole. A similar phenomenon has been observed in *Lactobacillus acidophilus*, which expresses a pH-inducible F_1F_0 -ATPase to extrude protons and maintain cytoplasmic pH (Kullen and Klaenhammer, 1999).

In summary, I have purified the Al-induced RMP51 proteins from the Al-resistant cultivar of wheat, PT741 and shown that the RMP51 band consists of the B subunit of the V-ATPase and the α and β subunits of the F_1F_0 -ATPase. Vacuolar ATPase activity increased to 183-324% of control, while levels of the transcript encoding the B subunit increased by 2.2x, from 0-100 μ M $AlCl_3$. Mitochondrial F_1F_0 -ATPase activity increased by 214-899%, with a constant level of the transcript encoding the α subunit, over the same range of Al concentrations. Increased V-ATPase and F_1F_0 -ATPase activities are observed only in PT741, suggesting that induction of V-ATPase and F_1F_0 -ATPase activities is not simply symptomatic of stress. These results allow me to put forward the hypothesis that induction of the V-ATPase and the F_1F_0 -ATPase plays a role in Al resistance in PT741. Moreover, the demonstration that tolerance to Al in PT741 is mediated by a different mechanism than is observed in other varieties of wheat (e.g. Maringa) suggests the possibility of combining these traits genetically to produce a cultivar with enhanced Al tolerance.

2.5 Literature Cited

- Altschul SF, Madden TL, Schäffer AA, Zhang J, Zhang Z, Miller W and Lipman DJ (1997)** Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res.* **25**:3389-3402
- Ames BN (1966)** Assay of inorganic phosphate, total phosphate and phosphatase. *Methods Enzymol* **8**: 115-118
- Aniol A (1984)** Induction of aluminum tolerance in wheat seedlings by low doses of aluminum in the nutrient solution. *Plant Physiol* **75**: 551-555
- Ballesteros E, Donaire JP and Belver A (1996)** Effects of salt stress on H⁺-ATPase and H⁺-PP_iase activities of tonoplast-enriched vesicles isolated from sunflower roots. *Physiol Plant* **97**: 259-268
- Basu A, Basu U and Taylor GJ (1994a)** Induction of microsomal membrane proteins in roots of an aluminum-resistant cultivar of *Triticum aestivum* L. under conditions of aluminum stress. *Plant Physiol* **104**: 1007-1013
- Basu U, Basu A and Taylor GJ (1994b)** Differential exudation of polypeptides by roots of aluminum-resistant and aluminum-sensitive cultivars of *Triticum aestivum* L. in response to aluminum stress. *Plant Physiol* **106**: 151-158
- Basu U, Godbold D and Taylor GJ (1994c)** Aluminum resistance in *Triticum aestivum* associated with enhanced exudation of malate. *J Plant Physiol* **144**: 747-753

- Basu U, Good AG and Taylor GJ (2001)** Transgenic *Brassica napus* plants overexpressing aluminium-induced mitochondrial manganese superoxide dismutase cDNA are resistant to aluminium. *Plant Cell Environ* (In Press)
- Basu U, McDonald-Stephens JL, Archambault DJ, Good AG, Briggs KG, Taing-Aung and Taylor GJ (1997)** Genetic and physiological analysis of doubled-haploid, aluminum resistant lines of wheat provide evidence for the involvement of a 23 kD, root exudate polypeptide in mediating resistance *Plant Soil* **196**(2): 283-288
- Berkelman T, Houtchens KA and Dupont FM (1994)** Two cDNA clones encoding isoforms of the B subunit of the vacuolar ATPase from barley roots. *Plant Physiol* **104**: 287-288
- Bradford M (1976)** A rapid and sensitive method for the quantification of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* **72**: 248-254
- Briskin DP, Leonard RT and Hodges TK (1987)** Isolation of the plasma membrane: Membrane markers and general principals. *In* L Packer, R Douce, eds, *Methods in Enzymology*. Vol 148. Academic Press, pp 542-558
- Campbell TA, Jackson PR and Xia ZL (1994)** Effects of aluminum stress on alfalfa root proteins. *J Plant Nutr* **17**(2/3): 461-471
- Chaumont F, Boutry M, Briquet M and Vassarotti A (1988)** Sequence of the gene encoding the mitochondrial F₁-ATPase alpha subunit from *Nicotiana plumbaginifolia*. *NAR* **16**: 6247

- Colombo R and Cerana R** (1993) Enhanced activity of tonoplast pyrophosphatase in NaCl-grown cells of *Daucus carota*. *J Plant Physiol* **142**: 226-229
- Cruz-Ortega R and Ownby JD** (1993) A protein similar to PR (pathogenesis-related) proteins is elicited by metal toxicity in wheat roots. *Physiol Plant* **89**: 211-219
- Cruz-Ortega R, Cushman JC and Ownby JD** (1997) cDNA clones encoding 1,3-beta-glucanase and a fimbrin-like cytoskeletal protein are induced by Al toxicity in wheat roots. *Plant Physiol* **114**(4): 1453-1460
- Cumming JR, Buckelew Cumming A and Taylor GJ** (1992) Patterns of root respiration associated with the induction of aluminum tolerance in *Phaseolus vulgaris* L. *J Exp Bot* **43**(253): 1075-1081
- Delhaize E, Higgins TJV and Randall PJ** (1991) Aluminum tolerance in wheat: analysis of polypeptides in the root apices of tolerant and sensitive genotypes. *In* RJ Wright, VC Baligar, RP Murrman, eds. *Plant-Soil Interactions at Low pH. Proceedings of the Second International Symposium on Plant-Soil Interactions at Low pH, 24-29 June 1990, Beckley, West Virginia, USA.* Kluwer Academic Publishers, Dordrecht, pp 1071-1079
- Delhaize E, Ryan PR and Randall PJ** (1993) Aluminum tolerance in wheat (*Triticum aestivum* L.). 2. Aluminum-stimulated excretion of malic acid from root apices. *Plant Physiol* **103**: 695-702

- Ezaki B, Gardner RC, Ezaki Y and Matsumoto H (2000)** Expression of aluminum-induced genes in transgenic Arabidopsis plants can ameliorate aluminum stress and/or oxidative stress. *Plant Physiol* **122**: 657-665
- Foy CD, Chaney RL and White MC (1978)** The physiology of metal toxicity in plants. *Annu Rev Plant Physiol* **29**: 511-566
- Hamel F, Breton C and Houde M (1998)** Isolation and characterization of wheat aluminum-regulated genes: possible involvement of aluminum as a pathogenesis response regulator. *Planta* **205**(4): 531-538
- Haug A (1984)** Molecular aspects of aluminum toxicity. *Crit Rev Plant Sci* **1**: 345-373
- Hurkman WJ and Tanaka CK (1986)** Solubilization of plant membrane proteins for analysis by two-dimensional gel electrophoresis. *Plant Physiol* **81**: 802-806
- Kasai M, Sasaki M, Tanakamaru S, Yamamoto Y and Matsumoto H (1993)** Possible involvement of abscisic acid in increases in activities of two vacuolar H⁺-pumps in barley roots under aluminum stress. *Plant Cell Physiol* **34**: 1335-1338
- Kasai M, Sasaki M, Yamamoto Y and Matsumoto H (1992)** Aluminum stress increases K⁺ efflux and activities of ATP- and PPi-dependent H⁺ pumps of tonoplast-enriched membrane vesicles from barley roots. *Plant Cell Physiol* **33**: 1035-1039

Kirsch M, Zhigang A, Viereck R, Low R and Rausch T (1996) Salt stress induces an increased expression of V-type H⁺-ATPase in mature sugar beet leaves.

Plant Mol Biol **32**: 543-547

Koyama H, Takit, E, Kawamura A, Hara T and Shibata D (1999) Over expression of mitochondrial citrate synthase gene improves the growth of carrot cells in Al-phosphate medium. Plant Cell Physiol **40**: 482-488

Kullen MJ and Klaenhammer TR (1999) Identification of the pH-inducible, proton-translocating F₁F₀-ATPase (*atpBEFHAGDC*) operon of *Lactobacillus acidophilus* by differential display: gene structure, cloning and characterization. Mol Microbiol **33**: 1152-1161

Laemmli UK (1970) Cleavage of the structural proteins during the assembly of the head of bacteriophage T4. Nature **227**: 680-685

Matsumoto H (1988) Inhibition of proton transport activity of microsomal membrane vesicles of barley roots by aluminium. Soil Sci Plant Nutr **34**: 499-526

Matsumoto H and Chung GC (1988) Increase in proton transport activity of tonoplast vesicles as an adaptive response of barley roots to NaCl stress. Plant Cell Physiol **29**: 1133-1140

Matsumoto H, Yamamoto Y and Kasai M (1992) Changes of some properties of the plasma membrane-enriched fraction of barley roots related to aluminum stress: membrane-associated ATPase, aluminum and calcium. Soil Sci Plant Nutr **38**: 411-419

Merril CR, Goldman D, Sedman SA and Ebert MH (1981) Ultrasensitive stain for proteins in polyacrylamide gels shows regional variation in cerebrospinal fluid proteins. *Science* **211**: 1437-1438

Miyasaka SC, Buta JG, Howell RK and Foy CD (1991) Mechanism of aluminum tolerance in snapbeans. Root exudation of citric acid. *Plant Physiol* **96**: 737-743

Morrissey JH (1981) Silver stain for proteins in polyacrylamide gels: a modified procedure with enhanced uniform sensitivity. *Anal Biochem* **117**: 307-310

Nakamura Y, Kasamo K, Shimosato N, Sakata M and Ohta E (1992) Stimulation of the extrusion of protons and H⁺-ATPase activities with the decline of pyrophosphatase activity of the tonoplast in intact mung bean roots under high NaCl stress and its relation to external levels of Ca²⁺ ions. *Plant Cell Physiol* **33**: 139-149

Ownby JD and Hruschka WR (1991) Quantitative changes in cytoplasmic and microsomal proteins associated with aluminum toxicity in two cultivars of winter wheat. *Plant Cell Environ* **14**: 303-309

Picton SJ, Richards KD and Gardner RC (1991) Protein profiles in root-tips of two wheat (*Triticum aestivum* L.) cultivars with differential tolerance to aluminium. In RJ Wright, VC Baligar, RP Murrmann, eds, *Plant-Soil Interactions at Low pH. Proceedings of the Second International Symposium on Plant-Soil Interactions at Low pH, 24-29 June 1990, Beckley, West Virginia, USA*. Kluwer Academic Publishers, Dordrecht, pp 1063-1070

Reuveni M, Bennett AB, Bressan RA and Hasegawa PM (1990) Enhanced H⁺ transport capacity and ATP hydrolysis activity of the tonoplast H⁺-ATPase after NaCl adaptation. *Plant Physiol* **94**: 524-530

Richards KD, Snowden KC and Gardner RC (1994) *Wali6* and *wali7*. Genes induced by aluminum in wheat (*Triticum aestivum* L.) roots. *Plant Physiol* **105**: 1455-1456

Richards KD, Schott EJ, Sharma YK, Davis KR and Gardner RC (1998) Aluminum induces oxidative stress genes in *Arabidopsis thaliana*. *Plant Physiol* **116**: 409-418

Rincon M and Gonzales RA (1991) Induction of protein synthesis by aluminium in wheat (*Triticum aestivum* L.) root tips. In RJ Wright, VC Baligar, RP Murrmann, eds, *Plant-Soil Interactions at Low pH. Proceedings of the Second International Symposium on Plant-Soil Interactions at Low pH, 24-29 June 1990, Beckley, West Virginia, USA. Kluwer Academic Publishers, Dordrecht.* pp 851-858

Sambrook J, Fritsch T and Maniatis T (1989) *Molecular Cloning: A Laboratory Manual*. 2nd Ed. Cold Spring Harbor Laboratory Press

Sasaki M, Kasai M, Yamamoto Y and Matsumoto H (1995) Involvement of plasma membrane potential in the tolerance mechanism of plant roots to aluminum toxicity. In RA Date *et al.*, eds. *Plant-Soil Interactions at Low pH. Proceedings of the Second International Symposium on Plant-Soil Interactions*

at Low pH, 24-29 June 1990, Beckley, West Virginia, USA. Kluwer Academic Publishers, Dordrecht, pp 285-290

Slaski JJ (1996) Aluminum resistance in wheat (*Triticum aestivum*) is associated with rapid, Al-induced changes in activities of glucose-6-phosphate dehydrogenase and 6-phosphoglucanate dehydrogenase in root apices. *Physiol Plant* **98**: 477-484

Snowden KC and Gardner RC (1993) Five genes induced by aluminum in wheat (*Triticum aestivum* L.) roots. *Plant Physiol* **103**: 855-861

Snowden KC, Richards KD and Gardner RC (1995) Aluminum-induced genes. Induction by toxic metals, low calcium, and wounding and pattern of expression in root tips. *Plant Physiol* **107**: 341-348

Somers DJ, Briggs KG and Gustafson JP (1996) Aluminum stress and protein synthesis in near isogenic lines of *Triticum aestivum* differing in aluminum tolerance. *Physiol Plant* **97**: 694-700

Stevens TH and Forgac M (1997) Structure, function and regulation of the vacuolar (H⁺)-ATPase. *Annu Rev Cell Dev Biol* **13**: 779-808

Takita E, Koyama H and Hara T (1999a) Organic acid metabolism in aluminum-phosphate utilizing cells of carrot (*Daucus carota* L.). *Plant Cell Physiol* **40**: 489-495

Takita E, Koyama H, Shirano Y, Shibata D and Hara T (1999b) Structure and expression of the mitochondrial citrate synthase gene in carrot cells utilizing Al-phosphate. *Soil Sci Plant Nutr* **45**: 197-205

Taylor GJ, Basu A, Basu U, Slaski JJ, Zhang G and Good A (1997) Al-induced, 51 kilodalton, membrane-bound proteins are associated with resistance to Al in a segregating population of wheat. *Plant Physiol* **114**: 363-372

Widell S, Asp H and Jensen P (1994) Activities of plasma membrane-bound enzymes isolated from roots of spruce (*Picea abies*) grown in the presence of aluminium. *Physiol Plant* **92**: 459-466

Zhang G, Hoddinott J and Taylor GJ (1994) Characterization of 1,3- β -glucan (callose) synthesis in roots of *Triticum aestivum* in response to aluminum toxicity. *J Plant Physiol* **144**: 229-234

Zhang W, Zhang F, Shen Z and Liu Y (1998) Changes of H⁺ pumps of tonoplast vesicle from wheat roots *in vivo* and *in vitro* under aluminum treatment and effect of calcium. *J Plant Nutr* **21**: 2515-2526

3. Vacuolar H⁺-ATPase, but not Mitochondrial F₁F₀-ATPase, is Required for Aluminum Resistance in *Saccharomyces cerevisiae*²

3.1 Introduction

Although many mechanisms have been proposed to account for aluminum (Al) resistance in crop species (see for example Taylor, 1991), few have been tested directly. I have shown that vacuolar ATPase and mitochondrial F₁F₀-ATPase activities are induced by Al stress in the Al-resistant cultivar of wheat, PT741 (Chapter 2, Hamilton *et al.*, 2001). Synthesis of RMP51 (consisting of the B subunit of the V-ATPase and the α and β subunits of the F₁F₀-ATPase) is induced by Al in a dose- and time-dependent manner, and protein levels decline after removal of Al (Basu *et al.*, 1994). Aluminum-induced synthesis of RMP51 segregates with Al resistance in progeny derived from a cross between sensitive and resistant cultivars (Taylor *et al.*, 1997). This suggests that induction of V-ATPase and F₁F₀-ATPase may be an adaptive trait involved in Al resistance.


To further our understanding of Al resistance, I have developed a testable

² A version of this chapter has been published. Hamilton CA, Good AG and Taylor GJ (2001) Vacuolar H⁺-ATPase, but not Mitochondrial F₁F₀-ATPase, is Required for Aluminum Resistance in *Saccharomyces cerevisiae*. FEMS Microbiol Lett **205**: 231-236

hypothesis describing the role of V-ATPase and F_1F_0 -ATPase in Al resistance (Fig 3-1). Based on my model, inhibition of the plasma membrane ATPase (P-ATPase) by Al would cause proton accumulation in the cytoplasm and decreased cytoplasmic pH. To counteract this effect, I predict an increase in activity of the vacuolar ATPase (V-ATPase) which would transport excess protons into the lumen of the vacuole, thus alleviating the stress caused by cytoplasmic acidosis. Increased V-ATPase activity would also contribute to energizing the tonoplast membrane. It is possible that maintaining the tonoplast membrane electrochemical potential could provide energy required for a secondary active transport process involved in Al resistance. The ATP required for these processes would be provided by increased activity of ATP synthase (F_1F_0 -ATPase). To test this model directly, I used yeast as a model system. Although yeast and plants do not necessarily respond to stress in a similar fashion, yeast has the advantages of rapid, simple transformation, short life cycle and amenability to manipulation. This allows rapid hypothesis testing before, or while, more arduous experiments in plants are undertaken.

In this chapter I examined the response of wild type yeast to Al and tested the effect of mutations in V-ATPase and F_1F_0 -ATPase subunits on Al resistance. If my model is correct, then inhibiting the activity of the V-ATPase should cause increased sensitivity to Al. Inhibiting the activity of the F_1F_0 -ATPase may have a lesser effect on Al resistance, since yeast, unlike plants, can produce ATP to support V-ATPase activity via fermentation.

Fig 3-1. Model representing the energy balance hypothesis for V-ATPase and F_1F_0 -ATPase-mediated Al resistance in wheat. (1) Inhibition of the plasma membrane ATPase (P-ATPase) by Al. (2) Proton accumulation in the cytoplasm causes cytoplasmic acidosis. (3) Increased activity of the vacuolar ATPase (V-ATPase) could transport excess protons into the lumen of the vacuole, alleviating stress caused by decreased cytoplasmic pH and energizing the tonoplast membrane. (4) It is possible that maintaining the tonoplast membrane potential could provide energy required for a secondary active transport process involved in a hypothetical Al resistance mechanism. (5) The energy required for these processes could be provided by increased activity of ATP synthase (F_1F_0 -ATPase).



3.2 Methods

3.2.1 Media and culture conditions

Yeast strains (Table 3-1) were maintained on standard media as described by Sherman (1991). For experiments using Al, a low pH, low phosphate (LPP) medium (Schott and Gardner, 1997) was used. This medium is synthetic complete (SC) minimal medium (0.67% yeast nitrogen base, 2% glucose) with K_2HPO_4 reduced to 100 μ M, KCl added to 4.9 mM and the pH reduced to 3.5.

3.2.2 ATPase activity measurements

Seed cultures (3 ml in SC) of wild-type yeast were inoculated from YPD plate stocks and incubated for 16-24 h at 30°C. Cells were harvested by centrifugation, washed three times and resuspended in 1 ml of sterile water. This cell suspension was used to inoculate triplicate 3 ml cultures of LPP medium (pH 3.5) containing 0-15 μ M $AlCl_3$ to a starting OD_{600} of approximately 0.01 and cultures were incubated for 16-24 h at 30°C (final OD_{600} approximately 0.9). Each 3 ml culture was harvested by centrifugation and spheroplasts prepared as described by Uchida *et al.* (1988). Cell suspensions were then sonicated at low intensity (Branson Sonifier, 10% output) for 30s to gently lyse cells. Cell debris was removed by centrifugation and the resulting supernatant used for enzyme assays. Quantitation of protein in cell extracts was performed using the Bradford assay (Bradford, 1976). Adenosine triphosphatase activity was then assayed as described by Briskin *et al.* (1987).

Table 3-1. Yeast strains used in this study

Strain	Genotype	Source
SF838-1D	<i>MATα</i> , <i>ade6</i> , <i>his4-519</i> , <i>leu2-3,112</i> , <i>ura3-52</i> , <i>pep4-3</i>	Rothman and Stevens, 1986
<i>vma1</i>	Isogenic to SF838-1D except <i>vma2Δ::LEU2</i>	Yamashiro <i>et al.</i> , 1990
<i>vma2</i>	Isogenic to SF838-1D except <i>vma1Δ::LEU2</i>	Kane <i>et al.</i> , 1990
<i>vma1/VMA1</i>	Isogenic to <i>vma1</i> except carrying wild-type <i>VMA1</i> on plasmid pRS316 (<i>URA3</i>)	This study
<i>vma2/VMA2</i>	Isogenic to <i>vma2</i> except carrying wild-type <i>VMA2</i> on plasmid pRS316 (<i>URA3</i>)	This study
W303-1A	<i>MATα</i> , <i>ade2-1</i> , <i>his3-1,15</i> , <i>leu2-3,112</i> , <i>trp1-1</i> , <i>ura3-1</i>	(Zhang <i>et al.</i> , 1999)
W303- $\Delta\alpha$	Isogenic to W303-1A except null deletion mutation in <i>ATP1</i>	(Zhang <i>et al.</i> , 1999)
W303- $\Delta\beta$	Isogenic to W303-1A except null deletion mutation in <i>ATP2</i>	(Zhang <i>et al.</i> , 1999)
W303- $\Delta\alpha$ / <i>ATP1</i>	Isogenic to W303- $\Delta\alpha$ except carrying wild-type <i>ATP1</i> on plasmid pRS314 (<i>TRP1</i>)	This study
W303- $\Delta\beta$ / <i>ATP2</i>	Isogenic to W303- $\Delta\beta$ except carrying wild-type <i>ATP2</i> on plasmid pRS316 (<i>URA3</i>)	This study

3.2.3 *Complementation of ATPase mutations*

Yeast strains carrying *LEU2* insertion mutations in *VMA1* and *VMA2*, as well as plasmids carrying the complementing genes, were provided by Dr Tom Stevens (University of Oregon). Strains carrying null deletion mutations of *ATP1* and *ATP2* and plasmids carrying the complementing genes were provided by Dr David Mueller (Chicago Medical School). In order to select for transformation of the plasmids in the appropriate mutant strains, *ATP1* and *VMA2* were recloned into pRS314 (*TRP1*) and pRS316 (*URA3*), respectively, using standard molecular biology techniques (Sambrook *et al.*, 1989). The complementing constructs were then transformed into appropriate yeast strains using the lithium acetate method (Geitz *et al.*, 1995).

3.2.4 *Aluminum dose response experiments*

Inoculum for each culture was prepared as described above and used to inoculate triplicate 3 ml cultures of SC medium containing 0-150 μM AlCl_3 to a starting OD_{600} of approximately 0.01. Cultures were incubated for 16-24 h at 30°C and the final OD_{600} of each culture was measured in a 96-well plate (Costar) using a Spectromax Plus plate reader and Softmax Pro v 1.1 software (Molecular Devices). For comparison, the OD_{600} values for each strain were expressed as a percentage of the control value for that strain.

3.2.5 *Immunoblotting*

Total yeast proteins were separated by SDS-PAGE (Laemmli, 1970) and electroblotted onto nitrocellulose (0.45 μm , BioRad Laboratories) membranes using the Mini Trans-Blot Cell (Bio-Rad Laboratories) according to the manufacturer's instructions. Membranes were blocked and probed as directed in the Lumi-glo Chemiluminescent Detection System manual (Kirkegaard and Perry Laboratories).

3.2.6 *RNA Isolation, Northern Hybridization and Analysis*

RNA was isolated from yeast cultures (approximately 5×10^7 cells) using the RNeasy Kit (Qiagen) according to the manufacturer's directions. Northern transfers were carried out using GeneScreen Plus (DuPont) membranes according to the manufacturer's directions. Probes for Northern blots were isolated from pRS316VMA2 and pRS316ATP1 by digestion with appropriate restriction enzymes (Amersham Pharmacia Biotech) and inserts were isolated from agarose gels using the QiaQuick gel extraction kit (Qiagen). Probes were then prepared by random priming (Sambrook *et al.*, 1989). Membranes were prehybridized, hybridized and washed according to the manufacturer's directions, followed by exposure to storage phosphor screens (Molecular Dynamics). Phosphor screens were developed using a Phosphorimager 445SI and the resulting images analyzed using ImagequaNT v 4.2 software. For figure preparation, representative lanes of triplicate samples were selected from the same blot and compiled using Adobe Photoshop^R v 5.5.

3.3 Results

3.3.1 Response of yeast ATPases to aluminum

To confirm that yeast is a suitable model for investigating the effect of Al on ATPases in plants, the responses of ATPases to Al in wild type yeast were examined (Fig 3-2). F_1F_0 -ATPase activity increased to $0.225 \text{ nmol Pi mg}^{-1} \text{ min}^{-1}$ (580% of control) at $2.5 \text{ }\mu\text{M AlCl}_3$. Although the Al-induced stimulation declined to $0.137 \text{ nmol Pi mg}^{-1} \text{ min}^{-1}$ (340% of control) from 5 to $10 \text{ }\mu\text{M AlCl}_3$, F_1F_0 -ATPase activity was still $0.151 \text{ nmol Pi mg}^{-1} \text{ min}^{-1}$ (380% greater than control) at $15 \text{ }\mu\text{M}$, similar to what has been observed in wheat (Hamilton *et al.*, 2001). Although the magnitude of the change was less, P-ATPase activity showed an effect similar to what has been observed in plants (Hamilton *et al.*, 2001; Matsumoto, 1988; Matsumoto *et al.*, 1992; Sasaki *et al.*, 1995; Widell *et al.*, 1994), decreasing to $0.064 \text{ nmol Pi mg}^{-1} \text{ min}^{-1}$ (54% of control) at $15 \text{ }\mu\text{M Al}$. Vacuolar ATPase activity remained essentially constant from 0- $10 \text{ }\mu\text{M AlCl}_3$. This is an interesting contrast to the Al-induced increase observed in wheat (Zhang *et al.*, 1998; Hamilton *et al.*, 2001) and barley (Kasai *et al.*, 1992, 1993).

To determine the mechanism by which ATPase activities change during Al stress, expression of V-ATPase and F_1F_0 -ATPase subunits was examined by Western and Northern analysis (Fig 3-3). Levels of the transcript encoding the F_1F_0 -ATPase α subunit increased to 201% of control levels by $7.5 \text{ }\mu\text{M AlCl}_3$, followed by a decline to 135% by $15 \text{ }\mu\text{M}$ (Fig 3-3A,B). This pattern closely resembles that observed for increased F_1F_0 -ATPase activity (Fig 3-2). Protein levels of the F_1F_0 -ATPase α and β

Fig 3-2. The effect of Al on cellular ATPase activities (F_1F_0 -ATPase, V-ATPase, P-ATPase) in wild type yeast. Values are means \pm SE (n=3). Results shown are representative of three independent trials.






Fig 3-3. The effect of Al on expression of V-ATPase and F₁F₀-ATPase. (A) Levels of the transcripts encoding the F₁F₀-ATPase α subunit and V-ATPase B subunit were measured by Northern analysis. (B) Relative abundance of transcripts as measured by densitometry. The ratio of mRNA/rRNA was measured at each concentration of AlCl₃ and compared to the control values for each transcript. (C) Determination of protein levels of V-ATPase (B subunit) and F₁F₀-ATPase (α and β subunits) by Western analysis. Cross-reacting bands were visualized using a chemiluminescent detection system. Samples shown are representative lanes selected from triplicates compiled using Adobe Photoshop^R v 5.5. Values shown are means \pm SE (n=3) and are representative of 2 independent trials.


A



B



C



subunits also increased with increasing AlCl_3 concentration (Fig 3-3C). Interestingly, levels of these subunits continue to increase past 5-7.5 μM AlCl_3 , the peak concentrations for F_1F_0 -ATPase activity and transcript levels.

Levels of the transcript encoding the V-ATPase B subunit remained nearly constant over the range of Al concentrations tested (Fig 3-3A,B), with only a slight increase to 123% of control at 5 μM AlCl_3 . The amount of protein also remained constant (Fig 3-3C), reflecting the constant level of V-ATPase activity observed during Al stress (Fig 3-2). In wheat, V-ATPase transcript, protein and activity increased in parallel with increasing concentrations of Al (Hamilton *et al.*, 2001, C.A. Hamilton, unpublished data).

3.3.2 V-ATPase mutants, but not F_1F_0 -ATPase mutants, are hypersensitive to Al

To test the hypothesis that V-ATPase and F_1F_0 -ATPase activities are involved in Al resistance, yeast mutants lacking subunits of these enzymes were tested to determine whether or not they are hypersensitive to Al. The relative sensitivities of V-ATPase mutants, their parent strain and complemented strains to Al were determined by observing their growth in liquid culture. Cultures were grown in SC medium (pH 4.5) since *vma1* and *vma2* mutant strains grew poorly in the low phosphate, low pH medium, LPP (Schott and Gardner, 1997), normally used for experiments involving Al. In the V-ATPase mutant strains (*vma1*, *vma2*), there was a striking 40% decrease in growth (from 0.85 and 1.09 control OD_{600} to 0.45 and 0.62 OD_{600} at 75 μM AlCl_3 , respectively) at all Al concentrations tested, while the parent

strain (control OD₆₀₀ 1.28) was unaffected (Fig 3-4a). The complemented strains (control OD₆₀₀ values of 1.15 and 1.19, respectively) were similarly unaffected by Al. This confirmed that the V-ATPase mutations are responsible for the Al sensitive phenotype of the *vma1* and *vma2* strains.


The relative sensitivities of F₁F₀-ATPase mutants, their parent strain and complemented strains to Al were also measured (Fig 3-4B). In contrast to V-ATPase mutants, F₁F₀-ATPase mutants did not show hypersensitivity to Al from 0-150 μM. To determine whether higher concentrations were required to observe phenotypic differences, levels of Al up to 400 μM were tested (Fig 3-4B, inset). No discernable differences were observed between the five strains.

3.4 Discussion


The responses of plants and yeast to Al are somewhat different. For example, Al toxicity in yeast can be overcome by overexpression of a Mg²⁺ transporter (MacDiarmid and Gardner, 1998), presumably due to inhibition of Mg²⁺ transport by Al³⁺. In contrast, although addition of Mg²⁺ (or other divalent cations such as Ca²⁺) alleviates Al³⁺ toxicity in wheat (Kinraide, 1998), reduced Mg²⁺ availability does not cause Al toxicity symptoms (Kinraide and Parker, 1987). Also, many gene products induced by Al in plants have no protective effect against Al when expressed in yeast (Ezaki *et al.*, 1999). This is likely because many Al-induced proteins are general stress response proteins (induced in response to oxidative stress and heat shock) which do not necessarily have a role in Al resistance. Therefore, physiological and

Fig 3-4. The effect of ATPase mutations on Al resistance in yeast. (a) Yeast strains carrying LEU2 insertion mutations in the *VMA1* and *VMA2* genes (*vma1*, *vma2*), their isogenic parent strain (SF838), and strains carrying wild type copies of the *VMA1* and *VMA2* genes on low copy plasmids (*vma1/VMA1*, *vma2/VMA2*) were grown for 16-24 h in SC minimal medium (pH 4.5) containing 0-150 μM AlCl_3 . (b) Yeast strains with null deletion mutations in the *atp1* and *atp2* genes (W303- $\Delta\alpha$, W303- $\Delta\beta$), their isogenic wild type parent strain (W303-1A) and strains carrying wild type copies of the *atp1* and *atp2* genes on low copy plasmids ($\Delta\alpha$ -*ATP1*, $\Delta\beta$ -*ATP2*) were grown for 16-24 h in SC minimal medium (pH 4.5) containing 0-150 μM AlCl_3 (the inset shows that similar results were observed up to 400 μM AlCl_3). Total growth (OD_{600}) of each culture was measured and compared to the control (0 AlCl_3) values for each strain. Values shown are means \pm SE (n=3). Results shown are representative of three independent trials.

A



B



structural differences between yeast and plants must be considered when analyzing data from this system.

It was recently shown by Ezaki *et al.* (1998) that yeast cells exposed to Al show irregular vacuolar morphology. The authors did not speculate as to the possible cause of this effect, but it is possible that Al associated with the yeast vacuole causes this phenotype. Aluminum uptake has been demonstrated in yeast (Jagannatha *et al.*, 1997; Ezaki *et al.*, 1999) and the giant algae *Chara corallina* (Taylor *et al.*, 2000), where it is associated mainly with the cell wall, but is also taken up into the vacuole. Ezaki *et al.* (1999) suggested that intracellular Al does not associate with the yeast vacuole based on observations of morin fluorescence, but low levels such as those measured in *Chara* ($<5 \mu\text{g m}^{-2}$, Taylor *et al.*, 2000) would not be detected using this method. It is possible that Al is sequestered in the vacuole by an active transport mechanism which relies on the tonoplast electrochemical potential. Once in the vacuole, Al could be effectively sequestered by complexing with polyphosphate (as is calcium) (Dunn *et al.*, 1994) or organic anions.

Here I have used the yeast model system for simple and direct testing of a hypothesis derived from higher plants. Based on experiments in plants, I hypothesized that the V-ATPase is involved in Al resistance, with the ATP required for its activity supplied by increased activity of ATP synthase. Although V-ATPase transcript and protein levels, as well as enzyme activity, are not affected by increasing Al concentrations in yeast as they are in plants, the observation that V-ATPase mutants are hypersensitive to Al supports my hypothesis. In wild-type yeast, F_1F_0 -

ATPase transcript, protein and activity levels all increase in response to Al, while F_1F_0 -ATPase mutants do not show an Al hypersensitive phenotype. It is possible that the demand for ATP can be met by fermentation, although ATP synthase would normally fulfill this requirement in wild-type yeast and plants.

3.5 Literature Cited

Basu A, Basu U and Taylor GJ (1994) Induction of microsomal membrane proteins in roots of an aluminum-resistant cultivar of *Triticum aestivum* L. under conditions of aluminum stress. *Plant Physiol* **104**: 1007-1013

Bradford M (1976) A rapid and sensitive method for the quantification of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* **72**: 248-254

Briskin DP, Leonard RT and Hodges TK (1987) Isolation of the plasma membrane: Membrane markers and general principals. *In* L Packer, R Douce, eds, *Methods in Enzymology*. Vol 148. Academic Press, pp 542-558

Dunn T, Gable K and Beeler T (1994) Regulation of cellular Ca^{2+} by yeast vacuoles. *J Biol Chem* **269**: 7273-7278

Ezaki B, Sivaguru M, Ezaki Y, Matsumoto H and Gardner RC (1999) Acquisition of aluminum tolerance in *Saccharomyces cerevisiae* by expression of the *BCB* or *NtGD11* gene derived from plants. *FEMS Microbiol Lett* **171**: 81-87

Ezaki B, Gardner RC, Ezaki Y, Kondo H and Matsumoto H (1998) Protective roles of two aluminum (Al)-induced genes, *HSP150* and *SEDI* of *Saccharomyces cerevisiae*, in Al and oxidative stresses. *FEMS Microbiol Lett* **159**: 99-105

- Geitz RD, Schiestl RH, Willems AR and Woods RA (1995)** Studies on the transformation of intact yeast cells by the LiAc/SS-DNA/PEG procedure. *Yeast* **11**:355-360
- Hamilton CA, Good AG and Taylor GJ (2001)** Induction of vacuolar ATPase and mitochondrial ATP synthase by aluminum in an aluminum-resistant cultivar of wheat. *Plant Physiol* **125**: 2068-2077
- Jagannatha Rao KS and Easwaran KRK (1997)** ²⁷Al-NMR studies of aluminum transport across yeast cell membranes. *Mol Cell Biochem* **175**: 59-63
- Kane PM, Yamashiro CT, Wolczyk DF, Neff N, Goebel M and Stevens TH (1990)** Protein splicing converts the yeast TFP1 gene product to the 69-kD subunit of the vacuolar H⁺-adenosine triphosphatase. *Science* **250**: 651-7
- Kasai M, Sasaki M, Yamamoto Y and Matsumoto H (1992)** Aluminum stress increases K⁺ efflux and activities of ATP- and PPI-dependent H⁺ pumps of tonoplast-enriched membrane vesicles from barley roots. *Plant Cell Physiol* **33**: 1035-1039
- Kasai M, Sasaki M, Tanakamaru S, Yamamoto Y and Matsumoto H (1993)** Possible involvement of abscisic acid in increases in activities of two vacuolar H⁺-pumps in barley roots under aluminum stress. *Plant Cell Physiol* **34**: 1335-1338
- Kinraide TB (1998)** Three mechanisms for the calcium alleviation of mineral toxicities. *Plant Physiol* **118**: 513-20

- Kinraide TB and Parker DR** (1987) Cation amelioration of aluminum toxicity in wheat. *Plant Physiol* **83**: 546-551
- Laemmli UK** (1970) Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* **227**:680-5
- Lai-Zhang J, Xiao Y and Mueller DM** (1999) Epistatic interactions of deletion mutants in the genes encoding the F₁-ATPase in yeast *Saccharomyces cerevisiae*. *EMBO J* **18**: 58-64
- MacDiarmid CW and Gardner RC** (1998) Overexpression of the *Saccharomyces cerevisiae* magnesium transport system confers resistance to aluminum ion. *J Biol Chem* **273**: 1727-1732
- Matsumoto H** (1988) Inhibition of proton transport activity of microsomal membrane vesicles of barley roots by aluminium. *Soil Sci Plant Nutr* **34**: 499-526
- Matsumoto H, Yamamoto Y and Kasai M** (1992) Changes of some properties of the plasma membrane-enriched fraction of barley roots related to aluminum stress: membrane-associated ATPase, aluminum and calcium. *Soil Sci Plant Nutr* **38**: 411-419
- Rothman JH and Stevens TH** (1986) Protein sorting in yeast: The localization determinant of yeast vacuolar carboxypeptidase Y resides in the propeptide. *Cell* **48**: 887-97
- Sambrook J, Fritsch EF and Maniatis T** (1989) *Molecular Cloning: A Laboratory Manual*, 2nd Ed, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY

- Sasaki M, Kasai M, Yamamoto Y and Matsumoto H** (1995) Involvement of plasma membrane potential in the tolerance mechanism of plant roots to aluminum toxicity. *In* RA Date *et al.*, eds. Plant-Soil Interactions at Low pH. Proceedings of the Second International Symposium on Plant-Soil Interactions at Low pH, 24-29 June 1990, Beckley, West Virginia, USA. Kluwer Academic Publishers, Dordrecht, pp 285-290
- Schott EJ and Gardner RC** (1997) Aluminum-sensitive mutants of *Saccharomyces cerevisiae*. *Mol Gen Genet* **254**: 63-72
- Sherman F** (1991) Getting started with yeast. *Methods Enzymol* **194**: 3-21
- Taylor GJ** (1991) Current views of the aluminum stress response; the physiological basis of tolerance. *Curr Topics Plant Biochem and Physiol* **10**: 57-93
- Taylor GJ, Basu A, Basu U, Slaski JJ, Zhang G and Good A** (1997) Al-induced, 51 kilodalton, membrane-bound proteins are associated with resistance to Al in a segregating population of wheat. *Plant Physiol* **114**: 363-372
- Taylor GJ, McDonald-Stephens JL, Hunter DB, Bertsch PM, Elmore D, Rengel Z and Reid RJ** (2000) Direct measurement of aluminum uptake and distribution in single cells of *Chara corallina*. *Plant Physiol* **123**: 987-996
- Uchida E, Ohsumi Y and Anraku Y** (1988) Purification of yeast vacuolar membrane H⁺-ATPase and enzymological discrimination of three ATP-driven proton pumps in *Saccharomyces cerevisiae*. *Methods Enzymol* **157**: 544-562

Widell S, Asp H and Jensen P (1994) Activities of plasma membrane-bound enzymes isolated from roots of spruce (*Picea abies*) grown in the presence of aluminium. *Physiol Plant* **92**: 459-466

Yamashiro CT, Kane PM, Wolczyk DF, Preston RA and Stevens TH (1990) Role of vacuolar acidification in protein sorting and zymogen activation: a genetic analysis of the yeast vacuolar proton-translocating ATPase. *Mol Cell Biol.* **10**: 3737-49

Zhang W, Zhang F, Shen Z and Liu Y (1998) Changes of H⁺ pumps of tonoplast vesicle from wheat roots *in vivo* and *in vitro* under aluminum treatment and effect of calcium. *J Plant Nutr* **21**: 2515-2526

4. Vacuolar H⁺-ATPase, but not Mitochondrial F₁F₀-ATPase, is Required for NaCl Tolerance in *Saccharomyces cerevisiae*³

4.1 Introduction

Salt tolerance is a complex trait involving responses to both osmotic and ionic stresses. The mechanisms by which organisms are able to tolerate high concentrations of salt in their environment have been extensively studied by mutational analysis of the budding yeast *Saccharomyces cerevisiae*. Data obtained thus far indicate that an array of signaling pathways activate salt-responsive genes involved in modulating plasma membrane polarization, Na⁺ influx, Na⁺ efflux and sequestration of Na⁺ in the vacuole.

One effect of salt stress is a hyperpolarization of the plasma membrane. A low affinity cation uptake current activated by membrane hyperpolarization (NSC1) has been observed in patch clamp studies (Bihler *et al.*, 1998) and may contribute to subsequent depolarization of the membrane. Also contributing to depolarization of the plasma membrane are protein kinases HAL4 and HAL5, the induction of which increases the selectivity of the high affinity TRK1/2 K⁺ channel (Mulet *et al.*, 1999). Influx of K⁺ and H⁺ (and perhaps Na⁺) through this channel dissipates the driving force for uptake of Na⁺ across the plasma membrane. The Ser-Thr protein

³ A version of this chapter has been submitted for publication. Hamilton CA, Taylor GJ and Good AG (2001) Vacuolar H⁺-ATPase, but not Mitochondrial F₁F₀-ATPase, is Required for NaCl Tolerance in *Saccharomyces cerevisiae*. FEMS Microbiol Lett (In Press)

phosphatase, calcineurin, is also required for increased selectivity of TRK1/2 (Mendoza *et al.*, 1994). Calcineurin is a heterotrimer composed of a catalytic subunit (encoded by either *CNA1* or *CNA2*), a Ca^{2+} -binding regulatory subunit (encoded by *CNB1*) and Ca^{2+} -calmodulin (Stemmer and Klee, 1994). Binding of Ca^{2+} /calmodulin to the catalytic subunit prevents its autoinhibitory activity (Cohen, 1989) so that calcineurin is free to interact with a variety of substrates within the cell, regulating the salt response.

Calcineurin is a central regulator of Na^+ influx and efflux. Sodium ions may enter the cell through the TRK1/2 channel or the NSC1 current. Under alkaline or neutral conditions, Na^+ is exported mainly through the action of PMR2/ENA1, a Na^+ -ATPase localized to the plasma membrane (Prior *et al.*, 1996). Expression and activity of this enzyme is controlled by many factors, highlighting its importance to salt tolerance. Activity of ENA1 is stimulated by calcineurin (Cunningham and Fink, 1996), HAL1,8,9 (Gaxiola *et al.*, 1992; Rios *et al.*, 1997; Mendizabal *et al.*, 1998), SNF1 (Rios *et al.*, 1997), and in *Schizosaccharomyces pombe*, AFT1, a transcription factor activated by the *STY1/WIS1* MAPK cascade (Nishikawa *et al.*, 1999). It is also indirectly activated by HAL3, a negative regulatory subunit of PPZ1/2 protein phosphatases, which normally inhibit ENA1 activity (Posas *et al.*, 1995). Carbon source also affects ENA1 activity. The presence of glucose inhibits SNF1 (an activator of ENA1) and activates protein kinase A, an inhibitor of ENA1 (Rios *et al.*, 1997). Under acidic conditions such as those used in this study, efflux of Na^+ is mediated mainly by the Na^+/H^+ antiporter, NHA1 (Prior *et al.*, 1996).

Another mechanism of salt tolerance in yeast is sequestration in the vacuole. Transport of Na^+ into the vacuole is mediated by *NHX1*, a Na^+/H^+ antiporter localized to a prevacuolar compartment and activated by cytoplasmic acidosis (Nass *et al.*, 1997; Nass and Rao, 1998). Other proteins required for *NHX1*-mediated salt tolerance are the chloride channel (*GEF1*) and the vacuolar ATPase (Gaxiola *et al.*, 1999).

Since transmembrane transport of Na^+ is a key component of salt tolerance, the objective of the present study was to characterize the responses of cellular ATPases (P-ATPase, F_1F_0 -ATPase and V-ATPase) to salt. The contributions of the V-ATPase and F_1F_0 -ATPase were of particular interest because the V-ATPase (with ATP provided by F_1F_0 -ATPase) creates the proton motive force (PMF) required for import of cations into the vacuole. Changes in ATPase transcript levels, protein and activity in response to salt were measured, and the phenotypes of V-ATPase and F_1F_0 -ATPase mutant and complemented strains were compared.

4.2 Materials and Methods

4.2.1 Media and culture conditions

Yeast strains (Table 3-1) were maintained on standard media as described by Sherman, 1991.

4.2.2 ATPase activity measurements

Seed cultures (3 ml in synthetic complete medium, SC) of wild-type yeast were inoculated from YPD plate stocks and incubated for 16-24 h at 30°C. Cells were

harvested by centrifugation, washed three times and resuspended in 1 ml of sterile water. This cell suspension was used to inoculate triplicate 3 ml cultures of SC medium containing 0-1000 mM NaCl to a starting OD₆₀₀ of approximately 0.01 and cultures were incubated for 16-24 h at 30°C (final OD₆₀₀ approximately 0.9). Each 3 ml culture was harvested by centrifugation and spheroplasts were prepared as described by Uchida *et al.*, 1988. Cell suspensions were then sonicated at low intensity (Branson Sonifier, 10% output) for 30 s to gently lyse cells. Cell debris was removed by centrifugation and the resulting supernatant used for enzyme assays. Quantitation of protein in cell extracts was performed using the Bradford assay (Bradford, 1976). Adenosine triphosphatase activity was then assayed as described by Briskin *et al.*, 1987.

4.2.3 *Complementation of ATPase mutations*

Yeast strains carrying *LEU2* insertion mutations in *vma1* and *vma2*, as well as plasmids carrying the complementing genes, were provided by Dr Tom Stevens (University of Oregon). Strains carrying null deletion mutations of *atp1* and *atp2* and plasmids carrying the complementing genes were provided by Dr David Mueller (Chicago Medical School). In order to select for transformation of the plasmids in the appropriate mutant strains, *ATP1* and *VMA2* were recloned into pRS314 (TRP1) and pRS316 (URA3) and transformed into appropriate yeast strains as described by Hamilton *et al.*, 2001 (Chapter 2).

4.2.4 *NaCl dose response experiments*

Inoculum for each culture was prepared as described above and used to inoculate triplicate 3 ml cultures of SC medium containing 0-1000 mM NaCl to a starting OD₆₀₀ of approximately 0.01. Cultures were incubated for 16-24 h at 30°C and the final OD₆₀₀ of each culture was measured in a 96-well plate (Costar) using a Spectromax Plus plate reader and Softmax Pro v 1.1 software (Molecular Devices). For comparison, the OD₆₀₀ values for each strain were expressed as a percentage of the control value for that strain. For halotolerance tests, 5 µL aliquots of serial 10⁻¹ dilutions of overnight cultures were spotted onto YPD medium containing 1 M NaCl, 1.5 M sorbitol, 100 mM LiCl or 50 µg ml⁻¹ hygromycin B and incubated at 30°C for 2-4 d.

4.2.5 *Immunoblotting*

Total yeast proteins were separated by SDS-PAGE and electroblotted onto nitrocellulose (0.45 µm, BioRad Laboratories) membranes using the Mini Trans-Blot Cell (Bio-Rad Laboratories) according to the manufacturer's instructions. Membranes were blocked and probed as directed in the Lumi-glo Chemiluminescent Detection System manual (Kirkegaard and Perry Laboratories). Polyclonal antibodies raised against the α and β subunits of the yeast F₁-ATPase were provided by Dr A. Lewin, University of Florida. Antibodies raised against the B subunit of the *Vigna radiata* V-ATPase were provided by Dr M. Maeshima, Nagoya University.

4.2.6 RNA Isolation, Northern Hybridization and Analysis

RNA was isolated from yeast cultures (approximately 5×10^7 cells) using the RNeasy Kit (Qiagen) according to the manufacturer's directions. Northern transfers were carried out using GeneScreen Plus (DuPont) membranes according to the manufacturer's directions. Probes for Northern blots were isolated from pRS316VMA2 and pRS316ATP1 by digestion with appropriate restriction enzymes (Amersham Pharmacia Biotech) and inserts were isolated from agarose gels using the QiaQuick gel extraction kit (Qiagen). Probes were then prepared by random priming. Membranes were prehybridized, hybridized and washed according to the manufacturer's directions, followed by exposure to storage phosphor screens (Molecular Dynamics). Phosphor screens were developed using a Phosphorimager 445SI and the resulting images analyzed using ImagequaNT v 4.2 software. For figure preparation, representative lanes of triplicate samples were selected from the same blot and compiled using Adobe Photoshop^R version 5.5.

4.3 Results

4.3.1 Response of yeast ATPases to NaCl

The responses of cellular ATPases (F_1F_0 -ATPase, P-ATPase and V-ATPase) to NaCl in wild type yeast were characterized by measuring ATPase activities, protein and transcript levels. Adenosine triphosphatase activities were measured after growth

in SC medium containing 0-1000 mM NaCl (Fig 4-1). F_1F_0 -ATPase activity gradually increased from $0.078 \text{ nmol Pi mg}^{-1} \text{ min}^{-1}$ to $0.154 \text{ nmol Pi mg}^{-1} \text{ min}^{-1}$ (197% of control), reaching its peak activity at 600 mM NaCl. P-ATPase activity remained essentially constant from 0-800 mM NaCl, but declined to $0.056 \text{ nmol Pi mg}^{-1} \text{ min}^{-1}$ (57% of control) by 1000 mM. At first sight this is surprising, since the main determinant of NaCl tolerance in yeast is ordinarily the ENA1 Na^+ efflux pump (a P-type ATPase). However, under the acidic conditions used here (pH 4.5), Na^+/H^+ antiport mediated by NHA1 would predominate (Prior *et al.*, 1996). The presence of glucose in the growth medium would also inhibit the activity of ENA1 by repression of SNF1, an activator of ENA1, and by activation of protein kinase A, an inhibitor of ENA1 (Rios *et al.*, 1997).

In contrast, vacuolar ATPase activity increased sharply from $0.103 \text{ nmol Pi mg}^{-1} \text{ min}^{-1}$ to $0.303 \text{ nmol Pi mg}^{-1} \text{ min}^{-1}$ (296% of control) from 0-200 mM NaCl, and decreased slightly at higher salt concentrations. However, even at 1000 mM NaCl, V-ATPase activity was still comparatively high, at $0.194 \text{ nmol Pi mg}^{-1} \text{ min}^{-1}$ (189% of control).

To determine the mechanism by which ATPase activities change during NaCl stress, expression of V-ATPase and F_1F_0 -ATPase subunits was examined by Northern and Western analysis (Fig 4-2). Levels of the transcript encoding the F_1F_0 -ATPase α subunit increased to 325% of the control level by 800 mM NaCl, followed by a decline to 257% of control by 1000 mM (Fig 4-2A,B). This pattern is similar to that observed for increased F_1F_0 -ATPase activity (Fig 4-1), with the exception that

Fig 4-1. The effect of NaCl on cellular ATPase activities (F_1F_0 -ATPase, P-ATPase, V-ATPase) in wild type yeast. Cultures were grown 16-24 h in the presence of 0-1000 mM NaCl in SC medium. Values shown are means \pm SE (n=3). Results shown are representative of three independent trials.







Fig 4-2. The effect of NaCl on expression of F₁F₀-ATPase and V-ATPase. (A) Levels of the transcripts encoding the F₁F₀-ATPase α subunit and V-ATPase B subunit were measured by Northern analysis. (B) Relative abundance of transcripts as measured by densitometry. The ratio of mRNA/rRNA measured at each concentration of NaCl was compared to the control value for each transcript. Values shown are means \pm SE (n=3). (C) Determination of protein levels of V-ATPase (B subunit) and F₁F₀-ATPase (α and β subunits) by Western analysis. Cross-reacting bands were visualized using a chemiluminescent detection system. Samples shown are representative lanes selected from triplicates and compiled using Adobe Photoshop^R v 5.5. For all panels, data shown are representative of 2 independent trials.




enzyme activity peaks at 600 mM while the transcript level peaks at 800 mM. Protein levels of the F_1F_0 -ATPase α and β subunits also increased with increasing NaCl concentration, as determined by Western analysis (Fig 4-2C). Interestingly, protein levels of these subunits peak at 600-800 mM NaCl, which are also the peak concentrations for F_1F_0 -ATPase transcript levels. Levels of the transcript encoding the V-ATPase B subunit also increased gradually over the range of NaCl concentrations tested (Fig 4-2A,B), reaching a peak (248% of control) at 600 mM NaCl. The amount of protein also increased in response to NaCl stress. Like the α and β subunits of the F_1F_0 -ATPase, protein levels of the V-ATPase B subunit continued to increase from 0-800 mM NaCl (Fig 4-2C), in contrast to the peak levels of enzyme activity and transcript abundance (200 mM and 600 mM NaCl, respectively).

4.3.2 *V-ATPase mutants, but not F_1F_0 -ATPase mutants, are hypersensitive to NaCl*


To test the hypothesis that V-ATPase and F_1F_0 -ATPase activities are involved in salt tolerance, yeast mutants lacking subunits of these enzymes were tested to determine whether or not they are hypersensitive to NaCl. The relative sensitivities of V-ATPase mutants, wild-type parent and complemented strains to NaCl were determined by observing growth in liquid culture. Final OD_{600} values at each NaCl concentration were related to the control values for each strain (Fig 4-3A). For each V-ATPase mutant strain (*vma1*, *vma2*), there was a 40-50% greater decrease in growth compared to wild-type at 600-1000 mM NaCl (from control OD_{600} values of

Fig 4-3. The effect of ATPase mutations on NaCl tolerance in yeast. (A) Yeast strains carrying LEU2 insertion mutations in the *VMA1* and *VMA2* genes (*vma1*, *vma2*), their isogenic parent strain (SF838), and strains carrying wild type copies of the *VMA1* and *VMA2* genes on low copy plasmids (*vma1/VMA1*, *vma2/VMA2*) were grown for 16-24 h in SC minimal medium (pH 4.5) containing 0-1000 mM NaCl. (B) Yeast strains with null deletion mutations in the *atp1* and *atp2* genes (W303- $\Delta\alpha$, W303- $\Delta\beta$), their isogenic wild type parent strain (W303-1A) and strains carrying wild type copies of the *atp1* and *atp2* genes on low copy plasmids ($\Delta\alpha$ -*ATP1*, $\Delta\beta$ -*ATP2*) were grown for 16-24 h in SC minimal medium (pH 4.5) containing 0-1000 mM NaCl. Total growth (OD_{600}) of each culture was measured and compared to the control (0 NaCl) values for each strain. Values shown are means \pm SE (n=3). Results shown are representative of three independent trials.

A



B



0.49 and 0.48 to OD₆₀₀ values of 0.08 and 0.05, for *vma1* and *vma2* respectively at 800 mM NaCl). In contrast, the complemented strains grew nearly as well as wild-type at these concentrations (control OD₆₀₀ values of 0.82, 0.73 and 0.76 and OD₆₀₀ values of 0.55, 0.46 and 0.46 at 800 mM NaCl for SF838, *vat1/VAT1* and *vat2/VAT2* respectively).


The salt-sensitive phenotype observed for the V-ATPase mutant strain, *vma1*, was specific to NaCl and LiCl (Fig 4-4). This mutant grew as well as the wild-type parent and complemented strains on YPD medium containing other osmotic (sorbitol) and ionic (hygromycin B) stress-inducing agents. In contrast, hypersensitivity of V-ATPase mutants was observed on YPD medium containing 1 M NaCl or 100 mM LiCl (Fig 4-4).

The relative sensitivities of F₁F₀-ATPase mutant strains, parent and complemented strains (W303-Δα, W303-Δβ, W303-1A, Δα-*ATP1* and Δβ-*ATP2*) to NaCl were also measured (Fig 4-3B). In contrast to the V-ATPase mutants, the F₁F₀-ATPase mutants were not hypersensitive to NaCl from 0-1000 mM (control OD₆₀₀ values of 0.87, 0.79, 0.73, 0.83, and 0.87 for W303-1A, W303-Δα, W303-Δβ, Δα-*ATP1* and Δβ-*ATP2*, respectively). Growth of the W303-Δβ mutant strain was indistinguishable from its parent and complemented strains in the presence of hygromycin B, LiCl, sorbitol and NaCl on YPD solid medium (Fig 4-4).

Fig 4-4. Halotolerance of V-ATPase and F₁F₀-ATPase mutant strains. Saturated cultures of wild-type parent (SF838, W303-1A), mutant (*vma1*, W303-Δβ) and complemented (*vma1/VMA1*, Δβ-*ATP2*) strains were serially diluted and 5 μL aliquots of 10⁻¹ to 10⁻⁵ dilutions were spotted onto YPD plates containing (A) no added compounds, (B) 1.5 M sorbitol, (C) 1 M NaCl, (D) 100 mM LiCl or (E) 50 μg ml⁻¹ hygromycin B. Results shown are representative of two independent trials.

A. Control

1 2 3 4 5



SF838
vma1
vma1/VMA1
W303-1A
W303-Δβ
Δβ-ATP2


B. 1.5 M Sorbitol

1 2 3 4 5



C. 1 M NaCl

1 2 3 4 5



SF838
vma1
vma1/VMA1
W303-1A
W303-Δβ
Δβ-ATP2

D. 100 mM LiCl

1 2 3 4 5



E. 50 μg/mL Hyg

1 2 3 4 5



SF838
vma1
vma1/VMA1
W303-1A
W303-Δβ
Δβ-ATP2

4.4 Discussion

This chapter highlights the importance of the vacuolar ATPase in yeast NaCl tolerance. In wild-type yeast, NaCl stress causes increased activity of the V-ATPase, with ATP presumably provided by increased activity of the F_1F_0 -ATPase. The V-ATPase is thought to contribute to NaCl tolerance by creating the PMF required for sequestration of Na^+ in the vacuole by the Na^+/H^+ antiporter, NHX1 (Nass *et al.*, 1997). The electrical potential created by the V-ATPase also creates a driving force for uptake of Cl^- through the chloride channel (GEF1). Therefore, it is possible that V-ATPase mutant strains are hypersensitive to NaCl because they are less able to effectively sequester Na^+ and Cl^- in the vacuole, allowing Na^+ to interact with sensitive cytoplasmic targets. This hypothesis is supported by recent experiments in which the vacuolar H^+ -pyrophosphatase (AVP1) from *Arabidopsis thaliana* was overexpressed in yeast and in *Arabidopsis*. Overexpression of this enzyme mimics overexpression of the V-ATPase, and in both species resulted in enhanced NaCl tolerance (Gaxiola *et al.*, 1999, 2001). Further support for the involvement of the vacuole in NaCl tolerance comes from the observation that overexpression of the vacuolar Na^+/H^+ antiporter also confers enhanced NaCl tolerance in yeast (Gaxiola *et al.*, 1999) and *Arabidopsis* (Apse *et al.*, 1999).

Interestingly, my observation that V-ATPase activity is stimulated by NaCl in wild-type yeast contrasts observations by Perzov *et al.* (2001). These authors examined vacuoles isolated from yeast cells (strain W303-1A) grown in YPD medium containing 800 mM NaCl and found that proton uptake activity was

decreased in comparison to activity in vacuoles isolated from cells grown in medium without NaCl. However, the differences in medium composition (YPD vs SC) and pH (neutral vs acidic) make direct comparison difficult, possibly highlighting the importance of environment in determining which salt tolerance mechanisms are active.

My observation that V-ATPase mutants were inhibited 40-50% more than the wild-type parent and complemented strains at 600-1000 mM NaCl is consistent with those obtained by Gaxiola *et al.* (1999), who found that overexpression of the *Arabidopsis* vacuolar H⁺-pyrophosphatase led to increased NaCl tolerance. Haro *et al.* (1993) examined the NaCl tolerance of yeast strains lacking the C subunit of the V-ATPase. Although my observation that V-ATPase mutants are hypersensitive to NaCl is in agreement with their results, they also reported that these mutants tolerate only 15 mM LiCl, contrary to our observations. I observed hypersensitivity at 100 mM LiCl but not at 50 mM LiCl (data not shown). It is possible that these differences arose due to the different media used (YPD vs minimal medium arginine phosphate) or because I performed growth tests on solid medium while Haro *et al.* used liquid medium. In my experience, V-ATPase mutant strains grow poorly in liquid culture because the cells aggregate and have a slow growth rate. These confounding effects, caused by the pleiotropic nature of V-ATPase mutations, are not apparent when these strains are grown on solid medium.

Although it was previously known from genetic studies that V-ATPase mutants are hypersensitive to NaCl stress (Haro *et al.*, 1993), a detailed study of the

effect of NaCl on ATPase activity and expression in wild-type yeast has not been described in the literature. This chapter showed that NaCl induces activities of V-ATPase and F_1F_0 -ATPase, and represses activity of the P-ATPase at high NaCl concentrations (1 M). Transcript and protein levels of the α and β subunits of the F_1F_0 -ATPase and the B subunit of the V-ATPase also increased with NaCl treatment. I have also shown that under the same conditions that normally cause induction of V-ATPase activity (up to 1 M NaCl in SC medium pH 4.5), V-ATPase mutant strains exhibit NaCl hypersensitivity compared to wild-type parent and complemented strains. This hypersensitivity is restricted to NaCl and LiCl, since no hypersensitivity is observed when the same strains are grown on sorbitol or hygromycin B. Since V-ATPase activity is required for NaCl tolerance, as well as Al tolerance as described in chapter 3, future testing of the energy balance hypothesis will involve measurement of Al and NaCl tolerance in transgenic plants.

4.5 Literature cited

- Anraku Y, Umemoto N, Hirata R and Ohya Y** (1992) Genetic and cell biological aspects of the yeast vacuolar H⁺-ATPase. *J Bioeng Biomembr* **24**: 395-405
- Apse MP, Aharon GS, Snedden WA and Blumwald E** (1999) Salt tolerance conferred by overexpression of a vacuolar Na⁺/H⁺ antiport in *Arabidopsis*. *Science* **285**: 1256-1258
- Bihler H, Slayman C and Bertl A** (1998) NSC1: A novel high current for cations in the plasma membrane of *Saccharomyces cerevisiae*. *FEBS Lett* **432**: 59-64
- Bradford M** (1976) A rapid and sensitive method for the quantification of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* **72**, 248-254
- Briskin DP, Leonard RT and Hodges TK** (1987) Isolation of the plasma membrane: Membrane markers and general principals. *In* L Packer, R Douce, eds, *Methods in Enzymology*. Vol 148. Academic Press, pp 542-558.
- Cohen P** (1989) The structure and function of protein phosphatases. *Annu Rev Biochem* **58**: 453-508
- Cunningham KW and Fink GR** (1996) Calcineurin inhibits VCX1-dependent H⁺-Ca²⁺ exchange and induces Ca²⁺-ATPases in *Saccharomyces cerevisiae*. *Mol Cell Biol* **16**: 2226-2237
- Gaxiola R, de Larrinoa IF, Villalba JM and Serrano R** (1992) A novel and conserved salt-induced protein is an important determinant of salt tolerance in yeast. *EMBO J* **11**: 3157-3164

- Gaxiola RA, Rao R, Sherman A, Grisafi P, Alper SL and Fink G (1999)** The *Arabidopsis thaliana* proton transporters, AtNhx1 and Avp1, can function in cation detoxification in yeast. *Proc Natl Acad Sci USA* **96**: 1480-1485
- Gaxiola RA, Li J, Undurraga S, Dang LM, Allen GJ, Alper SL and Fink GR (2001)** Drought- and salt-tolerant plants result from overexpression of the AVP1 H⁺-pump. *PNAS* **98**: 11444-11449
- Hamilton CA, Good AG and Taylor GJ (2001)** Induction of vacuolar ATPase and mitochondrial ATP synthase by aluminum in an aluminum-resistant cultivar of wheat. *Plant Physiol* **125**: 2068-2077
- Haro R, Banuelos MA, Quintero FJ, Rubio F and Rodriguez-Navarro A (1993)** Genetic basis of sodium exclusion and sodium tolerance in yeast. A model for plants. *Physiol Plant* **89**: 868-874
- Kane PM, Yamashiro CT, Wolczyk DF, Neff N, Goebel M and Stevens TH (1990)** Protein splicing converts the yeast TFP1 gene product to the 69-kD subunit of the vacuolar H⁺-adenosine triphosphatase. *Science* **250**: 651-657
- Lai-Zhang J, Xiao Y and Mueller DM (1999)** Epistatic interactions of deletion mutants in the genes encoding the F₁-ATPase in yeast *Saccharomyces cerevisiae*. *EMBO J* **18**: 58-64
- Mendizabal I, Rios G, Mulet JM, Serrano R and Larrinoa IF (1998)** Yeast putative transcription factors involved in salt tolerance. *FEBS Lett* **425**: 323-328

- Mendoza I, Rubio F, Rodriguez-Navarro A and Pardo J** (1994) The protein phosphatase calcineurin is essential for NaCl tolerance of *Saccharomyces cerevisiae*. *J Biol Chem* **269**: 8792-8796
- Mulet JM, Leube MP, Kron SJ, Rios G, Fink, GR and Serrano R** (1999) A novel mechanism of ion homeostasis and salt tolerance in yeast: The Hal4 and Hal5 protein kinases modulate the Trk1-Trk2 potassium transporter. *Mol Cell Biol* **19**: 3328-3337
- Nass R, Cunningham KW and Rao R** (1997) Intracellular sequestration of sodium by a novel Na⁺/H⁺ exchanger in yeast is enhanced by mutations in the plasma membrane H⁺-ATPase. *J Biol Chem* **272**: 26145-26152
- Nass R and Rao R** (1998) Novel localization of a Na⁺/H⁺ exchanger in a late endosomal compartment of yeast. *J Biol Chem* **273**: 21054-21060
- Perzov N, Padler-Karavani V, Nelson H and Nelson N** (2001) Features of V-ATPases that distinguish them from F-ATPases. *FEBS Lett* **504**: 223-228.
- Posas F, Camps M and Arino J** (1995) The PPZ protein phosphatases are important determinants of salt tolerance in yeast cells. *J Biol Chem* **270**: 13036-13041
- Prior C, Potier S, Souciet J-L and Sychrova H** (1996) Characterization of the *NHA1* gene encoding a Na⁺/H⁺ antiporter of the yeast *Saccharomyces cerevisiae*. *FEBS Lett* **387**: 89-93
- Rios G, Ferrando A and Serrano R** (1997) Mechanisms of salt tolerance conferred by overexpression of the *HAL1* gene in *Saccharomyces cerevisiae*. *Yeast* **13**: 515-528

Rothman JH and Stevens TH (1986) Protein sorting in yeast: The localization determinant of yeast vacuolar carboxypeptidase Y resides in the propeptide.

Cell **48**: 887-897

Sherman F (1991) Getting started with yeast. Methods Enzymol **194**: 3-21

Stemmer PM and Klee CB (1994) Dual calcium regulation of calcineurin by

calmodulin and calcineurin B. Biochemistry **33**: 6859-6866

Uchida E, Ohsumi Y and Anraku Y (1988) Purification of yeast vacuolar

membrane H⁺-ATPase and enzymological discrimination of three ATP-driven

proton pumps in *Saccharomyces cerevisiae*. Methods Enzymol **157**: 544-562

Yamashiro CT, Kane PM, Wolczyk DF, Preston RA and Stevens TH (1990) Role

of vacuolar acidification in protein sorting and zymogen activation: A genetic

analysis of the yeast vacuolar proton-translocating ATPase. Mol Cell Biol.

10: 3737-3749

5. Involvement of the Vacuolar H⁺-ATPase in Al and NaCl Tolerance in *Arabidopsis thaliana*

5.1 Introduction

Aluminum tolerance in *Arabidopsis thaliana* has been studied mainly from a genetic perspective. Mutagenesis has been used to generate Al-resistant and Al-sensitive mutants of *Arabidopsis*, some of which have been characterized. Larsen *et al.* (1997a) isolated 9 *als* (aluminum sensitive) mutants corresponding to 8 unique loci and 7 *alr* (aluminum resistant) mutants corresponding to 2 unique loci. One of the *als* mutants, *als-3*, caused inhibition of shoot development, leaf expansion and root growth in response to Al (Larsen *et al.*, 1997b). The genetic and physiological basis for this mutant phenotype has not been determined, and mutants representing the other 7 *als* loci have not yet been described in the literature. One of the *alr* loci, represented by the *alr-108*, *alr-128* and *alr-131* alleles, is associated with increased exudation of citrate, malate and pyruvate in response to Al, which may be caused by altered organic acid metabolism or transport (Larsen *et al.*, 1998). Increased exudation of organic anions is thought to contribute to detoxification of Al in the rhizosphere (Taylor, 1991). The other *alr* locus, represented by the *alr-104* allele, is associated with alkalization of the rhizosphere in response to Al. After Al

exposure, H^+ influx across the plasma membrane is increased 2-fold in the *alr-104* mutant compared to wild-type. The Al resistant phenotype is not observed in buffered solutions, suggesting that alkalinization of the rhizosphere is responsible for the Al resistance in this mutant (Degenhardt *et al.*, 1998).

Arabidopsis has also been used as model system to test whether Al-induced genes can confer Al-resistance when overexpressed. Ezaki *et al.* (2000) overexpressed 9 Al-induced genes from *Arabidopsis*, tobacco, wheat and yeast and tested the resistance of transgenic plants to Al, oxidative stress and other metals such as Cd, Cu, Na, and Zn. Of the 9 genes tested, 4 were associated with increased resistance to Al in transgenic lines of *Arabidopsis*. These were *NtPox* (peroxidase), *NtPARB* (glutathione S-transferase), *NtGDII* (GDP dissociation inhibitor) and *AtBCB* (blue copper binding protein), all of which have been implicated in response to oxidative stress. Overexpression of *NtPox*, *AtBCB* and *AtPox* also conferred resistance to diamide-induced oxidative stress, while overexpression of *NtPARB* conferred resistance to Cu and Na. A wheat cDNA encoding phosphatidylserine synthase, which conferred increased Al resistance when expressed in yeast, was also expressed in *Arabidopsis* (Delhaize *et al.*, 1999). Although expression of this enzyme is transcriptionally induced by Al in wheat root apices, the Al tolerance of transgenic *Arabidopsis* was not affected. These results suggest that there is not necessarily a link between Al-induction of genes and a role for their gene products in Al resistance, which may require appropriate regulation and localization of the Al-induced gene products or the presence of other enzymes.

Phenotype testing of yeast vacuolar ATPase mutants has shown that they are hypersensitive to Al and NaCl (Hamilton *et al.*, 2001a,b; Chapter Three; Chapter Four). This suggests that V-ATPase activity is required for Al and NaCl tolerance in yeast, as predicted by the energy balance hypothesis (Figure 3-1). To further test this hypothesis, I have examined the effect of altering V-ATPase expression in plants. Expression of a cDNA encoding the V-ATPase B subunit in the antisense orientation was used to downregulate V-ATPase expression in the model plant, *Arabidopsis thaliana*. This approach allowed me to directly test my hypothesis and further characterize the importance of the V-ATPase in mediating Al tolerance.

Although it would have been ideal to examine the effect of V-ATPase downregulation on Al tolerance in wheat, this was not practical due to the complexity of the wheat genome and the low efficiency of its *Agrobacterium*-mediated transformation (see for example Cheng *et al.*, 1997). *Arabidopsis* is commonly used as a model for plant biology because it has a comparatively small genome (125 Mb) which has been completely sequenced, it has been genetically well-characterized, a vast array of genetic resources are commercially available (<http://www.Arabidopsis.org>) and its small size allows simple manipulation of many plants on agar medium and in soil.

Due to the complex multisubunit structure of the V-ATPase (Stevens and Forgac, 1997; Chapter One), it was not feasible to construct transgenic plants overexpressing this enzyme. Therefore, the importance of V-ATPase activity in Al

tolerance was tested by expressing the cDNA encoding the B subunit in the antisense orientation. More specifically, I have used glucocorticoid-inducible expression of the V-ATPase B subunit cDNA in the antisense orientation to test the effect of downregulating V-ATPase activity on Al tolerance in *Arabidopsis*. If the energy balance hypothesis is correct, antisense expression should cause hypersensitivity to Al, as measured by inhibition of root elongation. If there is no discernable difference between transgenic and parent lines of *Arabidopsis*, then either the hypothesis can be rejected, which is unlikely based on our observations from wheat and yeast, or it must be modified to include other enzymes such as the vacuolar H⁺-pyrophosphatase, which may be able to counteract decreased V-ATPase activity.

5.2 Materials and Methods

5.2.1 Plant material and growth conditions

Seeds of *Arabidopsis thaliana* (all lines derived from ecotype C24) were surface sterilized in 20% (v/v) bleach, resuspended in 0.1% (w/v) agarose and plated on ½ MS medium (Murashige and Skoog basal medium, Sigma) containing 0.7% (w/v) phytigel, 2.5% (w/v) sucrose and 300 µg ml⁻¹ timentin (SmithKline Beecham, Oakville ON) to limit bacterial growth. Following cold stratification at 4°C for 2 days, seeds were transferred to a controlled environment chamber at 20°C with 16 h light and 8 h dark. Alternatively, seeds were sown in soil (Redi-Earth, Grace Horticultural products, Ajax ON) and grown under the same conditions described

above. Soil-grown plants were fertilized weekly with 5 mM KNO₃, 2 mM MgSO₄, 2 mM Ca(NO₃)₂, 0.5 mM Fe EDTA, 0.7 mM boric acid, 0.14 mM MnCl₂, 0.05 mM CuSO₄, 0.01 mM ZnSO₄, 0.007 mM NaMoO₄, 0.1 mM NaCl, 0.0001 mM CoCl₂, 2.5 mM potassium phosphate buffer pH 5.5 (<http://www.bch.msu.edu/pamgreen>).

5.2.2 Transformation constructs

A cDNA encoding the V-ATPase B subunit from *Arabidopsis* (*At57*) was provided by Dr M. Manolson (Hospital for Sick Children, Toronto ON, Manolson *et al.*, 1988). It was excised from the pBluescript cloning vector and ligated into the glucocorticoid-inducible binary vector, pTA7002 (Aoyama and Chua, 1997), in the antisense orientation using unique *Xho*I and *Spe*I restriction enzyme sites. The presence of the desired insert in the resulting construct, pTA-At57a, was confirmed by sequencing. The pTA-At57a construct was transformed into *Agrobacterium tumefaciens* GV3101 by electroporation (T=2.5 kV, R=129 Ω, S=1.44 kV, BioRad Gene Pulser) and the presence of the construct confirmed by PCR. Primers used were C6 (5'-CACGCAACATCTTCCAGTCG-3'), at the 3' end of the transgene, and P8 (5'-ATTTGGAGAGGAACACGCTGA-3'), downstream of the *Spe*I cloning site.

5.2.3 *Arabidopsis* transformation

The floral dip method of *Arabidopsis* transformation (Clough and Bent, 1998) was used, with minor modifications. Seeds of *Arabidopsis thaliana* ecotype C24 were sown in soil in 15 cm diameter pots as described above. Following germination, plants were thinned to 10 per pot and grown until emergence of primary inflorescences. The primary inflorescences were removed and secondary

inflorescences allowed to reach approximately 5-15 cm tall with numerous flowers and buds. *Agrobacterium tumefaciens* strain GV3101 (Koncz and Schell, 1986) carrying the binary vector of interest was grown overnight at 30°C in 500 ml YEP medium [10% (w/v) yeast extract, 10% (w/v) bacto-peptone, 5% (w/v) NaCl] containing 50 µg ml⁻¹ kanamycin. The cell pellet was harvested by centrifugation and resuspended in 10% (w/v) sucrose, 0.05% (v/v) Silwet L-77 (Osi Specialties, Danbury CT) to a final OD₆₀₀ of 0.8. The inflorescences were dipped into the *Agrobacterium* suspension for two 30 s intervals with gentle agitation. The floral dip was repeated one week later.

5.2.4 Selection and confirmation of transformants

T₁ seeds harvested from floral-dipped plants were sown on ½ MS medium containing 25 µg ml⁻¹ hygromycin B (to select for transformants) and 300 µg ml⁻¹ timentin as described above. Eight hygromycin-resistant seedlings (T₁) were transferred from tissue culture to soil after 4-6 weeks and matured in Arasystem 360 selfing tubes (BetaTech, Gent Belgium). T₂ seeds were harvested from individual T₁ plants to be used for experimental work. The presence of the transgene was tested by growth on medium containing hygromycin and by PCR, using the REDExtract-N-Amp plant PCR kit (Sigma) and primers specific to hygromycin phosphotransferase, P16 (5'-CATGTAGGAGGGCGTGGATA-3') and P17 (5'-CCTCCTCCATAACAAGCCAAC-3'), and specific to the transgene, C6 (5'-CACGCAACATCTTCCAGTCG-3') and P8 (5'-ATTTGGAGAGGAACACGCTGA-3').

5.2.5 *Transgene Copy Number Determination*

For each of the 8 transgenic lines, 24 three-week old T₂ seedlings grown on ½ MS medium were transferred to ½ MS medium containing 50 µg mL⁻¹ hygromycin B to determine the ratio of hygromycin-resistant to hygromycin-sensitive individuals. Plants were grown on hygromycin-containing medium for one week, when toxicity symptoms (chlorosis, death) were observed in control (C24) plants.

5.2.6 *Root elongation measurements*

Three-week old T₂ seedlings were transferred to ½ MS medium (1.4% phytigel) containing 0 or 30 µM dexamethasone (DEX) and either 100-500 µM AlCl₃, or 50-100 mM NaCl. The end of the longest root was marked on the plate with a marker and the plates were incubated vertically for 1 week (Ellis *et al.*, 1999). Root elongation from the starting mark was then measured. After the experiment, plants were transferred to ½ MS medium containing 50 µg mL⁻¹ hygromycin B to select for transgenic plants. Data points for nontransgenic plants were not included in the analysis. Plates containing 6 plants were used for each treatment and the experiment was performed three times.

5.3 **Results**

5.3.1 *The Effect of Al on Root Elongation in Wild-type Arabidopsis*

In order to quantify the effect of Al on wild-type *Arabidopsis*, the effect Al on root elongation was investigated. Three-week old seedlings of *Arabidopsis thaliana* ecotype C24 were transferred to ½ MS medium (pH 4.3) containing 0-1000 µM AlCl₃

and incubated vertically for one week. In the absence of Al, root length increased by 35.2 +/- 0.8 mm. Inhibition of root elongation by Al was observed at all concentrations tested, with virtually no elongation (0.8 +/- 0.1 mm) at 1000 μM AlCl_3 (Fig 5-1). In addition to inhibition of root elongation, concentrations of Al 500 μM or greater caused proliferation of lateral roots, which grew into the air. Based on these observations, Al concentrations of 100 μM and 500 μM were chosen for phenotype testing of transgenic plants. If plants expressing the B subunit of the V-ATPase in the antisense orientation are hypersensitive to Al, then the effect should be apparent at these concentrations.

5.3.2 *The Effect of NaCl on Root Elongation in Wild-type Arabidopsis*

The effect of NaCl on wild-type *Arabidopsis* was also investigated. Three-week old seedlings were transferred to $\frac{1}{2}$ MS (pH 4.3) containing 0-300 mM NaCl and incubated vertically for one week (Fig 5-2). In the absence of NaCl, root length increased by 31.6 +/- 1.2 mm. Inhibition of root elongation was observed at all concentrations of NaCl tested, with complete inhibition at 200 and 300 mM (0 mm). Plants grown on $\frac{1}{2}$ MS containing 200 mM NaCl showed minor chlorosis and club-shaped roots. At 300 mM NaCl, plants showed extensive chlorosis (data not shown). Since NaCl toxicity symptoms were severe at 200-300 mM NaCl, concentrations of 50 mM and 100 mM were chosen for phenotype testing of transgenic plants.

Fig 5-1. The effect of Al on root elongation in wild-type *Arabidopsis* (ecotype C24). Three-week old seedlings were grown vertically on ½ MS medium (pH 4.3) containing 0-1000 μM AlCl_3 for one week. Results shown are means \pm SE (n=6) and the results shown are representative of two independent experiments.





Fig 5-2. The effect of NaCl on root elongation in wild-type *Arabidopsis* (ecotype C24). Three-week old seedlings were grown vertically on ½ MS medium (pH 4.3) containing 0-300 mM NaCl for one week. Results shown are means +/- SE (n=6) and the results shown are representative of two independent experiments.



5.3.3 *Production of Transgenic Arabidopsis Expressing the V-ATPase B Subunit in the Antisense Orientation*

Since the V-ATPase is important in plant growth and development, a glucocorticoid-inducible promoter (Aoyama and Chua, 1997) was used to drive expression of the antisense RNA. The T-DNA of the binary vector pTA7002 (Fig 5-3) constitutively expresses a fusion protein (GVG) from the constitutive cauliflower mosaic virus 35S promoter. The fusion protein consists of the GAL4 DNA binding domain, the herpes viral protein VP16 transactivating domain and the receptor domain of the rat glucocorticoid receptor. The synthetic glucocorticoid, dexamethasone (DEX), binds to the GVG transcription factor and this complex activates transcription of the antisense RNA by binding to six copies of the GAL4 upstream activating sequence (6xUAS_{GAL}) located upstream of the transgene.

The antisense construct (Fig 5-3) was transformed into *Arabidopsis thaliana* ecotype C24 using the floral dip method (Clough and Bent, 1998). T₁ plants, selected by growth on medium containing hygromycin, were selfed and the resulting T₂ seeds were used for all experimental manipulations. To confirm that the T₂ plants were in fact transgenic, the presence of the hygromycin phosphotransferase gene and the transgene were determined by PCR. Genomic DNA was isolated from leaf discs of putative transgenic plants or parental plants for use in PCR reactions. The products of the PCR reactions were separated by agarose gel electrophoresis and visualized by staining with ethidium bromide (Fig 5-4). The 650 bp hygromycin phosphotransferase PCR product was present in all putative transgenic plants and

Fig 5-3. The T-DNA construct designed for glucocorticoid-inducible expression of the B subunit of the vacuolar ATPase (At57) in the antisense orientation (indicated by arrow). The NOS promoter controls expression of the selectable marker, hygromycin phosphotransferase (HPT). The constitutive cauliflower mosaic virus 35S promoter expresses a fusion protein (GVG) consisting of the GAL4 DNA binding domain, the herpes viral protein VP16 transactivating domain and the receptor domain of the rat glucocorticoid receptor, followed by the Rbc termination sequence. The synthetic glucocorticoid, dexamethasone (DEX), binds to the GVG transcription factor and this complex activates transcription of the antisense RNA by binding to six copies of the GAL4 upstream activating sequence (6xUAS_{GAL}) located upstream of the CMV 35S TATA box and the transgene. LB and RB denote the left and right borders of the T-DNA, respectively. The position of each component is given in bp. The diagram is not to scale.






Fig 5-4. Analysis of transgenic lines of *Arabidopsis* by PCR amplification of (A) a 650 bp fragment of hygromycin phosphotransferase and (B) a 598 bp fragment specific to the transgene. The positive control (+) was the plasmid used for transformation, pTA-At57a. Lane 1 is the parental line, C24, and lane 2 is the binary vector with no insert. Lanes 3-10 are independent transgenic lines 32, 33, 34, 36, 37, 38, 39 and 40, respectively. The results shown are representative of two independent experiments.



absent in the parent, C24 (Fig 5-4A). The 598 bp transgene-specific PCR product was present in all eight antisense lines tested and absent in the parent and the vector controls (Fig 5-4B). These results confirmed that all of the antisense lines contained the construct in the correct orientation and that the transgene was absent in the parent and vector control lines.

The copy number of T-DNA inserts was determined by calculating the ratio of hygromycin-resistant to hygromycin-sensitive plants in the T₂ generation. Twenty-four T₂ seedlings from each independent transgenic line were transferred from ½ MS medium to ½ MS medium containing 50 µg mL⁻¹ hygromycin B, which is toxic to parental plants. After 1 week, plants were scored as resistant or sensitive to hygromycin (Table 5-1). Hygromycin resistant plants were healthy and showed no apparent effects, while hygromycin sensitive plants were chlorotic and had ceased growth. The number of multiple T-DNA insertions in *Arabidopsis* was low. Of the eight lines tested here, four (33, 34, 38, 40) showed a 3:1 ratio, indicating that a single insertion event has occurred. Line 32 showed a 5:1 ratio, which also likely indicates that a single insertion event has occurred. Two lines (36 and 39) showed other ratios (11:1 and 24:0, respectively), indicating that multiple insertions have occurred. In line 37, which showed a ratio of approximately 1:1, homozygous transgenic individuals and nontransgenic individuals were likely killed by hygromycin. This may be the result of silencing of the hygromycin phosphotransferase gene. Transgenic lines 33, 34 and 40 were selected as the focus of phenotypic testing since they contained single copies of the T-DNA insert.

Table 5-1. Determination of copy number in transgenic *Arabidopsis thaliana* as measured by segregation of hygromycin resistance in T₂ progeny. Four week-old plants (24 from each line) were transferred to ½ MS medium containing 50 µg mL⁻¹ hygromycin B for one week and scored as hygromycin resistant (no symptoms) or hygromycin sensitive (chlorotic, no growth).

<i>Arabidopsis</i> Line	Hygromycin-Resistant T ₂ Progeny	Hygromycin-Sensitive T ₂ Progeny	Ratio	Copy Number
C24 (parent)	0	24	0:24	0
32	20	4	5:1	1*
33	18	6	3:1	1
34	18	6	3:1	1
36	22	2	11:1	2
37	13	11	1:1	1 (silenced)
38	18	6	3:1	1
39	24	0	24:0	multiple
40	18	6	3:1	1

*Since a 5:1 ratio is nearer to 3:1 than 15:1 (expected for a double insertion event), this line is presumed to have a single insertion of the hygromycin phosphotransferase gene.

5.3.4 *Phenotype Testing of Transgenic Arabidopsis*

To examine the effect of downregulating V-ATPase expression on Al and NaCl tolerance, root elongation was measured. As demonstrated in Fig 5-1 and Fig 5-2, root elongation in three-week old *Arabidopsis* seedlings showed a dose-dependent response to both Al and NaCl. Aluminum concentrations of 100 μM and 500 μM and NaCl concentrations of 50 mM and 100 mM were chosen as exposure conditions based on these results. Dexamethasone at 30 μM was also included as a variable since it was required to induce transgene expression.

As shown in Fig 5-5, 5-7 and Fig 5-8, transgenic lines of *Arabidopsis* showed dexamethasone-dependent hypersensitivity to Al. The selected transgenic lines (33, 34 and 40) exhibited no dexamethasone toxicity, simplifying comparison between treatments. Some lines (32 and 37) showed significant inhibition of root elongation in the presence of dexamethasone (21% and 30%, respectively), so they were not included in the present discussion. At 100 μM Al, transgenic line 33 was inhibited a further 34% (from 20.8 mm to 13.8 mm) by the addition of dexamethasone, while lines 34 and 40 were inhibited a further 32% (from 21.3 mm to 14.5 mm) and 12% (from 14.5 mm to 12.8 mm), respectively. At 500 μM Al, addition of dexamethasone caused a further 30% decrease in root elongation in line 33 (from 13.8 mm to 9.7 mm) and further decreases of 51% (from 14.2 mm to 7.0 mm) and 34% (from 10.5 mm to 7.0 mm), respectively, in lines 34 and 40. In contrast, root elongation in C24 was inhibited by only 5% with the addition of dexamethasone (from 14.5 mm to 13.8 mm)

at 100 μM Al and not at all at 500 μM Al. Analysis of variance determined that the lines examined (C24 compared to transgenic), the presence of Al and dexamethasone caused significant effects ($P < 0.0001$). There were also significant interactions ($P < 0.01$) between line x aluminum and line x dexamethasone (Table 5-2A). In summary, the transgenic lines exhibited dexamethasone-dependent Al hypersensitivity.

Although the phenotypic data suggests that the transgene is being expressed, levels of the sense and antisense transcripts must be measured to confirm that the phenotype is caused by downregulation of V-ATPase expression. These experiments are in progress. Measurement of V-ATPase activity in dexamethasone-treated transgenic plants would provide additional evidence that the antisense strategy is effective.

Compared to the results observed for Al (Fig 5-5, Fig 5-7, Fig 5-8), the effect of downregulating V-ATPase expression on NaCl tolerance was variable (Fig 5-6, Fig 5-7, Fig 5-8). At 50 mM NaCl, addition of dexamethasone caused a 1% increase in root elongation (from 21.7 mm to 22.0 mm) in line 33, and decreases of 39% (from 23.8 mm to 14.5 mm) in line 34 and 3% (from 17.5 mm to 17.3 mm) in line 40. In C24, addition of dexamethasone at 50 mM NaCl caused root elongation to decrease by 26% (from 18.8 mm to 14.0 mm). At 100 mM NaCl, dexamethasone caused no additional decrease in root elongation in C24. In the transgenic lines, addition of dexamethasone caused decreases of 10% (from 17.2 mm to 15.5 mm) in line 33, 38% (from 15.7 mm to 9.8 mm) in line 34 and 2% (from 10.7 mm to 10.5 mm) in line 40.

Fig 5-5. The effect of transgene expression on Al resistance in *Arabidopsis*, measured by root elongation in the presence and absence of 30 μM dexamethasone (DEX). Three-week old T_2 seedlings were transferred to $\frac{1}{2}$ MS medium (pH 4.3) containing 0 or 30 μM DEX and 0, 100 or 500 μM AlCl_3 and root elongation was measured after one week. Results shown are means \pm SE (n=6) and are representative of two independent experiments.




Fig 5-6. The effect of transgene expression on NaCl resistance in *Arabidopsis*, measured by root elongation in the presence and absence of 30 μ M dexamethasone (DEX). Three-week old T₂ seedlings were transferred to ½ MS medium (pH 4.3) containing 0 or 30 μ M DEX and 0, 50 or 100 mM NaCl and root elongation was measured after one week. Results shown are means \pm SE (n=6) and are representative of two independent experiments.




Fig 5-7. The effect of dexamethasone application on Al and NaCl resistance in wild-type *Arabidopsis thaliana* (ecotype C24). Three-week old T₂ seedlings were transferred to ½ MS medium (pH 4.3) containing 0 or 30 µM DEX and 0, 100 or 500 µM AlCl₃, and 50 or 100 mM NaCl for one week. Results shown are representative of two independent experiments.




Fig 5-8. The effect of transgene expression on Al and NaCl resistance in transgenic *Arabidopsis thaliana* (line 34). Three-week old T₂ seedlings were transferred to ½ MS medium (pH 4.3) containing 0 or 30 µM DEX and 0, 100 or 500 µM AlCl₃, and 50 or 100 mM NaCl for one week. Results shown are representative of two independent experiments.




Table 5-2. Three factor analysis of variance of root elongation measurements of transgenic and parent (C24) lines in the presence of dexamethasone and/or (A) Al (B) NaCl. P values < 0.05 are considered significant, as indicated by asterisks (*).

	Variable	P Value
(A)	line	$3.7 \times 10^{-5}*$
	aluminum	$< 1 \times 10^{-6}*$
	dexamethasone	$1.7 \times 10^{-5}*$
	line x aluminum	$1.4 \times 10^{-3}*$
	line x dexamethasone	$7.0 \times 10^{-3}*$
	aluminum x dexamethasone	2.0×10^{-1}
	line x aluminum x dexamethasone	6.0×10^{-1}
(B)	line	$< 1 \times 10^{-6}*$
	salt	$< 1 \times 10^{-6}*$
	dexamethasone	$1.7 \times 10^{-2}*$
	line x salt	$1.4 \times 10^{-3}*$
	line x dexamethasone	3.2×10^{-1}
	salt x dexamethasone	9.5×10^{-1}
	line x salt x dexamethasone	1.0×10^{-1}

Analysis of variance suggests that the effect of transgene expression on NaCl tolerance is more variable than the effect on Al tolerance. The lines examined (C24 compared to transgenic) and the presence of NaCl had significant effects ($P < 1 \times 10^{-6}$) on root elongation. Dexamethasone had a weaker, but significant, effect ($P < 0.05$). There was a significant interaction ($P < 0.01$) between line x NaCl, but no significant interaction between line x dexamethasone (Table 5-2B). Although not as convincing as the results observed for Al, the results from line 34 indicate that downregulation of V-ATPase expression could potentially cause NaCl hypersensitivity.

5.4 Discussion

Previous investigations of Al tolerance using transgenic technology in *Arabidopsis* have overexpressed many Al-induced genes in an effort to cause a phenotypic change (Ezaki *et al.*, 2000; Richards *et al.*, 1998). The moderate success of this approach indicates that the complex interactions involved in Al tolerance must be taken into consideration when designing transgenic experiments. The approach described here is hypothesis-driven, based on previous work in wheat (Hamilton *et al.*, 2001c; Chapter Two) and yeast (Hamilton *et al.*, 2001a, 2001b; Chapter Three, Chapter Four). Results from my previous work indicated that V-ATPase expression is induced by Al stress in an Al-resistant cultivar of wheat (Hamilton *et al.*, 2001c; Chapter Two). The V-ATPase is also required for Al and NaCl resistance in yeast (Hamilton *et al.*, 2001a, 2001b; Chapter Three, Chapter Four). Based on these

results, I predicted that downregulation of V-ATPase activity would cause Al and NaCl hypersensitivity in transgenic *Arabidopsis*.

According to the energy balance hypothesis (Fig 3-1), downregulating V-ATPase activity should cause hypersensitivity to Al. The results presented here are consistent with this hypothesis, since transgenic plants show dexamethasone-dependent Al hypersensitivity. It is unknown whether this hypersensitivity is caused by impairment of a pH homeostatic mechanism or by impaired Al compartmentation in the vacuole. Although it is tempting to speculate about Al sequestration, no conclusions can be drawn in the absence of data on Al localization. The V-ATPase may have a role in pH homeostasis during Al stress, similar to the *alr104* mutant characterized by Degenhardt *et al.* (1998). This mutant exhibits enhanced H⁺ influx across the plasma membrane in response to Al stress, resulting in alkalization of the rhizosphere. Increased activity of the V-ATPase could result in alkalization of the cytoplasm, which would reduce the solubility and/or toxicity of intracellular Al, and would also prevent acidosis caused by impairment of the plasma membrane H⁺-ATPase by Al (Matsumoto, 1988; Sasaki *et al.*, 1995; Widell *et al.*, 1994; Hamilton *et al.*, 2001a).

Unlike their response to Al, the response of the transgenic plants to NaCl was variable. Overexpression of the Na⁺/H⁺ antiporter (Apse *et al.*, 1999) or the vacuolar H⁺-pyrophosphatase (Gaxiola *et al.*, 2001) conferred increased NaCl tolerance in transgenic *Arabidopsis*. This suggests that maintenance of the tonoplast membrane potential contributes to NaCl tolerance by providing the necessary proton motive

force for sequestration of Na^+ in the vacuole. It is possible that the conditions used here were not optimal for observing a phenotypic difference, since Gaxiola *et al.* (2001) used a soil system and watered their transgenic plants with saline solutions. These conditions are different from the tissue culture conditions used here, so it would be interesting to test our transgenic plants in soil using their regimen. It would also be a powerful test of the energy balance hypothesis to use plants overexpressing the vacuolar H^+ -pyrophosphatase (Gaxiola *et al.*, 2001) in the assays described here, since one would predict that they would be tolerant to Al as well as to NaCl.

Since the vacuolar H^+ -ATPase is required for growth and development, the glucocorticoid-responsive transcriptional induction system was used to control antisense expression. This system has the advantages of localized, inducible expression of transgenes at variable levels. It has been used successfully to control expression of the luciferase gene (Aoyama and Chua, 1997) as well as many other genes (for example McNellis *et al.*, 1998; Lally *et al.*, 2001; Okamoto *et al.*, 2001; Pautot *et al.*, 2001) in transgenic *Arabidopsis*. However, this system has limitations. In certain transgenic lines, the glucocorticoid-inducible system can cause severe growth defects and induce defense-related genes in transgenic *Arabidopsis* (Kang *et al.*, 1999). This was not observed with the transgenic lines used here. This suggests that the transgenic lines used, as well as concentration and means of application (liquid medium, agar or spraying) of dexamethasone may be important variables to be considered when analyzing data from this system. It is also essential to include

dexamethasone treatment as an experimental control to rule out nonspecific dexamethasone-induced effects.

In summary, this chapter describes the design, construction and phenotypic testing of transgenic *Arabidopsis* expressing a cDNA encoding the B subunit of the V-ATPase in the antisense orientation. Induction of transgene expression by dexamethasone causes hypersensitivity to Al, as predicted by the energy balance hypothesis. Interestingly, transgenic plants showed a variable phenotype with respect to NaCl tolerance. Further investigation will need to determine why certain lines exhibit NaCl hypersensitivity while others do not so that the role of the V-ATPase in NaCl tolerance can be determined.

5.5 Literature Cited

- Aoyama T and Chua N-H** (1997) A glucocorticoid-inducible transcriptional induction system in transgenic plants. *Plant J* **11**: 605-612
- Apse MP, Aharon GS, Snedden WA and Blumwald E** (1999) Salt tolerance conferred by overexpression of a vacuolar Na⁺/H⁺ antiport in *Arabidopsis*. *Science* **285**: 1256-1258
- Cheng M, Fry JE, Pang S, Zhou H, Hironaka CM, Duncan DR, Conner TW and Wan Y** (1997) Genetic transformation of wheat mediated by *Agrobacterium tumefaciens*. *Plant Physiol* **115**: 971-980
- Clough SJ and Bent AF** (1998) Floral dip: A simplified method for *Agrobacterium*-mediated transformation of *Arabidopsis thaliana*. *Plant J* **16**: 735-743
- Degenhardt J, Larsen PB, Howell SH and Kochian LV** (1998) Aluminum resistance in the *Arabidopsis* mutant *alr-104* is caused by an aluminum-induced increase in rhizosphere pH. *Plant Physiol* **117**: 19-27
- Delhaize E, Hebb DM, Richards KD, Lin J-M, Ryan PR and Gardner RC** (1999) Cloning and expression of a wheat (*Triticum aestivum* L.) phosphatidylserine synthase cDNA. *J Biol Chem* **274**: 7082-7088
- Ellis MH, Dennis ES and Peacock WJ** (1999) *Arabidopsis* roots and shoots have different mechanisms for hypoxic stress tolerance. *Plant Physiol* **119**: 57-64

- Ezaki B, Gardner RC, Ezaki Y and Matsumoto H** (2000) Expression of aluminum-induced genes in transgenic *Arabidopsis* plants can ameliorate aluminum stress and/or oxidative stress. *Plant Physiol* **122**: 657-665
- de la Fuente JM, Ramirez-Rodriguez V, Cabrera-Ponce JL and Herrera-Estrella L** (1997) Aluminum tolerance in transgenic plants by alteration of citrate synthesis. *Science* **276**: 1566-1568
- Gaxiola RA, Li J, Undurraga S, Dang LM, Allen GJ, Alper SL and Fink GR** (2001) Drought- and salt-tolerant plants result from overexpression of the AVP1 H⁺-pump. *Proc Nat Acad Sci* **98**: 11444-11449
- Hamilton CA, Good AG and Taylor GJ** (2001a) Vacuolar H⁺-ATPase, but not mitochondrial ATP synthase, is required for aluminum tolerance in *Saccharomyces cerevisiae*. *FEMS Microbiol Lett* **205**: 231-236
- Hamilton CA, Taylor GJ and Good AG** (2001b) Vacuolar H⁺-ATPase, but not mitochondrial ATP synthase, is required for NaCl tolerance in *Saccharomyces cerevisiae*. *FEMS Microbiol Lett*. (In Press)
- Hamilton CA, Good AG and Taylor GJ** (2001c) Induction of vacuolar ATPase and mitochondrial ATP synthase by aluminum in an aluminum-resistant cultivar of wheat. *Plant Physiol* **125**: 2068-2077
- Kang H-G, Fang Y and Singh KB** (1999) A glucocorticoid-inducible transcription system causes severe growth defects in *Arabidopsis* and induces defense-related genes. *Plant J* **20**: 127-133

- Koncz C and Schell J** (1986) The promoter of T_L-DNA gene 5 controls the tissue-specific expression of chimaeric genes carried by a novel type of *Agrobacterium* binary vector. *Mol Gen Genet* **204**: 383-396
- Lally D, Ingmire P, Tong HY and He ZH** (2001) Antisense expression of a cell wall-associated protein kinase, WAK4, inhibits cell elongation and alters morphology. *Plant Cell* **13**: 1317
- Larsen PB, Tai C-Y, Howell SH and Kochian LV** (1996) *Arabidopsis* mutants with increased sensitivity to aluminum. *Plant Physiol* **110**: 743-751
- Larsen PB, Stenzler LM, Tai C-Y, Degenhardt J, Howell SH and Kochian LV** (1997a) Molecular and physiological analysis of *Arabidopsis* mutants exhibiting altered sensitivities to aluminum. *Plant Soil* **192**: 3-7
- Larsen PB, Kochian LV and Howell SH** (1997b) Al inhibits both shoot development and root growth in *als-3*, an Al-sensitive *Arabidopsis* mutant. *Plant Physiol* **114**: 1207-1214
- Larsen PB, Degenhardt J, Tai C-Y, Stenzler LM, Howell SH and Kochian LV** (1998) Aluminum-resistant *Arabidopsis* mutants that exhibit altered patterns of aluminum accumulation and organic acid release from roots. *Plant Physiol* **117**: 9-18
- Manolson MF, Ouellette BFF, Filion M and Poole RJ** (1988) cDNA sequence and homologies of the "57-kDa" nucleotide-binding subunit of the vacuolar ATPase from *Arabidopsis*. *J Biol Chem* **263**: 17987-17994

Matsumoto H (1988) Inhibition of proton transport activity of microsomal membrane vesicles of barley roots by aluminium. *Soil Sci Plant Nutr* **34**: 499-526

McNellis TW, Mudgett MB, Li K, Aoyama T, Horvath D, Chua NH and Staskawicz BJ (1998) Glucocorticoid-inducible expression of a bacterial avirulence gene in transgenic *Arabidopsis* induces hypersensitive cell death. *Plant J* **14**: 247-257

Okamoto H, Matsui M and Deng XW (2001) Overexpression of the heterotrimeric G-protein alpha-subunit enhances phytochrome-mediated inhibition of hypocotyl elongation in *Arabidopsis*. *Plant Cell* **13**: 1639-1652

Pautot V, Dockx J, Hamant O, Kronenberger J, Grandjean O, Jublot D and Traas J (2001) KNAT2: Evidence for a link between knotted-like genes and carpel development. *Plant Cell* **13**: 1719-1734

Richards KD, Schott EJ, Sharma YK, Davis KR and Gardner RC (1998) Aluminum induces oxidative stress genes in *Arabidopsis thaliana*. *Plant Physiol* **116**: 409-418

Sasaki M, Kasai M, Yamamoto Y and Matsumoto H (1995) Involvement of plasma membrane potential in the tolerance mechanism of plant roots to aluminum toxicity. *In* RA Date *et al.*, eds. *Plant-Soil Interactions at Low pH. Proceedings of the Second International Symposium on Plant-Soil Interactions at Low pH, 24-29 June 1990, Beckley, West Virginia, USA*. Kluwer Academic Publishers, Dordrecht, pp 285-290

Widell S, Asp H and Jensen P (1994) Activities of plasma membrane-bound enzymes isolated from roots of spruce (*Picea abies*) grown in the presence of aluminium. *Physiol Plant* **92**: 459-466

6. Concluding Discussion

In this thesis, the role of the vacuolar ATPase in aluminum resistance has been examined in wheat (*Triticum aestivum* L.), yeast (*Saccharomyces cerevisiae*) and thale cress (*Arabidopsis thaliana*). Earlier research from our laboratory identified two 51 kD, Al-induced proteins from an Al-resistant cultivar of wheat (PT741), which were newly synthesized upon administration of Al stress (Basu *et al.*, 1994). Induction of these proteins segregated with Al-resistance in a segregating population derived from a cross between Al-resistant PT741 and Al-sensitive Katepwa (Taylor *et al.*, 1997). These observations indicated that Al-induction of the 51 kD proteins may be an adaptive response involved in mediating Al resistance.

The 51 kD, Al-induced proteins were purified, sequenced and found to be subunits of the vacuolar H⁺-ATPase (V-ATPase) and the mitochondrial F₁F₀-ATPase. Their identities were further confirmed by measuring ATPase activity in Al treated roots of PT741, where Al induced both V-ATPase and F₁F₀-ATPase activities in a dose-dependent manner. Levels of the transcript encoding the B subunit of the V-ATPase also increased with Al treatment, although levels of the transcript encoding the α subunit of the F₁F₀-ATPase were unaffected by Al at the levels tested. Interestingly, increased V-ATPase and F₁F₀-ATPase activities were observed only in PT741 and not in the other Al-resistant and Al-sensitive cultivars of wheat tested. In contrast, inhibition of the plasma membrane H⁺-ATPase was observed in all cultivars. This suggests that induction of the V-ATPase and F₁F₀-ATPase, thought to be an

adaptive response to Al stress, is specific to this cultivar. This may prove useful in future breeding programs for Al resistance.

Based on the observation that the V-ATPase and the F_1F_0 -ATPase are induced by Al, a testable hypothesis describing a possible role for these enzymes in mediating Al resistance was developed (Fig 3-1). Briefly, this hypothesis predicts that the V-ATPase, using ATP generated by the F_1F_0 -ATPase, is responsible for preventing cytoplasmic acidosis caused by Al-inhibition of the plasma membrane H^+ -ATPase. In addition to protecting sensitive cytoplasmic proteins from pH changes, prevention of cytoplasmic acidosis is also required to allow continued tonoplast transmembrane transport. Whether this transport involves sequestration of an unknown Al species remains to be determined. Considering recent advances in analysis of Al uptake in single cells, it may be possible to address this question directly in the near future.

It is not presently possible to manipulate gene expression in monocots using transgenic technology without highly labour-intensive techniques. Yeast, however, is amenable to relatively simple genetic manipulation, so it was used as a model system to test my hypothesis. Yeast strains with knockout mutations in the genes encoding the A and B subunits of the V-ATPase, but not those with mutations in the genes encoding the α and β subunits of the F_1F_0 -ATPase, exhibited dramatic hypersensitivity to Al in comparison to the isogenic wild-type parent and complemented strains. This provided support for the hypothesis that V-ATPase activity is involved in mediating Al resistance, even in a different species. This hypothesis could also be tested using yeast strains overexpressing the vacuolar H^+ -

pyrophosphatase (Gaxiola *et al.*, 1999). Overexpression of this single protein is analogous to overexpression of the multisubunit V-ATPase. According to my hypothesis, yeast strains overexpressing this enzyme should exhibit enhanced Al resistance. Phenotype testing of this strain is in progress.

Characterization of wild-type yeast revealed that V-ATPase activity, transcript and protein levels were unchanged by the addition of Al, in contrast to the Al-induction of V-ATPase activity and transcript levels observed in wheat.

Mitochondrial F_1F_0 -ATPase activity was induced by Al, consistent with observations from wheat. The differences in V-ATPase expression in wheat and yeast are reminiscent of the V-ATPase activity differences between Al-resistant PT741 and Al-sensitive cultivars of wheat. This may lend further support to the hypothesis that V-ATPase activity is involved in Al resistance.

Genetic studies have shown that the yeast V-ATPase contributes to NaCl tolerance, presumably by establishing the tonoplast transmembrane potential required for sequestration of Na^+ in the vacuole. This has been convincingly demonstrated by overexpressing the *Arabidopsis* vacuolar H^+ -pyrophosphatase in yeast (Gaxiola *et al.*, 1999) and *Arabidopsis* (Gaxiola *et al.*, 2001), where it confers increased NaCl tolerance. My observation that yeast strains with knockout mutations in the genes encoding the A and B subunits of the V-ATPase are hypersensitive to NaCl lends further support to the argument that vacuole function is required for NaCl tolerance.

Although NaCl tolerance in yeast has been intensively investigated, the response of ATPases to NaCl in wild-type yeast had not been previously

characterized. Therefore, the scope of my research was expanded to include analysis of ATPase expression in response to NaCl stress with the hope that parallels could be drawn with Al stress. In contrast to the constant levels observed with increasing Al concentrations, V-ATPase activity, protein and transcript levels increased with the addition of NaCl. In another interesting contrast, induction of the F_1F_0 -ATPase was also observed, but to a lesser extent. The reason that F_1F_0 -ATPase expression is more highly induced by Al in yeast is unknown, but it is possible that additional energy is required to combat Al stress. In the case of NaCl, energy is required because to sequester Na^+ in the vacuole and to export Na^+ from the cell by active transport across the plasma membrane (Marquez and Serrano, 1996). Although NaCl induces F_1F_0 -ATPase expression, yeast strains with knockout mutations in the genes encoding the α and β subunits of the F_1F_0 -ATPase were not hypersensitive to NaCl. This suggests that for NaCl, like Al, the V-ATPase is more critical in stress resistance. The genetic approach used here may point towards a new direction in Al stress research. A program of systematically studying components thought to be involved in Al resistance using yeast knockout mutants has been proposed as a means of dissecting this complex stress response. This approach will allow us to investigate the involvement of individual components or pathways in Al resistance in a simple model system.

Since experiments performed with yeast supported my hypothesis that the V-ATPase is required for Al and NaCl tolerance, the next step was to test whether or not altering the expression of the V-ATPase could affect Al and NaCl tolerance in plants.

Arabidopsis thaliana was selected as a model plant for these experiments because it can be easily transformed, unlike wheat (Cheng *et al.*, 1997). Inhibition of V-ATPase expression was achieved using a glucocorticoid-inducible antisense approach, which has the advantage of temporal and spatial control of antisense RNA expression (Aoyama and Chua, 1997) and because phenotypic screening is relatively simple. Although antisense inhibition of V-ATPase expression was not measured directly, T₂ transgenic lines exhibited glucocorticoid-dependent Al hypersensitivity. Ongoing experiments are investigating V-ATPase expression by measuring enzyme activity, antisense and sense RNA levels, in dexamethasone-treated roots. Some lines were also hypersensitive to NaCl, although this aspect of their phenotype was not as clear. These results could be supplemented by testing the phenotype of transgenic *Arabidopsis* expressing the vacuolar H⁺-pyrophosphatase. Overexpression of this enzyme in *Arabidopsis*, which is analogous to overexpression of the V-ATPase, causes enhanced NaCl tolerance in transgenic lines (Gaxiola *et al.*, 2001). It would be informative to obtain these lines and test their Al and NaCl tolerance in the root elongation system.

In future, it would be interesting to determine the precise mechanism by which increased V-ATPase activity mediates Al resistance. Is V-ATPase activity required to allow transmembrane transport of an Al species, analogous to how it is involved in NaCl tolerance? It is also possible that the tonoplast transmembrane potential is required to allow transport of signaling molecules, such as calcium, which then trigger other downstream events. Finally, it is possible that increased V-ATPase

activity is simply a mechanism of pH control. Comparing vacuolar accumulation of Al in Al-resistant PT741 and Al-sensitive wheat cultivars could test the first of these possibilities. However, the rare ^{26}Al isotope and specialized microextraction techniques would need to be used to measure low levels of Al within root tip cells (Taylor *et al.*, 2000). Even with these techniques, the required sensitivity may not be available. Examining Ca^{2+} fluxes during Al stress could determine whether or not Ca^{2+} release from vacuolar stores exhibits a different pattern in PT741 than in other wheat cultivars. The simplest of these possibilities, that the V-ATPase acts as an intracellular “pH-stat”, could be tested using microelectrodes or indicator dyes to measure the cytoplasmic pH of individual cells of PT741 and other wheat cultivars during Al stress. The above list is not exhaustive, but it illustrates the additional layers of complexity which remain to be unraveled by future research. It is possible that one or more of these mechanisms, or a different mechanism not discussed here, is responsible for V-ATPase mediated Al-resistance.

In conclusion, in this thesis I have: 1) identified three Al-induced proteins as the B subunit of the vacuolar H^+ -ATPase and the α and β subunits of the mitochondrial F_1F_0 -ATPase, 2) demonstrated that V-ATPase and F_1F_0 -ATPase activities are induced by Al in Al-resistant PT741 and not in other Al-resistant or Al-sensitive wheat cultivars tested, 3) developed a testable hypothesis describing a possible mechanism by which increased V-ATPase and F_1F_0 -ATPase activities could mediate Al resistance, 4) verified this hypothesis using knockout mutants of yeast and 5) transgenic Arabidopsis and 6) expanded this hypothesis by analogy to NaCl

tolerance. Further support for my hypothesis awaits future phenotypic testing of Al resistance in yeast and transgenic plants overexpressing the vacuolar H⁺-pyrophosphatase, and determination of the mechanism by which increased V-ATPase activity mediates Al resistance.

6.1 Literature cited

- Aoyama T and Chua N-H** (1997) A glucocorticoid-inducible transcriptional induction system in transgenic plants. *Plant J* **11**: 605-612
- Basu A, Basu U and Taylor GJ** (1994) Induction of microsomal membrane proteins in roots of an aluminum-resistant cultivar of *Triticum aestivum* L. under conditions of aluminum stress. *Plant Physiol* **104**: 1007-1013
- Cheng M, Fry JE, Pang S, Zhou H, Hironaka CM, Duncan DR, Conner TW and Wan Y** (1997) Genetic transformation of wheat mediated by *Agrobacterium tumefaciens*. *Plant Physiol* **115**: 971-980
- Gaxiola RA, Li J, Undurraga S, Dang LM, Allen GJ, Alper SL and Fink GR** (2001) Drought- and salt-tolerant plants result from overexpression of the AVP1 H⁺-pump. *Proc Nat Acad Sci* **98**: 11444-11449
- Gaxiola RA, Rao R, Sherman A, Grisafi P, Alper SL and Fink GR** (1999) The *Arabidopsis thaliana* proton transporters, AtNhx1 and Avp1, can function in cation detoxification in yeast. *Proc Nat Acad Sci* **96**: 1480-1485
- Marquez JA and Serrano R** (1996) Multiple transduction pathways regulate the sodium-extrusion gene *PMR2/ENAI* during salt stress in yeast. *FEBS Lett* **382**: 89-92
- Taylor GJ, Basu A, Basu U, Slaski JJ, Zhang G and Good A** (1997) Al-induced, 51 kilodalton, membrane-bound proteins are associated with resistance to Al in a segregating population of wheat. *Plant Physiol* **114**: 363-372

Taylor GJ, McDonald-Stephens JL, Hunter DB, Bertsch PM, Elmore D, Rengel Z and Reid RJ (2000) Direct measurement of aluminum uptake and distribution in single cells of *Chara corallina*. *Plant Physiol* **123**: 987-996