

# CANADIAN THESES ON MICROFICHE

I.S.B.N.

## THESES CANADIENNES SUR MICROFICHE



National Library of Canada  
Collections Development Branch

Canadian Theses on  
Microfiche Service

Ottawa, Canada  
K1A 0N4

Bibliothèque nationale du Canada  
Direction du développement des collections

Service des thèses canadiennes  
sur microfiche

### NOTICE

The quality of this microfiche is heavily dependent upon the quality of the original thesis submitted for microfilming. Every effort has been made to ensure the highest quality of reproduction possible.

If pages are missing, contact the university which granted the degree.

Some pages may have indistinct print especially if the original pages were typed with a poor typewriter ribbon or if the university sent us a poor photocopy.

Previously copyrighted materials (journal articles, published tests, etc.) are not filmed.

Reproduction in full or in part of this film is governed by the Canadian Copyright Act, R.S.C. 1970, c. C-30. Please read the authorization forms which accompany this thesis.

**THIS DISSERTATION  
HAS BEEN MICROFILMED  
EXACTLY AS RECEIVED**

### AVIS

La qualité de cette microfiche dépend grandement de la qualité de la thèse soumise au microfilmage. Nous avons tout fait pour assurer une qualité supérieure de reproduction.

S'il manque des pages, veuillez communiquer avec l'université qui a conféré le grade.

La qualité d'impression de certaines pages peut laisser à désirer, surtout si les pages originales ont été dactylographiées à l'aide d'un ruban usé ou si l'université nous a fait parvenir une photocopie de mauvaise qualité.

Les documents qui font déjà l'objet d'un droit d'auteur (articles de revue, examens publiés, etc.) ne sont pas microfilmés.

La reproduction, même partielle, de ce microfilm est soumise à la Loi canadienne sur le droit d'auteur, SRC 1970, c. C-30. Veuillez prendre connaissance des formules d'autorisation qui accompagnent cette thèse.

**LA THÈSE A ÉTÉ  
MICROFILMÉE TELLE QUE  
NOUS L'AVONS REÇUE**



National Library  
of Canada

Bibliothèque nationale  
du Canada

Canadian Theses Division    Division des thèses canadiennes

Ottawa, Canada  
K1A 0N4

53927

0-315-06006-9

### PERMISSION TO MICROFILM — AUTORISATION DE MICROFILMER

• Please print or type — Écrire en lettres moulées ou dactylographier

Full Name of Author — Nom complet de l'auteur

JALETA ANNE GREGG

Date of Birth — Date de naissance

MAY 19, 1952

Country of Birth — Lieu de naissance

USA

Permanent Address — Résidence fixe

Dept of Animal Science  
University of Alberta  
Edmonton, Alberta T6G 2E3

Title of Thesis — Titre de la thèse

METABOLIC COMPONENTS OF THE MAINTENANCE  
ENERGY EXPENDITURE OF MAMMALIAN  
SKELETAL MUSCLE

University — Université

University of Alberta

Degree for which thesis was presented — Grade pour lequel cette thèse fut présentée

Ph. D.

Year this degree conferred — Année d'obtention de ce grade

1981

Name of Supervisor — Nom du directeur de thèse

L. P. MILLIGAN

Permission is hereby granted to the NATIONAL LIBRARY OF CANADA to microfilm this thesis and to lend or sell copies of the film.

The author reserves other publication rights, and neither the thesis nor extensive extracts from it may be printed or otherwise reproduced without the author's written permission.

L'autorisation est, par la présente, accordée à la BIBLIOTHÈQUE NATIONALE DU CANADA de microfilmer cette thèse et de prêter ou de vendre des exemplaires du film.

L'auteur se réserve les autres droits de publication; ni la thèse ni de longs extraits de celle-ci ne doivent être imprimés ou autrement reproduits sans l'autorisation écrite de l'auteur.

Date

October 8, 1981

Signature

Jaleta Anne Gregg

THE UNIVERSITY OF ALBERTA

METABOLIC COMPONENTS OF THE MAINTENANCE ENERGY EXPENDITURE  
OF MAMMALIAN SKELETAL MUSCLE

by



Valeta Anne Gregg

A THESIS

SUBMITTED TO THE FACULTY OF GRADUATE STUDIES AND RESEARCH  
IN PARTIAL FULFILMENT OF THE REQUIREMENTS FOR THE DEGREE

OF DOCTOR OF PHILOSOPHY

IN

ANIMAL BIOCHEMISTRY

DEPARTMENT OF ANIMAL SCIENCE

EDMONTON, ALBERTA

FALL, 1981

THE UNIVERSITY OF ALBERTA

RELEASE FORM

NAME OF AUTHOR Valeta Anne Gregg  
TITLE OF THESIS METABOLIC COMPONENTS OF THE MAINTENANCE  
ENERGY EXPENDITURE OF MAMMALIAN SKELETAL  
MUSCLE

DEGREE FOR WHICH THESIS WAS PRESENTED DOCTOR OF PHILOSOPHY  
YEAR THIS DEGREE GRANTED FALL, 1981

Permission is hereby granted to THE UNIVERSITY OF ALBERTA LIBRARY to reproduce single copies of this thesis and to lend or sell such copies for private, scholarly or scientific research purposes only.

The author reserves other publication rights, and neither the thesis nor extensive extracts from it may be printed or otherwise reproduced without the author's written permission.

(SIGNED) *Valeta Anne Gregg*....

PERMANENT ADDRESS:

*Dept. of Animal Science....  
University of Alberta....  
Edmonton, Alberta T6S 2E3.*

DATED *October 7*.....1981

THE UNIVERSITY OF ALBERTA  
FACULTY OF GRADUATE STUDIES AND RESEARCH

The undersigned certify that they have read, and recommend to the Faculty of Graduate Studies and Research, for acceptance, a thesis entitled METABOLIC COMPONENTS OF THE MAINTENANCE ENERGY EXPENDITURE OF MAMMALIAN SKELETAL MUSCLE submitted by Valeta Anne Gregg in partial fulfilment of the requirements for the degree of DOCTOR OF PHILOSOPHY in ANIMAL BIOCHEMISTRY.

..... *L.P. Miller* .....

Supervisor

..... *Mary Spencer* .....

..... *Gary Matheson* .....

..... *R.J. Christopherson* .....

..... *R.E. Howarth* .....

External Examiner

Date... *Oct 5, 1951* .....

## Abstract

The intent of this study was to gain a greater quantitative understanding of the costs of  $\text{Na}^+, \text{K}^+$ -ATPase activity and protein synthesis as components of the maintenance or background energy expenditure of mammalian muscle.

The rate of  $\text{O}_2$  consumption and its inhibition by ouabain were measured for intact and sliced mouse soleus and diaphragm muscles incubated in vitro in an  $\text{O}_2$  electrode system. Slicing lowered ( $p < 0.05$ ) the rate of  $\text{O}_2$  consumption of soleus and diaphragm muscles but did not cause a significant difference in the extent of ouabain inhibition of respiration. Ouabain caused a 19.7% inhibition of soleus muscle incubated in 1 mM  $\text{MgCl}_2$  buffer. The response of respiration to ouabain was abolished upon incubation in buffer containing 10 mM  $\text{MgCl}_2$ . Soleus muscle from mice that had been held at an ambient temperature of  $5^\circ\text{C}$  for 3 weeks had a greater ( $p < 0.001$ ) rate of  $\text{O}_2$  consumption than did the soleus muscle of mice held at  $24^\circ\text{C}$ . Increased  $\text{Na}^+, \text{K}^+$ -ATPase activity accounted for 20% of the cold-induced muscle thermogenesis.

An in vitro preparation of sternomandibularis muscle from sheep and cattle in which  $\text{O}_2$  availability and membrane potential were maintained was developed.  $\text{O}_2$  consumption and inhibition of respiration by ouabain were measured for sheep exposed to a warm ( $25^\circ\text{C}$ ) or cold ( $1^\circ\text{C}$ ) ambient temperature for 5 weeks and fed either at maintenance or at the same

level of intake. Cold exposed sheep had a whole body  $O_2$  consumption 24 or 42% greater ( $p < 0.05$ ) when fed at the same level as warm exposed sheep, or at maintenance (1400 g alfalfa pellets/d), respectively. Muscle from cold exposed sheep fed at either level of intake exhibited an  $O_2$  consumption rate 48% greater ( $p < 0.001$ ) than that of warm exposed sheep. Ouabain inhibited the muscle  $O_2$  consumption of cold exposed sheep by 41.0 to 45.0%, and of warm exposed sheep by 29.0 to 38.0%. Increased energy expenditure at the level of the  $Na^+, K^+$ -ATPase accounted for 50 to 80% of the cold-induced muscle thermogenesis.

Ouabain ( $10^{-5}M$ ) caused an average of 40% inhibition of sternomandibularis muscle respiration for dairy calves aged 10-21d and 7 months, and 7 month old calves from a double-muscled (DM) population of normal (control DM) and heavily muscled (extreme DM) phenotypes. Rate of  $O_2$  consumption was greatest ( $p < 0.001$ ) for muscle from 10-21d dairy calves and lowest ( $p < 0.05$ ) for control DM calves. The energy expenditure estimated to be required for peptide bond synthesis accounted for 2.0 to 3.3% of the  $O_2$  consumption of the muscle preparations. Rate of tyrosine release, considered to be an indicator of protein degradation, was greatest ( $p < 0.05$ ) for muscle from control and extreme DM calves; both dairy groups had similarly low rates of muscle tyrosine release.

Muscle  $O_2$  consumption was greater ( $p < 0.001$ ) for lambs at 2 weeks than at 7 weeks of age, and for ewes when

lactating than when dry. Ouabain ( $10^{-5}M$ ) inhibited muscle  $O_2$  consumption by an average 39% for all animals. Increased energy expenditure at the level of the  $Na^+,K^+$ -ATPase accounted for 40% of the increased  $O_2$  consumption rate of muscle from lambs at 2 weeks as contrasted to 7 weeks of age, and 60% of the increased  $O_2$  consumption rate of muscle due to lactation.



## Acknowledgements

I am deeply grateful to Dr. L. P. Milligan for his uncompromising support and dedication in providing the intellectual environment necessary for a successful and meaningful graduate program.

I wish to thank Dr. W. N. Garrett for his continual encouragement and inspiration.

It is with love and respect that I thank my parents, Capt. Richard W. Gregg and Maria R. C. de Gregg.

I thank the Department of Animal Science for use of the facilities.

## Table of Contents

Chapter	Page
Introduction .....	1
I. INHIBITION BY OUABAIN OF THE O <sub>2</sub> CONSUMPTION OF MOUSE (MUS MUSCULUS) SOLEUS AND DIAPHRAGM MUSCLES .....	3
A. Abstract .....	3
B. Introduction .....	3
C. Experimental .....	5
Measurement of ouabain-sensitive respiration of whole soleus and diaphragm muscles .....	5
Comparison of ouabain-sensitive respiration of whole and sliced muscles .....	6
Statistical analysis .....	6
D. Results and Discussion .....	7
E. Acknowledgements .....	10
F. References .....	10
II. INHIBITION OF Na <sup>+</sup> ,K <sup>+</sup> -ATPase OF INTACT MOUSE SOLEUS MUSCLE BY Mg <sup>2+</sup> .....	14
A. Abstract .....	14
B. Introduction .....	14
C. Experimental .....	16
Measurement of ouabain-sensitive respiration of intact soleus muscle in control and experimental buffers. ....	16
Statistical analysis .....	17
D. Results and Discussion .....	17
E. Acknowledgements .....	18
F. References .....	19

III.	O <sub>2</sub> CONSUMPTION AND Na <sup>+</sup> ,K <sup>+</sup> -ATPase ACTIVITY IN INTACT SOLEUS MUSCLE FROM COLD EXPOSED MICE .....	21
	A. Abstract .....	21
	B. Introduction .....	21
	C. Experimental .....	22
	Measurement of O <sub>2</sub> consumption and Na <sup>+</sup> ,K <sup>+</sup> -ATPase activity .....	22
	Statistical analysis .....	23
	D. Results and Discussion .....	23
	E. Acknowledgements .....	25
	F. References .....	26
IV.	ROLE OF NA <sup>+</sup> ,K <sup>+</sup> -ATPASE IN MUSCULAR ENERGY EXPENDITURE OF WARM AND COLD EXPOSED SHEEP .....	28
	A. Abstract .....	28
	B. Introduction .....	29
	C. Experimental .....	30
	Muscle preparation .....	30
	Electrophysiological studies .....	30
	Muscle characterization .....	31
	Measurement of sheep muscle respiration and response to ouabain .....	31
	Trial 1. Respiration and ouabain-sensitive respiration of muscle preparations from warm and cold exposed sheep fed at maintenance .....	32
	Trial 2. Respiration and ouabain-sensitive respiration of muscle preparations from warm and cold exposed sheep fed at the same level of feed intake .....	34
	Na <sup>+</sup> ,K <sup>+</sup> -ATPase Assay .....	34
	Statistical analysis .....	35
	D. Results .....	35

	Response of <del>sheep</del> muscle respiration to ouabain .....	35
	Trial 1. Respiration and ouabain-sensitive respiration of muscle preparations from warm and cold exposed sheep fed at maintenance .....	36
	Trial 2. Respiration and ouabain-sensitive respiration of muscle preparations from warm and cold exposed sheep fed at the same level of intake .....	37
	Na <sup>+</sup> ,K <sup>+</sup> -ATPase assay .....	38
	E. Discussion .....	39
	F. Acknowledgments .....	43
	G. References .....	43
V.	ENERGY COSTS OF NA <sup>+</sup> ,K <sup>+</sup> ATPase ACTIVITY AND PROTEIN SYNTHESIS IN MUSCLE FROM CALVES DIFFERING IN AGE AND BREED .....	52
	A. Abstract .....	52
	B. Introduction .....	53
	C. Experimental .....	54
	Animals .....	54
	Respiration and Na <sup>+</sup> ,K <sup>+</sup> -ATPase-dependent respiration .....	54
	Measurement of <sup>14</sup> C-phenylalanine incorporation .....	55
	Measurement of tyrosine release .....	57
	Statistical analysis .....	57
	D. Results .....	58
	O <sub>2</sub> consumption and Na <sup>+</sup> ,K <sup>+</sup> -ATPase activity ..	58
	Rates of <sup>14</sup> C-phenylalanine incorporation and tyrosine release .....	59
	E. Discussion .....	60
	F. Acknowledgements .....	66

G. References .....	66
VI. O <sub>2</sub> CONSUMPTION AND Na <sup>+</sup> , K <sup>+</sup> -ATPase-DEPENDENT RESPIRATION IN MUSCLE OF LAMBS AND LACTATING AND NON-LACTATING EWES .....	72
A. Abstract .....	72
B. Introduction .....	73
C. Experimental .....	73
Measurement of respiration and Na <sup>+</sup> , K <sup>+</sup> -ATPase-dependent respiration .....	74
Statistical analysis .....	75
D. Results .....	75
Comparisons of measurements during lactation and after weaning .....	76
E. Discussion .....	78
F. Acknowledgements .....	81
G. References .....	81
General Discussion and Conclusions .....	85

## List of Tables

Table		Page
I.1	Ouabain-sensitive respiration of whole soleus and diaphragm muscles .....	12
I.2	Comparison of ouabain-sensitive respiration of whole and sliced soleus and diaphragm muscles .....	13
II.1	Effect of $MgCl_2$ concentration on inhibition of respiration of intact mouse soleus muscle by ouabain .....	20
III.1	$O_2$ consumption and $Na^+, K^+$ -ATPase-dependent and independent respiration of intact soleus muscle from warm and cold exposed mice .....	27
IV.1	Whole animal and muscle respiration from sheep fed at maintenance .....	46
IV.2	Whole animal and muscle respiration from sheep fed at the same level of intake .....	47
V.1	In vitro $O_2$ consumption, ouabain inhibition and $Na^+, K^+$ -ATPase-dependent and independent respiration of calf muscle preparations .....	69
V.2	Rate of $^{14}C$ -phenylalanine incorporation into protein of calf muscle preparations .....	70
V.3	Rate of tyrosine release from calf muscle preparations .....	71
VI.1	$O_2$ consumption and proportion of respiration inhibited by ouabain for muscle from lambs and lactating and non-lactating ewes .....	83
VI.2	$Na^+, K^+$ -ATPase-dependent respiration and independent respiration for muscle from lambs and lactating and non-lactating ewes .....	84

## List of Figures

Figure	Page
IV.1 Inhibition of sheep muscle respiration by ouabain .....	48
IV.2 Respiration and ouabain-sensitive and insensitive respiration of muscle from warm and cold exposed sheep fed at maintenance or 1150 g alfalfa pellets/d .....	49
IV.3 Inhibition by ouabain of respiration of muscle from warm and cold exposed sheep fed at maintenance .....	50
IV.4 Inhibition by ouabain of respiration of muscle from warm and cold exposed sheep fed 1150 g alfalfa pellets/d .....	51

### Introduction

The concept of a constant maintenance requirement based on body weight has been a fundamental element of energy nutrition. This concept of a maintenance component of energy expenditure inevitably influences estimates of energetic efficiency in intact animals because the presumed maintenance energy requirement is first subtracted from energy intake and the difference is then used as the input component in assessment of energetic efficiency of production. In contrast to nutritional concepts associated with production, maintenance has not been identified in terms of metabolic events and it has certainly not been established that the metabolic events accounting for maintenance energy expenditures are constant in relation to body weight. The purpose of this study was to identify metabolic components of maintenance or background energy expenditure.

Metabolic processes which could intuitively be suggested to be part of the maintenance energy requirement include the continual background expenditures on active  $\text{Na}^+\text{-K}^+$  transport, catalyzed by the plasma membrane  $\text{Na}^+\text{,K}^+\text{-ATPase}$ , and on the protein synthesis that must occur during protein turnover in animals. It is not possible to measure  $\text{Na}^+\text{,K}^+\text{-ATPase}$  activity in vivo without disruption or destruction of supporting vital functions, therefore, efforts were focused on obtaining in vitro measurements of



Na<sup>+</sup>,K<sup>+</sup>-ATPase activity which would be indicative of the physiological role of the enzyme as a component of the energy expenditure of skeletal muscle. This involved development of an in vitro preparation in which membrane potential and O<sub>2</sub> availability to the sites of respiration were maintained. This preparation was then amenable to measurement of the proportion of respiration required to support active transport of Na<sup>+</sup>-K<sup>+</sup> using ouabain as a specific inhibitor and to support protein synthesis as measured by the rate of incorporation of <sup>14</sup>C-phenylalanine, and could be used in the examination of the effects of physiological state on these parameters.

# I. INHIBITION BY OUABAIN OF THE O<sub>2</sub> CONSUMPTION OF MOUSE (MUS MUSCULUS) SOLEUS AND DIAPHRAGM MUSCLES'

## A. Abstract

The rate of O<sub>2</sub> consumption of intact, or sliced soleus and diaphragm muscles was measured polarographically with lactate or glucose as the added substrate; the dimensions of the muscles were such that O<sub>2</sub> diffusion should not have limited respiration.

Ouabain (10<sup>-3</sup>M) inhibited the respiration of intact soleus and diaphragm muscles by 22 and 33% indicating very real importance of Na<sup>+</sup> + K<sup>+</sup> transport in the energy metabolism of these muscles. Slicing lowered the rate of O<sub>2</sub> uptake of soleus and diaphragm by 9 and 14%. Ouabain inhibition of respiration tended to be greater for sliced than for intact muscles but the effect of this method of preparation of tissue was not statistically (p>0.05) significant.

## B. Introduction

It is generally recognized that ATP production in intact coupled cells is regulated by ATP utilization (Ismail-Beigi, 1977); the availability of ADP and, consequently, the rate of free energy expenditure controls the rate of respiration (Chance & Williams, 1956). Active

-----  
'A slightly modified version of this chapter has been published. Gregg, V. A. & Milligan, L. P. (1980) Gen. Pharmac. 11, 323.

ion transport, particularly that catalyzed by the  $\text{Na}^+, \text{K}^+$ -ATPase of the plasma membrane, has been thought by some investigators to result in a significant portion of total ADP generation (Racker, 1976), accounting for 20-45% of the energy expenditure of resting cells (Whittam, 1964; Whittam & Blond, 1964). Ouabain, a specific inhibitor of  $\text{Na}^+, \text{K}^+$ -ATPase has been used in attempts to measure  $\text{Na}^+, \text{K}^+$ -ATPase-dependent respiration in coupled systems (Ismail-Beigi & Edelman, 1970). However, ensuing results on the importance of  $\text{Na}^+ + \text{K}^+$  transport as a component of energy expenditure have been contradictory. Work conducted with quarter-sections of rat diaphragm muscle indicated that  $\text{Na}^+ + \text{K}^+$  transport accounts for 40% of total cellular energy expenditures (Asano et al. 1976), while workers using perfused rat liver (Folke & Sestoft, 1977), isolated rat soleus muscle and adipose tissue (Chinet et al. 1977), have concluded that  $\text{Na}^+ + \text{K}^+$  transport accounts for no more than 6% of the energy expenditure of these tissues.

A study was undertaken to measure the  $\text{Na}^+, \text{K}^+$ -ATPase-dependent respiration of intact mouse (*Mus musculus*) diaphragm and soleus muscles under conditions in which  $\text{O}_2$  availability would not limit respiration. A second study was conducted to assess the response of these muscles to ouabain after the tissue was sliced.

### C. Experimental

Adult male mice, in the weight range of 24-41 g, were stunned by a blow to the head and bled from the neck. The intact soleus muscles were removed with care to minimize tissue damage following the procedure of Kohn and Clausen (1971). The diaphragm was dissected from the rib attachments and cut through the central connective tissue. This method produces few damaged muscle fibers relative to the number of intact fibers.

#### Measurement of ouabain-sensitive respiration of whole soleus and diaphragm muscles

The modified Krebs-Ringer bicarbonate buffer used as the incubation medium contained (mM): NaCl, 116.8; KCl, 5.9; NaHCO<sub>3</sub>, 25.0; MgSO<sub>4</sub>, 1.2; NaH<sub>2</sub>PO<sub>4</sub>, 1.2; CaCl<sub>2</sub>, 1.0; lactate, 1.0; pH 7.4. The tissue preparations were incubated individually in 6 ml buffer at 37°C for 1 h without (control) and with 10<sup>-3</sup>M ouabain. Incubation flask contents were quantitatively transferred to the O<sub>2</sub> electrode chamber and O<sub>2</sub> consumption was then measured with a YSI O<sub>2</sub> electrode. O<sub>2</sub> consumption values were obtained for the combined soleus muscles from two mice; values for the diaphragms of the two mice were obtained separately and averaged. The O<sub>2</sub> consumption of representative preparations was measured initially and at approximately 10 min intervals throughout the incubation period and was found to remain constant indicating adequacy of the experimental conditions

in maintaining tissue metabolism.

### Comparison of ouabain-sensitive respiration of whole and sliced muscles

The Na<sup>+</sup>-Ringer's buffer used as the incubation medium in this part of the study contained (mM): NaCl, 130.0; KCl, 5.0; NaH<sub>2</sub>PO<sub>4</sub>, 5.0; CaCl<sub>2</sub>, 1.0; glucose, 10.0; pH7.4. O<sub>2</sub> consumption values were obtained for whole and sliced muscle following the procedure used for the measurement of the ouabain-sensitive respiration of whole muscles. Soleus muscles were longitudinally sliced by hand using a razor blade into two or three sections; hemidiaphragm muscles were sliced into five or six sections.

### Statistical analysis

Results obtained for the ouabain-sensitive respiration of whole soleus and diaphragm muscle were analysed using a three-way analysis of variance with groups as a random source and tissue and incubation as fixed sources of variation. The comparisons of ouabain-sensitive respiration of whole and sliced soleus and diaphragm muscles, as well as that of whole muscles in two buffers, were analysed using four-way analysis of variance with treatment, tissue and inhibitor as fixed sources and groups nested within treatment a random source.

#### D. Results and Discussion

Although a variety of buffers have been used by previous investigators (Asano et al. 1976; Folke & Sestoft, 1977), an analysis of the values obtained in this experiment for respiration rates and inhibition of respiration by ouabain of whole soleus and diaphragm muscles showed that determinations conducted in Krebs-Ringer bicarbonate buffer with lactate as substrate were not significantly different from those conducted with Na<sup>+</sup>-Ringer's buffer with glucose as substrate. The respiration rates measured for whole diaphragm muscle, 4.3  $\mu\text{l O}_2/\text{mg}$  tissue dry wt/h in Krebs-Ringer bicarbonate buffer (Table I.1) and 4.5  $\mu\text{l O}_2/\text{mg}$  tissue dry wt/h in Na<sup>+</sup>-Ringer's buffer (Table I.2), were lower than the reports of 7.7 (Ismail-Beigi & Edelman, 1970) and 7.5  $\mu\text{l O}_2/\text{mg}$  tissue dry wt/h (Asano et al. 1976) for quarter sections of rat diaphragm muscle, but the mouse preparations likely include proportionally more connective tissue. The respiration rates measured for whole soleus muscle, 3.7  $\mu\text{l O}_2/\text{mg}$  tissue dry wt/h in Krebs-Ringer bicarbonate buffer (Table I.1) and 3.5  $\mu\text{l O}_2/\text{mg}$  tissue dry wt/h in Na<sup>+</sup>-Ringer's buffer (Table I.2) were slightly less than those measured for whole diaphragm muscle.

Inhibition of respiration of whole soleus and diaphragm muscles by ouabain was greater, but not significantly ( $p > 0.05$ ) so in Krebs-Ringer bicarbonate buffer (21.6 and 32.6%) (Table I.1) than in Na<sup>+</sup>-Ringer's buffer (14.3 and 15.6%) (Table I.2). The values determined in this study are

consistent with the 15.6-41.3% inhibition reported previously for rat diaphragm muscle (Ismail-Beigi & Edelman, 1970; Asano et al. 1976). The values for ouabain inhibition of respiration for whole soleus muscle in Krebs-Ringer buffer (Table I.1) were significantly less ( $p < 0.05$ ) than those of whole diaphragm muscles; this difference was not observed for the values determined in  $\text{Na}^+$ -Ringer's buffer (Table I.2).

The results of a comparison of respiration rates and of inhibition of respiration by ouabain for whole and sliced tissue are shown in Table I.2. Slicing lowered ( $p < 0.05$ )  $\text{O}_2$  consumption of soleus muscle from 3.5 to 3.2  $\mu\text{l O}_2/\text{mg}$  tissue dry wt/h and of diaphragm muscle from 4.5 to 3.8  $\mu\text{l O}_2/\text{mg}$  tissue dry wt/h (Table I.2). The inhibition of respiration by ouabain, while greater, was not significantly ( $p > 0.05$ ) different from that observed in whole tissue; inhibition was 21.9 to 14.3% in sliced and whole soleus muscle and 26.3 and 15.6% in sliced and whole diaphragm muscle (Table I.2).

The extent to which  $\text{Na}^+, \text{K}^+$ -ATPase acts as a pacemaker of energy metabolism in intact, coupled cells is determined by the magnitude of energy that must be expended to maintain intracellular ionic concentrations and counteract the tendency for external  $\text{Na}^+$  and internal  $\text{K}^+$  to be moved or to leak across the cell membrane in the direction of their diffusion gradients. Changes in the leakage rates would influence this expenditure. The ouabain-sensitive respiration values measured in this study for mouse soleus

and diaphragm muscles certainly support the importance of the role played by  $\text{Na}^+ + \text{K}^+$  transport in the energy expenditure of these tissues.

Chinet et al. (1977) suggested that the large ouabain-sensitive respiration values measured by others for cut or sliced muscle preparations were due to the stimulation of  $\text{Na}^+ + \text{K}^+$  transport above its basal level by the leakage of  $\text{Na}^+$  into the cytoplasm of the cut fibers. Clearly, our results do not support this argument. However, it is likely that the low ouabain-sensitive respiration values measured for whole organ preparations (Chinet et al. 1977; Folke & Sestoft, 1977) were obtained under conditions in which  $\text{O}_2$  availability limited respiration. The microcalorimetric studies of heat production by whole rat soleus muscle (Chinet et al. 1977) would have entailed a tissue thickness of 1.5-2.0 mm which exceeds the approximate 0.4 mm at which the rate of  $\text{O}_2$  diffusion would limit respiration, according to the method of calculation of Kleiber (1961). Folke & Sestoft (1977) concluded that during the perfusion of rat livers with ouabain there was loss of vascular integrity as shown by increased resistance to perfusion and the appearance of dark areas on the surface of the tissue; this suggests that there was likely impaired delivery of  $\text{O}_2$  to the sites of cellular utilization. Since mouse soleus and diaphragm muscles are <0.5 mm thick, our determinations of respiration rates of these whole muscles likely did not involve rate limitation by  $\text{O}_2$  diffusion.



Differences in species and tissue sensitivity to ouabain have been established (Schwartz et al. 1969; Allen & Schwartz, 1969). Rat  $\text{Na}^+, \text{K}^+$ -ATPase is particularly insensitive to ouabain largely as a result of instability of the enzyme-ouabain complex which dissociates relatively rapidly. If the mouse enzyme-ouabain complex also dissociates rapidly, the ouabain-sensitive respiration values obtained in this experiment would be minimal indications of active  $\text{Na}^+ + \text{K}^+$  transport in the energy metabolism of mouse soleus and diaphragm muscles.

#### E. Acknowledgements

Grateful acknowledgement is extended to the Natural Sciences and Engineering Research Council Canada for support in part of this research.

#### F. References

- Allen, J. C. & Schwartz, A. (1969) J. Pharmac. Exp. Ther. 168, 42.
- Asano, Y., Liberman, V. A. & Edelman, I. S. (1976) J. Clin. Invest. 57, 368.
- Chance, B. & Williams, G. R. (1956) Adv. Enzymol. 17, 65.
- Chinet, A., Clausen, T. & Girardier, L. (1977) J. Physiol. 265, 43.
- d

- Folke, M. & Sestoft, L. (1977) *J. Physiol.* 269, 407.
- Ismail-Beigi, F. (1977) *Curr. Top. Mem. Transp.* 9, 367.
- Ismail-Beigi, F. & Edelman, I. S. (1970) *J. gen. Physiol.* 57, 710.
- Ismail-Beigi, F. & Edelman, I. S. (1970) *Proc. Natl. Acad. Sci., USA* 67, 1071.
- Kleiber, M. (1961) *The Fire of Life*. New York, John Wiley & Sons, Inc., p.66.
- Kohn, P. G. & Clausen, T. (1971) *Biochim. Biophys. Acta.* 225, 277.
- Racker, E. (1976) *A New Look at Mechanisms in Bioenergetics*. Academic Press, New York.
- Schwartz A., Allen, J. C. & Harigaya, S. (1969) *J. Pharmac. exp. Ther.* 168, 31.
- Whittam, R. (1964) in *The Cellular Functions of Membrane Transport* (J.P. Hoffman, ed.), p. 139. Prentice-Hall, Engelwood Cliffs.
- Whittam, R. & Blond, D. M. (1964) *Biochem. J.* 92, 147.

Table 1.1 Ouabain-sensitive respiration of whole soleus and diaphragm muscles<sup>1</sup>

Muscle	n	Control	+Ouabain	% Inhibition
Soleus <sup>2</sup>	9	3.7 ± 0.1	2.9 ± 0.1	21.6
Diaphragm <sup>3</sup>	9	4.3 ± 0.1	2.9 ± 0.1	32.6

<sup>1</sup> Values expressed as  $\mu\text{l O}_2/\text{mg tissue dry wt/h} \pm \text{SEM}$ .

<sup>2</sup> Average dry wt. 1.4 mg.

<sup>3</sup> Average dry wt. 2.5 mg.

Table I.2 Comparison of ouabain-sensitive respiration of whole and sliced soleus and diaphragm muscles.

Treatment	Muscle	n	Control	+Ouabain	% Inhibition
Whole	Soleus <sup>1</sup>	10	3.5 ± 0.1	3.0 ± 0.1	14.3
	Diaphragm <sup>2</sup>	10	4.5 ± 0.1	3.8 ± 0.1	15.6
Sliced	Soleus <sup>1</sup>	10	3.2 ± 0.1	2.5 ± 0.1	21.9
	Diaphragm <sup>2</sup>	10	3.8 ± 0.1	2.8 ± 0.1	26.3

<sup>1</sup> Values expressed as  $\mu\text{l O}_2/\text{mg tissue dry wt/h} \pm \text{SEM}$ .  
<sup>2</sup> Average dry wt. 1.7 mg.  
<sup>3</sup> Average dry wt. 10.2 mg.

## II. INHIBITION OF $\text{Na}^+, \text{K}^+$ -ATPase OF INTACT MOUSE SOLEUS MUSCLE BY $\text{Mg}^{++}$

### A. Abstract

The effect of 10 mM  $\text{Mg}^{++}$  on the inhibition of respiration by ouabain was investigated with intact mouse soleus muscle preparations. Although ouabain caused a 19.7% inhibition of respiration of soleus muscle incubated in 1 mM  $\text{MgCl}_2$  buffer, the response of respiration to ouabain was abolished upon incubation in buffer containing 10 mM  $\text{MgCl}_2$ . Initial respiration rates were significantly decreased ( $p < 0.001$ ) in soleus muscle exposed to 10 mM, as contrasted to 1 mM  $\text{MgCl}_2$ .

### B. Introduction

Studies of active  $\text{Na}^+$  transport conducted with intact muscle preparations have yielded conflicting results concerning the importance of  $\text{Na}^+, \text{K}^+$ -ATPase ( $\text{Na}^+, \text{K}^+$ -dependent adenosine triphosphatase EC 3.6.1.3.) as a component of cellular energy expenditure. Microcalorimetric studies of the ouabain-sensitive heat production of intact rat (Chinet et al. 1977) and mouse soleus muscles (Biron et al. 1979) have led to the conclusion that active  $\text{Na}^+$  transport accounts for no more than 6-8% of resting heat production. However, measurements of the ouabain-sensitive component of

-----  
'A slightly modified version of this chapter has been published. Gregg, V. & Milligan, L. P. (1980) Biochem. Biophys. Res. Commun. 95, 608.

O<sub>2</sub> consumption of intact mouse soleus muscle have shown 14-22% of O<sub>2</sub> uptake to be due to the activity of Na<sup>+</sup>,K<sup>+</sup>-ATPase (Gregg & Milligan, 1980). In view of the key role proposed for Na<sup>+</sup>,K<sup>+</sup>-ATPase in thyroid thermogenesis (Ismail-Beigi & Edelman, 1970), cold-induced thermogenesis (Guernsey & Stevens, 1977), and as a primary mechanism in the development of obesity (Lin et al. 1978), it is imperative to resolve the current uncertainty concerning the physiological importance of the Na<sup>+</sup>,K<sup>+</sup>-ATPase in resting energy expenditure.

The microcalorimetric determinations of ouabain-sensitive heat production of mouse soleus muscle were conducted in a buffer containing 10 mM MgCl<sub>2</sub> to suppress a secondary rise in heat production which occurred following infusion with ouabain (Chinet et al. 1977; Biron et al. 1979). However, previous investigators have found Na<sup>+</sup>, K<sup>+</sup>-ATPase activity to be inhibited by a high concentration of Mg<sup>++</sup> (Bond & Hudgins, 1975; Schwartz et al. 1963). Thus, it has been suggested (Smith & Edelman, 1979) that inhibition by Mg<sup>++</sup> of the Na<sup>+</sup> pump before challenge with ouabain may have contributed to the lack of inhibition of heat production by ouabain reported by Chinet et al. (1977). The possibility that Mg<sup>++</sup> will negate inhibition of respiration by ouabain was investigated in this study.

### C. Experimental

Adult female mice, in the weight range of 20-25 g, were stunned by a blow to the head and bled from the neck. The intact soleus muscles were removed with care to minimize tissue damage following the procedure of Kohn and Clausen (1971).

#### Measurement of ouabain-sensitive respiration of intact soleus muscle in control and experimental buffers.

The modified Krebs-Ringer HEPES buffer used as the incubation medium contained (mM): NaCl, 116.8; KCl, 5.9; NaHCO<sub>3</sub>, 5.0; MgSO<sub>4</sub>, 1.2; NaH<sub>2</sub>PO<sub>4</sub>, 1.2; CaCl<sub>2</sub>, 1.0; HEPES (N-(2-hydroxymethylethyl piperazine-N'-2 ethanesulfonic acid), 10.0; MgCl<sub>2</sub>, 1.0 (control) or 10.0 (experimental); glucose, 5.0; pH 7.3-7.4. Both soleus muscles were utilized from each mouse. Ouabain-sensitive respiration was measured in both the control and experimental buffers for each animal. To achieve thorough oxygenation, muscle preparations were individually equilibrated and incubated in 600 ml beakers containing 25 ml of control or experimental buffer in a shaking water bath at 37°C. Muscle preparations were equilibrated in either control or experimental buffer for 10-20 min and then transferred to the O<sub>2</sub> electrode chamber. O<sub>2</sub> consumption was measured with a YSI O<sub>2</sub> electrode for 10-15 min. The O<sub>2</sub> content of the buffer did not fall below 85% of the initial air-saturated level during the period of measurement. The muscle preparations were then incubated in

control or experimental buffer containing  $10^{-3}$ M ouabain for 45 min and  $O_2$  consumption again measured. Muscle preparations from two mice were incubated in control or experimental buffers for 45 min without ouabain and found to maintain initial respiration rates throughout the incubation period.

#### Statistical analysis

Respiration rates were compared between treatment groups by the unpaired Student's t-test.

#### D. Results and Discussion

The initial respiration rate of soleus muscle in the control buffer ( $4.65 \pm 0.39 \text{ } \mu\text{l } O_2/\text{mg dry wt/h}$ ) (Table II.1), was higher than that measured previously for intact mouse soleus muscle ( $3.7 \pm 0.10 \text{ } \mu\text{l } O_2/\text{mg dry wt/h}$ ) (Gregg & Milligan, 1980), in a modified Krebs-Ringer bicarbonate buffer. The ouabain-sensitive component of the respiration of soleus muscle incubated in the control buffer was 19.7%, which is similar to the 14-22% inhibition of respiration reported previously for intact mouse soleus muscle (Gregg & Milligan, 1980). In contrast, using the experimental buffer, addition of ouabain resulted in a slight increase (6%) in the rate of  $O_2$  consumption (Table II.1). This finding is consistent with that obtained by Bond and Hudgins (1975), in which progressive inhibition of red blood cell  $Na^+, K^+$ -ATPase was produced with increasing concentrations of  $Mg^{++}$  greater



than 3.0 mM in a buffer with a high content of  $\text{Na}^+$  and a low content of  $\text{K}^+$ . Thus, inhibition of respiration by ouabain is abolished in muscle tissue exposed to buffer containing 10 mM  $\text{MgCl}_2$ .

The respiration rate of soleus muscle in the experimental buffer was significantly lower ( $p < 0.001$ ) than the initial respiration rate of soleus muscle in the control buffer. The inhibitory effect of  $\text{Mg}^{++}$  on respiration was significantly greater ( $p < 0.001$ ) than the inhibitory effect of ouabain on respiration. It is likely that high  $\text{Mg}^{++}$  is disruptive to other cell processes involved in energy transformations as well as to  $\text{Na}^+, \text{K}^+$ -ATPase although the mechanism of its inhibitory effect is not known.

We conclude that at least part of the lack of response to ouabain observed in microcalorimetric studies of active  $\text{Na}^+$  transport in intact muscle preparations (Chinet et al. 1977; Biron et al. 1979), was due to prior inhibition of the  $\text{Na}^+, \text{K}^+$ -ATPase by the high concentration of  $\text{Mg}^{++}$  included in the buffer rather than to an unimportant role of active  $\text{Na}^+$  transport in the energy expenditure of physiologically intact muscle preparations.

#### **E. Acknowledgements**

Partial financial support from the Natural Sciences and Engineering Research Council is gratefully acknowledged.

**F. References**

- Biron, R., Burger, A., Chinet, A., Clausen, T. & Dubois-Ferriere, R. (1979) *J. Physiol.* 297, 47.
- Bond, G. H. & Hudgins, P. M. (1975) *Biochem. Biophys. Res. Commun.* 66, 645.
- Chinet, A., Clausen, T. & Girardier, L. (1977) *J. Physiol.* 265, 43.
- Gregg, V. A. & Milligan, L. P. (1980) *Gen. Pharmac.* 11, 323.
- Guernsey, D. L. & Stevens, E. D. (1977) *Science* 196, 908.
- Ismail-Beigi, F. & Edelman, I. S. (1970) *Proc. Natl. Acad. Sci. USA* 67, 1071.
- Kohn, P. G., & Clausen, T. (1971) *Biochim. Biophys. Acta* 225, 277.
- Lin, M. H., Romsos, D. R., Aker, T. & Leveille, G. A. (1978) *Biochem. Biophys. Res. Commun.* 80, 398.
- Schwartz, A., Laseter, A. & Kraitz, L. (1963) *J. Cell. Comp. Physiol.* 62, 193.
- Smith, T. J. & Edelman, I. S. (1970) *Fed. Proc.* 38, 2150.

Table II.1 Effect of MgCl<sub>2</sub> concentration on inhibition of respiration<sup>1</sup> of intact mouse soleus muscle by ouabain.

MgCl <sub>2</sub> Concentration	n	Control	+Ouabain	% Change
1.0 mM	12	4.65 ± 0.39	3.75 ± 0.29	-19.7*
10.0 mM	12	3.17 ± 0.21	3.37 ± 0.25	+6.1

<sup>1</sup> Values expressed as  $\mu\text{l O}_2/\text{mg tissue dry wt/h} \pm \text{SEM}$ .

\* P<0.001

### III. O<sub>2</sub> CONSUMPTION AND Na<sup>+</sup>,K<sup>+</sup>-ATPase ACTIVITY IN INTACT SOLEUS MUSCLE FROM COLD EXPOSED MICE

#### A. Abstract

Rates of O<sub>2</sub> consumption and Na<sup>+</sup>,K<sup>+</sup>-ATPase activity were measured for intact soleus muscle preparations from warm and cold exposed mice. Cold exposure increased muscle O<sub>2</sub> consumption by 23% (p<0.001). The portion (13-14%) of respiration inhibited by 10<sup>-3</sup>M ouabain did not differ significantly between muscle from warm and cold exposed mice. Increased Na<sup>+</sup>,K<sup>+</sup>-ATPase activity accounted for 20% of the cold-induced muscle thermogenesis.

#### B. Introduction

In previous studies of active Na<sup>+</sup>-K<sup>+</sup> transport conducted with intact mouse soleus muscle, Na<sup>+</sup>,K<sup>+</sup>-ATPase (EC 3.6.1.3) activity accounted for 14 to 33% of muscle O<sub>2</sub> consumption (Gregg & Milligan, 1980a,b).

Cold exposure was shown to increase O<sub>2</sub> consumption in sliced preparations of mouse skeletal muscle (Stevens & Kido, 1974) and rat pectoral and diaphragm muscles (Guernsey & Stevens, 1977). Increased energy expenditure at the level of the Na<sup>+</sup>,K<sup>+</sup>-ATPase accounted for 31-83% of the cold-induced muscle thermogenesis. However, the cell damage resulting from slicing has been shown to cause a decrease in O<sub>2</sub> consumption and tended to cause an increase in the proportion of respiration inhibited by ouabain for soleus

and diaphragm muscles from mice (Gregg & Milligan, 1980a). Therefore, the use of sliced preparations may overestimate the role of the  $\text{Na}^+, \text{K}^+$ -ATPase in cold-induced muscle thermogenesis.

In this experiment, the effect of cold exposure on  $\text{O}_2$  consumption and  $\text{Na}^+, \text{K}^+$ -ATPase activity was studied with intact soleus muscle preparations from mice.

### C. Experimental

Adult male mice, in the weight range of 20-25g, were housed individually in plastic cages without bedding at either 24°C (warm) or 5°C (cold) for 3 weeks. Standard Purina laboratory chow and water were available ad libitum.

#### Measurement of $\text{O}_2$ consumption and $\text{Na}^+, \text{K}^+$ -ATPase activity

Mice were stunned by a blow to the head and bled from the neck. Intact soleus muscles were removed following the procedure of Kohn and Clausen (1971).

The buffer used as the incubation medium contained (mM):  $\text{NaCl}$ , 116.0;  $\text{KCl}$ , 5.9;  $\text{CaCl}_2$ , 1.0;  $\text{NaH}_2\text{PO}_4$ , 1.2;  $\text{MgSO}_4$ , 1.2;  $\text{NaHCO}_3$ , 10.0; HEPES (N-2-hydroxyethylpiperazine-N'-2-ethanesulfonic acid), 20.0 (pH 7.4); glucose, 10.0. Muscles were equilibrated and incubated individually in 600 ml beakers containing 25 ml of buffer in a shaking water bath at 37°C. Muscles were equilibrated in buffer for 5 min and then transferred to the  $\text{O}_2$  electrode chamber.  $\text{O}_2$  consumption was measured with a YSI

O<sub>2</sub> electrode for 10-15 min. The O<sub>2</sub> content of the buffer did not fall below 85% of the initial air-saturated level during the period of measurement. Muscles were then incubated in buffer containing 10<sup>-3</sup>M ouabain for 45 min and O<sub>2</sub> consumption was again measured. Percent inhibition of respiration by ouabain was calculated using the ratio of the difference between initial and post-incubation respiration rates to the initial respiration rate.

#### Statistical analysis

Results are expressed as the mean values and their standard errors. An unpaired Student's t test was used to establish significance of differences between the means for muscle O<sub>2</sub> consumption rate, percent inhibition of respiration by ouabain, and Na<sup>+</sup>,K<sup>+</sup>-ATPase-dependent and independent respiration.

#### D. Results and Discussion

Cold-exposure increased ( $p < 0.001$ ) muscle O<sub>2</sub> consumption by 23% (Table III.1). This result is consistent with the 9.2-55% increases in O<sub>2</sub> consumption of sliced skeletal muscle preparations from mice and rats when subjected to cold exposure (Stevens & Kido, 1974; Guernsey & Stevens, 1977; Guernsey & Whittow, 1981).

The proportion (13.0-14.1%) of respiration inhibited by ouabain did not differ significantly between treatment groups (Table III.1). This is at the lower end of the

14.3-21.6% inhibition range reported previously for intact mouse soleus muscle (Gregg & Milligan, 1980) and for sliced skeletal muscle preparations from warm and cold exposed mice (Stevens & Kido, 1974).

Muscle from cold exposed mice had a  $\text{Na}^+, \text{K}^+$ -ATPase-dependent respiration value 32% greater ( $p < 0.001$ ), and a  $\text{Na}^+, \text{K}^+$ -ATPase-independent value 20% greater ( $p < 0.001$ ) than those of warm exposed mice (Table III.1). Increased energy expenditure at the level of the  $\text{Na}^+, \text{K}^+$ -ATPase accounted for 20% of the increased muscle  $\text{O}_2$  consumption from cold exposed mice. This value is similar to that reported for sliced mouse skeletal muscle preparations (Stevens & Kido, 1974) but lower than the 54-83% values estimated for the proportion of cold-induced thermogenesis accounted for by  $\text{Na}^+, \text{K}^+$ -ATPase activity for sliced muscle preparations from cold exposed rats (Guernsey & Stevens, 1977; Guernsey & Whittow, 1981). We are not able to conclude that the role of the  $\text{Na}^+, \text{K}^+$ -ATPase in the increased muscle respiration induced by cold exposure differs for mice and rats because the extent to which  $\text{Na}^+-\text{K}^+$  transport was actually inhibited in the intact soleus muscle preparation under our experimental conditions is not known. The inhibition of respiration by ouabain is a function of both enzyme availability to ouabain and the equilibrium reached between formation and dissociation of the inhibitor-enzyme complex (Tobin & Brody, 1972) as well as the activity of  $\text{Na}^+, \text{K}^+$ -ATPase. Incomplete inhibition of the  $\text{Na}^+, \text{K}^+$ -ATPase

would underestimate the contribution of the  $\text{Na}^+, \text{K}^+$ -ATPase to the increased  $\text{O}_2$  consumption observed in the muscle of cold exposed mice. Although our current results indicate that a significant proportion (20%) of the increase in mouse muscle  $\text{O}_2$  consumption induced by cold exposure was in support of energy expenditure and heat production at the level of the  $\text{Na}^+, \text{K}^+$ -ATPase, we are not able to support the suggestion of Guernsey & Stevens (1977) that this is the major mechanism of increased thermogenesis.

In conclusion, cold exposure was shown to increase total  $\text{O}_2$  consumption of intact soleus muscle preparations from mice. Increased energy expenditure at the level of the  $\text{Na}^+, \text{K}^+$ -ATPase accounted for 20% of the cold-induced increase of muscle  $\text{O}_2$  consumption. It is not known how much of the  $\text{Na}^+, \text{K}^+$ -ATPase-independent respiration is due to incomplete inhibition of active  $\text{Na}^+-\text{K}^+$  transport in the intact mouse soleus muscle preparation or to as yet quantitatively unidentified energy expending metabolic processes.

#### **E. Acknowledgements**

Partial financial support from the Natural Sciences and Engineering Research Council is gratefully acknowledged.



**F. References**

- Gregg, V. A. & Milligan, L. P. (1980a) *Gen. Pharmac.* 11, 323.
- Gregg, V. & Milligan, L. P. (1980b) *Biochem. Biophys. Res. Commun.* 95, 608.
- Guernsey, D. L. & Stevens, E. D. (1977) *Science* 196, 908.
- Guernsey, D. L. & Whittow, C. G. (1981) *J. therm. Biol.* 6,7.
- Kohn, P. G. & Clausen, T. (1971) *Biochim. Biophys. Acta* 225, 277.
- Stevens, E. D. & Kido, M. (1974) *Comp. Biochem. Physiol.* 47A, 395.
- Tobin, T. & Brody, T.M. (1972) *Biochem. Pharmacol.* 21,1553.

Table III.1  $O_2$  consumption and  $Na^+, K^+$ -ATPase-dependent, and independent, respiration of intact soleus muscle from warm and cold exposed mice.

Group	Animals	Muscle $O_2$ consumption ( $\mu l O_2/mg/h$ )	% Inhibition of $O_2$ consumption by ouabain	$Na^+, K^+$ -ATPase-dependent $O_2$ consumption ( $\mu l O_2/mg/h$ )	$Na^+, K^+$ -ATPase-independent $O_2$ consumption ( $\mu l O_2/mg/h$ )
Warm	4	5.20 $\pm$ 0.25a	13.0 $\pm$ 3.8a	0.67 $\pm$ 0.03a	4.52 $\pm$ 0.22a
Cold	11	6.38 $\pm$ 0.39b	14.1 $\pm$ 2.3a	0.90 $\pm$ 0.05b	5.42 $\pm$ 0.33b

1.  $Na^+, K^+$ -ATPase-dependent respiration = total  $O_2$  consumption  $\times$  inhibition by ouabain.  
 2.  $Na^+, K^+$ -ATPase-independent respiration = total  $O_2$  consumption -  $Na^+, K^+$ -ATPase-dependent respiration.  
 3. Values expressed as means  $\pm$  S.E.  
 a, b Means within a column followed by different letters differ significantly ( $p < 0.05$ ).

#### IV. ROLE OF $\text{Na}^+, \text{K}^+$ -ATPASE IN MUSCULAR ENERGY EXPENDITURE OF WARM AND COLD EXPOSED SHEEP

##### A. Abstract

The role of  $\text{Na}^+, \text{K}^+$ -ATPase in the energy expenditure of sheep skeletal muscle and the influence of exposure to cold on this role was studied. An in vitro preparation of muscle was developed which achieved  $\text{O}_2$  availability and a functional membrane potential. A  $10^{-6}$  M concentration of ouabain yielded a maximum inhibition of respiration of  $38.9 \pm 1.8\%$  using muscle preparations from a random group of sheep. Whole body and muscle  $\text{O}_2$  consumptions and ouabain-sensitive muscle respiration were measured for warm and cold exposed sheep fed at maintenance or 1150 g alfalfa pellets/d. Cold exposure increased whole body and muscle  $\text{O}_2$  consumption. Inhibition of respiration by ouabain was  $37.6 \pm 1.2\%$  and  $41.0 \pm 3.6\%$  for warm and cold exposed sheep fed at maintenance, and  $28.5 \pm 4.0\%$  and  $45.0 \pm 4.0\%$  for warm and cold exposed sheep fed 1150 g alfalfa pellets/day. The increase in the ouabain-sensitive component of respiration accounted for 48-79% of the increased  $\text{O}_2$  consumption of muscle from cold exposed sheep. It was concluded that the  $\text{Na}^+, \text{K}^+$ -ATPase of sheep muscle is a major means of energy expenditure and has an important role in the increased thermogenesis, resulting from cold exposure.

## B. Introduction

The activity of  $\text{Na}^+, \text{K}^+$ -ATPase ( $\text{Na}^+, \text{K}^+$ -dependent adenosine triphosphatase EC 3.6.1.3) in counteracting the transmembrane movement of  $\text{Na}^+$  and  $\text{K}^+$  along their concentration gradients has been identified as an important component of cellular energy expenditure, causing 20-45% of the  $\text{O}_2$  uptake of resting cells (Whittam, 1961). Activation of  $\text{Na}^+, \text{K}^+$ -ATPase has also been suggested to be an important mechanism for heat production in the cold induced thermogenesis of small mammals (Guernsey & Stevens, 1977). The capacity of the  $\text{Na}^+, \text{K}^+$ -ATPase in basal energy expenditures and heat generation in cold induced thermogenesis was initially studied in sliced tissues from small mammals (see Himms-Hagen, 1976). Concern that damaged cells in the sliced tissue preparations may have yielded physiologically unrealistic impressions of *in vivo*  $\text{Na}^+, \text{K}^+$ -ATPase activity prompted the use of intact organ preparations including rat liver (Folke & Sestoft, 1977), isolated rat soleus muscle and adipose tissue (Chinet et al., 1977), and mouse soleus and diaphragm muscles (Biron et al., 1979; Gregg and Milligan, 1980). With the intent of achieving further definition of the metabolic components of resting energy expenditure and of energy expenditure in the cold, the objectives of this study were to measure  $\text{O}_2$  consumption and  $\text{Na}^+, \text{K}^+$ -ATPase activity in functionally intact sheep muscle preparations and to ascertain the influence of cold exposure of the donor animal, and the

consequent increase in metabolic rate, on these measurements. To achieve these objectives, an in vitro preparation of skeletal muscle was developed in which a functional membrane potential was maintained and there was availability of  $O_2$  to all sites of cellular utilization.

#### C. Experimental

##### Muscle preparation

Sheep were anesthetized with 2.0-2.5% halothane gas. A lengthwise incision was made along the neck exposing the sternomandibularis muscle. A longitudinal section of muscle, approximately 5 mm in diameter and 30 mm in length, was tied at each end with 5-0 braided silk suture and bluntly dissected loose in such a way as to prevent leakage from the tied muscle fibers when a cut was made distal to each tie. The muscle section was removed from the animal and placed in ice-cold buffer solution. With the use of a dissecting microscope, the interjacent fascia was removed and the section re-tied into up to six muscle preparations varying in length from 10-25 mm and less than 0.5 mm in thickness; the approximate calculated thickness at which the rate of  $O_2$  diffusion becomes limiting to respiration (Kleiber, 1961).

##### Electrophysiological studies

The muscle preparations were held on the stage of a fixed stage microscope. Medium was continuously changed by a flow through system supplying Krebs-Ringer bicarbonate

buffer oxygenated with 95%O<sub>2</sub>:5%CO<sub>2</sub> at 37°C (Bonkowski & Runion, 1976). The recording microelectrode was mounted on a Prior micromanipulator and cells were penetrated under direct observation. The glass microelectrodes were filled with 3M KCl and connected through a chlorided Ag wire to a WPI M750 electrometer. Electrode resistances were 20-40 MΩ. The D.C. potentials were observed on a Gould Digital Storage Oscilloscope OS4000.

#### Muscle characterization

Preparations of the sternomandibularis were characterized according to fiber type (Guth and Samaha, 1970). This method identifies muscle fiber types on the basis of qualitative differences in actomyosin ATPase content.

#### Measurement of sheep muscle respiration and response to ouabain

Muscle preparations were obtained from a random group of eight Suffolk crossbred sheep: four 3-5 year old non-pregnant, non-lactating ewes, two 3-4 year old wethers, and one 2 year old ram. One or two sections were taken per animal per surgery and two to four surgeries were performed per animal. The HEPES buffer used as the incubation medium contained (mM): NaCl, 116.8; KCl, 5.9; NaHCO<sub>3</sub>, 10.0; MgSO<sub>4</sub>, 1.2; NaH<sub>2</sub>PO<sub>4</sub>, 1.2; CaCl<sub>2</sub>, 1.0; HEPES (N-2-hydroxyethylpiperazine-N'-2-ethanesulfonic acid), 20.0

(pH 7.3-7.4); acetate, 5.0; glucose, 10.0. To achieve thorough oxygenation, muscle preparations were equilibrated and later individually incubated in 600 ml beakers containing 25 ml buffer in a shaking water bath at 37°C. Muscle preparations were transferred from the equilibration beaker to the electrode chamber and O<sub>2</sub> consumption measured with a YSI O<sub>2</sub> electrode. Each muscle preparation was incubated for 45 min in one concentration of ouabain within the range of 0 to 10<sup>-9</sup> M. The rate of O<sub>2</sub> consumption was measured at the end of the incubation period, following which the preparations were dried overnight at 80°C for tissue dry weight determination. Percent inhibition of respiration by ouabain was calculated using the ratio of the difference between initial and that after incubation to the initial respiration rate. Mean percent inhibition of respiration by ouabain was calculated for each concentration of ouabain in which muscle preparations were incubated. A dose response curve was constructed expressing inhibition as a percentage of the maximum inhibition.

**Trial 1. Respiration and ouabain-sensitive respiration of muscle preparations from warm and cold exposed sheep fed at maintenance**

Eight Suffolk crossbred female sheep, in the weight range of 33-37 kg, were shorn and randomly divided into two groups. Sheep were individually housed in metabolic crates at either 1°C (cold exposed) or 25°C (warm exposed) for 5

weeks prior to muscle sampling. Warm and cold exposed sheep were fed to maintain body weight. Feeding levels were 950 and 1450 g alfalfa pellets/d for sheep exposed to 25°C and 1°C, respectively. Whole animal O<sub>2</sub> consumption was measured using the respiratory gaseous exchange analysis system described by Young et al. (1975). O<sub>2</sub> consumption was measured at 25°C and 1°C for warm and cold exposed sheep, respectively, for a 30 min period 24 h after feeding. Three measurements were made per animal before, during, and after the week of muscle sampling. Respired gases were collected by ventilated hood. Surgery was performed on the cold exposed sheep under warm (25°C) and cold (1°C) conditions. One surgery was performed per animal yielding tissue used to obtain five observations of muscle O<sub>2</sub> consumption and one observation of inhibition of respiration by ouabain at each concentration of inhibitor. Values for percent inhibition of muscle respiration by ouabain were determined as described. Muscle preparations were exposed to ouabain in the range of 0 to 10<sup>-5</sup>M since ouabain concentrations of 10<sup>-6</sup>M or greater were shown to yield maximum inhibition of respiration for excised sheep muscle (Fig. 1). Dose response curves were constructed for each animal.



**Trial 2. Respiration and ouabain-sensitive respiration of muscle preparations from warm and cold exposed sheep fed at the same level of feed intake**

The same experimental outline was followed as in Trial 1 with the same experimental animals except that all sheep were given 1150g/day of the pelleted alfalfa diet and all surgeries were performed under warm (25°C) conditions.

#### **Na<sup>+</sup>,K<sup>+</sup>-ATPase Assay**

The Na<sup>+</sup>,K<sup>+</sup>-ATPase activity of muscle homogenates was measured for sheep of trials 1 and 2 according to a modification of the method of Lo et al.(1976). Samples of the sternomandibularis muscle, approximately 30 mg in weight, were taken from animals during surgery and immediately frozen in liquid N<sub>2</sub>. Frozen muscle samples were homogenized in 10 volumes of the homogenizing medium with a Tekmar homogenizer. Crude homogenate (0.1 ml) was added to 0.9 ml of the reaction mixture in the presence and absence of 0.02 M KCl. After incubation, the inorganic phosphate (Pi) content of the reaction mixture was determined by the method of Fiske & Subbarow (1925). Na<sup>+</sup>,K<sup>+</sup>-ATPase activity was measured as the difference between Pi generated in the presence and absence of K<sup>+</sup>. The protein concentration of the muscles was determined by the method of Lowry et al.(1951).

### Statistical analysis

Values for  $K_i$ , the concentration of ouabain giving one-half maximal inhibition of  $\text{Na}^+, \text{K}^+$ -ATPase-dependent respiration, were calculated from a least-squares regression line fitted through the linear portion of the individual dose response curves for the ouabain effect on muscle respiration. Values for  $K_i$  and means and their standard errors for total, ouabain-sensitive and ouabain-insensitive respiration, and  $\text{Na}^+, \text{K}^+$ -ATPase activities were evaluated for statistical differences by the unpaired Student's t-test (Steel & Torrie, 1960).

### D. Results

Halothane gas was used as the surgical anaesthetic since it has been shown in isolated nerve-muscle preparations that membrane potentials regain pre-exposure status shortly after removal of the volatile gas (MacGregor, 1978). Membrane potentials, measured during development of the intact muscle technique, were -50 mV, or more negative. The sternomandibularis of sheep is a predominantly red type muscle and contained 66% red and 33% white fibers; intermediate type fibers were not present.

### Response of sheep muscle respiration to ouabain

During the course of a day's measurements, some of the muscle preparations were held in the equilibration beaker for up to 3 h before measurement of initial rate of  $\text{O}_2$ .

consumption. No consistent decrease in the rate of  $O_2$  consumption occurred during this period. The mean initial respiration rate of the sheep muscle preparations was  $2.65 \pm 0.11 \text{ } \mu\text{l } O_2/\text{mg dry wt/h}$  ( $n=34$ ). The dose response curve constructed for ouabain inhibition of respiration of sheep muscle is shown in Fig. IV.1. The rate of  $O_2$  consumption of control muscle preparations incubated in buffer without ouabain for 45 min did not change ( $P>0.05$ ) from the initial rates of respiration. The lowest concentration of ouabain resulting in maximum inhibition of respiration was  $10^{-6}\text{M}$ . Values for inhibition of respiration at greater concentrations of ouabain were not different ( $P>0.05$ ) from those obtained with  $10^{-6}\text{M}$  ouabain. Inhibition by  $10^{-6}\text{M}$  ouabain was  $38.9 \pm 1.8\%$  of the total respiration of the muscle. The calculated  $K_i$  value was  $0.25 \text{ } \mu\text{M}$ .

#### **Trial 1. Respiration and ouabain-sensitive respiration of muscle preparations from warm and cold exposed sheep fed at maintenance**

The body weights of both the warm and cold exposed sheep were maintained throughout the experimental period. Whole animal  $O_2$  consumption rates were  $8.9 \pm 1.0$  and  $12.6 \pm 1.0 \text{ l } O_2/\text{h}$ , for warm and cold exposed sheep, respectively (Table IV.1). The increased metabolic rate of the cold exposed sheep was reflected in the initial respiration rates of their muscle preparations; the respiration rates were  $1.82 \pm 0.11$  and  $2.69 \pm 0.16 \text{ } \mu\text{l } O_2/\text{mg dry wt/h}$  for the

preparations from warm and cold exposed sheep, respectively. Ouabain-sensitive respiration was  $37.6 \pm 1.2\%$  and  $41.0 \pm 3.6\%$  of total respiration in the muscles of warm and cold exposed sheep, respectively. Both the ouabain-sensitive and ouabain-insensitive components of respiration of muscle preparations were significantly increased ( $P < .001$ ) by cold exposure of sheep fed at maintenance; there was an increase of 69% in the ouabain-sensitive component, while the ouabain-insensitive component increased 39% (Fig. IV.2). The increase in ouabain-sensitive respiration accounted for 48% of the cold induced increase in muscle respiration. The dose response curves for the warm and cold exposed groups are shown in Fig. IV.3.  $K_i$  values were calculated for each animal; the mean  $K_i$  did not differ significantly ( $p > 0.05$ ) between treatment groups. The overall mean  $K_i$  for the ouabain inhibition of muscle respiration for preparations from the warm and cold exposed sheep was  $0.19 \pm 0.06 \mu M$ .

#### **Trial 2. Respiration and ouabain-sensitive respiration of muscle preparations from warm and cold exposed sheep fed at the same level of intake**

The warm exposed sheep gained an average of 37 g/d and the cold exposed sheep lost an average of 33 g/d during the experimental period while receiving 1150g alfalfa pellets per day. Whole animal respiration rates were  $11.1 \pm 1.0$  and  $13.8 \pm 1.0$  l O<sub>2</sub>/h, for warm and cold exposed sheep, respectively (Table IV.2). Surgeries were performed under

warm (25°C) conditions since in two instances of doing surgery in the cold in Trial 1, difficulties of surgery were increased and no effect of surgery room temperature on initial and ouabain-sensitive respiration was observed. Initial respiration rates were  $2.09 \pm 0.17$  and  $3.09 \pm 0.19$   $\mu\text{l O}_2/\text{mg dry wt/h}$  for the muscle preparations from the warm and cold exposed sheep, respectively. Ouabain-sensitive respiration was  $28.5 \pm 4.0\%$  of total respiration in the warm exposed sheep and  $45.0 \pm 4.0\%$  in the cold exposed sheep. The ouabain-sensitive component of respiration of muscle preparations was increased 132% in cold exposed sheep, accounting for 79% of the total increase in muscle respiration due to cold exposure (Fig. IV.2). The ouabain-insensitive component of respiration did not differ ( $P>0.05$ ) between warm and cold exposed sheep fed at the same level of intake. The mean dose response curves for the warm and cold exposed groups are shown in Fig IV.4.  $K_i$  values did not differ ( $P>0.05$ ) between treatment groups; the overall mean  $K_i$  value of warm and cold exposed sheep fed at the same level of intake was  $0.20 \pm 0.03 \mu\text{M}$ .

#### **Na<sup>+</sup>,K<sup>+</sup>-ATPase assay**

In animals fed at maintenance and at the same level of intake, the mean Na<sup>+</sup>,K<sup>+</sup>-ATPase activity of muscle from warm exposed sheep ( $1.31 \pm 0.26 \mu\text{mole Pi/mg protein/h}$ ) did not differ ( $P>0.05$ ) from that of muscle from cold exposed sheep ( $1.48 \pm 0.26 \mu\text{mole Pi/mg protein/h}$ ).

## E. Discussion

The difficulty of obtaining tissue slice preparations of muscles from large animals which would not contain a large proportion of damaged cells prompted the development of a preparation that more closely resembled the *in vivo* muscle state, for example, by maintaining a functional membrane potential. The small size of this muscle preparation allowed repeated sampling of each animal and avoided problems associated with ouabain perfusion of whole organ preparations, such as loss of vascular integrity and oxygenation (Folke & Sestoft, 1977).

Ouabain, the specific inhibitor of  $\text{Na}^+, \text{K}^+$ -ATPase, inhibited 40% of the respiration of excised sheep muscle, supporting the role of  $\text{Na}^+, \text{K}^+$ -ATPase as a major component of cellular energy expenditure. This measurement is in agreement with values reported for the ouabain-sensitive respiration of rat diaphragm (Asano et al. 1976; Ismail-Beigi & Edelman, 1970) and mouse soleus and diaphragm muscles (Gregg & Milligan, 1980). The sigmoidal dose response curves obtained for ouabain inhibition of the  $\text{Na}^+, \text{K}^+$ -ATPase-dependent respiration of sheep muscle (Figs. IV.3, IV.4), are similar to the pattern reported for purified  $\text{Na}^+, \text{K}^+$ -ATPase preparations from other mammals (Charnock & Simonson, 1977, 1978). The  $K_i$  value of  $0.25 \mu\text{M}$  calculated for ouabain inhibition of respiration of muscle from a random group of sheep is within the range of  $0.11$ - $2.50 \mu\text{M}$  reported for purified  $\text{Na}^+, \text{K}^+$ -ATPase

preparations for several mammalian tissues (Charnock & Simonson, 1977, 1978; Tobin & Brody, 1972).

Cold induced thermogenesis, as studied in small mammals, is thought to be a consequence of increased heat production by brown adipose tissue (Horwitz, 1979). The increased thermogenesis induced by cold exposure appears to be mediated by a complex interaction involving neural and hormonal factors and includes increased thyroid activity. (Horwitz, 1979). In larger animals containing little, if any, brown adipose tissue, muscle becomes an important site of heat production when an increased thermal demand is imposed (Jansky, 1973). Indeed, we found that the rate of  $O_2$  uptake of muscle preparations from cold exposed sheep was greater than uptake by preparations from warm exposed sheep. It is then of importance to ascertain the mechanisms within the tissue by which the increased metabolic rate is achieved. The results obtained using functionally intact preparations indicate that 48-79% of the increased  $O_2$  consumption observed in muscle preparations from cold exposed sheep is attributable to increased  $Na^+$ ,  $K^+$ -ATPase activity. In principle, this is in agreement with the evidence derived with damaged muscle preparations that a major portion of the increased  $O_2$  consumption of muscle tissue from cold exposed animals results from increased energy expenditure by means of  $Na^+$ ,  $K^+$ -ATPase activity (Guernsey & Stevens, 1977; Stevens & Kido, 1974).

The mechanism of enhancement of  $\text{Na}^+, \text{K}^+$ -ATPase activity of muscle from cold exposed animals is not clear at this time. Stimulation of the activity of pre-existing enzyme or hormonal induction of increased enzyme synthesis or of the synthesis of a more active enzyme are possible ways of achieving the observed increase of the  $\text{Na}^+, \text{K}^+$ -ATPase activity of muscle from cold exposed animals. There is evidence of increased thyroid activity in sheep subjected to cold stress (Westra & Christopherson, 1976) and  $^3\text{H}$ -ouabain binding studies have indicated increased amounts of  $\text{Na}^+, \text{K}^+$ -ATPase in the muscles of rats upon treatment with thyroid hormone (Lin & Akera, 1978), thus one might expect increased amounts of  $\text{Na}^+, \text{K}^+$ -ATPase in the skeletal muscle of cold exposed sheep. However, overall  $\text{Na}^+, \text{K}^+$ -ATPase activities, measured in this study as an index of the maximum capacity of the enzyme system, did not differ significantly ( $p > 0.05$ ) between warm and cold exposed sheep. The present study does not provide conclusive evidence for resolution of the possibilities of increased activity of existent enzyme versus an increased amount of enzyme as the cause of the increase in  $\text{Na}^+, \text{K}^+$ -ATPase-dependent respiration in the muscle of cold exposed sheep.

The kinetic constant  $K_i$ , a measure of enzyme sensitivity to inhibitor, may be a useful tool in the assessment of changes in enzyme characteristics occurring with different physiological stresses. The lack of difference between calculated  $K_i$  values from warm and cold



exposed sheep fed at maintenance or at the same level of intake may indicate a lack of change in the nature of the enzyme in the cold exposed animals.

In designing a study of effects of cold stress on animals, an investigator is faced with the dilemma of providing additional dietary energy intake to the cold exposed animals to offset the imposed thermal demand, or of simply providing the same amount of feed to the two experimental groups. The former choice results in warm and cold exposed groups at differing levels of intake, as in Trial 1, while the latter choice results in warm and cold exposed groups that differ quantitatively in the energy they can expend on metabolic processes other than those expressly for heat production, as in Trial 2, in which the cold exposed group lost weight. Thus, to arrive at a reliable conclusion, it is of importance to have examined effects under both of the foregoing experimental circumstances. Considerable confidence arises upon having obtained similar findings regarding  $\text{Na}^+, \text{K}^+$ -ATPase activity under both experimental circumstances. It is concluded that expenditure of energy by  $\text{Na}^+, \text{K}^+$ -ATPase-catalyzed transport of  $\text{Na}^+$  and  $\text{K}^+$  across the plasma membrane, has a highly significant involvement in the basal energy expenditure of sheep muscle and that a major part of the increased respiration, and presumably heat generation, of muscle from cold exposed sheep was due to increased  $\text{Na}^+, \text{K}^+$ -ATPase activity.

## F. Acknowledgments

Grateful acknowledgement is extended to the Natural Sciences and Engineering Research Council Canada for support in part of this research; to Dr. W. F. Dryden for assistance in the measurement of membrane potentials, and to Dr. A. M. Nicol for assistance in the measurement of whole animal O<sub>2</sub> consumptions.

## G. References

Asano, Y., Liberman, U. A. & Edelman, I. S. (1976) J. Clin. Invest. 57, 368.

Biron, R., Burger, A., Chiner, A., Clausen, T. & Dubois-Ferriere, R. (1979) J. Physiol. 297, 47.

Bonkowski, L. & Runion, H. J. (1976) Experientia 32, 1619.

Charnock, J. S. & Simonson, L. P. (1977) Comp. Biochem. Physiol. 58B, 381.

Charnock, J.S. and Simonson, L. P. (1978) Comp. Biochem. Physiol. 59B, 223.

Chinet, A., Clausen, T. & Girardier, L. (1977) J. Physiol. 265, 43.

Fiske, C.R. & Subbarow, Y. (1925). J. Biol. Chem. 66, 375.

Folke, M. & Sestoft, L. (1977) J. Physiol. 269, 407.

Gregg, V.A. & Milligan, L. P. (1980) Gen. Pharmac. 11, 323.

- Guernsey, D.L. & Stevens, E. D. (1977) *Science* 196: 908-910
- Guth, L. & Samaha, F. J. (1970) *Exp. Neurol.* 28, 365.
- Himms-Hagen, J. (1976) *Ann. Rev. Physiol.* 38, 315.
- Horwitz, B.A. (1979) *Fed. Proc.* 38, 2170.
- Ismail-Beigi, F. & Edelman, I. S. (1970) *Proc. Nat. Acad. Sci. U.S.A.* 67, 1071.
- Jansky, L. (1973) *Biol. Rev.* 48, 85.
- Kleiber, M. (1961) *The Fire of Life*. New York, John Wiley & Sons, Inc., p. 66.
- Lin, M.H. & Akera, T. (1978) *J. Biol.Chem.* 253, 723.
- Lo, C.C., August, T. R., Liberman, U. A. & Edelman, I. S. (1976) *J. Biol. Chem.* 251, 7826.
- Lowry, O.H., Rosebrough, N. J., Farr, A. L. & Randal, R. (1951) *J. Biol. Chem.* 193, 265.
- MacGregor, M.H.G. (1978) The effects of three inhalation anaesthetics on neuromuscular transmission. Ph.D. Thesis, University of Glasgow, Scotland., pgs. 66-96.
- Steel, R. G. D. & Torrie, J. H. (1960) *Principles of Statistics*. New York: McGraw-Hill, Inc.
- Stevens, E.D. & Kido, M. (1974) *Comp. Biochem. Physiol.* 47A, 395.
- Tobin, T. & Brody, T. M. (1972) *Biochem. Pharmac.* 21, 1553.
- Westra, R. & Christopherson, R. J. (1976) *Can. J. Anim. Sci.* 56, 699.

Whittam, R. (1961) *Nature* 191, 603.

Young, B. A., Kerrigan, B. & Christopherson, R.J. (1975)  
*Can. J. Anim. Sci.* 55, 17.

Table IV.1 Whole animal and muscle respiration from sheep fed at maintenance<sup>1</sup>.

Treatment	Whole animal O <sub>2</sub> consumption (l O <sub>2</sub> /h)	Muscle O <sub>2</sub> consumption ( $\mu$ l O <sub>2</sub> /mg/h)	% Inhibition by ouabain of muscle O <sub>2</sub> consumption <sup>2</sup>
Warm	8.9 $\pm$ 1.0	1.82 $\pm$ 0.11	37.6 $\pm$ 1.2
Cold	12.6 $\pm$ 1.0*	2.69 $\pm$ 0.16**	41.0 $\pm$ 3.6

<sup>1</sup> Maintenance feed intakes were 950 and 1450 g alfalfa pellets/d for warm and cold adapted sheep, respectively.

<sup>2</sup> ouabain=10<sup>-6</sup>M.

\* P<0.05

\*\* P<0.001

Table IV.2 Whole animal and muscle respiration from sheep fed at the same level of intake<sup>1</sup>.

Treatment	Whole animal O <sub>2</sub> consumption (l O <sub>2</sub> /h)	Muscle O <sub>2</sub> consumption (μl O <sub>2</sub> /mg/h)	% Inhibition by ouabain of muscle O <sub>2</sub> consumption <sup>2</sup>
Warm	11.1 ± 1.0	2.09 <sup>b</sup> ± 0.17	28.5 ± 4.0
Cold	13.8 ± 1.0	3.09 ± 0.19**	45.0 ± 4.0**

<sup>1</sup> 1150 g alfalfa pellets/d

<sup>2</sup> ouabain=10<sup>-6</sup>M.

\*\* P<0.001




Figure IV.1. Relative inhibition of sheep muscle respiration by ouabain expressed as a percentage of inhibition at  $10^{-6}\text{M}$  ouabain (maximum inhibition). Values are means  $\pm$  S.E.




Figure IV.2. Total respiration and ouabain-sensitive (hatched) and insensitive (clear) respiration of muscle from warm and cold exposed sheep fed at maintenance or 1150 g alfalfa pellets/d. Values are means  $\pm$  S.E.






Figure IV.3. Relative inhibition by ouabain of respiration of muscle from warm and cold exposed sheep fed at maintenance. Inhibitions expressed as a percentage of inhibition at  $10^{-6}$ M ouabain (maximum inhibition). Values are means  $\pm$  S.E.




Figure IV.4. Relative inhibition by ouabain of respiration of muscle from warm and cold exposed sheep fed 1150 g alfalfa pellets/d. Inhibitions expressed as a percentage of inhibition at 10<sup>-6</sup>M ouabain (maximum inhibition). Values are means ± S.E.

V. ENERGY COSTS OF  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase ACTIVITY AND PROTEIN SYNTHESIS IN MUSCLE FROM CALVES DIFFERING IN AGE AND BREED

A. Abstract

An in vitro preparation was used to measure rates of  $\text{O}_2$  consumption,  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase-dependent respiration,  $^{14}\text{C}$ -phenylalanine incorporation and tyrosine release of skeletal (sternomandibularis) muscle from 10-21d and 7 month dairy calves, and control and extreme double-muscled (DM) calves. Rate of  $\text{O}_2$  consumption was greatest ( $p < 0.001$ ) for muscle from 10-21d dairy calves and lowest ( $p < 0.05$ ) for control DM calves. Ouabain ( $10^{-5}\text{M}$ ) caused a 40% inhibition of muscle respiration.  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase-dependent respiration was similar for muscle from all calf groups except 10-21d dairy calves which had a value 26% greater ( $p < 0.001$ ) than that of older dairy calves.  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase-independent respiration was 16% greater ( $p < 0.001$ ) for muscle from 10-21d than that of older dairy calves while muscle from extreme DM calves had a value 11% greater than that of control DM calves. The rate of  $^{14}\text{C}$ -phenylalanine incorporation was greater ( $p < 0.05$ ) for muscle from 10-21d dairy than from older dairy calves, similar between older dairy and control DM calves, and decreased ( $p < 0.05$ ) for extreme DM calves. Rate of tyrosine release was greatest ( $p < 0.05$ ) for muscle from control and extreme DM calves; both dairy groups had similarly low rates of muscle tyrosine release. The energy estimated to be required for peptide bond synthesis

accounted for 2.0 to 3.3% of the  $O_2$  consumption of the muscle preparations.

## B. Introduction

In order to more fully understand whole animal energy expenditure it is necessary first to identify the causes of metabolic energy expenditure, and then to determine their quantitative importance under a variety of physiological conditions.

Active  $Na^+$ - $K^+$  transport, that is, the activity of the plasma membrane  $Na^+$ , $K^+$ -ATPase (EC 3.6.1.3) in counteracting transmembrane movement of  $Na^+$  and  $K^+$  along their concentration gradients, has been suggested to be a major component of the energy expenditure of animals and has been estimated to account for 20-45% of the  $O_2$  uptake of resting cells (Whittam, 1961). Protein synthesis has also been suggested to be a major energy cost of animals, accounting for up to 30% of the heat production of cattle (Lobley et al. 1980). The extent to which energy expended by processes such as active  $Na^+$ - $K^+$  transport and protein synthesis can vary between animals and is influenced by genetic and environmental factors is not clear. Evidence that exposure of animals to a cold environment selectively increased energy expenditure at the level of the  $Na^+$ , $K^+$ -ATPase has been presented for muscle preparations from sheep (Chapter IV).

The objectives of this experiment were to obtain physiologically realistic estimates of the magnitudes of the energy costs of active  $\text{Na}^+$ - $\text{K}^+$  transport and protein synthesis in skeletal muscle from calves and to examine the effects of breed and age upon the relative costs of these two processes as components of background or maintenance energy expenditure of the tissue.

### C. Experimental

#### Animals

Muscle samples were obtained from six 7 month old male calves from a beef crossbred population selected for a high incidence of double-muscling (DM), two calves exhibited normal muscling (control DM) and four calves exhibited overt muscular hyperplasia (extreme DM); three 10-21d old male dairy (Holstein) calves; and three 7 month old male dairy calves. All animals were housed indoors in heated barns (approximately  $20^\circ\text{C}$ ) for at least 4 weeks prior to surgery. The 10-21d dairy calves were fed a milk replacer diet; 7 month calves were fed good quality grass hay free choice and a ration of barley-oat concentrate mix containing minerals. Water was available ad libitum. Animals were fasted overnight prior to surgery.

#### Respiration and $\text{Na}^+$ , $\text{K}^+$ -ATPase-dependent respiration

A section of the sternomandibularis muscle was taken from each animal and preparations made following the method

of Gregg & Milligan (Chapter IV). One surgery was performed per animal yielding tissue used to obtain four observations of muscle  $O_2$  consumption and one observation of inhibition by ouabain at each concentration of inhibitor. Immediately upon removal from the animal, muscle sections were placed in cooled ( $15^\circ\text{C}$ ) HEPES buffer containing 10 mM glucose and 5 mM acetate as substrates, and the small tied fibre bundles (approximately  $20.0 \times 0.5$  mm) were prepared at room temperature with the aid of a dissecting microscope. Following measurement of initial respiration rates (Chapter IV) in an  $O_2$  electrode system, the muscle preparations were incubated in buffer containing 0 to  $10^{-4}$  M ouabain. Respiration rates were then measured again. Percent inhibition of respiration by ouabain was calculated using the ratio of the difference between initial and post-incubation respiration rates to the initial respiration rate.

#### Measurement of $^{14}\text{C}$ -phenylalanine incorporation

Muscle sections were placed in HEPES buffer to which the following additions had been made: essential and non-essential amino acids, except phenylalanine, at the concentrations reported for sheep plasma (Bergman et al. 1974); phenylalanine, 500  $\mu\text{M}$ ; insulin, 0.1 unit per ml; chloramphenicol, 0.3 mg per l. Four muscle preparations from each animal were incubated in 3 ml of the complete HEPES buffer containing approximately 0.28  $\mu\text{Ci}$  of L-(U- $^{14}\text{C}$ )-

phenylalanine (Amersham Corp., Ontario) per ml, two preparations were incubated for 1.0 h and two for 2.5 h in a shaking water bath at 37°C. Preliminary studies established the rate of <sup>14</sup>C-phenylalanine incorporation to be linear for incubations periods of up to 3.0 h. At the end of the incubation period the muscle preparations were rinsed, blotted and weighed. They were then homogenized in 1 ml of cold 50% (w/v) trichloroacetic acid, centrifuged, and the precipitate washed according to the method of Fulks et al. (1975). Acid-precipitated pellets were combusted in a Beckman biological material oxidizer and <sup>14</sup>CO<sub>2</sub> collected in 10 mls of CO<sub>2</sub>-trapping cocktail (50%, v/v, toluene, 30% methyl cellosolve, 20% monoethanolamine, 5.0 g PPO, 0.2 g POPOP). Radioactivity was measured with a Searle Mark III liquid scintillation counter and counting efficiency determined with the channels-ratio method.

In a separate experiment, acid-precipitated material from muscle preparations incubated 2.5 h was dried, hydrolyzed and chromatographed according to the method of McBride et al. (1979). Sections of the thin layer chromatography plates were scraped into separate counting vials and counted as described above. Radioactivity was found to be present only at the position corresponding to the phenylalanine standard.

### Measurement of tyrosine release

The rate of tyrosine release from muscle was measured in the HEPES buffer used for the study of  $^{14}\text{C}$ -phenylalanine incorporation, in which was included 0.5 mM cycloheximide and from which tyrosine was omitted, according to the method of Fulks et al. (1975). Two muscle preparations from each animal were pre-incubated in 2 ml of buffer in a shaking water bath at  $37^\circ\text{C}$  for 0.5 h then transferred to similar flasks; one preparation was incubated for 0.5 h and the other for 1.5 h. At the end of the incubation period, muscle preparations were blotted and weighed. The amount of tyrosine in the buffer was measured fluorometrically by the method of Waalkes & Udenfriend (1957). Preliminary studies established the rate of tyrosine release from muscle preparations into buffer to be linear for incubation periods of up to 2.5 h after a 0.5 h pre-incubation period.

### Statistical analysis

Results are expressed as the mean values and their standard errors. An unpaired Student's t test was used to establish the significance of differences between the means for muscle  $\text{O}_2$  consumption, percent inhibition of respiration by  $10^{-5}\text{M}$  ouabain, and  $\text{Na}^+, \text{K}^+$ -ATPase-dependent and independent respiration. The rates of  $^{14}\text{C}$ -phenylalanine incorporation and tyrosine release were determined by least squares analysis. Differences between groups were located by the Student-Newman-Kuel range test (Steel & Torrie, 1960).



#### D. Results

##### O<sub>2</sub> consumption and Na<sup>+</sup>,K<sup>+</sup>-ATPase activity

Total muscle O<sub>2</sub> consumption and the proportion of respiration inhibited by 10<sup>-5</sup>M ouabain are shown in Table V.1.

The rate of muscle O<sub>2</sub> consumption was greatest ( $p < 0.05$ ) for 10-21d dairy calves and was reduced by 16% in the muscle from the older dairy calves. Among calves of similar age, dairy calves had a muscle O<sub>2</sub> consumption rate greater ( $p < 0.05$ ) than control DM calves.

Dose-response curves constructed for each calf group differed in absolute values for percent inhibition of respiration observed at each concentration of inhibitor but had a similar sigmoidal shape and a similar value for the lowest concentration of inhibitor yielding maximum inhibition (10<sup>-5</sup>M ouabain).

The proportion of respiration inhibited by ouabain ranged from 39.4 ± 2.9% for older dairy to 42.7 ± 1.4% for control DM calves. Na<sup>+</sup>,K<sup>+</sup>-ATPase-dependent respiration, the amount of O<sub>2</sub> calculated to have been consumed to support the ouabain-inhibitable portion of Na<sup>+</sup>,K<sup>+</sup>-ATPase activity, was statistically similar for muscle from all calf groups except 10-21d dairy calves which had a value 26% greater ( $p < 0.001$ ) than that of older dairy calves.

Na<sup>+</sup>,K<sup>+</sup>-ATPase-independent respiration, the residual portion of muscle O<sub>2</sub> consumption not accounted for by activity of the Na<sup>+</sup>,K<sup>+</sup>-ATPase, was greatest for muscle from

10-21d dairy calves. The value for muscle from 10-21d dairy calves was 16% greater ( $p < 0.001$ ) than that of older dairy calves. For calves of similar age,  $\text{Na}^+, \text{K}^+$ -ATPase-independent respiration was measured to be 11% greater ( $p < 0.05$ ) for muscle from extreme than that of control DM calves and 19% greater ( $p < 0.005$ ) for muscle from older dairy than that of control DM calves.

#### Rates of $^{14}\text{C}$ -phenylalanine incorporation and tyrosine release

Rates for  $^{14}\text{C}$ -phenylalanine incorporation into muscle protein are shown in Table V.2. The rate of  $^{14}\text{C}$ -phenylalanine incorporation into muscle differed ( $p < 0.05$ ) between all groups except between older dairy and control DM calves.  $K_s$  values ranged from 0.7 to 1.5%/d (Table V.2); muscle preparations from 10-21d dairy calves had a  $K_s$  value 50% greater than those from older dairy calves, and muscle preparations from control DM calves had a  $K_s$  value 54% higher than those from extreme DM calves. The  $K_s$  values for muscle preparations from older dairy and control DM calves were similar. The estimated cost of incorporation of amino acids into growing peptide chains ranged from 2.0 to 3.3% of the total in vitro  $\text{O}_2$  consumption of the muscle preparations (Table V.2).

Rates of release of tyrosine from muscle preparations are shown in Table V.3. The rate of tyrosine release from muscle differed ( $p < 0.05$ ) between all calf groups except

between 10-21d and older dairy calves. Both dairy groups had tyrosine release rates which were too low to allow accurate measurement. Muscle preparations from extreme DM calves had a greater ( $p < 0.05$ ) rate of tyrosine release (Table V.3) than those from control DM calves.

The fraction of muscle protein degraded per day (Kd) was calculated assuming a muscle tyrosine content of 2.8 mole % (Chang & Goldberg, 1978). Kd values thus calculated were 0.1%/d for 10-21d and older dairy calves, 0.8%/d for control DM calves, and 5.4%/d for extreme DM calves. It was assumed that tyrosine released from muscle originated from muscle protein since Fulks et al. (1975) have shown that under similar experimental conditions muscle free tyrosine content did not change during 3 h of incubation.

#### E. Discussion

Exposure of muscle to  $10^{-5}$ M ouabain resulted in an average of 40% for inhibition of respiration. This value is in agreement with values for inhibition of respiration by ouabain reported for skeletal muscle from sheep (Chapter IV), mice (Gregg & Milligan, 1980a), and rats (Asano et al., 1976; Ismail-Beigi & Edelman, 1970).

The rates of  $O_2$  consumption measured for calf muscle preparations were similar to those previously measured with like preparations of the same muscle for adult sheep (Chapter IV). A decrease of metabolic rate with age has been previously observed for cattle (Webster et al. 1974). The

greater rate of muscle  $O_2$  consumption measured for 10-21d than for older dairy calves resulted from increases in both the  $Na^+, K^+$ -ATPase-dependent and independent components of respiration. For calves of similar age, the differences measured for total  $O_2$  consumption between extreme and control DM calves and between dairy and control DM calves were due to an increased amount of  $O_2$  consumed in the  $Na^+, K^+$ -ATPase-independent component of respiration.

The use of ouabain to determine  $Na^+, K^+$ -ATPase-dependent respiration was criticized (Himms-Hagen, 1976) primarily on the basis that the decreased  $O_2$  consumption observed in the presence of ouabain may be due to altered intracellular  $Na^+$  and  $K^+$  concentrations and may, therefore, be secondary to the inhibition of the  $Na^+, K^+$ -ATPase. However, Asano et al. (1976) found measurements of the ouabain-inhibitable  $O_2$  consumption of rat skeletal muscle to be independent of a wide range of experimentally induced changes in intracellular  $Na^+$  and  $K^+$  concentrations. Also, inhibition by ouabain of the rate of glycolytic substrate-level phosphorylation in ascites tumor cells was shown to have resulted from direct inhibition of  $Na^+, K^+$ -ATPase rather than from a disturbance of glycolytic enzyme function through alterations of intracellular  $K^+$  concentrations (Scholnick et al. 1973). A second objection (Chinet et al. 1977) concerned the use of sliced tissue preparations for measurement of  $Na^+, K^+$ -ATPase-dependent respiration: rapid leakage of  $Na^+$  into the cells at sites of membrane damage may have occurred

and caused a non-physiological stimulation of the  $\text{Na}^+, \text{K}^+$ -ATPase leading to the large values measured for  $\text{Na}^+, \text{K}^+$ -ATPase-dependent respiration. Investigations conducted with intact organ perfusions have measured  $\text{Na}^+, \text{K}^+$ -ATPase-dependent respiration or heat production to account for less than 6% of total tissue  $\text{O}_2$  consumption or heat production (Chinet et al., 1977; Fólke & Sestoft, 1977). In contradiction to this criticism, Gregg & Milligan (1980b) have presented evidence that at least part of the low response to ouabain obtained in the microcalorimetric studies of perfused organs was due to prior inhibition of the enzyme by the experimental conditions.

The importance of tissue preparation and condition in achieving measurements indicative of physiological  $\text{O}_2$  consumption and  $\text{Na}^+, \text{K}^+$ -ATPase activity cannot be ignored. The muscle preparation used in this experiment was developed specifically to study the metabolism of muscle from large mammals under conditions which minimize cellular damage and maintain physiological characteristics such as membrane potential and oxygenation (Chapter IV). The inhibition of respiration induced by ouabain is a function of both enzyme availability to ouabain and the equilibrium reached between formation and dissociation of the inhibitor-enzyme complex (Tobin & Brody, 1972). The proportion of  $\text{Na}^+, \text{K}^+$ -ATPase activity actually inhibited in the muscle preparation under our experimental conditions is not known. Therefore, our measurements of the proportion of muscle  $\text{O}_2$  consumption

required for support of  $\text{Na}^+, \text{K}^+$ -ATPase activity must be considered as minimum estimates.

Accurate calculation of rates of protein synthesis from incorporation of  $^{14}\text{C}$ -phenylalanine into muscle protein requires that the specific activity of phenylalanine-tRNA be known. McKee et al. (1978) have shown for perfused rat heart that the specific activities of extracellular, intracellular, and tRNA-bound phenylalanine are the same when the perfusate phenylalanine concentration was 0.4 mM or greater. It was assumed that protein degradation that may have occurred during the period of incubation did not significantly influence the estimate of synthesis by causing cleavage of newly added phenylalanine from peptide chains.

The calculated values for the fraction of muscle protein synthesized per day ( $K_s$ ) (Table V.2) are in good agreement with  $K_s$  values reported for heifer and cow muscle (0.8-2.0%/d) by Lobley et al. (1980, 1978) as determined by in vivo constant infusion of  $^3\text{H}$ -tyrosine. The greater rate of muscle protein synthesis measured for 10-21d dairy than from older dairy calves is consistent with the higher  $K_s$  values for protein synthesis measured for immature in contrast to adult animals (see Garlick, 1980).

Although the rate of protein synthesis was 50% greater in muscle of 10-21d dairy calves than that of older dairy calves, the increased rate of peptide bond synthesis would account for only 9% of the difference in total muscle  $\text{O}_2$  consumption measured between the two dairy groups. It is not

known how much of the greater  $\text{Na}^+, \text{K}^+$ -ATPase-dependent respiration might have been in support of, or association with, the increased rate of protein synthesis as suggested by Reeds et al. (1980). Older dairy and control DM calf muscle did yield similar values for both the amount of  $\text{O}_2$  consumed in direct support of protein synthesis and  $\text{Na}^+, \text{K}^+$ -ATPase-dependent respiration. However, among 7 month old calves, the rate of protein synthesis in the muscle of control DM calves was 55% greater than in muscle of extreme DM calves while no difference was measured for  $\text{Na}^+, \text{K}^+$ -ATPase-dependent respiration between the two calf groups. These results do not appear to support a close association of increased rate of protein synthesis and of  $\text{Na}^+, \text{K}^+$ -ATPase activity. However, the extent to which  $\text{Na}^+, \text{K}^+$ -ATPase-dependent respiration is underestimated and the extent to which differences in protein degradation may have affected muscle  $\text{O}_2$  consumption are not known. Therefore, it is not possible to reach a conclusion regarding a relationship between rate of protein synthesis and  $\text{Na}^+, \text{K}^+$ -ATPase activity.

Rates of protein turnover should not be calculated from our results since the rates of  $^{14}\text{C}$ -phenylalanine incorporation and of tyrosine release were measured under different experimental conditions, the latter in a medium with tyrosine omitted and cycloheximide added. Nonetheless, estimated rates of protein degradation (Table V.3) were less than, or similar to, rates of protein synthesis (Table V.2)

for muscle preparations from all calf groups except extreme DM calves. The DM muscle preparations exhibited an enhanced rate of tyrosine release which is certainly suggestive of an increased capacity for protein degradation.

DM cattle appear to partition energy in a way that is different from non-double muscled cattle including altered potentials for protein and fat deposition (Holmes & Ashmore, 1972). The metabolic cause of these changes is not known. One might speculate, on the basis of the possibility of increased protein degradation, that muscles of extreme DM calves may expend more energy for protein turnover than those of control DM calves and this might account for the tendency for increased total muscle  $O_2$  consumption. In the animal, increased energy expenditure for such a maintenance function could result in a reduction of energy available for fat deposition.

The available evidence supports the importance of activity of  $Na^+, K^+$ -ATPase as a cause of cellular energy expenditure in muscle. While it would be invalid to conclude that the role of  $Na^+, K^+$ -ATPase is of the identical magnitude in vivo as measured in vitro, the comparison of in vitro and in vivo estimates for rate of protein synthesis does provide some confidence that at least what is measured for these muscle preparations is indicative of what occurs in the animal.

In conclusion, activity of the  $Na^+, K^+$ -ATPase accounted for a minimum of 40% of muscle  $O_2$  consumption. The energy



required for peptide bond synthesis was estimated to account for less than 4% of muscle O<sub>2</sub> consumption. The amount of muscle O<sub>2</sub> consumption due to activity of the Na<sup>+</sup>,K<sup>+</sup>-ATPase was similar for 7 month old calves of different breed backgrounds, and was increased for 10-21d old dairy calves. The amount of O<sub>2</sub> calculated to have been expended on peptide bond synthesis was increased 50% or greater for muscle from 10-21d dairy over that of older dairy calves, and from control DM over that of extreme DM calves. The increased amount of O<sub>2</sub> consumption expended in direct support of protein synthesis would not account for the difference in muscle O<sub>2</sub> consumption measured between the dairy groups.

#### **F. Acknowledgements**

Grateful acknowledgement is extended to the Natural Sciences and Engineering Research Council Canada for support in part of this research; to Dr. J. R. Thompson for assistance in measurement of rates of protein synthesis; and to Dr. D. Hadziyev of the Department of Food Science for access to a spectrophotofluorometer.

#### **G. References**

Asano, Y., Liberman, U.A. & Edelman, I.S. (1976) J. Clin. Invest. 57, 368.

Bergman, E.N., Kaufman, C. F., Wolff, J. E. & Williams, H.H.  
(1974) Am. J. Physiol. 226, 833.

Chang T.W. & Goldberg, A.L. (1978) J. Biol. Chem. 253, 3685.

Chinet, A., Clausen, T. & Girardier, L. (1977) J. Physiol.  
265, 43.

Folke, M. & Sestoft, L. (1977) J. Physiol. 269, 407.

Fulks, R.M., Li, J.B. & Goldberg, A.L. (1975) J. Biol. Chem.  
250, 290.

Garlick, P.J. (1980) in Protein Deposition in Animals,  
(Buttery, P.J. & Lindsay, D.B., eds.) London:  
Butterworths, p.51.

Gregg, V.A. & Milligan, L.P. (1980a) Gen. Pharmac. 11, 323.

Gregg, V. & Milligan, L.P. (1980b) Biochem. Biophys. Res.  
Comm. 95, 608.

Himms-Hagen, J. (1976) Ann. Rev. Physiol. 38, 315.

Holmes, J.H.G. & Ashmore, C.R. (1972) Growth 36, 351.

Ismail-Beigi, F. & Edelman, I.S. (1970) Proc. Natl. Acad.  
Sci. USA 67, 1071.

Lobley, G.E., Reeds, P.J. & Pennie, K. (1978) Br. J. Nutr.  
37, 96A.

Lobley, G.E., Milne, V., Lovie, J.M., Reeds, P.J. &  
Pennie, K. (1980) Br. J. Nutr. 43, 491.

McBride, R.W., Jolly, D.W., Kadis, B.M. & Nelson, T.E.  
(1979) J. Chromatog. 168, 290.

McKee, E. E., Cheung, J. Y., Rannels, E. & Morgan, H. E.  
(1978) J. Biol. Chem. 253, 1030.

Reeds, P.J., Cadenhead, A., Fuller, M.F., Lobley, G.E. &  
McDonald, J.D. (1980) Br. J. Nutr. 43, 445.

Scholnick, P., Lang, D. & Racker, E. (1973) J. Biol. Chem.  
248, 5175.

Steel, R.G.D. & Torrie, J.H. (1960) Principles and  
Procedures of Statistics. New York: McGraw-Hill, Inc.

Tobin, T. & Brody, T. M. (1972) Biochem. Pharmac. 21, 1553.

Waalkes, T.P. & Udenfriend, S. (1957) J. Lab. & Clin. Med.  
50, 733.

Webster, A. J. F., Brockway, J. M. & Smith, J. S. (1974)  
Anim. Prod. 19, 127.

Whittam, R. (1961) Nature 191, 603.

Table V.1 In vitro O<sub>2</sub> consumption, ouabain inhibition and Na<sup>+</sup>, K<sup>+</sup>-ATPase-dependent and independent respiration of calf muscle preparations.

Group	O <sub>2</sub> consumption (ul O <sub>2</sub> /mg/h)	% Inhibition of O <sub>2</sub> consumption by ouabain	Na <sup>+</sup> , K <sup>+</sup> -ATPase-dependent O <sub>2</sub> consumption (ul O <sub>2</sub> /mg/h)	Na <sup>+</sup> , K <sup>+</sup> -ATPase-independent O <sub>2</sub> consumption (ul O <sub>2</sub> /mg/h)
10-21d dairy	3.27 ± 0.27a	41.3 ± 4.5ab	1.36 ± 0.11a	1.93 ± 0.16a
7 month dairy	2.75 ± 0.27b	39.4 ± 2.9b	1.08 ± 0.11b	1.67 ± 0.16b
control DM	2.44 ± 0.24c	42.7 ± 1.4a	1.04 ± 0.12b	1.40 ± 0.16c
extreme DM	2.61 ± 0.27bc	40.6 ± 2.2ab	1.06 ± 0.10b	1.55 ± 0.14b

1. Na<sup>+</sup>, K<sup>+</sup>-ATPase-dependent respiration = total O<sub>2</sub> consumption × inhibition by ouabain.  
 2. Na<sup>+</sup>, K<sup>+</sup>-ATPase-independent respiration = total O<sub>2</sub> consumption - Na<sup>+</sup>, K<sup>+</sup>-ATPase-dependent respiration.  
 3. Values are expressed as mean ± S.E.  
 a, b, c Means within a column followed by different letters differ significantly (p < 0.05).

Table V.2 Rate of  $^{14}\text{C}$ -phenylalanine incorporation into protein of calf muscle preparations.

Group	Rate of $^{14}\text{C}$ -phenylalanine incorporation (nmole/mg/h) <sup>1</sup>	Ks <sup>2</sup> (%/d)	% O <sub>2</sub> consumption expended by protein synthesis <sup>3</sup>
10-21d dairy	0.023 ± 0.002a	1.5	3.3
7 month dairy	0.015 ± 0.002b	1.0	2.5
control DM	0.017 ± 0.002b	1.1	3.1
extreme DM	0.011 ± 0.001c	0.7	2.0

<sup>1</sup> Values expressed as mean ± S.E.

<sup>2</sup> Values for the fraction of muscle protein synthesized per day (Ks) were calculated assuming an average muscle amino acid molecular weight of 130 g/mole and a phenylalanine content of 3.0 mole % (Chang & Goldberg, 1978).

<sup>3</sup> The calculation of the fraction of muscle O<sub>2</sub> consumption expended for protein synthesis assumed 5 mole ATP to be required for incorporation of 1 mole of amino acid into a peptide chain and 1 mole O<sub>2</sub> to be required for synthesis of 5 mole ATP.

a, b, c Means within a column followed by different letters differ significantly (P<0.05).

Table V.3 Rate of tyrosine release from calf muscle preparations.

Group	nmole tyrosine/mg/h'
10-21d dairy	0.002 ± 0.009a
7 month dairy	0.002 ± 0.011a
control DM	0.016 ± 0.008b
extreme DM	0.078 ± 0.008c

' Values expressed as mean ± S.E.  
a,b,c Means within a column followed by different letters  
differ significantly (P<0.05).

VI. O<sub>2</sub> CONSUMPTION AND Na<sup>+</sup>, K<sup>+</sup>-ATPase-DEPENDENT RESPIRATION  
IN MUSCLE OF LAMBS AND LACTATING AND NON-LACTATING EWES

A. Abstract

The in vitro rate of O<sub>2</sub> consumption and the portion of respiration inhibited by ouabain were determined for sternomandibularis preparations from lambs, their lactating dams, and non-pregnant, non-lactating (control) ewes. Measurements were repeated after a 5 week interval at which time lambs had been weaned for 2 weeks and their dams were no longer lactating (dry).

Muscle from lambs at 2 weeks of age had an O<sub>2</sub> consumption rate 25% greater ( $p < 0.001$ ) than at 7 weeks of age and 49% greater ( $p < 0.001$ ) than that of control ewes; at 7 weeks of age lamb muscle O<sub>2</sub> consumption was 21% greater ( $p < 0.001$ ) than that of control ewes. Muscle preparations from lactating ewes had an O<sub>2</sub> consumption rate 35% greater ( $p < 0.001$ ) than dry and control ewes. The O<sub>2</sub> consumption rate of muscle preparations from control ewes did not differ significantly between sampling periods.

$10^{-5}$ M ouabain inhibited muscle O<sub>2</sub> consumption by 39%. Increased energy expenditure at the level of Na<sup>+</sup>,K<sup>+</sup>-ATPase accounted for 40% of the increased O<sub>2</sub> consumption rate of muscle from lambs of 2 weeks as contrasted to 7 weeks of age, and 60% of the increased O<sub>2</sub> consumption of muscle due to lactation.

## B. Introduction

To understand the factors which affect maintenance energy requirements at the metabolic level, it is necessary to identify specific components of background or maintenance energy expenditure and to quantify their relative contribution under a variety of different physiological conditions.

The activity of  $\text{Na}^+, \text{K}^+$ -ATPase in counteracting the transmembrane movement of  $\text{Na}^+$  and  $\text{K}^+$  along their concentration gradients has been identified to be a major source of energy expenditure in the muscle of sheep and calves (Chapters IV & V). The extent to which physiological, genetic and environmental factors influence  $\text{Na}^+, \text{K}^+$ -ATPase activity has not been established, but up to 80% of the increased *in vitro*  $\text{O}_2$  consumption for muscle from cold exposed sheep is due to increased energy expenditure at the level of the  $\text{Na}^+, \text{K}^+$ -ATPase (Chapter IV).

This study was undertaken to examine the effect of age and lactation on total *in vitro* muscle  $\text{O}_2$  utilization and the proportion of respiration used in support of the activity of  $\text{Na}^+, \text{K}^+$ -ATPase.

## C. Experimental

Muscle samples were obtained from four 2 week old nursing male lambs; their four lactating dams, 2-4 years of age; and four non-pregnant, non-lactating (control) ewes, 2-5 years of age. These sheep were all sampled 5 weeks later



after the lambs had been weaned for 2 weeks and their dams were no longer lactating. Animals were brought in from pasture 3-5 d prior to surgery and housed in a heated barn (approximately 20°C) where they were fed good quality grass hay free choice. Water was available at all times. Lactating and control ewes were fasted and nursing lambs separated from their dams overnight prior to surgery.

#### Measurement of respiration and $\text{Na}^+$ , $\text{K}^+$ -ATPase-dependent respiration

A section of the sternomandibular muscle was taken from each animal and preparation made following the method described in Chapter IV. Immediately upon removal from the animal the sections were placed in cooled (15°C) HEPES buffer containing 10.0 mM glucose and 5.0 mM acetate as substrates. Two to three small tied muscle fiber bundles were prepared (20.0 x 0.5 mm) at room temperature with the aid of a dissecting microscope. Initial respiration rates were measured for each muscle fiber bundle in an  $\text{O}_2$  electrode system (Chapter IV). The muscle preparations were incubated for 45 min in buffer containing  $10^{-5}\text{M}$  ouabain. Respiration rates were again measured. Percent inhibition of respiration by ouabain was calculated using the ratio of the difference between initial and that after incubation to the initial respiration rate.

### Statistical analysis

Rate of muscle  $O_2$  consumption, portion of respiration inhibited by ouabain, and  $Na^+, K^+$ -ATPase-dependent and independent respiration were analysed using a one-way analysis of variance with groups as fixed effects.

### D. Results

In the first period, muscle from lambs at 2 weeks of age had an  $O_2$  consumption rate greater ( $p < 0.001$ ) than that of lactating and control ewes (Table VI.1). The rate of  $O_2$  consumption for muscle from lactating ewes was 32% greater ( $p < 0.001$ ) than that of control ewes. Inhibition of respiration by ouabain ranged from 35.8 to 45.8% and was greatest ( $p < 0.001$ ) for muscle from lactating ewes.

In the second period, muscle from lambs at 7 weeks of age had an  $O_2$  consumption rate greater ( $p < 0.001$ ) than that of dry and control ewes (Table VI.2). The  $O_2$  consumption rates of muscle from dry and control ewes did not differ significantly. Inhibition of respiration by ouabain ranged from 35.2 to 41.7% and was lowest ( $p < 0.001$ ) for muscle from lambs at 7 weeks of age.

$Na^+, K^+$ -ATPase-dependent respiration, the amount of  $O_2$  calculated to have been expended by the muscle through  $Na^+, K^+$ -ATPase activity, was greater ( $p < 0.001$ ) for lambs at 2 weeks of age than lactating and control ewes (Table VI.2).  $Na^+, K^+$ -ATPase-dependent respiration for muscle from lactating ewes was 67% greater ( $p < 0.001$ ) than that of

control ewes.  $\text{Na}^+, \text{K}^+$ -ATPase-independent respiration, the residual portion of muscle  $\text{O}_2$  consumption not accounted for by activity of the  $\text{Na}^+, \text{K}^+$ -ATPase, was greater ( $p < 0.001$ ) for muscle from lambs at 2 weeks of age than that of lactating and control ewes.  $\text{Na}^+, \text{K}^+$ -ATPase-independent respiration for muscle from lactating ewes was 13% greater ( $p < 0.005$ ) than that of control ewes.

$\text{Na}^+, \text{K}^+$ -ATPase-dependent respiration did not differ significantly between lambs at 7 weeks of age and dry and control ewes.  $\text{Na}^+, \text{K}^+$ -ATPase-independent respiration was greater ( $p < 0.001$ ) for muscle from lambs at 7 weeks of age than from dry and control ewes. Muscle from dry and control ewes had similar values for  $\text{Na}^+, \text{K}^+$ -ATPase-independent respiration.

#### Comparisons of measurements during lactation and after weaning

Muscle from lambs at 2 weeks had an  $\text{O}_2$  consumption rate 25% greater ( $p < 0.001$ ) than at 7 weeks of age. The proportion of respiration inhibited by ouabain was similar for lambs at both ages. Muscle from lambs at 2 weeks of age had a 28% greater ( $p < 0.001$ ) value for  $\text{Na}^+, \text{K}^+$ -ATPase-dependent respiration than at 7 weeks of age, and a 23% greater  $\text{Na}^+, \text{K}^+$ -ATPase-independent respiration than at 7 weeks of age. Muscle from one of the lambs differed from samples from the other lambs with a greater ( $p < 0.05$ ) proportion of respiration inhibited by ouabain at 2 and 7 weeks of age,

and a greater ( $p < 0.05$ ) muscle  $O_2$  consumption rate at 7 weeks of age.

The  $O_2$  consumption rate of muscle from lactating ewes was 35% greater ( $p < 0.001$ ) than when the ewes were dry. The proportion of respiration inhibited by ouabain was also greater ( $p < 0.05$ ) for muscle taken from ewes when lactating than when dry. Muscle from lactating ewes had  $Na^+, K^+$ -ATPase-dependent and independent respiration values 50% ( $p < 0.001$ ) and 24% greater ( $p < 0.001$ ), respectively, than when dry. Muscle  $O_2$  consumption and the proportion of respiration inhibited by ouabain did not differ significantly between animals within groups in sampling periods.

The  $O_2$  consumption rate of muscle from control ewes remained constant between sampling periods. Inhibition of muscle respiration by ouabain increased ( $p < 0.05$ ) from 36.8 to 39.8% for the second sampling period. As a result,  $Na^+, K^+$ -ATPase-dependent and independent respiration differed by 8% ( $p < 0.05$ ) for muscle from control ewes between sampling periods. Muscle  $O_2$  consumption and the proportion of respiration inhibited by ouabain did not differ between animals within groups in sampling periods.

## E. Discussion

Skeletal muscle is an important site of energy expenditure, estimated to account for 26-62% of the whole body  $O_2$  consumption of dogs and small mammals (Martin & Fuhrman, 1955), and from measurements of cardiac output to body tissues (Christopherson et al. 1980), can be calculated to account for 37% of total cardiac output. Both the age related decrease in energy metabolism of sheep (Graham, 1967) and the increased maintenance energy requirements associated with lactation (Moe et al. 1972) were reflected in the in vitro  $O_2$  consumption rates of muscle from lambs and ewes. The decrease in muscle  $O_2$  consumption measured for lambs at 7 weeks as compared to 2 weeks of age, was the result of approximately equal proportional decreases in both the  $Na^+, K^+$ -ATPase-dependent and independent components of respiration. The effects of weaning and aging are not separated in the comparison of 7 and 2 week old lambs. The lower muscle  $O_2$  consumption of dry and control ewes than of 7 week lambs was due to a decrease in  $Na^+, K^+$ -ATPase-independent rather than dependent respiration suggesting that either aging per se, or aging after 7 weeks influences muscle energy expenditure differently than aging and weaning.

Within 2 weeks of weaning, the  $O_2$  consumption rate, portion of respiration inhibited by ouabain, and  $Na^+, K^+$ -ATPase-dependent and independent respiration of muscle from the ewes that had been lactating had decreased

to the values measured for muscle from control ewes. Lactating animals are known to have a greater feed intake than when dry; greater feed intake has been measured to cause increased  $O_2$  consumption as a result of a greater amount of  $Na^+, K^+$ -ATPase-independent respiration in muscle from sheep (Chapter IV). Since they were allowed voluntary intake, it is likely that lactating ewes had a greater feed intake than dry and control ewes and this may have contributed to the greater muscle  $O_2$  consumption of lactating than dry and control ewes. Nonetheless, of the increased muscle  $O_2$  consumption measured for ewes when lactating, 60% was accounted for by increased energy expenditure at the level of  $Na^+, K^+$ -ATPase. Of relevance is the observation that cold exposure selectively increases  $Na^+, K^+$ -ATPase activity in the muscle of sheep (Chapter IV) and small mammals (Stevens & Guernsey, 1974; Guernsey & Stevens, 1977). The mechanism for cold-induced enhancement of enzyme activity may be at least partially achieved by increased enzyme synthesis as a result of enhanced thyroid activity (Chapter IV). Clearly, the physiological objectives of lactation are different from cold exposure, the former being to achieve the synthesis of milk and the latter to counteract the heat demand of the environment. However, as in cold exposure, it is conceivable that the very considerable endocrine changes (Hart et al. 1978) that occur during lactation may cause the enhancement of  $Na^+, K^+$ -ATPase activity.

Ouabain inhibited muscle  $O_2$  consumption by an average of 89%, in agreement with values previously reported for muscle of sheep and cattle (Chapters IV & V). The magnitude of inhibition of respiration by ouabain is a function of the extent to which  $Na^+K^+$  transport does in fact account for  $O_2$  consumption and also both ouabain accessibility to the enzyme and the equilibrium reached between formation and dissociation of the inhibitor-enzyme complex (Tobin & Brody, 1972). The exact proportion of  $Na^+K^+$ -ATPase activity actually inhibited in our muscle preparations under the experimental conditions employed is not known. Therefore, our measurements of the proportion of muscle  $O_2$  consumption required for support of  $Na^+K^+$ -ATPase activity must be considered as minimum estimates. Underestimation of  $Na^+K^+$ -ATPase activity resulting from incomplete inhibition of the enzyme would overestimate  $Na^+K^+$ -ATPase-independent respiration. The metabolic processes which would account for the  $Na^+K^+$ -ATPase-independent component of muscle  $O_2$  consumption have not been identified. From estimates of the cost of protein synthesis in the energy metabolism of muscle from growing calves (Chapter V), the energy expended for peptide bond synthesis would appear to account for only a minor portion of  $Na^+K^+$ -ATPase-independent respiration.

In conclusion, lactation was shown to increase  $O_2$  consumption in muscle from ewes and to cause increased energy expenditure at the level of the  $Na^+K^+$ -ATPase. The mechanism for increased  $Na^+K^+$ -ATPase activity is not

Aging to 7 weeks plus weaning decreased muscle  $O_2$  consumption and  $Na^+, K^+$ -ATPase activity in lambs. The amount of muscle  $O_2$  consumption not explained by  $Na^+, K^+$ -ATPase activity is considerable and may be due to incomplete enzyme inhibition under the experimental conditions, or as yet unidentified energy consuming metabolic processes.

#### F. Acknowledgements

Grateful acknowledgement is extended to the Natural Sciences and Engineering Research Council Canada for support in part of this research.

#### G. References

- Christopherson, R. J., Gregory, N., Lister, D., & Wotton, S. (1980) *Can. J. Anim. Sci.* 60, 1058.
- Graham, N. McC. (1967) *Australian J. Agr. Res.* 18, 127.
- Guernsey, D. L. & Stevens, E. D. (1977) *Science* 196, 908.
- Hart, I. C., Bines, J. A., Morant, S. V. & Ridley, J. L. (1978) *J. Endocr.* 77, 333.
- Martin, A. W. & Fuhrman, F. A. (1955) *Physiol. Zool.* 28, 18.
- Moe, P. W., Flatt, W. P. & Tyrrell, H. F. (1972) *J. Dairy Sci.* 55, 945.
- Stevens, E. D. & Kido, M. (1974) *Comp. Biochem. Physiol.* 47A, 395.



Tobin, T. & Brody, T. M. (1972) *Biochem. Pharmac.* 21, 1553.

Table VI.1 O<sub>2</sub> consumption and proportion of respiration inhibited by ouabain for muscle from lambs and lactating and non-lactating ewes.

Group	Muscle O <sub>2</sub> consumption ( $\mu$ l O <sub>2</sub> /mg/h)	% Inhibition of respiration by ouabain
Period I		
2 week lambs	5.37 ± 0.33a	35.8 ± 2.3a
lactating ewes	4.75 ± 0.25b	45.8 ± 4.8b
control ewes	3.61 ± 0.20c	36.8 ± 3.2a
Period II		
7 week lambs	4.31 ± 0.29a	35.2 ± 3.0a
dry ewes	3.53 ± 0.20b	41.7 ± 3.9b
control ewes	3.56 ± 0.21b	39.8 ± 1.9b

Values expressed as mean ± S.E.  
 a,b,c Means within a column and within a measurement period followed with different letters differ significantly (p<0.05).

Table VI.2 Na<sup>+</sup>,K<sup>+</sup>-ATPase-dependent and independent respiration for muscle from lambs and lactating and non-lactating ewes.

Group	Na <sup>+</sup> ,K <sup>+</sup> -ATPase-dependent respiration (ul O <sub>2</sub> /mg/h) <sup>1</sup>	Na <sup>+</sup> ,K <sup>+</sup> -ATPase-independent respiration (ul O <sub>2</sub> /mg/h) <sup>2</sup>
Period I		
2 week lambs	1.94 ± 0.19a	3.43 ± 0.22a
lactating ewes	2.19 ± 0.16b	2.56 ± 0.13b
control ewes	1.31 ± 0.12c	2.30 ± 0.21c
Period II		
7 week lambs	1.51 ± 0.16a	2.80 ± 0.23a
dry ewes	1.46 ± 0.15a	2.07 ± 0.19b
control ewes	1.43 ± 0.12a	2.13 ± 0.13b

<sup>1</sup> Na<sup>+</sup>,K<sup>+</sup>-ATPase-dependent respiration = total O<sub>2</sub> consumption x inhibition by ouabain.  
<sup>2</sup> Na<sup>+</sup>,K<sup>+</sup>-ATPase-independent respiration = total O<sub>2</sub> consumption - Na<sup>+</sup>,K<sup>+</sup>-ATPase-dependent respiration.  
<sup>3</sup> Values expressed as mean ± S.E.  
**a,b,c** Means within a column followed by different letters differ significantly (p<0.05).

### General Discussion and Conclusions

This study confirmed previous findings in work conducted with sliced tissue preparations that active transport of  $\text{Na}^+$  and  $\text{K}^+$ , catalyzed by the plasma membrane  $\text{Na}^+, \text{K}^+$ -ATPase, is a major component of muscle energy expenditure. While sliced muscle exhibited a significant decrease in  $\text{O}_2$  consumption as compared to intact muscle, inhibition of respiration by ouabain was not substantially altered. The low response to ouabain of  $\text{O}_2$  consumption and heat production obtained in investigations conducted with intact organ preparations was shown to be due, at least in part, to prior inhibition of  $\text{Na}^+, \text{K}^+$ -ATPase by use of a buffer with a high content of  $\text{MgCl}_2$ .  $\text{Na}^+, \text{K}^+$ -ATPase activity accounted for 13-33% of mouse, and 29-46% of sheep and cattle skeletal muscle  $\text{O}_2$  consumption. The lower ouabain inhibition of respiration for mouse muscle than ruminant muscle may be a function of the enzyme-inhibitor interaction rather than a real difference in the actual proportion of muscle  $\text{O}_2$  consumption expended in support of active  $\text{Na}^+-\text{K}^+$  transport. Species differences in enzyme susceptibility to inhibition by ouabain have been shown to exist.  $\text{Na}^+, \text{K}^+$ -ATPase from rats has been measured to require a greater concentration of ouabain for maximal inhibition than that from dogs, pigs, sheep and guinea pigs. The actual proportion of the total  $\text{Na}^+-\text{K}^+$  transport inhibited by ouabain in the muscle preparations employed in this study is

In vitro estimates of protein synthesis for calf muscle preparations were in good agreement with measurements obtained in vivo by other workers. While the rate of protein synthesis was affected by age and breed, the amount of energy expended in support of peptide bond synthesis was a minor component of total energy expenditure by the sternomandibularis muscle.

Tyrosine release from muscle has been used by investigators as a measure of protein degradation. Significantly greater rates of tyrosine release were measured for muscle from control and extreme DM calves than for dairy calves, which may reflect differences in protein degradative capacity.

not known, but may not have been complete if the enzyme was not fully accessible to ouabain.

The amount of energy expended for  $\text{Na}^+, \text{K}^+$ -ATPase activity was responsive to physiological and environmental factors. Long term cold exposure was found to result in specific stimulation of  $\text{Na}^+, \text{K}^+$ -ATPase activity in muscle from sheep. More rapid cycling of  $\text{Na}^+$  and  $\text{K}^+$  between the extra- and intra-cellular space would be a futile cycle and yield heat production. Cold exposure of mice resulted in an increased muscle  $\text{O}_2$  consumption not due to specific enhancement of  $\text{Na}^+, \text{K}^+$ -ATPase activity. This may be a result of resistance of mouse  $\text{Na}^+, \text{K}^+$ -ATPase to inhibition by ouabain, or due to the presence of an alternate mechanism for cold-induced muscle thermogenesis. The mechanism of enhancement of  $\text{Na}^+, \text{K}^+$ -ATPase activity is not known, but was speculated to be due to either an increased amount of enzyme as a result of hormonal stimulation of protein synthesis, a changed enzyme efficiency, or a changed membrane permeability to  $\text{Na}^+$  and  $\text{K}^+$  resulting in increased substrate presentation to the enzyme.

Lactation was measured to result in greater muscle  $\text{O}_2$  consumption compared to non-lactating ewes, due to a specific enhancement of  $\text{Na}^+, \text{K}^+$ -ATPase activity. Young lambs and calves had a greater rate of muscle  $\text{O}_2$  consumption as contrasted to older animals. This was due to increases in both  $\text{Na}^+, \text{K}^+$ -ATPase-dependent and independent components of respiration.