

Interventions to prevent hypothermia at birth in preterm and/or low birthweight infants (Review)

McCall EM, Alderdice F, Halliday HL, Jenkins JG, Vohra S



**THE COCHRANE
COLLABORATION®**

This is a reprint of a Cochrane review, prepared and maintained by The Cochrane Collaboration and published in *The Cochrane Library* 2010, Issue 3

<http://www.thecochranelibrary.com>



Interventions to prevent hypothermia at birth in preterm and/or low birthweight infants (Review)
Copyright © 2010 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

TABLE OF CONTENTS

HEADER	1
ABSTRACT	1
PLAIN LANGUAGE SUMMARY	2
BACKGROUND	2
OBJECTIVES	4
METHODS	4
RESULTS	8
DISCUSSION	17
AUTHORS' CONCLUSIONS	19
ACKNOWLEDGEMENTS	19
REFERENCES	20
CHARACTERISTICS OF STUDIES	27
DATA AND ANALYSES	49
Analysis 1.1. Comparison 1 Plastic wrap versus routine care, Outcome 1 Core body temperature (°C) on admission to NICU or up to 2 hours after birth.	53
Analysis 1.2. Comparison 1 Plastic wrap versus routine care, Outcome 2 Core body temperature (°C) 1 hour after the initial admission temperature to the NICU was taken.	54
Analysis 1.3. Comparison 1 Plastic wrap versus routine care, Outcome 3 Hypothermia on admission to NICU: core body temperature < 36.5°C or skin temperature < 36°C.	54
Analysis 1.4. Comparison 1 Plastic wrap versus routine care, Outcome 4 Death within hospital stay.	55
Analysis 1.5. Comparison 1 Plastic wrap versus routine care, Outcome 5 Major brain injury.	56
Analysis 1.6. Comparison 1 Plastic wrap versus routine care, Outcome 6 Duration of oxygen therapy (days).	56
Analysis 1.7. Comparison 1 Plastic wrap versus routine care, Outcome 7 Duration of hospitalisation (days).	57
Analysis 1.9. Comparison 1 Plastic wrap versus routine care, Outcome 9 Apgar at 1 minute.	58
Analysis 1.11. Comparison 1 Plastic wrap versus routine care, Outcome 11 Apgar at 5 minutes.	59
Analysis 1.12. Comparison 1 Plastic wrap versus routine care, Outcome 12 First blood gas pH.	59
Analysis 1.13. Comparison 1 Plastic wrap versus routine care, Outcome 13 Bicarbonate (mmol/L).	60
Analysis 1.14. Comparison 1 Plastic wrap versus routine care, Outcome 14 First serum glucose concentration (mmol/L).	60
Analysis 1.15. Comparison 1 Plastic wrap versus routine care, Outcome 15 Intubation in delivery room.	61
Analysis 1.16. Comparison 1 Plastic wrap versus routine care, Outcome 16 Hyperthermia on admission to NICU: core body temperature > 37.5°C.	61
Analysis 2.1. Comparison 2 Plastic cap versus routine care, Outcome 1 Core body temperature (°C) on admission to NICU or up to 2 hours after birth.	62
Analysis 2.2. Comparison 2 Plastic cap versus routine care, Outcome 2 Core body temperature (°C) 1 hour after the initial admission temperature to the NICU was taken.	62
Analysis 2.3. Comparison 2 Plastic cap versus routine care, Outcome 3 Hypothermia on admission to NICU: core body temperature < 36.4 °C.	63
Analysis 2.4. Comparison 2 Plastic cap versus routine care, Outcome 4 Death within hospital stay.	63
Analysis 2.5. Comparison 2 Plastic cap versus routine care, Outcome 5 Major brain injury.	64
Analysis 2.6. Comparison 2 Plastic cap versus routine care, Outcome 6 Apgar score at 1 minute.	64
Analysis 2.7. Comparison 2 Plastic cap versus routine care, Outcome 7 Apgar score at 5 minutes.	65
Analysis 2.8. Comparison 2 Plastic cap versus routine care, Outcome 8 First blood gas pH.	65
Analysis 2.9. Comparison 2 Plastic cap versus routine care, Outcome 9 Bicarbonate (mmol/L).	66
Analysis 2.10. Comparison 2 Plastic cap versus routine care, Outcome 10 First serum glucose concentration (mmol/L).	66
Analysis 2.11. Comparison 2 Plastic cap versus routine care, Outcome 11 Intubation in delivery room.	67
Analysis 3.1. Comparison 3 Stockinet cap versus routine care, Outcome 1 Core body temperature (°C) on admission to NICU or up to 2 hours after birth.	67
Analysis 3.2. Comparison 3 Stockinet cap versus routine care, Outcome 2 Hypothermia on admission to NICU: core body temperature < 36.5°C or skin temperature < 36°C.	68
Analysis 4.1. Comparison 4 Skin-to-skin care versus routine care, Outcome 1 Hypothermia: skin temperature < 35.5°C for 2 consecutive recordings.	68

Analysis 4.2. Comparison 4 Skin-to-skin care versus routine care, Outcome 2 Hypoglycaemia: blood glucose level < 2.6 mmol/L.	69
Analysis 5.1. Comparison 5 Transwarmer mattress versus routine care, Outcome 1 Core body temperature (°C) on admission to NICU or up to 2 hours after birth.	69
Analysis 5.2. Comparison 5 Transwarmer mattress versus routine care, Outcome 2 Hypothermia on admission to NICU: core body temperature < 36.5 °C or skin temperature < 36°C.	70
WHAT'S NEW	70
HISTORY	70
CONTRIBUTIONS OF AUTHORS	71
DECLARATIONS OF INTEREST	71
SOURCES OF SUPPORT	72
INDEX TERMS	72

[Intervention Review]

Interventions to prevent hypothermia at birth in preterm and/or low birthweight infants

Emma M McCall¹, Fiona Alderdice², Henry L Halliday³, John G Jenkins⁴, Sunita Vohra⁵

¹School of Medicine, Dentistry & Biomedical Sciences, Queen's University Belfast, Belfast, UK. ²Nursing and Midwifery Research Unit, Queen's University Belfast, Belfast, UK. ³Perinatal Room, Royal-Jubilee Maternity Service, Belfast, UK. ⁴Paediatrics, Antrim Area Hospital, Antrim, UK. ⁵CARE Program, Department of Pediatrics, University of Alberta, Edmonton, Canada

Contact address: Emma M McCall, School of Medicine, Dentistry & Biomedical Sciences, Queen's University Belfast, Microbiology Building, Grosvenor Road, Belfast, Northern Ireland, BT12 6BN, UK. e.mccall@qub.ac.uk.

Editorial group: Cochrane Neonatal Group.

Publication status and date: New search for studies and content updated (conclusions changed), published in Issue 3, 2010.

Review content assessed as up-to-date: 7 February 2010.

Citation: McCall EM, Alderdice F, Halliday HL, Jenkins JG, Vohra S. Interventions to prevent hypothermia at birth in preterm and/or low birthweight infants. *Cochrane Database of Systematic Reviews* 2010, Issue 3. Art. No.: CD004210. DOI: 10.1002/14651858.CD004210.pub4.

Copyright © 2010 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

ABSTRACT

Background

Keeping vulnerable preterm infants warm is problematic even when recommended routine thermal care guidelines are followed in the delivery suite.

Objectives

To assess efficacy and safety of interventions designed for prevention of hypothermia in preterm and/or low birthweight infants applied within 10 minutes after birth in the delivery suite compared with routine thermal care.

Search methods

We used the standard search strategy of the Cochrane Neonatal Review Group (CNRG). The review was updated in October 2009.

Selection criteria

Trials using randomised or quasi-randomised allocations to test a specific intervention designed to prevent hypothermia, (apart from 'routine' thermal care) applied within 10 minutes after birth in the delivery suite to infants of < 37 weeks' gestational age or birthweight ≤ 2500 g.

Data collection and analysis

We used the methods of the CNRG for data collection and analysis.

Main results

1) Barriers to heat loss [5 studies; plastic wrap or bag (3), plastic cap (1), stockinet cap (1)]:

Plastic wraps or bags were effective in reducing heat losses in infants < 28 weeks' gestation (4 studies, n = 223; WMD 0.68 °C; 95% CI 0.45, 0.91), but not in infants between 28 to 31 week's gestation. Plastic caps were effective in reducing heat losses in infants < 29 weeks' gestation (1 study, n = 64; MD 0.80 °C; 95% CI 0.41, 1.19). There was insufficient evidence to suggest that either plastic wraps

or plastic caps reduce the risk of death within hospital stay. There was no evidence of significant differences in other clinical outcomes for either the plastic wrap/bag or the plastic cap comparisons. Stockinet caps were not effective in reducing heat losses.

2) External heat sources [2 studies; skin-to-skin (1), transwarmer mattress (1)]:

Skin-to-skin care (SSC) was shown to be effective in reducing the risk of hypothermia when compared to conventional incubator care for infants (1 study, n = 31; RR 0.09; 95% CI 0.01, 0.64). The transwarmer mattress reduced the incidence of hypothermia on admission to NICU in VLBW infants (1 study, n = 24; RR 0.30; 95% CI 0.11, 0.83).

Authors' conclusions

Plastic wraps or bags, plastic caps, SSC and transwarmer mattresses all keep preterm infants warmer leading to higher temperatures on admission to neonatal units and less hypothermia. However, the small numbers of infants and studies and the absence of long-term follow-up mean that firm recommendations for clinical practice cannot be given.

PLAIN LANGUAGE SUMMARY

Interventions to prevent hypothermia at birth in preterm and/or low birthweight infants

Preventing low body temperature at birth in premature and low birthweight infants may be important to survival and long-term outcome. Babies rely on external help to maintain body and skin temperature particularly in the first 12 hours of life. For vulnerable infants born prematurely or that are very small, abnormally low body temperature (hypothermia) is a world-wide issue across all climates and can lead to a variety of diseases and even death. Preventative action is taken by reducing heat loss and/or providing warmth using external heat sources. Precautionary steps routinely include a warm delivery room; drying the newborn immediately, especially the head; wrapping in pre-warmed dry blankets that cover the head; pre-warming surfaces and eliminating draughts. A review of seven studies involving 391 infants used additional preventative actions in the first 10 minutes of life to prevent problems with hypothermia. Results showed that the use of special plastic wraps or bags, plastic caps, heated mattresses and skin-to-skin contact kept the infants warmer than routine preventative action. Limitations included the small numbers of infants and studies included; variations in the methods and definitions of normal body temperature, routine care; and the use of different materials. Although this review confirmed that some of these measures are effective in preventing hypothermia, we do not yet know the long-term consequences of these interventions therefore the authors recommend that further research is carried out.

BACKGROUND

Keeping preterm infants sufficiently warm immediately after birth, especially during resuscitation, is problematic even when routine thermal care guidelines are followed. The newborn cannot shiver (Scopes 1963), and relies on interventions to protect it against exposure to cold. The ability to maintain an equilibrium between heat loss and heat gain (Bickmann 1992) despite variation in environmental temperatures is restricted during the first 12 hours of life (Smales 1978). After birth, deep body and skin temperature of the term newborn can drop at a rate of approximately 0.1 °C and 0.3 °C per minute respectively unless immediate action is taken (Adamsons 1965a). Although cold stress may be important for initiating breathing (Harned 1970) and induced cooling may help protect the brain of asphyxiated term newborns, prolonged exposure to cold should be avoided.

Description of the condition

Extended periods of cold stress can lead to harmful side effects, which include hypoglycaemia (Elliott 1957), respiratory distress (Pomerance 1974), hypoxia, metabolic acidosis (Gandy 1964), coagulation defects (Chadd 1972), delayed readjustment from fetal to newborn circulation (Stephenson 1970), acute renal failure, necrotizing enterocolitis, failure to increase weight or weight loss (Glass 1968) and in extreme cases death (Elliott 1957). Nayeri 2006 concluded that hypothermia at birth is one of the most significant risk factors causing death in newborn infants of all birthweights and gestational ages and particularly for vulnerable preterm infants (Costeloe 2000; CESDI 2003). Factors that increase the risk of hypothermia include prematurity, intrauterine growth retardation (Borse 1997; Hey 1975), asphyxia, certain congenital anomalies such as gastroschisis and damage to the central

nervous system (Bickmann 1992).

Rapid postnatal fall in body temperature is attributable to a combination of the physical characteristics of the infant (e.g. large surface area in relation to body weight and a thin layer of insulating fat) and environmental factors in the delivery suite. Extent of total heat loss and the four modes of heat exchange (conduction, convection, radiation and evaporation) are influenced by the ambient air temperature, pressure and relative humidity, and temperature of surrounding surfaces (Capobianco 1980; Thomas 1994). Increased rate of heat loss is mainly caused by evaporation of amniotic fluid from the skin surface (Hammarlund 1980) when the wet newborn moves from the warm environment of the uterus (Adamsons 1965a) into a cool, dry delivery suite.

In an attempt to maintain core body temperature within the normal range of 36.5 to 37.5 °C (skin temperature of 0.5 to 1.0 °C lower) (Hey 1970 ; Oliver 1965), the term infant responds mainly by production of heat from the breakdown of brown fat (Davis 1980) (non-shivering thermogenesis) (Stern 1970) and peripheral vasoconstriction. When skin temperature falls to 35 to 36 °C, non-shivering thermogenesis is initiated (Bruck 1961). The World Health Organisation classifies a core body temperature for newborns of 36 to 36.4 °C as mild hypothermia, 32 to 35.9 °C as moderate and < 32 °C as severe (WHO 1997). The preterm infant has the combined disadvantages of decreased fat for heat production and insulation, decreased glycogen stores, immature skin which increases water loss and poor vascular control. They experience even higher evaporative heat losses than term infants in the first day, especially at low ambient relative humidities (Hammarlund 1979). Five hundred and sixty calories of heat are lost for each millilitre of water which evaporates from the skin (Rutter 2000). Currently, there is no accepted formal definition of 'normal' temperatures for preterm infants and methods and accuracy of temperature measurement continue to be debated (Bailey 2000; Smith 2004).

The external (skin-environment) temperature gradient is pivotal in influencing the infant's response to cold (Adamsons 1965) and it is here that health professionals can intervene in the delivery suite to minimise the risk of hypothermia.

Standard care includes providing a warm delivery room at a minimum of 25 °C (although rarely achieved in practice) (WHO 1997), drying the infant thoroughly, immediately after birth (especially the head) (Bloom 1994), removing any wet blankets, wrapping in a prewarmed blanket, prewarming any contact surfaces, eliminating draughts and close proximity to outside walls (Capobianco 1980). If available, radiant warmers for resuscitation and stabilisation allow easy access and are effective in preventing heat losses, provided that the infant is immediately dried and placed under the prewarmed heater (Du 1969; Dahm 1972). Although the infant gains heat by radiation, there are increased potential losses through convection and evaporation and these losses are exacerbated if the infant is inadequately dried. Servocontrol is advantageous for the avoidance of overheating or underheating

if absorption of heat is being obstructed by coverings. Watkinson 2006 suggested two reasons why conventional thermal care has failed to prevent hypothermia at birth: 1) current recommended techniques are inadequate and 2) thermal management awareness needs to be increased among personnel carrying out the resuscitation of newborns. The latter has also been highlighted in reviews of practice for care of very immature infants during resuscitation and transfer (Lyon 2006).

Description of the intervention

Studies to investigate the effectiveness of additional measures to reduce heat loss in the immediate postnatal period fall into two groups:

- (1) barriers to heat loss;
- (2) external heat sources.

Interventions in the first group focus mainly on reducing evaporative heat losses (LeBlanc 1991) and have included wraps and/or headcoverings made from a variety of materials (Chaput 1979; Coles 1979; Holzman 1985 ; Lang 2004). Baum 1968 tested a polyester suit lined with aluminum, known as the 'silver swaddler', designed to prevent hypothermia by reducing all modes of heat transfer to the environment. This was effective for infants with birthweights > 3000 g but, since the material is opaque, it is not practical during resuscitation. Transparent plastic coverings such as bubble wrap (Besch 1971) and single layer gowns (Hobbs 1975) are effective in the delivery suite for full-term healthy newborn infants and those with birthweights > 2000 g respectively and where the intervention was polyethylene wrap for infants of < 33 weeks' gestation (Lenclen 2002). Hoods or heat shields that are not in contact with the infant's body have also been used in conjunction with a radiant warmer or incubator (Baumgart 1981; Bell 1980). Barrier creams, waxes or protective films such as Aquaphor® (Nopper 1996) have also been utilised to reduce heat losses in immature infants but are not normally applied within 10 minutes of birth.

Interventions in the second group have included heated mattresses (Almeida 2009) and as an alternative to radiant warmers, for a healthy term newborn skin-to-skin contact, (where the infant is thoroughly dried and placed on the mother's chest and abdomen with a light blanket around them), can reduce radiant and conductive heat loss and promote temperature stabilisation (Britton 1980; Christensson 1992). More recently, efforts to reduce the incidence of hypothermia on admission to neonatal units have included quality improvement initiatives using 'intervention bundles' such as staff education, consistent room air temperature, use of polyethylene bags and transfer in a warmed incubator (Kaplan 2009 ; Ho 2007).

All of these interventions have potential disadvantages; for example, Newton 2003 reported that significantly more infants (with gestational ages < 30 completed weeks) wrapped in polythene bags were hyperthermic (> 37 °C) when compared to unwrapped his-

torical controls. Brun 1997 noted that a chemical hot pack during resuscitation of a newborn infant resulted in third degree burns and recommended that these should not be used unless the peak temperature of the pack is < 44 °C.

How the intervention might work

Interventions should either decrease total heat losses or provide external heat without compromising accessibility during resuscitation and should have minimal side effects (such as hyperthermia, burns, maceration, or infection).

Why it is important to do this review

Neonatal hypothermia after birth is a world-wide issue (Costeloe 2000) across all climates (Christensson 1988; Johanson 1992; Tafari 1973; Lupton 2007; Kumar 2009) and, if prolonged, can lead to harm and in severe cases death. Silverman 1958 and Day 1964 showed that reducing heat losses in preterm infants in the first few days after birth increased survival rates. Knobel 2005 reported that chart review of 100 patients revealed that 93% of infants of < 1000 g birthweight had rectal temperature < 36.4°C on admission to neonatal intensive care. The Project 27/28 undertaken in England, Wales and Northern Ireland for a period of two years including all live born infants (n = 3522) with a gestation between 27⁰ and 28⁶ weeks (CESDI 2003) stated that some component of early thermal care was inadequate in over a third of infants and this was associated with the infants who died. Early intervention in the delivery suite, particularly for the preterm infant undergoing resuscitation (Lupton 2008) is therefore of high priority if hypothermia is to be prevented. Soll 2008 re-emphasised the need to address and understand the consequences of poor thermal care for the newborn infant in order to improve clinical outcomes. This review focuses on individual interventions applied within 10 minutes after birth in the delivery suite and is limited to preterm and/or low birthweight infants since these are most susceptible to the adverse effects of hypothermia. Longer term thermal management and spatial or environmental strategies for increasing warming are beyond the scope of this review. This is an update of a Cochrane review first published in 2005 and previously updated in 2008.

OBJECTIVES

PRIMARY OBJECTIVE

To assess the efficacy and safety of interventions, designed for the prevention of hypothermia in preterm and/or low birthweight infants, and applied within 10 minutes after birth in the delivery suite, compared with routine thermal care.

Subgroup analyses to be carried out by intervention, by birthweight/gestational age and by developmental status of the country of study, to determine whether effectiveness varies according to:

The interventions applied:

Birthweight and gestational age within the following categories:

- birthweight (< 1500 g), (1500 g to 2500 g)

- gestational age (< 28 wk), (28 to 32 wk) and (33 to 37 wk)

The developmental status of the country in which the trial was conducted based upon the UNICEF classification of a developing country (UNICEF 2002).

However, the subgroups reported in the included studies were not compatible with those pre-specified. Therefore, post facto subgroup analyses based on reported gestational age and birthweight subcategories were carried out where appropriate within each comparison group.

SECONDARY OBJECTIVES

To assess effects on complications associated with preterm birth, hypothermia, and adverse outcomes.

METHODS

Criteria for considering studies for this review

Types of studies

All trials using randomised or quasi-randomised allocation to test a specific intervention designed to prevent hypothermia immediately after birth.

Types of participants

Preterm infants of < 37 weeks' gestational age (according to best obstetric estimate at time of delivery) or low birthweight infants of ≤ 2500 g, where the intervention to prevent hypothermia is applied within 10 minutes after birth in the delivery suite. Both appropriate and small-for-gestational age infants were eligible.

Exclusions: Infants with major congenital malformations, especially abdominal wall defects.

Types of interventions

Any intervention applied within 10 minutes after birth in the delivery suite apart from ROUTINE THERMAL CARE, which was defined as any of the following routine practices: providing a warm delivery suite at a minimum of 25 °C (rarely achieved in practice), drying the infant immediately after birth, removing

any wet blankets and wrapping in a pre-warmed blanket, pre-warming any contact surfaces, avoiding draughts and, in developed countries, the use of radiant warmers or incubators. The control intervention comprised any elements of routine thermal care.

The interventions studied were to include:

(1) Barriers to heat loss applied to any part of the body of the preterm and/or low birthweight infant within 10 minutes after birth in the delivery suite

- Coverings such as transparent plastic wraps and bags made of low density polyethylene (LDPE) or linear low density polyethylene (LLDPE) or polyvinylidene chloride (PVDC)
- Semi-permeable membranes such as Opsite® or Tegaderm®
- Other additional swaddling materials or wraps (excluding delivery room blankets) such as the 'silver swaddler'.

(2) External heat sources (non-routine) initiated within 10 minutes after birth in the delivery suite

- Skin-to-skin care
- Heated/gel/chemical mattresses.

Types of outcome measures

Primary outcomes

The temperature of the infant taken on admission to the Neonatal Intensive Care Unit (NICU) or up to two hours after birth. Temperature was assessed as both continuous and dichotomous variables.

Rectal, axillary, oral or tympanic temperature measurements were accepted as equivalent core body temperature and abdominal skin temperature was accepted for skin temperature. Where both core temperature and skin temperature were recorded core temperature took priority. Where multiple temperatures were recorded (i.e. within different time frames up to two hours after birth) the lowest temperature recorded took priority.

A core body temperature of < 36.5 °C or a skin temperature of < 36 °C indicated the presence of hypothermia within the control and intervention groups.

For hypothermia, core body temperature and skin temperature subgroupings as defined by WHO 1997 were used to determine three levels of severity:

- mild hypothermia or cold stress: core body temperature 36 to 36.4 °C or skin temperature of 35.5 to 35.9 °C;
- moderate hypothermia: core body temperature 32 to 35.9 °C or skin temperature of 31.5 to 35.4 °C;
- severe hypothermia: core body temperature of < 32 °C or skin temperature < 31.5 °C.

Secondary outcomes

These were categorised as (1) morbidity and (2) adverse outcomes due to the intervention.

(1) Morbidity

- hypoglycaemia (defined by a blood glucose level of < 2.0 mmol/L within two hours of birth);
- respiratory distress syndrome (RDS) (defined by clinical signs of grunting, flaring, retractions, cyanosis in room air, tachypnoea and a radiological picture of reticulogranular mottling and air bronchogram);
- surfactant given at any time;
- intubation in delivery room;
- requirement for ventilation and duration of ventilation (days);
- length of stay (days);
- mortality: death within seven days, death within 28 days and/or death during hospital stay;
- Severe metabolic acidosis as defined by pH < 7.20 and/or base deficit > 10 mmol/L within the first three days of life;
- intraventricular haemorrhage (defined according to the criteria of Papile et al from head ultrasound performed before 14 days of life) (Lee 2000; Papile 1978);
- patent ductus arteriosus (defined by clinical diagnosis plus treatment with indomethacin or surgical ligation or both) (Lee 2000);
- chronic lung disease (defined by oxygen dependency at 36 weeks' postmenstrual age for an infant who was born at ≤ 32 weeks' gestation) (Lee 2000; Shennan 1988);
- necrotizing enterocolitis [defined according to the criteria of Bell et al (Bell 1978) stage 2 or higher and classified as medical (clinical symptoms and signs plus evidence of pneumatosis on abdominal radiographs) or surgical (histological evidence of NEC on surgical specimen of intestine)] (Lee 2000);
- acute renal failure [defined by a serum creatinine level of more than 1.5 mg/dL (Stapleton 1987) and oliguria (urine output < 1 ml/kg/hr)].

(2) Adverse outcomes due to the intervention

- hyperthermia (defined by an admission temperature to NICU or within two hours of birth of ≥ 38 °C);
- burns within three days of birth;
- maceration within three days of birth;
- skin or systemic infection secondary to intervention within the first week of birth (defined by a culture of pathogenic bacteria from normally sterile body tissue or fluid);
- antibiotic course of five days or more started within the first seven days of birth;
- interference with resuscitation and other practices (e.g. UV catheter placement for fluid replacement, chest tube insertion);
- fluid problems such as dehydration or fluid overload, electrolyte imbalance such as hypernatraemia (serum sodium > 150 mmol/L) or hyponatraemia (serum sodium < 130 mmol/L);
- any other unexplained adverse outcome attributed to the intervention within seven days of birth;

- negative psychological outcomes (perception of care by parents).

Search methods for identification of studies

The standard search strategy of the Cochrane Collaboration was used (Clarke 2002).

Electronic searches

We searched the following electronic databases: the Cochrane Central Register of Controlled Trials (CENTRAL, The Cochrane Library, Issue 4, 2009), the Database of Abstracts of Reviews of Effects (DARE 1994 to October 2009), MEDLINE (1950 to October 2009), CINAHL (1982 to October 2009), EMBASE (1974 to October 2009), conference/symposia proceedings using ZETOC (1993 to October 2009) and ISI proceedings (1990 to October 2009). In addition, OCLC WorldCat (October 2009) was searched and identified articles were cross referenced. No language restrictions were imposed. The MEDLINE search strategy is detailed below and similar search strategies were devised using appropriate terminology for each electronic database.

1. plastic barrier*.ab,ti.
2. polyethylene*.ab,ti.
3. (bubble wrap* or bubble-wrap* or bubblewrap*).ab,ti.
4. (clingfilm* or cling film* or cling-film*).ab,ti.
5. (plasticwrap* or plastic-wrap* or plastic wrap*).ab,ti.
6. exp Polyethylenes/
7. exp Polyvinyls/
8. (polyvinyl* or poly-vinyl*).ab,ti.
9. (low density polyethylene* or low-density polyethylene*).ab,ti.
10. (gladwrap* or glad wrap* or glad-wrap*).ab,ti.
11. (polybag* or poly bag* or poly-bag*).ab,ti.
12. (saranwrap* or saran-wrap* or saran wrap*).ab,ti.
13. transparent baby bag*.ab,ti.
14. baby bag*.ab,ti.
15. (headwrap* or polyester headwrap*).ti,ab.
16. silver swaddling*.ti,ab.
17. (silver and swaddling*).ti,ab.
18. swaddling*.ti,ab.
19. exp Incubators, Infant/
20. radiant warmer*.ti,ab.
21. exp Membranes, Artificial/
22. (semi-permeable membrane* or semipermeable membrane*).ti,ab.
23. exp POLYURETHANES/
24. polyurethane*.ti,ab.
25. kangaroo care*.ti,ab.
26. skin to skin contact*.ti,ab.
27. heat* mattress*.ti,ab.
28. chemical gel mattress*.ti,ab.
29. gel mattress*.ti,ab.

30. chemical mattress*.ti,ab.
31. head hood*.ti,ab.
32. head insulation*.ti,ab.
33. swaddl*.ti,ab.
34. plastic bag*.ti,ab.
35. early suck*.ti,ab.
36. Breast Feeding/
37. (breast feed* or breastfeed*).ti,ab.
38. skin-to-skin.ti,ab.
39. suckl*.ti,ab.
40. bath*.ti,ab.
41. radiant heater*.ti,ab.
42. radiant heat lamp*.ti,ab.
43. (mother* adj5 (bab* or infant*) adj5 (contact* or hold*)).ti,ab.
44. heated bed*.ti,ab.
45. resuscitat*.ti,ab.
46. exp RESUSCITATION/
47. (intervention* or procedure* or method*).ti,ab.
48. exp Infant, Low Birth Weight/
49. ((birth* or bab* or infant*) and (premature or pre-mature or preterm or pre-term or low weight or low birth weight or very low birth weight or VLBW or LBW)).ab,ti.
50. Infant, Newborn/
51. Body Temperature Regulation/ or Body Temperature/
52. HYPOTHERMIA/
53. (hypothermia or cold stress or heat loss or temperature regulation or body temperature).ab,ti.
54. temperature drop*.ti,ab.
55. heat loss.ti,ab.
56. or/1-47
57. or/48-50
58. or/51-55
59. 56 and 57 and 58

Searching other resources

We searched for cross references from included articles.

Data collection and analysis

We followed the standard method of The Cochrane Collaboration for conducting a systematic review as described in The Cochrane Reviewers' Handbook (Clarke 2002).

Selection of studies

The search strategy was designed and electronic databases searched in association with the Medical Faculty Librarians, Queen's University Belfast. At least three review authors independently assessed the full list of titles and abstracts for eligibility and the full texts of those considered to be relevant were retrieved. Reasons for exclu-

sion of studies were clearly stated by each review author. Formal translation of studies was not required.

Data extraction and management

The review authors separately extracted, assessed and coded all data for each study using a form that was designed specifically for this review. Any standard error of the mean was replaced by the corresponding standard deviation. Any disagreement was resolved by discussion.

Assessment of risk of bias in included studies

At least three independent review authors assessed those studies that fulfilled the criteria for inclusion for quality and risk of bias and extracted data using prepared proformas. Methodological quality and risk of bias was judged according to: (1) blinding of randomisation, (2) blinding of intervention, (3) completeness of follow-up and (4) blinding of outcome measurement. There was complete agreement among team members. Additional information was sought from investigators for four included trials (Vohra 2004a; Vohra 1999; Knobel 2005; Trevisanuto 2009b). Review authors were not blinded to authors or to institution. In two papers, a team member was also an author of a selected trial so that team member was excluded from the study appraisal process. The information retrieved is detailed in the Table of "Characteristics of included studies".

In addition, the following issues were evaluated and entered into the Risk of Bias Table:

1) Sequence generation (checking for possible selection bias). Was the allocation sequence adequately generated?

For each included study, we categorized the method used to generate the allocation sequence as:

- adequate (any truly random process e.g. random number table; computer random number generator);
- inadequate (any non random process e.g. odd or even date of birth; hospital or clinic record number);
- unclear.

(2) Allocation concealment (checking for possible selection bias). Was allocation adequately concealed?

For each included study, we categorized the method used to conceal the allocation sequence as:

- adequate (e.g. telephone or central randomization; consecutively numbered sealed opaque envelopes);
- inadequate (open random allocation; unsealed or non-opaque envelopes, alternation; date of birth);
- unclear.

(3) Blinding (checking for possible performance bias). Blinding of participants, personnel and outcome assessors: Was knowledge of the allocated intervention adequately prevented during the study? At study entry? At the time of outcome assessment?

For each included study, we categorized the methods used to blind study participants and personnel from knowledge of which in-

tervention a participant received. Blinding was assessed separately for different outcomes or classes of outcomes. We categorized the methods as:

- adequate, inadequate or unclear for participants;
- adequate, inadequate or unclear for personnel;
- adequate, inadequate or unclear for outcome assessors.

In some situations there may be partial blinding e.g. where outcomes are self-reported by unblinded participants but they are recorded by blinded personnel without knowledge of group assignment. Where needed "partial" was added to the list of options for assessing quality of blinding.

(4) Incomplete outcome data (checking for possible attrition bias through withdrawals, dropouts, protocol deviations). Were incomplete outcome data adequately addressed?

For each included study and for each outcome, we described the completeness of data including attrition and exclusions from the analysis. We noted whether attrition and exclusions were reported, the numbers included in the analysis at each stage (compared with the total randomized participants), reasons for attrition or exclusion where reported, and whether missing data were balanced across groups or were related to outcomes. Where sufficient information was reported or supplied by the trial authors, we re-included missing data in the analyses. We categorized the methods as:

- adequate (< 20% missing data);
- inadequate (\geq 20% missing data);
- unclear.

(5) Selective reporting bias. Are reports of the study free of suggestion of selective outcome reporting?

For each included study, we described how we investigated the possibility of selective outcome reporting bias and what we found. We assessed the methods as:

- adequate (where it is clear that all of the study's pre-specified outcomes and all expected outcomes of interest to the review have been reported);
- inadequate (where not all the study's pre-specified outcomes have been reported; one or more reported primary outcomes were not pre-specified; outcomes of interest are reported incompletely and so cannot be used; study fails to include results of a key outcome that would have been expected to have been reported);
- unclear.

(6) Other sources of bias. Was the study apparently free of other problems that could put it at a high risk of bias?

For each included study, we described any important concerns we had about other possible sources of bias (for example, whether there was a potential source of bias related to the specific study design or whether the trial was stopped early due to some data-dependent process). We assessed whether each study was free of other problems that could put it at risk of bias as: yes; no; or unclear.

If needed, we planned to explore the impact of the level of bias through undertaking sensitivity analyses.

Measures of treatment effect

Relative risk (RR) and 95% confidence limits were calculated for dichotomous outcomes. From the risk difference (RD), the number needed to treat (NNT) or the number needed to harm (NNH) and 95% confidence limits were calculated. Mean differences (MD) and 95% confidence limits were calculated for continuous outcomes.

Assessment of heterogeneity

We estimated the treatment effects of individual trials and examined heterogeneity between trials by inspecting the forest plots and quantifying the impact of heterogeneity using the I^2 statistic. If we detected statistical heterogeneity, we explored the possible causes (for example, differences in study quality, participants, intervention regimens, or outcome assessments) using *post hoc* sub group analyses.

Assessment of reporting biases

Where heterogeneity was identified, subgroup analyses were carried out for gestational age and birthweight. Had there been sufficient studies, sensitivity analyses would have provided valuable information on the role of variations in methodological quality, definitions of hypothermia, type of patient group and intervention.

Data synthesis

Meta-analysis was performed using Review Manager software (RevMan 5) supplied by the Cochrane Collaboration. For estimates of typical relative risk and risk difference, we used the Mantel-Haenszel method. For measured quantities, we used the inverse variance method. All meta-analyses were done using the fixed effect model.

Subgroup analysis and investigation of heterogeneity

Subgroup analyses to be carried out by intervention, by birthweight/gestational age and by developmental status of the country of study, to determine whether effectiveness varies according to: The interventions applied:

Birthweight and gestational age within the following categories:

- birthweight (< 1500 g), (1500 g to 2500 g)
- gestational age (< 28 wk), (28 to 32 wk) and (33 to 37 wk)

The developmental status of the country in which the trial was conducted based upon the UNICEF classification of a developing country (UNICEF 2002).

However, the subgroups reported in the included studies were not compatible with those pre-specified. Therefore, *post facto* subgroup analyses based on reported gestational age and birthweight sub-categories were carried out where appropriate within each comparison group.

Sensitivity analysis

Had there been more studies, the influence of the trial quality on the findings of the review would have been explored by conducting a sensitivity analysis of adequate versus unclear versus inadequate allocation concealment as outlined in The Cochrane Reviewers' Handbook (Clarke 2002).

RESULTS

Description of studies

See: [Characteristics of included studies](#); [Characteristics of excluded studies](#); [Characteristics of studies awaiting classification](#); [Characteristics of ongoing studies](#).

One hundred and sixty studies potentially eligible for inclusion in our review were identified. Of these, ninety-eight were randomised or quasi-randomised controlled studies and sixty-two were non-randomised studies. On assessment, eight studies were identified as fulfilling all of our criteria for inclusion. Ninety randomised or quasi randomised studies did not fulfill our criteria for inclusion and are detailed in the Table of "Characteristics of excluded studies". One randomised study is awaiting classification pending further details from the authors (Punnahitananda 2008). This study is detailed in the Table of "Characteristics of studies awaiting classification". We decided not to include in the Table of "Characteristics of excluded studies" non-randomised studies. Further details of these non-randomised trials can be obtained from the authors on request. In all, seven studies involving 400 randomised infants were included in this review: two theses (Brennan 1996; Roberts 1981), four published papers (Bergman 2004; Vohra 1999; Vohra 2004a; Knobel 2005) and one manuscript accepted for publication by the Journal of Pediatrics (Trevisanuto 2009b) and also available as a conference proceeding abstract.

Five studies compared barriers to heat loss to no barriers (Roberts 1981; Vohra 1999; Vohra 2004a; Knobel 2005; Trevisanuto 2009b) and two studies compared external heat sources (non-routine) to no external heat sources (Bergman 2004; Brennan 1996). There were three comparison groups within the barriers to heat loss category: plastic wrap or bag versus routine care, plastic cap versus routine care (Vohra 1999; Vohra 2004a; Knobel 2005; Trevisanuto 2009b) and stockinet cap versus routine care (Roberts 1981). There were also two comparison groups within the external heat source category: skin-to-skin care versus routine care (Bergman 2004) and transwarmer mattress versus routine care (Brennan 1996).

All interventions were applied immediately after birth in the delivery suite. Participants were categorised by gestational age (all preterm) in five studies and by birthweight (all low birthweight) in two studies. One study was conducted in a developing country, (South Africa) (Bergman 2004), and all other studies took

place in developed countries (USA: Brennan 1996; Roberts 1981; Knobel 2005, Canada: Vohra 1999; Vohra 2004a and Italy Trevisanuto 2009b). Routine external heat sources were utilised as part of care (control group and/or intervention group) in all studies: servo controlled incubator (Bergman 2004), radiant warmer (Brennan 1996; Vohra 1999; Vohra 2004a), warmer table (Knobel 2005) and radiant warmer ± transport incubator (Roberts 1981; Trevisanuto 2009b). The main outcome measure (temperature of the infant on admission to NICU or up to 2 hours after birth) was reported as a continuous variable in six studies and as a dichotomous variable only in one study (Bergman 2004). Six studies reported core body temperature (rectal:3, axillary:3) and in one study skin temperature was reported. In addition, four studies also provided dichotomous data pertaining to incidence of hypothermia in the intervention and control groups. Definitions of hypothermia were not consistent across studies. There was limited reporting of pre-specified secondary outcomes.

SUMMARY DESCRIPTIONS OF INDIVIDUAL STUDIES

Further details for each study are given in the Table of “Characteristics of included studies.”

BARRIERS TO HEAT LOSS

Roberts 1981

Participants

Forty inborn infants of 32 to 36 completed weeks' gestation born between 7 am to 5 pm on weekdays were included in this single centre study conducted at Parkland Memorial Hospital in Dallas, Texas, USA. Infants were excluded from the study if they were not appropriate for gestational age, had an Apgar score < 7 at 5 minutes, signs of central nervous system defect, developed sepsis or where the maternal temperature > 37.8 °C during labour.

Intervention Stockinet cap (CAP) - sterile headpiece made of stockinet that covers the forehead, the ears, underneath the occipital bone and all the area above this plane.

CAP group (n = 17): infants received a stockinet cap after delivery as soon as possible after being dried under a radiant warmer.

Control group (n = 23): infants received the same treatment as the intervention group but did not receive a stockinet cap.

All infants weighing < 2500 g were transported to the nursery in a transport incubator.

Outcomes

Core body temperature (axillary °C) was measured within 10 minutes of admission to the neonatal unit. The amount of heat lost from the time the axillary temperature was taken in the delivery room until the temperature was taken in the neonatal unit was also reported.

Vohra 1999

Participants

Sixty-two inborn infants of < 32 completed weeks' gestation, where the neonatal resuscitation team were present, were included in this single centre study conducted at the Grace Hospital (later the IWK Grace Health Centre) in Halifax, Nova Scotia, Canada.

Infants were excluded from the study if they had major congenital anomalies with open lesions and where the infant was considered “preivable” by the attending Obstetrician.

Intervention Polyethylene bag (WRAP) - 20cm x 50cm manufactured by Eastern Paper, a Division of EPC Industries.

WRAP group (n = 27): a transparent polyethylene bag was opened at resuscitation under a radiant warmer and the infant was placed on it from the shoulders down. Only the head of the infant was dried; the body was wrapped without drying.

Control group (n = 32): infants were managed with the protocol described by the Neonatal Resuscitation Program. Infants were routinely dried under the radiant warmer.

All infants were transferred to NICU in incubators.

Outcomes

Core body temperature (rectal °C) was measured on removal of the bag/wrap on admission to the neonatal unit. Incidence of mortality, hyperthermia, infection, skin maceration and interference with resuscitation were also reported.

Vohra 2004a

Participants

Fifty-five inborn infants of < 28 completed weeks' gestation, where the neonatal resuscitation team were present, were included in this single centre study conducted at McMaster University Medical Center in Hamilton, Ontario, Canada. Infants were excluded from the study if they had major congenital anomalies that were not covered by the skin and blistering skin conditions.

Intervention Polyethylene wrap (WRAP) - 20cm X 50cm manufactured by Eastern Paper, a Division of EPC Industries.

WRAP group (n = 26): infants were placed on polyethylene, wrapped from the neck down, only the head was dried and stabilised under a radiant warmer.

Control group (n = 27): infants were dried completely according to the International Guidelines for Neonatal Resuscitation and stabilised under a radiant warmer.

All infants were carried by one member of the neonatal team from the delivery room to the neonatal unit and placed in a single walled incubator with 60 percent humidity.

Outcomes

Core body temperature (rectal °C) on removal of the bag/wrap on admission to the neonatal unit, core body temperature (rectal °C) 1 hour later and incidence of mortality from all causes were reported. Secondary outcome measures have been updated as reported in the publication of the full manuscript (Vohra 2004a): Apgar score at 1 minute, Apgar score at 5 minutes, blood gas pH, Bicarbonate (mmol/L), Glucose mmol/L, hyperthermia (rectal °C > 37.5) and interference with resuscitation.

Knobel 2005

Participants

Eighty-eight infants of < 29 completed weeks' gestation were included in this single centre study conducted at Pitt County Memorial Hospital, Greenville, NC, USA. Infants were excluded from

the study if they had congenital anomalies with open lesions, meconium staining of the amniotic fluid or if they were considered pre-viable or subsequent assessment indicated that the infant was ≥ 29 completed weeks' gestation.

Intervention Polyurethane bag (WRAP) - 19" X 18" sterile isolation transport bag DeRoyal REF30-5510.

WRAP group (n = 41): infants were placed immediately into a polyurethane bag, on a radiant warmer bed, while still wet, up to their necks. The head and face were dried and infants were resuscitated according to the guidelines for Neonatal Resuscitation.

Control group (n = 47): infants were resuscitated according to these guidelines without the bags.

All infants were taken to the neonatal unit on a warmer table, (with the heat off) with warm blankets on top of the infant.

Outcomes

Core body temperature (rectal °C) and incidence of hypothermia (rectal °C < 36.4) on admission to the neonatal unit, incidence of mortality, major brain injury, duration of oxygen therapy and hospitalisation, incidence of hyperthermia and interference with assessment or resuscitative interventions.

[Trevisanuto 2009b](#)

Participants

Ninety-six infants of < 29 completed weeks' gestation were included in this single centre study conducted at the Pediatric Department, Medical School, University of Padua, Azienda Ospedakiera di Padova, Padua, Italy. Infants were excluded if they had congenital anomalies with open lesions (e.g. gastroschisis, meningomyelocele) and if the delivery was not attended by the neonatal team.

Intervention Polyethylene cap (CAP) - (Degusto Dolomiti, Belluno, Italy)

CAP group (n = 32): polyethylene caps were placed on the radiant warmer bed prior to delivery. The heads of the infants were covered with a polyethylene cap immediately after birth; only the head was dried.

Active comparator group (n = 32): polyethylene bags (Vedovato SNC, Camposampiero, Padova, Italy, isolation transport bag 35 cm by 40 cm) were placed on the radiant warmer bed prior to delivery. The infants were placed into the bag while still wet, up to the neck; only the head was dried.

Control group (n = 32): the infants were placed in pre-warmed towels after drying, according to International Guidelines for Neonatal Resuscitation. All infants were stabilised in the delivery room under radiant warmers and a transport incubator was used to transfer all infants from delivery room into the NICU where they were then placed in a double walled incubator and at this point, the cap or the wrap were removed.

Outcomes

Core body temperature (axillary °C) and incidence of hypothermia (axillary temperature < 36.4°C) on admission to the neonatal unit (immediately after cap and wrap removal) and again one hour later, incidence of: mortality prior to hospital discharge, major

brain injury and requirement for endotracheal intubation at birth, Apgar scores, blood gas analysis and serum glucose concentration on NICU admission.

EXTERNAL HEAT SOURCES (NON-ROUTINE)

[Bergman 2004](#)

Participants

Thirty-five inborn infants delivered, (excluding caesarean sections), having a birthweight between 1200 and 2199 g were included in this study conducted at two secondary level referral hospitals (Mowbray Maternity Hospital and Karl Brenar Hospital) in Cape Town, South Africa. Infants were excluded from the study if they had a birthweight below 1200 g or above 2199 g, an Apgar score below six at five minutes, congenital malformations detected at birth or where the mother was too severely ill to be able to look after herself or the infant, was known to have a positive HIV status or had given up the infant for adoption.

Intervention: Skin-to-skin care (SSC)

All infants were delivered onto a theatre cloth on the mother's abdomen/chest, dried, assessed, then placed on the mother's naked chest and covered with double layered cotton cloth. The bed was then placed in the semi-Fowlers position.

SSC group (n = 18): the infant remained in skin-to-skin contact, was secured to the mother's chest, hips and arms were flexed and placed in a 'frog' position.

Control group (n = 13): the infant was immediately transferred to a pre-warmed servo-controlled closed incubator which remained with the mother in the delivery ward for the first hour. A cap, booties and heat shield were applied if the infant skin temperature fell below 36 °C.

Outcomes

Infant stability was measured in terms of a set of pre-defined physiological parameters. Observations for heart rate, oxygen saturation and body temperature (skin °C) were recorded at five minute intervals in the first half hour, then 15 minute intervals. Continuous observations for apnoea and signs of respiratory distress, and blood sugar were measured at one, three and six hours.

[Brennan 1996](#)

Participants

Twenty-four inborn infants having a birthweight ≤ 1500 g were included in this single centre study conducted at Shands Hospital located in north-central Florida, USA. Infants were excluded from the study where the maternal temperature at delivery was > 38.0 °C, the infant had an obvious neural tube defect, an omphalocele, gastroschisis, extrophy of the bladder, cloacal exstrophy, any other open lesion that would cause greater than normal heat loss at delivery, sacral teratoma, or any other lesion that would prevent the infant from being positioned and evaluated in the normal manner.

Intervention: Transwarmer Infant Transport Mattress (TM) manufactured by Prism Technologies, San Antonio, Texas. These are filled with a gel of sodium acetate, water and thickeners. Once activated they heat to approximately 40 °C for two hours.

TM group (n = 12): the mattress was activated and placed on the

radiant warmer surface with blankets on top of it when delivery was imminent. The newborn infant was immediately placed on the blankets and dried. Blankets were removed and the infant was then placed directly onto the warming mattress.

Control group (n = 12): infants received the same care as the intervention group but without the warming mattress.

Both groups were evaluated and resuscitated according to the Neonatal Resuscitation Program and transported to the neonatal unit on the radiant warmer surface. The infants in the intervention group remained on the mattress until the admission temperatures were taken.

Outcomes

Core body temperature (axillary °C) was measured as soon as possible after arrival at the neonatal unit. Incidence of interference with resuscitation was also monitored. Individual patient data was reported in this study.

Risk of bias in included studies

BARRIERS TO HEAT LOSS

The generation of allocation sequence was adequate (computer randomisation or lot drawing) in four studies and unclear in one. Allocation concealment was adequate in four studies using varying degrees of sealed/opaque/double enclosed envelopes and in one study the method was unclear. However, two studies [Vohra 2004a](#) and [Trevisanuto 2009b](#) used computer generated random sequence balanced in blocks of four and six subjects respectively. There is some potential for inadequate allocation concealment, therefore, selection bias cannot be ruled out. No attempt was reported in any of the studies to blind participants or care givers to the intervention. However, lack of blinding is often not feasible for non-pharmacological interventions [Boutron 2008](#). The majority of outcome measures were objective and so were less likely to be biased than subjective outcome measures. The primary outcome (temperature of the infant on admission to NICU or up to two hours after birth) was measured using digital thermometers in all studies. No studies reported any attempt to blind the outcome assessors or data analysis team to the intervention, therefore, potential biases cannot be ruled out.

Follow-up was complete in three studies. In the remaining two studies [Vohra 1999](#) (3 of 62 or 5%) and [Vohra 2004a](#) (2 of 55 or 4%) of infants randomised were lost to follow-up. Reasons for loss to follow-up were reported adequately in each study. There was insufficient information to assess whether these studies were free from selective reporting.

Other potential bias for barriers to heat loss:

In two studies, [Knobel 2005](#) and [Vohra 1999](#) the authors attempted to allay fears of potential confounding factors (other biases) by statistically adjusting for this and found that the main conclusions remained unchanged.

The [Knobel 2005](#) study attempted to maintain delivery suite temperature at 26°C for all preterm deliveries, but actual temperatures

ranged from 18.9 °C to 31.1 °C. On post hoc analysis warmer delivery suite temperatures were associated with higher admission temperatures, but only the subgroup of infants who were both delivered in warm rooms and placed in plastic bags had a mean temperature > 36.4 °C. There was, however, no significant differences between intervention and control group for mean delivery room temperature. After controlling for delivery suite temperature the mean admission temperature to NICU in the intervention group was still 0.6 °C higher than controls.

There was some imbalance in birthweight between study groups in [Vohra 1999](#). However the birthweight adjusted difference in mean rectal temperature of 1.54°C for the smaller group remained significant. For infants < 28 completed weeks birthweight (g) was mean (914) SD (163) for the plastic wrap group and mean (742) SD (206) for the non-wrap group, therefore, results could be potentially biased towards the wrap group since the non-wrap group were smaller and, therefore, more vulnerable to heat loss.

The [Trevisanuto 2009b](#) study reported that delivery to admission time was significantly shorter in the control group than in the plastic wrap group. This measure could be an indicator of potentially how unwell an infant is and to see if the team behaved differently towards intervention infants than control infants. If the plastic wrap infants took longer to get to the NICU and therefore these infants had a greater chance of being chilled then the plastic wrap measured estimate of treatment effect is likely to be a conservative one. [Trevisanuto 2009b](#) also reported that delivery suite temperature was not monitored throughout the study. Standard environmental temperature of delivery suites within the institution was reported to be 24°C. The authors felt that this may have negatively influenced the findings of the study.

The [Roberts 1981](#) study reported some imbalance in mean delivery suite axillary temperatures for infants < 2000 g between study groups. Analysis of covariance showed that the delivery suite axillary temperature had a significant effect on the infant axillary temperature on admission to the NICU. When the two groups were statistically equalised with respect to delivery suite axillary temperature there were no significant differences between the two interventions.

EXTERNAL HEAT SOURCES (NON-ROUTINE)

The generation of allocation sequence was adequate (computer randomisation or lot drawing) in both studies, however, the method of allocation concealment for each study was unclear. No attempt was reported in any of the studies to blind participants or care givers to the intervention. However, lack of blinding is often not feasible for non-pharmacological interventions [Boutron 2008](#). The majority of outcome measures were objective and so were less likely to be biased than subjective outcome measures. The primary outcome (temperature of the infant on admission to NICU or up to two hours after birth) was measured using a digital thermometer in [Brennan 1996](#) and a dedicated and regularly calibrated monitor (Dash 3000) in [Bergman 2004](#). Follow-up was complete in one study. In the remaining study [Bergman 2004](#) (4 of

35 or 11%) of infants randomised were lost to follow-up. Reasons for loss to follow-up were adequately reported. Only one study [Bergman 2004](#) reported any attempt to blind the outcome assessors to the intervention where the principal investigator and data analysis team were blinded. There was insufficient information to assess whether these studies were free of selective reporting.

Other potential bias for external heat sources:

For the [Bergman 2004](#) study there was a potential for selection bias since the assigned research nurse was unavailable for 99 potentially eligible mother-infant dyads. The aetiology of these infants may have differed from those studied. In addition, recruitment for this study was terminated based on significant results after an interim analysis was conducted.

In all, no studies in either category fulfilled all the methodological quality criteria. Risk of bias across studies is considered to be low to moderate. Further details for each study are given in the Table of “Characteristics of included studies.”

Effects of interventions

RESULTS OF META-ANALYSES

Seven studies were identified which fulfilled our criteria for inclusion giving a total of 400 infants randomised and 391 infants completing the studies. Five studies compared barriers to heat loss to no barriers giving a total of 341 infants randomised and 336 infants completing the studies ([Roberts 1981](#); [Vohra 1999](#); [Vohra 2004a](#); [Knobel 2005](#); [Trevisanuto 2009b](#)). Two studies compared external heat sources (non-routine) to no external heat sources giving a total of 59 infants randomised and 55 infants completing the studies ([Brennan 1996](#); [Bergman 2004](#)). There were three comparisons within the barriers to heat loss category: plastic wrap or bag versus routine care ([Vohra 1999](#); [Knobel 2005](#); [Vohra 2004a](#)), plastic cap versus routine care ([Trevisanuto 2009b](#)) and stockinet cap versus routine care ([Roberts 1981](#)). There were also two comparisons group within the external heat source category: skin-to-skin care versus routine care ([Bergman 2004](#)) and transwarmer mattress versus routine care ([Brennan 1996](#)). There were insufficient studies to carry out subgroup analysis according to developmental status of the country.

BARRIERS TO HEAT LOSS

PLASTIC WRAP VERSUS ROUTINE CARE (COMPARISON 1):

Primary outcomes

Core body temperature (°C) of the infant taken on admission to NICU or up to 2 hours after birth (Outcome 1.1):

Four studies comprising 264 infants ([Vohra 1999](#); [Vohra 2004a](#); [Knobel 2005](#); [Trevisanuto 2009b](#)) reported core body temperature (rectal °C or axillary °C) on admission to NICU. Each individual study showed a significant effect in favour of the intervention (plastic wrap) group for infants with a gestational age < 28 completed weeks. Although the infants in the [Knobel 2005](#) and [Trevisanuto 2009b](#) studies had a gestational age < 29 completed

weeks, these infants were included in the < 28 completed week subgroup for meta-analysis and are referred to as such in the text unless all the included studies participants were of < 29 completed weeks' gestational age.

Overall

For infants with a gestational age < 32 completed weeks, a statistically significant difference was shown in core body temperature on admission to NICU favouring the intervention (plastic wrap) group when compared to those who received routine care immediately after birth in the delivery suite (four studies, n = 264; WMD 0.57°C 95% CI 0.37, 0.77).

However, the overall test for homogeneity and for the subgroup of infants with a gestational age < 28 completed weeks failed with an I² value of 75%. We note the most likely reason for this failure is the contribution of the small [Vohra 1999](#) study for the subgroup of infants < 28 completed weeks which gives rise to a 95% CI range somewhat outside the range suggested by the other studies. Rather than excluding this study purely on statistical grounds the possible reasons for this heterogeneity are explored in the discussion section of this review. The outcomes reported below suggest that the heterogeneity issue has been dealt with by dividing the population into the two gestational age subgroups: < 28 completed weeks and 28 to 31 completed weeks.

Outcome 1.1.1: For infants with a gestational age < 28 completed weeks a statistically significant difference was shown in core body temperature on admission to NICU favouring the intervention (plastic wrap) group when compared to those who received routine care immediately after birth in the delivery suite (four studies, n = 223; WMD 0.68 °C; 95% CI 0.45, 0.91).

Outcome 1.1.2: For infants with a gestational age of 28 to 31 completed weeks there were no statistically significant differences between the two interventions (plastic wrap and routine care) for core body temperature on admission to NICU (one study, n = 41; MD 0.17 °C; 95% CI -0.27, 0.61).

Core body temperature taken 1 hour after initial admission temperature to the NICU was taken (Outcome 1.2):

This outcome was reported in one small study ([Vohra 2004a](#)) in terms of core body temperature (rectal °C) one hour after the initial admission temperature was taken and in an additional study in terms of core body temperature (axillary °C) ([Trevisanuto 2009b](#)). This was not pre-defined at review protocol stage. The reason stated for collection of this outcome measure was to ascertain whether the intervention (plastic wrap) prevented rather than delayed the postnatal fall in body temperature immediately after birth.

Outcome 1.2.1 For infants with a gestational age < 28 completed weeks a statistically significant difference in the effect was shown for core body temperature one hour after the initial admission temperature to NICU was taken favouring the intervention (plastic wrap) group when compared to those who received routine care immediately after birth in the delivery suite (two studies, n= 117; WMD 0.40 °C; 95% CI 0.16, 0.65). The test for homogeneity

passed with an I^2 value of 20%.

Hypothermia on admission to NICU (core body temperature < 36.5 °C or skin temperature < 36 °C) (Outcome 1.3):

Outcome 1.3.1 [Knobel 2005](#) and [Trevisanuto 2009b](#) in addition to reporting core body temperature on admission to NICU as a continuous variable, also provided data in a dichotomous format in terms of incidence of hypothermia in intervention and control groups (plastic wrap and routine care). Hypothermia was defined in both studies as a core body temperature (rectal or axillary) < 36.4 °C on admission to NICU. For infants of gestational age < 29 completed weeks, plastic wrap significantly reduces the risk of hypothermia on admission to NICU (two studies, $n = 152$; RR 0.66, 95% CI 0.51, 0.84; RD -0.27; 95% CI -0.41, -0.13). This finding is consistent with those for the outcome measure 01.01.01. Four infants would need to be wrapped in plastic in order to prevent one infant from becoming hypothermic (NNT 4, 95% CI 2 to 8).

Secondary outcomes

Mortality (death prior to hospital discharge) (Outcome 1.4):

Four studies ([Vohra 1999](#); [Vohra 2004a](#); [Knobel 2005](#); [Trevisanuto 2009b](#)) reported this outcome in terms of death within hospital stay. Mortality figures for the [Vohra 2004a](#) study included two infants who died in the delivery suite. The test for homogeneity passed with an I^2 value of 0%.

Outcome 1.4.1: For infants with a gestational age < 28 completed weeks the results of the meta-analysis are in the direction in favour of the intervention group (plastic wrap). However, this is being influenced by the ([Vohra 1999](#)) study with five events in the control group and showing borderline significance. Overall there is insufficient evidence to suggest that plastic wrap reduces the risk of death within hospital stay for this group of infants (four studies, $n = 225$; RR 0.66; 95% CI 0.35, 1.24; RD -0.06 ; 95% CI -0.15, 0.03).

Outcome 1.4.2: For infants with a gestational age of 28 to 31 completed weeks there were no deaths within hospital stay in either group. Data were provided from one study [Vohra 1999](#) ($n = 41$ infants).

Major brain injury (defined as sonographic evidence of intraventricular haemorrhage with ventricular dilatation, parenchymal haemorrhagic infarction or periventricular leukomalacia) (Outcome 1.5):

This outcome was reported in two studies ([Knobel 2005](#); [Trevisanuto 2009b](#)) but it was not pre-defined at review protocol stage.

Outcome 1.5.1: For infants of gestational age < 29 completed weeks there was no evidence of a difference in risk of major brain injury (two studies, $n = 152$; RR 1.10; 95% CI 0.41, 2.98; RD 0.01; 95% CI -0.08, 0.10). The test for homogeneity passed with an I^2 value=0%.

Duration of oxygen therapy (days) (Outcome 1.6):

This outcome was reported in one study ([Knobel 2005](#)) but it was not pre-defined at review protocol stage.

Outcome 1.6.1: For infants of gestational age < 29 completed weeks there were no statistically significant differences between the two interventions (plastic wrap and routine care) for duration of oxygen therapy (one study, $n = 88$; MD -6.51 days; 95% CI -23.30, 10.28).

Duration of hospitalisation (days) (Outcome 1.7):

This outcome was reported in one study ([Knobel 2005](#)).

Outcome 1.7.1: For infants of gestational age < 29 completed weeks there were no statistically significant differences between the two interventions (plastic wrap and routine care) for duration of hospitalisation (one study, $n = 88$; MD -5.49 days; 95% CI -19.93, 8.95).

Apgar score at one minute (Outcome 1.8):

This outcome was reported as a median value with interquartile ranges in [Vohra 2004a](#) and as a mean value with SD in [Trevisanuto 2009b](#) but it was not pre-defined at review protocol stage.

Outcome 1.8.1: For infants of gestational age < 28 completed weeks, median Apgar scores at one minute were comparable in the intervention and control group (one study, $n = 53$; $P = 0.6$).

Apgar score at one minute (Outcome 1.9):

Outcome 1.9.1: For infants of gestational age < 29 completed weeks there were no statistically significant differences between the two interventions (plastic wrap and routine care) for Apgar scores at one minute (one study, $n = 64$; MD 0.20; 95% CI -0.83, 1.23).

Apgar score at five minutes (Outcome 1.10):

This outcome was reported as a median value with interquartile ranges in ([Vohra 2004a](#)) and as a mean value with SD in [Trevisanuto 2009b](#) but it was not pre-defined at review protocol stage.

Outcome 1.10.1: For infants of gestational age < 28 completed weeks, median Apgar scores were comparable in the intervention and control group (one study, $n = 53$; $P = 0.9$).

Apgar score at five minutes (Outcome 1.11):

Outcome 1.11.1: For infants of gestational age < 29 completed weeks there were no statistically significant differences between the two interventions (plastic wrap and routine care) for Apgar score at five minutes (one study, $n=64$; MD 0.40; 95% CI -0.19, 0.99).

First blood gas pH (Outcome 1.12):

This outcome was reported in two studies ([Vohra 2004a](#); [Trevisanuto 2009b](#)). This was not pre-specified at review protocol stage.

Outcome 1.12.1: For infants of gestational age < 28 completed weeks there were no statistically significant differences between the two interventions (plastic wrap and routine care) for first blood gas pH (two studies, $n = 117$; MD 0.01 ; 95% CI -0.02, 0.04). The test for homogeneity failed with an I^2 value of 70%. Since pH is on a logarithmic scale it would be more appropriate to measure this outcome as a median value and range or interquartile range (IQR) or to avoid its use as a continuous variable such as that pre-specified at review protocol stage: severe metabolic acidosis as defined by pH < 7.20 and/or base deficit > 10 mmol/L within the first three days of life.

Bicarbonate (mmol/L) (Outcome 1.13):

This outcome was reported in two studies (Vohra 2004a; Trevisanuto 2009b). This was not pre-specified at review protocol stage.

Outcome 1.13.1: For infants of gestational age < 28 completed weeks there were no statistically significant differences between the two interventions (plastic wrap and routine care) for bicarbonate (two studies, n = 117; WMD 0.31 mmol/L; 95% CI -0.72, 1.35).

First serum glucose concentration (mmol/L) (Outcome 1.14):

This outcome was reported in two studies (Vohra 2004a; Trevisanuto 2009b). This was not our pre-specified definition of hypoglycaemia at review protocol stage.

Outcome 1.14.1: For infants of gestational age < 28 completed weeks there were no statistically significant differences between the two interventions (plastic wrap and routine care) for first serum glucose concentration (two studies, n = 117; WMD 0.01 mmol/L; 95% CI , -0.36, 0.39).

Intubation in delivery room (Outcome 1.15):

This outcome was reported in one study (Trevisanuto 2009b).

Outcome 1.15.1: For infants of gestational age < 29 completed weeks, there was no evidence of a difference in risk of intubation in delivery room (one study, n = 64; RR 1.00; 95% CI 0.63, 1.58; RD 0.00; 95% CI -0.24, 0.24).

Other secondary outcomes

The following secondary outcome measures were not reported (as pre-defined at review protocol stage) for any of the included studies for this comparison group: hypoglycaemia, respiratory distress syndrome, surfactant given at any time, requirement for ventilation, duration of ventilation, severe metabolic acidosis, intraventricular haemorrhage, patent ductus arteriosus, chronic lung disease, necrotising enterocolitis, acute renal failure, burns, antibiotics or negative psychological outcomes.

Adverse occurrences

There were no occurrences of adverse events attributable to the intervention (i.e. hyperthermia, infection, skin maceration or interference with resuscitation) for infants in the intervention and control groups in the Vohra 1999 study. Knobel 2005 reported that one participant in the intervention group was hyperthermic with a rectal admission temperature of 38.3°C; the delivery room environmental temperature was 26.7 °C. Knobel 2005 also reported that the clear, pliable polyurethane bag did not interfere with either assessment (visualization, auscultation, palpation) or resuscitative interventions. Vohra 2004a reported that the wrap procedure was accepted by the neonatal staff and did not interfere with resuscitation in the delivery room. Two infants in the intervention group (Vohra 2004a) were reported to have a rectal temperature on admission above 37.5 °C. Two infants in the plastic wrap group (active comparator) (Trevisanuto 2009b) were reported to have an axillary admission temperature > 37.5 °C (37.6 °C and 38 °C respectively).

Hyperthermia on admission to NICU: core body temperature > 37.5°C (Outcome 1.16):

A clear definition of hyperthermia was reported in only one study (Vohra 2004a). This was not the pre-specified definition of hyperthermia at review protocol stage.

Outcome 1.16.1: For infants of gestational age < 28 completed weeks, there was no significant difference in risk of hyperthermia between those infants who received plastic wrap and those who received routine care (one study, n = 53; RR 4.82, 95% CI 0.24, 95.88; RD 0.07; 95% CI -0.04, 0.19).

PLASTIC CAP VERSUS ROUTINE CARE (COMPARISON 2):**Primary outcomes**

Core body temperature (°C) of the infant taken on admission to NICU or up to 2 hours after birth (Outcome 2.1):

One study Trevisanuto 2009b reported core body temperature (axillary °C) on admission to NICU.

Outcome 2.1.1: For infants with a gestational age < 29 completed weeks a statistically significant difference in the effect was shown for core body temperature on admission to NICU favouring the intervention (plastic cap) group when compared to those who received routine care immediately after birth in the delivery suite (one study, n = 64; MD 0.80 95% CI 0.41, 1.19).

Core body temperature taken 1 hour after initial admission temperature to the NICU was taken (Outcome 2.2):

This outcome was reported in one study in terms of core body temperature (axillary °C) (Trevisanuto 2009b). This was not pre-defined at review protocol stage.

Outcome 2.2.1: For infants with a gestational age < 29 completed weeks a statistically significant difference in the effect was shown for core body temperature taken one hour after initial admission temperature to the NICU was taken favouring the intervention (plastic cap) group when compared to those who received routine care immediately after birth in the delivery suite (one study, n = 64; MD 0.80 95% CI 0.46, 1.14).

Hypothermia on admission to NICU (core body temperature < 36.5 °C or skin temperature < 36 °C) (Outcome 2.3):

Trevisanuto 2009b in addition to reporting core body temperature on admission to NICU as a continuous variable, also provided data in a dichotomous format in terms of incidence of hypothermia in intervention and control groups (plastic cap and routine care). Hypothermia was defined as a core body temperature (axillary) < 36.4 °C on admission to NICU.

Outcome 2.3.1: For infants of gestational age < 29 completed weeks, plastic cap significantly reduces the risk of hypothermia on admission to NICU (one study, n = 64; RR 0.48, 95% CI 0.32, 0.73; RD -0.47; 95% CI -0.67, -0.27).

This finding is consistent with those for the outcome measure 02.01.01. Two infants would need to wear a plastic cap in order to prevent one infant from becoming hypothermic (NNT 2, 95% CI 2 to 4).

Mortality (death prior to hospital discharge) (Outcome 2.4):

Trevisanuto 2009b reported this outcome in terms of death within

hospital stay.

Outcome 2.4.1: For infants of gestational age < 29 completed weeks there is insufficient evidence to suggest that plastic cap reduces the risk of death within hospital stay for this group of infants (one study, n = 64; RR 1.50; 95% CI 0.27, 8.38; RD 0.03; 95% CI -0.10, 0.16).

Major brain injury (defined as sonographic evidence of intraventricular haemorrhage with ventricular dilatation, parenchymal haemorrhagic infarction or periventricular leukomalacia) (Outcome 2.5):

This outcome was reported in one study (Trevisanuto 2009b) but it was not pre-defined at review protocol stage.

Outcome 2.5.1: For infants of gestational age < 29 completed weeks there was no evidence of a difference in risk of major brain injury (one study, n = 64; RR 1.50; 95% CI 0.27, 8.38; RD 0.03; 95% CI -0.10, 0.16).

Apgar score at one minute (Outcome 2.6):

This outcome was reported as a mean value with SD in Trevisanuto 2009b but it was not pre-defined at review protocol stage.

Outcome 2.6.1: For infants of gestational age < 29 completed weeks there were no statistically significant differences between the two interventions (plastic cap and routine care) for Apgar scores at one minute (one study, n=64; MD 0.80; 95% CI -0.21, 1.81).

Apgar score at five minutes (Outcome 2.7):

This outcome was reported as a mean value with SD in Trevisanuto 2009b but it was not pre-defined at review protocol stage.

Outcome 2.7.1: For infants of gestational age < 29 completed weeks a statistically significant difference in the effect was shown for Apgar score at five minutes favouring the intervention (plastic cap) group when compared to those who received routine care immediately after birth in the delivery suite (one study, n=64; MD 0.70 95% CI 0.08, 1.32).

First blood gas pH (Outcome 02.08):

This outcome was reported in one study (Trevisanuto 2009b). This was not pre-specified at review protocol stage.

Outcome 02.08.01: For infants of gestational age < 29 completed weeks there were no statistically significant differences between the two interventions (plastic cap and routine care) for first blood gas pH (one study, n = 64; MD 0.01 ; 95% CI -0.03 0.05). Since pH is on a logarithmic scale it would be more appropriate to measure this outcome as a median value and range or interquartile range (IQR) or to avoid its use as a continuous variable such as that pre-specified at review protocol stage: severe metabolic acidosis as defined by pH < 7.20 and/or base deficit > 10 mmol/L within the first three days of life.

Bicarbonate (mmol/L) (Outcome 2.9):

This outcome was reported in one study (Trevisanuto 2009b).

Outcome 2.9.1: For infants of gestational age < 29 completed weeks there were no statistically significant differences between the two interventions (plastic cap and routine care) for bicarbonate (one study, n = 64; MD 1.00 mmol/L; 95% CI -0.25, 2.25).

First serum glucose concentration (mmol/L) (Outcome 2.10):

This outcome was reported in one study (Trevisanuto 2009b). This was not our pre-specified definition of hypoglycaemia at review protocol stage.

Outcome 2.10.1: For infants of gestational age < 29 completed weeks there were no statistically significant differences between the two interventions (plastic cap and routine care) for first serum glucose concentration (one study, n = 64; MD 0.10 mmol/L; 95% CI, -0.42, 0.62).

Intubation in delivery room (Outcome 2.11):

This outcome was reported in one study (Trevisanuto 2009b).

Outcome 2.11.1: For infants of gestational age < 29 completed weeks, there was no evidence of a difference in risk of intubation in delivery room (one study, n = 64; RR 0.82; 95% CI 0.49, 1.37; RD -0.09; 95% CI -0.34, 0.15).

Other secondary outcomes

The following secondary outcome measures (as pre-defined at review protocol stage) were not reported for this comparison group: hypoglycaemia, respiratory distress syndrome, surfactant given at any time, requirement for ventilation, duration of ventilation, severe metabolic acidosis, intraventricular haemorrhage, patent ductus arteriosus, chronic lung disease, necrotizing enterocolitis or acute renal failure.

Adverse occurrences

The following adverse outcome measures (as pre-defined at review protocol stage) were not reported for this comparison group: hyperthermia, burns, infection, antibiotics, skin maceration, interference with resuscitation or negative psychological outcomes for infants in the intervention and control group.

STOCKINET CAP VERSUS ROUTINE CARE (COMPARISON 3):

Primary outcomes

Core body temperature (°C) on admission to NICU or up to 2 hours after birth (Outcome 3.1):

Roberts 1981 reported core body temperature (axillary °C) on admission to NICU in a study of 40 infants. This study reported figures for all infants with a gestational age 32 to 36 completed weeks, and also for the subgroup of infants < 2000 g birthweight. We have tried to disaggregate the data using the information available, in order to obtain data for the subgroup of infants weighing 2000 g or more at birth. These subgroup analyses by birthweight class were not pre-specified in the protocol for this review.

Overall:

The I² value of 65.8% indicates that there is a moderate degree of heterogeneity of effect across these two birthweight subgroups. Overall, the results show no statistically significant difference in the effects of the two interventions (stockinet cap versus routine care) on core body temperature on admission to NICU (one study, n = 40; MD 0.15 °C; 95% CI -0.18, 0.48).

Outcome 3.1.1: For infants with a birthweight < 2000 g a borderline statistically significant difference was shown in core body

temperature on admission to NICU favouring the intervention (stockinet cap) group when compared to those who received routine care (one study, n = 10; MD 0.70 °C; 95% CI -0.01, 1.41).

Outcome 3.1.2: For infants with a birthweight \geq 2000 g there were no statistically significant differences between the two interventions (stockinet cap versus routine care) for core body temperature on admission to NICU (one study, n = 30; MD 0.00 °C; 95% CI -0.37, 0.37).

Hypothermia on admission to NICU (core body temperature < 36.5 °C or a skin temperature < 36 °C) (Outcome 3.2):

Outcome 3.2.1: Roberts 1981, in addition to reporting core body temperature on admission to NICU as a continuous variable, also provided data in a dichotomous format in terms of incidence of hypothermia in both groups (stockinet cap and routine care). Hypothermia was defined in this study as a core body temperature (axillary) < 36 °C on admission to the NICU. For infants with a gestational age 32 to 36 completed weeks (birthweight range 1360 to 2965 g), there was no significant difference in risk of hypothermia (as defined by the study) between those infants who received stockinet cap and those who received routine care (one study, n = 40; RR 0.90; 95% CI 0.48, 1.71; RD -0.05; 95% CI -0.36, 0.26).

Secondary outcomes

The following secondary outcome measures (as pre-defined at review protocol stage) were not reported for this comparison group: hypoglycaemia, respiratory distress syndrome, surfactant given at any time, intubation in delivery room, requirement for ventilation, duration of ventilation, length of stay, mortality, severe metabolic acidosis, intraventricular haemorrhage, patent ductus arteriosus, chronic lung disease, necrotizing enterocolitis, acute renal failure and adverse events due to the intervention (hyperthermia, burns, maceration, skin or systemic infection, antibiotics, interference with resuscitation and other practices, fluid problems, negative psychological outcomes).

EXTERNAL HEAT SOURCES (NON-ROUTINE)

SKIN-TO-SKIN CARE VERSUS ROUTINE CARE (COMPARISON 4):

Primary outcomes

Hypothermia (skin temperature < 35.5 °C for two consecutive recordings) (Outcome 4.1):

Bergman 2004 reported this outcome in terms of skin temperature remaining below 35.5 °C for two consecutive readings (five minute intervals for the first hour, thereafter 15 minute intervals during the six hour observation period). This outcome was not pre-specified at review protocol stage.

Outcome 4.1.1: For infants with a birthweight between 1200 and 2199 g evidence suggests that skin-to-skin contact significantly reduces the risk of hypothermia (as defined by the study) within 6 hours of birth when compared to conventional incubator care (one study, n = 31; RR 0.09; 95% CI 0.01, 0.64; RD -0.56; 95% CI -0.84, -0.27). Two infants would need to receive skin-to-skin contact in order to prevent one infant from becoming hypothermic

(NNT 2, 95% CI 1 to 4).

Secondary outcomes

Hypoglycaemia (blood glucose < 2.6 mmol/L) (Outcome 4.2): Bergman 2004 reported this outcome which was confirmed by laboratory estimation within the 6 hour observation period where blood glucose was measured by heel prick at one, three and six hours. This was not the pre-specified definition of hypoglycaemia at review protocol stage.

Outcome 4.2.1: For infants with a birthweight between 1200 and 2199 g there is no significant difference in risk of hypoglycaemia (as defined by the study) between those infants who received skin-to-skin contact and those who received conventional incubator care (one study, n = 31; RR 0.24; 95% CI 0.03, 2.06; RD -0.18; 95% CI -0.43, 0.08).

Other secondary outcomes

This study reported the following additional outcome measures: heart rate below 100, or above 180 beats per minute for two consecutive recordings, apnoea longer than 20 s, oxygen saturation below 87% for two consecutive recordings despite supplementation with nasal prong oxygen, FiO₂ up to 0.60 and CPAP up to 5 cm water, and overall data (transfers to NICU, exceeded parameters, mean SCRIP score within first 6 hours and number of perfect scores, mean SCRIP score in the sixth hour and number of perfect scores).

The following secondary outcome measures (as pre-defined at review protocol stage) were not reported for this comparison group: respiratory distress syndrome, surfactant given at any time, intubation in delivery room, requirement for ventilation, duration of ventilation, length of stay, mortality, severe metabolic acidosis, intraventricular haemorrhage, patent ductus arteriosus, chronic lung disease, necrotizing enterocolitis or acute renal failure.

Adverse occurrences

Bergman 2004 reported that “there were no adverse events related to the intervention”.

TRANSWARMER (SODIUM ACETATE) MATTRESS VERSUS ROUTINE CARE (COMPARISON 5):

Primary outcomes

Core body temperature (°C) on admission to NICU or up to two hours after birth (Outcome 5.1):

Brennan 1996 reported core body temperature (axillary °C) on admission to NICU in a study of 24 infants.

Outcome 5.1.1: For infants with a birthweight \leq 1500 g a statistically significant difference in effect was shown for core body temperature on admission to NICU favouring the intervention (transwarmer mattress) group when compared to those who received routine care immediately after birth in the delivery suite (one study, n = 24; MD 1.60 °C; 95% CI 0.83, 2.37).

Hypothermia on admission to NICU (core body temperature < 36.5°C or skin temperature < 36°C) (Outcome 5.2):

Brennan 1996 in addition to reporting core body temperature on admission to NICU as a continuous variable, also provided data

in a dichotomous format in terms of incidence of hypothermia in the intervention and control groups (transwarmer mattress and routine care). Hypothermia was defined in this study as a core body temperature (axillary °C) < 36.5 on admission to NICU.

Outcome 5.2.1: For infants with a birthweight ≤ 1500 g evidence suggests that the transwarmer mattress significantly reduces the risk of hypothermia on admission to NICU (one study, $n = 24$; RR 0.30 95% CI 0.11, 0.83; RD -0.58; 95% CI -0.91, -0.26). Two infants would need to receive a sodium acetate mattress in order to prevent one infant from becoming hypothermic (NNT 2, 95% CI 1 to 4).

Secondary outcomes

The following secondary outcome measures (as pre-defined at review protocol stage) were not reported for this comparison group: hypoglycaemia, respiratory distress syndrome, surfactant given at any time, intubation in delivery room, requirement for ventilation, duration of ventilation, length of stay, mortality, severe metabolic acidosis, intraventricular haemorrhage, patent ductus arteriosus, chronic lung disease, necrotizing enterocolitis, acute renal failure, hyperthermia, burns, maceration, skin or systemic infection, antibiotics, fluid problems or negative psychological outcomes.

Adverse occurrences

Brennan 1996 reported that the intervention did not at any time interfere with the care of the infants.

DISCUSSION

Hypothermia (body temperature below normal) on admission to neonatal units is a problem world-wide across all climates, particularly for small infants and those born too early. Early intervention in the delivery room is vital. This review focused on individual interventions to prevent hypothermia applied immediately at birth apart from 'routine' care in comparison to 'routine care' only. The studies fell into two major groups: barriers to prevent heat losses and additional external heat sources. The strengths of the review lie in the fact that a comprehensive literature search was undertaken encompassing both published and unpublished studies. In addition, strict inclusion criteria were adhered to ensuring that the review was focused on interventions that were applied within the first 10 minutes after birth for preterm and/or low birthweight infants as opposed to bigger infants or long-term thermal management. Limitations of the review were dictated by the number and size of the relevant studies and by the lack of reported data for the pre-specified secondary outcomes.

Seven studies were identified which fulfilled our criteria for inclusion giving a total of 400 infants randomised and 391 infants completing the studies. Five studies compared barriers to heat loss to no barriers (Roberts 1981; Vohra 1999; Vohra 2004a; Knobel 2005; Trevisanuto 2009b) and two studies compared external heat sources (non-routine) to no external heat sources (Brennan 1996;

Bergman 2004). There were three comparisons within the barriers to heat loss category: plastic wraps or bags versus routine care (Vohra 1999; Vohra 2004a; Knobel 2005), plastic cap versus routine care (Trevisanuto 2009b) and stockinet cap versus routine care (Roberts 1981). There were also two comparisons within the external heat source category: skin-to-skin care versus routine care (Bergman 2004) and transwarmer (sodium acetate) mattress versus routine care (Brennan 1996). Each technique was successful to a greater or lesser degree in increasing temperature on admission to NICU.

Barriers to heat loss

Plastic wraps or bags were effective in reducing heat losses for the younger group of infants with gestational ages of < 28 completed weeks. The results showed that there was a high level of heterogeneity within this subgroup of infants. Possible causes may lie within the variations in methods used in the four studies (Vohra 1999; Vohra 2004a; Knobel 2005; Trevisanuto 2009b). The authors of these studies were contacted and potential differences were explored. In both the Vohra 1999 and Vohra 2004a studies the study sample represented all preterm infants within the eligible gestational age range born at the institution. The intervention consisted of a polyethylene bag which was cut into a wrap, only the head of the infant was dried, the body was wrapped from the neck down, a radiant warmer was used, and no parental consent was sought although institutional consent was given. No attempt was made to change standard delivery room temperatures during the studies. Infants were transferred to the NICU in incubators in the first study (Vohra 1999) and in the arms of one of the neonatal team in the second (Vohra 2004a). In the Trevisanuto 2009b study where possible, parental consent was obtained before delivery; the remaining parents received a parent information form after initial stabilisation of their baby. The intervention was also polyethylene bags and the infants were also placed into the bag while still wet up to their necks and only the head was dried and a radiant warmer was used. No attempt was made to change standard delivery room temperatures during the study. Infants were also transferred to the NICU in a transport incubator.

In contrast, in Knobel 2005, the study sample excluded infants for whom no consent was given (and who might have been smaller or more ill and thus more prone to heat loss); and the intervention consisted of polyurethane bag with a drawstring, and the body was wrapped from the neck down while still wet. Attempts to control delivery room temperatures (Knobel 2005), as recommended by the World Health Organisation, failed and resulted in a wide spread of delivery room temperatures during the study period. Infants were transferred to the NICU down a very long, draughty hallway on an open warmer with its heat off with warm blankets over the infants. All four studies followed the NRP protocol and exclusion criteria were similar. An additional point of interest was

the reported imbalance of birthweight between study groups in the [Vohra 1999](#) study. The birthweight-adjusted difference in rectal temperatures for infants with a gestational age of < 28 completed weeks remained significant but decreased from 1.90 °C, SD(0.43 °C) to 1.54 °C, SD(0.42 °C).

There was limited reporting of pre-specified secondary outcomes for this group of studies. There was insufficient evidence to suggest that plastic barriers reduce the risk of death within hospital stay for the younger infants and no deaths were reported for the older group of infants. Similarly, there was no evidence of a significant difference in major brain injury, mean duration of oxygen therapy or hospitalisation, for infants with a gestational age < 29 completed weeks ([Knobel 2005](#)) nor median Apgar score at one and five minutes, first serum glucose concentrations, first blood pH, bicarbonate and hyperthermia for infants < 28 completed weeks gestational age ([Vohra 2004a](#); [Trevisanuto 2009b](#)) mean Apgar score at one minute and five minutes and intubation in delivery room for infants with a gestational age < 29 completed weeks ([Trevisanuto 2009b](#)).

Plastic caps were also effective in reducing heat losses for the younger group of infants with gestational ages of < 29 completed weeks. There was limited reporting of pre-specified secondary outcomes for this comparison group. There was insufficient evidence to suggest that plastic caps reduce the risk of death within hospital stay and there was no significant difference in major brain injury, Apgar score at one minute, first blood pH, bicarbonate and intubation in delivery room. However, a statistically significant difference in effect was shown for Apgar at five minutes favouring the plastic cap group when compared to those who received routine care immediately after birth in the delivery suite.

Where the barrier to heat loss was stockinet caps, a borderline statistically significant difference in temperature on admission to NICU in favour of the intervention group was shown for those infants with a birthweight < 2000 g but no difference for those infants \geq 2000 g. This finding is consistent with those reported by [Greer 1988](#) in which various head coverings were compared under radiant warmers for infants > 2500 g. Where head coverings were applied within one minute of birth, results showed that infants wearing 'stockinettes' had lower mean core body temperatures at 5, 15 and 30 minutes after delivery than either the hatless group or the group wearing an insulated fabric bonnet. As a result, 'stockinettes' were not recommended for use in conjunction with a radiant warmer.

External heat sources (non-routine)

Skin-to-skin care was shown to be effective in reducing the risk of hypothermia when compared to conventional incubator care for infants with a birthweight between 1200 and 2199 g and the transwarmer mattress significantly kept infants \leq 1500 g warmer and reduced the incidence of hypothermia on admission to NICU. There was no evidence that skin-to-skin care reduced the risk of

hypoglycaemia. There was limited reporting of secondary outcomes.

Overall

Despite the variations in interventions applied, definitions of 'routine care', definitions of hypothermia and groups of infants included, across all studies there is a similar pattern emerging showing that infants in the intervention group are significantly warmer (or show a non-significant trend in that direction) when compared to infants receiving 'routine care'. There is also an indication from these studies that the effect is greater in the lightest and most immature infants. Babies of < 28 weeks or those weighing \leq 1500 g appeared to derive most benefit from interventions in the delivery suite to prevent hypothermia. These are also the infants most likely to suffer from the adverse effects of hypothermia and in whom further studies should be undertaken.

There was limited reporting of pre-specified secondary outcomes across all studies. In the [Vohra 1999](#) study, there were no occurrences of adverse events attributable to the intervention (i.e. hyperthermia infection, skin maceration or interference with resuscitation) for infants in the intervention and control groups. [Knobel 2005](#) reported that one participant in the intervention group was hyperthermic with a rectal admission temperature of 38.3°C; the delivery room environmental temperature was 26.7 °C and [Trevisanuto 2009b](#) reported two infants in the plastic wrap group with axillary admission temperatures of 37.6 °C and 38 °C respectively. [Knobel 2005](#) also reported that the clear, pliable polyurethane bag did not interfere with either assessment (visualization, auscultation, palpation) or resuscitative interventions. [Vohra 2004a](#) reported that the wrap procedure was accepted by the neonatal staff and did not interfere with resuscitation in the delivery room. Two infants in the intervention group ([Vohra 2004a](#)) were reported to have a rectal temperature on admission above 37.5 °C. [Bergman 2004](#) reported that "there were no adverse events related to the intervention" and [Brennan 1996](#) stated that the intervention did not at any time interfere with the care of the infants.

[Knobel 2005a](#) in a national survey of newborn intensive care units reported that twenty percent of responding level three neonatal intensive care units used occlusive material instead of drying preterm infants in the delivery room. They also found that only one of the one-hundred and twenty-five responding neonatal intensive care units reported any adverse effects from using wrap or bags in the delivery room. This unit reported that a single infant was hyperthermic on admission to the neonatal intensive care unit but did not report the actual infant temperature.

More recently, the International Liaison Committee on Resuscitation (ILCOR) consensus statement recommends that plastic bags or plastic wrapping under radiant heat be considered as standard techniques to maintain temperature ([ILCOR 2006](#)). National and

international agencies responsible for the development of neonatal resuscitation guidelines have a challenging role when research evidence is emerging, but long-term safety data are not yet available. While the interventions studied in this review offer some short-term benefit regarding heat loss prevention in vulnerable preterm and/or low birthweight infants, their effect on morbidity, mortality and their long-term safety remain unknown. We recommend these additional data are sought, before these interventions are recommended for routine use. Therefore, monitoring (for both benefits and for risks of potential adverse events) should continue in neonatal units where such interventions are adopted as routine practice since wide variation in clinical practice has been reported (Knobel 2005a). In addition, Clarification on 'normal' temperatures for these populations of infants is also essential as is better data correlating axillary versus rectal versus other temperatures.

Paradoxically hypothermia is more of a problem in the developing world where climates are generally warmer (Kumar 2009). Only one of the included studies was conducted in a developing country, South Africa (Bergman 2004). Although South Africa is classified as developing, the units participating in this study carried out in Cape Town provided secondary level care for inborn babies and is therefore not truly representative of the developing world. The plastic wraps utilised in the ongoing Heat Loss Prevention study Vohra 2004 cost approximately 157 Canadian Dollars for 5000 wraps which when calculated per infant is approximately three cents therefore these may be an affordable option for preterm infants in low resource settings.

Further studies are needed of interventions such as plastic bag wrapping for infants under 30 weeks' gestation with adequate sample size to assess both short-term and long-term neurodevelopmental outcomes. Such a multicentre study (HeLP) is currently underway in partnership with the Vermont Oxford Network (n = 1600). The primary study outcome is all cause mortality while secondary outcomes include neurodevelopmental status at 18 months corrected age and safety data. This study will yield important prospective data on 'normal' temperatures in this population, including correlating axillary and rectal temperature.

AUTHORS' CONCLUSIONS

Implications for practice

Health professionals strive to minimise risk of cold stress for newborn infants immediately after birth by adhering to practice guidelines such as: providing a warm delivery room at a minimum of 25 °C (although this is difficult to achieve in practice), drying the infant, removing any wet blankets and wrapping in a prewarmed blanket, prewarming any contact surfaces, avoiding draughts and in developed countries, using radiant warmers or incubators. Despite this hypothermia remains a world-wide problem, especially for small infants and those born too early (Costeloe 2000). Any

additional intervention designed for prevention of hypothermia in very preterm and/or low birthweight infants, and applied within 10 minutes after birth in the delivery suite, compared with routine thermal care may be beneficial in practice. The interventions in this review (plastic wraps and bags, plastic caps, skin-to-skin contact, and transwarmer mattresses) keep infants warmer and lead to higher temperatures on admission to the NICU and to decreased incidence of hypothermia. As hypothermia increases the risk of morbidity and mortality in preterm infants, consideration should be given to using these interventions in the delivery suite. However, given the small numbers of infants and the small numbers of studies in this review and the fact that no long-term data on follow-up are available, firm recommendations for clinical practice cannot be given at this time.

Implications for research

Due to the small numbers and size of trials in this review there is a need to conduct high quality randomised controlled trials where the specifically designed intervention to prevent hypothermia is applied within the first few minutes after birth in the delivery suite. These studies should be large enough to look at long-term neurodevelopmental outcomes and could also include economic evaluations to assess feasibility for use in poorer countries where cost is a fundamental concern. We report in the Table of "Characteristics of ongoing studies" that a large multicentre randomised controlled trial of heat loss prevention (HeLP) in the delivery room commenced in September 2004 (Vohra 2004). This trial will include infants < 28 weeks' gestational age, the intervention will be polyethylene skin wrap and the study will look at mortality as its primary outcome. Eighteen-month neurodevelopmental follow-up is also planned.

ACKNOWLEDGEMENTS

Mr Diarmuid Kennedy, Ms Angela Thompson and Ms Alex McIlroy, Faculty Librarians - Medicine and Health Sciences, Queen's University Medical Library, Mulhouse Building, Belfast, BT12 6DP.

Mr Mike Stevenson, Senior Lecturer in Medical Statistics - Epidemiology & Public Health, Queen's University Belfast, Mulhouse Building, Grosvenor Road, Belfast BT12 6BJ.

Ms Yolanda Montagne, Trials Search Co-ordinator, Cochrane Neonatal Review Group.

Ms Suzanne Sweeney, Texas Woman's University Library, Texas Woman's University, Box 23717, TWU Station, Denton, Texas 76204.

The Cochrane Neonatal Review Group has been funded in part with Federal funds from the Eunice Kennedy Shriver National

Institute of Child Health and Human Development National Institutes of Health, Department of Health and Human Services, USA, under Contract No. HHSN267200603418C.

REFERENCES

References to studies included in this review

- Bergman 2004** {published data only}
Bergman NJ, Linley LL, Fawcus SR. Randomized controlled trial of skin-to-skin contact from birth versus conventional incubator for physiological stabilization in 1200- to 2199-gram newborns. *Acta Paediatrica* 2004;**93**:779–85.
- Brennan 1996** {published data only}
Brennan AB. *Effect of sodium acetate transport mattresses on admission temperatures of infants <= 1500 grams [dissertation]*. Gainesville (FL): University of Florida, 1996.
- Knobel 2005** {published data only}
Knobel RB, Wimmer JE, Holbert D. Heat loss prevention for preterm infants in the delivery room. *Journal of Perinatology* 2005;**25**:304–8.
- Roberts 1981** {published data only}
Roberts JR. *Use of a stockinet cap on premature infants after delivery [dissertation]*. Denton (TX): Texas Woman's University, 1981.
- Trevisanuto 2009b** {published and unpublished data}
Trevisanuto D, Dogioni N, Parlotto M, Micaglio M, Zanardo V. Heat loss prevention in delivery room: a randomized controlled trial of polyethylene caps in very preterm infants. Pediatric Academic Societies' Annual Meeting. May 2-5, 2009, Baltimore (MD). Pediatric Academic Societies, 2009.
- Vohra 1999** {published data only}
Vohra S, Frent G, Campbell V, Abbott M, Whyte R. Effect of polyethylene occlusive skin wrapping on heat loss in very low birth weight infants at delivery: a randomized trial. *Journal of Pediatrics* 1999;**134**:547–51.
- Vohra 2004a** {published data only}
Vohra S, Roberts R, Zhang B, Janes M, Schmidt B. Heat loss prevention (HELP) in the delivery room: a randomized controlled trial of polyethylene occlusive skin wrapping in very preterm infants. *The Journal of Pediatrics* 2004;**145**:750–3.

References to studies excluded from this review

- Ammari 2009** {published data only}
Ammari A, Schulze KF, Ohira-Kist K, Kashyap S, Fifer WP, Myers M, Sahni R. Effects of body position on thermal, cardiorespiratory and metabolic activity in low birthweight infants. *Early Human Development* 2009;**85**:497–501.
- Anderson 2003** {published data only}
Anderson GC, Chiu S H, Dombrowski M A, Swinth J Y, Albert J M, Wada N. Mother-newborn contact in a

randomised trial of kangaroo (skin-to- skin) care. *Journal of Obstetric, Gynecologic, & Neonatal Nursing* 2003;**32**:604–11.

- Andrade 2005** {published data only}
Andrade I. Suckling of the premature newborn child: comparison between the kangaroo with traditional care [Succao do recém-nascido prematuro: comparacao do metodo mae-canguro com os cuidados tradicionais]. *Revista Brasileira de Saude Materno Infantil* 2005;**5**:61–9.
- Baum 1968** {published data only}
Baum JD, Scopes JW. The silver swaddler. Device for preventing hypothermia in the newborn. *Lancet* 1968;**1**:672–3.
- Bell 1983** {published data only}
Bell EF, Rios GR. A double-walled incubator alters partition of body heat loss of premature infants. *Pediatric Research* 1983;**17**:135–40.
- Bergstrom 2005** {published data only}
Bergstrom A, Byaruhanga R, Okong P. The impact of newborn bathing on the prevalence of neonatal hypothermia in Uganda: a randomized, controlled trial. *Acta Paediatrica* 2005;**94**:1462–7.
- Besch 1971** {published data only}
Besch NJ, Perlstein PH, Edwards NK, Keenan WJ, Sutherland JM. The transparent baby bag. A shield against heat loss. *New England Journal of Medicine* 1971;**284**:121–4.
- Bier 1996** {published data only}
Bier JA, Ferguson AE, Morales Y, Liebling JA, Archer D, Oh W, et al. Comparison of skin-to-skin contact with standard contact in low-birth-weight infants who are breast-fed. *Archives of Pediatrics & Adolescent Medicine* 1996;**150**:1265–9.
- Boo 2007** {published data only}
Boo N Y, Chew E L. A randomised control trial of clingfilm for prevention of hypothermia in term infants during phototherapy. *Singapore Medical Journal* 2007;**47**:757–62.
- Brice 1981** {published data only}
Brice JEH, Rutter N, Hull D. Reduction of skin water loss in the newborn II. Clinical trial of two methods in very low birthweight babies. *Archives of Disease in Childhood* 1981;**56**:673–5.
- Bystrova 2003** {published data only}
Bystrova K, Widstrom AM, Matthiesen AS, Ransjo-Arvidson, Wells-Nystrom B, Wassberg C, et al. Skin-to-skin contact may reduce negative consequences of “the stress of being born”: a study on temperature in newborn infants, subjected to different ward routines in St. Petersburg. *Acta Paediatrica* 2003;**92**:320–6.

- Bystrova 2007** *{published data only}*
Bystrova K, Matthiesen AS, Vorontsov I, Widstrom AM, Ransjo-Arvidson AB, Uvnas-Moberg K. Maternal axillar and breast temperature after giving birth: effects of delivery ward practices and relation to infant temperature. *Birth* 2007;**34**:291–300.
- Carfoot 2005** *{published data only}*
Carfoot S, Williamson P, Dickson R. A randomised controlled trial in the North of England examining the effects of skin-to-skin care on breast feeding. *Midwifery* 2005;**21**:71–9.
- Cattaneo 1998** *{published data only}*
Cattaneo A, Davanzo R, Worku B, Surjono A, Escheverria M, Bedri A, et al. Kangaroo mother care for low birthweight infants: a randomized controlled trial in different settings. *Acta Paediatrica* 1998;**87**:976–85.
- Chaput 1979** *{published data only}*
Chaput de Saintonge DM, Cross KW, Hathorn MK, Lewis SR, Stothers JK. Hats for the newborn infant. *British Medical Journal* 1979;**2**:570–1.
- Charpak 1997** *{published data only}*
Charpak N, Ruiz-Pelaez JG, Figueroa de CZ, Charpak Y. Kangaroo mother versus traditional care for newborn infants <= 2000 grams: a randomized, controlled trial. *Pediatrics* 1997;**100**:682–8.
- Cheah 2000** *{published data only}*
Cheah FC, Boo NY. Risk factors associated with neonatal hypothermia during cleaning of newborn infants in labour rooms. *Journal of Tropical Pediatrics* 2000;**46**:46–50.
- Christensson 1992** *{published data only}*
Christensson K, Siles C, Moreno L, Belaustequi A, De La Fuente P, Lagercrantz H, et al. Temperature, metabolic adaptation and crying in healthy full-term newborns cared for skin-to-skin or in a cot. *Acta Paediatrica* 1992;**81**:488–93.
- Christensson 1995** *{published data only}*
Christensson K, Cabrera T, Christensson E, Uvnas Moberg K, Winberg J. Separation distress call in the human neonate in the absence of maternal body contact. *Acta Paediatrica* 1995;**84**:468–473.
- Christensson 1996** *{published data only}*
Christensson K. Fathers can effectively achieve heat conservation in healthy newborn infants. *Acta Paediatrica* 1996;**85**:1354–60.
- Christensson 1998** *{published data only}*
Christensson K, Bhat GJ, Amadi BC, Eriksson B, Hojer B. Randomised study of skin-to-skin versus incubator care for rewarming low-risk hypothermic neonates. *Lancet* 1998;**352**:1115.
- Chwo 1999** *{published data only}*
Chwo MJ. *Early kangaroo care for 34-36 week preterm infants: effects on temperature, weight, behavior, and acuity [dissertation]*. Cleveland(OH): Case Western Reserve University, 1999.
- Chwo 2002** *{published data only}*
Chwo MJ, Anderson GC, Good M, Dowling DA, Siau SH, Cho DM. A randomized controlled trial of early kangaroo care for preterm infants: effects on temperature, weight, behaviour and acuity. *Journal of Nursing Research* 2002;**10**:129–42.
- Coles 1979** *{published data only}*
Coles EC, Valman HB. Hats for the newborn infant. *British Medical Journal* 1979;**2**:734–5.
- Dahm 1972** *{published data only}*
Dahm LS, James LS. Newborn temperature and calculated heat loss in the delivery room. *Pediatrics* 1972;**49**:504–13.
- Daniel 2004** *{published data only}*
Daniel LM. Prevention of neonatal hypothermia. Proceedings of the 13th congress of the Federation-of-Asia-and-Oceania-Perinatal-Societies (FAOPS 2004) APR 14–18, 2004 (Kuala Lumpur, Malaysia). 2004.
- Dannaway 2009** *{published data only}*
Dannaway D, Simon PL. Warming mattress versus polyethylene wrapping to prevent hypothermia in preterm newborns. ClinicalTrials.gov identifier:NCT00603837. ClinicalTrials.gov. U.S. National Institute of Health, 12 October 2009.
- Darmstadt 2007** *{published data only}*
Darmstadt GL, Kumar V, Yadav R, Singh V, Singh P, Mohanty S, et al. Introduction of community-based skin-to-skin care in rural Uttar Pradesh, India. *Journal of Perinatology* 2006;**26**:597–604.
- Darmstadt 2008** *{published data only}*
Darmstadt GL, Saha SK, Ahmed AS, Ahmed S, Chowdhury MAK, Law PA, et al. Effect of skin barrier therapy on neonatal mortality rates in preterm infants Bangladesh: a randomized, controlled, clinical trial. *Pediatrics* 2008;**121**:522–9.
- Day 1964** *{published data only}*
Day LD, Caliguiri L, Kamenski C, Ehrlich F. Body temperature and survival of premature infants. *Pediatrics* 1964;**34**:171–81.
- Dodd 2003** *{published data only}*
Dodd VL. *Effect of kangaroo care in preterm infants [dissertation]*. Storrs (CT): The University of Connecticut, 2003.
- Erlandsson 2007** *{published data only}*
Erlandsson K, Dsilna A, Fagerberg I, Christensson K. Skin-to-skin care with the father after cesarean birth and its effect on newborn crying and prefeeding behavior. *Birth* 2007;**34**:105–14.
- Fallis 2006** *{published data only}*
Fallis WM, Hamelin K, Symonds J, Wang X. Maternal and newborn outcomes related to maternal warming during cesarean delivery. *Journal of Obstetric, Gynecologic and Neonatal Nursing* 2006;**35**:324–31.
- Fardig 1980** *{published data only}*
Fardig JA. A comparison of skin-to-skin contact and radiant heaters in promoting neonatal thermoregulation. *Journal of Nurse-Midwifery* 1980;**25**:19–28.

- Ferber 2004** *{published data only}*
Ferber SG, Makhoul IR. The effect of skin-to-skin contact (kangaroo care) shortly after birth on the neurobehavioral responses of the term newborn: a randomized, controlled trial. *Pediatrics* 2004;**113**:858–65.
- Gray 2004** *{published data only}*
Gray PH, Paterson S, Finch G, Hayes M. Cot-nursing using a heated, water-filled mattress and incubator care: a randomized clinical trial. *Acta Paediatrica* 2004;**93**:350–5.
- Green-Abate 1994** *{published data only}*
Green-Abate C, Tafari N, Rao MR, Yu K, Clemens JD. Comparison of heated water-filled mattress and space-heated room with infant incubator in providing warmth to low birthweight newborns. *International Journal of Epidemiology* 1994;**23**:1226–33.
- Greer 1988** *{published data only}*
Greer PS. Head coverings for newborns under radiant warmers. *Journal of Obstetric, Gynecological and Neonatal Nursing* 1988;**17**:265–71.
- Grover 1994** *{published data only}*
Grover G, Berkowitz CD, Lewis RJ, Thompson M, Berry L, Seidel J. The effects of bundling on infant temperature. *Pediatrics* 1994;**94**:669–73.
- Gulezian 1980** *{published data only}*
Gulezian GZ. *Effect of skin-to-skin contact on transitional newborns infants' temperature [dissertation]*. Chicago (IL): University of Illinois at the Medical Center, 1980.
- Harrison 2004** *{published data only}*
Harrison M. *Kangaroo mother care: an innovative way of care for infants [dissertation]*. Hattiesburg (MS): University of Southern Mississippi, 2004.
- Hellin Martinez 2000** *{published data only}*
Heillin Martinez MJ, Lopez Terol E, Perez Lopez M, Vidagany Escrig I. Maintain the temperature of a healthy newborn infant. *Revista Rol de Enfermeria* 2000;**23**:63–5.
- Hobbs 1975** *{published data only}*
Hobbs JF, MacKuanying N, Eidelman I, Schneider KM. A new transparent insulating infant gown to maintain thermal stability. *Pediatric Research* 1975;**9**:366 [Abstract 658].
- Holzman 1985** *{published data only}*
Holzman IR. A method to maintain infant temperature. *American Journal of Diseases of Children* 1985;**139**:390–2.
- Horn 2002** *{published data only}*
Horn EP, Schroeder F, Gottschalk A, Sessler DI, Hiltmeyer N, Standl T, et al. Active warming during cesarean delivery. *Anesthesia and Analgesia* 2002;**94**:409–14.
- Huang 2002** *{published data only}*
Huang L, Chwo M, Chu D, Chang Y. Effects of very early kangaroo care on infants' extrauterine adaptation. *Journal of Nursing (China)* 2002;**49**:37–51.
- Huang 2006** *{published data only}*
Huang YY, Huang CY, Lin SM, Wu SC. Effect of very early kangaroo care on extrauterine temperature adaptation in newborn infants with hypothermia problems. *Hu Li Tsa Chih - Journal of Nursing* 2006;**53**:41–8.
- Johanson 1992** *{published data only}*
Johanson RB, Spencer SA, Rolfe P, Jones P, Malla DS. Effect of post-delivery care on neonatal body temperature. *Acta Paediatrica* 1992;**81**:859–63.
- Kadam 2005** *{published data only}*
Kadam S, Binoy S, Kanbur W, Mondkar JA, Fernandez A. Feasibility of kangaroo mother care in Mumbai. *The Indian Journal of Pediatrics* 2005;**72**:35–8.
- Kaushal 2005** *{published data only}*
Kaushal M, Agarwal R, Aggarwal R, Singal A, Upadhyay M, Sirmivas V, et al. Cling wrap, an innovative intervention for temperature maintenance and reduction of insensible water loss in very low-birthweight babies nursed under radiant warmers: a randomized, controlled trial. *Annals of Tropical Paediatrics* 2005;**25**:111–8.
- Kumar 2008** *{published data only}*
Kumar V, Mohanty S, Kumar A, Misra RP, Santosham S, Baqui AH, et al. The Saksham Study Group. Effect of community-based behaviour change management on neonatal mortality in Shivgarh, Uttar Pradesh, India: a cluster-randomised controlled trial. *The Lancet* 2008;**372**:1151–1162.
- Legault 1993** *{published data only}*
Legault M, Goulet C. Comparative study of two methods of holding premature infants: the kangaroo method versus traditional method [Etude comparative de deux methodes de sortie du premature: methode kangourou versus methode traditionnelle]. *Canadian Journal of Nursing Research* 1993;**25**:67–80.
- Ludington-Hoe 1994** *{published data only}*
Ludington-Hoe S, Thompson C, Swinth J, Hadeed AJ, Anderson GC. Kangaroo care: research results and practice implications and guidelines. *Neonatal Network* 1994;**13**:19–27.
- Ludington-Hoe 2000** *{published data only}*
Ludington-Hoe SM, Nguyen N, Swinth JY, Satyshur RD. Kangaroo care compared to incubators in maintaining body warmth in preterm infants. *Biological Research for Nursing* 2000;**2**:60–73.
- Ludington-Hoe 2004** *{published data only}*
Ludington-Hoe SM, Anderson GC, Swinth JY, Thompson C, Hadeed AJ. Randomized controlled trial of kangaroo care: cardiorespiratory and thermal effects on healthy preterm infants. *Neonatal Network* 2004;**23**:39–48.
- Marks 1985** *{published data only}*
Marks KH, Devenyi AG, Bello ME, Nardis EE, Seaton JF, Ultman JS. Thermal head wrap for infants. *Journal of Pediatrics* 1985;**107**:956–9.
- Mathew 2008** *{published data only}*
Mathew B, Lakshminrusimha S, Carrion V. Vinyl bag versus thermal mattress to prevent hypothermia in extremely low birthweight (ELBW) infants. 2008 Pediatric Academic Societies' Annual Meeting. May 2–6 2008, Honolulu, Hawaii. Pediatric Academic Societies, 2008.

- Mazurek 1999** *{published data only}*
Mazurek T, Mikiel-Kostyra K, Mazur J, Wieczorek P, Radwanska B, Pachuta-Weigier L. Influence of immediate newborn care on infant adaptation to the environment [Wplyw postepowania z noworodkiem bezposrednio po porodzie na cechy jego adaptacji do srodowiska]. *Medycyna Wieku Rozwojowego* 1999;**3**:215–24.
- Medves 2004** *{published data only}*
Medves JM, O'Brien B. Effect of bather and location of first bath on maintaining thermal stability in newborns. *Journal of Obstetric, Gynecologic and Neonatal Nursing* 2004;**33**: 175–82.
- Meyer 2001** *{published data only}*
Meyer MP, Payton MJ, Salmon A, Hutchinson C, de Klerk A. A clinical comparison of radiant warmer and incubator care for preterm infants from birth to 1800 grams. *Pediatrics* 2001;**108**:395–401.
- Meyer 2007** *{published data only}*
Meyer MP, Bold GT. Admission temperatures following radiant warmer or incubator transport for preterm infants < 28 weeks: a randomised study. *Archives of Disease in Childhood Fetal and Neonatal Edition* 2007;**92**:F295–7.
- Miles 2006** *{published data only}*
Miles R, Cowan F, Glover V, Stevenson J, Modi N. A controlled trial of skin-to-skin contact in extremely preterm infants. *Early Human Development* 2006;**82**:447–55.
- Monterosso 1999** *{published data only}*
Monterosso L, Percival P, Cole J, Evans SF. Effect of nappy liners on temperature stability in very preterm infants. *Journal of Paediatrics and Child Health* 1999;**35**:363–6.
- Moore 2007** *{published data only}*
Moore ER, Anderson GC. Randomized controlled trial of very early mother-infant skin-to-skin contact and breastfeeding status. *Journal of Midwifery & Women's Health* 2007;**52**:116–25.
- Nopper 1996** *{published data only}*
Nopper AJ, Horri KA, Sookdeo-Drost S, Wang TH, Mancini AJ, Lane AT. Topical ointment therapy benefits premature infants. *Journal of Pediatrics* 1996;**128**:660–9.
- Nuntnarumit 2004** *{published data only}*
Nuntnarumit P, Deesomchok A. Efficacy of polyethylene occlusive skin wrapping with plastic bag in preventing hypothermia in infants less than 32 weeks' gestational age at delivery. 2004 Pediatric Academic Societies' Annual Meeting, May 1-4, 2004, San Francisco, (CA). Pediatric Academic Societies, 2004.
- Omene 1978** *{published data only}*
Omene JA, Diejomaoh FM, Faal M, Diakparomre MA, Obiaya M. Heat loss in Nigerian newborn infants in the delivery room. *International Journal of Gynaecology and Obstetrics* 1978–79;**16**:300–2.
- Pattinson 2005** *{published data only}*
Pattinson RC, Arsalo I, Bergh AM, Malan AF, Patrick M, Phillips N. Implementation of kangaroo mother care: a randomized trial of two outreach strategies. *Acta Paediatrica* 2005;**94**:924–7.
- Punthmatharith 2001** *{published data only}*
Punthmatharith B. *Randomized controlled trial of early kangaroo (skin-to-skin) care: effects on maternal feelings, maternal-infant interaction and breastfeeding success in Thailand [dissertation]*. Cleveland (OH): Case Western Reserve University, 2001.
- Raman 1992** *{published data only}*
Raman S, Shahla A. Temperature drop in normal term newborn infants born at the University Hospital, Kuala Lumpur. *Australian and New Zealand Journal of Obstetrics and Gynaecology* 1992;**32**:117–9.
- Ramanathan 2001** *{published data only}*
Ramanathan K. Kangaroo mother care in very low birth weight infants. *Indian Journal of Pediatrics* 2001;**68**: 1019–23.
- Rao 2008** *{published data only}*
Rao S, Udani R, Nanavanti R. Kangaroo mother care low birth weight infants: a randomized controlled trial. *Indian Pediatrics* 2008;**45**:17–23.
- Roberts 2000** *{published data only}*
Roberts KL, Paynter C, McEwan B. A comparison of kangaroo mother care and conventional cuddling care. *Neonatal Network* 2000;**19**:31–5.
- Rojas 2001** *{published data only}*
Rojas MA, Kaplan M, Mayes L, Quevedo ME, Foster LB, Sherwonit E, et al. Traditional holding (TH) and skin-to-skin care (SSC) for newborn infants <= 1500 grams. A randomized controlled trial. *Pediatric Research* 2001;**49**: 360A.
- Ruiz 1998** *{published data only}*
Ruiz JG. Kangaroo mother versus 'traditional' care for newborn infants <= 2000 grams: a randomised control trial. *Journal of Clinical Epidemiology* 1998;**51** Suppl:125.
- Sankaranarayanan 2005** *{published data only}*
Sankaranarayanan K, Mondkar JA, Chauhan MM, Mascarenhas BM, Mainkar AR, Salvi RY. Oil massage in neonates: an open randomized controlled study of coconut versus mineral oil. *Indian Pediatrics* 2005;**42**:877–84.
- Sarman 1989** *{published data only}*
Sarman I, Tunell R. Providing warmth for preterm babies by a heated, water filled mattress. *Archives of Disease in Childhood Fetal and Neonatal Edition* 1989;**64**:29–33.
- Sarman 1992** *{published data only}*
Sarman I. Thermal responses and heart rates of low-birth-weight premature babies during daily care on a heated, water-filled mattress. *Acta Paediatrica* 1992;**81**:15–20.
- Short 1998** *{published data only}*
Short MA. A comparison of temperature in VLBW infants swaddled versus unswaddled in a double-walled incubator in skin control mode. *Neonatal Network* 1998;**17**:25–31.
- Sloan 1994** *{published data only}*
Sloan NL, Camacho WL, Rojas EP, Stern C, and Maternidad Isidro Ayora Study Team. Kangaroo mother method: randomised controlled trial of an alternative method of care for stabilised low-birthweight infants. *The Lancet* 1994;**344**:782–5.

Syfrett 1993 {published data only}

Syfrett BE. *Very early and virtually continuous kangaroo care for 34 to 36 week gestation preterm infants [dissertation]*. Tallahassee (FL): University of Florida, 1993.

Trevisanuto 2009a {published data only}

Trevisanuto D, Fasson A, Doglioni N, Biason S, Micaglio M, Zanardo V. A new device (cicogna) for transferring the neonate in the delivery room setting: a randomised, controlled trial. *The Journal of Maternal-Fetal and Neonatal Medicine* 2009;**22**:148–51.

Tsogt 2005 {published data only}

Tsogt B, Maniseki-Holland S, Pollock J, Blair P, Fleming P. The development of thermoregulation in a harsh environment: a prospective controlled study of the effects of swaddling on infants' thermal balance in a Mongolian winter. 2005 The Neonatal Societies Meeting Bristol, 30th June - 1st July 2005. 2005.

Vaidya 2005 {published data only}

Vaidya K, Sharma A, Dhungel S. Effect of early mother-baby close contact over the duration of exclusive breastfeeding. *Nepal Medical College Journal* 2005;**7**:138–40.

van den Bosch 1990 {published data only}

van den Bosch CA, Bullough CH. Effect of early suckling on term neonates' core body temperature. *Annals of Tropical Paediatrics* 1990;**10**:347–53.

van den Bosch 1996 {published data only}

van den Bosch CA, Nhlane C, Kazembe P. Trial of polythene tobacco-wrap in prevention of hypothermia in neonates less than 1500 grams. *Tropical Doctor* 1996;**26**:26–8.

Villalon 1992 {published data only}

Villalon HU, Alvarez PC, Barria EH, Caneleo DH, Carrillo LM, Duran SG. Effect of early skin-to-skin contact on temperature regulation, heart rate, and respiratory rate in healthy, full term newborns [Contacto percoz piel a piel: efecto sobre los parametros fisiologicos en las cuatro horas posteriores al parto en recién nacidos de termino sanos]. *Revista Chilena de Pediatria* 1992;**63**:140–4.

Worku 2005 {published data only}

Worku B, Kassie A. Kangaroo mother care: a randomized controlled trial on effectiveness of early kangaroo mother care for the low birthweight infants in Addis Ababa, Ethiopia. *Journal of Tropical Pediatrics* 2005;**51**:93–7.

Yeh 1980 {published data only}

Yeh TF, Vohra S, Lillien LD, Matwynshym J, Srinivasan G, Pildes RS. Oxygen consumption and insensible water loss in premature infants in single versus double-walled incubators. *Journal of Pediatrics* 1980;**97**:967–71.

Yokoyama 2009 {published data only}

Yokoyama K, Suzuki M, Shimada Y, Matsushima T, Bito H, Sakamoto A. Effect of administration of pre-warmed intravenous fluids on the frequency of hypothermia following spinal anesthesia for Cesarean delivery. *Journal of Clinical Anesthesia* 2009;**21**:242–8.

References to studies awaiting assessment**Punnahitananda 2008 {published data only}**

Punnahitananda S, Thaithumyanon P. Efficacy of polyethylene plastic wrap for the prevention of hypothermia during immediate postnatal period in preterm infants: a randomized controlled trial. 2008 Pediatric Academic Societies' Meeting. May 2–6, 2008, Honolulu, Hawaii. Pediatric Academic Societies, 2008.

References to ongoing studies**Vohra 2004 {unpublished data only}**

Vohra S. Multi centered randomised controlled trial of heat loss prevention (HeLP) in delivery room. Personal communication 08 July 2004 and September 2007.

Additional references**Adamsons 1965**

Adamsons K, Gandy GM, James LS. The influence of thermal factors upon oxygen consumption of the newborn human infant. *Journal of Pediatrics* 1965;**66**:495–508.

Adamsons 1965a

Adamsons K, Towell M E. Thermal homeostasis in the fetus and newborn. *Anesthesiology* 1965;**26**:531–48.

Almeida 2009

Almedia PG, Chandley J, Davis J, Harrigan RC. Use of heated gel mattress and its impact on admission temperature of very low birth-weight infants. *Advances in Neonatal Care* 2009;**9**:34–9.

Bailey 2000

Bailey J, Rose P. Temperature measurement in the preterm infant: A literature review. *Journal of Neonatal Nursing* 2000;**6**:28–32.

Baumgart 1981

Baumgart S, Engle WD, Fox WW, Polin RA. Effect of heat shielding on convective and evaporative heat losses and on radiant heat transfer in the premature infant. *Journal of Pediatrics* 1981;**99**:948–56.

Bell 1978

Bell MJ, Ternberg JL, Feigin RD, Keating JP, Marshall R, Barton L, et al. Neonatal necrotizing enterocolitis. Therapeutic decisions based on clinical staging. *Annals of Surgery* 1978;**187**:1–7.

Bell 1980

Bell EF, Weinstein MR, Oh W. Heat balance in premature infants: comparative effects of convectively heated incubator and radiant warmer, with and without plastic heat shield. *Journal of Pediatrics* 1980;**96**:460–5.

Bickmann 1992

Buczowski-Bickmann MK. Thermoregulation in the neonate and the consequences of hypothermia. *CRNA: The Clinical Forum for Nurse Anesthetists* 1992;**3**:77–82.

Bloom 1994

Bloom RS, Cropley C, the AHA/AAP Neonatal Resuscitation Steering Committee, editors. *Textbook of neonatal resuscitation*. 3rd Edition. Dallas (TX): American Heart Association, 1994:2–9.

Borse 1997

Borse N, Deodhar J, Pandit AN. Effects of thermal environment on neonatal thermoregulation. *Indian Pediatrics* 1997;**34**:718–20.

Boutron 2008

Boutron I, Moher D, Altman DG, Schultz KF, Ravard P, for the CONSORT Group. Extending the CONSORT statement to randomized trials of nonpharmacologic treatment: explanation and elaboration. *Annals of Internal Medicine* 2008;**148**:295–309.

Britton 1980

Britton GR. Early mother-infant contact and infant temperature stabilization. *JOGN Nursing* 1980;**9**:84–6.

Bruck 1961

Bruck K. Temperature regulation in the newborn infant. *Biology of the Neonate* 1961;**3**:65–119.

Brun 1997

Brun C, Stokvad P, Alsbjorn BF. Burn wounds after resuscitation of a newborn girl [Brandsar efter genoplivning af en nyfodt pige]. *Ugestrift for Læger* 1997;**159**:6531–2.

Capobianco 1980

Capobianco JA. Keeping the newborn warm: how to safeguard the infant against life-threatening heat loss. *Nursing* 1980;**10**:64–7.

CESDI 2003

CESDI (Confidential Enquiry into Stillbirths, Deaths in Infancy). *PROJECT 27/28. An enquiry into quality of care and its effect on the survival of babies born at 27–28 weeks.* London, United Kingdom: Maternal and Child Health Consortium, 2003.

Chadd 1972

Chadd MA, Gray OP. Hypothermia and coagulation defects in the newborn. *Archives of Disease in Childhood* 1972;**47**: 819–21.

Christensson 1988

Christensson K, Ransjo-Arvidson AB, Kakoma C, Lungu F, Darkwah G, Chikamata D, et al. Midwifery care routines and prevention of heat loss in the newborn: a study in Zambia. *Journal of Tropical Pediatrics* 1988;**34**:208–12.

Clarke 2002

Clarke M, Oxman AD, editors. *Cochrane Reviewers' Handbook 4.1.5 [updated April 2002].* Cochrane Database of Systematic Reviews. Oxford: Update Software, 2, issue 2002.

Costeloe 2000

Costeloe K, Hennessy E, Gibson AT, Marlow N, Wilkinson AR. The EPICure study: outcomes to discharge from hospital for infants born at the threshold of viability. *Pediatrics* 2000;**106**:659–71.

Davis 1980

Davis V. The structure and function of brown adipose tissue in the neonate. *Journal of Obstetric, Gynecologic & Neonatal Nursing* 1980;**9**:368–72.

Du 1969

Du JN, Oliver TK Jr. The baby in the delivery room. A suitable microenvironment. *Journal of the American Medical Association* 1969;**207**:1502–4.

Elliott 1957

Elliott RI, Mann TP. Neonatal cold injury due to accidental exposure to cold. *Lancet* 1957;**272**:229–34.

Gandy 1964

Gandy GM, Adamsons K, Cunningham N, Silverman WA, James LS. Thermal environment and acid-base homeostasis in human infants during the first few hours of life. *Journal of Clinical Investigation* 1964;**43**:751–8.

Glass 1968

Glass L, Silverman WA, Sinclair JC. Effects of the thermal environment on cold resistance and growth of small infants after the first week of life. *Pediatrics* 1968;**41**:1033–46.

Hammarlund 1979

Hammarlund K, Sedin G. Transepidermal water loss in newborn infants. III. Relation to gestational age. *Acta Paediatrica Scandinavica* 1979;**68**:795–801.

Hammarlund 1980

Hammarlund K, Nilsson GE, Oberg PA, Sedin G. Transepidermal water loss in newborn infants. V. Evaporation from the skin and heat exchange during the first hours of life. *Acta Paediatrica Scandinavica* 1980;**69**: 385–92.

Harned 1970

Harned HS Jr, Herrington RT, Ferreiro JI. The effects of immersion and temperature on respiration in newborn lambs. *Pediatrics* 1970;**45**:598–605.

Hey 1970

Hey EN, Katz G. The optimal thermal environment for naked babies. *Archives of Disease in Childhood* 1970;**45**: 328–34.

Hey 1975

Hey E. Thermal neutrality. *British Medical Bulletin* 1975; **31**:69–74.

Ho 2007

Ho TQ, Lee HC, Rhine WD. The impact of a quality improvement project to reduce hypothermia in premature newborns. 2007 Pediatric Academic Societies' Annual Meeting. May 5–8, 2007, Toronto, Canada. Pediatric Academic Societies, 2007.

ILCOR 2006

The International Liaison Committee on Resuscitation. The International Liaison Committee on Resuscitation (ILCOR) consensus on science with treatment recommendations for pediatric and neonatal patients: neonatal resuscitation. *Pediatrics* 2006;**117**:e978–88.

Kaplan 2009

Kaplan HC, Breig R, Meek-Silvers D, Hallett-Voss C, Corcoran N, Falciglia H, et al. Intervention to reduce delivery room cold stress in very low birthweight infants. Pediatric Academic Societies' Annual Meeting. May 2–5, 2009, Baltimore (MD). Pediatric Academic Societies, 2009.

Knobel 2005a

Knobel RB, Vohra S, Lehmann CU. Heat loss prevention in the delivery room for preterm infants: a national survey of newborn intensive care units. *Journal of Perinatology* 2005; **25**:514–8.

Kumar 2009

Kumar V, Shearer JC, Kumar A, Darmstadt GL. Neonatal hypothermia in low resource settings: a review. *Journal of Perinatology* 2009; **29**:401–12.

Lang 2004

Lang N, Bromiker R, Arad I. The effect of wool versus cotton head covering and length of stay with mother following delivery on infant temperature. *International Journal of Nursing Studies* 2004; **41**:843–6.

Laptook 2007

Laptook A, Salhab W, Bhaskar B and the Neonatal Research Network. Admission temperature of low birth weight infants: predictors and associated morbidities. *Pediatrics* 2007; **119**:643–9.

Laptook 2008

Laptook AR, Watinson M. Temperature management in the delivery room. *Seminars in Fetal and Neonatal Medicine* 2008; **13**:383–91.

LeBlanc 1991

LeBlanc MH. Thermoregulation: incubators, radiant warmers, artificial skins and body hoods. *Clinics in Perinatology* 1991; **18**:403–22.

Lee 2000

Lee SK, McMillan DD, Ohlsson A, Pendray M, Synnes A, Whyte R et al and the Canadian NICU Network. Variations in practice and outcomes in the Canadian NICU Network 1996-1997. *Pediatrics* 2000; **106**:1070–9.

Lenclen 2002

Lenclen R, Mazraani M, Jugie M, Couderc S, Hoenn C, Carbajal R, et al. Use of a polyethylene bag: a way to improve the thermal environment of the premature newborn at the delivery room [Utilisation d'un sac en polyethylene: un moyen d'ameliorer l'environnement thermique du premature en salle de naissance]. *Archives de Pediatrie* 2002; **9**:238–44.

Lyon 2006

Lyon A. Applied physiology: temperature control in the newborn infant. *Current Paediatrics* 2006; **16**:386–92.

Nayeri 2006

Nayeri F, Nili F. Hypothermia at birth and its associated complications in newborns: a follow up study. *Iranian Journal of Public Health* 2006; **35**:48–52.

Newton 2003

Newton T, Watkinson M. Preventing hypothermia at birth in preterm babies: at a cost of overheating some?. *Archives of Disease in Childhood Fetal and Neonatal Edition* 2003; **88**: F256.

Niermeyer 2000

Niermeyer S, Kattwinkel J, Van Reempts P, Nadkarni V, Phillips B, Zideman D, et al. International Guidelines for

Neonatal Resuscitation: an excerpt from the Guidelines 2000 for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care: International Consensus on Science. Contributors and Reviewers for the Neonatal Resuscitation Guidelines. *Pediatrics* 2000; **106**:E29.

Oliver 1965

Oliver TK Jr. Temperature regulation and heat production in the newborn. *Pediatric Clinics of North America* 1965; **12**: 765–79.

Papile 1978

Papile LA, Burstein J, Burstein R, Koffler H. Incidence and evolution of subependymal and intraventricular hemorrhage: a study of infants with birth weights less than 1500 gm. *Journal of Pediatrics* 1978; **92**:529–34.

Pomerance 1974

Pomerance JJ, Madore C. Effect of temperature on survival of infants with RDS. *Pediatric Research* 1974; **8**:449 [Abstract].

Rutter 2000

Rutter N. Clinical consequences of an immature barrier. *Seminars in Neonatology* 2000; **5**:281–7.

Scopes 1963

Scopes JW, Tizard JP. The effect of intravenous noradrenaline on the oxygen consumption of new-born mammals. *Journal of Physiology* 1963; **165**:305–26.

Shennan 1988

Shennan AT, Dunn MS, Ohlsson A, Lennox K, Hoskins EM. Abnormal pulmonary outcomes in preterm infants: prediction from oxygen requirement in the neonatal period. *Pediatrics* 1988; **82**:527–32.

Silverman 1958

Silverman WA, Fertig JW, Berger AP. The influence of the thermal environment upon the survival of newly born premature infants. *Pediatrics* 1958; **22**:876–86.

Smales 1978

Smales ORC, Kime R. Thermoregulation in babies immediately after birth. *Archives of Disease in Childhood* 1978; **53**:58–61.

Smith 2004

Smith LS. Temperature monitoring in newborns: A comparison of thermometry and measurement sites. *Journal of Neonatal Nursing* 2004; **10**:157–65.

Soll 2008

Soll RF. Heat loss prevention in neonates. *Journal of Perinatology* 2008; **28**:S57–S59.

Stapleton 1987

Stapleton FB, Jones DP, Green RS. Acute renal failure in neonates: incidence, etiology and outcome. *Pediatric Nephrology* 1987; **1**:314–20.

Stephenson 1970

Stephenson JM, Du JN, Oliver TK Jr. The effect of cooling on blood gas tensions in newborn infants. *Journal of Pediatrics* 1970; **76**:848–52.

Stern 1970

Stern L. The newborn infant and his thermal environment. *Current Problems in Pediatrics* 1970;**1**:1–29.

Tafari 1973

Tafari N, Olsson E. Neonatal cold injury in the tropics. *Ethiopian Medical Journal* 1973;**11**:57–65.

Thomas 1994

Thomas K. Thermoregulation in neonates. *Neonatal Network* 1994;**13**:15–25.

UNICEF 2002

The United Nations Children's Fund (UNICEF). *The State of the World's Children Report 2003 - Child Participation*. New York: UNICEF, 2002.

Watkinson 2006

Watkinson M. Temperature control of premature infants in the delivery room. *Clinics in Perinatology* 2006;**33**:43–53.

WHO 1997

Department of Reproductive Health and Research (RHR), World Health Organisation. *Thermal protection of the*

newborn: a practical guide (WHO/RHT/MSM/97.2). Geneva: World Health Organisation, 1997.

References to other published versions of this review**McCall 2005**

McCall EM, Alderdice FA, Halliday HL, Jenkins JG, Vohra S. Interventions to prevent hypothermia at birth in preterm and/or low birthweight babies. *Cochrane Database of Systematic Reviews* 2005, Issue 1. [DOI: 10.1002/14651858.CD004210.pub2]

McCall 2008

McCall EM, Alderdice FA, Halliday HL, Jenkins JG, Vohra S. Interventions to prevent hypothermia at birth in preterm and/or low birthweight infants. *Cochrane Database of Systematic Reviews* 2008, Issue 1. [DOI: 10.1002/14651858.CD004210.pub3]

* Indicates the major publication for the study

CHARACTERISTICS OF STUDIES

Characteristics of included studies *[ordered by study ID]*

Bergman 2004

Methods	<p>Randomised. Multicentre (2). Blinding of randomisation: unclear. Generation of allocation sequence: adequate, (computer minimisation method). Allocation concealment: unclear, (“... a computer minimisation method determined allocation in a concealed manner” , “.. second hospital... minimisation factors and subsequent allocation by the computer were exchanged by mobile telephone, with no delays”) . Blinding of intervention: participants: no/unfeasible, care givers: no/unfeasible. Blinding of outcome measurement: yes, after the pilot phase, the principal investigator and data analyses team were blinded. A statistician, blinded to the allocation, performed the statistical analyses. Complete follow-up: no, (31 of 35 infants randomised completed the study, one infant was excluded after 2 hours when the mother required a caesarean section for a second twin, three further infants 'bailed out' within the first 3 hr requiring additional respiratory support and moved to neonatal intensive care) Room temperature, maternal temperature and incubator temperature were recorded. Ethical approval was obtained. Informed consent.</p>
Participants	<p>Infants delivered with a birthweight 1200 to 2199 g. Thirty-five infants randomised: intervention group (n = 21), control group (n =14), 31 completed study: intervention group (n = 18), control group (n =13). Setting: South Africa. Exclusion criteria: Mother: 1) delivered outside the unit, 2) had a caesarean section, 3) too severely ill to be able to look after themselves or their infants, 4) known to have positive HIV status, 5) giving their babies up for adoption. Infant: 1) birthweight below 1200 g or above 2199 g, 2) Apgar score below 6 at 5 min, 3) congenital malformations detected at birth Intervention group (skin-to-skin care) characteristics: Mothers: gravidity mean (1.9), parity mean (1.8), race African (8), race Coloured (12), smoking in pregnancy (7), alcohol use(6), opiate in labour (9), Oxytocin in labour (3), hypertension (3), antepartum steroids (5). Infants: male (12), resuscitation (4), birthweight (g) mean (1813) SD(260), gestational age (wks) mean (34.2) SD(1.9), appropriate for GA (13), small for GA (7) Control group (conventional care) characteristics: Mothers: gravidity mean (1.9), parity mean (1.6), race African (4), race Coloured (10), smoking in pregnancy (4), alcohol use (2), opiate in labour (6), Oxytocin in labour (1), hypertension (1), antepartum steroids (3). Infants: male: (7), resuscitation baby (1), birthweight (g) mean (1866) SD(258), gestational age (wks) mean (35.3) SD(1.9), appropriate for GA (10), small for GA (5)</p>

Interventions	<p>SKIN-TO-SKIN CARE</p> <p>All infants were delivered onto a theatre cloth on the mother's abdomen/chest , dried gently and assessed. They were then placed naked on the mother's naked chest and covered with double layered cotton cloth. The bed was placed in semi-Fowlers position</p> <p>Intervention group: The infant remained in skin-to-skin contact, was secured to the mother's chest, the hips were flexed and placed in a 'frog position', arms also flexed</p> <p>Control group: The infant was immediately transferred to a prewarmed servocontrolled closed incubator which remained with the mother in delivery ward for the first hour. If the infant temperature was < 36 °C, a cap and booties were applied along with a heat shield placed over the infant. If this was insufficient a sheet of plastic was framed over the foot end of the heat shield and the outlet of the warm air funnelled over the infant</p>	
Outcomes	<p>Infant stability in terms of a set of predetermined physiological parameters. Observations for heart rate, respiratory rate, oxygen saturation and temperature were recorded at 5 minute intervals in the first hour, then 15 minute intervals. Continuous observations for apnoea and signs of respiratory distress, blood sugar measured at 1, 3 and 6hr</p> <p>Primary Outcomes:</p> <ol style="list-style-type: none"> 1) neonatal intensive care admissions 2) number of exceeded parameters 3) composite stabilization score (SCRIP) (1-6h) 4) composite stabilization score (SCRIP) (6th h). <p>Secondary outcomes:</p> <ol style="list-style-type: none"> 1) skin temperature degrees centigrade remaining below 35.5 °C for 2 consecutive readings 2) heart rate below 100 or above 180 beats per minutes for 2 consecutive readings 3) apnoea longer than 20 seconds 4) oxygen saturation below 87% for 2 consecutive recordings, despite supplementation with nasal prong oxygen, FiO2 up to 0.6 and CPAP up to 5cm water 5) blood glucose below 2.6 mmol/L reading confirmed by laboratory 	
Notes		
Risk of bias		
Item	Authors' judgement	Description
Adequate sequence generation?	Yes	Computer minimisation method.
Allocation concealment?	Unclear	Quote: "a computer minimisation method determined allocation in a concealed manner", "second hospital ... minimisation factors and subsequent allocation by the computer were exchanged by mobile telephone, with no delays"

Bergman 2004 (Continued)

Blinding? All outcomes	No	Blinding of participants: no/unfeasible. Blinding of care givers: no/unfeasible. Blinding of outcome measurement: yes, after the pilot phase, the principal investigator and data analyses team were blinded. A statistician, blinded to the allocation, performed the statistical analyses
Incomplete outcome data addressed? All outcomes	Yes	Complete follow-up: no Thirty - one of 35 infants randomised completed the study, one infant was excluded after 2 hours when the mother required a caesarean section for a second twin, three further infants 'bailed out' within the first 3 hrs requiring additional respiratory support and moved to neonatal intensive care
Free of selective reporting?	Unclear	Unclear.
Free of other bias?	No	Quote: "The possibility of selection bias exists: many prematurely born infants delivered unexpectedly (without a nurse researcher available), and the aetiology of such premature delivery may differ from those studied" Recruitment for this study was terminated after interim analysis was carried out, based on significant results

Brennan 1996

Methods	Randomised. Single centre. Blinding of randomisation: unclear. Generation of allocation sequence: adequate, (lot drawing - pulling cards from an envelope in which 24 cards had been placed. Equal numbers of cards had either 'mattress' or 'no mattress' written on them). Allocation concealment: unclear. Blinding of intervention: participants: no/unfeasible, care givers: no/unfeasible. Blinding of outcome measurement: no. Complete follow-up: yes, all 24 infants randomised completed the study Gestational age, birthweight, and the time temperature taken after birth were recorded. The proposal for the study underwent a full review by the Institutional Review Board of the University of Florida, Health Science Centre. Informed consent.
---------	--

Participants	<p>Infants delivered with a birthweight \leq 1500 g. Twenty - four infants randomised: intervention group (n = 12), control group (n = 12). Setting: USA. Exclusion criteria: 1) the maternal temperature at delivery was greater than 38.0 °C 2) the infant had an obvious neural tube defect 3) the infant had an omphalocele, gastroschisis, exstrophy of the bladder, cloacal exstrophy, or any other open lesion than would cause greater than normal heat loss at delivery, 4) the infant had a sacral teratoma, or any other lesion that would prevent the infant from being positioned and evaluated in the normal manner All infants characteristics: gestational age range (wks) 24 to 32, birthweight range (g) 531 to 1498. Intervention group (transwarmer mattress) characteristics: gestational age (wks) mean (27.8), birthweight (g) mean (1033) Control group characteristics: gestational age (wks) mean (27.3), birthweight (g) mean (1027)</p>	
Interventions	<p>TRANSWARMER INFANT TRANSPORT MATTRESS (Prism Technologies, San Antonio, Texas). These mattresses are filled with a gel of sodium acetate, water and thickeners. Once activated they heat to approximately 40 degrees centigrade for 2 hours Intervention group: The NICU nurse activated the mattress when delivery was imminent and placed it on the radiant warmer surface with blankets on top of it and the newborn infant was immediately placed on the blankets, dried and the blankets removed, then placed directly onto the warming mattress Control group: Infants received the same care as the intervention group but without the warming mattress Both groups were evaluated and resuscitated according to the Neonatal Resuscitation Program and transported to NICU on the radiant warmer surface. The infants in the intervention group remained on the mattress until the admission temperatures were taken in the NICU</p>	
Outcomes	<p>Primary outcomes: 1) core body temperature °C (axillary) was measured using the IVAC 2080A Temp Plus 11 (IVAC, San Diego, California) as soon as possible after arrival at NICU Secondary outcomes: 1) interference to resuscitation.</p>	
Notes	<p>Individual patient data was reported in this study therefore dichotomous data for hypothermia on admission to NICU for the control and intervention group was calculated</p>	
Risk of bias		
Item	Authors' judgement	Description
Adequate sequence generation?	Yes	Generation of allocation sequence: adequate, (lot drawing - pulling cards from an envelope in which 24 cards had been placed. Equal numbers of cards had either

Brennan 1996 (Continued)

		'mattress' or 'no mattress' written on them)
Allocation concealment?	Unclear	Unclear.
Blinding? All outcomes	No	Blinding of participants: no/unfeasible. Blinding of care givers: no/unfeasible. Blinding of outcome measurement: no.
Incomplete outcome data addressed? All outcomes	Yes	Complete follow-up: yes, all 24 infants randomised completed the study
Free of other bias?	Yes	

Knobel 2005

Methods	<p>Randomised. Single centre. Blinding of randomisation: yes. Generation of allocation sequence: unclear. Allocation concealment : adequate, (opaque envelopes). Blinding of intervention: participants: no/unfeasible, care givers: no/unfeasible. Blinding of outcome measurement: no. Complete follow-up: yes, all 88 infants randomised completed the study Delivery to admission time was also recorded. Parental consent was sought if the mother was expected to deliver before completing 29 weeks' gestation</p>
Participants	<p>Infants delivered with a gestational age < 29 completed weeks. Eighty - eight infants randomised: intervention group (n = 41), control group (n = 47). Setting: USA. Exclusion criteria: 1)congenital anomalies with open lesions (e.g. gastroschisis, meningomyelocele), 2) resuscitation not undertaken because of previability, 3) meconium staining of amniotic fluid, 4) subsequent assessment indicated that infant had a gestational age \geq 29 completed weeks Intervention group (Polyurethane bag) characteristics: birthweight (g) mean (918) SD (259), gestational age (wks) mean (26.5), SD (1.4) Control group (conventional care) characteristics: birthweight (g) mean (850) SD (253) , gestational age (wks) mean (26.1) SD (1.4)</p>
Interventions	<p>POLYURETHANE BAG - measured 19" x 18" DeRoyal REF30-5510 Intervention group: Infants were placed immediately into a polyurethane bag, while still wet, up to their necks and resuscitated according to the guidelines for Neonatal Resuscitation Control group: Controls were resuscitated according to these guidelines without bags Both groups were taken to the neonatal intensive care unit on a warmer table, (with the heat off) with warm blankets on top of the infant</p>

Knobel 2005 (Continued)

Outcomes	<p>Primary outcome:</p> <ol style="list-style-type: none"> 1) core body temperature °C (rectal) on admission to the NICU 2) hypothermia on admission to the NICU defined by a core body temperature (rectal) < 36.4 °C <p>Secondary outcomes:</p> <ol style="list-style-type: none"> 1) mortality 2) major brain injury 3) duration of oxygen therapy 4) duration of hospitalisation 5) hyperthermia 6) interference with assessment and resuscitation. 	
Notes		
Risk of bias		
Item	Authors' judgement	Description
Adequate sequence generation?	Unclear	Unclear.
Allocation concealment?	Yes	Opaque envelopes.
Blinding? All outcomes	No	Blinding of participants: no/unfeasible. Blinding of care givers: no/unfeasible. Blinding of outcome measurement: no.
Incomplete outcome data addressed? All outcomes	Yes	Complete follow-up: yes, all 88 infants randomised completed the study
Free of other bias?	No	This study attempted to maintain delivery suite temperature at 26°C for all preterm deliveries, but actual temperatures ranged from 18.9 to 31.1°C. On post hoc analysis warmer delivery suite temperatures were associated with higher admission temperatures, but only the subgroup of infants who were both delivered in warm rooms and place in plastic bag had a mean temperature > 36.4 °C. There was however no significant differences between intervention and control group for mean delivery room temperature. After controlling for delivery suite temperature the mean admission temperature in the intervention group was still 0.6 °C higher than controls

Roberts 1981

Methods	<p>Randomised. Single centre. Stratified randomisation by gestational age 32 to 34 and 35 to 36 completed weeks' gestation. Blinding of randomisation: unclear. Generation of allocation sequence: adequate, (lot drawing - index cards were utilised to pre-arranged randomisation - random number tables). Allocation concealment: unclear. Blinding of intervention: participants: no/unfeasible, care givers: no/unfeasible. Blinding of outcome measurement: no Complete follow-up: yes, all 40 infants randomised completed the study Delivery room, transport incubator, highest maternal temperatures, birthweight, gestational age, method of delivery, time elapsed from birth to onset of drying and time elapsed from birth to arrival in nursery were recorded. Informed consent.</p>
Participants	<p>Infants delivered from 7am until 5 pm Monday through Friday with a gestational age of 32 to 36 completed weeks. Forty infants randomised: intervention group (n = 17), control group (n = 23). Setting: USA. Criteria for exclusion: 1) infant not appropriate birthweight for gestational age, 2) Apgar score @ 5 min < 7, 3) signs of central nervous system defect, 4) infant dropped from study on development of sepsis, 5) maternal temperature ≥ 37.8 °C, 6) delivered outside of delivery room Intervention group (stockinet cap) characteristics: male (4), female (13), white (6), black (5), Mexican - American (6), vaginal delivery (15), caesarean section (2), gestational age (wks) (32 to 34 (5), 35 to 36 (12)), gestational age (wks) mean (34.9), birthweight (g) range (1420 to 2890), birthweight (g) mean (2253.4), Apgar score @ 5 min equal to 9 (16), infants transported in incubator (7), highest maternal temperature (degrees centigrade) range (36.6 to 37.7) mean (37.1) Control group (conventional care) characteristics: male (12), female (11), white (8), black (13), Mexican - American (2), vaginal delivery (18), caesarean section (5), gestational age (wks) (32 to 34 (4), 35 to 36 (19)), gestational age (wks) mean (35.2), birthweight (g) range (1360 to 2965), birthweight (g) mean (2276.8), Apgar score @ 5 min equal to 9 (22), infants transported in incubator (12), temperature (°C) transport incubator range (33 to 37) mean (35.2), highest maternal temperature (°C) range (36.2 to 37.8) mean (37.2)</p>
Interventions	<p>STOCKINET CAP - a sterile headpiece made of stockinet (a material used to protect skin under orthopedic casts) which covers the forehead, the ears, underneath the occipital bone, and all the area above this plane Intervention group: Infants received a stockinet cap after delivery as soon as possible after being dried under a radiant warmer Control group: Infants received the same treatment as the intervention group but did not receive a stockinet cap All infants weighing < 2500 g were transported to the nursery in a transport incubator</p>

Roberts 1981 (Continued)

Outcomes	<p>Primary outcomes: 1) core body temperature °C (axillary) within 10 minutes of admission to NICU (IVAC electric thermometer Model 821) Core body temperatures °C (axillary) in the delivery room were also recorded to enable calculation of the “amount of heat lost from the time the axillary temperature was taken in the delivery room until the axillary temperature was taken in the nursery” Secondary outcomes: None reported.</p>	
Notes		
Risk of bias		
Item	Authors' judgement	Description
Adequate sequence generation?	Yes	Lot drawing - index cards were utilised to pre-arranged randomisation - random number tables
Blinding? All outcomes	No	Blinding of participants: no/unfeasible. Blinding of care givers: no/unfeasible. Blinding of outcome measurement: no.
Incomplete outcome data addressed? All outcomes	Yes	Complete follow-up: yes, all 40 infants randomised completed the study
Free of other bias?	No	For infants < 2000 g the mean delivery suite temperatures were different for the intervention and control groups

Trevisanuto 2009b

Methods	<p>Randomised. Single centre. Blinding of randomisation: adequate. Generation of allocation sequence: adequate (computer-generated, randomisation sequence balanced in blocks of six subjects). Allocation concealment: adequate (double-enclosed, opaque, sealed and sequentially numbered envelopes). Blinding of intervention: participants: no/unfeasible, care givers: no/unfeasible. Blinding of outcome measurement: no. Complete follow-up: yes, all 96 infants randomised completed the study This study was approved by the Azienda Ospedaliera - Padova Research Ethics Committee Mothers axillary temperature was recorded 30 minutes within delivery. Delivery suite temperature was not monitored throughout the study. Delivery suite temperature is normally 24° C. Time from delivery to NICU admission (minutes) was recorded Where possible, parental consent was obtained before delivery; the remaining parents received a parent information form after initial stabilisation of the infant</p>
Participants	<p>Inborn infants with a gestational age of < 29 completed weeks 96 infants randomised: intervention group (n = 32), active comparator (n = 32), control group (n = 32) Setting: Italy Exclusion criteria: 1) infants with congenital anomalies with open lesions (e.g. gastroschisis, meningomyelocele) 2) infants whose delivery was not attended by the neonatal team Intervention group (polyethylene cap) characteristics: Mothers: ethnicity white (31), ethnicity black (1), age (years) mean (32) SD (7), antenatal steroids (31), temperature (°C) mean (36.4) SD (0.4), vaginal delivery (7), caesarean section (25) Infants: gestational age (wks) mean (26.1) SD (1.4), birthweight (g) mean (834) SD (246), female (18), multiple birth (5) Active comparator (polyethylene bag) characteristics: Mothers: ethnicity white (30), ethnicity black (2), age (years) mean (33) SD (5), antenatal steroids (30), temperature (°C) mean (36.4) SD (0.5), vaginal delivery (8), caesarean section (24) Infants: gestational age (wks) mean (25.8) SD (1.5), birthweight (g) mean (800) SD (223), female (19), multiple birth (6) Control group (conventional care) characteristics: Mothers: ethnicity white (32), ethnicity black (0), age (years) mean (32) SD (4), antenatal steroids (30), temperature (°C) mean (36.4) SD (0.6), vaginal delivery (8), caesarean section (24) Infants: gestational age (wks) mean (26.3) SD (1.0), birthweight (g) mean (813) SD (225), female (16), multiple birth (7)</p>
Interventions	<p>POLYETHYLENE CAPS (Degusto Dolomiti, Belluno, Italy). Intervention group: Polyethylene caps were placed on the radiant warmer bed prior to delivery. The head of the infant was covered with a polyethylene cap immediately after birth; only the head was dried Active comparator group:</p>

	<p>Polyethylene bags (Vedovato SNC, Camposampiero, Padova, Italy, isolation transport bag 35cm by 40 cm) were placed on the radiant warmer bed prior to delivery. The infants were placed into the bag while still wet, up to the neck; only the head was dried</p> <p>Control group: The infants were placed in pre-warmed towels after drying, according to International Guidelines for Neonatal Resuscitation</p> <p>All infants were stabilised in the delivery room under radiant warmers (Babytherm® 8004/8010, Dräger Medizintechnik GmbH, Lubeck, Germany). A transport incubator (MOG® 500, Ginevri, Tecnologie Biomediche, Cecchina, Roma, Italy) was used to transfer all infants from delivery room into the NICU; where they were placed in a double walled incubator at 35°C temperature and 70% humidity (Dräger Medical AG& Co. KGaA, Lubeck, Germany)</p> <p>At this point, the cap or the wrap were removed.</p>	
Outcomes	<p>Primary outcomes:</p> <ol style="list-style-type: none"> 1) core body temperature °C (axillary) was measured with a digital thermometer (Terumo® Digital Clinical Thermometer C202, Terumo Corporation, Tokio, Japan) on admission to the NICU (immediately after cap and wrap removal) and again one hour later 2) hypothermia, defined as axillary temperature less than 36.4°C, on NICU admission was also evaluated <p>Secondary outcomes:</p> <ol style="list-style-type: none"> 1) mortality prior to hospital discharge 2) presence of major brain injury 3) tracheal intubation at birth 4) Apgar scores 5) blood gas analysis 6) serum glucose concentration on NICU admission. 	
Notes		
Risk of bias		
Item	Authors' judgement	Description
Adequate sequence generation?	Yes	Computer-generated, randomisation sequence balanced in blocks of six subjects
Allocation concealment?	Yes	Double-enclosed, opaque, sealed and sequentially numbered envelopes
Blinding? All outcomes	No	Blinding of participants: no/unfeasible. Blinding of care givers: no/unfeasible. Blinding of outcome measure: no.
Incomplete outcome data addressed? All outcomes	Yes	Complete follow-up: yes, all 96 infants randomised completed the study

Trevisanuto 2009b (Continued)

Free of other bias?	No	Delivery to admission time was significantly shorter in the control group than in the plastic wrap group Delivery room temperature was not monitored. Standard environmental temperature of delivery suite is 24°C
---------------------	----	---

Vohra 1999

Methods	<p>Randomised. Single centre. Prognostic stratification was by gestational age 23 to 27 and 28 to 31 completed weeks' gestation. Blinding of randomisation: yes. Generation of allocation sequence: adequate, (computer generated random sequence). Allocation concealment: adequate, (double-enclosed, opaque, sealed, and sequentially numbered envelopes). Blinding of intervention: participants: no/unfeasible, care givers: no/unfeasible. Blinding of outcome measurement: no. Complete follow-up: no, (59 of 62 infants randomised completed the study, 1 infant was excluded as no rectal temperature was recorded, two infants were excluded due to an imperforate anus) Delivery room, nursery temperatures and other potential confounding variables, (birthweight, maternal temperature, illnesses, medications, length of second stage of labour, length of time to arrival at nursery) were recorded. Ethics approval was obtained. No informed consent. Institutional consent was given.</p>
Participants	<p>Infants delivered with a gestational age of less than 32 completed weeks. Sixty-two infants randomised: 59 completed study, intervention group (n = 27), control group (n = 32). Setting: Canada. Exclusion criteria: 1) major congenital anomalies with open lesions, 2) infants whom the attending obstetrician considered "preivable" Intervention group (polyethylene wrap) characteristics < 28 wks: birthweight (g) mean (914) SD (163), gestational age (wk) mean (26.1) SD (1.4) Apgar score @ 1min mean (5.0) SD (3.0), Apgar score @ 5 min mean (6.7) SD (1.7), ruptured membranes (min) mean (186) SD (471), length 2nd stage (min) mean (26) SD (55) Control group (conventional care) characteristics < 28 wks: birthweight (g) mean (742) SD (206), gestational age (wks) mean (25.7) SD (1.5), Apgar score @ 1 min mean (4.3) SD (2.1), Apgar score @ 5 min mean (7.1) SD (1.4), ruptured membranes (min) mean (10) SD (21), length 2nd stage (min) (18) SD (24) Intervention group (polyethylene wrap) characteristics ≥ 28 wks: birthweight (g) mean (1251) SD (282), gestational age (wk) mean (29.6) SD (1.1) Apgar score @ 1min mean (6.5) SD (1.8), Apgar score @ 5 min mean (8.5) SD (1.1), ruptured membranes (min) mean (105) SD (310), length 2nd stage (min) mean (15) SD (25)</p>

Vohra 1999 (Continued)

	Control group (conventional care) characteristics \geq 28 wks: birthweight (g) mean (1265) SD (206), gestational age (wks) mean (29.4) SD (1.5), Apgar score @ 1 min mean (6.0) SD (2.1), Apgar score @ 5 min mean (8.0) SD (1.4), ruptured membranes (min) mean (108) SD (21), length 2nd stage (min) (13) SD (24)	
Interventions	<p>POLYETHYLENE BAG - measured 20 cm x 50 cm and was manufactured by Eastern Paper, a Division of EPC Industries</p> <p>Intervention group: A transparent polyethylene bag was opened at resuscitation under a radiant warmer and the infant was placed on it from the shoulders down. Only the head of the infant was dried; the body was wrapped without drying</p> <p>Control group: Controls were managed with the protocol described by the Neonatal Resuscitation Program. Infants were routinely dried under the radiant warmer Infants were transferred to the neonatal unit in an incubator</p>	
Outcomes	<p>Primary outcome: 1) core body temperature °C (rectal) was measured with a digital rectal thermometer on removal of the bag/wrap on admission to NICU</p> <p>Secondary outcomes: 1) mortality 2) hyperthermia 3) infection 4) skin maceration 5) interference with resuscitation.</p>	
Notes	Forty-two infants who met the inclusion criteria were not enrolled in the study reasons were as follows: still birth (1), 34 wks gestational age (2), failure to reach delivery in time (8), failure of staff to pick up envelope on the way to the delivery room (31). The non-enrolled infants did not differ significantly from those in the study population	
Risk of bias		
Item	Authors' judgement	Description
Adequate sequence generation?	Yes	Computer generated random sequence.
Allocation concealment?	Yes	Double-enclosed, opaque, sealed and sequentially numbered envelopes
Blinding? All outcomes	No	Blinding of participants: no/unfeasible. Blinding of care givers: no/unfeasible. Blinding of outcome measurement: no.
Incomplete outcome data addressed? All outcomes	Yes	Complete follow-up: no, (59 of 62 infants randomised completed the study, one infant was excluded as no rectal temperature was recorded, two infants were excluded due to an imperforate anus

Vohra 1999 (Continued)

Free of other bias?	No	There was some imbalance in birthweight between study groups. For infants < 28 completed weeks' gestational age birthweight was 914 g; SD 163 for the plastic wrap group and 742 g; SD 206 for the non wrap group
---------------------	----	---

Vohra 2004a

Methods	<p>Randomised. Single centre. Blinding of randomisation: yes. Generation of allocation sequence: adequate, (computer generated random sequence balanced in blocks of 4 subjects). Concealment of allocation: adequate, (double enclosed, opaque, sealed and sequentially numbered envelopes). Blinding of intervention: participants: no/ unfeasible, care givers: no/unfeasible. Blinding of outcome measurement: no. Complete follow-up: no, (53 of 55 infants randomised completed the study, 2 infants died in the delivery room) Multiple eligible births were separately randomised. Institutional consent was given.</p>
Participants	<p>Infants delivered with a gestational age less than 28 completed weeks where the birth was attended by the neonatal team. 55 infants randomised: intervention group (n = 28), control group (n = 27), 53 completed study: intervention group (n = 26), control group (n = 27). Setting: Canada. Exclusion criteria: 1) neonatal team did not attend the delivery, 2) major congenital anomalies that were not covered by skin e.g. gastroschisis, meningomyelocele, 3) blistering skin conditions Intervention group (polyethylene wrap) characteristics: female (14), birthweight (g) mean (858) SD (199), gestational age (wks) mean (26) SD (1.5). Control group (conventional care) characteristics: female (13), birthweight (g) mean (825) SD (270), gestational age (wks) mean (26) SD (1.4)</p>
Interventions	<p>POLYETHYLENE WRAP - measured 20 cm x 50 cm and was manufactured by Eastern Paper, a Division of EPC Industries LTD Intervention group: Infant placed on polyethylene, wrapped from the neck down, only the head was dried, stabilised under radiant warmer Control group: Control infants were dried completely according to the International Guidelines for Neonatal Resuscitation and stabilised under a radiant warmer All infants were carried by one member of the neonatal team from the delivery suite to the neonatal unit and placed in a single walled incubator with 60 percent humidity</p>

Vohra 2004a (Continued)

Outcomes	<p>Primary outcomes: 1) core body temperature °C (rectal) on admission to NICU following removal of wrap</p> <p>Secondary outcomes: 1) core body temperature °C (rectal) taken one hour later. 2) mortality - death before discharge.</p> <p>Further secondary outcomes are reported for Apgar scores, blood gas pH, Bicarbonate (mmol/L), Glucose (mmol/L), hyperthermia and interference with resuscitation</p>
Notes	<p>One hundred and seventeen infants were screened for the study. Twenty-one were excluded for the following reasons: neonatal team did not attend the delivery (13), parents refused to participate (5), and presence of congenital malformations (3). Baseline characteristics of excluded infants and their mothers were not different from those included in the study</p>

Risk of bias

Item	Authors' judgement	Description
Adequate sequence generation?	Yes	Computer generated random sequence balanced in blocks of four subjects
Allocation concealment?	Yes	Double enclosed, opaque, sealed and sequentially numbered envelopes
Blinding? All outcomes	No	Blinding of participants: no/ unfeasible. Blinding of care givers: no/unfeasible. Blinding of outcome measurement: no.
Incomplete outcome data addressed? All outcomes	Yes	Complete follow-up: no, (53 of 55 infants randomised completed the study, two infants died in the delivery room)
Free of other bias?	Yes	

Characteristics of excluded studies [ordered by study ID]

Study	Reason for exclusion
Ammari 2009	Intervention was not applied immediately at birth (within 10 mins) in the delivery suite
Anderson 2003	No thermal outcome measures were reported.
Andrade 2005	No thermal outcome measures were reported. Query re: randomisation
Baum 1968	Participants had a birthweight greater \geq 6 lb.

(Continued)

Bell 1983	Intervention was not applied immediately at birth (within 10 mins) in the delivery suite
Bergstrom 2005	Participants were normal infants, mean gestational age at delivery : 38 wks (Intervention group), 38.4 wks control group. Intervention not strictly for the prevention of hypothermia immediately at birth
Besch 1971	Participants had a birthweight > 2000 g.
Bier 1996	Intervention was not applied immediately at birth (within 10 mins) in the delivery suite
Boo 2007	Intervention was not applied immediately at birth (within 10 mins) in the delivery suite. Participants were normal term infants undergoing phototherapy
Brice 1981	Intervention was not applied immediately at birth (within 10 mins) in the delivery suite
Bystrova 2003	Participants had a birthweight > 2500 g.
Bystrova 2007	Participants were term.
Carfoot 2005	Participants were term.
Cattaneo 1998	Intervention was not applied immediately at birth (within 10 mins) in the delivery suite
Chaput 1979	Participants had a birthweight > 2000 g.
Charpak 1997	Intervention was not applied immediately at birth (within 10 mins) in the delivery suite. No thermal outcome measures were reported
Cheah 2000	Participants were term.
Christensson 1992	Participants were term.
Christensson 1995	Participants were term.
Christensson 1996	Participants were term delivered by caesarean section.
Christensson 1998	Intervention was not applied immediately at birth (within 10 mins) in the delivery suite
Chwo 1999	Intervention was not applied immediately at birth (within 10 mins) in the delivery suite but on the day following birth
Chwo 2002	Intervention was not applied immediately at birth (within 10 mins) in the delivery suite but on the day following birth
Coles 1979	Participants were term.
Dahm 1972	Participants were term.

(Continued)

Daniel 2004	Unable to obtain a copy of the abstract.
Dannaway 2009	This study compared two interventions applied after delivery: warming mattress (sodium acetate warming blanket) to polyethylene wrap. This comparison is not within the scope of this review
Darmstadt 2007	Intervention was not applied immediately at birth (within 10 mins) in the delivery suite
Darmstadt 2008	Primary outcome measure was rate of nosocomial infection. Hypothermia was reported as a morbidity among neonates who died
Day 1964	Intervention was not applied immediately at birth (within 10 mins) in the delivery suite
Dodd 2003	No thermal outcome measures were reported.
Erlandsson 2007	Participants were term delivered by caesarean section. No thermal outcome measures were reported
Fallis 2006	Participants were term. Intervention was maternal warming during caesarean section
Fardig 1980	Participants were term.
Ferber 2004	Participants were term. Intervention was not applied immediately at birth (within 10 mins) in the delivery suite. No thermal outcome measures were reported
Gray 2004	Intervention was not applied immediately at birth (within 10 mins) in the delivery suite. Participants were at least 3 days old
Green-Abate 1994	Intervention was not applied immediately at birth (within 10 mins) in delivery suite. Participants were less than 1 week old
Greer 1988	Participants were term and had a birthweight > 2500 g.
Grover 1994	Participants were term and the intervention was not applied immediately at birth (within 10 mins) in the delivery suite. Participants were aged 11 to 95 days
Gulezian 1980	Insufficient information - unable to obtain a copy of the abstract
Harrison 2004	Insufficient information - unable to obtain a copy of the abstract
Hellin Martinez 2000	Participants were term.
Hobbs 1975	Participants were term.
Holzman 1985	Participants were term.
Horn 2002	Query if participants were term as infants were delivered by elective caesarean section. The author was contacted for verification. Intervention was active warming during caesarean section

(Continued)

Huang 2002	Participants had a birthweight \geq 2500 g.
Huang 2006	Participants were term. Intervention was not applied immediately at birth (within 10 mins) in the delivery suite
Johanson 1992	Participants were term with a mean gestational age 39.1 weeks. Randomised controlled intervention study compared to a prospective observational study
Kadam 2005	Intervention was not applied immediately at birth (within 10 mins) in the delivery suite. (Verified by author)
Kaushal 2005	Intervention was not applied immediately at birth (within 10 mins) in the delivery suite
Kumar 2008	This cluster-randomised controlled efficacy trial looked at three community-based interventions which are outside the scope of this review
Legault 1993	Intervention was not applied immediately at birth (within 10 mins) in the delivery suite. Participants were aged 1 week or more
Ludington-Hoe 1994	Intervention was not applied immediately at birth (within 10 mins) in the delivery suite. Participants in intervention group averaged 18 days old and control participants averaged 13 days old
Ludington-Hoe 2000	Participants had a postnatal age of 6 to 53 days on enrolment to the study therefore the intervention was not applied immediately at birth (within 10 mins) in the delivery suite
Ludington-Hoe 2004	Intervention was not applied immediately at birth (within 10 mins) in the delivery suite
Marks 1985	Participants had a gestational age of 36 plus or minus one week. Intervention was not applied immediately at birth (within 10 mins) in the delivery suite. Participants were aged 14 to 68 days
Mathew 2008	This study compared vinyl bag (Vi-Drape) to thermal mattress (Transwarmer). Both groups underwent drying of the head and placement of a stocking cap. This comparison is not within the scope of this review
Mazurek 1999	Participants were term.
Medves 2004	Participants were term.
Meyer 2001	Intervention was not applied immediately at birth (within 10 mins) in the delivery suite. Intervention applied on arrival at NICU
Meyer 2007	This study compared plastic wrap plus radiant warmer to plastic wrap plus incubator for transport to NICU. All infants were wrapped under a radiant warmer at birth but it is unlikely that assignment to incubator or radiant warmer took place within 10 minutes of birth for all infants. This comparison is not within the scope of this review
Miles 2006	Intervention was not applied immediately at birth (within 10 mins) in the delivery suite. No thermal outcome measures were reported

(Continued)

Monterosso 1999	Intervention was not applied immediately at birth (within 10 mins) in the delivery suite. Participants were at least one week of age at time of study
Moore 2007	Participants were term. No thermal outcome measures were reported
Nopper 1996	Intervention was not applied immediately at birth (within 10 mins) in the delivery suite. Participants enrolled within the first 96 hours after birth
Nuntnarumit 2004	This study compared plastic wrap and plastic bag to plastic wrap only. Plastic wrap only was not considered to be 'routine care' as pre-specified in the review inclusion criteria
Omene 1978	Participants were term.
Pattinson 2005	Intervention was an educational package (with/without facilitation visits) on the implementation of kangaroo mother care
Punthmatharith 2001	Thermal outcome measures were not reported.
Raman 1992	Participants were term.
Ramanathan 2001	Intervention was not applied immediately at birth (within 10 mins) in the delivery suite. Thermal outcome measures were not reported
Rao 2008	Intervention was not applied immediately at birth (within 10 mins) in the delivery suite
Roberts 2000	Intervention was not applied immediately at birth (within 10 mins) in the delivery suite. Participants were born at 30 or more weeks' gestation or corrected age
Rojas 2001	Intervention was not applied immediately at birth (within 10 mins) in the delivery suite. Thermal outcome measures were not reported
Ruiz 1998	Intervention not applied immediately at birth (within 10 mins) in the delivery suite. Thermal outcome measures were not reported
Sankaranarayanan2005	Intervention was not applied immediately at birth (within 10 mins) in the delivery suite
Sarman 1989	Intervention was not applied immediately at birth (within 10 mins) in the delivery suite
Sarman 1992	Intervention was not applied immediately at birth (within 10 mins) in the delivery suite. Participants were 3 to 32 days of age during the study
Short 1998	Intervention was not applied immediately at birth (within 10 mins) in the delivery suite. Participants day of life 32.9 plus or minus 15.9 days
Sloan 1994	Intervention was not applied immediately at birth (within 10 mins) in the delivery suite. No thermal outcome measures were reported

(Continued)

Syfrett 1993	Intervention was not applied immediately at birth (within 10 mins) in the delivery suite
Trevisanuto 2009a	Participants were mainly term infants delivered by caesarean section. Primary outcome measure was the time elapsed from the cord clamping to the positioning of the infant under the radiant warmer for infants managed with the 'cicogna' system and those who received standard care
Tsogt 2005	Participants were term. Intervention was not applied immediately at birth (within 10 mins) in the delivery suite
Vaidya 2005	Participants were term. Intervention was not applied immediately at birth (within 10 mins) in the delivery suite. No thermal outcomes were measured
van den Bosch 1990	Participants were term.
van den Bosch 1996	Intervention was not applied immediately at birth (within 10 mins) in the delivery suite. Intervention applied after 48 hours in the NICU
Villalon 1992	Participants were full term.
Worku 2005	Intervention was not applied immediately at birth (within 10 mins) in the delivery suite
Yeh 1980	Intervention was not applied immediately at birth (within 10 mins) in the delivery suite
Yokoyama 2009	Participants were term infants delivered by elective cesarean section

Characteristics of studies awaiting assessment [ordered by study ID]

Punnahitananda 2008

Methods	Randomised. Single centre. Blinding of randomisation: need further information. Generation of allocation sequence: need further information. Allocation concealment: need further information. Blinding of intervention: need further information. Blinding of outcome measurement: need further information. Complete follow-up: no, 130 infants randomised, 122 completed the study, need further information
Participants	Inborn infants of 34 weeks' gestation or less. 130 infants randomised, 122 completed the study: intervention group (n = 61), control group (n = 61) Setting: Thailand. Exclusion criteria: need further information. Infant characteristics: need further information.
Interventions	POLYETHYLENE PLASTIC WRAP/BAG. Intervention group: Infants were placed in polyethylene plastic bags immediately after birth in delivery room, leaving only the head

Punnahitananda 2008 (Continued)

	uncovered and were kept in plastic bags for three hours. Any resuscitation treatment was carried out with the bags covering the bodies Control group: Infants were dried and resuscitated per standard protocol before transferred to nursery All infants were stabilised under radiant warmers.
Outcomes	Primary outcomes: 1) core body temperature degrees centigrade (rectal) on admission to the nursery and hourly there after for three hours using a digital thermometer 2) hypothermia on admission to the nursery and during the three hour study period (definition of hypothermia not given)
Notes	Unable to contact author. Further information required for: definition of hypothermia and incidence, data for rectal temperature on admission to nursery for infants of 31-34 weeks' gestation and information regarding the methodological quality

Characteristics of ongoing studies [ordered by study ID]

Vohra 2004

Trial name or title	Multicentred randomised controlled trial of heat loss prevention (HeLP) in the delivery room
Methods	The randomised controlled trial (RCT) is a two-intervention, parallel design with a 24 month accrual period. Preterm infants meeting the eligibility criteria will be randomly assigned within three gestational age strata (less than 24 weeks, 24+0 to 25+6 weeks gestation, and 26+0 to 27+6 weeks gestation) to either the occlusive skin wrap group or the standard of care (non-wrap) group
Participants	Infants less than 28 weeks' gestation.
Interventions	Polyethylene occlusive skin wrap applied immediately following birth in the delivery suite
Outcomes	Mortality, axillary temperature, rectal temperature (in some) and clinically important variables including: APGAR scores, incidence of acidosis, hypotension, hypoglycaemia, seizures, patent ductus arteriosus, respiratory distress syndrome/chronic lung disease, necrotizing enterocolitis, intraventricular haemorrhage. Also length of stay and cause of death
Starting date	Infant enrolments began in December 2004.
Contact information	Sunita Vohra MD FRCPC MSc Director CARE Program for Integrative Health and Healing Director PedCAM Research and Education Network Professor, Department of Pediatrics, Faculty of Medicine and School of Public Health University of Alberta, Edmonton AB, Canada. Email: svohra@ualberta.ca
Notes	This multicentre trial will involve 40 centres, Canada, US and UK, all part of the Vermont-Oxford Network. This study has received funding from the Canadian Institute of Health Research (CIHR).Forty-five centers

Vohra 2004 (Continued)

are currently involved in this study

DATA AND ANALYSES

Comparison 1. Plastic wrap versus routine care

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Core body temperature (°C) on admission to NICU or up to 2 hours after birth	4	264	Mean Difference (IV, Fixed, 95% CI)	0.57 [0.37, 0.77]
1.1 < 28 completed weeks' gestational age	4	223	Mean Difference (IV, Fixed, 95% CI)	0.68 [0.45, 0.91]
1.2 28 to 31 completed weeks' gestational age	1	41	Mean Difference (IV, Fixed, 95% CI)	0.17 [-0.27, 0.61]
2 Core body temperature (°C) 1 hour after the initial admission temperature to the NICU was taken	2	117	Mean Difference (IV, Fixed, 95% CI)	0.40 [0.16, 0.65]
2.1 < 28 completed weeks' gestational age	2	117	Mean Difference (IV, Fixed, 95% CI)	0.40 [0.16, 0.65]
3 Hypothermia on admission to NICU: core body temperature < 36.5°C or skin temperature < 36°C	2	152	Risk Ratio (M-H, Fixed, 95% CI)	0.66 [0.51, 0.84]
3.1 < 28 completed weeks' gestational age	2	152	Risk Ratio (M-H, Fixed, 95% CI)	0.66 [0.51, 0.84]
4 Death within hospital stay	4	266	Risk Ratio (M-H, Fixed, 95% CI)	0.66 [0.35, 1.24]
4.1 < 28 completed weeks' gestational age	4	225	Risk Ratio (M-H, Fixed, 95% CI)	0.66 [0.35, 1.24]
4.2 28 to 31 completed weeks' gestational age	1	41	Risk Ratio (M-H, Fixed, 95% CI)	Not estimable
5 Major brain injury	2	152	Risk Ratio (M-H, Fixed, 95% CI)	1.10 [0.41, 2.98]
5.1 < 28 completed weeks' gestational age	2	152	Risk Ratio (M-H, Fixed, 95% CI)	1.10 [0.41, 2.98]
6 Duration of oxygen therapy (days)	1	88	Mean Difference (IV, Fixed, 95% CI)	-6.51 [-23.30, 10.28]
6.1 < 28 completed weeks' gestational age	1	88	Mean Difference (IV, Fixed, 95% CI)	-6.51 [-23.30, 10.28]
7 Duration of hospitalisation (days)	1	88	Mean Difference (IV, Fixed, 95% CI)	-5.49 [-19.93, 8.95]
7.1 < 28 completed weeks' gestational age	1	88	Mean Difference (IV, Fixed, 95% CI)	-5.49 [-19.93, 8.95]
8 Apgar at 1 minute			Other data	No numeric data
8.1 < 28 completed weeks' gestational age			Other data	No numeric data
9 Apgar at 1 minute	1	64	Mean Difference (IV, Fixed, 95% CI)	0.20 [-0.83, 1.23]
9.1 < 28 completed weeks' gestational age	1	64	Mean Difference (IV, Fixed, 95% CI)	0.20 [-0.83, 1.23]
10 Apgar at 5 minutes			Other data	No numeric data

10.1 < 28 completed weeks' gestational age			Other data	No numeric data
11 Apgar at 5 minutes	1	64	Mean Difference (IV, Fixed, 95% CI)	0.40 [-0.19, 0.99]
11.1 < 28 completed weeks' gestational age	1	64	Mean Difference (IV, Fixed, 95% CI)	0.40 [-0.19, 0.99]
12 First blood gas pH	2	117	Mean Difference (IV, Fixed, 95% CI)	0.01 [-0.02, 0.04]
12.1 < 28 completed weeks' gestational age	2	117	Mean Difference (IV, Fixed, 95% CI)	0.01 [-0.02, 0.04]
13 Bicarbonate (mmol/L)	2	117	Mean Difference (IV, Fixed, 95% CI)	0.31 [-0.72, 1.35]
13.1 < 28 completed weeks' gestational age	2	117	Mean Difference (IV, Fixed, 95% CI)	0.31 [-0.72, 1.35]
14 First serum glucose concentration (mmol/L)	2	117	Mean Difference (IV, Fixed, 95% CI)	0.01 [-0.36, 0.39]
14.1 < 28 completed weeks' gestational age	2	117	Mean Difference (IV, Fixed, 95% CI)	0.01 [-0.36, 0.39]
15 Intubation in delivery room	1	64	Risk Ratio (M-H, Fixed, 95% CI)	1.0 [0.63, 1.58]
15.1 < 28 completed weeks' gestational age	1	64	Risk Ratio (M-H, Fixed, 95% CI)	1.0 [0.63, 1.58]
16 Hyperthermia on admission to NICU: core body temperature > 37.5°C	1	53	Risk Ratio (M-H, Fixed, 95% CI)	4.82 [0.24, 95.88]
16.1 < 28 completed weeks' gestational age	1	53	Risk Ratio (M-H, Fixed, 95% CI)	4.82 [0.24, 95.88]

Comparison 2. Plastic cap versus routine care

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Core body temperature (°C) on admission to NICU or up to 2 hours after birth	1	64	Mean Difference (IV, Fixed, 95% CI)	0.80 [0.41, 1.19]
1.1 < 29 completed weeks' gestational age	1	64	Mean Difference (IV, Fixed, 95% CI)	0.80 [0.41, 1.19]
2 Core body temperature (°C) 1 hour after the initial admission temperature to the NICU was taken	1	64	Mean Difference (IV, Fixed, 95% CI)	0.80 [0.46, 1.14]
2.1 < 29 completed weeks' gestational age	1	64	Mean Difference (IV, Fixed, 95% CI)	0.80 [0.46, 1.14]
3 Hypothermia on admission to NICU: core body temperature < 36.4 °C	1	64	Risk Ratio (M-H, Fixed, 95% CI)	0.48 [0.32, 0.73]
3.1 < 29 completed weeks' gestational age	1	64	Risk Ratio (M-H, Fixed, 95% CI)	0.48 [0.32, 0.73]
4 Death within hospital stay	1	64	Risk Ratio (M-H, Fixed, 95% CI)	1.5 [0.27, 8.38]
4.1 < 29 completed weeks' gestational age	1	64	Risk Ratio (M-H, Fixed, 95% CI)	1.5 [0.27, 8.38]
5 Major brain injury	1	64	Risk Ratio (M-H, Fixed, 95% CI)	1.5 [0.27, 8.38]

5.1 < 29 completed weeks' gestational age	1	64	Risk Ratio (M-H, Fixed, 95% CI)	1.5 [0.27, 8.38]
6 Apgar score at 1 minute	1	64	Mean Difference (IV, Fixed, 95% CI)	0.80 [-0.21, 1.81]
6.1 < 29 completed weeks' gestational age	1	64	Mean Difference (IV, Fixed, 95% CI)	0.80 [-0.21, 1.81]
7 Apgar score at 5 minutes	1	64	Mean Difference (IV, Fixed, 95% CI)	0.70 [0.08, 1.32]
7.1 < 29 completed weeks' gestational age	1	64	Mean Difference (IV, Fixed, 95% CI)	0.70 [0.08, 1.32]
8 First blood gas pH	1	64	Mean Difference (IV, Fixed, 95% CI)	0.01 [-0.03, 0.05]
8.1 < 29 completed weeks' gestational age	1	64	Mean Difference (IV, Fixed, 95% CI)	0.01 [-0.03, 0.05]
9 Bicarbonate (mmol/L)	1	64	Mean Difference (IV, Fixed, 95% CI)	1.0 [-0.25, 2.25]
9.1 < 29 completed weeks' gestational age	1	64	Mean Difference (IV, Fixed, 95% CI)	1.0 [-0.25, 2.25]
10 First serum glucose concentration (mmol/L)	1	64	Mean Difference (IV, Fixed, 95% CI)	0.10 [-0.42, 0.62]
10.1 < 29 completed weeks' gestational age	1	64	Mean Difference (IV, Fixed, 95% CI)	0.10 [-0.42, 0.62]
11 Intubation in delivery room	1	64	Risk Ratio (M-H, Fixed, 95% CI)	0.82 [0.49, 1.37]
11.1 < 28 completed weeks' gestational age	1	64	Risk Ratio (M-H, Fixed, 95% CI)	0.82 [0.49, 1.37]

Comparison 3. Stockinet cap versus routine care

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Core body temperature (°C) on admission to NICU or up to 2 hours after birth	1	40	Mean Difference (IV, Fixed, 95% CI)	0.15 [-0.18, 0.48]
1.1 < 2000 g birthweight	1	10	Mean Difference (IV, Fixed, 95% CI)	0.70 [-0.01, 1.41]
1.2 > or = 2000 g birthweight	1	30	Mean Difference (IV, Fixed, 95% CI)	Not estimable
2 Hypothermia on admission to NICU: core body temperature < 36.5°C or skin temperature < 36°C	1	40	Risk Ratio (M-H, Fixed, 95% CI)	0.90 [0.48, 1.71]
2.1 32 to 36 completed weeks' gestational age	1	40	Risk Ratio (M-H, Fixed, 95% CI)	0.90 [0.48, 1.71]

Comparison 4. Skin-to-skin care versus routine care

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Hypothermia: skin temperature < 35.5°C for 2 consecutive recordings	1	31	Risk Ratio (M-H, Fixed, 95% CI)	0.09 [0.01, 0.64]
1.1 1200 g to 2199 g birthweight	1	31	Risk Ratio (M-H, Fixed, 95% CI)	0.09 [0.01, 0.64]
2 Hypoglycaemia: blood glucose level < 2.6 mmol/L	1	31	Risk Ratio (M-H, Fixed, 95% CI)	0.24 [0.03, 2.06]
2.1 1200 g to 2199 g birthweight	1	31	Risk Ratio (M-H, Fixed, 95% CI)	0.24 [0.03, 2.06]

Comparison 5. Transwarmer mattress versus routine care


Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Core body temperature (°C) on admission to NICU or up to 2 hours after birth	1	24	Mean Difference (IV, Fixed, 95% CI)	1.60 [0.83, 2.37]
1.1 < or = 1500 g birthweight	1	24	Mean Difference (IV, Fixed, 95% CI)	1.60 [0.83, 2.37]
2 Hypothermia on admission to NICU: core body temperature < 36.5 °C or skin temperature < 36°C	1	24	Risk Ratio (M-H, Fixed, 95% CI)	0.3 [0.11, 0.83]
2.1 < or = 1500 g birthweight	1	24	Risk Ratio (M-H, Fixed, 95% CI)	0.3 [0.11, 0.83]

Analysis 1.1. Comparison 1 Plastic wrap versus routine care, Outcome 1 Core body temperature (°C) on admission to NICU or up to 2 hours after birth.

Review: Interventions to prevent hypothermia at birth in preterm and/or low birthweight infants

Comparison: 1 Plastic wrap versus routine care

Outcome: 1 Core body temperature (C) on admission to NICU or up to 2 hours after birth




Analysis 1.2. Comparison 1 Plastic wrap versus routine care, Outcome 2 Core body temperature (°C) 1 hour after the initial admission temperature to the NICU was taken.

Review: Interventions to prevent hypothermia at birth in preterm and/or low birthweight infants

Comparison: 1 Plastic wrap versus routine care

Outcome: 2 Core body temperature (C) 1 hour after the initial admission temperature to the NICU was taken




Analysis 1.3. Comparison 1 Plastic wrap versus routine care, Outcome 3 Hypothermia on admission to NICU: core body temperature < 36.5°C or skin temperature < 36°C.

Review: Interventions to prevent hypothermia at birth in preterm and/or low birthweight infants

Comparison: 1 Plastic wrap versus routine care

Outcome: 3 Hypothermia on admission to NICU: core body temperature < 36.5C or skin temperature < 36C




Analysis 1.4. Comparison 1 Plastic wrap versus routine care, Outcome 4 Death within hospital stay.

Review: Interventions to prevent hypothermia at birth in preterm and/or low birthweight infants

Comparison: 1 Plastic wrap versus routine care

Outcome: 4 Death within hospital stay




Analysis 1.5. Comparison 1 Plastic wrap versus routine care, Outcome 5 Major brain injury.

Review: Interventions to prevent hypothermia at birth in preterm and/or low birthweight infants

Comparison: 1 Plastic wrap versus routine care

Outcome: 5 Major brain injury




Analysis 1.6. Comparison 1 Plastic wrap versus routine care, Outcome 6 Duration of oxygen therapy (days).

Review: Interventions to prevent hypothermia at birth in preterm and/or low birthweight infants

Comparison: 1 Plastic wrap versus routine care

Outcome: 6 Duration of oxygen therapy (days)




Analysis 1.7. Comparison 1 Plastic wrap versus routine care, Outcome 7 Duration of hospitalisation (days).

Review: Interventions to prevent hypothermia at birth in preterm and/or low birthweight infants

Comparison: 1 Plastic wrap versus routine care

Outcome: 7 Duration of hospitalisation (days)



Analysis 1.8. Comparison 1 Plastic wrap versus routine care, Outcome 8 Apgar at 1 minute.

Apgar at 1 minute


Study	Group	N	Median	Interquartile range	P value
< 28 completed weeks' gestational age					
Vohra 2004a	Intervention	27	6	2 - 6	0.6
Vohra 2004a	Control	26	6	3 - 6	

Analysis 1.9. Comparison 1 Plastic wrap versus routine care, Outcome 9 Apgar at 1 minute.

Review: Interventions to prevent hypothermia at birth in preterm and/or low birthweight infants

Comparison: 1 Plastic wrap versus routine care

Outcome: 9 Apgar at 1 minute



Analysis 1.10. Comparison 1 Plastic wrap versus routine care, Outcome 10 Apgar at 5 minutes.

Apgar at 5 minutes


Study	Group	N	Median	Interquartile Range	P value
< 28 completed weeks' gestational age					
Vohra 2004a	Intervention	27	7	6 - 8	0.9
Vohra 2004a	Control	26	7	7 - 8	0.9

Analysis 1.11. Comparison 1 Plastic wrap versus routine care, Outcome 11 Apgar at 5 minutes.

Review: Interventions to prevent hypothermia at birth in preterm and/or low birthweight infants

Comparison: 1 Plastic wrap versus routine care

Outcome: 11 Apgar at 5 minutes




Analysis 1.12. Comparison 1 Plastic wrap versus routine care, Outcome 12 First blood gas pH.

Review: Interventions to prevent hypothermia at birth in preterm and/or low birthweight infants

Comparison: 1 Plastic wrap versus routine care

Outcome: 12 First blood gas pH




Analysis 1.13. Comparison 1 Plastic wrap versus routine care, Outcome 13 Bicarbonate (mmol/L).

Review: Interventions to prevent hypothermia at birth in preterm and/or low birthweight infants

Comparison: 1 Plastic wrap versus routine care

Outcome: 13 Bicarbonate (mmol/L)




Analysis 1.14. Comparison 1 Plastic wrap versus routine care, Outcome 14 First serum glucose concentration (mmol/L).

Review: Interventions to prevent hypothermia at birth in preterm and/or low birthweight infants

Comparison: 1 Plastic wrap versus routine care

Outcome: 14 First serum glucose concentration (mmol/L)




Analysis 1.15. Comparison 1 Plastic wrap versus routine care, Outcome 15 Intubation in delivery room.

Review: Interventions to prevent hypothermia at birth in preterm and/or low birthweight infants

Comparison: 1 Plastic wrap versus routine care

Outcome: 15 Intubation in delivery room




Analysis 1.16. Comparison 1 Plastic wrap versus routine care, Outcome 16 Hyperthermia on admission to NICU: core body temperature > 37.5°C.

Review: Interventions to prevent hypothermia at birth in preterm and/or low birthweight infants

Comparison: 1 Plastic wrap versus routine care

Outcome: 16 Hyperthermia on admission to NICU: core body temperature > 37.5C




Analysis 2.1. Comparison 2 Plastic cap versus routine care, Outcome 1 Core body temperature (°C) on admission to NICU or up to 2 hours after birth.

Review: Interventions to prevent hypothermia at birth in preterm and/or low birthweight infants

Comparison: 2 Plastic cap versus routine care

Outcome: 1 Core body temperature (C) on admission to NICU or up to 2 hours after birth




Analysis 2.2. Comparison 2 Plastic cap versus routine care, Outcome 2 Core body temperature (°C) 1 hour after the initial admission temperature to the NICU was taken.

Review: Interventions to prevent hypothermia at birth in preterm and/or low birthweight infants

Comparison: 2 Plastic cap versus routine care

Outcome: 2 Core body temperature (C) 1 hour after the initial admission temperature to the NICU was taken




Analysis 2.3. Comparison 2 Plastic cap versus routine care, Outcome 3 Hypothermia on admission to NICU: core body temperature < 36.4 °C.

Review: Interventions to prevent hypothermia at birth in preterm and/or low birthweight infants

Comparison: 2 Plastic cap versus routine care

Outcome: 3 Hypothermia on admission to NICU: core body temperature < 36.4 C




Analysis 2.4. Comparison 2 Plastic cap versus routine care, Outcome 4 Death within hospital stay.

Review: Interventions to prevent hypothermia at birth in preterm and/or low birthweight infants

Comparison: 2 Plastic cap versus routine care

Outcome: 4 Death within hospital stay




Analysis 2.5. Comparison 2 Plastic cap versus routine care, Outcome 5 Major brain injury.

Review: Interventions to prevent hypothermia at birth in preterm and/or low birthweight infants

Comparison: 2 Plastic cap versus routine care

Outcome: 5 Major brain injury




Analysis 2.6. Comparison 2 Plastic cap versus routine care, Outcome 6 Apgar score at 1 minute.

Review: Interventions to prevent hypothermia at birth in preterm and/or low birthweight infants

Comparison: 2 Plastic cap versus routine care

Outcome: 6 Apgar score at 1 minute




Analysis 2.7. Comparison 2 Plastic cap versus routine care, Outcome 7 Apgar score at 5 minutes.

Review: Interventions to prevent hypothermia at birth in preterm and/or low birthweight infants

Comparison: 2 Plastic cap versus routine care

Outcome: 7 Apgar score at 5 minutes




Analysis 2.8. Comparison 2 Plastic cap versus routine care, Outcome 8 First blood gas pH.

Review: Interventions to prevent hypothermia at birth in preterm and/or low birthweight infants

Comparison: 2 Plastic cap versus routine care

Outcome: 8 First blood gas pH




Analysis 2.9. Comparison 2 Plastic cap versus routine care, Outcome 9 Bicarbonate (mmol/L).

Review: Interventions to prevent hypothermia at birth in preterm and/or low birthweight infants

Comparison: 2 Plastic cap versus routine care

Outcome: 9 Bicarbonate (mmol/L)




Analysis 2.10. Comparison 2 Plastic cap versus routine care, Outcome 10 First serum glucose concentration (mmol/L).

Review: Interventions to prevent hypothermia at birth in preterm and/or low birthweight infants

Comparison: 2 Plastic cap versus routine care

Outcome: 10 First serum glucose concentration (mmol/L)




Analysis 2.1.1. Comparison 2 Plastic cap versus routine care, Outcome 11 Intubation in delivery room.

Review: Interventions to prevent hypothermia at birth in preterm and/or low birthweight infants

Comparison: 2 Plastic cap versus routine care

Outcome: 11 Intubation in delivery room




Analysis 3.1. Comparison 3 Stockinet cap versus routine care, Outcome 1 Core body temperature (°C) on admission to NICU or up to 2 hours after birth.

Review: Interventions to prevent hypothermia at birth in preterm and/or low birthweight infants

Comparison: 3 Stockinet cap versus routine care

Outcome: 1 Core body temperature (C) on admission to NICU or up to 2 hours after birth




Analysis 3.2. Comparison 3 Stockinet cap versus routine care, Outcome 2 Hypothermia on admission to NICU: core body temperature < 36.5°C or skin temperature < 36°C.

Review: Interventions to prevent hypothermia at birth in preterm and/or low birthweight infants

Comparison: 3 Stockinet cap versus routine care

Outcome: 2 Hypothermia on admission to NICU: core body temperature < 36.5C or skin temperature < 36C




Analysis 4.1. Comparison 4 Skin-to-skin care versus routine care, Outcome 1 Hypothermia: skin temperature < 35.5°C for 2 consecutive recordings.

Review: Interventions to prevent hypothermia at birth in preterm and/or low birthweight infants

Comparison: 4 Skin-to-skin care versus routine care

Outcome: 1 Hypothermia: skin temperature < 35.5C for 2 consecutive recordings




Analysis 4.2. Comparison 4 Skin-to-skin care versus routine care, Outcome 2 Hypoglycaemia: blood glucose level < 2.6 mmol/L.

Review: Interventions to prevent hypothermia at birth in preterm and/or low birthweight infants

Comparison: 4 Skin-to-skin care versus routine care

Outcome: 2 Hypoglycaemia: blood glucose level < 2.6 mmol/L




Analysis 5.1. Comparison 5 Transwarmer mattress versus routine care, Outcome 1 Core body temperature (°C) on admission to NICU or up to 2 hours after birth.

Review: Interventions to prevent hypothermia at birth in preterm and/or low birthweight infants

Comparison: 5 Transwarmer mattress versus routine care

Outcome: 1 Core body temperature (C) on admission to NICU or up to 2 hours after birth




Analysis 5.2. Comparison 5 Transwarmer mattress versus routine care, Outcome 2 Hypothermia on admission to NICU: core body temperature < 36.5 °C or skin temperature < 36°C.

Review: Interventions to prevent hypothermia at birth in preterm and/or low birthweight infants

Comparison: 5 Transwarmer mattress versus routine care

Outcome: 2 Hypothermia on admission to NICU: core body temperature < 36.5 C or skin temperature < 36C



WHAT'S NEW

Last assessed as up-to-date: 7 February 2010.

Date	Event	Description
10 February 2010	New search has been performed	This updates the review "Interventions to prevent hypothermia at birth in preterm and/or low birthweight infants" published in The Cochrane Database of Systematic Reviews, Issue 1, 2008 (McCall 2008). One additional eligible study has been included. Data are included in the plastic wrap versus routine care comparison group and a new comparison group 'Plastic cap versus routine care' has been added
10 February 2010	New citation required and conclusions have changed	New citation. Conclusions not changed.

HISTORY

Protocol first published: Issue 2, 2003

Review first published: Issue 1, 2005

Date	Event	Description
3 July 2008	Amended	Converted to new review format.
13 September 2007	New search has been performed	<p>This review updates the existing review of “Interventions to prevent hypothermia at birth in preterm and/or low birthweight babies”, published in The Cochrane Library, Issue 1, 2005 (McCall 2005).</p> <p>Literature searches were updated to July week 4, 2007. No additional studies fulfilled the criteria for inclusion in this review. Secondary outcome measures for Vohra 2004a were updated as a result of publication of the full manuscript.</p> <p>There were no changes to the reviewer’s conclusions nor to implications for practice or research</p>
13 September 2007	New citation required but conclusions have not changed	Substantive amendment.

CONTRIBUTIONS OF AUTHORS

The review was conceived by the Northern Ireland Neonatal Intensive Care Outcomes Research and Evaluation Group (NICORE).

The review was co-ordinated and the manuscript prepared in RevMan5 by the contact reviewer (EC). The search strategy was designed by EC and the QUB Medical Faculty Librarian. The data extraction proforma was drafted by EC and edited by FA, HH, JJ and SV.

FA, HH, JJ, SV and EC screened abstracts and papers against the inclusion criteria, appraised the quality and extracted the data. Entry onto RevMan was carried out by EC.

HH, JJ and SV provided clinical input at all stages of the review process. FA and HH also provided methodological input.

All reviewers were involved equally in the development of the content of the final manuscript.

DECLARATIONS OF INTEREST

SV is principal investigator on two studies that met the review eligibility criteria. This team member was excluded from the study appraisal process for these studies.

SV is also co-leading the current multicentre HeLP study.

SOURCES OF SUPPORT

Internal sources

- Northern Ireland Neonatal Outcomes Research and Evaluation Group (NICORE), UK.

External sources

- Research & Development Office - Northern Ireland, UK.

INDEX TERMS

Medical Subject Headings (MeSH)

*Infant, Low Birth Weight; Hypothermia [*prevention & control]; Infant, Newborn; Infant, Premature; Infant, Premature, Diseases [*prevention & control]; Perinatal Care [methods]; Randomized Controlled Trials as Topic

MeSH check words

Humans