

National Library  
of Canada

Bibliothèque nationale  
du Canada

Canadian Theses Service / Service des thèses canadiennes

Ottawa, Canada  
K1A 0N4

## NOTICE

The quality of this microform is heavily dependent upon the quality of the original thesis submitted for microfilming. Every effort has been made to ensure the highest quality of reproduction possible.

If pages are missing, contact the university which granted the degree.

Some pages may have indistinct print especially if the original pages were typed with a poor typewriter ribbon or if the university sent us an inferior photocopy.

Previously copyrighted materials (journal articles, published tests, etc.) are not filmed.

Reproduction in full or in part of this microform is governed by the Canadian Copyright Act, R.S.C. 1970, c. C-30.

## AVIS

La qualité de cette microforme dépend grandement de la qualité de la thèse soumise au microfilmage. Nous avons tout fait pour assurer une qualité supérieure de reproduction.

S'il manque des pages, veuillez communiquer avec l'université qui a conféré le grade.

La qualité d'impression de certaines pages peut laisser à désirer, surtout si les pages originales ont été dactylographiées à l'aide d'un ruban usé ou si l'université nous a fait parvenir une photocopie de qualité inférieure.

Les documents qui font déjà l'objet d'un droit d'auteur (articles de revue, tests publiés, etc.) ne sont pas microfilmés.

La reproduction, même partielle, de cette microforme est soumise à la Loi canadienne sur le droit d'auteur, SRC 1970, c. C-30.

THE UNIVERSITY OF ALBERTA

FROM O,S-DIETHYL THIOLMALONATE TO  $\beta$ -HYDROXYPROPIONATE  
AND ACRYLATE DERIVATIVES.

'SYNTHETIC STUDIES OF CLOVANE-DIOL.

BY

W. M. NJUE

A THESIS

SUBMITTED TO THE FACULTY OF GRADUATE STUDIES AND RESEARCH  
IN PARTIAL FULFILLMENT OF THE REQUIREMENTS FOR THE DEGREE  
OF DOCTOR OF PHILOSOPHY

DEPARTMENT OF CHEMISTRY

EDMONTON, ALBERTA

SPRING, 1988

Permission has been granted  
to the National Library of  
Canada to microfilm this  
thesis and to lend or sell  
copies of the film.

The author (copyright owner)  
has reserved other  
publication rights, and  
neither the thesis nor  
extensive extracts from it  
may be printed or otherwise  
reproduced without his/her  
written permission.

L'autorisation a été accordée  
à la Bibliothèque nationale  
du Canada de microfilmer  
cette thèse et de prêter ou  
de vendre des exemplaires du  
film.

L'auteur (titulaire du droit  
d'auteur) se réserve les  
autres droits de publication;  
ni la thèse ni de longs  
extraits de celle-ci ne  
doivent être imprimés ou  
autrement reproduits sans son  
autorisation écrite.

ISBN: 0-315-42961-5

THE UNIVERSITY OF ALBERTA

RELEASE FORM

NAME OF AUTHOR: WILSON M. NJUE

TITLE OF THESIS:

PART I. FROM O,S-DIETHYL THIOLMALONATE TO  $\beta$ -  
HYDROXYPROPIONATE AND ACRYLATE-DERIVATIVES.

PART II. SYNTHETIC STUDIES OF CLOVANE-DIOL.

DEGREE FOR WHICH THESIS WAS PRESENTED: Ph.D.

YEAR THIS DEGREE GRANTED: 1988

Permission is hereby granted to THE UNIVERSITY OF  
ALBERTA LIBRARY to reproduce single copies of this  
thesis and to lend or sell such copies for private,  
scholarly or scientific research purposes only.

The author reserves other publication rights, and  
neither the thesis nor extensive extracts from it may  
be printed or otherwise reproduced without the author's  
written permission.

(Signed) .....

PERMANENT ADDRESS:

P.O. Box 53405  
Nairobi, Kenya

DATED MARCH 21 1988

THE UNIVERSITY OF ALBERTA  
FACULTY OF GRADUATE STUDIES AND RESEARCH

The undersigned certify that they have read, and  
recommend to the Faculty of Graduate Studies and Research,  
for acceptance, a thesis entitled

FROM O,S-DIETHYL THIOLMALONATE TO  $\beta$ -HYDROXY  
PROPIONATE AND ACRYLATE DERIVATIVES.

SYNTHETIC STUDIES OF CLOVANE-DIOL.

submitted by W. M. NJUE in partial fulfillment of the  
requirements for the degree of Doctor of Philosophy in  
Chemistry.

H.J. Liu, Supervisor

E.V. Black

George Kotowich

John M. Brumley

John H. Dill

Glenn Tolosa

External Examiner

MAR 21 1988

Date .....

To


My late parents, my wife Jane Rose,  
my son Munene and daughter Mukami.

ABSTRACT


The first part of the thesis describes a convenient method for the preparation of  $\beta$ -hydroxypropionate and acrylate derivatives using O,S-diethyl thiolmalonate (23, Part I) as a common starting material. As shown in Scheme II, Part I (p. 11), thiolester 23 was found to undergo alkylation with a variety of alkyl halides under mild conditions. Selective reduction of the thiolester group of the alkylation products gave rise to ethyl  $\beta$ -hydroxypropionates. Subsequent dehydration resulted in the formation of ethyl acrylate derivatives. The use of thiolester 23 as a convenient source of a  $\beta$ -hydroxypropionate carbanion in alkylation reactions has therefore been established in this work.

The second part of this thesis describes the preparation of bicyclic keto ester 101 (Part II), a potential synthetic precursor to clovane-diol (1, Part II), starting from 10-camphorsulfonic acid via the route outlined in Scheme XV, Part II (p. 139). Fusion of 10-camphorsulfonic acid (35) with potassium hydroxide followed by esterification gave methyl campholenate (40) which was subjected to photooxygenation to furnish enone ester 42. The methyldene group was removed by epoxidation followed by treatment with sodium hydroxide. Esterification of the

resulting carboxylic acid gave rise to keto ester 53 which was subjected to Michael addition with methyl acrylate to give keto diester 73. Subsequent Dieckmann condensation afforded diketo ester 73 which was methylated to give keto ester 78. Thioacetalization followed by Michael addition of the resulting thioacetal 92 furnished keto diester 101.



23



101

1

### ACKNOWLEDGEMENTS

It gives me great pleasure to express my utmost gratitude to my research director, Professor H.J. Liu, for his outstanding guidance, comments and encouragement throughout this work. His personal involvement and assistance during the preparation of this thesis are immensely appreciated.

I am greatly thankful to the technical staff of this department for recording all the spectra as well as micro-analyses and X-ray analyses. I greatly appreciated the proof reading of J. Nyangulu and P. McKenzie. Special thanks are extended to D. Dowhaniuk for typing the thesis. Last but not the least, I thank my wife and friends for their encouragement and emotional support during all these years.

TABLE OF CONTENTS

	<u>Page</u>
ABSTRACT.....	v
ACKNOWLEDGEMENTS.....	vii
LIST OF TABLES.....	ix
LIST OF FIGURES.....	x
LIST OF SCHEMES.....	xi
PART I. FROM O,S-DIETHYL THIOLMALONATE TO β-HYDROXYPROPIONATE AND ACRYLATE DERIVATIVES.....	1
INTRODUCTION.....	2
RESULTS AND DISCUSSION.....	17
EXPERIMENTAL.....	43
PART II. SYNTHETIC STUDIES OF CLOVANE-DIOL.....	77
INTRODUCTION.....	78
RESULTS AND DISCUSSION.....	95
EXPERIMENTAL.....	141
REFERENCES.....	178

LIST OF TABLES

	<u>Page</u>
I. Alkylation reactions of O,S-diethyl thiomalonate .....	28
II. Selective reduction of alkylation products to $\beta$ -hydroxypropionates! .....	35
III. Dehydration of ethyl $\beta$ -hydroxy- propionates!.....	41

LIST OF FIGURES

	<u>Page</u>
• Figure 1 .....	127

LIST OF SCHEMES

	<u>Page</u>
PART I	
Scheme I .....	9
Scheme II .....	11
PART II	
Scheme I .....	80
Scheme II .....	81
Scheme III .....	83
Scheme IV .....	86
Scheme V .....	88
Scheme VI .....	99
Scheme VII .....	91
Scheme VIII .....	92
Scheme IX .....	93
Scheme X .....	96
Scheme XI .....	100
Scheme XII .....	108
Scheme XIII .....	112
Scheme XIV .....	122
Scheme XV .....	139

PART I.

FROM O,S-DIETHYL THIOLMALONATE  
TO  $\beta$ -HYDROXYPROPIONATE AND ACRYLATE DERIVATIVES.

## INTRODUCTION


A thiolester is a versatile functional group which has been used in organic synthesis to facilitate the formation of carbon-carbon bonds and the preparation of a number of important functionalities. The first thiolester was reported in 1875 by Muhler<sup>1</sup> who prepared ethyl thiol-acetate from acetyl chloride and ethanethiol. In the next seventy years or so several additional methods were reported: Wallach and Bleibtreu<sup>2</sup> illustrated the formation of thiolacetates by the hydrolysis of thiolacetonilides. Obermeyer<sup>3</sup> obtained methyl thiolacetate and thiolisobutyrate by the reaction of the corresponding acid chlorides and lead methylmercaptides. Later, Wheeler<sup>4</sup> prepared ethyl thiobenzoate by reacting ethyl bromide with potassium thiobenzoate.

Very little attention was drawn to the chemical properties of thioesters until the mid-1940's when Jeger and coworkers<sup>5</sup> examined the potential use of thioesters in the modification of steroids. They showed that Raney nickel promoted reductive desulfurization of the thioester functional group into a primary alcohol. In an attempt to extend this desulfurizing action of Raney nickel to the synthesis of aldehydes, Wolfrom and Karabinos<sup>6</sup> also investigated the reduction of thio-

esters. They succeeded in the preparation of benzaldehyde and propionaldehyde from the corresponding thioesters. A couple of years later, McIntosh *et al.*<sup>7,8</sup> reexamined the reduction of thioesters and observed that, whereas the standard Raney nickel effected the conversion of thioesters to the corresponding alcohols, deactivated Raney nickel (e.g. in boiling acetone) gave rise to aldehydes as major products.


Prior to 1970, the progress in the chemistry of thioesters was rather slow.<sup>9</sup> Since then the use of thioesters has grown significantly due to their widely spread applications as useful intermediates in organic synthesis.<sup>10</sup>

In 1974 Mukayama and coworkers<sup>11</sup> prepared thioesters in high yields by treatment of trimethylsilyl sulfides with carboxylic esters in the presence of aluminum chloride (Eq. 1).



Mukayama<sup>12</sup> also showed that onium salts of aza arenes (e.g. 2-halogenated pyridinium salts 1) could be used to activate carboxylic acids (Eq. 2). The intermediates 2 thus formed reacted readily with a variety of thiols under

practically neutral conditions giving high yields of the corresponding thioesters.



In recent years, several methods have been developed for the direct transformation of carboxylic acids to the corresponding thioesters using various activating agents. *N,N*-Dimethylphosphoramidic dichloride (**3**)<sup>13</sup> and phenylphosphorodichloride (**4**)<sup>14</sup> have been shown in our laboratory to be excellent activating agents which facilitate the formation of thioesters under very mild conditions. Other activating agents successfully studied include diethylphosphorocyanide (**5**),<sup>15</sup> diphenylphosphorazidate (**6**),<sup>16</sup> *N,N*-bis(2-oxo-3-oxazolidinyl)phos-

phorodiamidic chloride (7)<sup>17</sup> and 1-fluoro-2,4,6-trinitrobenzene (8).<sup>18</sup>




3





4



5



6




8





A number of indirect methods for the preparation of thioesters have also been developed. Masamune

et al.<sup>19,20</sup> prepared S-t-butyl thioesters 9 from the corresponding acids by treatment of intermediate acid chlorides (Eq. 3) or mixed acid anhydrides (Eq. 4) with thallous 2-methylpropane-2-thiolate. In another method, Ueda and coworkers<sup>21</sup> employed S- and N-acyl derivatives (10 and 11) of 2-mercaptopbenzoxazole as intermediates.

These compounds were found to be highly reactive acylating




agents, apparently due to the electron withdrawing character of the benzoxazole ring system. Imidazolides 12 and 1,2,4-triazolides 13 were also shown to be excellent acylating agents which gave rise to thioesters in high yields upon treatment with mercaptans.<sup>22</sup>






Thiolester has emerged as a highly useful functional group in organic synthesis. Based on its interesting reactivities, a large number of synthetic methods have been developed recently. Several examples are illustrated below.

The best known example is probably the use of the hydroxy thiolester system to induce the formation of medium and large lactone rings,<sup>23-27</sup> which are present in a variety of natural products.<sup>24, 28, 29</sup> Dithioesters have been shown to undergo Dieckmann condensation under conditions substantially milder than those required for the oxygen analogues. The cyclization of S,S-diethyl dithioladipate (14),<sup>30</sup> for instance, occurred smoothly within 2 h when treated with sodium hydride in 1,2-dimethoxyethane at room temperature in the presence of a catalytic amount of ethanethiol to give a 91% yield of, S-ethyl 2-cyclopentanonecarbonylthioate (15).




14




Interestingly, unlike simple thioesters which gave rise to alcohols or aldehydes on treatment with Raney nickel,

$\beta$ -keto thioesters were found to undergo complete dealkylthiolcarbonylation (e.g. 16 + 17).<sup>30,31</sup>

Extrapolation of these results suggests that Raney nickel reduction of dithiomalonate derivatives 18 could lead to the formation of ethanol derivatives 19. This expectation was realized experimentally and S,S'-diethyl dithiomalonate 20 was used successfully as an ethanol carbanion equivalent in alkylation and Michael reactions.<sup>32,33</sup>




Another useful property of the thioester group is its high reactivity towards metal hydride reducing agents. Fujita *et al.*<sup>34</sup> observed that the highly reactive thioester 21 derived from 2-mercaptopthiazoline could be reduced to the alcohol level by sodium borohydride. It




was later found that the reduction of ordinary thioesters was equally facile and could be selectively accomplished in the presence of commonly encountered acid derivatives such as amides, esters and nitriles.<sup>35</sup>

The ease of reduction of thioester group with sodium borohydride suggests a number of interesting possibilities for its use as a latent hydroxymethyl unit in synthesis, especially when such a unit, with or without protection, can not be directly involved in a synthetic transformation. Accordingly, a convenient synthetic approach to  $\beta$ -hydroxypropionitrile and acrylonitrile derivatives has been developed involving alkylation, Michael addition or Knoevenagel-type reactions of cyanothiolacetate, as the initiating step.<sup>36,37</sup> As shown in Scheme I, the synthetic approach is greatly facilitated because of the ability of the thioester group present in the starting material to serve both as an activating group for the carbon-carbon


Scheme I



bond formation and as a convenient source of the hydroxymethyl moiety. The overall transformation can be considered as the replacement of one or two  $\alpha$ -protons of  $\beta$ -hydroxypropionitrile (or acrylonitrile) by electrophiles using cyanothiolacetate as a  $\beta$ -hydroxypropionitrile carb-anion equivalent. This method, which in essence provides a convenient means for the incorporation of a highly functionalized isopropyl unit, has been successfully applied as a key step to the synthesis of  $\alpha$ -costal (22).<sup>38</sup>




In continuation of our investigation of the application of the thiolester group as a latent hydroxymethyl unit in synthesis, we have studied the use of O,S-diethyl thiomalonate (23) as a convenient source of  $\beta$ -hydroxypropionate carbanion to facilitate the synthesis of  $\beta$ -hydroxypropionate and acrylate derivatives. As shown in Scheme II, thiomalonate 23 is expected to undergo substitution reactions with electrophiles such as alkyl halides ( $23 \rightarrow 24$ ). Selective reduction of the thioester

Scheme II


group (24 + 25) would provide a simple entry to  $\beta$ -hydroxypropionates. In case of monosubstituted  $\beta$ -hydroxypropionates, subsequent dehydration should lead to acrylate formation.

The preparation of  $\beta$ -hydroxypropionate derivatives by displacement of an  $\alpha$ -hydrogen of the parent molecule and its homologues has been the subject of considerable interest. Schlessinger and Herrmann<sup>39</sup> observed that treatment of methyl  $\beta$ -hydroxypropionate with two equivalents of lithium diisopropylamide gave rise to the corresponding dianion (26). They also showed that the dianion so generated reacted with electrophiles in a highly regioselective manner involving predominantly the

carbanion. The generality of the method is apparent from the formation of a large number of substituted  $\beta$ -hydroxypropionates 27 in high yields using various alkyl halides and dimethyl disulfide. This direct substitution



26




27


was further investigated by Fräter<sup>40,41</sup> and Seebach,<sup>42,43</sup> involving homologous  $\beta$ -hydroxypropionates. In their studies, a high degree of stereoselectivity (wherever applicable) was also observed. For example, methylation of the dianion generated from (3S)-ethyl 3-hydroxybutanoate (28) with lithium diisopropylamide gave rise to a 67% yield of (2S,3S)- (29) and (2R,3S)-ethyl 3-hydroxy-2-methylbutanoate (30) in the ratio of 19:1 in favour of the former product. The predominant formation of the



28





29





30

threo isomer has been consistently observed for a number of alkylation reactions. This method has been used in the synthesis of lavandulol (31),<sup>44</sup> a monoterpenoid alcohol, and compound 32,<sup>45</sup> the enantiomer of the pheromone isolated from smaller European bark beetle (Scolytus multistriatus).



The above alkylation procedure leading to  $\beta$ -hydroxy-propionate derivatives has the obvious advantage of being direct and stereoselective. Unfortunately, the method has the following practical limitations. The generation of a dianion requires a powerful base which is not compatible with many frequently encountered functional groups.


Furthermore, the introduction of a second substituent to the  $\alpha$ -carbon (e.g. 33  $\rightarrow$  34) was shown to be difficult. It



appears that the reaction proceeds only with methyl and allyl halides. Even with these highly reactive reagents, the yields of the products were only modest (~50%).<sup>40</sup>



Alkylation of an intact unit of acrylate has also been reported.<sup>46,47</sup> Lithium  $\alpha$ -ethoxycarbonylvinyl(hex-1-ynyl)cuprate (35) was prepared from ethyl  $\alpha$ -bromoacrylate and lithium hexynylmethylcuprate. This complex was found to undergo alkylation with allyl halides to give products

36 in modest yields (50%–70%).<sup>46</sup> This reagent, however,



has only modest reactivity and it does not undergo reactions with alkyl halides nor with benzyl bromide. In approaches to synthesis of acrylate derivatives, several acrylate equivalents have also been used successfully.

Ficini and others<sup>48–52</sup> prepared cuprates 37 from  $\alpha$ -bromoacrolein diethyl acetal (38). These copper reagents were found to react with a number of electrophiles. The addition of reagent 39 to the enone 40, for example,




39


gave rise to Michael adduct 41 in 93% yield.<sup>49</sup> Subsequent hydrolysis and oxidation led to the formation of acrylic acid derivative 42. In another approach, methyl  $\beta$ -(N,N-dimethylamino)propionate (43) was used as a latent acrylate unit.<sup>53</sup> Treatment of 43 with lithium diisopropylamide followed by alkylation of the resulting enolate ion gave rise to monosubstituted products 44. Alkylating agents used include methyl, allyl and several other primary alkyl halides. The yields were in the range of 57% to 88%. N-Methylation of  $\beta$ -aminopropionates 44 followed by base-induced elimination resulted in the formation of 2-substituted acrylates 45.

The first part of this thesis describes a new method for the preparation of  $\beta$ -hydroxypropionates and acrylate derivatives using O,S-diethyl thiolmalonate as a convenient source of masked ethyl  $\beta$ -hydroxypropionate carb-anion. This method, which has been developed on the basis of the principle outlined in Scheme II, is operationally simple and apparently general. Furthermore, it makes use of mild reaction conditions which are compatible with many

commonly encountered functional groups. These salient features warrant its broad utility in organic synthesis.




40

 $\text{Me}_3\text{SiO}^-$  $\text{CH}(\text{OC}_2\text{H}_5)_2$ 

42

43



44



45

## RESULTS AND DISCUSSION

O,S-Diethyl thiolmalonate (23), used in the present studies, was prepared from the commercially available ethyl malonyl chloride (46) by treatment with an excess of ethanethiol (1.5 equivalents) at room temperature for



23

46


24 h. The O,S-diethyl thiolmalonate (23) thus obtained was purified by distillation at 60°C/1 torr followed by flash chromatography<sup>54</sup> on silica gel using a solution of 20% ether in petroleum ether as an eluent. Pure thiolmalonate 23 was isolated in near quantitative yield and could be stored in the refrigerator for a prolonged period of time without apparent decomposition. Thiolmalonate 23 displayed absorption bands in the ir spectrum at 1742 and 1688  $\text{cm}^{-1}$  for the O- and S-ester functionalities, respectively. These characteristic absorption bands were generally observed for all the substitution products (vide infra) derived from this compound with small differences in exact positions. The nmr spectrum showed a singlet at  $\delta$  3.57 for the methylene hydrogens adjacent to the carbonyl groups. In addition, two pairs of ethyl signals

were observed, one at  $\delta$  4.20 (q, 2H,  $J = 8$  Hz) and 1.29 (t, 3H,  $J = 8$  Hz), and the other at  $\delta$  2.95 (q, 2H,  $J = 8$  Hz) and 1.28 (t, 3H,  $J = 8$  Hz). The former pair was assigned to the ethoxy moiety and the latter to the thioethoxy group. These ethyl signals were commonly observed for all the compounds prepared from 23 in the current studies. The mass spectrum of thiolmalonate 23 showed a molecular ion peak at m/z 176.0509 in accord with the required formula of  $C_7H_{12}O_3S$ .

The feasibility of the general synthetic approach to  $\beta$ -hydroxypropionate and acrylate derivatives (Scheme II) depends on the ability of thiolmalonate 23 to undergo reaction with various electrophiles such as alkyl halides. The methylene protons of thioester 23, which are adjacent to the two electron-withdrawing carbonyl groups, are expected to be strongly acidic. The acidity of 23, which remains to be determined, should be comparable to that of diethyl malonate which has a pKa value of 13.3. Indeed thiolmalonate 23 could easily be deprotonated with a variety of bases. The carbanion thus generated was found to undergo alkylation with a variety of alkyl halides. Details are described below. Also discussed are successful transformations of the alkylation products to  $\beta$ -hydroxypropionates and acrylic esters.

1. Alkylation of O,S-diethyl thiolmalonate (23)




Thiolmalonate 23 could easily be deprotonated with sodium hydride. When the anion was subjected to reactions with alkyl halides in tetrahydrofuran at temperatures ranging from 20°C to ~70°C, alkylation products were obtained in good yields. For monoalkylation, the incorporation of a simple alkyl group could be controlled by the use of stoichiometric amounts (or slight excess) of the reagents. Thus, treatment of thiolmalonate 23 with 1.1 equivalents of sodium hydride and 1.0 equivalent of methyl iodide in tetrahydrofuran at room temperature for 3-h gave rise to methyl derivative 47 in 80% yield. A small amount (11%) of dialkylation product 48 was also



obtained. The ir and nmr spectra of compound 47 showed characteristic signals (*vide supra*) for the ester and thioester groups. The nmr spectrum also displayed a quartet at δ3.65 for one hydrogen and a methyl doublet at δ 1.50, each with a coupling constant of 7.5 Hz. The structure was further confirmed by the mass spectrum showing a molecular ion peak at 190.0657. With more

reactive alkyl halides the formation of dialkylation products in small quantity could not be suppressed.

Alkylation of 23 with benzyl bromide ( $20^{\circ}\text{C}$ , 4 h) gave rise to a 72% yield of the desired product 49 along with 2% of the dialkylation product 50. Similarly, treatment of the sodium salt of 23 with methyl 3-bromopropionate gave a mixture of mono- and dialkylation products 51 and 52 in



8:1 ratio in a total yield of 74%. Compounds 49 and 51 were readily identified on the basis of the spectral data. Other than the signals generally observed for thioester and ester groups, the ir spectrum of 49 showed additional absorption bands at  $1604$  and  $1584\text{ cm}^{-1}$  for the phenyl group. In the nmr spectrum a 5H multiplet at  $\delta$   $7.22$ , a 1H triplet ( $J = 8$  Hz) at  $\delta$   $3.84$  and two 1H doublets ( $J = 8$  Hz each) at  $\delta$   $3.22$  and  $3.25$  could readily



be assigned to the phenylethyl unit. In the case of compound 51, the nmr spectrum displayed the following additional signals: a methyl signal at  $\delta$  3.68, a 3H triplet ( $J = 7$  Hz) at  $\delta$  3.65, a 2H triplet ( $J = 7$  Hz) at  $\delta$  2.38 and a 2H doublet of triplets ( $J = J' = 7$  Hz) at  $\delta$  2.24. These signals were due to the propionate unit present in the molecule. In confirmation of the structural assignment, compound 49 showed a molecular ion peak at 266.0975. The molecular ion peak of compound 51 was not detected. Instead, an ion peak appeared at m/z 231.0686. ( $M^+ - 31$ ) due to the loss of a methoxy unit.

With less reactive alkylating agents, the alkylation proceeded much slower as expected. At the same time, under the conditions applied, the dialkylation was not detected. For example, the alkylation of thiolmalonate 23 with ethyl iodide, carried out at room temperature in tetrahydrofuran gave a 90% yield of the desired ethyl derivative 53 as the sole product after 20 h. The nmr spectrum of the compound showed the presence of a n-propyl unit with a methyl triplet at  $\delta$  0.96 ( $J = 7.5$  Hz), a methylene quintet at  $\delta$  1.98 ( $J = 7.5$  Hz) and a methine triplet at  $\delta$  3.46 ( $J = 7.5$  Hz). In the mass spectrum the expected molecular ion peak was observed at m/z 204.0824. The reaction of thiolmalonate



**23** and 1-iodopropane was carried out under similar conditions. After 8 h, compound **54** was produced in 82% yield. The nmr spectrum showed signals at  $\delta$  0.92 (t, 3H,  $J = 8$  Hz), 1.35 (m, 2H,  $J = 7$  Hz), 1.90 (q, 2H,  $J = 7$  Hz) and 3.54 (t, 1H,  $J = 7$  Hz) due to a 1,1-disubstituted butane. In agreement with the structural assignment, the mass spectrum displayed a molecular ion peak at m/z 218.0974.

The alkylation of thiolmalonate **23** with isopropyl iodide was found to be extremely slow at room temperature. At refluxing temperature, however, the reaction occurred quite readily to give a 91% yield of compound **55** which showed, in the nmr spectrum, a multiplet at  $\delta$  2.48 and a doublet ( $J = 9$  Hz) at  $\delta$  3.28 integrating to one hydrogen each, and two methyl doublets ( $J = 7$  Hz each) at  $\delta$  0.96 and 0.98. In the mass spectrum, a molecular ion peak appeared at m/z 218.0978 as expected.



Sequential treatment of thiolester 23 with sodium hydride and n-octyl bromide in refluxing tetrahydrofuran for 24 h gave rise to compound 56 in 56% yield along with a 29% recovery of the starting material. When the same reaction was carried out in the presence of a catalytic amount of sodium iodide (0.1 eq.), an improved yield (84%) of 56 was obtained with complete consumption of starting material. The nmr spectrum of this compound showed a methyl triplet ( $J = 7$  Hz) at  $\delta$  0.87 and an unresolved multiplet centered at  $\delta$  1.28 for a total of eighteen protons. Other signals observed include three quartets at  $\delta$  1.90, 2.92 and 4.20, each with a coupling constant of 7 Hz and integrating to two protons, and a triplet at  $\delta$  3.54 also with a coupling constant of 7 Hz for the methine proton. The assigned structure was confirmed by the mass spectrum which showed a peak at  $m/z$  288.1760 for the molecular ion.




Dialkylation of thiolmalonate 23 was found to be equally facile. For the incorporation of two identical substituents, the reaction could be easily carried out using adequate amount of alkylating agent, and base. The conversion of 23 to dimethyl derivative 48, for example, was effected by treatment of the former compound with 3.0

equivalents of sodium hydride\* and 2.1 equivalents of methyl iodide in tetrahydrofuran for 4 h. The product 48 thus obtained in 97% yield showed characteristic absorption bands in the ir spectrum for the ester and thioester groups at  $1739\text{ cm}^{-1}$  and  $1682\text{ cm}^{-1}$ , which were generally observed for this series of compounds. Other than those characteristic signals due to ethoxy and thioethyl moieties (vide supra), the nmr spectrum displayed a singlet at  $\delta 1.48$  for the gem-dimethyl. The structure was further verified by the mass spectrum showing a molecular ion peak at m/z 204:0818. Dialkylation of thioester 23 was also carried out with ethyl iodide, n-propyl iodide, n-butyl iodide using similar reaction conditions. The expected products 57, 58 and 59 were isolated in consistently high yields (81-90%). In all these cases, the structures were readily deduced on the basis of the spectral data.\*\*


---

\* When a lesser amount of sodium hydride was used, the reaction was found to be incomplete even after an extended period of time. These results suggested that the percentage of sodium hydride used was lower than indicated.

\*\* See Experimental for details.




Reaction of thioester 23 with an alkyl dihalide could, in principle, lead to the formation of a cyclic compound. This cyclization process has been demonstrated with 1,5-diiodopentane. Treatment of 23 with sodium hydride (3.0 eq.) and 1,5-diiodopentane (1.2 eq.) in tetrahydrofuran at room temperature for 24 h gave rise to cyclic compound 60 in 56% yield along with recovered starting material (>20%) and a small amount of unidentified



products. Compound 60 showed a molecular ion peak at m/z 244.1133 in the mass spectrum and three broad multiplets centered at  $\delta$  1.48 (6H), 1.90 (2H) and 2.15 (2H) in the nmr spectrum due to the cyclohexyl hydrogens. In an



attempt to improve the yield of 60, the reaction was carried out using 4.0 equivalents of sodium hydride. In this case, the starting material was completely consumed in 30 h and the desired product was isolated in 67% yield. Analysis of the remaining material indicated the



presence of three additional products. These compounds were identified as uncyclized iodo compound 61 (10%), dialkylation product 62 (4%) and dimeric compound 63 (20%). Iodo compound 61 showed, in the mass spectrum, a molecular ion peak at  $m/z$  372.0259 and, in the nmr spectrum, a triplet at 3.18 (2H,  $J = 7$  Hz) for the methylene group bearing the iodine. On treatment with sodium hydride (2.0 equivalents) in tetrahydrofuran for 24 h, this compound was further cyclized to give an additional amount of 60 in 97% yield. The nmr

spectrum of 62 displayed a 4H triplet at  $\delta$  3.18 for the iodo methylene groups. Its structure was further confirmed by mass spectrum showing the required molecular ion peak at m/z 568.1192. A consistent molecular ion peak was also observed for the dimeric compound 63.

For the preparation of O,S-diethyl thiolmalonates possessing two non-identical substituents, the introduction of the second group could also be carried out with ease as exemplified below. Treatment of ester 53 with n-propyl iodide (1.2 eq.) and sodium hydride (1.5 eq.) in refluxing tetrahydrofuran for 2 h gave rise to the desired product 64 in 91% yield. Under similar conditions the reaction of 53 with isopropyl iodide afforded an 87% yield of compound 65. These products were readily identified spectroscopically.\*



The results obtained from alkylation reactions are further summarized in Table I.

---

\* See Experimental for details.

Table I (continued):

Entry	Alkylating Agent (Equiv.)	Base (Equiv.)	Solvent	Time (h)	Temp. °C	Products	Recovered Yield <sup>a</sup> Thiolester	
							mono	di
7	(CH <sub>3</sub> ) <sub>2</sub> CHI	NaH (1.5)	THF	20	reflux	R=(CH <sub>3</sub> ) <sub>2</sub> CH, R'=H	91	-
8	CH <sub>3</sub> (CH <sub>2</sub> ) <sub>7</sub> Br	NaH (1.3)	THF	24	reflux	R=CH <sub>3</sub> (CH <sub>2</sub> ) <sub>7</sub> , R'=H	55	-
9	CH <sub>3</sub> (CH <sub>2</sub> ) <sub>7</sub> Br	NaH (1.5)	THF	24	reflux	56	70	19
10	CH <sub>3</sub> CH <sub>2</sub> I	NaI (cat.)			r.t.	R=R'=CH <sub>3</sub> CH <sub>2</sub>	-	82
11	*CH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> I	NaH (3.0)	THF	12	r.t.	R=R'=CH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub>	-	90
12	CH <sub>3</sub> (CH <sub>2</sub> ) <sub>3</sub> I	NaH (3.0)	THF	24	r.t.	R=R'=CH <sub>3</sub> (CH <sub>2</sub> ) <sub>3</sub>	-	81
13	I(CH <sub>2</sub> ) <sub>5</sub> I	NaH (3.0)	THF	24	r.t.	R=I(CH <sub>2</sub> ) <sub>5</sub> , R',R'=(CH <sub>2</sub> ) <sub>5</sub>	13	56
						R'=H	60	20
							62	

continued... ↗


Table I (continued):

Entry	Alkylating Agent (Equiv.)	Base (Equiv.)	Time (h)	Temp. °C	Products		Recovered % Yield <sup>a</sup>		
					mono	di	mono	di	23
14	I(CH <sub>2</sub> ) <sub>5</sub> I (1.2)	NaH (4.0) THF	30	r.t.	61	60	10	67	-
15	O(CH <sub>2</sub> ) <sub>5</sub> I EtOC-CHCSET <sub>6</sub>	NaH (1.5) THF	24	r.t.	60	-	-	97	-
16	CH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> I (1.2)	NaH (1.5) THF	20	reflux	R=CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> OC-COSCH <sub>2</sub> CH <sub>3</sub> CH <sub>3</sub> CH <sub>2</sub> OC-COSCH <sub>2</sub> CH <sub>3</sub>	53	91	-	-
17	(CH <sub>3</sub> ) <sub>2</sub> CHI	NaH (1.5) THF	20	reflux	R=(CH <sub>3</sub> ) <sub>2</sub> CH	65	87	-	-

<sup>a</sup> Yields were based on the amount of starting material used.

2. Selective reduction of alkylation products to  $\beta$ -hydroxypropionates

For the preparation of  $\beta$ -hydroxypropionates via O,S-diethyl thiolmalonate derivatives, attempts were initially made to reduce the thioester group selectively to the alcohol level using sodium borohydride. Previously, it was shown in our laboratory that the thioester group could be reduced preferentially in the presence of an O-ester moiety with sodium borohydride. Unfortunately, in the present case, the reduction was found to be non-selective. Under various conditions both thioester and ester groups were invariably reduced. For example, the reduction of benzyl compound 49 with sodium borohydride in ethanol at 0°C for 14 h resulted in the formation of a ca. 3:1 mixture of  $\beta$ -hydroxypropionate 66 and  $\beta$ -hydroxythiol-propionate 67 in 70% yield along with a 20% yield of diol 68. The unexpected reduction of O-ester group with sodium



borohydride could be attributed to its enhanced reactivity due to the presence of an electron-withdrawing group.

In search of adequate methods for the required selective reduction, the use of Raney nickel as reducing agent was examined. Previously, it was shown that Raney nickel reduction of  $\beta$ -keto thioesters resulted in complete removal of the thioester group.<sup>30</sup> It was also found that similar reduction of dithiolmalonates gave rise to ethanol derivatives resulting from the removal of one thioester group and the reduction of the remaining one to the alcohol level.<sup>31</sup> To our delight however, in the present case, the thioester group of thiomalonate was reduced to the alcohol level without apparent cleavage of the carbon-carbon bond. This was probably because, unlike ketone and thioester, an O-ester functional group was insufficient to activate the carbon-carbon bond cleavage. Thus, the thioester group in thiomalonate behaved normally. More importantly, Raney nickel could serve as effective reagent for the desired conversion of thiomalonates to  $\beta$ -hydroxypropionates. In a typical experiment, thiomalonate 49 was dissolved in benzene, and W-2 Raney nickel,<sup>55</sup> which was prewashed with benzene to remove ethanol, was added. The solution was then stirred at room temperature for 4 h. This was followed by careful filtration and extensive washing of the residue with ethanol. Bulb-to-bulb distillation of the crude product gave  $\beta$ -hydroxy propionate<sup>66</sup> in 93% yield. The ir


spectrum showed characteristic absorption bands at 3445 (hydroxyl), 1732 (ester), 1580 and 1600  $\text{cm}^{-1}$  (phenyl) and the mass spectrum displayed the expected molecular ion peak at m/z 208.1100. In the nmr spectrum the aromatic hydrogen atoms appeared as a multiplet at  $\delta$  7.34 whereas the ethyl ester displayed a triplet at  $\delta$  1.22 and a quartet at  $\delta$  4.15, each with a coupling constant of 7 Hz. Two broad doublets were observed at  $\delta$  2.93 and 3.72. The former was due to the benzylic hydrogens and the latter to the methylene group bearing the hydroxyl group. A broad singlet at  $\delta$  2.22 and a multiplet at  $\delta$  2.63 could be attributed to the hydroxyl group and the methine proton respectively.

The procedure for selective reduction proved to be general. A total of twelve compounds were examined. The results are compiled in Table II. In all the cases studied, the thiolmalonate was readily reduced with Raney nickel under virtually neutral conditions to give the corresponding  $\beta$ -hydroxypropionate in greater than 79% yield.

Several observations are noteworthy. As expected, the rate of reaction is dependent on the reactivity of the Raney nickel used. When the freshly prepared Raney nickel was applied, the reaction was complete within 8 h even with highly hindered thiolmalonates such as 58, 59 and 60 (Table II, Entries 10-12). On the other hand, when Raney

nickel was stored in ethanol over a period of time prior to use, the reaction was found to be rather slow. In several cases the starting material was not completely consumed even over an extended reaction time. Although the reaction could be forced into completion by introduction of an additional amount of Raney nickel, the yield of product was generally inferior. Thus, it is desirable to carry out the reaction with freshly prepared Raney nickel.

For Raney nickel desulfurization in general, extensive washing of Raney nickel residue is essential in order to obtain a high yield of the product. In the present case, as in many other cases, ethanol was found to be a highly effective solvent for washing. Furthermore, while products with high molecular weight could be efficiently purified by bulb-to-bulb distillation, the purification of the more volatile  $\beta$ -hydroxypropionates such as 69, 70 and 72 (Table II, Entries 2, 3 and 5) was



best carried out using flash chromatography in order to avoid substantial loss of material.

**Table II.** Selective Reduction Products to  $\beta$ -Hydroxypropiionates.

Entry	Substrate	Time (h)	Product	% Yield
1	R=PhCH <sub>2</sub> , R'=H <b>49</b>	4	R=PhCH <sub>2</sub> , R'=H <b>66</b>	93
2	R=CH <sub>3</sub> , R'=H <b>47</b>	4	R=CH <sub>3</sub> , R'=H <b>69</b>	95
3	R=R'=CH <sub>3</sub> <b>48</b>	4	R=R'=CH <sub>3</sub> <b>70</b>	0
4	R=CH <sub>3</sub> OCH <sub>2</sub> CH <sub>2</sub> , R'=H <b>51</b>	4	R=CH <sub>3</sub> OCH <sub>2</sub> CH <sub>2</sub> , R'=H <b>71</b>	85
5	R=CH <sub>3</sub> CH <sub>2</sub> , R'=H <b>53</b>	4	R=CH <sub>3</sub> CH <sub>2</sub> , R'=H <b>72</b>	79
6	R=CH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> , R'=H <b>54</b>	4	R=CH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> , R'=H <b>73</b>	86
7	R=(CH <sub>3</sub> ) <sub>2</sub> CH, R'=H <b>55</b>	4	R=(CH <sub>3</sub> ) <sub>2</sub> CH, R'=H <b>74</b>	98
8	R=CH <sub>3</sub> (CH <sub>2</sub> ) <sub>7</sub> , R'=H <b>56</b>	6	R=CH <sub>3</sub> (CH <sub>2</sub> ) <sub>7</sub> , R'=H <b>75</b>	98


continued...

Table II (Continued):

Entry	Substrate	Time (h)	Product	Yield
9	$R=R'=CH_3CH_2$ 57	4	$R=R'=CH_3CH_2$ 76	98
10	$R=R'=CH_3CH_2CH_2$ 58	6	$R=R'=CH_3CH_2CH_2$ 77	95
11	$R=R'=CH_3(CH_2)_3$ 59	8	$R=R'=CH_3(CH_2)_3$ 78	91
12	$R, R'=(CH_2)_5$ 60	6	$R, R'=(CH_2)_5$ 79	96

It is also worth pointing out that a large number of  $\beta$ -hydroxypropionates did not show molecular ion peaks in the mass spectra. Instead, a predominant fragment at  $M^+ - 30$  corresponding to the loss of a  $\text{CH}_2\text{O}$  unit was observed. For example, compound 69 (Table II, Entry 2) displayed an intense fragment at  $m/z$  102.0686 instead of a molecular ion peak (expected at  $m/z$  132.0874). The loss of a  $\text{CH}_2\text{O}$  unit could be readily rationalized by invoking a McLafferty rearrangement<sup>56</sup> as shown below with compound 69.

69.



3. Transformation of  $\beta$ -hydroxypropionates to acrylate derivatives

Two methods were successfully studied for the dehydration of  $\beta$ -hydroxypropionates to give the corresponding acrylates. These are illustrated below with specific examples.


Indirectly, compound 71 was mesylated with an excess of methanesulfonyl chloride and triethylamine in dichloromethane at room temperature for 6 h. The crude product, without purification, was subjected to treatment with 1,8-diazabicyclo[5.4.0]undec-7-ene in refluxing benzene for



5 h. The acrylate 80 thus obtained in 76% yield displayed, in the ir spectrum, two ester absorption bands at 1740 and 1718  $\text{cm}^{-1}$ , as well as a double bond absorption at 1636  $\text{cm}^{-1}$ . In the nmr spectrum, the methyl ester showed a singlet at  $\delta$  3.68 and the ethyl ester a pair of mutually coupled signals at  $\delta$  1.25 and 4.20, each with a coupling constant of 7 Hz. The ethylene unit appeared as a pair of triplets ( $J = 7$  Hz) at  $\delta$  2.38 and 2.28. The

latter signal was broadened due to allylic coupling with vinyl hydrogens which appeared at  $\delta$  5.60, a doublet with a coupling constant of 1 Hz, and  $\delta$  6.20, a broad singlet.

The structural assignment was confirmed by the mass spectrum exhibiting a molecular ion peak at 186.0892. As a further example, the dehydration of compound 66 was also done via the intermediacy of mesylate 81 to give compound 82 in 96% yield. This acrylate derivative showed,



in the ir spectrum, absorption bands at 3040-3100 (aromatic), 1718 (ester carbonyl), 1636 (C=C),  $1605 \text{ cm}^{-1}$  and 1586 (phenyl), 745 and  $701 \text{ cm}^{-1}$  (monosubstituted benzene). In the nmr spectrum, the ethyl ester showed signals at  $\delta$  1.26 and 4.18, each with a coupling constant of 7 Hz. A broad singlet at  $\delta$  3.64 was due to benzylic protons. Two doublets, one at  $\delta$  5.45 ( $J = 1 \text{ Hz}$ ) and the other at  $\delta$  6.22 ( $J = 1 \text{ Hz}$ ) were attributed to the vinylic protons. The aromatic protons appeared as a multiplet at  $\delta$  7.27. The structural assignment was confirmed by the mass spectrum which showed a molecular ion peak at m/z 190.0901.

In 1971, Alexandre and Rousessac<sup>57</sup> reported a facile method for the dehydration of  $\beta$ -hydroxy ketones using di-cyclohexylcarbodiimide in the presence of cuprous chloride. This method proved to be especially useful in the preparation of unstable conjugated enones and has been extended to the dehydration of  $\beta$ -hydroxypropionitriles.<sup>37</sup> Application of this method to compound 75 gave rise to a 92% yield of compound 83 which showed



absorption bands at 1719 (ester carbonyl) and  $1625\text{ cm}^{-1}$  ( $\text{C=C}$ ) in the ir spectrum, and a molecular ion peak at  $m/z$  212.1778 in the mass spectrum. The presence of  $\alpha$ -substituted ethyl acrylate unit was evident from the nmr spectrum which showed a 3H triplet ( $J = 7\text{ Hz}$ ) at  $\delta$  1.25, a 2H quartet ( $J = 7\text{ Hz}$ ) at 4.20, a 1H doublet of doublets ( $J = J' = 1.5\text{ Hz}$ ) at  $\delta$  5.50 and a 1H doublet ( $J = 1.5\text{ Hz}$ ) at  $\delta$  6.12. The above results are further outlined in Table III.

In conclusion, a convenient method for the preparation of  $\beta$ -hydroxypropionate and acrylate derivatives has been developed using O,S-diethyl thiomalonate 23 as a

Table III. Dehydration of Ethyl  $\beta$ -Hydroxypropionates.

Entry	Alcohol	Reagent (Equiv.)	Ethyl Acrylate	Yield
1	$\text{CH}_3\text{CH}_2\overset{\text{O}}{\underset{\text{C}}{\text{  }}}\text{CHCH}_2\text{Ph}$ $\text{CH}_2\text{OH}$	(i) $\text{MsCl}^a$ (3.0) Et <sub>3</sub> N (5.0) (ii) DBU, C <sub>6</sub> H <sub>6</sub> <sup>b</sup>	$\text{CH}_3\text{CH}_2\overset{\text{O}}{\underset{\text{C}}{\text{  }}}\text{CH}_2\text{COCH}_2\text{Ph}$ $\text{CH}_2\text{O}$	96 82
2	$\text{CH}_3\text{CH}_2\overset{\text{O}}{\underset{\text{C}}{\text{  }}}\text{CHCH}_2\text{CH}_2\text{COCH}_3$ $\text{CH}_2\text{OH}$	(i) $\text{MsCl}^a$ (3.0) Et <sub>3</sub> N (5.0) (ii) DBU, C <sub>6</sub> H <sub>6</sub> <sup>b</sup>	$\text{CH}_3\text{CH}_2\overset{\text{O}}{\underset{\text{C}}{\text{  }}}\text{CH}_2\text{CH}_2\text{COCH}_3$ $\text{CH}_2\text{O}$	76 80
3	$\text{CH}_3\text{CH}_2\overset{\text{O}}{\underset{\text{C}}{\text{  }}}\text{CH}_2\text{CH}_2\text{COCH}_3$ $\text{CH}_2\text{OH}$	DCC <sup>c</sup> (2.0)	$\text{CH}_3\text{CH}_2\overset{\text{O}}{\underset{\text{C}}{\text{  }}}\text{CH}_2\text{CH}_2\text{COCH}_3$ $\text{CH}_2\text{O}$	92 83

a Dichloromethane was used as solvent.

b At reflux.

c The reaction was performed in refluxing ether in the presence of a trace amount of copper(II) chloride.

41

common starting material. The method is operationally simple and apparently general. All the transformations could be done under rather mild conditions which would not affect most of the other functionalities frequently encountered in organic synthesis.

## EXPERIMENTAL

### General

Elemental analyses were performed by the micro-analytical laboratory of this department. Infrared spectra were recorded on Perkin-Elmer model 457 or Nicolet 7-199 FT-IR spectrophotometer. Unless otherwise stated, ir samples were run as thin films. Proton nuclear magnetic resonance (nmr) spectra were recorded on a Bruker WH-200, WH-400, or AM-300 spectrometer and, except where stated, were obtained on solutions in deuteriochloroform using tetramethylsilane as internal reference.

The following abbreviations were used: s = singlet, d = doublet, t = triplet, q = quartet, m = multiplet and br = broad. Mass spectra (ms) were recorded on Kratos A.E.I. model MS-50 or MS-12 mass spectrometers. Whenever possible the progress of reaction was monitored by thin layer chromatography using one or more of the following for visualization: UV absorption by fluorescence quenching; I<sub>2</sub>-staining; (1:1) methanol/H<sub>2</sub>SO<sub>4</sub> spray with charring; acidic solution of vanillin in methanol; aq. solution of 3% phosphomolybolic acid and 0.5% ceric sulfate in 3% H<sub>2</sub>SO<sub>4</sub>. Concentration of solvent systems used in column chromatography are given by volumes, e.g.

10% ether in petroleum ether, means 10 parts of ether by volume to 90 parts of petroleum ether by volume. One or more of the following solvent systems was used:

EtOAc/petroleum ether (most generally satisfactory);<sup>58</sup> CH<sub>2</sub>Cl<sub>2</sub>/petroleum ether; EtOAc/CH<sub>2</sub>Cl<sub>2</sub>; and ether/petroleum ether (for less polar mixtures). All spray reagents were prepared and used as described by Kreibs et al.<sup>59</sup> The flash chromatography was executed according to Still et al.<sup>54</sup> and "Dry-column" flash chromatography according to Harwood.<sup>60</sup> The reactions were done under a positive pressure of dry argon.

### Materials

Ether and benzene used for reactions were freshly distilled from lithium aluminium hydride. Tetrahydrofuran and 1,2-dimethoxyethane were distilled from sodium and benzophenone. Toluene and ethyl acetate were distilled from calcium hydride. Absolute ethanol was obtained by distilling 95% ethanol over magnesium turnings in the presence of a few crystals of iodine. Argon was passed through a purification train of Fieser's solution,<sup>61</sup> concentrated sulfuric acid and potassium hydroxide pellets. The Fieser's solution was prepared by adding sodium dithionite to a cold solution of sodium hydroxide

followed by the sodium salt of anthraquinone B sulfonic acid. Merck 60GF254 silica gel was used for thin layer chromatography. Silica gel, type 60, 230-400 mesh, was used for flash chromatography and type 60, 70-230 mesh was used for gravity chromatography. Unless otherwise stated, anhydrous magnesium sulfate was used for drying organic solutions. Solutions were concentrated under reduced pressure. Commercially available sodium hydride (80% dispersion in oil) was washed several times with petroleum ether before use. Commercially available iodomethane, iodoethane, 1-iodopropane, 2-iodopropane, 1-iodobutane, benzyl bromide, methyl  $\beta$ -bromopropionate, 1,5-diiodopentane and 1-bromoocetane were used without further purification.

#### W-2 Raney nickel

A solution of sodium hydroxide (190 g) in distilled water (750 mL) was cooled in an ice-bath. Nickel-aluminium alloy (150 g) was added in small portions with stirring. During the addition, the temperature of the mixture was not allowed to rise above 25°C. After the addition, the mixture was allowed to stand at room temperature until the evolution of hydrogen stopped. The mixture was heated on a steam bath with stirring for 12 h. The solvent was decanted and the residue washed

thoroughly with distilled water until neutral. The residue was further washed with absolute ethanol ( $3 \times 250$  mL) and stored under ethanol.

#### O,S-Diethyl thiolmalonate (23)

A solution of ethanethiol (3.03 mL, 41 mmol) in tetrahydrofuran (10 mL) was added dropwise to a solution of ethyl malonyl chloride (4.74 mL, 31 mmol) in tetrahydrofuran (30 mL) at  $0^\circ\text{C}$  under an argon atmosphere. The reaction mixture was allowed to stand at room temperature for 24 h. Most of the solvent was evaporated, and the residue distilled at  $60^\circ\text{C}/1$  torr. The distillate was further purified by flash chromatography on silica gel using 20% ether in petroleum ether as eluent to afford thiolmalonate 23 (5.34 g, 96% yield): ir 1742 (ester carbonyl) and  $1648 \text{ cm}^{-1}$  (thioester carbonyl); nmr  $\delta$  1.28 (t, 3H,  $J = 8$  Hz,  $-\text{SCH}_2\text{CH}_3$ ), 1.29 (t, 3H,  $J = 8$  Hz,  $-\text{OCH}_2\text{CH}_3$ ), 2.95 (q, 2H,  $J = 8$  Hz,  $-\text{SCH}_2\text{CH}_3$ ), 3.57 (s, 2H,  $-\text{COCH}_2\text{CO}-$ ) and 4.20 (q, 2H,  $J = 8$  Hz,  $-\text{OCH}_2\text{CH}_3$ ); ms  $M^+$  176.0509 (calcd. for  $\text{C}_7\text{H}_{12}\text{O}_3\text{S}$ : 176.0507).

#### Addition of iodomethane to O,S-diethyl thiolmalonate (23)

A solution of O,S-diethyl thiolmalonate (270 mg, 1.54 mmol) in tetrahydrofuran (15 mL) was added dropwise to sodium iodide (50.6 mg, 1.06 mmol) in tetrahydrofuran (10 mL) at  $0^\circ\text{C}$ . The solution was stirred under argon for

10 min. A solution of iodomethane (0.095 mL, 1.53 mmol) in tetrahydrofuran (1.0 mL) was then added and the mixture was stirred for 3 h. The reaction was quenched with ice-cold 1 N aqueous hydrochloric acid solution and extracted with dichloromethane (3 x 20 mL). The extracts were combined, washed with saturated sodium chloride solution, dried, filtered and concentrated. Flash chromatography of the residue ~~on~~ silica gel eluting with 20% ether in petroleum ether gave the following products.

Monoalkylated product **47** as a colourless oil (233 mg, 80% yield): ir 1749 (ester carbonyl) and 1688  $\text{cm}^{-1}$  (thiolester carbonyl); nmr  $\delta$  1.25 (t, 3H,  $J = 8$  Hz,  $-\text{SCH}_2\text{CH}_3$ ), 1.26 (t, 3H,  $J = 8$  Hz,  $-\text{OCH}_2\text{CH}_3$ ) 1.50 (d, 3H,  $J = 7.5$  Hz,  $-\text{CHCH}_3$ ), 2.96 (q, 2H,  $J = 8$  Hz,  $-\text{SCH}_2\text{CH}_3$ ), 3.65 (q, 1H,  $J = 7.5$  Hz,  $-\text{CHCH}_3$ ), and 4.20 (q, 2H,  $J = 8$  Hz,  $-\text{OCH}_2\text{CH}_3$ ); ms  $M^+$  190.0657 (calcd. for  $\text{C}_8\text{H}_{14}\text{O}_3\text{S}$ : 190.0663). Anal. Calcd. for  $\text{C}_8\text{H}_{14}\text{O}_3\text{S}$ : C 50.53, H 7.37, S 16.84. Found: C 50.48, H 7.77, S 16.73. Dialkylated product **48** as a colourless oil (34.70 mg, 11% yield): ir 1739 (ester carbonyl) and 1682  $\text{cm}^{-1}$  (thiolester carbonyl); nmr  $\delta$  1.25 (t, 3H,  $J = 7$  Hz,  $-\text{SCH}_2\text{CH}_3$ ), 1.26 (t, 3H,  $J = 7$  Hz,  $-\text{OCH}_2\text{CH}_3$ ), 1.48 (s, 6H,  $-\text{C}(\text{CH}_3)_2$ ), 2.90 (q, 2H,  $J = 7$  Hz,  $-\text{SCH}_2\text{CH}_3$ ) and 4.18 (q, 2H,  $J = 7$  Hz,  $-\text{OCH}_2\text{CH}_3$ ); ms  $M^+$  204.0818 (calcd. for  $\text{C}_9\text{H}_{16}\text{O}_3\text{S}$ : 204.0820). Anal. Calcd.

for  $C_9H_{16}O_3S$ : C 52.94, H 7.89, S 15.69; Found: C 52.95, H 7.92, S 15.68.

#### Dimethylation of 23

A solution of O,S-diethyl thiolmalonate (901 mg, 5.46 mmol) in tetrahydrofuran (10 mL) was added dropwise to sodium hydride (491 mg, 16.37 mmol) in tetrahydrofuran (10 mL) at 10°C. The solution was stirred under an argon atmosphere for 20 min. Then a solution of iodomethane (0.714 mL, 11.47 mmol) in tetrahydrofuran (2 mL) was added and the mixture was stirred at room temperature for 4 h. The reaction mixture was quenched with ice-cold 1 N hydrochloric acid solution and then extracted with dichloromethane ( $3 \times 25$  mL). The extracts were combined, washed with saturated sodium chloride solution, dried, filtered and concentrated. The residue was purified by flash chromatography on silica gel, eluting with 10% ether in petroleum ether to afford adduct 48 (1.08 g, 97% yield). The spectral results (ir,  $^1H$  nmr and ms) were identical to those already obtained (vide supra).

#### Alkylation of 23 with benzyl bromide

A solution of thioester 23 (3.335 g, 18.95 mmol) in dry tetrahydrofuran (10 mL) was added slowly to a suspension of sodium hydride (770 mg, 25.67 mmol) in tetrahydrofuran (25 mL) at 0°C. The solution was stirred

under an argon atmosphere for 20 min. A solution of benzyl bromide (2.48 mL, 20.85 mmol) in tetrahydrofuran (5 mL) was added. The mixture was stirred at room temperature for 4 h. The reaction was quenched with ice-cold 1 N hydrochloric acid solution then extracted with dichloromethane (4 x 25 mL). The extracts were combined, washed with saturated sodium chloride solution, dried, filtered and concentrated. The residue was chromatographed on silica gel, eluting with 20% ether in petroleum ether to afford pure adduct 49 (3.607 g, 72%): ir 1742 (ester carbonyl), 1684 cm<sup>-1</sup> (thiolester carbonyl), 1604 and 1584 cm<sup>-1</sup> (aromatic C=C); nmr δ 1.20 (t, 3H, J = 7 Hz, -SCH<sub>2</sub>CH<sub>3</sub>), 1.22 (t, 3H, J = 7 Hz, -OCH<sub>2</sub>CH<sub>3</sub>), 2.90 (q, 2H, J = 7 Hz, -SCH<sub>2</sub>CH<sub>3</sub>), 3.22, 3.25 (both d, 1H each, J = 8 Hz each, -CH<sub>2</sub>Ph), 3.84 (t, 1H, J = 8 Hz, -COCHCO-), 4.15 (q, 2H, J = 7 Hz, -OCH<sub>2</sub>CH<sub>3</sub>) and 7.22 (m, 5H, aromatic protons); ms M<sup>+</sup> 266.0975 (calcd. for C<sub>14</sub>H<sub>18</sub>O<sub>3</sub>S, 266.0977). Anal. Calcd. for C<sub>14</sub>H<sub>18</sub>O<sub>3</sub>S: C 63.13, H 6.80, S 12.02; Found: C 63.07, H 6.80, S 11.83. Further elution with the same solvent system afforded the dibenzyl adduct 50 (166 mg, 2% yield): ir 1741 (ester carbonyl), 1684 (thiolester carbonyl), 1604 and 1586 cm<sup>-1</sup> (phenyl); nmr δ 1.22 (t, 3H, J = 7 Hz, -SCH<sub>2</sub>CH<sub>3</sub>), 1.23 (t, 3H, J = 7 Hz, -OCH<sub>2</sub>CH<sub>3</sub>), 2.92 (q, 2H, J = 7 Hz, -SCH<sub>2</sub>CH<sub>3</sub>), 3.34 (s, 4H, 2 x -CH<sub>2</sub>Ph), 4.16 (q, 2H, J = 7 Hz, -OCH<sub>2</sub>CH<sub>3</sub>) and 7.23

(m, 10H, aromatic protons); ms  $M^+$  356.1454 (calcd. for  $C_{21}H_{24}O_3S$ : 356.1446).

Further elution with 20% ether in petroleum ether afforded unreacted 23 (523 mg, 16% recovery).

Addition of methyl 3-bromopropionate to O,S-diethyl

thiolmalonate (23)

A solution of thiolester 23 (1.89 g, 10.45 mmol) in dry tetrahydrofuran (10 mL) was added slowly to a suspension of sodium hydride (424 mg, 14.13 mmol) in tetrahydrofuran (15 mL) at 0°C. The solution was stirred under an argon atmosphere for 10 min. A solution of methyl 3-bromopropionate (1.26 mL, 11.54 mmol) in tetrahydrofuran (5 mL) was added. The mixture was stirred at room temperature for 12 h. The reaction mixture was quenched with ice-cold 1 N hydrochloric acid solution then extracted with dichloromethane (3 x 25 mL). The extracts were combined, washed with saturated sodium chloride solution, dried, filtered and concentrated. The residue was chromatographed on silica gel, eluting with 20% ether in petroleum ether to afford pure adduct 51 (1.63 g, 66% yield): ir 1741 (ester carbonyl) and 1683  $\text{cm}^{-1}$  (thiolester carbonyl); nmr  $\delta$  1.25 (t, 3H,  $J = 7$  Hz,  $-\text{SCH}_2\text{CH}_3$ ), 1.26 (t, 3H,  $J = 7$  Hz,  $-\text{OCH}_2\text{CH}_3$ ), 2.24 (q, 2H,  $J = 7$  Hz,  $-\text{CHCH}_2-$ ), 2.38 (t, 2H,

$J = 7$  Hz,  $-\text{CH}_2\text{CH}_2\text{CO}-$ ), 2.92 (q, 2H,  $J = 7$  Hz,  $-\text{SCH}_2\text{CH}_3$ ), 3.65 (t, 1H,  $J = 7$  Hz,  $-\text{COCHCO}-$ ), 3.68 (s, 3H,  $-\text{OCH}_3$ ) and 4.20 (q, 2H,  $J = 7$  Hz,  $-\text{OCH}_2\text{CH}_3$ ); ms m/z 231.0686 ( $M^+ - 31$ , calcd. for  $\text{C}_{10}\text{H}_{15}\text{O}_4\text{S}$ : 231.0691) and 201.0758 ( $M^+ - 61$ , calcd. for  $\text{C}_9\text{H}_{13}\text{O}_5$ : 201.0762); Anal. Calcd. for  $\text{C}_{11}\text{H}_{18}\text{O}_5\text{S}$ : C 50.37, H 6.92, S 12.21; Found: C 50.24, H 7.04, S 11.98.

Further elution with the same solvent system afforded the dialkylated compound 52 (301 mg, 8% yield): ir 1740 (ester carbonyl) and 1684  $\text{cm}^{-1}$  (thiolester), nmr  $\delta$  1.26 (t, 6H,  $J = 7$  Hz,  $-\text{SCH}_2\text{CH}_3$ ,  $-\text{OCH}_2\text{CH}_3$ ); 1.84 (t, 4H, 7 Hz, 2 x  $-\text{CH}_2\text{CH}_2\text{CO}-$ ), 2.40 (t, 4H,  $J = 7$  Hz, 2 x  $-\text{CH}_2\text{CH}_2\text{CO}-$ ), 2.92 (q, 2H,  $J = 7$  Hz,  $-\text{SCH}_2\text{CH}_3$ ), 3.68 (s, 6H, 2 x  $-\text{OCH}_3$ ), 4.20 (q, 2H,  $J = 7$  Hz,  $-\text{OCH}_2\text{CH}_3$ ); ms m/z 317.1063 ( $M^+ - 31$ , calcd. for  $\text{C}_{14}\text{H}_{21}\text{O}_6\text{S}$ : 317.1059), and 287.1136 ( $M^+ - 61$ , calcd. for  $\text{C}_{13}\text{H}_{19}\text{O}_7$ : 287.1131). Further elution using the same solvent system afforded the unreacted thiolester 23 (221 mg, 12% recovery).

#### Addition of iodoethane to 23

A solution of thiolmalonate 23 (903 mg, 5.13 mmol) in tetrahydrofuran (5 mL) was added dropwise to a suspension of sodium hydride (184 mg, 6.12 mmol) in tetrahydrofuran (10 mL) at 0°C. The reaction mixture was stirred under an argon atmosphere for 10 min. The solution of iodoethane

(0.451 mL, 5.64 mmol) in tetrahydrofuran (5 mL) was added and the mixture stirred at room temperature for 14 h. The reaction mixture was quenched with ice cold 1 N aqueous hydrochloric acid solution and extracted with dichloromethane ( $2 \times$  20 mL). The extracts were combined, washed with saturated sodium chloride solution, dried, filtered and concentrated. Chromatography of the residue on the silica gel, eluting with 10% ether in petroleum ether gave adduct 53 (942 mg, 90% yield): ir  $1742\text{ cm}^{-1}$  (ester carbonyl) and  $1648\text{ cm}^{-1}$  (thiolester carbonyl); nmr  $\delta$  0.96 (t, 3H,  $J = 7.5\text{ Hz}$ ,  $-\text{CH}_2\text{CH}_3$ ), 1.25 (t, 3H,  $J = 8\text{ Hz}$ ,  $-\text{SCH}_2\text{CH}_3$ ), 1.26 (t, 3H,  $J = 8\text{ Hz}$ ,  $-\text{OCH}_2\text{CH}_3$ ), 1.98 (m, 2H,  $-\text{CH}_2\text{CH}_3$ ), 2.92 (q, 2H,  $J = 8\text{ Hz}$ ,  $-\text{SCH}_2\text{CH}_3$ ), 3.46 (t, 1H,  $J = 8\text{ Hz}$ ,  $-\text{COCHCO}-$ ) and 4.20 (q, 2H,  $J = 8\text{ Hz}$ ,  $-\text{OCH}_2\text{CH}_3$ ); ms  $M^+$  204.0824 (calcd. for  $\text{C}_9\text{H}_{16}\text{O}_3\text{S}$ : 204.0820); Anal. Calcd. for  $\text{C}_9\text{H}_{16}\text{O}_3\text{S}$ : C 52.92, H 7.89, S 15.68; Found: C 52.89, H 7.94, S 15.69.

#### Monoalkylation of 23 with 1-iodopropane

A solution of O,S-diethyl thiolmalonate (919 mg, 5.22 mmol) in dry tetrahydrofuran (10 mL) was added dropwise to a suspension of sodium hydride (188 mg, 6.26 mmol) in tetrahydrofuran (10 mL) at  $0^\circ\text{C}$ . The solution was stirred under an argon atmosphere until the evolution of hydrogen gas stopped. A solution of 1-iodopropane (0.560 mL, 5.74

mmol) in tetrahydrofuran (5 mL) was added and the mixture was stirred for 8 h at room temperature. The reaction mixture was quenched with ice-cold 1 N hydrochloric acid solution and extracted with dichloromethane (3 x 15 mL). The extracts were combined, washed with saturated sodium chloride solution, dried, filtered and concentrated. The crude material was chromatographed on silica gel, eluting with 10% ether in petroleum ether to afford pure adduct 54 (933 mg, 82% yield): ir 1739 (ester carbonyl) and 1685 cm<sup>-1</sup> (thiolester carbonyl); nmr δ 0.92 (t, 3H, J = 8 Hz, -(CH<sub>2</sub>)<sub>2</sub>CH<sub>3</sub>), 1.25 (t, 3H, J = 8 Hz, -SCH<sub>2</sub>CH<sub>3</sub>), 1.26 (t, 3H, J = 8 Hz, -OCH<sub>2</sub>CH<sub>3</sub>), 1.35 (m, 2H, J = 8 Hz, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>), 1.90 (m, 2H, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>), 2.92 (q, 2H, J = 8 Hz, -SCH<sub>2</sub>CH<sub>3</sub>), 3.54 (t, 1H, J = 7 Hz, -COCHCO-), and 4.20 (q, 2H, J = 8 Hz, -OCH<sub>2</sub>CH<sub>3</sub>); ms M<sup>+</sup> 218.0974 (calcd. for C<sub>10</sub>H<sub>18</sub>O<sub>2</sub>S: 218.0976). Anal. Calcd. for C<sub>10</sub>H<sub>18</sub>O<sub>3</sub>S: C 55.02, H 8.25, S 14.62; Found: C 54.82, H 8.22, S 14.58.

Further elution afforded the unreacted 23 (82 mg, 9% recovery).

#### Addition of 2-iodopropane to 23

A solution of thiolester 23 (793 mg, 4.51 mmol) in dry tetrahydrofuran (10 mL) was added dropwise to a suspension of sodium hydride (162 mg, 5.43 mmol) in

tetrahydrofuran (10 mL) at 0°C. The solution was stirred under an argon atmosphere until the evolution of hydrogen gas stopped. A solution of 2-iodopropane (0.493 mL, 4.96 mmol), in tetrahydrofuran (5 mL) was added. The mixture was heated at 50°C for 8 h. The reaction mixture was cooled to room temperature and then quenched with ice-cold hydrochloric acid solution (1 N) followed by extraction with dichloromethane (3 x 25 mL). The extracts were combined, washed with saturated sodium chloride solution, dried, filtered and concentrated. The residue was chromatographed on silica gel, eluting with 10% ether in petroleum ether to afford pure alkylation adduct 55 (615 mg, 63% yield): ir 1743 (ester carbonyl) and 1686  $\text{cm}^{-1}$  (thiol ester carbonyl); nmr  $\delta$  0.96 (d, 3H, J = 7 Hz, -CH(CH<sub>3</sub>)CH<sub>3</sub>), 0.98 (d, 3H, J = 7 Hz, -CH(CH<sub>3</sub>)CH<sub>3</sub>), 1.25 (t, 3H, J = 7 Hz, -SCH<sub>2</sub>CH<sub>3</sub>), 1.26 (t, 3H, J = 7 Hz, -OCH<sub>2</sub>CH<sub>3</sub>), 2.48 (m, 1H, -CH(CH<sub>3</sub>)<sub>2</sub>), 2.92 (q, 2H, J = 7 Hz, -SCH<sub>2</sub>CH<sub>3</sub>), 3.28 (d, 1H, J = 9 Hz, -COCHCO) and 4.20 (q, 2H, J = 7 Hz, -OCH<sub>2</sub>CH<sub>3</sub>); ms  $\ddot{\text{M}}^+$  218.0978 (calcd. for C<sub>10</sub>H<sub>18</sub>O<sub>3</sub>S: 218.0977). Anal. Calcd. for C<sub>10</sub>H<sub>18</sub>O<sub>3</sub>S: C 55.0, H 8.30, S 14.69; Found: C 54.95, H 8.29, S 14.44. Further elution with the same solvent system afforded the unreacted thiolester 23 (214 mg, 26% recovery).

Addition of 1-bromoocetane to O,S-diethyl thiolmalonate

(23)

A. Without sodium iodide

A solution of thiolester 23 (2.40 g, 13.64 mmol) in freshly distilled tetrahydrofuran (15 mL) was added slowly to a suspension of sodium hydride (522 mg, 18.40 mmol) in tetrahydrofuran (15 mL) at 0°C. The solution was stirred under an argon atmosphere for 20 min. A solution of 1-bromoocetane (2.59 mL, 14.99 mmol) in tetrahydrofuran (5 mL) was added. The mixture was heated at reflux for 24 h. The mixture was cooled, quenched with ice-cold 1 N hydrochloric acid solution and extracted with dichloromethane (4 x 25 mL). The extracts were combined, washed with saturated sodium chloride solution, dried, filtered and concentrated. The residue was chromatographed on silica gel, eluting with 10% ether in petroleum ether to afford pure adduct 56 (2.17 g, 55% yield): ir 1744 (ester carbonyl) and 1689 cm<sup>-1</sup> (thiolester carbonyl); nmr δ 0.87 (t, 3H, J = 7 Hz, -(CH<sub>2</sub>)<sub>7</sub>CH<sub>3</sub>), 1.28 (m, 18H, -OCH<sub>2</sub>CH<sub>3</sub>, -SCH<sub>2</sub>CH<sub>3</sub>, -(CH<sub>2</sub>)<sub>6</sub>CH<sub>3</sub>), 1.90 (q, 2H, J = 7 Hz, -CHCH<sub>2</sub>-), 2.92 (q, 2H, J = 7 Hz, -SCH<sub>2</sub>CH<sub>3</sub>), 3.54 (t, 1H, J = 7 Hz, -COCH<sub>2</sub>CO-), 4.20 (q, 2H, J = 7 Hz, -OCH<sub>2</sub>CH<sub>3</sub>); ms M<sup>+</sup> 288.1760 (calcd. for C<sub>15</sub>H<sub>28</sub>O<sub>3</sub>S: 288.1759). Anal. Calcd. C<sub>15</sub>H<sub>28</sub>O<sub>3</sub>S: C 62.46, H 9.78, S 11.11; Found: C 62.45, H 9.72, S 11.10.

Further elution using the same solvent system afforded unconsumed O,S-diethyl thiolmalonate (579 mg, 24% recovery).

B. With sodium iodide

A solution of O,S-diethyl thiolmalonate (458 mg, 2.60 mmol) in freshly distilled tetrahydrofuran (5 mL) was added slowly to a suspension of sodium hydride (101 mg, 3.36 mmol) in tetrahydrofuran (10 mL), 0°C. The solution was stirred under an argon atmosphere for 10 min. A solution of 1-bromoocetane (0.517 mL, 2.98 mmol) in tetrahydrofuran (5 mL) was added followed by a catalytic amount of sodium iodide (39.3 mg, 0.26 mmol). The mixture was heated at reflux for 24 h. Usual work-up and purification as in method A afforded ester 56 (525 mg, 70% yield) and unreacted 23 (86 mg, 19% recovery). The ir, <sup>1</sup>H nmr and mass spectra were identical to those reported previously (*vide supra*).

Dialkylation of O,S-diethyl thiolmalonate (23) with iodoethane

A solution of O,S-diethyl thiolmalonate (844 mg, 4.80 mmol) in dry tetrahydrofuran (10 mL) was added dropwise to a suspension of sodium hydride (460 mg, 15.33 mmol) in tetrahydrofuran (10 mL) at 0°C. The solution was stirred under an argon atmosphere until the evolution of hydrogen

stopped. Then a solution of iodoethane (1.23 mL, 15.38 mmol) in tetrahydrofuran (5 mL) was added and stirring was continued at room temperature for 6 h. The reaction mixture was quenched with ice-cold 1 N hydrochloric acid solution, extracted with dichloromethane (3 x 25 mL) and then the extracts washed with saturated sodium chloride solution. The combined extracts were dried, filtered and concentrated. Chromatography of the residue on silica gel eluting with 10% ether in petroleum ether gave compound 57 (917 mg, 82% yield). ir 1739 (ester carbonyl) and 1685  $\text{cm}^{-1}$  (thiolester carbonyl); nmr  $\delta$  0.93 (t, 6 H, J = 8 Hz, 2 x  $-\text{CH}_2\text{CH}_3$ ), 1.26 (t, 3H, J = 8 Hz,  $-\text{SCH}_2\text{CH}_3$ ), 1.26 (t, 3H, J = 8 Hz,  $-\text{OCH}_2\text{CH}_3$ ), 2.02 (q, 4H, J = 8 Hz, 2 x  $-\text{CH}_2\text{CH}_3$ ), 2.90 (q, 2H, J = 8 Hz,  $-\text{SCH}_2\text{CH}_3$ ) and 4.20 (q, 2H, J = 8 Hz,  $-\text{OCH}_2\text{CH}_3$ ); ms  $M^+$  232.1141 (calcd. for  $\text{C}_{11}\text{H}_{20}\text{O}_3\text{S}$ : 232.1133). Anal. Calcd. for  $\text{C}_{11}\text{H}_{20}\text{O}_3\text{S}$ : C 56.90, H 8.68, S 13.80; Found: C 57.05, H 8.71, S 13.45.

Dialkylation of O,S-diethyl thiomalonate (23) with 1-iodopropane

A solution of O,S-diethyl thiomalonate (745 mg, 4.23 mmol) in dry tetrahydrofuran (10 mL) was added dropwise to a suspension of sodium hydride (406 mg, 13.53 mmol) in tetrahydrofuran (10 mL) at 0°C. The solution was stirred

under an argon atmosphere for 20 min. Then a solution of 1-iodopropane (1.03 mL, 10.56 mmol) was added and stirring was continued at room temperature for 12 h. The reaction mixture was quenched with ice-cold 1 N hydrochloric acid solution and then extracted with dichloromethane (3 x 25 mL). The extracts were combined, washed with saturated sodium chloride solution, dried, filtered and concentrated. The crude material was chromatographed on silica gel, eluting with 10% ether in petroleum ether to afford pure adduct 58 (989 mg, 90% yield): ir 1739 (ester carbonyl) and 1685  $\text{cm}^{-1}$  (thioester carbonyl); nmr  $\delta$  0.92 (t, 6H,  $J = 7$  Hz,  $2 \times -(\text{CH}_2)_2\text{CH}_3$ ), 1.21 (t, 3H,  $J = 7$  Hz,  $-\text{SCH}_2\text{CH}_3$ ), 1.22 (t, 3H,  $J = 7$  Hz,  $-\text{OCH}_2\text{CH}_3$ ), 1.29 (m, 4H,  $2 \times -\text{CH}_2\text{CH}_2\text{CH}_3$ ), 1.92 (t, 4H,  $J = 7$  Hz,  $2 \times -\text{CH}_2\text{CH}_2\text{CH}_3$ ), 2.90 (q, 2H,  $J = 7$  Hz,  $-\text{SCH}_2\text{CH}_3$ ) and 4.18 (q, 2H,  $J = 7$  Hz,  $-\text{OCH}_2\text{CH}_3$ ); ms  $M^+$  260.1444 (calcd. for  $\text{C}_{13}\text{H}_{24}\text{O}_3\text{S}$ : 260.1446); Anal. Calcd. for  $\text{C}_{13}\text{H}_{24}\text{O}_3\text{S}$ : C 60.0, H 9.23, S 12.31; Found: C 59.89, H 9.16, S 12.07.

#### Dialkylation of 23 with 1-iodobutane

A solution of O,S-diethyl thiomalonate (560 mg 3.18 mmol) in dry tetrahydrofuran (5 mL) was added dropwise to a suspension of sodium hydride (305 mg, 10.17 mmol) in tetrahydrofuran (10 mL) at 0°C. The solution was stirred under a argon atmosphere until evolution of hydrogen

stopped. A solution of 1-iodobutane (0.905 mL, 7.95 mmol) in tetrahydrofuran (5 mL) was added. The mixture was stirred at room temperature for 24 h. The reaction mixture was quenched with ice-cold 1 N hydrochloric acid solution then extracted with dichloromethane (3 x 25 mL). The extracts were combined, washed with saturated sodium chloride solution, dried, filtered and concentrated. The crude product was distilled under vacuum and then chromatographed on silica gel, eluting with 10% ether in petroleum ether to afford pure adduct 59 (741 mg, 81% yield); ir 1739 (ester carbonyl) and 1686 cm<sup>-1</sup> (thioester carbonyl); nmr δ 0.92 (t, 6H, J = 7 Hz, 2 x -(CH<sub>2</sub>)<sub>3</sub>CH<sub>3</sub>), 1.25 (m, 14H, 2 x -CH<sub>2</sub>(CH<sub>2</sub>)<sub>2</sub>CH<sub>3</sub>, -OCH<sub>2</sub>CH<sub>3</sub>, -SCH<sub>2</sub>CH<sub>3</sub>), 1.92 (t, 4 H, J = 7 Hz, 2 x -CH<sub>2</sub>(CH<sub>2</sub>)<sub>2</sub>CH<sub>3</sub>), 2.90 (q, 2H, J = 7 Hz, -SCH<sub>2</sub>CH<sub>3</sub>), 4.18 (q, 2H, J = 7 Hz, -OCH<sub>2</sub>CH<sub>3</sub>); ms M<sup>+</sup> 288.1759 (calcd. for C<sub>15</sub>H<sub>28</sub>O<sub>3</sub>S: 288.1759). Anal. Calcd. for C<sub>15</sub>H<sub>28</sub>O<sub>3</sub>S: C 62.46, H 9.78, S 11.10; Found: C 62.37, H 9.74, S 10.91.

#### Alkylation of 23 with 1,5-diiodopentane

##### A. With 3 equiv. of NaH

A solution of thioester 23 (985 mg, 5.61 mmol) dry tetrahydrofuran (10 mL) was added slowly to a suspension of sodium hydride (506 mg, 16.87 mmol) in tetrahydrofuran (10 mL) at 0°C. The solution was stirred

under an argon atmosphere for 20 min. A solution of 1,5-diiodopentane (1.09 mL, 7.33 mmol) in tetrahydrofuran (10 mL) was added. The mixture was stirred at room temperature for 24 h. The reaction mixture was then quenched with ice-cold 1 N hydrochloric acid solution and extracted with dichloromethane (3 x 25 mL). The extracts were combined, washed with saturated sodium chloride solution, dried, filtered and concentrated. The residue was chromatographed on silica gel, eluting with 10% ether in petroleum ether to afford ester **60** (769 mg, 56% yield): ir 1735 (ester carbonyl) and 1687  $\text{cm}^{-1}$  (thioester carbonyl); nmr  $\delta$  1.23 (t, 3H,  $J = 7$  Hz,  $-\text{SCH}_2\text{CH}_3$ ), 1.24 (t, 3H,  $J = 7$  Hz,  $-\text{OCH}_2\text{CH}_3$ ), 1.48 (br, m, 6H, cyclohexyl protons), 1.90 (br, m, 2H, cyclohexyl protons), 2.15 (br, m, 2H, cyclohexyl protons), 2.88 (q, 2H,  $J = 7$  Hz,  $-\text{SCH}_2\text{CH}_3$ ) and 4.22 (q, 2H,  $J = 7$  Hz,  $-\text{OCH}_2\text{CH}_3$ ); ms  $M^+$  244.1133 (calcd. for  $\text{C}_{12}\text{H}_{20}\text{O}_3\text{S}$ : 244.1133). Anal. Calcd. for  $\text{C}_{12}\text{H}_{20}\text{O}_3\text{S}$ : C 59.02, H 8.20, S 13.14; Found: C 59.08, H 8.18, S 13.10.

Further elution using the same solvent system afforded iodo compound **61** (278 mg, 13% yield): ir 1735 (ester carbonyl) and 1644  $\text{cm}^{-1}$  (thioester carbonyl); nmr  $\delta$  1.23 (t, 3H,  $J = 7$  Hz,  $-\text{SCH}_2\text{CH}_3$ ), 1.24 (t, 3H,  $J = 7$  Hz,  $-\text{OCH}_2\text{CH}_3$ ), 2.90 (q, 2H,  $J = 7$  Hz,  $-\text{SCH}_2\text{CH}_3$ ), 3.18 (t, 2H,  $J = 7$  Hz,  $-\text{CH}_2\text{I}$ ), 3.55 (t, 1H,  $J = 7$  Hz,  $-\text{COCHCO}-$ ) and

4.20 (q, 2H,  $J = 7$  Hz,  $-\text{OCH}_2\text{CH}_3$ ); ms  $M^+$  372.0259 (calcd.

for  $\text{C}_{12}\text{H}_{21}\text{O}_3\text{SI}$ : 372.0258). Anal. Calcd. for

$\text{C}_{12}\text{H}_{21}\text{O}_3\text{SI}$ : C 38.71, H 5.65, S 8.60, I 34.09; Found: C

38.43, H 5.38, S 8.51, I 33.8I.

Unconsumed thiolester 23 was also recovered (194 mg, 20% recovery).

B. With 4 equiv. NaH

A solution of thiolmalonate 23 (853 mg, 4.85 mmol) in tetrahydrofuran (10 mL) was added slowly to a suspension of sodium hydride (582 mg, 19.4 mmol) in tetrahydrofuran at 0°C. The solution was stirred under an argon atmosphere for 20 min. A solution of 1,5-diiodopentane (0.792 mL, 5.33 mmol) in tetrahydrofuran (15 mL) was added. The mixture was stirred at room temperature for 30 h. Usual work up and purification as in method A.

afforded adduct 60 (793 mg, 67% yield) and iodo-adduct 61 (186 mg, 10% yield). Spectral results (ir,  $^1\text{H}$  nmr and ms) were identical to those obtained by method A (vide supra). Further elution using 20% ether in petroleum ether produced the following compounds in order of chromatographic separation. The diiodo adduct 62 (123 mg, 4% yield): ir 1736 (ester carbonyl) and 1642 (thiolester carbonyl); nmr 1.22 (t, 3H,  $J = 7$  Hz,  $-\text{SCH}_2\text{CH}_3$ ), 1.23 (t, 3H,  $J = 7$  Hz,  $-\text{OCH}_2\text{CH}_3$ ), 2.92 (q, 2H,  $J = 7$  Hz,  $-\text{SCH}_2\text{CH}_3$ ),

3.18 (t, 4H, J = 7 Hz, 2 x -OCH<sub>2</sub>CH<sub>3</sub>) and 4.20 (q, 2H, J = 7 Hz, -OCH<sub>2</sub>CH<sub>3</sub>); ms M<sup>+</sup> 567.9974 (calcd. for C<sub>17</sub>H<sub>30</sub>O<sub>3</sub>Si<sub>2</sub>: 568.0009). Adduct 63 (256 mg, 20%): ir 1736 (ester carbonyl) and 1640 cm<sup>-1</sup> (thiol ester carbonyl); nmr δ 1.20 (t, 6H, J = 7 Hz, 2 x -SCH<sub>2</sub>CH<sub>3</sub>), 1.22 (t, 6H, J = 7 Hz, 2 x -OCH<sub>2</sub>CH<sub>3</sub>), 2.85 (q, 4H, J = 7.5 Hz, 2 x -SCH<sub>2</sub>CH<sub>3</sub>), 3.57 (t, 2H, J = 7 Hz, 2 x -COCHCO-) and 4.22 (q, 4H, J = 7.5 Hz, -OCH<sub>2</sub>CH<sub>3</sub>), ms M<sup>+</sup> 420.1643 (calcd. for C<sub>19</sub>H<sub>32</sub>O<sub>6</sub>S<sub>2</sub>: 420.1640).

Reaction of iodo adduct 61 with sodium hydride

A solution of iodo compound 61 (381 mg, 1.02 mmol) in tetrahydrofuran (10 mL) was added to a suspension of sodium hydride (61.5 mg, 2.05 mmol) in tetrahydrofuran (5 mL). The solution was stirred at room temperature under an argon atmosphere for 24 h. The reaction mixture was quenched with ice-cold 1 N hydrochloric acid solution then extracted with dichloromethane (3 x 20 mL). The extracts were combined, washed with saturated sodium chloride solution, dried, filtered, and concentrated. The residue was purified by flash chromatography eluting with 10% ether in petroleum ether to afford pure ester 60 (243 mg, 97% yield). The spectral results (ir, <sup>1</sup>H nmr and ms) were identical to those already reported (vide supra).

Addition of 1-iodopropane to thiolester 53

A solution of thiolester 53 (347 mg, 1.70 mmol) in tetrahydrofuran (2 mL) was added dropwise to a suspension of sodium hydride (77 mg, 2.57 mmol) in tetrahydrofuran (10 mL) at 0°C. The reaction mixture was stirred under an argon atmosphere at room temperature for 5 min. The solution of 1-iodopropane (0.20 mL, 2.05 mmol) in tetrahydrofuran (3 mL) was added and the mixture refluxed for 20 h. The mixture was cooled then quenched with ice-cold 1 N aqueous hydrochloric acid solution and extracted with dichloromethane. The extracts were combined, washed with saturated sodium chloride solution, dried, filtered and concentrated. Bulb-to-bulb distillation of the crude product at 48°C/1 torr afforded compound 64 (382 mg, 91% yield): ir 1740 (ester carbonyl) and 1648  $\text{cm}^{-1}$  (thiolester carbonyl); nmr  $\delta$  0.83 (t, 3H, J = 7 Hz,  $-\text{CH}_3$ ), 0.95 (t, 3H, J = 7 Hz,  $-\text{CH}_3$ ), 1.22 (t, 3H, J = 7 Hz,  $-\text{SCH}_2\text{CH}_3$ ), 1.25 (t, 3H, J = 7 Hz,  $-\text{OCH}_2\text{CH}_3$ ), 1.92 (t, 2H, J = 8 Hz,  $-\text{CH}_2-$ ), 2.00 (q, 2H, J = 7 Hz,  $-\text{CH}_2\text{CH}_3$ ), 2.90 (q, 2H, J = 7 Hz,  $-\text{SCH}_2\text{CH}_3$ ) and 4.20 (q, 2H, J = 7 Hz,  $-\text{OCH}_2\text{CH}_3$ ); ms  $M^+$  246.1293 (calcd. for  $\text{C}_{12}\text{H}_{22}\text{O}_3\text{S}$ : 246.1290).

Addition of 2-iodopropane to adduct 53

A solution of thiolester 53 (402 mg, 1.97 mmol) in tetrahydrofuran (3 mL) was added dropwise to a suspension of sodium hydride (89 mg, 2.97 mmol) in tetrahydrofuran (10 mL) at 0°C. The reaction mixture was stirred under an argon atmosphere at room temperature for 5 min. A solution of 2-iodopropane (0.24 mL, 2.40 mmol) in tetrahydrofuran (3 mL) was added and the mixture refluxed for 20 h. The mixture was cooled then quenched with ice-cold 1 N aqueous hydrochloric acid solution and extracted with dichloromethane. The extracts were combined, washed with saturated sodium chloride solution, dried, filtered and concentrated. Bulb-to-bulb distillation of the crude product at 45°C/1 torr afforded compound 65 (426 mg, 87% yield): ir 1740 (ester carbonyl) and 1646  $\text{cm}^{-1}$  (thiolester carbonyl); nmr  $\delta$  0.86 (t, 3H, J = 7 Hz,  $-\text{CH}_2\text{CH}_3$ ), 0.94 (d, 3H, J = 7 Hz,  $\text{CH}_3\text{CHCH}_3$ ), 1.02 (d, 3H, J = 7 Hz,  $\text{CH}_3\text{CHCH}_3$ ), 1.25 (t, 3H, J = 7 Hz,  $-\text{SCH}_2\text{CH}_3$ ), 1.28 (t, 3H, J = 7 Hz,  $-\text{OCH}_2\text{CH}_3$ ), 1.98 (m, 1H, J = 7 Hz,  $-\text{CH}-$ ), 2.07 (m, 1H, J = 7 Hz,  $-\text{CHHCH}_3$ ), 2.36 (m, 1H, J = 7 Hz,  $-\text{CHHCH}_3$ ), 2.92 (q, 2H, J = 7 Hz,  $-\text{SCH}_2\text{CH}_3$ ) and 4.20 (q, 2H, J = 7 Hz,  $-\text{OCH}_2\text{CH}_3$ ); ms  $M^+$  246.1298 (calcd. for  $\text{C}_{12}\text{H}_{22}\text{O}_3\text{S}$ : 246.1290).

Raney nickel reduction of adduct 47

A solution of ester 47 (387 mg, 2.04 mmol) in benzene (5 mL) was added to a suspension of Raney nickel (5 mL, settled volume) in benzene (10 mL). The mixture was stirred under an argon atmosphere at room temperature for 4 h and then filtered. The residue was washed thoroughly with benzene and then absolute ethanol (3 x 4 mL). The filtrate was then concentrated. Chromatography of crude material on silica gel, eluting with 20% ether in petroleum ether afforded the hydroxy ester 69 (225 mg, 95% yield): ir 1733 (ester carbonyl) and 3440  $\text{cm}^{-1}$  (alcohol); nmr  $\delta$  1.15 (d, 3H,  $J = 8$  Hz,  $-\text{CH}_3$ ), 1.26 (t, 3H,  $J = 8$  Hz,  $-\text{CH}_2\text{CH}_3$ ), 2.67 (m, 1H,  $-\text{COCHCH}_2\text{OH}$ ), 3.70 (m, 2H,  $-\text{CH}_2\text{OH}$ ), and 4.15 (q, 2H,  $J = 8$  Hz,  $-\text{OCH}_2\text{CH}_3$ ); ms m/z 102.0686 ( $M^+$  -30; calcd. for  $\text{C}_5\text{H}_{10}\text{O}_2$ : 102.0681). Anal. Calcd. for  $\text{C}_6\text{H}_{12}\text{O}_3$ : C 54.79, H 9.87; Found: C 54.79, H 9.83.

Reduction of compound 49 with Raney nickel

To a solution of ester 49 (1.827 g, 6.87 mmol), in benzene (20 mL) was added freshly prepared Raney nickel (10 mL, settled volume) in benzene (10 mL). The mixture was stirred under argon atmosphere at room temperature for 4 h and then filtered. The residue was thoroughly washed with benzene and then ethanol (3 x 4 mL). The filtrate was then concentrated. Flash chromatography of the

residue on silica gel, eluting with 20% ether in petroleum ether followed by bulb-to-bulb distillation, at 60°C/1 torr afforded the hydroxy ester 66 (1.327 g, 93% yield): ir 3445 (hydroxyl), 1732 (ester), 1580 and 1600  $\text{cm}^{-1}$  (phenyl); nmr  $\delta$  1.22 (t, 3H,  $J = 7$  Hz,  $-\text{OCH}_2\text{CH}_3$ ), 2.22 (br, s, 1H, -OH) 2.63 (m, 1H,  $-\text{COCH}_2\text{OH}$ ), 2.93 (br, d, 2H,  $J = 7.5$  Hz,  $-\text{CH}_2\text{Ph}$ ), 3.72 (br, d,  $J = 7.5$  Hz, 2H,  $-\text{CH}_2\text{OH}$ ) and 4.15 (q, 2H,  $J = 7$  Hz,  $-\text{OCH}_2\text{CH}_3$ ); ms  $M^+$  208.1100 (calcd. for  $\text{C}_{12}\text{H}_{16}\text{O}_3$ : 208.1099). Anal. Calcd. for  $\text{C}_{12}\text{H}_{16}\text{O}_3$ : C 69.21, H 7.69; Found: C 69.23, H 7.52.

#### Reduction of adduct 48 with Raney nickel

A solution of compound 48 (424 mg, 2.08 mmol) in benzene (5 mL) was added to a suspension of Raney nickel (5 mL, settled volume) in benzene (10 mL). The mixture was stirred under an argon atmosphere at room temperature for 4 h and then filtered. The residue was thoroughly washed with benzene and then absolute ethanol ( $2 \times 5$  mL). The filtrate was then concentrated. Chromatography of the crude material on silica gel, eluting with 20% ether in petroleum ether, afforded the hydroxy ester 70 (294 mg, 97% yield): ir 3445 (alcohol) and 1736 (esters); nmr  $\delta$  1.15 (s, 6H,  $-\text{C}(\text{CH}_3)_2$ ), 1.26 (t, 3H,  $J = 7.5$  Hz,  $-\text{OCH}_2\text{CH}_3$ ), 2.22 (br, s, -OH), 2.50 (br, s,  $-\text{CH}_2\text{OH}$ ) and

4.16 (q, 2H,  $J = 7.5$  Hz,  $-\text{OCH}_2\text{CH}_3$ ); ms m/z 116.0836 ( $M^+ - 30$ , calcd. for  $\text{C}_6\text{H}_{12}\text{O}_2$ : 116.0837).

#### Reduction of ester 51

A solution of adduct 51 (1.562 g, 5.87 mmol) in benzene (5 mL) was added to a suspension of Raney nickel (8 mL, settled volume) in benzene (20 mL). The mixture was stirred under an argon atmosphere at room temperature for 4 h and then filtered. The residue was thoroughly washed with benzene (3 mL) and ethanol (4 x 5 mL). The filtrate was concentrated. Bulb-to-bulb distillation at 68°C/1 torr afforded hydroxy ester 71 (1.04 g, 85% yield): ir 3440 (hydroxyl) and 1736  $\text{cm}^{-1}$  (ester carbonyl); nmr  $\delta$  1.28 (t, 3H,  $J = 7$  Hz,  $-\text{OCH}_2\text{CH}_3$ ), 1.64 (br, s, 1H, -OH), 1.96 (q, 2H,  $J = 7$  Hz,  $-\text{CHCH}_2\text{CH}_2-$ ), 2.38 (t, 2H,  $J = 7$  Hz,  $-\text{CH}_2\text{CO}-$ ), 2.62 (m, 1H,  $-\text{COCHCH}_2\text{OH}$ ), 3.68 (s, 3H,  $-\text{OCH}_3$ ), 3.78 (br, t, 2H,  $J = 6$  Hz,  $-\text{CH}_2\text{OH}$ ) and 4.20 (q, 2H,  $J = 7$  Hz,  $-\text{OCH}_2\text{CH}_3$ ); ms m/z 173.0811 ( $M^+ - 31$ , calcd. for  $\text{C}_8\text{H}_{13}\text{O}_4$ : 173.0814). Anal. Calcd. for  $\text{C}_9\text{H}_{16}\text{O}_6$ : C 52.94, H 7.84; Found: C 53.02, H 7.89.

#### Reduction of adduct 53

A solution of adduct 53 (541 mg, 2.65 mmol) in benzene (10 mL) was added to a suspension of Raney nickel (7 mL, settled volume) in benzene. The mixture was stirred under an argon atmosphere for 4 h and then

filtered. The residue was washed thoroughly with benzene (3 mL) and ethanol (3 x 4 mL). The filtrate was then concentrated. Chromatography of the crude material on silica gel, eluting with 20% ether in petroleum ether afforded the hydroxy ester 72 (306 mg, 79% yield): ir 1729 (ester carbonyl) and 3441  $\text{cm}^{-1}$  (alcohol); nmr  $\delta$  0.97 (t, 3H, J = 8 Hz,  $-\text{CH}_2\text{CH}_3$ ), 1.28 (t, 3H, J = 8 Hz,  $-\text{OCH}_2\text{CH}_3$ ), 1.63 (m, 2H,  $-\text{CH}_2\text{CH}_3$ ), 2.10 (br, s, 1H, -OH), 2.52 (m, 1H, -CH-), 3.75 (br, d, 2H,  $-\text{CH}_2\text{OH}$ ) and 4.20 (q, 2H, J = 8 Hz,  $-\text{OCH}_2\text{CH}_3$ ); ms m/z 116.0839 ( $M^+-31$ , calcd. for  $C_6\text{H}_{12}\text{O}_2$ : 116.0834). Anal. Calcd. for  $C_7\text{H}_{14}\text{O}_3$ : C 57.51, H 9.65; Found: C 57.43, H 9.72.

#### Reduction of adduct 54

A solution of adduct 54 (347 mg, 1.59 mmol) in benzene (5 mL) was added to a suspension of Raney nickel (5 mL, settled volume) in benzene. The mixture was stirred under an argon atmosphere at room temperature for 4 h and then filtered. The residue was washed thoroughly with benzene and then ethanol (2 x 5 mL). The filtrate was then concentrated. Flash chromatography of the crude material on silica gel eluting with 20% ether in petroleum ether afforded the hydroxy ester 73 (219 mg, 86%): ir 3440 (alcohol) and 1733  $\text{cm}^{-1}$  (ester carbonyl); nmr  $\delta$  0.92 (t, 3H, J = 7 Hz,  $-(\text{CH}_2)_2\text{CH}_3$ ), 1.30 (t, 3H, J = 7 Hz,

$-\text{OCH}_2\text{CH}_3$ ), 1.56 (m, 4H,  $-(\text{CH}_2)_2\text{CH}_3$ ), 2.22 (br, s, 1H,  $-\text{OH}$ ), 2.58 (m, 1H,  $-\text{OCHCH}_2\text{OH}$ ), 3.75 (br, d, 2H,  $-\text{CH}_2\text{OH}$ ) and 4.20 (q, 2H,  $J = 7$  Hz,  $-\text{OCH}_2\text{CH}_3$ ); ms  $M^+$  160.1098 (calcd. for  $\text{C}_8\text{H}_{16}\text{O}_3$ : 160.1099). Anal. Calcd. for  $\text{C}_8\text{H}_{16}\text{O}_3$ : C 59.98, H 10.07; Found: C 59.92, H 10.11.

#### Reduction of thiolester 55

A solution of thiolester 55 (224 mg, 1.27 mmol) in benzene (5 mL) was added to a suspension of Raney nickel (4 mL, settled volume) in benzene (10 mL). The mixture was stirred under an argon atmosphere at room temperature for 4 h and then filtered. The residue was washed thoroughly with benzene and then ethanol ( $2 \times 3$  mL). The filtrate was then concentrated. Flash chromatography of the crude product on silica gel eluting with 20% ether in petroleum ether afforded the hydroxy ester 74 (153 mg, 98% yield). ir 3440 (alcohol) and  $1732 \text{ cm}^{-1}$  (ester carbonyl); nmr  $\delta$  0.96 (d, 3H,  $J = 7$  Hz,  $-\text{CH}(\text{CH}_3)\text{CH}_3$ ), 0.98 (d, 3H,  $J = 7$  Hz,  $-\text{CH}(\text{CH}_3)\text{CH}_3$ ), 1.20 (t, 3H,  $J = 7$  Hz,  $-\text{OCH}_2\text{CH}_3$ ), 2.38 (m, 1H,  $-\text{CH}(\text{CH}_3)_2$ ), 3.50 (q, 1H,  $J = 7$  Hz,  $-\text{COCHCH}_2\text{OH}$ ), 3.75 (d, 2H,  $J = 7$  Hz,  $-\text{CH}_2\text{OH}$ ) and 4.21 (q, 2H,  $J = 7$  Hz,  $-\text{OCH}_2\text{CH}_3$ ); ms  $M^+$  160.1098 (calcd. for  $\text{C}_8\text{H}_{16}\text{O}_3$ : 160.1099).

Reduction of ester 56 with Raney nickel

To a slurry of Raney nickel (7 mL, settled volume) in benzene (15 mL) was added a solution of adduct 56 (1.07 g, 3.72 mmol) in benzene (10 mL). The mixture was stirred under an argon atmosphere at room temperature for 6 h and then carefully filtered. The residue was thoroughly washed with benzene and then ethanol (2 x 4 mL). The filtrate was then concentrated. Flash chromatography of the residue on silica gel, eluting with 20% ether in petroleum ether afforded the hydroxy ester 75 (847 mg, 98%): ir 3450 (alcohol) and 1736  $\text{cm}^{-1}$  (ester carbonyl); nmr  $\delta$  0.86 (t, 3H,  $J = 7$  Hz,  $-(\text{CH}_2)_7\text{CH}_3$ ), 1.28 (m, 15H,  $-\text{OCH}_2\text{CH}_3$ ,  $-\text{CH}_2(\text{CH}_2)_6\text{CH}_3$ ), 1.62 (m, 2H,  $-\text{CHCH}_2(\text{CH}_2)_6\text{CH}_3$ ), 2.56 (m, 1H,  $J = 7$  Hz,  $-\text{COCHCH}_2\text{OH}$ ), 3.73 (br, d,  $J = 7$  Hz,  $-\text{CH}_2\text{OH}$ ) and 4.18 (q, 2H,  $J = 7$  Hz,  $-\text{OCH}_2\text{CH}_3$ ); ms  $M^+$  230.1881 (calcd. for  $\text{C}_{13}\text{H}_{26}\text{O}_3$ : 230.1882). Anal. Calcd. for  $\text{C}_{13}\text{H}_{26}\text{O}_3$ : C 60.00, H 10.07; Found: C 60.31, H 10.33.

Reduction of adduct 57 to hydroxy ester 76

A solution of adduct 57 (423 mg, 1.82 mmol) in benzene (5 mL) was added to a suspension of Raney nickel (6 mL, settled volume) in benzene. The mixture was stirred under an argon atmosphere at room temperature for 3 h and then filtered. The residue was washed thoroughly with benzene and then absolute ethanol (2 x 5 mL). The

filtrate was then concentrated. Chromatography of crude material on silica gel, eluting with 20% ether in petroleum ether afforded the hydroxy ester 76 (310 mg, 98%): ir 3440 (hydroxy) and 1728  $\text{cm}^{-1}$  (ester carbonyl); nmr  $\delta$  0.89 (t, 6H,  $J = 7.5$  Hz, 2 x  $-\text{CH}_2\text{CH}_3$ ), 1.29 (t, 3H,  $J = 7.5$  Hz,  $-\text{OCH}_2\text{CH}_3$ ), 1.62 (q, 4H,  $J = 7.5$  Hz, 2 x  $-\text{CH}_2\text{CH}_3$ ), 3.68 (s, 2H,  $-\text{CH}_2\text{OH}$ ) and 4.18 (q, 2H,  $J = 7.5$  Hz,  $-\text{OCH}_2\text{CH}_3$ ); ms m/z 144.1149 ( $M^+ - 30$ , calcd. for  $\text{C}_8\text{H}_{16}\text{O}_2$ : 144.1150). Anal. Calcd. for  $\text{C}_9\text{H}_{18}\text{O}_3$ : C 66.63, H 11.18; Found: C 66.52, H 11.13.

#### Reduction of thiolester 58 with Raney nickel

A solution of thiolester 58 (277 mg, 1.03 mmol) in benzene (5 mL) was added to a suspension of freshly prepared Raney nickel (4 mL, settled volume) in benzene (10 mL). The mixture was stirred under an argon atmosphere at room temperature for 6 h and then filtered. The residue was thoroughly washed with benzene and ethanol (2 x 5 mL). The filtrate was then concentrated. Flash chromatography of the crude product on silica gel, eluting with 20% ether in petroleum ether afforded the hydroxy ester 77 (211 mg, 95% yield): ir 1728  $\text{cm}^{-1}$  (ester carbonyl) and 3440  $\text{cm}^{-1}$  (alcohol); nmr  $\delta$  0.88 (t, 6H,  $J = 7$  Hz, 2 x  $-(\text{CH}_2)_2\text{CH}_3$ ), 2.20 (br, s, 1H,  $-\text{OH}$ ), 3.68 (br, d, 2H,  $J = 3.5$  Hz,  $-\text{CH}_2\text{OH}$ ) and 4.15 (t, 2H,  $J = 7$  Hz,

$-\text{OCH}_2\text{CH}_3$ ; ms  $M^+$  202.1566 (calcd. for  $\text{C}_{11}\text{H}_{22}\text{O}_3$ : 202.1569).

Reduction of thiolester 59 with Raney nickel

A solution of thiolester 59 (682 mg, 2.37 mmol) in benzene (6 mL) was added to a suspension of freshly prepared Raney nickel (7 mL, settled volume) in benzene (15 mL). The reaction mixture was stirred under an argon atmosphere at room temperature for 6 h and then filtered. The residue was thoroughly washed with benzene (4 mL) and ethanol (3 x 5 mL). The filtrate was concentrated. Flash chromatography of the crude product on silica gel, eluting with 20% ether in petroleum ether afforded the hydroxy ester 78 (496 mg, 91% yield): ir 3440 (hydroxy) and 1728  $\text{cm}^{-1}$  (ester carbonyl); nmr 0.86 (t, 6H,  $J = 7$  Hz,  $2 \times -(\text{CH}_2)_3\text{CH}_3$ ), 2.22 (br, s, 1H, -OH), 3.67 (br, d, 2H,  $J = 3.5$  Hz,  $-\text{CH}_2\text{OH}$ ) and 4.20 (t, 2H,  $J = 7$  Hz,  $-\text{OCH}_2\text{CH}_3$ ); ms  $M^+$  230.1881 (calcd. for  $\text{C}_{13}\text{H}_{26}\text{O}_3$ : 230.1882).

Raney nickel reduction of thiolester 60

A solution of adduct 60 (1.869 g, 4.10 mmol) in benzene (10 mL) was added to a suspension of freshly prepared Raney nickel (10 mL, settled volume) in benzene (20 mL). The reaction mixture was stirred under an argon atmosphere at room temperature for 6 h and then filtered.

carefully. The residue was thoroughly washed with benzene (2 x 3 mL) and ethanol (4 x 5 mL). The filtrate was then concentrated. Flash chromatography of the residue on silica gel, eluting with 20% ether in petroleum ether afforded pure hydroxy ester 79 (1.306 g, 96% yield): ir 1725, 1732 (ester carbonyl), 3456 and 3466  $\text{cm}^{-1}$  (alcohol); nmr 1.26 (t, 3H,  $J = 7$  Hz,  $-\text{CH}_2\text{CH}_3$ ), 1.42 (m, 10H, cyclohexyl protons), 2.16 (br, s, 1H,  $-\text{OH}$ ), 3.62 (s, 2H,  $-\text{CH}_2\text{OH}$ ) and 4.18 (q, 2H,  $J = 7$  Hz,  $-\text{OCH}_2\text{CH}_3$ ); ms  $M^+$  186.1255 (calcd. for  $\text{C}_{10}\text{H}_{18}\text{O}_3$ : 186.1256). Anal. Calcd. for  $\text{C}_{10}\text{H}_{18}\text{O}_3$ : C 64.49, H 9.74; Found: C 64.44, H 9.63.

#### Dehydration of hydroxy ester 71 via mesylate

To a solution of hydroxy ester 71 (846 mg, 4.15 mmol) in freshly distilled dichloromethane (20 mL) was added methanesulfonyl chloride (0.963 mL, 12.44 mmol) and the mixture was stirred for 10 min at 0°C under an argon atmosphere. Triethylamine (2.89 mL, 20.73 mmol) was added slowly via a syringe and the mixture continued stirring for 6 h at room temperature. The reaction mixture was concentrated under reduced pressure and resulting yellow solids were redissolved in benzene (25 mL). 1,8-Diaza-bicyclo[5.4.0]undec-7-ene (1.55 mL) was added and the mixture heated at reflux under an argon atmosphere for 5 h. The mixture was then cooled at room temperature,

acidified with 1 N hydrochloric acid solution and extracted with dichloromethane ( $4 \times 25$  mL). The extracts were combined and washed successfully with aqueous saturated sodium carbonate solution and saturated sodium chloride solution. The combined extract was dried, filtered, and concentrated. The crude product was purified by flash chromatography on silica gel (10% ether in petroleum ether as eluent) to give olefin 80 (585 mg, 76% yield): ir 1740 and 1718 (ester carbonyls),  $1635\text{ cm}^{-1}$  ( $\text{C}=\text{C}$ ); nmr  $\delta$  1.25 (t, 3H,  $J = 7$  Hz,  $-\text{OCH}_2\text{CH}_3$ ), 2.28 (br, t, 2H,  $J = 7$  Hz,  $-\text{CH}_2\text{C}=\text{}$ ), 2.38 (t, 2H,  $J = 7$  Hz,  $-\text{CH}_2\text{CH}_2\text{CO}-$ ), 3.68 (s, 3H,  $-\text{OCH}_3$ ), 4.20 (q, 2H,  $J = 7$  Hz,  $-\text{OCH}_2\text{CH}_3$ ), 5.60 (d, 1H,  $J = 1$  Hz,  $=\text{CHH}$ ) and 6.20 (br, s, 1H,  $=\text{CHH}$ ); ms  $M^+$  186.0892 (calcd. for  $\text{C}_9\text{H}_{14}\text{O}_4$ : 186.0892). Anal. Calcd. for  $\text{C}_9\text{H}_{14}\text{O}_4$ : C 58.06, H 7.53; Found: C 57.93, H 7.59.

#### Dehydration of hydroxy ester 66 via mesylate

To a solution of hydroxy ester 66 (1.027 g, 4.94 mmol) in freshly distilled dichloromethane (20 mL) was added methanesulfonyl chloride (1.15 mL, 14.86 mmol) and the mixture stirred for 10 min at 0°C under an argon atmosphere. Triethylamine (3.44 mL, 24.68 mmol) was added slowly via a syringe and the mixture was stirred for 6 h at room temperature. The reaction mixture was

then concentrated under reduced pressure and the resulting yellow solid was redissolved in benzene (25 mL). 1,8-Diazabicyclo[5.4.0]undec-7-ene (2.95 mL) was added and the mixture heated at reflux under an argon atmosphere for 5 h. The mixture was then cooled to room temperature, poured into 1 N hydrochloric acid solution and extracted with dichloromethane (3 x 25 mL). The extracts were combined and washed successfully with aqueous saturated sodium carbonate solution and saturated sodium chloride solution. The combined extract was dried, filtered, and concentrated. The residue was purified by flash chromatography on silica gel (10% ether in petroleum ether) to give olefin 82 (901 mg, 96% yield): ir 3040-3100 (aromatic), 1718 (ester carbonyl), 1636 (C=C), 1605, 1586 (phenyl), 745 and 701  $\text{cm}^{-1}$  (monosubstituted benzene); nmr  $\delta$  1.26 (t, 3H,  $J = 7$  Hz,  $-\text{OCH}_2\text{CH}_3$ ), 3.64 (br, s, 2H,  $-\text{CH}_2\text{Ph}$ ), 4.18 (q, 2H,  $J = 7$  Hz;  $-\text{OCH}_2\text{CH}_3$ ), 5.45 (d, 1H,  $J = 1$  Hz,  $=\text{CHH}$ ), 6.22 (d, 1H,  $J = 1$  Hz,  $=\text{CHH}$ ) and 7.27 (m, 5H, aromatic protons); ms  $M^+$  190.0994 (calcd. for  $\text{C}_{12}\text{H}_{14}\text{O}_2$ : 190.0994).

Direct dehydration of hydroxy ester 75 by using dicyclohexylcarbodiimide

A solution of hydroxy ester 76 (713 mg, 3.10 mmol), dicyclohexylcarbodiimide (1.28 g, 6.20 mmol) and a


catalytic amount of copper(I) chloride (5 mg) in dry ether (20 mL) was heated at reflux for 12 h under an argon atmosphere. The mixture was then cooled to room temperature, diluted with ether and filtered. Concentration of the filtrate, followed by purification of the crude product by flash chromatography on silica gel using 20% ether in petroleum ether as an eluent gave olefin 83 (609 mg, 92% yield) as a colourless oil: ir 1719 (ester carbonyl) and  $1625\text{ cm}^{-1}$  ( $\text{C}=\text{C}$  stretching); nmr  $\delta$  0.86 (t, 3H,  $J = 7$  Hz,  $-\text{CH}_3$ ), 1.26 (t, 3H,  $J = 7$  Hz,  $-\text{OCH}_2\text{CH}_3$ ), 2.25 (br, t, 2H,  $J = 7$  Hz,  $-\text{CH}_2-\overset{\text{C}}{=}$ ), 4.20 (q, 2H,  $J = 7$  Hz,  $-\text{OCH}_2\text{CH}_3$ ), 5.50 (dd, 1H,  $J = J' = 1.5$  Hz,  $=\text{CHH}$ ), 6.12 (d, 1H,  $J = 1.5$  Hz,  $=\text{CHH}$ ); ms  $M^+$  212.1778 (calcd. for  $\text{C}_{13}\text{H}_{24}\text{O}_2$ : 212.1776).

**PART II.**

**SYNTHETIC STUDIES OF CLOVANE-DIOL.**

## INTRODUCTION

(+)-Clovane- $2\beta,9\alpha$ -diol (4,4,8-trimethyltricyclo-[6.3.1.0<sup>1,5</sup>]dodecane- $2\beta,9\alpha$ -diol) (1) is widely distributed in nature. This compound along with (-)-caryophyllene (2) and (-)-isocaryophyllene (3), were found to be the main constituents of the oil of cloves (Eugenia caryophyllata).<sup>62</sup> These sesquiterpenes were also found to occur in African capaiba oil (Oxystigma manii Harms)<sup>63</sup> and in French lavender oil.<sup>64</sup> Clovane- $2\beta,9\alpha$ -diol (1) was reported to occur also in essential oils of Mentha piperita.<sup>65</sup> Dev. and coworkers<sup>66</sup> have isolated this diol.




together with another complex blend of sesquiterpenes from oleoresin of Dipterocarpus pilosus, a tall tree occurring gregariously in Assam, India. In 1979 they also noted the presence of compound 1 in the oleoresin from Hardwickia pinnata,<sup>67</sup> a large handsome tree growing wildly in evergreen forests of Western Ghats of India. The tree, on tapping, yields in large quantities a dark or reddish brown oleoresin. Other natural sources of clovane-diol



(1) include Salvia canariensis,<sup>68</sup> the Brazilian shrub Bacharris megapotamica spren.,<sup>69</sup> and several species of the genus Viguiera<sup>70,71</sup> such as linearis, excelsa, oaxama, and hypagyrea.

The structure of clovane-diol (1) was first elucidated by Barton and coworkers mainly by chemical degradation according to Scheme I.<sup>72</sup> The assigned structure was later confirmed by spectroscopic methods.<sup>66,69</sup>

Barton and coworkers also proposed the biosynthesis of clovane-diol (1) as illustrated in Scheme II. This compound is believed to be produced biosynthetically from farnesyl pyrophosphate (4) via the intermediacy of caryophyllene (2) and the corresponding oxide 5.


The conversion of caryophyllene oxide (5),<sup>72,75</sup> both from natural sources and from epoxidation of caryophyllene (2),<sup>65,73</sup> to clovane-diol (1) has been realized chemically.<sup>74,76</sup> The former compound on mild acid


**Scheme I**

Scheme I (cont'd)Scheme II

treatment resulted in the rearrangement to give 1. Based on this finding, the configuration of compound 1 was deduced.

Although there has not been a direct attempt reported for the synthesis of clovane-diol (1), several formal synthesis via caryophyllene (2) have appeared. In the first synthesis of caryophyllene (2) in racemic form (Scheme III) by Corey and coworkers<sup>77,78</sup> in 1963, bicyclic ketone 6, readily obtained by irradiation of a mixture of 2-cyclohexenone and isobutylene, was chosen as the key intermediate. Carbomethylation followed by methylation gave rise to keto ester 7, which was then treated with the lithio salt of propargyl aldehyde dimethyl acetal to give alcohol 8. Hydrogenation followed by oxidation in aqueous acid resulted in the formation of lactone 9. This compound was subjected to Dieckmann condensation and the resulting keto ester 10 was hydrolyzed and decarboxylated to give ketone alcohol 11. Raney nickel reduction of 11 gave a pair of epimeric alcohols 12 and 13 in ca. 1:1 ratio. Upon exposure to the basic conditions, the tosylate of the latter alcohol 13 underwent a 1,3-glycol type rearrangement to give the desired enone 14, which was further epimerized to the trans-isomer 15. Wittig reaction of 15 led to the completion of the first total synthesis of ( $\pm$ )-caryophyllene (2).

**Scheme III**


Scheme III (cont'd)

- a.  $h\nu$ , pentane,  $-40^\circ C$ . b.  $(CH_3O)_2CO$ , NaH, dioxane. c. MeI, NaH, dioxane. d.  $Li^+ - CCCH(OCH_3)_2$ , THF,  $0^\circ C$ . e.  $H_2$ , 5% Pd-C,  $CH_3OH$ .
- f.  $CrO_3$ , aq.  $CH_3CO_2H$ . g.  $CH_3SOCH_2^- Na^+$ , DMSO. h. aq. NaOH,  $40^\circ C$ ; Py,  $110 - 115^\circ C$ . i. Ra-Ni,  $H_2$ . j. p-TsCl, Py,  $22^\circ C$ .
- k.  $Na^+ - CH_2SOCH_3$ , DMSO,  $25^\circ C$ . l.  $t-BuO^- Na^+$ , DMSO,  $25^\circ C$ .
- m.  $CH_2=P(Ph)_3$ , DMSO, R. T.



Some twenty years later the use of compound 7 in the synthesis of caryophyllene (2) and isocaryophyllene (3) was reinvestigated by Oishi and coworkers.<sup>79</sup> By a sequence of reactions including decarboxylation, enol acetate formation, ozonolysis, side chain extension and modification according to Scheme IV, a mixture of two isomeric olefins 16 and 17 was obtained. The former was converted to ( $\pm$ )-caryophyllene (2) via two cyclization reactions; the first one to induce the carbon-sulfur bond formation (16  $\rightarrow$  18) and the second one to effect the formation of the required carbocyclic system (19 20). Application of the same synthetic sequence to the *cis*-isomer 17 resulted in the synthesis of ( $\pm$ )-isocaryophyllene (3). These authors also devised a new strategy for the synthesis of compound 25 using cyclobutene 21 as the starting material (Scheme V). This compound was subjected to Michael addition with ethyl (phenylsulfonyl)acetate. Subsequent modification of the functionalities of the adduct 22 afforded sulfone 23 which was alkylated with chloride 24 to give compound 25, the desired synthetic precursor of caryophyllene.

In 1974, Bertland and Gras<sup>80</sup> made use of the photocyclization of optically active allene 26 and dimethyl ketene as the key step in the synthesis of (+)-isocaryophyllene. Compound 26 was prepared in five steps from


Scheme IV



**Scheme IV (cont'd).**



- a. 1 N NaOH, 35 - 45°C. b.  $\text{CH}_3\text{COOCH}=\text{C}(\text{CH}_3)_2$ , PhH, reflux .  
 c.  $\text{O}_3$ ,  $\text{CH}_3\text{OH}$ , -78°C. d. BuLi, THF, 0°C to R.T. e.  $(\text{COCl})_2$ ,  
 benzene, R.T. to 65°C. f. 9 - Ph(NH $\text{CH}_3$ )(SCH $_2$ CH $_2$ CN),  
 $\text{K}_2\text{CO}_3$ , THF, 0°C. g.  $(\text{CH}_3)_2\text{S}$ ,  $\text{BF}_3 \cdot \text{Et}_2\text{O}$ ,  $\text{Ac}_2\text{O}$ , THF, 0°C.  
 h.  $\text{K}_2\text{CO}_3$ ,  $\text{CH}_3\text{OH}$ . i.  $\text{CH}_3\text{SO}_2\text{Cl}$ ,  $\text{Et}_3\text{N}$ ,  $\text{CH}_2\text{Cl}_2$ , 0°C.  
 j. t-BuO $^-$ K $^+$ , t-BuOH. k.  $\text{NaIO}_4$ ,  $\text{CH}_3\text{OH}$ , R.T. l. LDA, THF,  
 HMPA, -78°C. m. Al - Hg,  $\text{CH}_3\text{OH}$  -  $\text{H}_2\text{O}$ . n. t-BuO $^-$ K $^+$ , t-BuOH.  
 o.  $\text{CH}_2=\text{P}(\text{Ph})_3$ , THF.

**Scheme V**


- a.  $\text{EtO}_2\text{CCH}_2\text{SO}_2\text{Ph}$ ,  $\text{t-BuOK}$ ,  $\text{THF}$ , R.T. b.  $2\%$   $\text{KOH}$ ,  $\text{H}_2\text{O} - \text{CH}_3\text{CH}_2\text{OH}$ , reflux ;  $\text{Py}$ , reflux . c.  $\text{K}_2\text{CO}_3$ ,  $(\text{CH}_3)_2\text{SO}_4$ , acetone, reflux . d.  $\text{LiAlH}_4$ ,  $\text{Et}_2\text{O}$ . e.  $\text{t-Bu}(\text{CH}_3)_2\text{SiCl}$ , imidazole,  $\text{DMF}$ , R.T. f. 24 ,  $\text{HMPA}$ ,  $n\text{-BuLi}$ ,  $\text{THF}$ . g.  $n\text{-Bu}_4\text{NF}$ ,  $\text{THF}$ . h.  $5\%$   $\text{Na} - \text{Hg}$ ,  $\text{NaH}_2\text{PO}_4$ . i. Jones reagent,  $0^\circ\text{C}$ .

1,5-octadiene as shown in Scheme VI. The deoxygenation of photo-adduct 27 via the corresponding alcohol gave a mixture of isomeric olefins 28 and 29 in 35:65 ratio. The former was converted to the latter by treatment with carbowax at 150°C. Hydroboration of compound 29 followed by oxidation resulted in the formation of ketone 30 which was pyrolyzed to give enone 31. Wittig reaction of this compound afforded the enantiomer of the natural isocaryophyllene (3).

A very similar approach (Scheme VII) has also been used by an Indian group<sup>81</sup> in their synthesis of racemic isocaryophyllene (3). More recently, the same compound was prepared by McMurry and Miller<sup>82</sup> starting from ethyl geranylacetate (32). As shown in Scheme VIII the synthetic approach is highly efficient. Unfortunately, the trans double bond required for caryophyllene could not be retained as intended during the key coupling reaction of 33.

In approaches to the synthesis of (+)-clove-diol (1), campholenic acid (34) presents itself as a highly attractive potential precursor. As can be seen in Scheme IX, this chiral substrate, which is readily accessible from the ammonium salt of (-)-10-camphorsulfonic acid (35) by fusion with potassium hydroxide pellets,<sup>83</sup> embodies nicely the complete A ring and part of B' ring.

Scheme VI




a.  $\text{CHCl}_3$ ,  $\text{Na}^+\text{:OCH}_3$ . b. Li,  $t\text{-BuOH}$ . c.  $t\text{-BuO}^- \text{K}^+$ , pentane,  $\text{CHBr}_3$ ,  $-20^\circ\text{C}$ .


d.  $n\text{-BuLi}$ ,  $-40^\circ\text{C}$ . e.  $\text{LiAlH}_4$ . f.  $p\text{-TsCl}$ , benzene. g.  $\text{LiAlH}_4$ , THF, reflux.

h. carbowax 20M,  $150^\circ\text{C}$ . i.  $\text{B}_2\text{H}_6$ ;  $\text{H}_2\text{O}_2$ , NaOH. j.  $\text{CrO}_3$ . k.  $360^\circ\text{C}$ .

l.  $\text{Ph}_3\text{P}=\text{CH}_2$ , DMSO.

Scheme VII


- a.  $\text{CBr}_4$ ,  $\text{CH}_3\text{Li}$ ,  $-65^\circ\text{C}$ . b.  $\text{B}_2\text{H}_6$  - THF;  $\text{CrO}_3$ . c.  $(\text{CH}_3)_2\text{C}=\text{CH}_2$ ,  $\text{h}\nu$ ,  
 $-40$  –  $-60^\circ\text{C}$ , pentane. d.  $\text{CH}_2\text{I}_2$ ,  $\text{Zn}$  -  $\text{Cu}$ . e.  $t\text{-BuO}^- \text{K}^+$ ,  $\text{DMSO}$ ,  $400^\circ\text{C}$ .  
f.  $\text{Ph}_3\text{P}=\text{CH}_2$ ,  $\text{DMSO}$ .

**Scheme IX**

(bold-faced portion) of the target compound with the correct chirality. Furthermore, the functionalities of this molecule are situated adequately at the strategic positions facilitating the construction of the required framework. Based on this strategy, a project directed towards the total synthesis of natural clovane-diol (**1**) was initiated in our laboratory several years ago.

The second part of this thesis describes the synthesis, from camphorsulfonic acid, of compound **101**, a potential precursor for clovane-diol (**1**).

94




101

## RESULTS AND DISCUSSION

In our approach towards the synthesis of clovane-diol (1), a bicyclic keto ester 36 was visualized as the key intermediate. This compound represents the AB ring system of the target molecule and contains appropriate functionalities for further manipulation. From a retrosynthetic analysis (Scheme X) it was recognized that keto ester 36 could arise from diketo ester 37 by methylation and deoxygenation. The latter compound in turn could be prepared by Dieckmann condensation of diester 38. The required propionate side chain could be installed by Michael reaction on compound 39. This compound could arise from 10-camphorsulfonic acid (35) via campholenic acid (34) by photooxygenation (34 → 39).

10-Camphorsulfonic acid is readily available in levorotatory, dextrorotatory and racemic forms. For synthesis of clovane-diol (1) with the natural configuration, L-10-camphorsulfonic acid is required as starting material. At the outset of the present studies, this compound was about four times more expensive than the corresponding racemate and yet was not immediately available in our laboratory. Consequently, for preliminary exploration, the results of which are discussed below, the racemic form was used.

**Scheme X**


When the sodium salt of ( $\pm$ )-10-camphorsulfonic acid (35) was fused with potassium hydroxide, ( $\pm$ )-campholenic acid (34) was produced in 78% yield. Esterification of 34 using potassium carbonate and methyl iodide in acetone gave the corresponding ester 40 in 98% yield. The



40


structure of the ester was readily established from its spectral data. The ir spectrum showed the ester absorption band at  $1748\text{ cm}^{-1}$ . In the nmr spectrum, the singlets at  $\delta$  3.68, 1.62, 0.98 and 0.78 were assigned to the methyl ester, vinylic methyl and gem-dimethyl respectively. The broad multiplet at  $\delta$  5.22 was readily assigned to the vinylic proton. The mass spectrum displayed a molecular ion peak at  $m/z$  182.1305 which was consistent with the molecular formula  $C_{11}H_{18}O_2$  which was further confirmed by the elemental analysis.

Photooxygenation of ester 40 in methanol using methylene blue as a photosensitizer<sup>84</sup> followed by reduction of the resulting hydroperoxides with sodium borohydride gave an epimeric mixture of allylic alcohols 41 in 66% yield. In the ir spectrum characteristic



absorption bands were observed at 3440 (alcohol), 3106, 1655 (olefin) and 1748 cm<sup>-1</sup> (ester). The ratio of the two epimeric alcohols was found to be approximately 7:3 as determined by hmr spectrum which displayed two pairs of methyl singlets, one at δ 1.12 and 0.82 and the other at δ 1.02 and 0.92. In agreement with the structural assignment, broad singlets at δ 5.12 and 4.92 for the vinylic protons and a singlet at 3.66 for the methyl ester were also observed. The methine proton adjacent to the hydroxyl group appeared at δ 4.95 as a multiplet. The molecular ion peak at m/z 198.1232 in the mass spectrum confirmed the chemical formula of C<sub>14</sub>H<sub>24</sub>O<sub>3</sub>.

Swern oxidation<sup>85</sup> of the mixture of the epimeric alcohols 41 with oxalyl chloride, dimethyl sulfoxide, and triethylamine gave enone 42 in 85% yield. The ir spectrum




showed absorptions at  $1748\text{ cm}^{-1}$  for the ester group,  $1729\text{ cm}^{-1}$  for the unsaturated ketone and  $1634\text{ cm}^{-1}$  for the carbon-carbon double bond. The nmr spectrum displayed two broad singlets at  $\delta$  5.98 and 5.20 for the enone protons. Two doublets of doublets appeared at  $\delta$  2.54 (1H,  $J = 18$ ,  $J' = 7\text{ Hz}$ ) and 2.06 (1H,  $J = 18$ ,  $J' = 11\text{ Hz}$ ) due to protons  $\alpha$  to the ketone carbonyl. The singlets at  $\delta$  1.22 and 1.00 were due to the gem-dimethyl group. The molecular formula  $C_{11}H_{16}O_3$  was verified by the presence of a molecular ion peak at  $m/z$  196.1305 in the mass spectrum and by the elemental analysis.

Later, it was also found that enone 42 could be prepared in one step from 40 using the procedure of Mihelich and Eickhoff.<sup>86</sup> Ester 40 was photoxxygenated in dichloromethane in the presence of pyridine, acetic anhydride and 4-dimethylaminopyridine using 5,10,15,20-tetraphenyl-


21H,23H-porphine (TPP) as a photosensitizer. Under these conditions enone 42 was obtained directly in 76% yield.

As shown in Scheme XI the formation of 42 could be rationalized by invoking the intermediacy of a peracetate which gave rise to the observed product upon elimination of a molecule of acetic acid.


With enone 42 in hand we set out to investigate the incorporation of a propionate unit in order to prepare a compound of type 38. The Michael addition of enone 42 to

Scheme XI

ethyl acrylate was examined first under a variety of conditions. Unfortunately, in no case was the desired Michael adduct formed and the only product observed was compound 43 resulting from dimerization of the starting material via a Michael reaction. The structure of the dimer was readily identified. The ir spectrum showed a broad carbonyl band at  $1734 \text{ cm}^{-1}$  and an absorption at  $1638 \text{ cm}^{-1}$  indicative of an olefin. The nmr spectrum displayed two one-proton singlets at  $\delta$  5.98 and 5.28 characteristic



43




44

of the geminal protons of the methyldene group. The methoxy groups appeared at  $\delta$  3.76 and 3.72 as singlets. Four additional methyl singlets were observed at  $\delta$  1.26, 1.18, 1.03 and 0.96. The mass spectrum showed a molecular ion peak at  $m/z$  392.2646 consistent with molecular formula  $\text{C}_{22}\text{H}_{32}\text{O}_6$ . Attempted alkylation using enamine 44, which was prepared by treatment of enone 40 with pyrrolidine and a catalytic amount of p-toluenesulfonic acid in refluxing benzene, was equally unsuccessful, resulting in the formation of a complex mixture which could not be characterized.



In the alternative approach to the preparation of compound 38, formation of enol ester 45 was attempted. Claisen rearrangement of this compound in the presence of an alcohol is expected to give the desired diester 38. Thus, enone 42 was treated with sodium acetate in acryloyl chloride at reflux for 24 h. Three compounds, two neutral

and one acidic, were produced. The neutral compounds were readily identified as the desired enol ester **45** (18% yield) and the corresponding chloride **46** (9% yield). The



former showed, in the ir spectrum, ester absorptions at  $1737\text{ cm}^{-1}$  and double bonds at  $1630$  and  $1600\text{ cm}^{-1}$ . The nmr spectrum displayed signals at  $\delta$   $6.54$  (dd,  $H_a$ ,  $J = 18\text{ Hz}$ ,  $J' = 2\text{ Hz}$ ),  $6.23$  (dd,  $H_b$ ,  $J = 18\text{ Hz}$ ,  $J' = 11\text{ Hz}$ ) and  $5.97$  (dd,  $H_c$ ,  $J = 11\text{ Hz}$ ,  $J' = 2\text{ Hz}$ ) for the vinylic protons of the  $\alpha,\beta$ -unsaturated ester. The other vinylic protons appeared at  $\delta$   $4.92$  ( $H_d$ ),  $4.73$  ( $H_e$ ) and  $5.92$  (br.,  $H_f$ ) as singlets. The singlets at  $\delta$   $1.22$  and  $1.04$  were observed for the gem-dimethyl group. The mass spectrum showed a molecular ion peak at  $m/z$   $250.1202$  corresponding to the molecular formula  $C_{14}H_{18}O_4$ . The chloro compound **46** showed absorption bands at  $1769$  (enol ester),  $1738$  (ester),  $1628$  and  $1602\text{ cm}^{-1}$  (olefin) in the ir spectrum. The nmr

spectrum showed one-proton singlets at  $\delta$  5.92, 4.94 and 4.73 for the vinylic protons. The signal at  $\delta$  3.84 (t, 2H,  $J = 7$  Hz) was due to the methylene protons adjacent to the chlorine atom. The gem-dimethyl group appeared at  $\delta$  1.20 and 1.02 as singlets. The mass spectrum gave a molecular ion peak at 286.0972, 288.0964 consistent with molecular formula  $C_{14}H_{19}O_4Cl$ . The acidic material, which was produced in 57% yield, was initially thought to be the desired compound **47** resulting from concomitant Claisen rearrangement of acylation product **45**. This acid turned out to be the hydrolyzed Diels-Alder adduct between acryloyl chloride and enone **42**. Esterification with potassium carbonate and methyl iodide in acetone afforded the corresponding methyl ester **48**, which showed in the ir spectrum a strong carbonyl absorption at  $1737\text{ cm}^{-1}$  (ester) and an olefinic absorption at  $1640\text{ cm}^{-1}$ . The absence of




any vinylic protons in the nmr spectrum strongly suggested the structure to be **48**. This was in accord with the

appearance of a pair of doublets of doublets at  $\delta$  3.74 ( $J = 8, J' = 5$  Hz) and 3.72 ( $J = 8, J' = 3$  Hz) due to the methylene protons adjacent to the enol ether oxygen as well as four methyl singlets at  $\delta$  3.70, 3.65, 0.92 and 0.75. The overall incorporation of a methyl acrylate moiety was further confirmed by the mass spectrum in which a molecular ion peak was found at m/z 282.1469. The structure of this compound was further verified by  $^{13}\text{C}$  nmr spectrum which showed the signals at  $\delta$  173.93, 171.71, 121.29 and 116.54, clearly indicating the presence of two ester carbonyls and a double bond. Although the spectral data indicated that the Diels-Alder adduct was a single stereoisomer, the stereochemistry could not be deduced unambiguously.

When the above enol ester formation was carried out in refluxing acryloyl chloride using potassium carbonate as the base, the formation of the Diels-Alder product 48 was completely suppressed and the enol ester 45 was isolated in 30% yield along with a 12% yield of 46. In this case a large amount of unreacted material was recovered even after a long reaction period of time. Further improvement of the reaction was effected by increasing the reaction temperature. Thus, treatment of enone 42 with acryloyl chloride and potassium carbonate in refluxing toluene over an extended period (72 h) gave rise



to enol ester 45 and chloride 46 in 53% and 6% yield respectively. Under these conditions, two new compounds were also formed in small quantities. Structures 49 and 50 were tentatively assigned to these compounds mainly on



the basis of the mass spectra which showed molecular ion peaks at m/z 446.3507 and m/z 482.2385. Further improvement on the formation of the desired enol ester 45 was attempted without success. When sodium carbonate was used as a base similar results were obtained, while a complex mixture was produced on carrying out the reaction in refluxing xylene.

With enol ester 45 in hand, we turned our attention to the Claisen rearrangement which could in principle lead to the formation of ketene 51. Furthermore, when the reaction is carried out in the presence of an alcohol the

corresponding ester is expected. Thus, enol ester 45 was subjected to rearrangement in refluxing toluene-methanol and xylene-methanol. Under these conditions, however, the starting material was recovered intact. On the other hand, when compound 45 was heated in refluxing xylene-pentanol, the desired diester 52 was formed. However, the







yield was found to be very low. After 48 h, only 16-33% yield of 52 was produced, with the balance of material being mainly the unreacted starting compound. Diester 52 showed in the ir spectrum a broad carbonyl band centered at  $1738\text{ cm}^{-1}$  and an absorption at  $1642\text{ cm}^{-1}$  indicative of an olefin. The nmr spectrum displayed two one-proton singlets at  $\delta$  6.40 and 5.22 characteristic of geminal protons of the methylidene group. The triplet at  $\delta$  3.97 integrating to two protons was attributed to the methylene protons adjacent to the oxygen. The methyl ester protons appeared as a singlet at  $\delta$  3.70. Two other methyl singlets and a methyl triplet ( $J = 7\text{ Hz}$ ) were obtained at

$\delta$  1.20, 1.02 and 0.92 respectively. The mass spectrum showed a molecular ion peak at m/z 338.2093 indicating the molecular formula of  $C_{19}H_{30}O_5$ . Attempts to improve the yield of diester 52 by increasing the reaction time and by the use of xylene and cyclohexanol were unsuccessful.


Surprisingly, no reaction was observed using the latter combination. Although the formation of the desired diester 52 could be effected via the Claisen rearrangement, the best yield obtained was only 33% from enol ester 45. This yield was not synthetically useful and an alternative route was sought.

In view of the problems associated with the methylidene group in compound 42, we decided to remove it prior to further elaboration. The cleavage of the methylidene group was initially attempted via a combination of hydration and retro-Aldol reaction. In practice, this proved not to be viable. Under no conditions applied, could the desired keto ester 53 be generated.




While this work was in progress, the elimination of the methylidene group of a closely related system 54 was investigated in our laboratory.<sup>87</sup> It was found that compound 54 could be converted to the corresponding epoxides 55 by treatment with hydrogen peroxide and lithium hydroxide in methanol. Interestingly, when the mixture of the epimeric epoxides was heated with sodium hydroxide in methanol at reflux, keto ether 56 was produced in 68% yield, presumably via a mechanistic pathway illustrated in Scheme XII. This procedure was

Scheme XII




applied to compound 42. On addition of a premixed solution of aqueous lithium hydroxide (0.2 eq.) and 30% hydrogen peroxide to a methanolic solution of enone 42 at 0°C followed by warming up to room temperature, a mixture of epimeric epoxides (57) (1:1) was formed in 86% yield after 3 h. The sequence of addition was found to be crucial to the success of the epoxidation. Sequential addition of lithium hydroxide and hydrogen peroxide without premixing gave rise to a large amount of methoxy ketone 58 as a by-product, apparently produced by the 1,4-addition of methoxide to the enone moiety. The nmr spectrum of epoxides



57 displayed protons on the epoxide ring at  $\delta$  3.15 and 2.85 as doublets with a coupling constant of 6 Hz each for one isomer and at  $\delta$  3.00 and 2.96 also as doublets ( $J = 6$  Hz) for the other isomer. A pair of singlets at  $\delta$  3.76 and 3.72 integrating to three protons each was due to methyl ester groups. Gem-dimethyl groups appeared as singlets at  $\delta$  1.04, 0.98, 1.00 and 0.94. The ir spectrum showed absorption bands at 1742 (five membered ring).

ketone) and  $1738\text{ cm}^{-1}$  (ester). A molecular ion peak at m/z 212.1043 was consistent with molecular formula  $\text{C}_{11}\text{H}_{16}\text{O}_4$ .

Heating a solution of epoxides 57 and sodium hydroxide (2.5 eq.) in methanol-water (10:1) at reflux for 2 to 3 days afforded an acidic material which was esterified immediately with potassium carbonate and methyl iodide in acetone. Two products were isolated, the desired keto ester 53 in 18-27% yield and a by-product 59 in nearly equal amount. The ir spectrum of 53 showed an absorption band at  $1740\text{ cm}^{-1}$  due to the five membered ring ketone and the ester. The nmr spectrum displayed a singlet at  $\delta$  2.14 for the neopentyl protons adjacent to the carbonyl. Three additional singlets were observed at  $\delta$  3.70, 1.18, and 0.92 for the methyl groups. The mass spectrum showed a molecular ion peak at m/z 184.1097 corresponding to the molecular formula  $\text{C}_{10}\text{H}_{16}\text{O}_3$ . The




59

major by-product was identified as the rearranged keto ester **59** on the basis of the following spectral data. The ir spectrum showed diagnostic absorption bands at 1735 (ester), 1710 and 1620  $\text{cm}^{-1}$  ( $\beta$ -methoxy  $\alpha,\beta$ -unsaturated ketone). The nmr spectrum showed a vinyl proton as a singlet at  $\delta$  7.15. Four methyl signals appeared as sharp singlets at  $\delta$  3.86 (methyl enol ether), 3.72 (-COOCH<sub>3</sub>), 1.32 and 1.04 (gem-dimethyl). Exact mass measurement revealed a molecular weight of 226.1204 in agreement with the required molecular formula of C<sub>12</sub>H<sub>18</sub>O<sub>4</sub>. The formation of compound **59** could be attributed to a 1,3-glycol type cleavage (**57**  $\rightarrow$  **60**), followed by a Claisen-type cyclization (**60**  $\rightarrow$  **61**) as shown in Scheme XIII. Methylation of diketo acid **61** would lead to the observed product **59**.


In order to improve the transformation of epoxides **57** to keto ester **53**, several modifications were examined.

When the reaction was carried out at 0°C, keto ester **59** was isolated as the only product after esterification.

The best results were obtained when the reaction was performed in the following manner. To the epoxides **57** in refluxing methanol, an aqueous solution of sodium hydroxide in methanol was added. After ca. 3 days and the ensuing esterification, the desired ester was formed in 54% yield along with a 20% yield of the rearranged

Scheme XIII

product 59. Other attempts were also made to effect the transformation of epoxides 57 to 53. It was envisaged that zinc reduction of 57 could lead to the  $\beta$ -hydroxy ketone 62 which might then undergo retro-aldo reaction to give ketone 53. Unfortunately, zinc reduction of epoxides 57 in refluxing acetic acid gave mainly methyl ketone 63 (33% yield) and only a small amount of the desired hydroxy ketone 62 (14% yield). The latter showed in the ir



spectrum broad bands centered at  $3540\text{ cm}^{-1}$  for the hydroxyl group and  $1740\text{ cm}^{-1}$  for the ketone and ester.


The mass spectrum displayed a molecular ion peak at  $m/z 214.1206$  which was consistent with molecular formula  $C_{11}H_{18}O_4$ . The nmr spectrum showed a methoxy singlet at  $\delta 3.70$  which was partially superimposed with the multiplet centered at  $\delta 3.72$  for the methylene protons adjacent to the hydroxyl group. Only a pair of singlets appeared at  $\delta 1.06$  and  $0.84$  for the gem-dimethyl group suggesting a single stereoisomer. The stereochemistry however could not be readily assigned on the basis of the available data. The methyl ketone 63 was probably produced by dehydration of compound 62 followed by further reduction of the resulting enone (i.e. 42). The absorption at  $1739\text{ cm}^{-1}$  in the ir spectrum was due to ketone and ester functional groups. The nmr spectrum displayed a quartet ( $J = 8.5\text{ Hz}$ ) at  $\delta 2.04$  for the methine proton adjacent to the carbonyl group. The doublet ( $J = 8.5\text{ Hz}$ ) at  $\delta 0.96$  was due to the newly introduced methyl group. Other methyl signals were observed at  $\delta 3.70, 1.12$  and  $0.85$ ,

each as a singlet. The mass spectrum showed a molecular ion peak at m/z 198.1262 confirming the molecular formula.

When the zinc reaction was carried out in refluxing benzene, an improved yield (38%) of hydroxy ketone 62 was realized along with recovery of an approximately equal amount of the starting material. The retro-aldol reaction of the hydroxy ketone 62 was examined using sodium methoxide in refluxing benzene. The reaction gave rise to a complex mixture which could not be easily identified.

Due to these findings the zinc reduction approach to ketone 53 was abandoned.

As an alternative, epoxide 57 was subjected to treatment with sodium hydride (2 eq.) in benzene at reflux, in an attempt to prepare the formyl ketone 64 which could then be deformylated to ketone 53.




64

Interestingly, the latter compound was formed directly but in low yield (18%) and a substantial amount of the starting material was recovered. The yield of 53 could


not be improved to a greater extent even with a larger excess of sodium hydride. Likely, the transformation was induced by the small amount of sodium hydroxide present in the reaction mixture.

For the incorporation of the required propionate side chain, keto ester 53 was subjected to Michael addition with ethyl acrylate (1.5 eq.) in 1,2-dimethoxyethane at room temperature in the presence of sodium hydride for 24 h. Good yields (83-93%) of the desired diester 65 could be obtained when less than 300 mg of the starting





ketone was used. The ir spectrum of 65 displayed diagnostic absorption bands at  $1739\text{ cm}^{-1}$  for the ester and ketone carbonyl groups. In the nmr spectrum, the ethyl group appeared at  $\delta$  4.12 (q, 2H,  $J = 8\text{ Hz}$ ) and 1.32 (t, 3H,  $J = 8\text{ Hz}$ ). The methylene protons  $\alpha$  to the ketone appeared as a singlet at  $\delta$  2.18. Three methyl singlets were obtained at  $\delta$  3.72, 1.08 and 0.94. The mass spectrum showed a molecular ion peak at  $m/z$  284.1617 indicating the



molecular formula  $C_{15}H_{24}O_5$ . Dieckmann condensation of diester 65 was effected by sodium methoxide (prepared in situ from methanol and sodium hydride) in refluxing



toluene for 24 h. The diketone ester 66 thus obtained showed in its mass spectrum a molecular ion peak at  $m/z$  252.1201 ( $C_{14}H_{20}O_4$ ). The ir spectrum showed absorptions at 1740 (cyclopentanone), 1656 and 1601  $\text{cm}^{-1}$  (enolized  $\beta$ -keto ester). The nmr spectrum displayed a chelated enol proton at  $\delta$  13.54 as a singlet. The signals at  $\delta$  4.08 (q, 2H,  $J = 8$  Hz) and 1.28 (t, 3H,  $J = 8$  Hz) were due to the ethyl group of the ester. The compound appeared to be a single stereoisomer as only one set of three singlets was observed at  $\delta$  2.18, 1.22 and 1.00. The former could be attributed to the methylene group adjacent to the five-membered ketone carbonyl and the other signals were due to the gem-dimethyl group. Since subsequent transformation employs conditions under which enolization of the five-membered ring ketone would also be effected, the ring junction stereochemistry of compound 66 was not determined.




In principle under the described conditions keto diester 65 could also undergo other modes of cyclization leading to compounds 67 and 68, involving enolate ion of the ketone carbonyl. To confirm the structural assignment, keto ester 66 was prepared in an unambiguous manner as follows. Thiolketalization of 65 with 1,2-ethanedithiol and boron trifluoride etherate in methylene chloride at room temperature for 6 h gave rise to the corresponding thiolketal 69 in 82% yield. The nmr spectrum showed a multiplet at  $\delta$  3.33 and integrating to four protons for the methylene protons of thiolketal group. The mass spectrum showed a molecular ion peak at m/z 360.1432 indicating the chemical formula  $C_{17}H_{28}S_2O_4$ . The thiolketal 69 was subjected to Dieckmann condensation in 1,2-dimethoxyethane using sodium hydride and methanol and a catalytic amount of potassium hydride. After 5 h at reflux,  $\beta$ -keto ester 70 was isolated in 78% yield. The ir spectrum showed an intense and broad absorption band at



$1735\text{ cm}^{-1}$  and two weak bands at  $1655$  and  $1610\text{ cm}^{-1}$  indicating the compound was mainly in the ketone form. The nmr spectrum displayed a quartet at  $\delta 4.10$  and a triplet at  $\delta 1.27$  for the ethyl group of the ester. A multiplet at  $\delta 3.27$  was due to the methylene protons adjacent to the sulfur atoms. A molecular ion peak at  $m/z 328.1169$  in the mass spectrum verified the chemical formula of  $C_{16}H_{24}S_2O_3$ . Dethiolketalization<sup>88</sup> of keto ester 70, effected by mercuric chloride and calcium carbonate in refluxing aqueous acetonitrile for 4 h, gave keto ester 66 (45% yield) identical in all respects with the one obtained previously.

While the sodium hydride induced Michael addition of keto ester 53 to ethyl acrylate (vide supra) proceeded smoothly when small amount of material was used, for reasons that remain unknown, the yield of product decreased substantially when reaction was scaled up. For instance, when 76 mg of keto ester 53 was used, a 93%

yield of diester 65 was obtained. The yield dropped to 83% when 216 mg of compound 53 was applied. When the Michael reaction was attempted in gram-scale under the described conditions at room temperature in 1,2-dimethoxyethane, it was consistently observed that the reaction ceased after the initial formation of the product. In an attempt to promote the reaction by heating the mixture to reflux, extensive alkylation occurred and triester 71 was isolated in 76% yield as the only




product. In order to circumvent this problem the use of lithium diisopropylamide as the base was investigated.

When keto ester 53 was treated with lithium diisopropylamide (prepared in situ from diisopropylamine and methyllithium) at -78°C in tetrahydrofuran followed by addition of ethyl acrylate, the desired product 65 was formed after 1 h at -78°C and 4 h at room temperature.


The yield was 86% when 303 mg of the starting material was used. Disappointingly, when the reaction was carried out with ca. 500 mg of the material, the yield of the desired

product dropped again rapidly to ca. 40% along with substantial recovery of the starting material. After some further preliminary investigation, it was observed, to our delight, that the reaction took a different course when performed in the presence of a small amount of hexamethylphosphoramide. In addition to the formation of diesters 72 (methyl acrylate was used as the reagent), subsequent Dieckmann cyclization occurred simultaneously to give a single keto ester 73 existing completely in the



corresponding enol form. The best results were obtained when keto ester 53 was subjected to treatment with lithium diisopropylamide (1.1 eq.) in tetrahydrofuran in the presence of hexamethylphosphoramide (1.1 eq.) at -78°C followed by addition of methyl acrylate (1.2 eq.) at -40°C and by allowing the mixture to react at 0°C for 5 h.


Under these conditions,  $\beta$ -keto ester 73 was produced in 48% yield, along with 23% of diesters 72 (7:3) and 18% of by-product 74. Diesters 72 could readily be cyclized to provide an additional amount of  $\beta$ -keto ester 73 (94%



yield) when further exposed to lithium diisopropylamide in tetrahydrofuran at room temperature for 6 h. The ir spectrum of diesters 72 and keto ester 73 were found to be similar to those obtained for the corresponding ethyl compounds 65 and 66. In the mass spectra, each showed a molecular ion peak with fourteen mass units less, while in the nmr spectrum, the ethyl ester signals observed for 65 and 66 were replaced by the methoxy singlets (four singlets at  $\delta$  3.75, 3.72, 3.70, 3.68 for 72 and a singlet at  $\delta$  3.77 for 73). Diketo ester 74 showed in the ir spectrum a broad band at  $1739\text{ cm}^{-1}$  for the esters and the five membered ring ketone. The absorption at  $1713\text{ cm}^{-1}$  was assigned to the six membered ring ketone. In the nmr spectrum five singlets appeared at  $\delta$  3.75, 3.65 (methoxy groups), 2.30 (methylene protons of the cyclopentanone moiety), 1.18 and 0.92 (gem-dimethyl). The mass spectrum showed a molecular ion peak at m/z 324.1573 in agreement with the molecular formula  $C_{17}H_{24}O_6$ .

Concurrent to the studies described immediately above, the alkylation of keto ester 53 with  $\beta$ -bromo-propionate was also examined. Sequential treatment of 53 with near stoichiometric amounts of lithium diisopropylamide and methyl 3-bromopropionate in tetrahydrofuran in the presence of hexamethylphosphoramide (1.0 eq.), initially at  $-78^{\circ}\text{C}$  then at room temperature for 10 h gave rise to keto ester 73 and diesters 72 in ca. 1:2 ratio and in a total yield of 66%. Both this reaction and the Michael reaction using hexamethylphosphoramide were shown to be reproducible regardless of the reaction scale. Of these two procedures the latter is preferred since it provides directly a higher yield of the cyclic material.


Scheme XIV



For the construction of the C ring from compound 73, two approaches are readily conceivable (Scheme XIV). In one approach the ester group is to be used as the angular methyl group and a suitable substituent will be introduced ( $73 \rightarrow 75$ ) to facilitate the C ring formation. In the other, a methyl group will be incorporated ( $73 \rightarrow 76$ ) and the ester moiety be used as part of ring C. The viability of these approaches will depend on the stereochemical outcome of the alkylation step (i.e.  $73 \rightarrow 75$  and  $73 \rightarrow 76$ ). Both of these approaches were examined and the results are as follows.

Compound 73 was found to undergo both methylation with methyl iodide and Michael addition with acrolein.

Treatment of 73 with ca. one equivalent each of acrolein and 1,4-diazabicyclo[2.2.2]octane in 1,2-dimethoxyethane at room temperature for 4 h gave keto aldehyde 77 in 90%




77


yield. The ir spectrum of 77 showed absorption bands at 2860, 2720 (aldehyde), 1740 (ester and cyclopentanone) and 1713  $\text{cm}^{-1}$  (aldehyde and cyclohexanone). The nmr spectrum displayed a broad singlet at  $\delta$  9.66 for the aldehyde proton. Sharp methyl singlets were observed at  $\delta$  3.72 (methyl ester), 1.08 and 1.02 (gem-dimethyl group). The regiochemistry of the Michael adduct was substantiated by the presence of a two-proton singlet at  $\delta$  2.18 assigned to the cyclopentanone methylene group and the disappearance of the signal corresponding to the enol proton. The mass spectrum displayed a molecular ion peak at m/z 294.1466. The spectral data also suggested that the compound was a single stereoisomer. The indicated stereochemistry was deduced from the results obtained for the subsequent methylation reaction of 73 (see below). Since this compound likely possesses the incorrect stereochemistry, further elaboration was not carried out.

The initially attempted methylation, using sodium hydride as a base, resulted in complete recovery of the starting material. When potassium hydride was employed, the reaction of 73 with methyl iodide in tetrahydrofuran at room temperature occurred to give the desired methylation product in 82% yield. However, dimethylation also occurred to the extent of 8-10%. This side reaction could be suppressed completely by the use of potassium carbonate.

as the base and acetone as the solvent.<sup>89</sup> A 82% yield of two products was obtained in the ratio of 2:1. These compounds were not readily separable. However, by fractional crystallization the major component (m.p. 75-76°C, petroleum ether-ethyl acetate) could be isolated in pure form. Treatment of this compound with potassium carbonate in refluxing acetone for 24 h gave rise to a new mixture consisting of the starting material and the 'minor' compound in ca. 1:3 ratio with the latter predominating. These observations clearly indicated that the two products produced by the methylation reaction were isomeric at an epimerizable center(s). Since the



78




79

structure 78 could be assigned to the major isomer on the basis of the spectral data, it follows that the minor isomer should possess the structure 79. Diketo ester 78 showed absorption bands at 1741 and 1710 cm<sup>-1</sup> in the ir spectrum due to the ester and ketone functionalities. The

mass spectrum showed a molecular ion peak at m/z 252.1363 in agreement with molecular formula  $C_{14}H_{20}O_4$ . The nmr spectrum displayed four methyl singlets at  $\delta$  3.74, 1.43, 1.17 and 1.07. By decoupling experiments it was possible to assign all the remaining protons as follows:  $\delta$  2.54 (dd,  $J = 15$ ,  $J' = 5$  Hz,  $H_B$ ), 2.48 (dd,  $J = 15$ ,  $J' = 13$  Hz,  $H_a$ ), 2.46 (ddd,  $J = J' = 13$ ,  $J'' = .5$  Hz,  $H_d$ ), 2.35 (dd,  $J = 19$ ,  $J' = 2$  Hz,  $H_g$ ), 2.26 (t,  $J = 13$  Hz,  $H_e$ ), 2.16 (d,  $J = 19$  Hz,  $H_h$ ), 2.14 (dd,  $J = 13$ ,  $J' = 5$  Hz,  $H_f$ ) and 1.90 (ddd,  $J = J' = 13$ ,  $J'' = 5$  Hz,  $H_c$ ).

Since both of the ring junction protons  $H_c$  and  $H_d$  appeared as a doublet of doublets of doublets with two large coupling constants of 13 Hz each and a small coupling of 5 Hz, a trans-ring junction could readily be assigned. In order to determine the stereochemistry of the remaining center, the following noe experiments were carried out. When  $C_4$  methyl singlet ( $\delta$  1.43) was irradiated, the doublet of doublets of doublets at  $\delta$  2.46 ( $H_d$ ) and the doublet of doublets at  $\delta$  2.48 ( $H_a$ ) each showed a 4% enhancement. An 8% enhancement was observed for  $H_f$  which appeared at  $\delta$  2.14 as a doublet of doublets while the doublet of doublets of doublets at  $\delta$  1.90 ( $H_c$ ) was unaffected. These observations strongly suggested that the methyl group was cis to  $H_a$   $H_d$  and  $H_f$  and trans to  $H_e$ . Consequently, the stereochemistry of compound 78 was







The three dimensional X-ray crystallographical  
of structure of diketo ester 78


Fig 1

deduced. This assignment was further confirmed by a single X-ray diffraction analysis. A computer generated perspective drawing of this compound is shown in Fig. 1.

Both diketo esters 78 and 79 are synthetically useful, since they differ from each other only in stereochemistry at an epimerizable center. Since the trans compound 78 was obtained in larger quantity and in pure form, it was used for the present studies. Prior to the incorporation of ring C, the removal of the six membered ketone carbonyl was examined. In principle, this could be effected by selective reduction of the less hindered five membered ring ketone (**78** → **80**), thioacetalization (**80** → **81**), desulfurization (**81** → **82**) and regeneration of the cyclopentanone (**82** → **83**). Thus, diketo ester **78** was





**80****81****82****83**

subjected to reduction with lithium tri-*t*-butoxyaluminium hydride (1.2 eq.)<sup>61</sup> in tetrahydrofuran at room temperature. After 12 h, the reaction was complete and two products were formed in 97% yield. To our surprise, the products were shown to be keto alcohols **84**, resulting from the selective reduction of what appeared to be the more hindered six membered ring ketone. That




the cyclohexanone moiety was selectively reduced was indicated by absence of the characteristic cyclohexanone absorption at  $\sim 1710\text{ cm}^{-1}$  in the ir spectrum. Instead, the products displayed absorption bands at  $3480\text{ cm}^{-1}$  for the hydroxyl group and  $1740\text{ cm}^{-1}$  for the carbonyls. In the nmr spectrum a broad triplet appeared at  $\delta 3.94$  for the proton adjacent to the hydroxyl group. A pair of mutually coupled doublets ( $J = 13\text{ Hz}$  each) at  $\delta 2.18$  and  $2.09$  integrating to one proton each was assigned to the methylene group attached to the ketone carbonyl. A singlet at  $\delta 3.74$  was due to the methoxy group. Two sets of methyl

signals at  $\delta$  1.12, 1.08, 1.03 (major) and  $\delta$  1.14, 1.06 and 1.00 clearly indicated the presence of two alcohols (3:1). The mass spectrum showed a molecular ion peak at m/z 254.1522 confirming the chemical formula  $C_{14}H_{22}O_4$ . To verify the structural assignment, the mixture of keto alcohols 84 was converted to thioketals 85 (90% yield) using boron trifluoride etherate and 1,2-ethanethiol in dichloromethane. Dethioketalization with Raney nickel in ethanol followed by oxidation of the resulting hydroxy esters 86 with pyridinium chlorochromate<sup>90</sup> and sodium acetate in dichloromethane gave a single product (81% yield) which was identified as ketone 87. It showed in the ir spectra.



an absorption band at  $1709\text{ cm}^{-1}$  diagnostic of a cyclohexanone moiety. The isomeric ketone 88 was not detected.


In order to prepare keto ester 88, the mixture of keto alcohols 84 was ketalized by transketalization with 2-ethyl-2-methyl-1,2-dioxolane and a catalytic amount of p-toluenesulfonic acid in refluxing benzene. Ketals 89 were produced in 80% yield. The mixture of isomeric ketals displayed a carbonyl band at  $1738\text{ cm}^{-1}$  and a hydroxy absorption at  $3480\text{ cm}^{-1}$  in the ir spectrum. The nmr spectrum showed multiplets at  $\delta$  3.87 for the ketal moiety and 3.97 for the methine proton adjacent to the hydroxy group. Two sets of methyl signals for the epimeric alcohols were observed at  $\delta$  3.70, 1.22, 1.02, 0.94, and  $\delta$  3.68, 1.20, 1.00, and 0.93. The mass spectrum showed a molecular ion peak at m/z 298.1781 indicating the chemical formula  $C_{16}H_{26}O_5$ .




Sequential treatment of ketal alcohols 89 with sodium hydride, carbon disulfide and methyl iodide in tetrahydrofuran gave rise to the corresponding xanthates 90 in

87% yield. The ir spectrum showed a strong absorption band at  $1220\text{ cm}^{-1}$  due to the carbon-sulfur double bond.

The nmr spectrum showed a singlet at  $\delta$  2.48 and integrating to three protons for the methyl group attached to the sulfur atom.




When the mixture of xanthates 90 was heated with tri-n-butyltin hydride and a catalytic amount of azobisisobutyronitrile<sup>91</sup> in degassed toluene under an argon atmosphere at reflux for 4 h, ketal 91 was isolated as a single compound in 78% yield. The ir spectrum displayed an absorption band at  $1734\text{ cm}^{-1}$  for an ester. In the nmr spectrum, a multiplet at  $\delta$  3.87 was due to the protons attached to the dioxolane ring. A set of four methyl singlets appeared at  $\delta$  3.68, 1.21, 1.02 and 0.94. The mass spectrum showed a molecular ion peak at 282.1831 indicating a molecular formula  $C_{16}H_{26}O_4$ . Exposure of 91 to moist acetone and p-toluenesulfonic acid furnished keto ester 88 in 94% yield. The ir spectrum showed an



absorption band at  $1741\text{ cm}^{-1}$  for the ester and the five membered ring ketone. The nmr spectrum displayed a singlet at  $\delta 2.19$  for the methylene protons neighboring the ketone moiety. Four other singlets were observed at  $\delta 3.68$  ( $-\text{COOCH}_3$ ), 1.20, 1.16 and 1.04 for the methyl groups. A molecular ion peak at  $m/z 238.1562$  in the mass spectrum confirmed the molecular formula  $C_{14}H_{22}O_3$ .

A shorter and more effective method has also been developed for the transformation of diketo ester 78 to keto ester 88. In this two-step procedure, diketone 78 was selectively thioketalized with 1,2-ethanedithiol and boron trifluoride etherate in methylene chloride at  $0^\circ\text{C}$ .






The ir spectrum of thioketal **92** thus obtained in 86% yield showed the absence of cyclohexanone absorption suggesting the six membered ring ketone was selectively ketalized.

The mass spectrum displayed a molecular ion peak at m/z 328.1166 corresponding to the molecular formula  $C_{16}H_{24}O_3S_2$ .


Subsequent desulfurization of **92** using Raney nickel in ethanol afforded keto ester **88** in 52% yield.

For the construction of C ring, two approaches were considered. In the first one, the methyl ester moiety in **88** was to be used as a handle for the incorporation of a suitable side chain as shown in structure **93**. Cyclization via an intramolecular Michael addition could lead to the required C ring (**93**  $\rightarrow$  **94**). In the second approach, introduction of a propionate unit at the ring junction could give rise to diester **95**. Dieckmann condensation of


**93****94****95**

**95** followed by decarboxylation could then provide the desired diketone **94**. Both of these approaches were

attempted. Reduction of ketal 91 with sodium bis(2-methoxyethoxy)aluminium hydride in ether at room temperature for 2 h afforded hydroxy ketal 96 in 95% yield. Aldehyde 97 was prepared in 78% yield by treatment of the alcohol 96 with pyridinium chlorochromate and sodium acetate in dichloromethane at room temperature for





96



97

1.5 h. The mass spectrum of the compound suggested the molecular formula to be  $C_{15}H_{24}O_3$  with a molecular ion peak at m/z 252.1536. The ir spectrum showed absorption bands at 2830, and  $1739\text{ cm}^{-1}$  characteristic of a aldehyde moiety. The nmr spectrum displayed a broad singlet at  $\delta$  9.38 readily attributed to the aldehyde proton. Other easily identifiable signals were observed at  $\delta$  3.86 (ketal), 1.10, 1.04 and 0.96 (methyls). The conversion of the aldehyde group to a vinyl ketone required for the desired intermediate 93 proved difficult. Disappointingly, attempts to introduce the vinyl group by a Grignard

reaction were unsuccessful. Neither vinylolithium nor the corresponding magnesium bromide reagent reacted with aldehyde **97** at 0°C. At room temperature the reaction of **97** with vinylolithium gave a complex mixture from which no identifiable compounds could be isolated. In case of vinylmagnesium bromide, the products appeared less complex. However, the desired compound **98** could not be detected. Instead, an impure material which showed characteristics of the rearrangement product **99** was isolated in small amount.


**98****99**

As a consequence of the above unsuccessful attempts, we turned our attention to the alternative approach for ring C formation. To prepare the desired compound **95**, keto ester **88** was subjected to treatment with methyl acrylate and potassium hydride in tetrahydrofuran. At 0°C no reaction occurred and the starting material was


recovered intact. When the reaction was carried out at room temperature, the starting material was consumed completely after 14 h and a major product was formed.

Disappointingly, the product which was found to be non-acidic and less polar than the starting material on silica gel, showed spectroscopic properties totally inconsistent with the desired structure 95. Although the ir showed a carbonyl absorption at  $1739\text{ cm}^{-1}$  which was expected for a five membered ketone and the ester groups, this band was rather sharp and not particularly intense. This observation coupled with the absence of any methoxy signals in the nmr spectrum suggested that the ester group had been eliminated from the starting material. The loss of a carbomethoxy was further suggested by the mass spectrum which displayed a peak at  $m/z 180.2230$  for the composition  $C_{12}H_{20}O$ . At the present, we are unable to arrive at a structure for this compound which would satisfy both the available spectral data and a reasonable mechanistic pathway for its formation. One possible pathway leading to this unidentified product could involve an intra-molecular Claisen-type cyclization ( $95 \rightarrow 100$ ) as the initiating step. In order to circumvent this possible problem, the ester 92 was subjected to treatment with potassium hydride and methyl acrylate in tetrahydrofuran at room temperature. To our delight the desired addition

product 101 was formed albeit in low yield (19% with recovery of starting material) even after a long reaction period of 32 h. The mass spectrum which showed the molecular ion peak at 414.1535 confirmed molecular formula  $C_{20}H_{30}O_5S_2$ . The nmr spectrum showed two methoxy signals at  $\delta$  3.72 and 3.68. Three methyl signals appeared at  $\delta$  1.32, 1.25 and 1.14. The multiplet at  $\delta$  3.25 was due to methylene protons next to the sulfur atoms. These spectral data suggested that the adduct was formed as a single stereoisomer. However, the stereochemistry could not be unambiguously deduced. The indicated stereochemistry was tentatively assigned based on the expected addition of methyl acrylate from the less hindered side of the precursor. Currently, we are examining more suitable conditions in order to improve the yield of compound 101.



100




101


In conclusion, starting from 10-camphorsulfonic acid (35), keto diester 101 has been prepared in eleven steps by the synthetic sequence summarized in Scheme XV. This

compound is considered to be a potentially useful synthetic precursor of clovane-diol (1). Desulfurization of 101 followed by Dieckmann condensation of the resulting ester 95 should lead to tricyclic compound 94 after decarbomethoxylation. This compound could be further elaborated to give the target molecule by modification of the existing functionalities.

Scheme XV



Scheme XV (cont'd)



a. KOH, heat. b.  $K_2CO_3$ ,  $CH_3I$ , acetone. c.  $O_2$ ,  $Ac_2O$ , Py, DMAP,  
 $CH_2Cl_2$ , h.v. d.  $H_2O_2$  / OH. e.  $NaOH$ ,  $CH_3OH$ , reflux. f.  $K_2CO_3$ ,  $CH_3I$ ,  
acetone. g. LDA, -78°C to R.T.,  $CH_2=CHCO_2CH_3$ . h.  $Na^+ OCH_3$ ,  
toluene, reflux. i.  $K_2CO_3$ ,  $CH_3I$ , acetone. j.  $BF_3 \cdot Et_2O$ ,  $HSCH_2CH_2SH$ ,  
0°C. k. KH,  $CH_2=CHCO_2CH_3$ , THF, R.T.

## EXPERIMENTAL

### General

For general remarks see the Experimental Section of Part I of this thesis. Carbon-13 nuclear magnetic resonance ( $^{13}\text{C}$  nmr) spectra were recorded on a Bruker WH-200 or WH-400 spectrometer and were obtained on solutions in deuteriochloroform. Crystalline samples were recrystallized and liquid samples were subjected to distillation before submitting for elemental analysis.

### Materials

Solvents were purified as follows: triethylamine, pyridine, diisopropylamine and acetonitrile by distillation over calcium hydride; dimethyl sulfoxide and hexamethylphosphoramide by distillation over calcium hydride at reduced pressure and stored over 3A molecular sieves; dichloromethane was distilled over phosphorus pentoxide; acetone was treated with potassium permanganate, dried over anhydrous potassium carbonate and distilled. Boron trifluoride etherate was distilled over calcium hydride according to the procedure of Brown.<sup>92</sup> Other solvents were distilled by the process described in

the Experimental Section of Part I. dl-Camphorsulfonic acid was obtained from Aldrich Chemical Company.

(±)-Campholenic acid (34)

A. From dl-10-camphorsulfonic acid (35)

Potassium hydroxide (242 g, 4.3 mol) was fused in a porcelain dish. The heat source was removed and immediately powdered dl-10-camphorsulfonic acid (100 g, 0.43 mol) was added slowly with vigorous stirring (the reaction is exothermic) over a period of 25 min. The colour of the molten mixture was chocolate brown. The molten mass was allowed to cool to room temperature and then dissolved in water (500 mL). The resulting solution was washed with dichloromethane (3 x 300 mL) and the aqueous fraction acidified with ice-cold 6 N aqueous hydrochloric acid. The acidified solution was extracted with dichloromethane (3 x 200 mL). The combined organic extract was dried, filtered and concentrated. The crude residue was distilled to give pure acid 34 (52 g, 71% yield): b.p. 87-89°C/0.5 torr; ir 3500 (acid) and 1719 cm<sup>-1</sup> (carbonyl); nmr δ 10.90 (br, s, 1H, -CO<sub>2</sub>H), 5.20 (m, 1H, -CH=), 1.60 (br, s, 3H, CH<sub>3</sub>-C=), 0.99 (s, 3H, -CH<sub>3</sub>) and 0.81 (s, 3H, -CH<sub>3</sub>); ms M<sup>+</sup> 168.1142 (calcd. for C<sub>10</sub>H<sub>16</sub>O<sub>2</sub>: 168.1150). Anal. Calcd. for C<sub>10</sub>H<sub>16</sub>O<sub>2</sub>: C 71.38, H 9.59; Found: 71.40, H 9.68.

B. From dl-10-camphorsulfonic acid sodium salt

To the fused potassium hydroxide (115 g, 2.05 mol) in a porcelain casserole, was added slowly with stirring dl-10-camphorsulfonic acid sodium salt (100 g, 0.39 mol). The heat source was carefully controlled to prevent the charring of the molten mass. After the addition (ca. 20 min), the molten mass was allowed to cool to room temperature and then dissolved in water (500 mL). The resulting solution was washed with ether and the aqueous fraction acidified with ice-cold 6 N hydrochloric acid solution then extracted with dichloromethane. The extracts were combined, dried, filtered and concentrated. The crude residue was distilled as before to give acid 34 (58 g, 78% yield). The spectral data were identical in all respects to those obtained previously (vide supra).

Methyl campholenate (40)

To a solution of campholenic acid 34 (45.0 g, 0.28 mol) in acetone (50 mL), anhydrous potassium carbonate (89.7 g, 0.65 mol) was added. After stirring under an argon atmosphere for 1 h, methyl iodide (33.3 mL, 0.53 mmol) was added and the mixture stirred for an additional 18 h. The reaction mixture was then filtered. The solid material was dissolved in water and extracted with

dichloromethane. The dichloromethane extract and acetone filtrate were combined. After most of the solvent had been evaporated, the mixture was taken up in dichloromethane (200 mL) and washed with ice-cold 2 N hydrochloric acid solution, water and saturated aqueous sodium chloride solution. The organic fraction was dried, filtered and concentrated. The residue was distilled to afford the pure ester 40 (48 g, 98% yield): 83-86°C/0.5 torr; ir 1748 cm<sup>-1</sup> (ester); nmr δ 5.22 (br, s, 1H, =CH), 3.68 (s, 3H, -COOCH<sub>3</sub>), 1.62 (br, s, 3H, =CCH<sub>3</sub>), 0.98 (s, 3H, -CH<sub>3</sub>) and 0.78 (s, 3H, -CH<sub>3</sub>); ms M<sup>+</sup> 182.1305 (calcd. for C<sub>11</sub>H<sub>18</sub>O<sub>2</sub>: 182.1307). Anal. Calcd. for C<sub>11</sub>H<sub>18</sub>O<sub>2</sub>: C 72.45, H 9.95; Found: C 72.12, H 9.82.

4-Carbometoxymethyl-3,3-dimethyl-2-methylidenecyclopentanol (41)

A solution of the ester 40 (5.0 g, 27.5 mmol) and methylene blue (200 mg) in methanol (250 mL) was irradiated with two 200 W tungsten light bulbs for 72 h. During this period a moderate stream of oxygen was bubbled through the solution. The solution was cooled at 0°C and sodium borohydride (0.39 g, 10.3 mmol) was added. After stirring for 1 h, it was poured into ice-cold 2 N aqueous hydrochloric acid solution (300 mL) and extracted with chloroform (3 x 100 mL). The combined chloroform extract

was dried, filtered and concentrated. The residue was chromatographed on silica gel, eluting with 15% ethyl acetate in petroleum ether to give a 7:3 mixture of two epimeric allylic alcohols **41** (3.6 g, 66% yield): ir 3440 (alcohol), 3106, 1650 (olefin) and 1746 cm<sup>-1</sup> (ester); nmr δ 5.16, 5.10, s(br, s, total 1H, -CHH) and 5.02, 4.92 (br, s, total 1H, =CHH); 4.40 (m, 1H, -CHOH) 3.66 (s, 3H, -CO<sub>2</sub>CH<sub>3</sub>), 1.12, 1.02, 0.92 and 0.82 (s, total 6H, 2 x -CH<sub>3</sub>); ms M<sup>+</sup> 198.1256 (calcd. for C<sub>11</sub>H<sub>18</sub>O<sub>3</sub>: 198.1256).

4-Carbomethoxymethyl-3,3-dimethyl-2-methylideneencyclo-  
pentanone (42)

A solution of dimethyl sulfoxide (14.1 mL, 0.20 mol) in dichloromethane (25 mL) was added dropwise to a solution of oxalyl chloride (6.9 mL, 79 mmol) at -78°C under an argon atmosphere. The mixture was stirred for 15 min after which a solution of hydroxy esters **41** (11.2 g, 56 mmol) in dichloromethane (25 mL) was added dropwise. The mixture was stirred for 1 h at -78°C. Triethylamine (25 mL) in dichloromethane (25 mL) was added dropwise and the reaction was allowed to warm up to room temperature overnight. It was poured into ice-cold water and the organic fraction separated, washed with water, dried, filtered and concentrated. The residue was chromatographed on silica gel. Elution with 5% ethyl acetate in

petroleum ether afforded the pure enone **42** (9.4 g, 85%):

ir 1748 (ester), 1729 ( $\alpha,\beta$ -unsaturated ketone), 1634 and 1605  $\text{cm}^{-1}$  (C=C); nmr  $\delta$  5.98 (br, s, 1H;  $=\text{CHH}$ ); 5.20 (br, s, 1H,  $=\text{CHH}$ ), 3.68 (s, 3H,  $-\text{CO}_2\text{CH}_3$ ), 2.54 (dd, 1H,  $J = 18$ ,  $J' = 7$  Hz,  $-\text{COCHH}-$ ), 2.06 (dd, 1H,  $J = 18$ ,  $J' = 11$  Hz,  $-\text{COCHH}-$ ), 1.22 (s, 3H,  $-\text{CH}_3$ ) and 1.00 (s, 3H,  $-\text{CH}_3$ ) <sup>MS</sup> M<sup>+</sup> 196.1099 (calcd. for  $\text{C}_{11}\text{H}_{16}\text{O}_3$ : 196.1099. Anal. Calcd. for  $\text{C}_{11}\text{H}_{16}\text{O}_3$ : C 67.30, H 8.22; Found: C 67.20, H 8.12.

#### Enone ester **42** from modified photo-oxygenation reaction

The reactor was charged with a solution of ester **40** (5.0 g, 27.5 mmol) in dichloromethane (20 mL), acetic anhydride (2.74 mL, 29 mmol), pyridine (1.11 mL, 13.7 mmol), 4-dimethylaminopyridine (60 mg, 0.5 mmol) and 5,10,15,20-tetraphenyl-21H,23H-porphine (15 mg). The reaction was irradiated with two 200 W tungsten light bulbs for 24 h. During this period a moderate stream of oxygen was bubbled through the solution. The reaction mixture was then let to stand without radiation for 2 h.

The reaction mixture was then diluted with ether and washed with saturated aqueous sodium bicarbonate solution, 1 N hydrochloric acid solution, saturated cupric sulfate solution and saturated sodium chloride solution. The organic extract was dried, filtered and concentrated.

Flash chromatography of the residue on silica gel, eluting

with 10% ethyl acetate in petroleum ether gave enone 42 (3.87 g, 72% yield). The spectral data were consistent with those obtained previously. (*vide supra*).

Reaction of enone ester 42 with acryloyl chloride

Enone ester 42 (300 mg, 1.53 mmol), sodium acetate (310 mg, 3.73 mmol) and acryloyl chloride (4 mL) were stirred at room temperature for 1 h then refluxed for an additional 24 h under an argon atmosphere. The reaction was cooled to room temperature. Aqueous 1 N sodium hydroxide solution was added then extracted with dichloromethane and ether. The organic extracts were then washed with concentrated sodium chloride solution, dried, filtered and evaporated. Flash chromatography of the residue on silica gel, eluting with 15% ethyl acetate in petroleum ether gave the required enol ester 45 (64 mg, 18%): ir 1737 (ester), 1630 and 1600  $\text{cm}^{-1}$  (C=C); nmr  $\delta$  6.54 (dd, 1H, J = 18, J' = 2 Hz, vinyl proton H<sub>a</sub>), 6.23 (dd, 1H, J = 18, J' = 11 Hz, vinyl proton H<sub>b</sub>), 5.97 (dd, 1H, J = 11 Hz, J' = 2 Hz, vinyl proton H<sub>c</sub>), 4.73 (s, 1H, H<sub>d</sub>), 4.94 (s, 1H, H<sub>e</sub>), 5.92 (br, s, vinyl proton H<sub>f</sub>), 3.73 (s, 3H, -COOCH<sub>3</sub>), 1.22 (s, 3H, -CH<sub>3</sub>) and 1.04 (s, 3H, -CH<sub>3</sub>); ms M<sup>+</sup> 250.1202 (calcd. for C<sub>14</sub>H<sub>18</sub>O<sub>4</sub>: 250.1205). Further elution with the same solvent system gave chloro compound 46 (38 mg, 9% yield): ir 1769 (enol ester), 1738

(ester), 1628 and 1602  $\text{cm}^{-1}$  ( $\text{C}=\text{C}$ ); nmr  $\delta$  5.92 (s, 1H,  $=\text{CH}_2$ ), 4.94 (s, 1H,  $=\text{CH}_2$ ), 4.73 (br, s, 1H,  $=\text{CH}-$ ), 3.84 (t, 2H,  $J = 7$  Hz,  $-\text{CH}_2\text{Cl}$ ), 3.70 (s, 3H,  $-\text{COOCH}_3$ ), 2.98 (t, 2H,  $J = 7$  Hz,  $-\text{CH}_2\text{CH}_2\text{Cl}$ ), 1.20 (s, 3H,  $-\text{CH}_3$ ) and 1.02 (s, 3H,  $-\text{CH}_3$ ); ms  $M^+$  286.0972 and 288.0967 (calcd. for  $\text{C}_{14}\text{H}_{19}\text{O}_4\text{Cl}$ : 286.0972). The aqueous layer was acidified with aqueous 1 N hydrochloric acid then extracted with dichloromethane. The extracts were combined, dried and concentrated to give a crude acid (236 mg, 57% yield). A mixture of the acid and potassium carbonate (280 mg, 2.03 mmol) in acetone (10 mL) was stirred at room temperature under argon atmosphere for 40 min. Methyl iodide (0.27 mL, 4.4 mmol) was added and the mixture stirred for an additional 6 h. The reaction mixture was acidified with aqueous 1 N hydrochloric acid solution then extracted with dichloromethane to give ester 48 (243 mg, 98% yield) after concentration and purification by flash chromatography (15% ethyl acetate in petroleum ether): ir  $\nu$  1737 (ester) and 1640  $\text{cm}^{-1}$  ( $\text{C}=\text{C}$ ); nmr  $\delta$  3.70 (s, 3H,  $-\text{COOCH}_3$ ), 3.74 (dd, 1H,  $J = 8$ ,  $J' = 5$  Hz,  $-\text{CHHO}-$ ), 3.72 (dd, 1H,  $J = 8$ ,  $J = 3$  Hz,  $-\text{CHHO}-$ ), 1.90 (d, 2H,  $J = 7$  Hz,  $-\text{CH}_2-$ ), 0.92 (s, 3H,  $-\text{CH}_3$ ) and 0.75 (s, 3H,  $-\text{CH}_3$ );  $^{13}\text{C}$  nmr  $\delta$  173.93, 171.71, 121.29, 116.54, 58.87, 51.69, 51.41, 50.13, 46.53, 41.65, 34.72,

32.2850, 17.5608, 16.0795 and 9.4641; ms  $M^+$  282.1469

(calcd. for  $C_{15}H_{22}O_5$ : 282.1477).

Reaction of 42 and acryloyl chloride in refluxing toluene

A mixture of enone 42 (386 mg, 1.98 mmol), potassium carbonate (200 mg, 1.45 mmol) and acryloyl chloride (3 mL) in toluene (6 mL) was refluxed under an argon atmosphere for 72 h. After this period an additional amount of acryloyl chloride (1 mL) in toluene was added and the mixture continued refluxing for another 6 h. The reaction mixture was cooled and washed with ice-cold 1 N hydrochloric acid solution and saturated sodium chloride solution. The organic extract was dried, filtered and concentrated. Flash chromatography of the residue on silica gel (15% ethyl acetate in petroleum ether) gave the required enol ester 45 (238 mg, 53% yield), chloro compound 46 (35 mg, 6% yield) and another mixture (62 mg) which was difficult to purify.

Claissen type rearrangement of enol ester 45

Enol ester 45 (124 mg, 0.5 mmol) in xylene (3 mL) was refluxed under an argon atmosphere for 1 h. After this period, 1-pentanol (1 mL, 92 mmol) was added and refluxing was continued for an additional 48 h. The solution was cooled to room temperature and acidified with ice cold 1 N hydrochloric acid solution then extracted with dichloro-

methane. The organic extracts were combined, dried, filtered and concentrated. The residue was chromatographed on silica gel eluting with 15% ethyl acetate in petroleum ether to give diester 52 (56 mg, 33% yield): ir  $1738\text{ cm}^{-1}$  (ester) and  $1642\text{ cm}^{-1}$  ( $\text{C}=\text{C}$ ); nmr  $\delta$  6.40 (s, 1H,  $=\text{CHH}$ ), 5.22 (s, 1H,  $=\text{CHH}$ ), 3.97 (t, 2H,  $J = 7\text{ Hz}$ ,  $-\text{OCH}_2-$ ), 3.70 (s, 3H,  $-\text{COOCH}_3$ ) 1.20 (s, 3H,  $-\text{CH}_3$ ), 1.04 (s, 3H,  $-\text{CH}_3$ ) and 0.92 (t, 3H,  $J = 7\text{ Hz}$ ,  $-\text{CH}_2\text{CH}_3$ ); ms  $M^+$  338.2093 (calcd. for  $\text{C}_{19}\text{H}_{30}\text{O}_5$ : 338.2093). Further elution with the same solvent system gave unreacted enol ester 45 (51 mg, 41% recovery).

5-Carbomethoxymethyl-4,4-dimethyl-7-oxo-1-oxaspiro[2.4]-heptane (57)

A solution of enone 42 (3.18 g, 16 mmol) in methanol (50 mL) was stirred at  $0^\circ\text{C}$  for 15 min. A solution of hydrogen peroxide (30% aqueous solution, 5.5 mL, 49 mmol) and lithium hydroxide monohydrate (105 mg, 2.4 mmol) was added slowly. The reaction mixture turned yellow then colourless. It was stirred at  $0^\circ\text{C}$  for 20 min then at room temperature under an atmosphere of argon for 3 h. Ice cold water and diluted hydrochloric acid solution were added and then the solution extracted with methyl chloride. The organic solution was dried, filtered and concentrated to give a 1:1 mixture of epimeric epoxides 57

(2.97 g, 86% yield). Flash chromatography of the mixture on silica gel eluting with 25% ethyl acetate in petroleum ether gave one of the epoxides: ir 1742 (ketone) and 1738  $\text{cm}^{-1}$  (ester); nmr  $\delta$  3.76 (s, 3H,  $-\text{COOCH}_3$ ), 3.15 (d, 1H,  $J = 6$  Hz,  $-\text{OCHH}-$ ), 2.85 (d, 1H,  $J = 6$  Hz,  $-\text{OCHH}-$ ), 1.04 (s, 3H,  $-\text{CH}_3$ ) and 0.98 (s, 3H,  $-\text{CH}_3$ ); ms  $M^+$  212.1043 (calcd. for  $C_{11}\text{H}_{16}\text{O}_4$ : 212.1049). Anal. Calcd. for  $C_{11}\text{H}_{16}\text{O}_4$ : C 62.23, H 7.60; Found: C 62.20, H 7.72. Further elution with the same solvent system gave the other epoxide 57: ir 1742 (ketone) and 1736  $\text{cm}^{-1}$  (ester); nmr  $\delta$  3.72 (s, 3H,  $-\text{COOCH}_3$ ), 3.02 (d, 1H,  $J = 6$  Hz,  $-\text{OCHH}-$ ), 2.96 (d, 1H,  $J = 6$  Hz,  $-\text{OCHH}-$ ), 1.00 (s, 3H,  $-\text{CH}_3$ ) and 0.94 (s, 3H,  $-\text{CH}_3$ ); ms  $M^+$  212.1046 (calcd. for  $C_{11}\text{H}_{16}\text{O}_4$ : 212.1049).

4-Carbomethoxymethyl-3,3-dimethylcyclopentanone (53) and 5-carbomethoxymethyl-6,6-dimethyl-3-methoxy-2-cyclohexone (59)

To a hot solution of epoxides 57 (8.17 g, 38.5 mmol) in methanol (25 mL) was added slowly a solution of sodium hydroxide (5.30 g, 133 mmol) in methanol (35 mL). The mixture was refluxed for 2 h. Water (5 mL) was added and refluxing was continued for an additional 72 h. The solution was allowed to cool to room temperature, acidified with 6 N hydrochloric acid solution and

extracted with methylene chloride. The organic extracts were dried, filtered and concentrated giving a crude acid (5.84 g): ir 3500-2500 (acid), 1737 (ketone), and 1709  $\text{cm}^{-1}$  (acid). To a solution of this acid in acetone (50 mL), anhydrous potassium carbonate (16.1 g, 116.5 mmol) was added. The mixture was stirred at room temperature under an argon atmosphere for 1 h and methyl iodide (6.4 mL, 103 mmol) added. After a gentle reflux overnight, the reaction mixture was poured into ice-cold water and extracted with methylene chloride. The organic solution was dried, filtered and concentrated. Flash chromatography of the residue on silica gel, eluting with 15% ethyl acetate in petroleum ether gave ester 53 (3.82 g, 54% yield): ir 1740  $\text{cm}^{-1}$  (ketone and ester);  $^1\text{H}$  nmr,  $\delta$  3.70 (s, 3H,  $-\text{COOCH}_3$ ), 2.14 (s, 2H,  $-\text{COCH}_2-\text{C}(\text{CH}_3)_2$ ), 1.18 (s, 3H,  $-\text{CH}_3$ ) and 0.92 (s, 3H,  $-\text{CH}_3$ ); ms  $M^+$  184.1097 (calcd. for  $\text{C}_{10}\text{H}_{16}\text{O}_3$ : 184.1099). Anal. Calcd. for  $\text{C}_{10}\text{H}_{16}\text{O}_3$ : C 65.19, H 8.75; Found: C 65.24, H 8.58. Further elution with the same solvent system gave compound 59 (1.72 g, 20% yield): ir 1730 (ester), 1710 (ketone) and 1620  $\text{cm}^{-1}$  ( $=\text{C}=\text{C}$ ); nmr  $\delta$  7.15 (s, 1H,  $=\text{CH}$ ), 3.86 (s, 3H,  $-\text{COOCH}_3$ ), 3.72 (s, 3H,  $-\text{OCH}_2$ ), 2.52 (m, 2H,  $-\text{CH}_2-$ ), 2.30 (m, 2H,  $-\text{CH}_2-$ ), 2.10 (dd, 1H,  $J = 11$ ,  $J' = 7$  Hz,  $=\text{CCHH}-$ ), 1.32 (s, 3H,  $-\text{CH}_3$ ) and 1.04 (s, 3H,  $-\text{CH}_3$ ); ms  $M^+$  226.1204 (calcd. for  $\text{C}_{12}\text{H}_{18}\text{O}_4$ : 226.1205).  $^{13}\text{C}$  nmr:  $\delta$  203.33, 172.13,

154.70, 123.39, 61.15, 50.52, 42.02, 40.68, 40.57, 33.23,  
24.75 and 20.14.

When the reaction was carried out at 0°C, keto ester 59 was isolated as the only product.

2-Hydroxymethyl-3,3-dimethyl-4-carbomethoxymethylcyclopentanone (62) and keto ester 63

To a solution of epoxides 57 (145 mg, 0.68 mmol) in acetic acid (5 mL) was added zinc metal (133 mg, 2.04 mmol). The reaction was refluxed for 2 h under an argon atmosphere. The mixture was cooled to room temperature then filtered. The filtrate was diluted with ether, washed with aqueous sodium bicarbonate solution, aqueous 1 N hydrochloric acid and water. The organic extract was dried, filtered and concentrated. Flash chromatography of the residue on silica gel, eluting with 20% ethyl acetate in petroleum ether gave keto ester 63 (44 mg, 33% yield): ir (neat) 1739 cm<sup>-1</sup> (ketone and ester); nmr δ 3.70 (s, 3H, -COOCH<sub>3</sub>), 2.04 (q, 1H, J = 8.5 Hz, -CHCO-), 1.12 (s, 3H, -CH<sub>3</sub>), 0.96 (d, 3H, J = 8.5, CH<sub>3</sub>) and 0.85 (s, 3H, -CH<sub>3</sub>); ms M<sup>+</sup> 198.1262 (calcd. for C<sub>11</sub>H<sub>18</sub>O<sub>3</sub>: 198.1265). Further elution using the same solvent system afforded hydroxy ester 62 (20 mg, 14% yield): ir (neat) 3540 (alcohol), 1740 (ketone and ester); nmr δ 3.70 (m, 5H, -CH<sub>2</sub>OH, -COOCH<sub>3</sub>), 1.06 (s, 3H, -CH<sub>3</sub>) and 0.84 (s, 3H,

$-\text{CH}_3$ ); ms  $M^+$  214.1206 (calcd. for  $\text{C}_{11}\text{H}_{18}\text{O}_4$ : 214.1205).

Unreacted epoxides 57 were recovered (74 mg, 51% recovery).

#### Hydroxy ester 62

To a solution of epoxides 57 (452 mg, 213 mmol) in benzene (10 mL) was added acetic acid (1 mL) and zinc metal (418 mg, 6.40 mmol). The reaction was refluxed for 2 h under an atmosphere of argon. The extraction and purification were performed as before. Hydroxy ester 62 (172 mg, 38% yield) and epoxides 57 (207 mg, 49% recovery) were obtained. All the spectra data were identical to the ones obtained previously.

#### Treatment of epoxides 57 with sodium hydride

To a suspension of sodium hydride (80% dispersion in oil; 202 mg, 6.73 mmol), freed from mineral oil by washing with petroleum ether ( $3 \times 5$  mL), in benzene (10 mL), a solution of epoxides 57 (1.1 g, 5.2 mmol) in benzene (5 mL) was added. After stirring for 1 h under an argon atmosphere at room temperature, the reaction mixture was heated at reflux for an additional 14 h. Ice-cold water and diluted hydrochloric acid solution were added and the mixture extracted with methylene chloride. The combined organic extracts were washed with saturated aqueous sodium chloride solution, dried, filtered and concentrated.

Flash chromatography of the residue on silica gel, eluting with 15% ethyl acetate in petroleum ether gave keto ester 53 (167 mg, 18% yield) and unreacted epoxides 57 (810 mg, 74% recovery). The spectral data were identical to those obtained previously (*vide supra*).

2-Carboethoxyethyl-3-carbomethoxymethyl-4,4-dimethylcyclopentanone (65)

To sodium hydride (36.4 mg, 1.21 mmol) in 1,2-dimethoxyethane (5 mL) was added keto ester 53 (216 mg, 1.17 mmol). The mixture was stirred at room temperature for 1 h under an argon atmosphere. Ethyl acrylate (0.15 mL, 1.38 mmol) was then added and stirring continued for additional 24 h. The mixture was taken up in dichloromethane (5 mL) and washed with dilute hydrochloric acid, water and saturated aqueous sodium chloride solution. The organic layer was dried, filtered and concentrated. Flash chromatography of the residue on silica gel, eluting with 15% ethyl acetate in petroleum ether gave diester 65 (276 mg, 83% yield); ir  $1739\text{ cm}^{-1}$  (esters and ketone); nmr,  $\delta$  4.12 (q, 2H,  $J = 8$  Hz,  $-\text{COOCH}_2\text{CH}_3$ ), 3.72 (s, 3H,  $-\text{COOCH}_3$ ), 2.18 (s, 2H,  $-\text{COCH}_2\text{C}-$ ), 1.32 (t, 3H,  $J = 8$  Hz,  $-\text{COOCH}_2\text{CH}_3$ ), 1.08 (s, 3H,  $-\text{CH}_3$ ) and 0.94 (s, 1H,  $-\text{CH}_3$ ); ms  $M^+$  284.1624 (calcd. for  $C_{15}\text{H}_{24}\text{O}_5$ : 284.1624).

2,5-Dicarboethoxyethyl-3-carbomethoxymethyl-1,4-dimethyl-cyclopentanone (71)

To a suspension of sodium hydride (111 mg, 3.7 mmol) in 1,2-dimethoxyethane (10 mL) was added keto ester 53 (566 mg, 3.1 mmol). The mixture was stirred at room temperature for 20 min under an argon atmosphere. After this period, the temperature was raised to 60°C and ethyl acrylate (0.44 mL, 4.1 mmol) added dropwise. The mixture was stirred for additional 3 h, cooled to room temperature, taken up in dichloromethane and washed with 1 N hydrochloric acid solution and saturated aqueous sodium chloride solution. The organic layer was dried, filtered and concentrated. Flash chromatography of the residue on silica gel, eluting with 20% ethyl acetate in petroleum ether gave triester 71 (898 mg, 76% yield); ir (neat) 1740 cm<sup>-1</sup> (ketone and esters); nmr δ 4.12 (q, 4H, J = 8 Hz, 2 x -COO-CH<sub>2</sub>CH<sub>3</sub>), 3.65 (s, 3H, -COOCH<sub>3</sub>), 2.27 (t, 1H, J = 6.5 Hz, -COCH-) and 1.22-1.98 (set of seven peaks, 12H, 2 x -CH<sub>3</sub>, 2 x -OCH<sub>2</sub>CH<sub>3</sub>); ms M<sup>+</sup> 384.2148 (calcd. for C<sub>20</sub>H<sub>32</sub>O<sub>7</sub>: 384.2148).

2-Carboethoxyethyl-3-carbomethoxymethyl-1,1-ethylene-dithio-4,4-dimethylcyclopentane (69)

At 0°C, to a solution of diester 65 (227 mg, 1.23 mmol) in dichloromethane (5 mL) were sequentially added

1,2-ethanedithiol (0.2 mL, 2.61 mmol) and boron trifluoride etherate (0.1 mL, 0.8 mmol). After stirring under an argon atmosphere for 6 h, ice-cold 1 N aqueous potassium hydroxide solution was added and the resulting mixture extracted with methylene chloride. The organic extracts were washed with water, dried, filtered and concentrated. Flash chromatography of the residue on silica gel, eluting with 10% ethyl acetate in petroleum ether gave thioketal **69** (235 mg, 82% yield): ir (neat) 1737 cm<sup>-1</sup> (esters); nmr δ 4.14 (q, 2H, J = 7.5 Hz, -O-CH<sub>2</sub>CH<sub>3</sub>), 3.70 (s, 3H, -COOCH<sub>3</sub>), 3.33 (m, 4H, -SCH<sub>2</sub>CH<sub>2</sub>S-), 2.28 (s, 2H, -CH<sub>2</sub>C-), 1.28 (t, 3H, J = 7.5 Hz, -OCH<sub>2</sub>CH<sub>3</sub>), 1.04 (s, 3H, -CH<sub>3</sub>) and 0.94 (s, 3H, -CH<sub>3</sub>); ms M<sup>+</sup> 360.1432 (calcd. for C<sub>17</sub>H<sub>28</sub>S<sub>2</sub>O<sub>4</sub>: 360.1429).

4-Carboethoxy-7,7-ethylenedithio-9,9-dimethylbicyclo-[4.3.0]nonan-3-one (**70**)

To a suspension of sodium hydride (80% dispersion in oil, 46 mg, 1.53 mmol) in toluene (10 mL), a solution of thioester **69** (429 mg, 1.19 mmol) in toluene (1 mL) and two drops of methanol were added. The reaction mixture was then refluxed for 5 h under an argon atmosphere. The reaction mixture was allowed to cool to room temperature, acidified with dilute aqueous hydrochloric acid solution and extracted with dichloromethane. The organic extracts

were dried, filtered and concentrated. Flash chromatography of the residue on silica gel, eluting with 5% ethyl acetate in petroleum ether afforded thiolester **70** (266 mg, 68% yield); ir 1735 (ketone and ester), 1656 and 1610  $\text{cm}^{-1}$  ( $\beta$ -hydroxy  $\alpha,\beta$ -unsaturated ester); nmr  $\delta$  4.10 (q, 2H,  $J = 7$  Hz,  $-\text{O}-\text{CH}_2\text{CH}_3$ ), 3.27 (m, 4H,  $-\text{SCH}_2\text{CH}_2\text{S}-$ ), 2.26 (s, 2H,  $-\text{CH}_2\text{C}-$ ), 1.27 (t, 3H,  $J = 7$  Hz,  $-\text{OCH}_2\text{CH}_3$ ), 1.08 (s, 3H,  $-\text{CH}_3$ ) and 0.92 (s, 3H,  $-\text{CH}_3$ ); ms  $M^+$  328.1169 (calcd. for  $\text{C}_{16}\text{H}_{24}\text{O}_3\text{S}_2$ : 328.1167).

#### 4-Carboethoxy-9,9-dimethylbicyclo[4.3.0]nona-3,7-dione

(66)

A solution of the dithiane **70** (219 mg, 0.68 mmol) in aqueous 80% acetonitrile (2 mL) was added at 25°C to a stirring solution of mercuric chloride (399 mg, 1.47 mmol), calcium carbonate (147 mg, 1.47 mmol) in aqueous 80% acetonitrile (5 mL). The mixture was heated at reflux under an argon atmosphere for 4 h. The solution was then cooled and filtered. The precipitate was washed thoroughly with 1:1 hexane/dichloromethane. The organic phase of the filtrate was washed with 5 M aqueous ammonium acetate, water, saturated sodium chloride solution then dried, filtered and concentrated. The residue was purified by flash chromatography over silica gel, eluting with 10% ethyl acetate in petroleum ether to afford diketo

ester 66: (75.7 mg, 45% yield); ir 1738 (carbonyl), 1656 and 1601 cm<sup>-1</sup> (enolized  $\beta$ -keto ester); nmr  $\delta$  4.08 (q, 2H, J = 7.5 Hz, -O-CH<sub>2</sub>CH<sub>3</sub>), 2.18 (s, 2H, -CH<sub>2</sub>C-), 1.28 (t, 3H, J = 7.5 Hz, -OCH<sub>2</sub>CH<sub>3</sub>), 1.22 (s, 3H, -CH<sub>3</sub>) and 1.00 (s, 3H, -CH<sub>3</sub>); ms M<sup>+</sup> 252.1362 (calcd. for C<sub>14</sub>H<sub>20</sub>O<sub>4</sub>: 252.1361).

2-Carboethoxyethyl-3-carbomethoxymethyl-4,4-dimethyl pentanone (65) Using LDA as a base

A solution of 1.35 M methylolithium (1.46 mL, 1.97 mmol) in hexane was added at -78°C to a solution of diisopropylamine (0.279 mL, 1.99 mmol) in dry tetrahydrofuran (10 mL) under an argon atmosphere. After stirring for 40 min, a solution of keto ester 53 (303 mg, 1.65 mmol) in 2 mL of tetrahydrofuran was added dropwise and stirring was continued for an additional 1 h. Ethyl acrylate (0.214 mL, 1.98 mmol) was added and the reaction mixture allowed to warm up to room temperature over 4 h period. The mixture was taken up in dichloromethane then washed with aqueous 1 N hydrochloric acid solution and aqueous saturated sodium chloride solution. The organic layer was dried, filtered and concentrated. On purification by flash chromatography (silica gel, 15% ethyl acetate in petroleum ether), diester 65 (401 mg, 86% yield) was produced. The ir, nmr and mass spectra were identical to those obtained previously.

2-Carbomethoxyethyl-3-carbomethoxymethyl-4,4-dimethyl-  
cyclopentanone (72) and 4-carbomethoxy-9,9-dimethyl-  
bicyclo[4.3.0]nona-3,7-dione (73)

A solution of 1.20 M methyllithium (2.34 mL, 2.81 mmol) in hexane was added at -78°C to a solution of diisopropylamine (0.393 mL, 2.80 mmol) in 10 mL of dry tetrahydrofuran under an argon atmosphere. After stirring for 40 min, a solution of keto ester 53 (430 mg, 2.34 mmol) in 2 mL of tetrahydrofuran was added dropwise followed by hexamethylphosphoramide (0.488 mL, 2.80 mmol) and stirring was continued for 1 h. Methyl acrylate (0.231 mL, 2.56 mmol) was added and the reaction mixture let to warm up to 0°C over a 5 h period. The reaction mixture was taken up in dichloromethane, washed with aqueous 1 N hydrochloric acid solution and saturated aqueous sodium chloride solution. The organic layer was dried, filtered and concentrated. Purification by flash chromatography on silica gel, eluting with 10% ethyl acetate in petroleum ether gave diketo ester 73 (192 mg, 35% yield). ir 1738 cm<sup>-1</sup> (carbonyl), 1659 and 1604 cm<sup>-1</sup> ( $\beta$ -hydroxy  $\alpha,\beta$ -unsaturated ester); nmr  $\delta$  13.56 (s, 1H,  $=C-OH$ ), 3.77 (s, 3H,  $-COOCH_3$ ), 2.12 (s, 2H,  $-COCH_2C-$ ), 1.20 (s, 3H,  $-CH_3$ ) and 1.02 (s, 3H,  $-CH_3$ ); <sup>13</sup>C nmr:  $\delta$  215.9022, 172.7809, 172.0691, 96.6322, 53.7887, 51.3241, 47.2978, 46.9170, 35.4030, 29.7600, 27.7923, 22.2262,

21,7813; ms  $M^+$  238.1205 (calcd. for  $C_{13}H_{18}O_4$ :

238.12038). Anal. Calcd. for  $C_{13}H_{18}O_4$ : C 65.51, H 7.56;

Found: C 65.29, H 7.42. Further elution using 10% ethyl

acetate in petroleum ether gave keto diester 72 (273 mg,

43%): ir 1739  $\text{cm}^{-1}$  (ketone and ester); nmr  $\delta$  3.75, 3.72,

3.70 and 3.68 (s, total 6H, 2 x  $-\text{COOCH}_3$ ), 2.18 (s, 2H,

$-\text{COCH}_2\text{C}-$ ), 1.14, 1.13, 0.95 and 0.94 (s, total 6H, 2 x

$-\text{CH}_3$ ); ms  $M^+$  270.1467 (calcd. for  $C_{14}H_{22}O_5$ : 270.1467).

Compounds 72, 73 and 4-carbomethoxyethyl-4-carbomethoxy-  
9,9-dimethylbicyclo[4.3.0]nona-3,7-dione (74)

A solution of 1.35 M methylolithium (7.40 mL, 9.98 mmol) in hexane was added at -78°C to a solution of diisopropylamine (1.40 mL, 9.98 mmol) in 25 mL of dry tetrahydrofuran under an argon atmosphere. After stirring for 40 min, a solution of keto ester 53 (1.67 g, 9.08 mmol) in tetrahydrofuran (5 mL) was added dropwise followed by hexamethylphosphoramide (1.74 mL, 9.98 mmol) then warmed up to -40°C. Methyl acrylate (0.90 mL, 9.98 mmol) was added and the reaction mixture left stirring for 2 h. After this period, the mixture was again warmed up to 0°C over a 4 h period. The mixture was taken up in dichloromethane, washed thoroughly with ice-cold aqueous 1 N hydrochloric acid and saturated sodium chloride solution. The organic layer was dried, filtered and

concentrated. Flash chromatography of the residue on silica gel, eluting with 10% ethyl acetate in petroleum ether gave: diketo ester 73 (1.03 g, 48% yield) and keto diester 72 (569 mg, 23% yield). The spectra data were identical to those obtained previously (vide supra).

Further elution using 15% ethyl acetate in petroleum ether gave diketo ester 74 (247 mg, 18% yield): ir 1739 (esters and five membered ketone) and 1713  $\text{cm}^{-1}$  (cyclohexanone); nmr  $\delta$  3.75 (s, 1H,  $-\text{COOCH}_3$ ), 3.65 (s, 3H,  $-\text{COOCH}_3$ ), 2.30 (s, 2H,  $-\text{COCH}_2\text{C}-$ ), 1.18 (s, 3H,  $-\text{CH}_3$ ) and 0.92 (s, 3H,  $-\text{CH}_3$ ); ms  $M^+$  324.1573 (calcd. for  $\text{C}_{17}\text{H}_{24}\text{O}_6$ : 324.1572).

#### Keto diester 72 from methyl $\beta$ -bromopropionate

A solution of 1.18 M methyl lithium (6.01 mL, 7.09 mmol) in hexane was added at  $-78^\circ\text{C}$  to a solution of diisopropylamine (0.99 mL, 7.09 mmol) in 20 mL of dry tetrahydrofuran under an argon atmosphere. After stirring for 1 h, a solution of keto ester 53 (870 mg, 4.73 mmol) in tetrahydrofuran (3 mL) was added dropwise followed by hexamethylphosphoramide (1.23 mL, 7.09 mmol) and the mixture was stirred for 1 h. Methyl  $\beta$ -bromopropionate (0.671 mL, 6.15 mmol) was added and the mixture warmed up to room temperature over a period of 10 h. The mixture was taken up in dichloromethane, washed with aqueous 2 N hydrochloric acid solution and saturated solution of sodium

chloride. The organic layer was dried, filtered and concentrated. Purification by flash chromatography over silica gel, eluting with 10% ethyl acetate in petroleum ether gave diketo ester 73 (232 mg, 14% yield), keto ester 53 (342 mg, 39% recovery) and keto diester 72 (523 mg, 42% yield). All the spectral data were identical to those recorded previously (*vide supra*).

4-Carbomethoxy-4-formylethyl-9,9-dimethylbicyclo[4.3.0]-nona-3,7-dione (77)

To a solution of diketo ester 73 (37 mg, 0.16 mmol) in 1,2-dimethoxyethane (2 mL) was added 1,4-diazabicyclo[2.2.2]octane (20.3 mg, 0.18 mL). After stirring for 20 min under an argon atmosphere at room temperature, acrolein (0.0143 mL, 0.21 mmol) was added. Stirring was continued for 4 h. The reaction mixture was taken up in ether (5 mL) and washed with aqueous 1 N hydrochloric acid solution. The organic layer was dried, filtered and concentrated. Flash chromatography of the residue on silica gel eluting with 15% ethyl acetate in petroleum ether afforded adduct 77 (41 mg, 90% yield): ir 2860, 2720 (aldehyde), 1740 (ester, cyclopentanone) and 1713  $\text{cm}^{-1}$  (aldehyde and cyclohexanone); nmr  $\delta$  9.66 (br, s, 1H, -CHO), 3.72 (s, 3H, -COOCH<sub>3</sub>), 1.08 (s, 3H, -CH<sub>3</sub>) and 1.02

(s, 3H,  $-\text{CH}_3$ ); ms  $M^+$  294.1466 (calcd. for  $\text{C}_{16}\text{H}_{22}\text{O}_5$ : 294.1467).

4-Carbomethoxy-4,9,9-trimethylbicyclo[4.3.0]nona-3,7-dione  
(78 and 79)

Potassium hydride (35% dispersion in mineral oil, 272 mg, 2.37 mmol) was washed ( $3 \times 3$  mL) under an argon atmosphere with petroleum ether and then tetrahydrofuran (10 mL) added. Diketo ester 73 (502 mg, 2.11 mmol) in tetrahydrofuran (2 mL) was added via a glass syringe and the resulting mixture stirred for 25 min at 0°C. The enolate was treated with iodomethane (0.146 mL, 2.35 mmol) then stirred for 8 h at room temperature. The reaction mixture was diluted with water then extracted with ether. The organic extracts were combined, washed with saturated sodium chloride solution, dried, filtered and concentrated. Flash chromatography of the residue on silica gel eluting with 15% ethyl acetate in petroleum ether gave a mixture of diastereomers 78 and 79 (ca. 2:1 ratio by nmr analysis; 492 mg, 93% yield): ir 1741 (cyclopentanone and ester) and  $1710\text{ cm}^{-1}$  (cyclohexanone); ms  $M^+$  252.1361 (calcd. for  $\text{C}_{14}\text{H}_{20}\text{O}_4$ : 252.1363). Anal. Calcd. for  $\text{C}_{14}\text{H}_{20}\text{O}_4$ : C 66.67, H 7.94; Found: 66.53, H 7.84. Diketo ester 78 (m.p., 75–76°C) was separated from 79 by recrystallization from ethyl acetate/petroleum

ether. The following nmr data was obtained for 78:

$\delta$  3.74 (s, 3H, -COOCH<sub>3</sub>), 2.54 (dd, J = 15, J' = 5 Hz, 1H<sub>b</sub>), 2.48 (dd, J = 15, J' = 13 Hz, 1H<sub>a</sub>), 2.35 (dd, J = 19, J' = 13 Hz, 1H<sub>g</sub>), 2.26 (t, J = 13 Hz, 1H<sub>e</sub>), 2.46 (ddd, J = J' = 13, J'' = 5 Hz, 1H<sub>d</sub>), 1.90 (ddd, J = J' = 13, J = 5 Hz, 1H<sub>c</sub>), 1.43 (s, 3H, -CH<sub>3</sub>), 1.17 (s, 3H, -CH<sub>3</sub>), 1.07 (s, 3H, -CH<sub>3</sub>).

Diketo esters 78 and 79 by use of potassium carbonate as a base

Potassium carbonate (582 mg, 4.21 mmol) in acetone (10 mL) was stirred for 40 min at room temperature under argon atmosphere. Sequentially keto ester 73 (334 mg, 0.40 mmol) in acetone (2 mL) and iodomethane (0.174 mL, 2.79 mmol) were added and the reaction mixture continued stirring for 24 h. After this period ice-cold dilute hydrochloric acid solution was added and then extracted with dichloromethane. The organic extracts were combined, dried, filtered and concentrated. The residue was chromatographed on silica gel eluting with 15% ethyl acetate in petroleum ether to afford diastereomeric esters 78 and 79 (293 mg, 82% yield). The unreacted 73 was also obtained (43 mg, 15% recovery). All the spectra data were identical to those obtained previously (vide supra).

Conversion of 78 to 79

To a solution of diketo ester 78 (122 mg, 0.48 mmol) in acetone (4 mL) was added potassium carbonate (134 mg, 0.97 mmol). The mixture was refluxed for 20 h under an atmosphere of argon. Aqueous 1 N hydrochloric acid solution was added and the resulting solution extracted with dichloromethane. The organic extracts were dried, filtered and concentrated. The residue was chromatographed on silica gel. Elution with a solution of 10% ethyl acetate in petroleum ether gave a mixture of epimers 78 and 79 (ca: 1.3 ratio by nmr analysis, 113 mg, 93% yield).

4-Carbomethoxy-3-hydroxy-4,9,9-trimethylbicyclo-[4.3.0]nonan-7-one (84)

A solution of diketo ester 78 (80.6 mg, 0.320 mmol) and lithium tri-tert-butoxyaluminium hydride (97.5 mg, 383 mmol) in tetrahydrofuran/ether (1:3, 4 mL) was stirred at room temperature under an atmosphere of argon for 12 h. Water was added and the resulting solution extracted with dichloromethane. The extracts were dried, filtered and concentrated. The residue was chromatographed on silica gel eluting with 15% ethyl acetate in petroleum ether to afford hydroxy esters 84 (78.68 mg, 97% yield): ir 3480 (alcohol) and 1740  $\text{cm}^{-1}$  (ketone and ester); nmr  $\delta$  3.94

(br, t, 1H, -CH-OH), 3.74 (s, 3H, -COOCH<sub>3</sub>), 2.18 (d, 1H, J = 13 Hz, -COCHH-), 2.09 (d, 1H, J = 13 Hz, -COCHH-), 1.14, 1.12, 1.08, 1.06, 1.03, 1.00 (s, total 9H, 3 x -CH<sub>3</sub>); ms M<sup>+</sup> 254.1522 (calcd. for C<sub>14</sub>H<sub>22</sub>O<sub>4</sub>: 254.1517).

4-Carbomethoxy-7,7-ethylenedithio-4,9,9-trimethylbicyclo-[4.3.0]nonan-3-ol (85)

At 0°C, to a solution of hydroxy esters 84 (78 mg, 0.307 mmol) in dichloromethane (3 mL) was added sequentially, 1,2-ethanedithiol (0.039 mL, 0.465 mmol) and boron trifluoride etherate (0.045 mL, 0.395 mmol). The resulting mixture was stirred at room temperature overnight under an argon atmosphere. Ice-cold aqueous 1 N potassium hydroxide solution was added and the resulting mixture extracted with methylene chloride. The organic extracts were washed with saturated sodium chloride solution, dried, filtered and concentrated. Flash chromatography of the residue on silica gel, eluting with 15% ethyl acetate in petroleum ether gave thioketal 85 (91.9 mg, 90% yield): ir 3480 (alcohol) and 1731 cm<sup>-1</sup> (ester); nmr δ 3.97 (br, m, 1H, -CH<sub>2</sub>CHOH), 3.70 (s, 3H, -COOCH<sub>3</sub>), 3.20 (m, 4H, -SCH<sub>2</sub>CH<sub>2</sub>S-), 1.20 (s, 3H, -CH<sub>3</sub>), 1.02 (s, 3H, -CH<sub>3</sub>) and 0.95 (s, 3H, -CH<sub>3</sub>); ms M<sup>+</sup> 330.1322 (calcd. for C<sub>16</sub>H<sub>26</sub>S<sub>2</sub>O<sub>3</sub>: 330.1325). Anal. Calcd. C 58.18, H 7.88, S 19.39; Found: C 58.03, H 7.97, S 19.33.

3-Carbomethoxy-4-hydroxy-3,7,7-trimethylbicyclo[4.3.0]-  
nonane (86)

Freshly prepared Raney nickel (0.5 mL, settled volume), was added to ester 85 (52 mg, 0.158 mmol) in ethanol (98%, 3 mL). The reaction mixture was stirred at room temperature under an argon atmosphere for 2 h. The reaction mixture was carefully filtered and residue washed with ethanol (2 x 1 mL). The filtrate was concentrated, then purified by flash chromatography on silica gel, eluting with 5% ethyl acetate in petroleum ether to afford hydroxy ester 86 (26 mg, 61% yield): ir 3480 (alcohol) and 1731 cm<sup>-1</sup> (ester); <sup>1</sup>nmr δ 3.97 (br, d, 1H, -CH<sub>OH</sub>), 3.67 (s, 3H, -COOCH<sub>3</sub>), 2.92 (br, s, 1H, -OH), 1.94 (dd, 1H, J = 13.5 Hz, J = 3.5 Hz, -CHHC(CH<sub>3</sub>)CO-), 1.16 (s, 3H, -CH<sub>3</sub>), 0.94 (s, 3H, -CH<sub>3</sub>) and 0.77 (s, 3H, -CH<sub>3</sub>); ms M<sup>+</sup> 240.1729 (calcd. for C<sub>14</sub>H<sub>24</sub>O<sub>3</sub>: 240.1725).

4-Carbomethoxy-4,9,9-trimethylbicyclo[4.3.0]nonan-3-one  
(87)

To a solution of hydroxy ester 86 (30 mg, 0.125 mmol) in dichloromethane (3 mL) were added pyridinium chlorochromate (40.4 mg, 0.186 mmol) and sodium acetate (0.46 mg, 3.5 μmol). The resulting mixture was stirred at room temperature under an argon atmosphere for 3 h. The mixture was filtered and the solid residue washed with

ether ( $3 \times 1$  mL). The filtrate was concentrated and purified by column chromatography eluting with 5% ethyl acetate in petroleum ether to give keto ester 87 (24 mg, 81% yield): ir 1739 (ester) and  $1709\text{ cm}^{-1}$  (cyclohexanone); nmr  $\delta$  3.65 (s, 3H,  $-\text{COOCH}_3$ ), 1.23 (s, 3H,  $-\text{CH}_3$ ), 1.16 (s, 3H,  $-\text{CH}_3$ ) and 1.04 (s, 3H,  $-\text{CH}_3$ ), ms  $M^+$  238.1569 (calcd. for  $C_{14}\text{H}_{22}\text{O}_3$ : 238.1569).

4-Carbomethoxy-7,7-ethylenedioxy-4,9,9-trimethylbicyclo-[4.3.0]nonan-3-ol (89)

p-Toluenesulfonic acid monohydrate (31 mg, 0.16 mmol) was dried by refluxing in benzene with azeotropic removal of water. To a solution of dry p-toluenesulfonic acid in benzene (2 mL), 2-methyl-2-ethyl-1,3-dioxolane (5 mL) and a solution of hydroxy ester 84 (278 mg, 1.09 mmol) in benzene (2 mL) were added. The mixture was heated at 80-90°C (oil bath temperature) for 4 h. After cooling to 0°C, 10% aqueous sodium bicarbonate solution was added and the mixture extracted with ether. The organic extracts were washed with water and saturated aqueous sodium chloride solution, dried, filtered and concentrated.

Flash chromatography of the residue on silica gel, eluting with 10% ethyl acetate in petroleum ether gave ketal 89 (261 mg, 80% yield): ir 3480 (alcohol), 1738 (ester), 1120, 1170 and  $1190\text{ cm}^{-1}$  (ketal); nmr  $\delta$  3.97 (m, 1H,

-CHOH) 3.87 (m, 4H, -OCH<sub>2</sub>CH<sub>2</sub>O-), 3.70 (s, 3H, -COOCH<sub>3</sub>), 3.68 (s, 3H, -COOCH<sub>3</sub>), 1.22, 1.20, 1.02, 1.00, 0.94, 0.93 (s, total 9H, 3 x -CH<sub>3</sub>); ms M<sup>+</sup> 298.1781 (calcd. for C<sub>16</sub>H<sub>26</sub>O<sub>5</sub>: 298.1781); Anal. Calcd. C 64.43, H 8.72; Found: C 64.41, H 8.72.

3-Carbomethoxy-9,9-ethylenedioxy-4-methylmercaptothiono-carboxyloxy-3,7,7-trimethylbicyclo[4.3.0]nonane (90)

At 0°C, to a suspension of sodium hydride (80% dispersion in oil, 119 mg, 3.97 mmol) in freshly distilled 1,2-dimethoxyethane (5 mL) was slowly added under an argon atmosphere a solution of alcohols **89** (236 mg, 0.79 mmol) in 1,2-dimethoxyethane (3 mL). After stirring for 10 min carbon disulfide (0.476 mL, 7.91 mmol) and methyl iodide (0.493 mL, 7.92 mmol) were added. The reaction was stirred at room temperature for 5 h and then cooled to 0°C. Water was added slowly to the reaction mixture and the resulting solution extracted with dichloromethane. Organic layer was washed with water, saturated aqueous sodium chloride then dried, filtered and concentrated. Column chromatography of the residue on silica gel, eluting with 7% ethyl acetate in petroleum ether gave pure **90** (268 mg, 87% yield): ir 1738 (ester), 1220, 1249 (C=S) and 1056 cm<sup>-1</sup> (C-O-C); nmr δ 3.88 (m, 4H, -OCH<sub>2</sub>CH<sub>2</sub>O-), 3.72 (s, 3H, -COOCH<sub>3</sub>), 2.48 (s, 3H, -SCH<sub>3</sub>), 1.24 (s, 3H,

$-\text{CH}_3$ ), 1.00 (s, 3H,  $-\text{CH}_3$ ) and 0.94 (s, 3H,  $-\text{CH}_3$ ); ms  $M^+$  388.1378 (calcd. for  $\text{C}_{18}\text{H}_{28}\text{S}_2\text{O}_5$ : 388.1376).

3-Carbomethoxy-9,9-ethylenedioxy-3,7,7-trimethylbicyclo-[4.3.0]nonane (91) from xanthate 90

Xanthate 90 (260 mg, 0.61 mmol), azobisisobutyronitrile (50 mg) and tri-n-butyltin hydride (0.37 mL, 1.34 mmol) were dissolved in freshly distilled toluene (7 mL). The solution was degassed by bubbling argon through it for 20 min. The resulting solution was refluxed for 3 h at 120°C (oil bath temperature). The reaction mixture was cooled to room temperature and concentrated. The residue on column chromatography over silica gel eluting with 5% ethyl acetate in petroleum ether gave pure 91 as colourless oil (147 mg, 78% yield): ir 1734  $\text{cm}^{-1}$  (ester); nmr  $\delta$  3.87 (m, 4H,  $-\text{OCH}_2\text{CH}_2\text{O}-$ ), 3.68 (s, 3H,  $-\text{COOCH}_3$ ), 1.78 (d, 1H,  $J = 8$  Hz,  $-\text{CHHC}-$ ), 1.72 (d,  $J = 8$  Hz,  $-\text{CHHC}-$ ), 1.21 (s, 3H,  $-\text{CH}_3$ ), 1.02 (s, 3H,  $-\text{CH}_3$ ) and 0.94 (s, 3H,  $-\text{CH}_3$ ); ms  $M^+$  282.1831 (calcd. for  $\text{C}_{16}\text{H}_{26}\text{O}_4$ : 282.1830); Anal. Calcd. for  $\text{C}_{16}\text{H}_{26}\text{O}_4$ : C 68.09, H 9.28; Found: C 68.34, H 9.47.

4-Carbomethoxy-4,9,9-trimethylbicyclo[4.3.0]nonan-7-one (88)

To a solution of p-toluenesulfonic acid monohydrate (40.5 mg, 0.21 mmol) in acetone (3 mL) were added ketal 91

(48.0 mg, 0.17 mmol) and 3 drops of water under an argon atmosphere. The reaction mixture was stirred at room temperature for 2 h. Aqueous 10% sodium bicarbonate solution was added and the resulting mixture extracted with ether. The organic extracts were washed with water and saturated sodium chloride solution, dried, filtered and concentrated. The residue on column chromatography over silica gel, eluting with 5% ethyl acetate in petroleum ether gave pure 88 (38.2 mg, 94%): ir 1741 cm<sup>-1</sup> (ester); nmr 3.68 (s, 3H, -COOCH<sub>3</sub>), 2.19 (s, 2H, -COCH<sub>2</sub>C-), 1.20 (s, 3H, -CH<sub>3</sub>), 1.16 (s, 3H, -CH<sub>3</sub>) and 1.04 (s, 3H, -CH<sub>3</sub>); ms M<sup>+</sup> 238.1562 (calcd. C<sub>14</sub>H<sub>22</sub>O<sub>3</sub>: 238.1565). Anal. Calcd. C 70.56, H 9.24; Found: C 70.54, H 9.16.

4-Carbomethoxy-3,3-ethylenedithiol-4,9,9-trimethylbicyclo-[4.3.0]nonan-7-one (92).

At 0°C, to a solution of diketo ester 78 (315 mg, 1.25 mmol) in dichloromethane (10 mL) were added sequentially, 1,2-ethanedithiol (0.105 mL, 1.25 mmol), boron trifluoride etherate (0.15 mL, 1.24 mmol). The reaction mixture was stirred at this temperature for 8 h under an argon atmosphere. Ice-cold aqueous 1 N potassium hydroxide solution was added and the resulting mixture extracted with methylene chloride. The organic extracts

were washed with saturated sodium chloride solution, dried, filtered and concentrated. Flash chromatography on silica gel, eluting with 10% ethylacetate in petroleum ether gave thioketal 92 (282 mg, 69% yield): ir 1740  $\text{cm}^{-1}$  (ester and ketone); nmr  $\delta$  3.72 (s, 3H, -COOCH<sub>3</sub>), 3.24 (m, 4H, -SCH<sub>2</sub>CH<sub>2</sub>S-), 2.22 (s, 2H, -COCH<sub>2</sub>C-), 1.48 (s, 3H, -CH<sub>3</sub>), 1.06 (s, 3H, -CH<sub>3</sub>) and 0.90 (s, 3H, -CH<sub>3</sub>); ms M<sup>+</sup> 328.1166 (calcd. for C<sub>16</sub>H<sub>24</sub>O<sub>3</sub>S<sub>2</sub>: 328.1166). Further elution using the same solvent system afforded a diketal (126 mg, 25% yield): ir 1731  $\text{cm}^{-1}$  (ester); nmr  $\delta$  3.70 (s, 3H, -COOCH<sub>3</sub>), 3.24 (m, 8H, -SCH<sub>2</sub>CH<sub>2</sub>S-), 1.48 (s, 3H, -CH<sub>3</sub>), 1.06 (s, 3H, -CH<sub>3</sub>) and 0.90 (s, 3H, -CH<sub>3</sub>); ms M<sup>+</sup> 404.0975 (calcd. for C<sub>18</sub>H<sub>28</sub>S<sub>4</sub>O<sub>2</sub>: 404.0974).

#### Keto ester 88 from thioketal 92

Freshly prepared Raney nickel (5 mL, settled volume) was added to thioketal 92 (308 mg, 0.94 mmol) in 98% ethanol (10 mL). The reaction mixture was stirred at room temperature under an argon atmosphere for 2 h. The mixture was then carefully filtered and residue washed with ethanol (2 x 2 mL). The filtrate was concentrated, then purified by flash chromatography eluting with 5% ethyl acetate in petroleum ether to afford keto ester 88 (115 mg, 52% yield). The spectra were identical to those previously obtained (*vide supra*).

9,9-Ethylenedioxy-3-hydroxymethyl-3,7,7-trimethylbicyclo-[4.3.0]nonane (96)

At 0°C, to a solution of 3.4 M sodium bis(2-methoxyethoxy)aluminium hydride (Red Al) (0.146 mL, 0.496 mmol) in ether (2 mL) was added ketal 91 (46.6 mg, 0.165 mmol) under an argon atmosphere. The reaction mixture was stirred at room temperature for 2 h. It was then cooled to 0°C and ice-cold water added dropwise. The resulting solution was extracted with dichloromethane and ether. The combined extracts were dried, filtered and concentrated. Column chromatography of the residue on silica gel, eluting with 15% ethyl acetate in petroleum ether gave hydroxy ketal 96 (39.8 mg, 95% yield): ir 3480 cm<sup>-1</sup> (alcohol); nmr δ 3.87 (m, 4H, -OCH<sub>2</sub>CH<sub>2</sub>O-), 3.76 (br, s, 2H, -CH<sub>2</sub>OH); 1.78 (s, 2H, -CH<sub>2</sub>C(CH<sub>3</sub>)<sub>2</sub>-), 1.16 (s, 3H, -CH<sub>3</sub>), 1.02 (s, 3H, -CH<sub>3</sub>) and 0.94 (s, 3H, -CH<sub>3</sub>); ms M<sup>+</sup> 254.1881 (calcd. for C<sub>15</sub>H<sub>26</sub>O<sub>3</sub>: 254.1881).

9,9-Ethylenedioxy-3-formyl-3,7,7-trimethylbicyclo[4.3.0]nonane (97)

A solution of pyridinium chlorochromate (44.6 mg, 0.207 mmol) and sodium acetate (3.4 mg, 0.041 mmol) in dry dichloromethane (3 mL) was stirred at room temperature under an argon atmosphere for 20 min. Hydroxy ketal 96 (35 mg, 0.138 mmol) was added and stirring continued for

1.5 h. The reaction mixture was filtered and washed with petroleum ether (2 mL). The filtrate was then concentrated. Column chromatography of the residue on silica gel, eluting with 15% ethyl acetate in petroleum ether afforded ketal aldehyde 97 (27 mg, 78% yield): ir 2830, (aldehyde) and 1739  $\text{cm}^{-1}$  (carbonyl); nmr  $\delta$  9.38 (br, s, -CHO), 3.86 (m, 4H, -OCH<sub>2</sub>CH<sub>2</sub>O-), 1.76 (s, 2H, -CH<sub>2</sub>C(CH<sub>3</sub>)<sub>2</sub>-), 1.10 (s, 3H, -CH<sub>3</sub>), 1.04 (s, 3H, -CH<sub>3</sub>) and 0.96 (s, 3H, -CH<sub>3</sub>); ms M<sup>+</sup> 252.1724 (calcd. for C<sub>15</sub>H<sub>24</sub>O<sub>3</sub>: 252.1725).

4-Carbomethoxy-6-carbomethoxyethyl-3,3-ethylenedithio-4,9,9-trimethylbicyclo[4.3.0]nonan-7-one (101)

A. Using sodium hydride as base

To a suspension of sodium hydride (80% dispersion in oil, 21.1 mg, 0.7 mmol) in 5 mL of tetrahydrofuran was added a solution of ketal 92 (192 mg, 0.5 mmol) in tetrahydrofuran (1 mL). The mixture was stirred for 30 min at room temperature under an argon atmosphere. Methyl acrylate (0.063 mL, 0.7 mmol) and a drop of tert-butanol were added and stirring was continued for 36 h. Water was added and the resulting solution extracted with dichloromethane. The extract was washed with aqueous saturated sodium chloride solution, dried, filtered and concentrated. Column chromatography of the residue on

silica gel, eluting with 10% ethyl acetate in petroleum ether gave pure starting material 92 (163 mg, 85% recovery). The ir, nmr and mass spectra were identical to those previously obtained. Further elution using the same solvent system afforded diester 101 (17 mg, 7% yield): ir 1739 cm<sup>-1</sup> (ketone and esters); nmr δ 3.72 (s, 3H, -COOCH<sub>3</sub>), 3.68 (s, 3H, -COOCH<sub>3</sub>), 3.25 (m, 4H, -SCH<sub>2</sub>CH<sub>2</sub>S-), 2.22 (m, 2H, -COCH<sub>2</sub>C-), 1.34 (s, 3H, -CH<sub>3</sub>), 1.25 (s, 3H, -CH<sub>3</sub>) and 1.14 (s, 3H, -CH<sub>3</sub>); ms M<sup>+</sup> 414.1535 (calcd. for C<sub>20</sub>H<sub>30</sub>S<sub>2</sub>O<sub>5</sub>: 414.1535).

B. Using potassium hydride as base

Potassium hydride (35% dispersion in mineral oil, 61.2 mg, 0.53 mmol) was washed with petroleum ether (3 x 3 mL) and tetrahydrofuran (5 mL) added. A solution of ketal 92 (146 mg, 0.45 mmol) and a drop of tert-butanol were added and the solution stirred under an atmosphere of argon at room temperature for 30 min. Methyl acrylate (0.048 mL, 0.53 mmol) was added and the solution stirred for 36 h. Ice-cold water was added and the resulting mixture extracted with dichloromethane. The extracts were washed with aqueous saturated sodium chloride solution, dried, filtered and concentrated. Flash chromatography of the residue on silica gel eluting with 10% ethyl acetate in petroleum ether afforded pure starting material 92

(105 mg, 72% recovery). Further elution using the same solvent system afforded the desired diester 101 (35 mg, 19% yield). The spectral data were identical to those reported previously.

## REFERENCES

1. W. Muhler, Ann., 176, 181 (1875).
2. O. Wallach and H. Bleibtreu, Ber., 12, 1061 (1879).
3. J. Obermeyer, Ber., 20, 2918 (1887).
4. H.L. Wheeler, Am. Chem. J., 24, 69 (1900).
5. O. Jeger, J. Norymberski, S. Szipilfogel, and V. Prelog, Helv. Chim. Acta, 29, 684 (1946).
6. M.L. Wolfrom and J.V. Karabinos, J. Am. Chem. Soc., 68, 1455 (1946).
7. A.V. McIntosh, Jr., E.M. Meinzer, and R.H. Levin, J. Am. Chem. Soc., 70, 2955 (1948).
8. A.V. McIntosh, Jr., A.M. Searcy, E.M. Meinzer, and R.H. Levin, J. Am. Chem. Soc. 71, 3317 (1949).
9. For a brief review, see L. Field, Synthesis, 101 (1972).
10. For a review, see Org. Compounds of Sulfur and Tellurium, Specialist Periodical Reports, Chemical Society, 4, 14 (1977).
11. T. Mukayama, T. Takeda, and K. Atsumi, Chem. Lett., 187 (1974).
12. T. Mukaiyama, Angew. Chem. Int. Ed. Engl. 18, 707 (1979).
13. H.J. Liu, S.P. Lee, and W.H. Chan, Synth. Commun. 9, 91 (1979).

14. H.J. Liu and S.I. Sabesan, Can. J. Chem., 58, 2645 (1980).
15. S. Yamada, Y. Yokoyama, and T. Shioiri, J. Org. Chem., 39, 3302 (1974).
16. A. Arieta, T. Garcia, J.M. Lago, and C. Palomo, Synth. Commun., 13 (6), 471 (1983).
17. M. Sekine, A. Kume, and T. Hata, Tetrahedron Lett., 3617 (1981).
18. S. Kim and S. Yang, Chem. Lett., 133 (1981).
19. S. Masamune, C.U. Kim, K.E. Wilson, G.O. Spessard, P.E. Georgiou, and G.S. Bates, J. Am. Chem. Soc., 97, 3512 (1975).
20. S. Masamune, S. Kamata, J. Diakur, Y. Sugihara, and G.S. Bates, Can. J. Chem., 53, 3693 (1975).
21. M. Ueda, K. Seki, and Y. Imai, Synthesis, 991 (1981).
22. H.J. Gais, Angew. Chem. Int. Ed. Engl., 16, 244 (1977).
23. S. Masamune, H. Yamamoto, S. Kamata, and A. Fukuzawa, J. Am. Chem. Soc., 97, 3513 (1975). }
24. K.C. Nicolau, Tetrahedron, 33, 683 (1977).
25. T.G. Back, Tetrahedron, 33, 3041 (1977).
26. E.J. Corey and N.C. Nicolau, J. Am. Chem. Soc., 46, 5614 (1974).
27. E.J. Corey, D.J. Brunelle, and P.J. Stork, Tetrahedron Lett., 3405 (1976).

28. H. Gerlach and A. Thalmann, *Helv. Chim. Acta*, **57**, 293 (1974).
29. H. Gerlach, K. Ortle, and A. Thalmann, *Helv. Chim. Acta*, **59**, 755 (1976).
30. H.J. Liu and H.K. Lai, *Tetrahedron Lett.*, 1193 (1979).
31. H.J. Liu, H.K. Lai, and S.K. Attah-Poku, *Tetrahedron Lett.*, 4121 (1979).
32. H.J. Liu and H.K. Lai, *Can. J. Chem.*, **57**, 2522 (1979).
33. H.J. Liu and I.V. Oppong, *Can. J. Chem.*, **60**, 94 (1982).
34. Y. Nagao, K. Kawabata, and E. Fujita, *Chem. Commun.*, (1978).
35. H.J. Liu, R.R. Bukownik, and P.R. Pednekar, *Synth. Commun.* **11**, 599 (1981).
36. H.J. Liu and H. Wynn, *Tetrahedron Lett.*, **23**, 3151 (1982).
37. H.J. Liu and H. Wynn, *Can. J. Chem.*, **64**, 649 (1986).
38. H.J. Liu and H. Wynn, *Can. J. Chem.*, **64**, 658 (1986).
39. J.L. Herrmann and R.H. Schlessinger, *Tetrahedron Lett.*, 2429 (1973).
40. G. Fräter, *Helv. Chim. Acta*, **62**, 2825 (1979); **62**, 2829 (1979).
41. G. Fräter, *Tetrahedron Lett.*, **22**, 425 (1981).
42. D. Seebach and D. Wasmuth, *Helv. Chim. Acta*, **63**, 197 (1980).

43. J.D. Aebi, M.A. Sutter, D. Wasmuth, and D. Seebach, Liebigs Ann. Chem., 2114 (1983).
44. A. Kramer and H. Pfander, Helv. Chim. Acta, 65, 495 (1982).
45. G. Fräter, U. Müller, and W. Gunther, Tetrahedron, 40, 1269 (1984).
46. J.P. Marino and D.M. Floyd, J. Am. Chem. Soc., 96, 7138 (1974).
47. C.G. Gordon-Gray and C.G. Whiteley, J. Chem. Soc., Perkin Trans. 1, 2040 (1977).
48. J. Ficini and J.C. Depezay, Tetrahedron Lett., 4797 (1969).
49. R.K. Boeckman, Jr. and M. Ramaiah, J. Org. Chem., 42, 1581 (1977).
50. J.C. Depezay and Y.L. Merrev, Tetrahedron Lett., 2751 (1978).
51. J.P. Marino and J.S. Fatina, Tetrahedron Lett., 3901 (1975).
52. P.A. Grieco, C.L.J. Wang, and G. Majetich, J. Org. Chem., 41, 726 (1976).
53. L.-C. Yu and B. Helquist, Tetrahedron Lett., 3423 (1978).
54. W.C. Still, M. Kahn, and A. Mitra, J. Org. Chem., 43, 2923 (1978).

55. R. Mazingo, *Organic Synthesis, Coll. Vol. 3*, 188 (1955).
56. R. McLafferty, *Anal. Chem.*, **28**, 306 (1956).
57. C. Alexandre and F. Rousset, *Bull. Soc. Chim. Fr.*, 1837 (1971).
58. D.F. Faber, *J. Org. Chem.*, **47**, 1351 (1982).
59. K.B. Kreibs, D. Heusser, and H. Wimmer, in "Thin layer chromatography: A Laboratory Handbook", 2nd Ed., E. Stahl, Ed., Springer-verlag, New York, 1969, P. 854.
60. L.M. Harwood, *Aldrichim. Acta*, **18**, 25 (1985).
61. L.F. Fieser and M. Fieser, "Reagents for Organic Synthesis", Vol. 1, Wiley and Sons Inc., New York, 1967.
62. J. Simonsen and D.H.R. Barton, "The Terpenes", Vol. III, 39 (1952).
63. C.F. Seidel, P.H. Müller, and H. Schinzz, *Helv. Chim. Acta*, **27**, 738 (1944).
64. G. Dupont, R. Dulou, and P. Nafta, *Bull. Soc. Chim.*, **15**, 990 (1948).
65. W. Treibs and G. Lössner, *Leibigs Ann.*, **634**, 124 (1960).
66. A.S. Gupta and S. Dev, *Tetrahedron*, **27**, 635 (1971).
67. R. Misra, R.C. Pandev, and S. Dev, *Tetrahedron*, **35**, 2301 (1979).

68. A.G. Gonzalez, B.M. Fraga, J.G. Luis, and A.G. Ravelo, *An. Quim.*, 70, 701 (1975).
69. B.B. Jarvis, N.B. Pena, S.N. Comezoglu, and M.M. Rao, *Phytochemistry*, 25, 533 (1986).
70. G. Delgado, L. Alvarez, A.R. de Vivar, *Phytochemistry*, 24, 2973 (1985).
71. G. Delgado, H. Cardenas, G. Pelaez, A.R. de Vivar, and R.P. Mirada, *J. Nat. Prod.* 47, 1042 (1984).
72. A. Aebei, D.H.R. Barton, A.W. Burgstahler, and A.S. Lindsey, *J. Chem. Soc.*, 4654 (1954).
73. A. Aebei, D.H.R. Barton, A.S. Lindsey, *J. Chem. Soc.*, 3124 (1953).
74. E.W. Warnhoff, *Can. J. Chem.*, 42, 1664 (1964).
75. A. Aebei, D.H.R. Barton, and A.S. Lindsey, *Chemistry and Industry*, 478 (1953).
76. N.W. Atwater and E.B. Reid, *Chemistry and Industry*, 278, 688 (1953).
77. E.J. Corey, R.B. Mitra, and H. Uda, *J. Am. Chem. Soc.*, 85, 3622 (1963).
78. E.J. Corey, R.B. Mitra, and H. Uda, *J. Am. Chem. Soc.*, 86, 485 (1964).
79. Y. Ohtsuka, S. Nituma, H. Tadokoro, T. Hayashi, and T. Oishi, *J. Org. Chem.*, 49, 2326 (1984).
80. M. Bertland and J.-L. Gras, *Tetrahedron*, 30, 793 (1974).

81. A. Kumar, A. Singh, and D. Devaprabhakara, Tetrahedron Lett., 2177 (1976).
82. J.E. McMurry and D.V. Miller, Tetrahedron Lett., 24, 1885 (1983).
83. H.J. Liu and W.H. Chan, Can. J. Chem., 60, 1081 (1982).
84. W. Eemical, D.R. Kearns, and P. Radlick, J. Am. Chem. Soc., 91, 3396, (1969); 91, 7771 (1969).
85. A.J. Mancuso, D.S. Brownfaith, and D. Swern, J. Org. Chem., 44, 4148 (1979).
86. E.D. Mihelich and D.J. Eickhoff, J. Org. Chem., 48, 4135 (1983).
87. H.J. Liu and M. Llinas-Brunet, Can. J. Chem., In press.
88. B.M. Trost, M. Poeckel, and L.M. Leichter, J. Am. Chem. Soc., 97, 2224 (1975).
89. G.A. Tolstikov, M.S. Miftakhov, and F.A. Akbutina, J. Org. Chem. U.S.S.R., Engl. Ed., 21, 611 (1985).
90. E.J. Corey and J.W. Suggs, Tetrahedron Lett., 2647 (1975).
91. D.H.R. Barton and S.W. McCombie, J. Chem. Soc., Chem. Commun., 1574 (1975).
92. G. Zweifel, H.C. Brown, Organic Reactions, 13, 28 (1963).