

1 **Comparative Assessment of qPCR Enumeration Methods that Discriminate Between Live**
2 **and Dead *Escherichia coli* O157:H7 on Beef**

3
4
5 Anna M. Laidlaw^{a,b}, Michael G. Gänzle^{b,c}, Xianqin Yang^a,

6
7 ^a Agriculture and Agri-Food Canada Lacombe Research and Development Centre, 6000 C & E
8 Trail, Lacombe, Alberta, Canada T4L 1W1

9 ^bDepartment of Agricultural, Food and Nutritional Science, University of Alberta, 4-10
10 Agriculture/Forestry Centre, Edmonton, AB, T6G 2P5, Canada

11 ^c Hubei University of Technology, College of Bioengineering and Food Science, Wuhan, P.R.
12 China.

13
14
15 Corresponding Author:

16 Xianqin Yang

17 Tel: 1 403 782 8119 Fax: 1 403 782 6120

18 Email: xianqin.yang@canada.ca

20 Abstract

21 Quantitative Polymerase Chain Reaction (qPCR) is a molecular method commonly used to
22 detect and quantify bacterial DNA on food but is limited by its inability to distinguish between
23 live and dead cell DNA. To overcome this obstacle, propidium monoazide (PMA) alone or with
24 deoxycholate (DC) was used to prevent dead cell detection in qPCR. qPCR methods were used to
25 detect strains of *Escherichia coli* O157, which can cause infection in humans with an infectious
26 dose of less than 10 cells. A 5 strain *E. coli* O157:H7 cocktail was inoculated onto beef steaks and
27 treated with interventions used in meat facilities (lactic acid (5%), peroxyacetic acid (200 ppm) or
28 hot water (80°C for 10 s)). Treatment of PMA or PMA + DC was applied to samples followed by
29 DNA extraction and quantification in qPCR. RNA was also quantified in addition to conventional
30 plating. For lactic acid intervention, qPCR DNA quantification of *E. coli* O157:H7 yielded $6.59 \pm$
31 0.21 and 6.30 ± 0.11 log gene copy #/cm² for control and lactic acid samples, respectively and after
32 treatment with PMA or PMA + DC this was further reduced to 6.31 ± 0.21 and 5.58 ± 0.38 ,
33 respectively. This trend was also observed for peroxyacetic acid and hot water interventions. In
34 comparison, RNA quantification yielded 7.65 ± 0.13 and 7.02 ± 0.38 log reverse transcript/cm²
35 for rRNA control and lactic acid samples, respectively, and for plating (LB), 7.51 ± 0.06 and 6.86
36 ± 0.32 log CFU/cm², respectively. Our research determined that treatment of PMA + DC in
37 conjunction with qPCR prevented dead cell DNA detection. However, it also killed cells injured
38 from intervention that may have otherwise recovered. RNA quantification was more laborious and
39 results had higher variability. Overall, quantification with conventional plating proved to be the
40 most robust and reliable method for live EHEC detection on beef.

41 Keywords: Propidium Monoazide, Deoxycholate, qPCR, Detection, Beef, *Escherichia coli*
42 O157:H7

43 **1. Introduction**

44 Molecular detection methods have become a mainstay in assessment of microbiological
45 composition and activity within microbial ecosystems, including food safety assessment, as they
46 are faster and more selective than labor-intensive and time-consuming conventional plating
47 methods (Ge and Meng 2009). Quantitative polymerase chain reaction (qPCR) is one of the most
48 commonly used methods of gene quantification, and is a validated method for pathogen detection
49 in food (Health Canada 2018; Wong and Medrano 2005). However, qPCR is limited in that it does
50 not discriminate between DNA from live and dead cells in a food sample, which may lead to false
51 positive results and lead to unnecessary recalls of food products (Bae and Wuertz 2009). Current
52 methods for pathogen detection use an enrichment step to prevent detection of dead cells (Ge and
53 Meng, 2009). Enrichment of samples also reduces the detection limit from > 1000 cells / g (qPCR
54 only) to less than 1 cell per 400 g (qPCR after enrichment); however, an enrichment increases the
55 time required for detection and does not allow quantitative results to be obtained.

56 Monoazide DNA binding dyes can enter dead cells with compromised membranes, intercalate
57 with DNA bases and form covalent bonds upon exposure to light (Takahashi et al. 2017; Taskin
58 et al. 2011). This modification of DNA prevents its amplification in qPCR. Ethidium monoazide
59 (EMA) has previously been investigated for this purpose, but was found to cause significant DNA
60 loss from viable cells after extraction (Flekna et al. 2007; Nocker et al. 2006a). Propidium
61 monoazide (PMA) penetrates into cells with compromised membranes without affecting viable
62 cells with intact membranes (Pan and Breidt 2007). However, PMA is excluded from dead cells
63 with intact membranes and therefore PMA-qPCR may overestimate the number of viable cells
64 (Løvdal et al. 2011; Nocker et al. 2007; Pacholewicz et al. 2013). To better select for live cells in
65 qPCR, a sodium deoxycholate treatment prior to PMA application can enhance its penetration due

66 to the emulsifier's ability to permeabilize intact membranes of dead cells (Yang et al. 2011). While
67 *E. coli* is highly tolerant to bile salts, the effect of deoxycholate on injured cells is unknown. An
68 alternative is quantification of RNA using qPCR as an indicator of cell viability as it degrades
69 more rapidly upon cell death than DNA, with mRNA degrading more rapidly than rRNA (Keer
70 and Birch 2003). However, the low stability of RNA also increases the risk of degradation of the
71 target molecule during sample preparation.

72 Ruminants including cattle are a major reservoir of enterohaemorrhagic *E. coli* (EHEC)
73 (Croxen et al., 2013); accordingly, foodborne illness caused by EHEC has often been linked to the
74 consumption of contaminated beef (Blagojevic et al. 2012; Rangel et al. 2005). Interventions
75 implemented in beef packing plants to control EHEC on carcasses include lactic acid and
76 peroxyacetic acid washes as well as hot water or steam application (Bacon et al. 2000). *E. coli*
77 O157, the prototype serogroup of EHEC, is regulated in Canada and the US and current methods
78 of detection of *E. coli* O157 in food include enrichment in conjunction with qPCR. However,
79 qPCR does not differentiate between live cells and cells killed the use of interventions, limiting its
80 use when both live and dead cells are present. This investigation aimed to compare four different
81 PCR-based detection methods to determine which method most accurately quantifies live cells of
82 *E. coli* O157, in comparison to viable plate counts as reference method for bacterial viability.
83 Reverse-transcription (RT-)qPCR of mRNA and rRNA as well as PMA or PMA and deoxycholate
84 treatment prior to DNA quantification in qPCR were evaluated. Quantification experiments were
85 performed on beef steaks inoculated with a cocktail of *E. coli* O157:H7 strains, and treated with
86 lactic acid, peroxyacetic acid or hot water as a decontamination treatments.

87 **2. Material and Methods**

88 **2.1. Preparation of Inoculum**

89 Five Shiga toxin (stx) negative *E. coli* O157:H7 (02-0628, 02-1840, 02-0304, 02-0627, 00-
90 3581; Luciano et al. 2011) were grown individually overnight in 9 mL of Luria-Bertani (LB;
91 Oxoid, Mississauga, Ontario, Canada) broth in 15 mL glass tubes at 37°C and 250 rpm. The next
92 day, cells of each strains were harvested by centrifugation, cell pellets were washed twice with 9
93 mL of 0.1% (w/v) peptone water (Difco, Becton-Dickinson, Sparks, Maryland, USA) and then
94 combined to create a 5 strain cocktail of *E. coli* O157:H7.

95 **2.2. Primers**

96 Quantification of DNA in qPCR was done with the species specific primers URL-301 (5'-TGT
97 TAC GTC CTG TAG AAA GCC C-3') and URR-432 (5'-AAA ACT GCC TGG CAC AGC AAT
98 T-3') that target *uidA* coding for β -glucuronidase activity of *E. coli* (Bej et al. 1991). For
99 quantification of rRNA in reverse transcription qPCR (RT-qPCR) primers targeting partial 16S
100 rRNA, HDAf (5'-ACT CCT ACG GGA GGC AGC AGT-3') and HDAr (5'-GTA TTA CCG CGG
101 CTG CTG GCA C-3') were used (Walter et al. 2000, Wang et al. 2013). For quantification of
102 mRNA in RT-qPCR, primers targeting glyceraldehyde-3-phosphate dehydrogenase were selected:
103 gapAf (5'-GTT GAC CTG ACC GTT CGT CT-3') and gapAr (5'-ACG TCA TCT TCG GTG
104 TAG CC-3'). While not all primers were species specific, all primers amplified the respective
105 target in *E. coli* and the beef samples were essentially free of other bacteria after decontamination
106 of the surface.

107 **2.3. Preparation of Beef Steaks for Intervention and Control Samples**

108 For each trial, fresh beef (eye of round) was purchased from a local grocery store. It was then
109 boiled in water for 2 min to decontaminate surface, the cooked outer layer (2 mm) was removed
110 with a sterile knife and then the clean beef cut was sliced into 2 cm thick steaks with surface area
111 of 88 cm². One mL of the 5 strain cocktail of *E. coli* O157:H7, corresponding to an inoculum of

112 10^7 to 10^8 cfu/cm², was inoculated onto the top surface of a beef steak in a biosafety cabinet which
113 was immediately spread evenly across the surface of the steak with a sterile plastic spreader. Steaks
114 with inoculum were left to air-dry in the biosafety cabinet for 30 min to allow for attachment.
115 Three trials were performed per intervention and for each trial, eight steaks were inoculated with
116 the 5 strain *E. coli* O157:H7 cocktail: four for application of interventions (2 x 0 min and 2 x 30
117 min sampling) and four for inoculated control (no application of intervention; 2 x 0 min and 2 x
118 30 min sampling). The sampling times were selected to allow for a comparison of the effect of
119 different interventions on cell viability, injury and recovery after immediate application and after
120 a 30 min holding time. One steak was left uninoculated as a control sample per trial to confirm
121 that the steak did not contain detectable bacteria or bacterial nucleic acids, where no intervention
122 was applied. The control steak was immediately placed in 99 mL of 100 mM PBS buffer (includes
123 buffering agents monopotassium phosphate and disodium phosphate) in a stomacher bag,
124 manually massaged for 2 min and suspension plated on LB and Violet Red Bile (VRB, Oxoid)
125 agars. Plates were incubated at 30°C for 24 h but all counts were below the detection limit of 100
126 CFU/g, corresponding to fewer than 2 colonies per plate. Samples of one mL were kept for DNA
127 qPCR analysis; similar to plating, quantification yielded negative result denoted by either
128 amplification at or above 35 Ct with species specific primers (*uidA*) for DNA qPCR or 10 Ct above
129 last sample's Ct value for non-specific primers (*HDA*, *gapA*) used in both RNA RT-qPCR
130 experiments.

131 **2.4. Lactic Acid or Peroxyacetic Acid Intervention**

132 Prior to application of interventions, inoculated and control steaks were placed in separate
133 aluminum trays (23 x 15 x 1.5 cm). A 5% (w/v) lactic acid (Sigma-Aldrich, Mississauga, Ontario,
134 Canada) or 200 mg/L peroxyacetic acid (Inspexx, Ecolab Inc., St. Paul, Minnesota, USA) solution

135 was prepared and connected to the equipment used for application of antimicrobials; this
136 equipment has been previously described in detail (Youssef et al. 2012). The aluminum trays
137 containing steaks were placed on the stainless steel conveyor belt that carried trays under a spray
138 nozzle, exposing steaks to a 5% lactic acid spray or 200 mg/L peroxyacetic acid spray at a volume
139 of 0.4 mL of fluid per cm² steak surface. The steaks were sampled immediately (0 min) or after 30
140 min of holding at room temperature.

141 **2.5. Hot Water Intervention**

142 To simulate spraying with hot water, which is used in beef processing as a pathogen
143 intervention method, a 4 L beaker was filled with 2.5 L of sterile water and brought to 80°C on a
144 hot plate. Once the target temperature was reached, steaks were submerged individually for 10 s
145 using sterile stainless steel tongs and immediately removed and sampled at 0 or 30 min.

146 **2.6. Sampling of Steaks after Intervention**

147 Each steak was placed in a sterile stomacher bag (VWR International, Radnor, Pennsylvania,
148 USA) with 100 mL of buffer. PBS was used to neutralize lactic acid or hot water intervention
149 samples and neutralizing buffer (includes buffering agent monopotassium phosphate and reducing
150 agent sodium thiosulfate) was used for peroxyacetic acid intervention-treated samples to neutralize
151 acidulant as well as quench oxidant. Steaks in buffer were then massaged manually in stomacher
152 bags for 2 min. From sample suspension, 15 x 1 mL aliquots were transferred to separate 2 mL
153 transparent Eppendorf tubes. Tubes were centrifuged at 10,000 *x g* for 5 min, supernatants
154 discarded, cells washed in 1 mL peptone water, centrifuged and resuspended in 300 µL peptone
155 water. Each tube was designated to one of 5 different sampling groups for subsequent treatment
156 application, in triplicate: DNA extraction (control; no PMA or deoxycholate), PMA treatment and
157 DNA extraction, PMA with deoxycholate treatment and DNA extraction, RNA extraction, plating

158 (Figure 1). Samples for RNA extraction were further washed with RNAprotect Bacteria Reagent
159 (Qiagen, Mississauga, Ontario, Canada) according to the manufacturer's instructions to stabilize
160 and protect RNA from degradation, the resulting pellet was kept frozen (-20°C) until use.

161 **2.7. PMA and Deoxycholate Treatment**

162 For PMA treatment of samples, a 20 mM stock solution of PMA (Biotium, Inc. Hayward,
163 California, USA) was stored in dark at 4°C until use. For each sample suspension (300 µL) in 2
164 mL Eppendorf tubes, 1.5 µL of 20 mM PMA was added and vortexed thoroughly. Tubes were
165 placed on ice horizontally in the dark for 5 min followed by exposure to a 650-W halogen lamp
166 for 2 min, held 20 cm above the tubes, while slowly rocking samples back and forth to ensure
167 uniform exposure. Samples were then removed from light, centrifuged at 8,000 x g for 3 min, the
168 supernatant was discarded, and the cells were washed with peptone water. The cell pellet was
169 stored at -20°C until further analysis.

170 For samples treated with both PMA and deoxycholate, deoxycholate was applied to samples
171 before PMA treatment using a protocol previously optimized by Yang et al. (2011). A 1% (w/v)
172 solution of deoxycholate (Sigma-Aldrich) was stored at -20°C until use. Sample suspensions (300
173 µL) in 2 mL Eppendorf tubes were centrifuged at 8,000 x g for 3 min, and the supernatants were
174 discarded. The pellets were resuspended with 300 µL 1% deoxycholate and incubated in a rocking
175 water bath at 37°C for 30 min. After incubation, the suspension was treated with PMA as described
176 above.

177 **2.8. Enumeration of *E. coli* O157:H7 on Beef Steaks**

178 Viable cell counts were enumerated by surface plating on the day of treatment. Ten-fold
179 dilutions were prepared in sterile 0.1% peptone water from original sample suspensions of beef

180 steaks in stomacher bags with buffer, and 100 μ L of dilutions estimated to provide 30-300 colonies
181 per plate was used to manually plate onto LB and VRB agar. Plates were incubated at 30°C for 24
182 h and enumerated next day. Uninjured cells (capable of reproduction without repair of cellular
183 injury) were enumerated on VRB agar, a medium which prevents the growth of injured cells with
184 compromised outer membrane due to the presence of bile salts (Gänzle and Vogel, 2001). All
185 viable (live) cells (capable of reproduction, including cells that require repair of cellular injury
186 prior to reproduction) were enumerated on LB agar, a non-selective medium (Scheusner et al.
187 1971).

188 **2.9. DNA Extraction and Quantification in qPCR**

189 DNA was extracted from samples (DNeasy blood and tissue kit; Qiagen) following instructions
190 for Gram-negative bacteria with modifications. Cells suspended in buffer ATL (supplemented with
191 proteinase K) were incubated for 2 h, and DNA was eluted from mini columns twice with 100 μ L
192 of buffer AE. For quantification in qPCR, primers URL-301 and URR-432 were used to target
193 *uidA* gene (Sigma-Aldrich). qPCR master mix was prepared as follows per reaction: 12.5 μ L of
194 SYBR green (Qiagen), 2.5 μ L each forward and reverse primer, and 2.5 μ L nuclease free water,
195 with 5 μ L template DNA added to 20 μ L of master mix for a total volume of 25 μ L per reaction.
196 A standard curve was prepared from purified PCR product of *E. coli* O157:H7 (00-3581) DNA
197 (QIAquick PCR Purification Kit; Qiagen), which was diluted to 100 mg/L and then further diluted
198 in serial ten-fold dilutions. Six dilutions were used to generate a standard curve in the range of
199 DNA concentrations in the samples. Nuclease free water served as negative control for all PCR
200 reactions; uninoculated steaks served as negative control for any native *E. coli* that may be already
201 present on steak. A 7500 Fast Real-Time PCR system (Applied Biosystems) was used for
202 quantification of DNA by one initial denaturation of template DNA and activation of Taq DNA

203 polymerase at 95°C for 10 min, followed by 40 cycles of denaturation at 95°C for 30 s, primer
204 annealing at 60°C for 1 min, extension at 72°C for 1 min. Quantification data was acquired from
205 7500 Software (version 2.0.5, Applied Biosystems) and analyzed. Melting curves of final products
206 verified no formation of non-specific product or primer dimers. Each sample was quantified in
207 duplicate per qPCR plate.

208 **2.10. RNA Extraction and Quantification in RT-qPCR**

209 RNA was extracted from samples (RNeasy Mini Kit; Qiagen) and reverse transcribed to cDNA
210 following instructions stated in the Kit's manual (QuantiTect Reverse Transcription Kit; Qiagen).
211 Two sets of primers were selected for RNA analysis, one targeting partial 16S rRNA (HDAf and
212 HDAr; Sigma-Aldrich) and the second targeting the glyceraldehyde-3-phosphate dehydrogenase
213 (GAPDH) (gapAf and gapAr; Sigma-Aldrich). Stocks were prepared for each primer set with
214 sterile nuclease free water. Master mix recipe per reaction was prepared as follows: 12.5 µL SYBR
215 green (Qiagen), 0.5 µL each forward and reverse primers, 10.5 µL nuclease free water, and 1 µL
216 sample cDNA. A standard curve was developed by extracting genomic DNA from an overnight
217 culture of *E. coli* O157:H7 (00-3581) using Wizard Genomic DNA purification kit (Promega).
218 Extracted genomic DNA was run in qPCR with respective primers (HDA or *gapA*) and the PCR
219 product was purified according to purification kit instructions (QIAquick PCR Purification Kit,
220 Qiagen) and diluted to 100 mg/L. For each primer set, this concentration was further diluted ten-
221 fold (up to 10⁻⁸) to generate a standard curve. Negative controls included nuclease free water,
222 samples after treatment with genomic DNA Wipeout but before reverse transcription, and isolation
223 of RNA from uninoculated steaks. The Reverse Transcription qPCR (RT-qPCR) method was one
224 initial denaturation of template DNA and activation of Taq DNA polymerase at 95°C for 5 min,
225 followed by 40 cycles of denaturation at 95°C for 30 s, primer annealing at 60°C for 30 s, extension

226 at 72°C for 30 s. Data was acquired from 7500 software (version 2.0.5, Applied Biosystems) and
227 analyzed. Melting curves of final products were used to verify that non-specific products or primer
228 dimers did not form. Samples were run in triplicate reaction per plate.

229 **2.11. Statistical Analysis**

230 Statistical Analysis Systems (SAS Institute, Inc., Cary, NC) was used to evaluate data by two
231 way analysis of variance (ANOVA) followed by pair-wise comparison of means from different
232 treatment groups with Bonferroni adjustment. A probability (p) of <0.05 was considered
233 statistically significant. Three independent trials were performed for each intervention, with
234 triplicate samples per trial. Linear regressions were calculated in SigmaPlot 12.5.

235 **3. Results**

236 **3.1. Viable Plate Counts and Sublethal Injury on Treated and Untreated Beef.**

237 Cell counts for untreated, inoculated beef were on average 7.27 ± 0.15 log CFU/cm² (mean \pm
238 standard deviation); the difference of plate counts between LB and VRB, which represents the sub-
239 lethally injured population, on average was 0.33 ± 0.18 log CFU/cm². After intervention with lactic
240 acid and hot water, the difference in plate counts on LB and VRB increased, indicating that the
241 number of injured cells had increased. However, the average proportion of injured cells was not
242 increased by peroxyacetic acid treatment compared to the control (Table 1). Comparison of post-
243 intervention counts to the controls indicated that the percentage of dead cells in samples was 68%,
244 45% and 90% for meat treated by lactic acid, peroxyacetic acid and hot water, respectively.

245 **3.2. Quantification of *E. coli* O157:H7 on Beef by qPCR After Lactic Acid Intervention**

246 To determine the method that most accurately quantifies viable *E. coli* O157:H7 on beef
247 steak after lactic acid intervention, five different detection methods were compared. Figures 2A
248 and B correlate viable plate counts of *E. coli* O157:H7 after lactic acid intervention to the gene

249 copy numbers obtained by quantification of microbial DNA or RNA. Control samples represent
250 steaks that were inoculated but not treated; control quantification represents DNA quantification
251 without PMA or PMA-DC. Lactic acid intervention reduced viable cell counts by about 1 log
252 CFU/cm² (Table 1), however, this reduction was not significant. Interestingly, significant
253 reduction in the log gene copy #/cm² of *E. coli* O157:H7 was determined by qPCR only in samples
254 treated with PMA combined with deoxycholate. However, reductions observed at 0 and 30 min
255 indicate that the combination treatment not only prevented amplification of DNA from cells killed
256 by intervention, but it also prevented the amplification of DNA from sub-lethally injured cells
257 when compared to plate count data. This explains why reductions were significant as compared
258 to plating. Accordingly, the slope of the regression line correlating cell counts and gene copy
259 numbers was < 1.0 for qPCR and PMA-qPCR quantification but > 1 for PMA-deoxycholate qPCR.
260 Quantification with RT-qPCR targeting rRNA or mRNA provided comparable results (Figure 2B).

261 **3.3. Quantification of *E. coli* O157:H7 on Beef by qPCR After Peroxyacetic Acid**

262 **Intervention**

263 Similarly, after peroxyacetic acid intervention of steaks inoculated with *E. coli* O157:H7, five
264 different detection methods were compared to determine the most accurate method to quantify
265 viable cells. Figures 3 A and B correlate quantification of *E. coli* O157:H7 on beef after
266 peroxyacetic acid intervention. Enumeration on LB agar was compared to DNA or RNA
267 quantification in qPCR or RT-qPCR, respectively. After peroxyacetic acid intervention, plate
268 counts and the number of gene copies were not reduced at 0 or 30 min sampling times. The
269 reduction of cell counts or of the copy number of nucleic acids after combination treatment of
270 PMA with deoxycholate was also not significant. This is likely because this intervention did not
271 cause the same degree of sub-lethal injury as the other two interventions.

272 **3.4. Quantification of *E. coli* O157:H7 on Beef by qPCR After Hot Water Intervention**

273 The same five different detection methods were compared to determine the most accurate
274 method to quantify live cells after hot water intervention on beef steaks inoculated with *E. coli*
275 O157:H7. The correlation of quantification of *E. coli* O157:H7 on beef after hot water intervention
276 by means of enumeration on LB agar as compared to DNA or RNA quantification in qPCR or RT-
277 qPCR is presented in Figures 4 A and B, respectively. Enumeration by plating, qPCR or RT-qPCR
278 determined that the number of viable *E. coli* O157:H7 on steaks at both 0 and 30 min were not
279 significantly reduced by hot water intervention. However, PMA with deoxycholate treatment and
280 qPCR quantification reduced the log gene copy #/cm² at 0 and 30 min. The significant reduction
281 is likely attributed the degree of sub-lethal injury that the intervention caused, and that this
282 population of cells was sensitive to the combination of PMA with deoxycholate treatment, which
283 then led to the prevention of amplification in qPCR. Accordingly, the slope of the regression line
284 correlating plate counts to gene copy numbers was 1.26 for quantification with PMA-deoxycholate
285 qPCR but only about 0.5 for quantification with qPCR and PMA-qPCR (Fig. 4A). Quantification
286 with RT-qPCR targeting rRNA or mRNA provided comparable results with a slope of 1.1 and 1.3,
287 respectively (Fig. 4B).

288 **3.5. *E. coli* O157:H7 Recovered on LB and VRB from Beef Steaks at 30 min After**

289 **Intervention and After Treatment of Cells with PMA or PMA with Deoxycholate**

290 qPCR quantification indicated that the combination of PMA and deoxycholate treatment
291 permeabilized viable *E. coli* cells to PMA after intervention when compared to control and cell
292 count enumeration data by plating. To determine whether treatment with PMA and PMA-DC
293 affects viability of *E. coli*, the organism was enumerated after intervention treatments on beef
294 steaks and PMA or PMA-deoxycholate treatment (Table 1). PMA did not affect the viability of *E.*

295 *coli* (Table 1). Treatment with both PMA and deoxycholate reduced the viability of *E. coli*; this
296 reduction was significant after treatment of *E. coli* that were previously exposed to hot water
297 washes on beef. Taken together, these data suggest that PMA-deoxycholate may permeabilize and
298 kill *E. coli* when cells were sublethally injured by intervention treatments.

299 **4. Discussion**

300 This study compared five different detection methods for the quantification of viable *E. coli*
301 O157 on beef, and after intervention treatments that are currently used in the meat industry. The
302 application of lactic acid, peroxyacetic acid and hot water interventions to beef steaks created a
303 heterogeneous population of viable, sub-lethally injured, and dead cells. These populations are
304 highly suitable for testing detection methods for the ability to discriminate between live and dead
305 bacterial cells. Lactic acid and hot water interventions were more lethal to *E. coli* O157:H7 on beef
306 than peroxyacetic acid, which is consistent with previous studies (Castillo et al. 1998; Ellebracht
307 et al. 2005; Gill and Bryant 2000; King et al. 2005; Youssef et al. 2012). Experiments reported in
308 this study were obtained with aseptic meat samples without background microbiota and primers
309 that were not specific to *E. coli* O157. Equivalent results can be expected if primer sets with more
310 specific targets are used to specifically detect *E. coli* O157 or STEC in commercial meat samples.

311 Overall, the cell counts obtained by plating on LB agar were similar to RT-qPCR based
312 quantification that targeted rRNA; mRNA copy numbers and DNA copy numbers were generally
313 lower. This discrepancy reflects that mRNA is less stable than rRNA and that *E. coli* cells contain
314 several thousand ribosomes (Keer and Birch 2003). The decrease of viable cell counts after
315 intervention treatments was reflected by a reduced copy number obtained by RT-qPCR. However,
316 qPCR targeting DNA did not discriminate between viable and dead cells, though plating does
317 (Sidari et al. 2011). Furthermore, rRNA provided a more robust estimation than mRNA, which is

318 present as a lower percentage of total RNA and is more susceptible to degradation as compared to
319 rRNA (Kushner 2002).

320 Enumeration of viable cells by surface plating indicated that lactic acid was more lethal than
321 peroxyacetic acid; this difference was not observed by RT-qPCR, demonstrating that conventional
322 plating provided the most accurate estimate of viable cells. Hot water intervention resulted in a
323 greater reduction in viable cells from the plate count quantification method and both RNA RT-
324 qPCR quantification methods. Hot water washes acted immediately while lactic acid acted more
325 slowly. The antimicrobial effect of lactic acid is based on penetration of cells and lowering
326 intracellular pH (Alakomi et al. 2000) while heat leads to immediate disruption of membranes and
327 irreversible denaturation of proteins (Lee and Kaletunc 2002).

328 PMA treatment does not affect viability of cells (Elizaquivel et al. 2012; Nocker et al. 2006b,
329 Yang et al. 2011; this study). PMA-qPCR, reduced the gene copy numbers detected in beef samples
330 but the difference between qPCR and PMA-qPCR was small and remained unchanged by
331 bactericidal intervention treatments (Figs. 2, 3, and 4). A high number of dead cells may inhibit
332 PMA crosslinking of DNA by light and thus reduce the correlation to viable plate counts (Løvdal
333 et al. 2011; Pan and Breidt 2007; Wagner et al. 2008; Yañez et al. 2011). To counter this, it has
334 been suggested that incubating samples for a longer period of time and at higher temperatures to
335 increase PMA penetration into dead cells (Nkuipou-Kenfack et al. 2013).

336 Several past studies compare PMA-qPCR to quantification of live cells of foodborne pathogens
337 by plate counts. Løvdal et al. (2011) applied lethal heat treatment of 80°C for 15 min to *Listeria*
338 *innocua* cells in broth and found large discrepancies between quantification methods: plating
339 determined cells counts below the detection limit whereas PMA-qPCR quantified 8 log cells/mL

340 Enumeration of *Campylobacter jejuni* on raw chicken carcasses after different slaughterhouse
341 processing steps determined that PMA-qPCR consistently yielded up to a 1-1.5 log CFU/mL
342 higher cell count than plating (Pacholewicz et al. 2013). After severe heat treatment of *C. jejuni* in
343 chicken rinse at 95°C for 5 min, PMA-qPCR and plating both indicated that cells counts were
344 below the detection limit (Josefsen et al. 2010). Lastly, Ju et al. (2016) quantified live *E. coli*
345 O157:H7 on lettuce. Plating indicated that *E. coli* O157:H7 did not survive on leaf surfaces and
346 was reduced by 10⁷ cells within 96 h; however, RT-qPCR and PMA-qPCR indicated a reduction
347 of 10² and 10³ cells, respectively. Overall, a majority of studies indicate that PMA-qPCR can lead
348 to false positive results. This conforms to the observation that the structurally related propidium
349 iodide is excluded by a substantial proportion of dead cells after lethal interventions (Ulmer et al.,
350 2000). Some studies defined PI or PMA excluding but non-culturable cells “viable but not
351 culturable”; however, sublethally injured cells of *E. coli* are readily recovered by cultivation media
352 containing pyruvate or catalase as scavengers for oxidative stress (Afari and Hunc, 2018; Mizunoe
353 et al., 1999). We used LB as reference medium for detection of viable cells and thus did not account
354 for those cells that require extended recovery times or addition of pyruvate or catalase to recover
355 from injury.

356 The use of deoxycholate prior to PMA treatment was proposed to increase cell membrane
357 permeability to permeabilize all dead cells (Yang et al., 2014). In suspensions where *E. coli* cells
358 were killed after heat treatment with no viable cells remaining, treatment with PMA prior to qPCR
359 significantly decreases amplification of dead cells in qPCR as compared to PMA alone (Yang et
360 al. 2014). *E. coli* cells that were killed by 72°C or less were permeable to PMA but the addition of
361 1% deoxycholate treatment rendered a large portion of these cells permeable to the dye (Yang et
362 al. 2011). After lactic acid intervention, membranes of dead and injured *E. coli* cells that were

363 impermeable to PMA were rendered permeable to the dye after treatment with 1% deoxycholate
364 treatment (Wang et al. 2014). Injured cells were resuscitated in nutrient rich broth for 2 h prior to
365 deoxycholate treatment which restored injured cell membrane barrier properties. This could be a
366 possible solution to injured cell sensitivity to PMA with deoxycholate treatment. However, since
367 our study was performed in a more complex system rather than in broth or buffer, this could prove
368 to be more complicated and warrants further investigation. Zhou et al. (2016) optimized the
369 concentration of deoxycholate for use with *Cronobacter sakazakii* in powdered infant formula.
370 The authors found that heat injured cells were not affected by a concentration of 0.08%
371 deoxycholate and determined that at this concentration, the accuracy of viable cell detection when
372 combined with PMA and qPCR was increased. Sodium dodecyl sulfate has also been used to aid
373 in PMA penetration into dead cells and shows little impact on cell viability. While this could offer
374 an alternative emulsifier to use with PMA to increase the accuracy of live cell detection in qPCR,
375 it still requires further research in different food systems (Dong et al. 2018, Takahashi et al. 2017).
376 Our work indicates that combination treatment of PMA with deoxycholate prevents amplification
377 of DNA from sublethally injured cells, and could lead to false negative results.

378 The inability to discriminate between DNA in live and dead cells in qPCR is a problem that
379 extends beyond *E. coli* O157:H7 in meat, this combination treatment could be a solution in other
380 areas of research. This includes determination of viable probiotics in the gut or viable lactic acid
381 bacteria or bifidobacteria in fermented foods (Cocolin et al. 2011; Garcia-Cayuela et al. 2009;
382 Rantsiou and Cocolin 2006; Villarreal et al. 2013). However, it is important to note that
383 concentration of treatment should be adjusted to target microorganism, as Gram-positive
384 microorganisms have different resistance to deoxycholate than Gram-negative microorganisms
385 (Nkuipou-Kenfack et al. 2013).

386 **5. Conclusions**

387 This study compared five different methods to determine the most accurate quantification of
388 viable *E. coli* O157:H7 on beef steaks after intervention. PMA treatment with qPCR did not
389 prevent DNA amplification from all dead cells in samples after intervention and therefore this
390 treatment does not resolve the issue of false positive result in qPCR. However, it does prevent
391 amplification of some dead cell DNA and therefore increases the accuracy of viable cell
392 quantification compared to qPCR alone. In contrast, treatment of cells with PMA with
393 deoxycholate prevented amplification of DNA from all dead cells but does so at the cost of sub-
394 lethally injured cells since deoxycholate supported PMA penetration into injured cells which may
395 have otherwise recovered. This makes DNA inaccessible for qPCR quantification and would lead
396 to false negative result. Both rRNA and mRNA quantification in qPCR had more variability and
397 was not as sensitive for quantifying viable cells as plate counts. While PMA with deoxycholate
398 may not be suitable for pathogen detection in food where sublethal injury of cells is of concern, it
399 could be used in other areas of research that aim to exclude all dead cells at the expense of injured
400 cells. On the other hand, if research aims to amplify all viable cells including the injured cell
401 population at the cost of also amplifying some dead cell DNA, PMA or rRNA with qPCR detection
402 methods could be used. Overall, conventional plating provided the most sensitive and reliable
403 quantification of viable *E. coli* O157:H7 on beef.

404 **Acknowledgements**

405 The authors would like to thank Agriculture and Agri-Food Canada A-Base Funding for
406 providing project support. Funding source had no involvement in study design, collection, analysis
407 and interpretation of data, in writing of the report or in submission of the article.

408 **References**

- 409 Afari, G.K., Hunc, Y.C. 2018. Detection and verification of the viable but nonculturable
410 (VBNC) state of *Escherichia coli* O157:H7 and *Listeria monocytogenes* using flow
411 cytometry and standard plating. *Journal of Food Science*, *in press* doi: 10.1111/1750-
412 3841.14203
- 413 Alakomi, H., Skyttä, E., Saarela, M., Helander, I.M., 2005. Lactic acid permeabilizes gram-
414 negative bacteria by disrupting the outer membrane. *Applied and Environmental*
415 *Microbiology* 66, 2000–2005.
- 416 Bae, S. Wuertz, S., 2009. Discrimination of viable and dead fecal *Bacteroidales* bacteria by
417 quantitative PCR with propidium monoazide. *Applied and Environmental Microbiology* 75,
418 2940–2944.
- 419 Bacon, R., Belk, K., Sofos, J., Clayton, R., Reagan, J., Smith, G., 2000. Microbial populations on
420 animal hides and beef carcasses at different stages of slaughtering in plants employing
421 multiple-sequential interventions for decontamination. *Journal of Food Protection* 63, 1080-
422 1086.
- 423 Bej, A., DiCesare J, Haff, L., Atlas, R., 1991. Detection of *Escherichia coli* and *Shigella* spp in
424 water by using the polymerase chain-reaction and gene probes for *uid*. *Applied and*
425 *Environmental Microbiology* 57, 1013–1017.
- 426 Blagojevic, B., Antic, D., Ducic, M., Buncic, S., 2012. Visual cleanliness scores of cattle at
427 slaughter and microbial loads on the hides and the carcasses. *Veterinary Record* 170, 563–
428 563.
- 429 Castillo, A., Lucia, L., Goodson, K., Savell, J., Acuff, G., 1998. Use of hot water for beef carcass
430 decontamination. *Journal of Food Protection* 61, 19-25.
- 431 Cocolin, L., Dolci, P., Rantsiou, K., 2011. Biodiversity and dynamics of meat fermentations: The
432 contribution of molecular methods for a better comprehension of a complex ecosystem.
433 *Meat Science* 89, 296–302.
- 434 Croxen, M. A., Law, R. J., Scholz, R., Keeney, K.M., Wlodarska, M., Finlay, B.B. (2013).
435 Recent advances in understanding enteric pathogenic *Escherichia coli*. *Clinical*
436 *Microbiology Reviews* 26, 822–880.
- 437 Dong, L., Liu, H., Meng, L., Xing, M., Wang, J., Wang, C., Chen, H., Zheng, N. (2018).
438 Quantitative polymerase chain reaction coupled with sodium dodecyl sulfate and propidium
439 monoazide for detection of viable *Staphylococcus aureus* in milk. *Journal of Dairy Science*
440 101, 1-8.

441 Elizaquível, P., Sánchez, G., Aznar, R., 2012. Quantitative detection of viable foodborne *E. coli*
442 O157 : H7 , *Listeria monocytogenes* and *Salmonella* in fresh-cut vegetables combining
443 propidium monoazide and real-time PCR. *Food Control* 25, 704–708.

444 Ellebracht, J.W., King, D.A., Castillo, A., Lucia, L.M., Acuff, G.R., Harris, K.B., Savell, J.W.,
445 2005. Evaluation of peroxyacetic acid as a potential pre-grinding treatment for control of
446 *Escherichia coli* O157 : H7 and *Salmonella* Typhimurium on beef trimmings. *Meat Science*
447 70, 197–203.

448 Flekna, G., Polonca, S., Smulders, F.J.M., Mo, S.S., Hein, I., 2007. Insufficient differentiation of
449 live and dead *Campylobacter jejuni* and *Listeria monocytogenes* cells by ethidium
450 monoazide (EMA) compromises EMA / real-time PCR. *Research in Microbiology* 158,
451 405–412.

452 Gänzle, M.G., Vogel, R.F. 2001. On-line fluorescence determination of pressure mediated outer
453 membrane damage in *Escherichia coli*. *Systematic and Applied Microbiology* 24, 477-
454 485. Garcia-Cayuela, T., Tabasco, R., Pela, C., Requena, T., 2009. Simultaneous detection
455 and enumeration of viable lactic acid bacteria and bifidobacteria in fermented milk by using
456 propidium monoazide and real-time PCR. *International Dairy Journal* 19, 405–409.

457 Ge, B. Meng, J., 2009. Advanced technologies for pathogen and toxin detection in foods: current
458 applications and future directions. *Journal of the Association for Laboratory Automation* 14,
459 235–241.

460 Gill, C., Bryant, J., 2000. The effects on product of a hot water pasteurizing treatment applied
461 routinely in a commercial beef carcass dressing process. *Food Microbiology* 17, 495–504.

462 Health Canada, 2018. Laboratory procedures for the microbiological analysis of foods. The
463 Compendium of Analytical Methods. Volume 3. Accessed on May 4, 2018. [http://www.hc-
464 sc.gc.ca/fn-an/res-rech/analy-meth/microbio/volume3-eng.php](http://www.hc-sc.gc.ca/fn-an/res-rech/analy-meth/microbio/volume3-eng.php)

465 Josefsen, M.H., Lo, C., Hansen, T.B., Christensen, L.S., Olsen, J.E., Hoorfar, J., 2010. Rapid
466 quantification of viable *Campylobacter* bacteria on chicken carcasses , using real-time PCR
467 and propidium monoazide treatment , as a tool for quantitative risk assessment. *Applied and
468 Environmental Microbiology* 76, 5097–5104.

469 Keer, J.T., Birch, L., 2003. Molecular methods for the assessment of bacterial viability. *Journal
470 of Microbiological Methods* 53, 175–183.

471 King, D.A., Lucia, L.M., Castillo, A., Acuff, G.R., Harris, K.B., Savell, J.W., 2005. Evaluation
472 of peroxyacetic acid as a post-chilling intervention for control of *Escherichia coli* O157 :
473 H7 and *Salmonella* Typhimurium on beef carcass surfaces. *Meat Science* 69, 401–407.

474 Kushner, S., 2002. mRNA decay in *Escherichia coli* comes of age. *Journal of bacteriology* 184,
475 4658–4665.

- 476 Lee, J., Kaletunc, G., 2002. evaluation of the heat inactivation of *Escherichia coli* and
 477 *Lactobacillus plantarum* by differential scanning calorimetry. Applied and Environmental
 478 Microbiology 68, 5379–5386.
- 479 Løvdal, T., Befring, M., Björkblom, B., Møller, S.G., 2011. Propidium monoazide combined
 480 with real-time quantitative PCR underestimates heat-killed *Listeria innocua*. Journal of
 481 Microbiological Methods 85, 164–169.
- 482 Mizunoe, Y., Wai, S.N., Takade, A., Yoshida, S.I. Restoration of culturability of starvation-
 483 stressed and low-temperature-stressed *Escherichia coli* O157 cells by using H₂O₂-degrading
 484 compounds. Archives of Microbiology 172 , 63–67.
- 485 Nocker, A., Camper, A.K., 2006a. Selective removal of dna from dead cells of mixed bacterial
 486 communities by use of ethidium monoazide. Applied and Environmental Microbiology 72,
 487 1997–2004.
- 488 Nocker, A., Cheung, C., Camper, A.K., 2006b. Comparison of propidium monoazide with
 489 ethidium monoazide for differentiation of live vs . dead bacteria by selective removal of
 490 DNA from dead cells. Journal of Microbiological Methods 67, 310–320.
- 491 Nocker, A., Sossa, K.E., Camper, A.K., 2007. Molecular monitoring of disinfection efficacy
 492 using propidium monoazide in combination with quantitative PCR. Journal of
 493 Microbiological Methods 70, 252–260.
- 494 Nkuipou-kenfack, E., Engel, H., Fakih, S., Nocker, A., 2013. Improving efficiency of viability-
 495 PCR for selective detection of live cells. Journal of Microbiological Methods 93, 20–24.
- 496 Pacholewicz, E., Swart, A., Lipman, L.J., Wagenaar, J., Havelaar, A.H., Duim, B., 2013.
 497 Propidium monoazide does not fully inhibit the detection of dead *Campylobacter* on broiler
 498 chicken carcasses by qPCR. Journal of Microbiological Methods 95, 32–38.
- 499 Pan, Y., Breidt, F., 2007. Enumeration of viable *Listeria monocytogenes* cells by real-time PCR
 500 with propidium monoazide and ethidium monoazide in the presence of dead cells. Applied
 501 and Environmental Microbiology 73, 8028–8031.
- 502 Rangel, J.M., Sparling, P.H., Crowe, C., Griffin, P.M., Swerdlow, D.L., 2005. 1982 – 2002.
 503 epidemiology of *Escherichia coli* O157:H7 outbreaks, United States, 1982-2002. Emerging
 504 Infectious Diseases 11, 603-609.
- 505 Rantsiou, K., Cocolin, L., 2006. New developments in the study of the microbiota of naturally
 506 fermented sausages as determined by molecular methods: a review. International Journal of
 507 Food Microbiology 108, 255–267.
- 508 Scheusner, S., Busta, F., Speck, M., 1971. Inhibition of injured *Escherichia coli* by several
 509 selective agents. Applied Microbiology 21, 46-49.


- 510 Sidari, R., Caridi, A., 2011. Methods for detecting Enterohaemorrhagic *Escherichia coli* in food.
511 Food Reviews International 27, 134–153.
- 512 Takahashi, H., Gao, Y., Miya, S., Kuda, T., Kimura, B. Discrimination of live and dead cells of
513 *Escherichia coli* using propidium monoazide after sodium dodecyl sulfate treatment. Food
514 Control 71, 79-82.
- 515 Taskin, B., Gozen, A.G., Duran, M., 2011. Selective quantification of viable *Escherichia coli*
516 bacteria in biosolids by quantitative PCR with propidium monoazide modification. Applied
517 and Environmental Microbiology 77, 4329–4335.
- 518 Ulmer, H.M., Gänzle, M.G., Vogel, R.F. 2000. Effects of high pressure on survival and
519 metabolic activity of *Lactobacillus plantarum*. Applied and Environmental Microbiology
520 66, 3966-3973.
- 521 Villarreal, M., Padilha, M., Diogo, A., Vieira, S., Franco, B., Martinez, R., Saad, S., 2013.
522 advantageous direct quantification of viable closely related probiotics in petit-suisse cheeses
523 under in vitro gastrointestinal conditions by propidium monoazide - qPCR. PLOS one 8, 1–
524 11.
- 525 Wagner, A.O., Malin, C., Knapp, B.A., Illmer, P., 2008. removal of free extracellular dna from
526 environmental samples by ethidium monoazide and propidium monoazide. Applied and
527 Environmental Microbiology 74, 2537–2539.
- 528 Walter, J., Tannock, G., Tisala-Timisiaarvi, A., Rodtong, S., Loach, D., Munroe, K., Alatosava,
529 T., 2000. Detection and identification of gastrointestinal *Lactobacillus* species by using
530 denaturing gradient gel electrophoresis and species-specific PCR primers. Applied and
531 Environmental Microbiology 66, 297-303.
- 532 Wang, H., Gill, C., Yang, X., 2014. Development of a real-time PCR procedure for
533 quantification of viable *Escherichia coli* in populations of *E. coli* exposed to lactic acid, and
534 the acid tolerance of verotoxigenic *E. coli* (VTEC) from cattle hides. Food Control 43, 104-
535 109.
- 536 Wang Y, Ametaj B, Ambrose D, Gänzle M, 2013. Characterisation of the bacterial microbiota of
537 the vagina of dairy cows and isolation of pediocin-producing *Pediococcus acidilactici*.
538 BMC Microbiol. 13:19.
- 539 Wong, M.L., Medrano, J.F., 2005. Real-time PCR for mRNA quantitation. BioTechniques 1 39,
540 1–11.
- 541 Yáñez, M.A., Nocker, A., Soria-soria, E., Múrtula, R., Martínez, L., Catalán, V., 2011.
542 Quantification of viable *Legionella pneumophila* cells using propidium monoazide
543 combined with quantitative PCR. Journal of Microbiological Methods 85, 124–130.
- 544 Yang, X., Badoni, M., Wang, H., Gill, C.O., 2014. Effects of mild and pasteurizing heat

- 545 treatments on survival of generic and verotoxigenic *Escherichia coli* from beef enrichment
546 cultures. Food Control 39, 100–104.
- 547 Yang, X., Badoni, M., Gill, C.O., 2011. Use of propidium monoazide and quantitative PCR for
548 differentiation of viable *Escherichia coli* from *E. coli* killed by mild or pasteurizing heat
549 treatments. Food Microbiology 28, 1478–1482.
- 550 Youssef, M.K., Yang, X., Badoni, M., Gill, C.O., 2012. Effects of spray volume, type of surface
551 tissue and inoculum level on the survival of *Escherichia coli* on beef sprayed with 5% lactic
552 acid. Food Control 25, 717–722.
- 553 Zhou, B. Chen, B., Wu, X., Li, F., Yu, P., Aguilar, Z., Wei, H., Xu, H., 2016. A new application
554 of a sodium deoxycholate-propidium monoazide quantitative PCR assay for rapid and
555 sensitive detection of viable *Cronobacter sakazakii* in powdered infant formula. Journal of
556 Dairy Science 99, 9550-9559.
- 557
- 558

Table 1. Enumeration of *E. coli* O157:H7 (Log CFU/cm²) Recovered on LB and VRB agars from Beef Steaks held for 30 min after Interventions of Lactic Acid (LA), Peroxyacetic Acid (PA) and Hot Water (HW) and after Treatment of cells with PMA or PMA with Deoxycholate.

Sample	Intervention					
	Lactic Acid		Peroxyacetic Acid		Hot Water	
	LB	VRB	LB	VRB	LB	VRB
Control	7.24 ± 0.08	6.85 ± 0.27	7.00 ± 0.40	6.70 ± 0.70	7.35 ± 0.09 ^a	7.03 ± 0.18 ^a
Intervention	6.41 ± 0.14	5.27 ± 0.69	6.48 ± 0.13	6.08 ± 0.57	6.19 ± 0.54 ^{ab}	5.65 ± 0.62 ^{ab}
Control + PMA	6.65 ± 0.30	6.45 ± 0.17	6.98 ± 0.45	6.63 ± 0.79	7.57 ± 0.16 ^a	7.11 ± 0.46 ^a
Intervention + PMA	5.91 ± 0.54	5.44 ± 0.45	6.29 ± 0.61	5.75 ± 1.17	6.16 ± 0.64 ^{ab}	5.88 ± 0.77 ^{ab}
Control + PMA + DC	6.57 ± 0.09	6.15 ± 0.70	6.41 ± 0.16	6.22 ± 0.25	6.46 ± 0.22 ^{ab}	6.09 ± 0.48 ^{ab}
Intervention + PMA + DC	5.58 ± 0.47	4.85 ± 0.94	5.70 ± 0.41	5.62 ± 0.31	5.20 ± 0.72 ^b	4.85 ± 0.68 ^b

Values are means ± standard deviation, n=3. Values with superscripts are significantly different among values in the same column; those that do not share a common superscript are significantly different ($p < 0.05$).



Note: Leave 1 steak uninoculated for quality control purposes. Do not apply intervention; place in buffer, massage and only sample for control plating and control DNA/qPCR analysis.

Figure 1. Experimental flow chart per trial.




Figure 2. (A) Correlation of cell counts on LB agar & qPCR DNA quantification of *E. coli* O157:H7 recovered from beef after lactic acid (LA) intervention & PMA or PMA + deoxycholate (DC) treatment. Circle, control; triangle, PMA; square, PMA + DC; black, control 0 min; white, LA 0 min; dark grey, control 30 min; light grey, LA 30 min. Treatments plotted at log CFU/cm² of respective control or LA sample at 0 or 30 min. Values are means of three trials with technical repeat (n=3), standard error of means of LB \leq 0.32; DNA \leq 0.72. †Significantly different log gene copy #/cm² from Control at 0 & 30 min ($p < 0.05$) *Significantly different log gene copy #/cm² from Control, Control + PMA, Control + PMA + DC and LA at both 0 & 30 min ($p < 0.05$) Correlation coefficients r^2 of the linear regressions were 0.94 (control), 0.88 (PMA) and 0.94 (PMA + DC). (B) Correlation of cell counts on LB agar & RTqPCR RNA quantification of *E. coli* O157:H7 recovered from beef after LA intervention. Diamond, rRNA; triangle, mRNA; black, control 0 min; white, LA 0 min; dark grey, control 30 min; light grey, LA 30 min. Treatments plotted at log CFU/cm² of respective control or LA sample at 0 or 30 min. Values are means of

three trials (n=3) with technical repeat, standard error of means of rRNA ≤ 0.68 ; mRNA ≤ 0.65 .

Correlation coefficients r^2 of the linear regressions were 0.49 (rRNA) and 0.55 (mRNA).




Figure 3. (A) Correlation of cell counts on LB agar & qPCR DNA quantification of *E. coli* O157:H7 recovered from beef after peroxyacetic acid (PA) intervention and PMA or PMA with deoxycholate treatment. Circle, control; triangle, PMA; square, PMA with deoxycholate; black, control 0 min; white, PA 0 min; dark grey, control 30 min; light grey, PA 30 min. Treatments plotted at log CFU/cm² of respective control or PA sample at 0 or 30 min. Values are means of three trials with technical repeat (n=3), standard error of means of LB \leq 0.39; DNA \leq 0.68. Correlation coefficients r^2 of the linear regressions were 0.83 (control), 0.87 (PMA) and 0.94 (PMA + DC). **(B)** Correlation of cell counts on LB agar & RTqPCR RNA quantification of *E. coli* O157:H7 recovered from beef after PA intervention. Diamond, rRNA; triangle, mRNA; black, control 0 min; white, PA 0 min; dark grey, control 30 min; light grey, PA 30 min. Treatments plotted at log CFU/cm² of respective control or PA sample at 0 or 30 min. Values are means of three trials with technical repeat (n=3), standard error of means of rRNA \leq 0.35; mRNA \leq 1.12. Correlation coefficients r^2 of the linear regressions were 0.87 (rRNA) and 1.00 (mRNA).




Figure 4. (A) Correlation of cell counts on LB agar & qPCR DNA quantification of *E. coli* O157:H7 recovered from beef after hot water (HW) intervention & PMA or PMA with deoxycholate treatment. Circle, control; triangle, PMA; square, PMA with deoxycholate; black, control 0 min; white, HW 0 min; dark grey, control 30 min; light grey, HW 30 min.. Treatments plotted at log CFU/cm² of respective control or HW sample. Values are means of three trials with technical repeat (n=3), standard error of means of LB \leq 0.54; DNA \leq 0.93. *Significantly different log gene copy #/cm² from Control, Control + PMA, Control + PMA + DC, HW, HW + PMA at 0 and 30 min ($p < 0.05$). Correlation coefficients r^2 of the linear regressions were 0.99 (control), 0.93 (PMA) and 0.92 (PMA + DC). (B) Correlation of cell counts on LB agar & RTqPCR RNA quantification of *E. coli* O157:H7 recovered from beef after HW intervention. Diamond, rRNA; triangle, mRNA; black, control 0 min; white, HW 0 min; dark grey, control 30 min; light grey, HW 30 min. Treatments plotted at log CFU/cm² of respective control or HW sample at 0 or 30 min. Values are means of three trials with technical repeat (n=3), standard error of means of rRNA

≤ 1.48 ; mRNA ≤ 1.20 . Correlation coefficients r^2 of the linear regressions were 0.66 (rRNA) and 0.75 (mRNA).