

## INFORMATION TO USERS

This manuscript has been reproduced from the microfilm master. UMI films the text directly from the original or copy submitted. Thus, some thesis and dissertation copies are in typewriter face, while others may be from any type of computer printer.

**The quality of this reproduction is dependent upon the quality of the copy submitted.** Broken or indistinct print, colored or poor quality illustrations and photographs, print bleedthrough, substandard margins, and improper alignment can adversely affect reproduction.

In the unlikely event that the author did not send UMI a complete manuscript and there are missing pages, these will be noted. Also, if unauthorized copyright material had to be removed, a note will indicate the deletion.

Oversize materials (e.g., maps, drawings, charts) are reproduced by sectioning the original, beginning at the upper left-hand corner and continuing from left to right in equal sections with small overlaps. Each original is also photographed in one exposure and is included in reduced form at the back of the book.

Photographs included in the original manuscript have been reproduced xerographically in this copy. Higher quality 6" x 9" black and white photographic prints are available for any photographs or illustrations appearing in this copy for an additional charge. Contact UMI directly to order.

**UMI<sup>®</sup>**

Bell & Howell Information and Learning  
300 North Zeeb Road, Ann Arbor, MI 48106-1346 USA  
800-521-0600



**University of Alberta**

**Cross-talk Between Nuclear Receptors in the Transcriptional Regulation of the Genes  
Encoding Peroxisomal Fatty-Acyl-CoA Oxidase and Enoyl-CoA Hydratase/3-  
Hydroxyacyl-CoA Dehydrogenase**

by

Christopher J. Winrow



A thesis submitted to the Faculty of Graduate Studies and Research in partial fulfillment of  
the requirements for the degree of Doctor of Philosophy

Department of Cell Biology and Anatomy

Edmonton, Alberta

Spring 1999



National Library  
of Canada

Acquisitions and  
Bibliographic Services

395 Wellington Street  
Ottawa ON K1A 0N4  
Canada

Bibliothèque nationale  
du Canada

Acquisitions et  
services bibliographiques

395, rue Wellington  
Ottawa ON K1A 0N4  
Canada

*Your file Votre référence*

*Our file Notre référence*

The author has granted a non-exclusive licence allowing the National Library of Canada to reproduce, loan, distribute or sell copies of this thesis in microform, paper or electronic formats.

The author retains ownership of the copyright in this thesis. Neither the thesis nor substantial extracts from it may be printed or otherwise reproduced without the author's permission.

L'auteur a accordé une licence non exclusive permettant à la Bibliothèque nationale du Canada de reproduire, prêter, distribuer ou vendre des copies de cette thèse sous la forme de microfiche/film, de reproduction sur papier ou sur format électronique.

L'auteur conserve la propriété du droit d'auteur qui protège cette thèse. Ni la thèse ni des extraits substantiels de celle-ci ne doivent être imprimés ou autrement reproduits sans son autorisation.

0-612-39606-1

**University of Alberta**

**Library Release Form**

**Name of Author:** Christopher J. Winrow


**Title of Thesis:** Cross-talk Between Nuclear Receptors in the Transcriptional Regulation of the Genes Encoding Peroxisomal Fatty-Acyl-CoA Oxidase and Enoyl-CoA Hydratase/3-Hydroxyacyl-CoA Dehydrogenase.

**Degree:** Doctor of Philosophy

**Year this Degree Granted:** 1999

Permission is hereby granted to the University of Alberta to reproduce single copies of this thesis and to lend or sell such copies for private, scholarly, or scientific research purposes only.

The author reserves all other publication and other rights in association with the copyright in the thesis, and except as hereinbefore provided, neither the thesis nor any substantial portion thereof may be printed or otherwise reproduced in any material form whatever without the author's prior written permission.






Christopher J. Winrow  
#205, 2 Ackerman St.  
Picton, Ontario  
Canada K0H 2T0

Date: JANUARY 26 1999

University of Alberta

Faculty of Graduate Studies and Research

The undersigned certify that they have read, and recommend to the Faculty of Graduate Studies and Research for acceptance, a thesis entitled: Cross-talk Between Nuclear Receptors in the Transcriptional Regulation of the Genes Encoding Peroxisomal Fatty-Acyl-CoA Oxidase and Enoyl-CoA Hydratase/3-Hydroxyacyl-CoA Dehydrogenase, submitted by Christopher J. Winrow in partial fulfillment of the requirements for the degree of Doctor of Philosophy.

  
\_\_\_\_\_  
Dr. Richard A. Rachubinski  
\_\_\_\_\_  
Dr. Luis B. Agellon  
\_\_\_\_\_  
Dr. Luc Berthiaume  
\_\_\_\_\_  
Dr. Tom C. Hobman  
\_\_\_\_\_  
Dr. Ellen K. Shibuya  
\_\_\_\_\_  
for Dr. Barry M. Forman

Date: JANUARY 26, 1999

## ABSTRACT

Peroxisomes are organelles which are responsible for the  $\beta$ -oxidation of long chain fatty acids and the metabolism of hydrogen peroxide. The metabolic activity of peroxisomes can be stimulated by administration of a structurally diverse group of compounds collectively termed peroxisome proliferators. These agents include fibrate family hypolipidemic drugs, antidiabetic thiazolidinediones, eicosanoids, as well as various naturally occurring and synthetic mono- and polyunsaturated fatty acids. The transcription of genes encoding enzymes responsible for peroxisomal  $\beta$ -oxidation is regulated by a group of nuclear receptors known as peroxisome proliferator-activated receptors (PPARs). PPARs form heterodimers with the 9-*cis*-retinoic acid receptor (RXR $\alpha$ ) and bind to cognate peroxisome proliferator-response elements upstream of target genes. PPARs govern the regulation of a number of genes involved in lipid metabolism and homeostasis including those encoding the peroxisomal  $\beta$ -oxidation enzymes fatty acyl-CoA oxidase and enoyl-CoA hydratase/3-hydroxyacyl-CoA dehydrogenase, enzymes which carry out  $\omega$ -hydroxylation, as well as apolipoproteins AI and CIII and fatty acid binding protein. The focus of my research has been to investigate the interplay between nuclear receptors and auxiliary factors in directing gene expression, with particular attention to the regulation of peroxisomal  $\beta$ -oxidation genes via the peroxisome proliferator signalling pathway. The results of my studies have established the involvement of hepatocyte nuclear factor-4, thyroid hormone receptor  $\alpha$ , retinoid-like Z receptor, RXR $\alpha$  and PPAR $\alpha$  in both the positive and negative transcriptional regulation of genes encoding core enzymes responsible for fatty acid metabolism. Calreticulin is a Ca<sup>2+</sup>-binding protein chaperone, suggested to be a general regulator of nuclear receptor activity. The results I have

obtained indicate that although calreticulin interacts with nuclear receptors *in vitro*, it does not have a general effect on all classes of nuclear receptors in a cellular context. My research has demonstrated multilevel control by distinct signalling networks in the regulation of genes involved in peroxisomal  $\beta$ -oxidation of fatty acids, thereby providing suitable targets for pharmacological intervention and contributing to our basic understanding of nuclear receptor signalling and the peroxisome proliferator response.



## **ACKNOWLEDGEMENTS**

I would first like to thank my supervisor, Rick Rachubinski for his helpful guidance, enthusiastic support and friendship. You have provided me with insight and direction, while at the same time fostering my growth as an independent researcher.

I wish to acknowledge my friends and colleagues in the “Rach Lab”, both the proliferators: Sandra Marcus, Pam Lagali and Altaf Kassam, and the biogenesis group: Rachel Szilard, Gary Eitzen, Jenn Smith, Melchior Evers, Vladimir Titorenko, Cleofe Hurtado, Gareth Lambkin and Trevor Brown. From adventures in the wilds of Alberta, to navigating the perils of the lab as a rookie, your help and friendship has made my time as a grad student a fun and rewarding experience. I would also like to recognize the contributions of my supervisory committee members, Manijeh Pashar, Steve Rice, Luc Berthiaume, Tom Hobman, Ellen Shibuya, Lou Agellon, Barry Marc Forman and our collaborators at McMaster University, John Capone, Kenji Miyata and John Hunter. Your ideas and opinions have helped to guide my research. In addition, I am thankful for the financial support of the Alberta Heritage Foundation for Medical Research and the Heart and Stroke Foundation of Canada.

I would finally like to express my sincere appreciation for the support which my family (the Winrows, Legates and Crodens), and my wife, Dana, have given me over the years. Your words and love have carried me through the tough times, and rejoiced in my accomplishments.

To all of these, I give my heartfelt thanks.

## TECHNICAL ACKNOWLEDGEMENTS

I am grateful for the efforts of those who have made the day to day laboratory operations run smoothly. My thanks go to Kris Decker, Rachel Oates and Gareth Lambkin for their assistance. I would especially like to recognize the contributions of Eileen Reklow, whose expertise in tissue culture has been invaluable.

My thanks are extended to my co-authors on the publications included in my thesis. In particular I wish to acknowledge the contributions of S.L. Marcus, K.S. Miyata and B. Zhang on the HNF-4 work in Chapter 2; S.L. Marcus and K.S. Miyata on the calreticulin research in Chapter 3; A. Kassam, J. Hunter, K.S. Miyata and S.L. Marcus with the TR research described in Chapter 4. Thank you for your insight and assistance.

The support of our collaborators who have provided reagents and plasmids is greatly appreciated, as are the contributions of others at the University of Alberta who have shared their techniques and experience with me.

## TABLE OF CONTENTS

<b>Chapter 1</b>	<b>-</b>	<b>Introduction</b>	<b>1</b>
1.1		Introductory Statement	2
1.2		Peroxisome Proliferators	3
1.3		Peroxisomal $\beta$ -Oxidation	6
1.4		Nuclear Receptor Response Elements	10
1.4.1		Peroxisome Proliferator Response Elements (PPRE)	10
1.4.2		Regulation of Genes Encoding Peroxisomal $\beta$ -Oxidation Enzymes	12
1.5		PPARs and the Nuclear Receptor Superfamily	13
1.5.1		General Structure of Nuclear Receptors	14
1.5.2		Dimerization Requirements	17
1.5.3		Orphan Receptors and the Role of Ligand Binding	18
1.6		Peroxisome Proliferator Activated Receptors (PPAR)	18
1.6.1		PPAR $\alpha$	19
1.6.2		PPAR $\gamma$	20
1.6.3		PPAR $\delta$	21
1.7		Hepatocyte Nuclear Factor-4 (HNF-4)	22
1.8		Thyroid Hormone Receptors (TR)	23
1.9		Retinoid-like Z Receptors/Retinoid Orphan Receptors (RZR/ROR)	24
1.10		Calreticulin	26
1.11		Nuclear Receptor Co-Regulatory Factors	27
1.11.1		The Activation Function Region	28
1.11.2		Nuclear Receptor Co-Activators	29
1.11.3		Nuclear Receptor Co-Repressors	31
1.11.4		Model of Co-Activator/Co-Repressor Interactions	33

1.12 Nuclear Receptor Interplay	33
1.13 Aims of This Project	37
<b>Chapter 2 - Transactivation of the Peroxisome Proliferator-Activated Receptor is Differentially Modulated by Hepatocyte Nuclear Factor-4</b>	<b>39</b>
2.1 Abstract	40
2.2 Introduction	41
2.3 Materials and Methods	44
2.3.1 Plasmids	44
2.3.2 Transfections and Measurement of Luciferase Activity	45
2.3.3 In Vitro Transcription/Translation	47
2.3.4 Electrophoretic Mobility Shift Analysis	47
2.4 Results	48
2.4.1 HNF-4 Interacts with the AOx- and HD-PPREs	48
2.4.2 HNF-4 Represses rPPAR-dependent Induction from the AOx-PPRE both in the Presence and Absence of a Potent Peroxisome Proliferator	54
2.4.3 HNF-4 Potentiates Activity of rPPAR on the HD-PPRE in the Presence of a Peroxisome Proliferator	56
2.4.4 HNF-4 Does Not Cooperate with RXR $\alpha$ or rPPAR for DNA Binding	59
2.5 Discussion	61
<b>Chapter 3 - Calreticulin Modulates the In Vitro DNA Binding But Not The In Vivo Transcriptional Activation by Peroxisome Proliferator-Activated Receptor/Retinoid X Receptor Heterodimers</b>	<b>66</b>
3.1 Abstract	67
3.2 Introduction	68
3.3 Materials and Methods	70

3.3.1	Plasmids, Cell Culture, and Transfection	70
3.3.2	Electrophoretic Mobility Shift Assays	71
3.4	Results	71
3.4.1	Calreticulin Inhibits PPAR/RXR Binding to DNA	71
3.4.2	Calreticulin Shows Little or No Effect on Transactivation by PPAR/RXR In Vivo	72
3.5	Discussion	79
<b>Chapter 4</b>	<b>- Interplay of the Peroxisome Proliferator-Activated Receptor and Thyroid Hormone Receptor Signalling Pathways in Regulating Peroxisome Proliferator-Responsive Genes</b>	<b>82</b>
4.1	Introduction	83
4.2	Materials and Methods	85
4.2.1	Plasmids	85
4.2.2	In Vitro Transcription/Translation	86
4.2.3	Gel Retardation Analysis	86
4.2.4	Transfection and Luciferase Assay	87
4.3	Results	88
4.3.1	TR $\alpha$ and TR $\alpha$ /RXR $\alpha$ Heterodimers Bind to the AOx-PPRE	88
4.3.2	TR $\alpha$ Competes with PPAR for Binding to the AOx-PPRE	95
4.3.3	Unliganded TR $\alpha$ Stimulates Transactivation by PPAR/RXR $\alpha$ Heterodimers	98
4.4	Discussion	107
<b>Chapter 5</b>	<b>- Cross-talk Between Orphan Nuclear Hormone Receptor RZR<math>\alpha</math> and Peroxisome Proliferator-Activated Receptor <math>\alpha</math> in Regulation of the Peroxisomal Hydratase-Dehydrogenase Gene</b>	<b>111</b>
5.1	Abstract	112
5.2	Introduction	113

5.3	Materials and Methods	115
5.3.1	Plasmids	115
5.3.2	In Vitro Transcription/Translation	116
5.3.3	Electrophoretic Mobility Shift Analysis	116
5.3.4	Transient Transfections and Measurement of Luciferase Activity	117
5.4	Results	118
5.4.1	RZR $\alpha$ Binds as a Monomer to the HD-PPRE but not the AOx-PPRE	118
5.4.2	RZR $\alpha$ and PPAR $\alpha$ /RXR $\alpha$ Bind to Different Regions of the HD-PPRE In Vitro	125
5.4.3	RZR $\alpha$ Potentiates Transactivation From the HD-PPRE by PPAR $\alpha$	125
5.4.4	Exogenous RXR $\alpha$ Abrogates the Potentiation Effect of RZR $\alpha$ on Transactivation by PPAR $\alpha$ From the HD-PPRE	134
5.5	Discussion	138
<b>Chapter 6</b>	<b>- General Discussion</b>	<b>142</b>
6.1	Discussion	143
6.1.1	Influence of HNF-4 on the AOx- and HD-PPREs	144
6.1.2	Calreticulin and PPAR $\alpha$ /RXR $\alpha$ Activity	149
6.1.3	The Effects of TR $\alpha$ on Peroxisome Proliferator Signalling	152
6.1.4	RZR $\alpha$ , PPAR $\alpha$ and RXR $\alpha$ Interplay	158
6.2	General Model of Nuclear Receptor Interactions with the AOx- and HD-PPREs	164
6.3	Summary	169
6.4	Conclusions	171
6.5	Bibliography	173

## LIST OF FIGURES

Figure 1-1	Comparison of peroxisomal and mitochondrial $\beta$ -oxidation pathways.	8
Figure 1-2	Nuclear receptor structure and comparison of PPAR isoforms.	15
Figure 1-3	Model of nuclear receptor accessory factor interactions.	34
Figure 2-1	Comparison of the AOx-PPRE and the HD-PPRE to the consensus HNF-4 binding site.	50
Figure 2-2	Endogenous HNF-4 related proteins of rat hepatoma H4IIEC3 cells bind to the AOx- and HD-PPREs.	51
Figure 2-3	HNF-4 synthesized <i>in vitro</i> binds to the AOx- and HD-PPREs.	53
Figure 2-4	HNF-4 antagonizes rPPAR-mediated induction of an AOx-PPRE reporter gene.	55
Figure 2-5	HNF-4 differentially modulates the function of rPPAR on the HD-PPRE.	57
Figure 2-6	HNF-4 alters transactivation from the AOx- and HD-PPREs in the presence of rPPAR and RXR $\alpha$ .	58
Figure 2-7	HNF-4 does not cooperate with rPPAR or RXR $\alpha$ for DNA binding.	60
Figure 3-1	The DNA-binding domains of nuclear hormone and steroid receptors contain sequences homologous to the calreticulin binding site.	73
Figure 3-2	Calreticulin inhibits PPAR/RXR heterodimer binding to the AOx-PPRE <i>in vitro</i> .	74
Figure 3-3	Calreticulin fails to repress the <i>in vivo</i> transactivation from the AOx-PPRE by PPAR/RXR.	76
Figure 3-4	Calreticulin represses transactivation from the GRE by GR.	78
Figure 4-1	Interaction of TR $\alpha$ with a synthetic DR4 element.	89
Figure 4-2	TR $\alpha$ binds to the AOx-PPRE on its own and in association with RXR $\alpha$ .	91
Figure 4-3	TR $\alpha$ is present in complexes formed on the AOx-PPRE.	93

Figure 4-4	TR $\alpha$ binds to the AOx-PPRE in a sequence-specific manner.	94
Figure 4-5	TR $\alpha$ competes with PPAR/RXR $\alpha$ for binding to the AOx-PPRE.	96
Figure 4-6	Effects of increasing amounts of RXR $\alpha$ and PPAR on the formation of TR $\alpha$ -dependent protein/DNA complexes.	97
Figure 4-7	TR $\alpha$ enhances transactivation by PPAR/RXR $\alpha$ <i>in vivo</i> .	99
Figure 4-8	TR $\alpha$ -mediated stimulation of Wy-14,643-dependent activation requires the presence of PPAR and RXR $\alpha$ .	102
Figure 4-9	TR $\alpha$ inhibits transactivation in the presence of T3.	104
Figure 4-10	Liganded TR $\alpha$ also inhibits Wy-14,643-mediated transactivation by PPAR/RXR $\alpha$ .	106
Figure 5-1	Comparison of the sequence of the RZR $\alpha$ consensus binding site to sequences within the AOx- and HD-PPREs.	119
Figure 5-2	RZR $\alpha$ binds as a monomer to the HD-PPRE but not the AOx-PPRE.	120
Figure 5-3	RZR $\alpha$ monomer binds to the HD-PPRE in a sequence-specific manner.	122
Figure 5-4	PPAR $\alpha$ /RXR $\alpha$ and RZR $\alpha$ bind independently to the HD-PPRE.	123
Figure 5-5	RZR $\alpha$ and PPAR $\alpha$ /RXR $\alpha$ do not affect each other's binding to the HD-PPRE.	126
Figure 5-6	RZR $\alpha$ monomer and PPAR $\alpha$ /RXR $\alpha$ heterodimer bind to different sites within the HD-PPRE <i>in vitro</i> .	128
Figure 5-7	RZR $\alpha$ potentiates transactivation by PPAR $\alpha$ from the HD-PPRE but not the AOx-PPRE.	131
Figure 5-8	RZR $\alpha$ alone weakly activates transcription from the HD-PPRE.	133
Figure 5-9	Melatonin does not stimulate transactivation by PPAR $\alpha$ /RZR $\alpha$ from the HD-PPRE.	135
Figure 5-10	RZR $\alpha$ does not potentiate transactivation by PPAR $\alpha$ from the HD-PPRE in the presence of exogenous RXR $\alpha$ .	136
Figure 6-1	Summary of results from studies of RZR $\alpha$ effects on peroxisome proliferator-mediated transactivation from the HD-PPRE.	159



Figure 6-2	Possible roles for RZR $\alpha$ in directing peroxisome proliferator-mediated transactivation from the HD-PPRE.	161
Figure 6-3	Overview of nuclear receptor crosstalk in the transactivation from the AOx- and HD-PPREs.	165

## LIST OF ABBREVIATIONS

ACTR	activator of the thyroid and retinoic acid receptors
AF	activation function
AOx	fatty acyl-CoA oxidase
AR	androgen receptor
ARE	adipocyte lipid-binding protein response element
BTM	basal transcription machinery
CARLA	co-activator-dependent receptor ligand assay
CBP	CREB-binding protein
COUP-TF	chicken ovalbumin upstream promoter transcription factor
CPS	carbamoyl phosphate synthetase
CREB	cAMP response element-binding protein
CRT	calreticulin
CoA	coenzyme A
CMV	cytomegalovirus
CYP	cytochrome P450
DBD	DNA-binding domain
DMEM	Dulbecco's modified Eagle's medium
DR <sub>n</sub>	direct repeat of TGACCT-like motifs separated by <i>n</i> nucleotide(s)
DTT	dithiothreitol
EDTA	ethylenediamine tetraacetic acid
EMSA	electrophoretic mobility shift analysis
ER	estrogen receptor
GR	glucocorticoid receptor
GRE	glucocorticoid response element
GRIP-1	GR-interacting protein 1
HD	enoyl-CoA hydratase/3-hydroxyacyl-CoA dehydrogenase
HMG-CoA	3-hydroxy-3-methylglutaryl-CoA
HNF	hepatocyte nuclear factor
LBD	ligand-binding domain
LTB <sub>4</sub>	leukotriene B <sub>4</sub>
<i>luc</i>	luciferase gene
LXR	liver X receptor
MAP kinase	mitogen-activated protein kinase
NCoA	nuclear receptor co-activator
NCoR	nuclear receptor co-repressor
PBP	PPAR $\gamma$ -binding protein
p/CAF	p300/CBP-associated factor
p/CIP	p300/CBP co-integrator associate protein
PGC	PPAR $\gamma$ -co-activator
PPAR	peroxisome proliferator-activated receptor
PPRE	peroxisome proliferator-response element
RAC3	receptor-associated co-activator 3

RAR	retinoic acid receptor
RIP-140	receptor-interacting protein 140
ROR	retinoid-like orphan receptor
RXR $\alpha$	9- <i>cis</i> retinoic acid receptor
RZR	retinoid-like Z receptor
SDS	sodium dodecyl sulfate
SMRT	silencing mediator for RARs and TRs
SRC-1	steroid receptor co-activator-1
STAT	signal transducer and activator of transcription
T3	3,3',5-triiodothyronine
TIF-2	transcriptional intermediary factor 2
TRAM	TR-activator molecule
TR	thyroid hormone receptor
TRE	thyroid hormone response element
VDR	vitamin D3 receptor

# **CHAPTER 1**

## **INTRODUCTION**

## 1.1 Introductory Statement

Peroxisomes are organelles bounded by a single membrane that are responsible for a myriad of cellular functions, including the  $\beta$ -oxidation of fatty acids, the degradation of  $H_2O_2$  and other toxins, gluconeogenesis, as well as cholesterol and bile acid biosynthesis (Bentley et al., 1993; Hiltunen et al., 1996; Mannaerts and Van Veldhoven, 1993; Tolbert, 1981). Over 30 years ago, it was observed that certain agents, later named peroxisome proliferators, could increase both the number and metabolic capacity of peroxisomes (Hess et al., 1965; Paget, 1963; Svoboda and Azarnoff, 1966). Within the past decade, it was determined that novel members of the nuclear receptor superfamily, termed peroxisome proliferator-activated receptors (PPAR), are central mediators of peroxisomal metabolism through transcriptional regulation of genes encoding the core peroxisomal enzymes (Issemann and Green, 1990; Latruffe and Vamecq, 1997; Schoonjans et al., 1996). Substantial roles for PPARs in the regulation of cell growth and development, lipid metabolism and storage, and signal transduction have since been established (for reviews see Desvergne et al., 1998; Reddy and Chu, 1996). PPARs are ligand-modulated transcription factors that direct gene expression through interactions with peroxisome proliferator-response elements (PPRE) located upstream of target genes (Forman et al., 1996; Latruffe and Vamecq, 1997; Schoonjans et al., 1997). We have endeavoured to explore the basic mechanisms by which nuclear receptors cross-talk in order to modulate the transcriptional activity of the genes encoding the first two enzymes of the peroxisomal  $\beta$ -oxidation pathway in response to peroxisome proliferators.

## 1.2 Peroxisome Proliferators

Peroxisome proliferators were first identified in studies of fibrate drugs, in which it was observed that administration of clofibrate and other hypolipidemic agents caused an induction of peroxisomes in rodent hepatocytes (Hess et al., 1965; Paget, 1963; Svoboda and Azarnoff, 1966). Peroxisome proliferators constitute a class of structurally diverse agents that have marked effects at both the cellular level and in the context of the whole animal. The effects observed in rodents exposed to peroxisome proliferators include hepatomegaly, hepatocarcinogenesis and the induction of genes for both peroxisomal and non-peroxisomal enzymes (Hawkins et al., 1987; Hijikata et al., 1987; Lock et al., 1989; Reddy and Lalwai, 1983; Reddy et al., 1986b). Although chronic exposure to peroxisome proliferators leads to hepatocarcinogenesis in rodents, humans appear to be refractory to the effects of these agents (Bentley et al., 1993; Bieri and Lhuguenot, 1993; Cattley et al., 1998). Peroxisome proliferators are classified as non-genotoxic hepatocarcinogens, as they do not directly interact with, or damage DNA (Shaw and Jones, 1994). Administration of peroxisome proliferators to responsive cells results in a dramatic increase in the absolute number and metabolic activity of peroxisomes (Hawkins et al., 1987; Lazarow and De Duve, 1976; Osumi and Hashimoto, 1978). In particular, the transcription of genes encoding the enzymes responsible for peroxisomal  $\beta$ -oxidation is stimulated (Hijikata et al., 1990; McQuaid et al., 1987; Osumi, 1993; Ozasa et al., 1985; Reddy et al., 1986a; Thangada et al., 1989). Peroxisome proliferators include fibrate hypolipidemic drugs, non-steroidal anti-inflammatory agents, eicosanoids, phthalate ester plasticizers, thiazolidinedione antidiabetic compounds, chlorinated hydrocarbons, as well as naturally occurring and synthetic mono- and

polyunsaturated fatty acids (Forman et al., 1997a; Kliewer et al., 1997; Krey et al., 1997; Lee et al., 1997; Lehmann et al., 1997; Reddy and Chu, 1996; Youssef and Badr, 1998; Zhou and Waxman, 1998). Their actions are mediated by the PPARs (Issemann and Green, 1990). The ability of such structurally diverse compounds to activate PPARs, and the failure to identify directly-interacting PPAR agonists led to hypothesis that peroxisome proliferators are reduced to a common metabolite which activates PPARs.

Recently, 15-deoxy- $\Delta^{12,14}$ -prostaglandin J<sub>2</sub> and the thiazolidinedione, BRL49653, have been shown to bind directly to, and to activate the  $\gamma$  isoform of PPAR (Kliewer et al., 1997; Lehmann et al., 1995). Leukotriene B<sub>4</sub> and the hypolipidemic drug, Wy-14,643, have been identified as *bona fide* ligands for the PPAR $\alpha$  isoform (Forman et al., 1997a; Kliewer et al., 1997; Krey et al., 1997; Schoonjans et al., 1997). An *in vitro* conformation-based assay that monitors PPAR-DNA binding has shown that specific fatty acids, eicosanoids and fibrate hypolipidemic drugs also act directly as ligands for PPAR $\alpha$  and/or PPAR $\delta$  (Forman et al., 1997a). Additionally, an *in vitro* study employing a high-affinity radiolabelled ligand for PPAR $\alpha$  and PPAR $\gamma$  demonstrated the binding of specific mono- and polyunsaturated fatty acids to both these receptors (Kliewer et al., 1997). This study also showed that the eicosanoid 8(*s*)-hydroxyeicosatetraenoic acid binds selectively to PPAR $\alpha$  whereas 15-deoxy- $\Delta^{12,14}$ -prostaglandin J<sub>2</sub> binds directly to PPAR $\gamma$  (Kliewer et al., 1997). Structural analysis has indicated that the large, ligand-binding pocket of PPARs may account for the capacity of these receptors to directly interact with a relatively large number of structurally distinct ligands (Nolte et al., 1998).

Although chronic exposure to peroxisome proliferators leads to hepatocarcinogenesis

in most rodents, exposure to these agents has not been shown to induce hepatomegaly or hepatocarcinogenesis in humans (Bentley et al., 1993; Bieri and Lhuguenot, 1993; Cattley et al., 1998). Guinea pigs are also not susceptible to the tumorigenic effects of peroxisome proliferators (Bell et al., 1998; Bentley et al., 1993; Lake and Gray, 1985). This lack of a carcinogenic response may be due to the reduced level of PPAR $\alpha$  found in humans and guinea pigs (Bell et al., 1998; Cattley et al., 1998; Palmer et al., 1998). Guinea pig PPAR $\alpha$  is functional, as it has been shown to be capable of inducing transcription from a PPRE-reporter construct in response to peroxisome proliferators (Bell et al., 1998; Tugwood et al., 1998). PPAR $\alpha$  is present at a 10-fold lower level in human liver than in responsive species, and human PPAR $\alpha$  may contain a splice variant form that decreases its proliferative effects (Palmer et al., 1998).

Several models have been proposed by which peroxisome proliferators may induce hepatocarcinogenesis (Bentley et al., 1993; Christensen et al., 1998; Peters et al., 1997a; Rao and Reddy, 1991). In rodents, the hallmarks of peroxisome proliferators are their ability to increase the levels of fatty acyl-CoA oxidase, peroxisomal fatty acid oxidation and excess production of H<sub>2</sub>O<sub>2</sub> and to stimulate cell proliferation and the expression of PPAR $\alpha$  (Cattley et al., 1998). The elevation of H<sub>2</sub>O<sub>2</sub> without a concomitant rise in H<sub>2</sub>O<sub>2</sub>-degrading peroxisomal catalase may result in oxidative DNA damage (Rao and Reddy, 1991; Reddy and Rao, 1989; Reddy et al., 1986b). Peroxisome proliferators have also been found to be mitogenic, since they increase DNA synthesis in primary hepatocyte cultures in a glucocorticoid-dependent manner (Plant et al., 1998). It appears that stimulation of the glucocorticoid receptor dramatically increases the transcription of the PPAR $\alpha$  gene, and as



a result, makes cells more responsive to peroxisome proliferators (Plant et al., 1998). The recent observation that PPAR inhibits *bcl-2* expression furthers the hypothesis that peroxisome proliferators act as carcinogens by suppressing apoptosis (Christensen et al., 1998; Schulte-Hermann et al., 1991). Direct evidence for the involvement of PPAR $\alpha$  in mediating peroxisome proliferator-induced hepatocarcinogenesis comes from studies of transgenic mice in which PPAR $\alpha$  has been eliminated. In these studies, 100% of wild-type mice exposed to the peroxisome proliferator Wy-14,643 developed multiple hepatocellular neoplasms, whereas none of the PPAR $\alpha$  *-/-* mice were affected under the same conditions (Peters et al., 1997a).

### 1.3 Peroxisomal $\beta$ -Oxidation


Peroxisomal  $\beta$ -oxidation is the main metabolic pathway for the degradation of fatty acids with chain length greater than 18 carbon atoms (*i.e.* C18), typically referred to as very long-chain fatty acids (Mannaerts and van Veldhoven, 1996). This pathway is also employed to metabolize medium and long chain fatty acids with chain lengths between C10-C18, especially in conditions where the capacity of mitochondrial  $\beta$ -oxidation is exceeded (Hashimoto, 1996; Lehninger et al., 1993; Mannaerts and van Veldhoven, 1996; Tolbert, 1981). Peroxisomal  $\beta$ -oxidation is incomplete since fatty acids shorter than C8 cannot be oxidized in peroxisomes. The C8-fatty acids generated by peroxisomal  $\beta$ -oxidation are converted to carnitine esters by peroxisomal carnitine acyl-transferase and directed to the mitochondria for further oxidation (Mannaerts and van Veldhoven, 1996).

The peroxisomal  $\beta$ -oxidation cycle consists of three core enzymes: fatty acyl-CoA

oxidase (AOx), enoyl-CoA hydratase/3-hydroxyacyl-CoA dehydrogenase bifunctional enzyme (HD), and 3-ketoacyl-CoA thiolase. Peroxisomes also contain catalase for the metabolism of  $H_2O_2$  (Hashimoto, 1996; Lehninger et al., 1993). Peroxisomal and mitochondrial  $\beta$ -oxidation of fatty acids both require fatty acyl-CoA as the initial substrate; however, the two forms of  $\beta$ -oxidation are carried out by distinct enzymes and differ in several important respects (see Fig. 1-1) (Eaton et al., 1996; Hashimoto, 1996; Lehninger et al., 1993; Mannaerts and van Veldhoven, 1996). In the first step of peroxisomal  $\beta$ -oxidation, electrons transferred to FAD are directly passed to  $O_2$ , yielding  $H_2O_2$ , which is then broken down to  $H_2O$  and  $O_2$  by peroxisomal catalase. In mitochondria, electrons transferred to FAD proceed through the respiratory chain via the electron-transferring flavoprotein in the mitochondrial inner membrane to yield  $H_2O$  and ATP. Another difference between these two forms of fatty acid  $\beta$ -oxidation is that the NADH produced through the action of enoyl-CoA hydratase/3-hydroxyacyl CoA dehydrogenase bifunctional protein is not oxidized within peroxisomes and must be shuttled outside and eventually passed on to the mitochondria, which are capable of oxidizing NADH. Similarly, acetyl-CoA produced in peroxisomes is exported and can be further oxidized by the citric acid cycle in mitochondria. Peroxisomal  $\beta$ -oxidation is preferential for medium-, long- and very long-chain fatty acids (C10-C30), whereas mitochondria carry out the  $\beta$ -oxidation of short-, medium- and long-chain fatty acids (up to C18) (Eaton et al., 1996; Hashimoto, 1996; Lehninger et al., 1993; Mannaerts and van Veldhoven, 1996). In mitochondria, each of the  $\beta$ -oxidation enzymes is a soluble, separate protein; however, the enzymes of the peroxisomal  $\beta$ -oxidation cascade form a multifunctional protein complex (Lehninger et al., 1993). Since plant mitochondria do not perform  $\beta$ -

Mitochondrial

Peroxisomal



**Figure 1-1 Comparison of peroxisomal and mitochondrial  $\beta$ -oxidation pathways.**

oxidation, plants contain specialized peroxisomes referred to as glyoxysomes that are responsible for the  $\beta$ -oxidation of stored lipids and the conversion of triacylglycerols to glucose and essential metabolites (Hashimoto, 1996; Kindl, 1993; Lehninger et al., 1993). It is thought that the main function of mitochondrial  $\beta$ -oxidation in animals is the production of ATP, and that peroxisomal  $\beta$ -oxidation is important for the metabolism of toxic very long-chain fatty acids, medium chain dicarboxylic acids, cholesterol and unsaturated fatty acids that are not effectively oxidized in the mitochondria (Hashimoto, 1996; Mannaerts and van Veldhoven, 1996).

Administration of peroxisome proliferators increases the metabolic activity of peroxisomes and dramatically alters the peroxisomal enzyme content in a PPAR-dependent manner. In particular, the levels of the core enzymes of the peroxisomal  $\beta$ -oxidation cycle, AOX, HD and 3-ketoacyl-CoA thiolase, are significantly increased in response to peroxisome proliferators (Hijikata et al., 1990; McQuaid et al., 1987; Osumi, 1993; Ozasa et al., 1985; Reddy et al., 1986a; Thangada et al., 1989). Interestingly, the level of peroxisomal catalase is not concomitantly increased with the increases in the core  $\beta$ -oxidation enzymes. This dysregulation is thought to lead to an increase in highly reactive  $H_2O_2$  radicals, and has been proposed to be responsible for the hepatocarcinogenic effects of peroxisome proliferators through oxidative DNA damage (Rao and Reddy, 1991; Reddy and Rao, 1989; Reddy et al., 1986b). Given that many fatty acids function as peroxisome proliferators, this provides a feedback loop to stimulate  $\beta$ -oxidation and  $\omega$ -hydroxylation in response to elevated fatty acid levels, and to govern the inhibition of  $\beta$ -oxidation in conditions where fatty acid levels are reduced.

## 1.4 Nuclear Receptor Response Elements

Members of the nuclear receptor superfamily mediate their actions on transcription by binding to response elements located within the promoter regions of target genes (Evans, 1988; Gronemeyer, 1992; Keightley, 1998; Meier, 1997; Parker, 1993; Umesono and Evans, 1989). PPARs belong to the retinoid/vitamin D<sub>3</sub>/thyroid receptor subfamily of nuclear receptors that recognize core binding sites consisting of repeats of the consensus sequence TGACCT (Latruffe and Vamecq, 1997; Schoonjans et al., 1996). Several layers of specificity are contained within the arrangement of nuclear receptor response elements, and include different requirements in terms of the orientation, spacing, core consensus sequence and sequences flanking the TGACCT-like repeats (Castelein et al., 1997; IJpenberg et al., 1997; Juge-Aubry et al., 1997; Mader et al., 1993; Olson and Koenig, 1997; Osada et al., 1997; Umesono et al., 1991).

### *1.4.1 Peroxisome Proliferator Response Elements*

PPAR heterodimerizes with 9-*cis*-retinoic acid receptor (RXR $\alpha$ ) and binds preferentially to PPREs that contain 1 nucleotide spacing between direct tandem repeats, referred to as DR1 spacing (Chu et al., 1995a; Gearing et al., 1993; Keller et al., 1993; Kliewer et al., 1992b). It has been reported that PPAR/RXR heterodimers favour PPREs in which the separating nucleotide is A or T (Johnson et al., 1996; Juge-Aubry et al., 1997; Palmer et al., 1995). The absolute requirement for orientation and spacing varies widely between different nuclear receptors (Glass, 1994; Green, 1993; Mader et al., 1993). The thyroid hormone (TR) and retinoic acid receptors (RAR) bind to response elements which

typically contain DR3, 4 or 5 spacing. TR also recognizes sites consisting of palindromic or inverted palindromic repeats with variations in spacing (Desvergne, 1994; Lazar et al., 1991b; Ribeiro et al., 1995b; Tsai and O'Malley, 1994). Other nuclear receptors exhibit greater promiscuity in binding to response elements. The chicken ovalbumin upstream promoter transcription factor (COUP-TF) represses transactivation by binding to a large number of differentially spaced and oriented repeats (Cooney et al., 1993; Cooney et al., 1992; Miyata et al., 1993). This variation in binding preferences accounts for the interplay between different nuclear receptors through competition for shared target sites. Additionally some nuclear receptors can accommodate alterations in the core TGACCT-like motif and will bind readily to response elements that harbor variations in the core sequence (Green, 1993; Meier, 1997). PPREs have been characterized that contain minor variations in sequence, yielding a broad consensus site of TGAC/ACR (Ijpenberg et al., 1997; Juge-Aubry et al., 1997; Palmer et al., 1995).

It is becoming more apparent that the sequences adjacent to the core sites within PPREs exhibit a role in directing the binding of PPAR/RXR heterodimers to these sites (Castelein et al., 1997; Ijpenberg et al., 1997; Juge-Aubry et al., 1997; Osada et al., 1997; Palmer et al., 1995). Studies of a number of natural and synthetic PPREs that examined additional sequence requirements and heterodimer polarity determined that PPAR binds to the 3' TGACCT-like site and RXR binds to the 5' repeat (Chu et al., 1995a; Ijpenberg et al., 1997; Juge-Aubry et al., 1997). This is the opposite binding arrangement of most other RXR heterodimerization partners such as TR and RAR, and may be due to intrinsic differences within specific response elements and to sites within the heterodimerization interfaces of

PPAR and RXR. Interestingly, the sequence flanking the 3' TGACCT-like repeat has been demonstrated to direct differential binding of distinct PPAR isoforms. PPAR $\gamma$  binds to, and transactivates readily from, the adipocyte lipid-binding protein AP2 response element (ARE6)-PPRE, whereas PPAR $\alpha$  is unable to associate with this response element (Juge-Aubry et al., 1997). Replacement of only the ARE6-PPRE 3' flanking region with that of 3-hydroxy-3-methyl-glutaryl-CoA (HMG-CoA) synthase-PPRE enables the binding of PPAR $\alpha$  to this hybrid response element (Juge-Aubry et al., 1997). The arrangement of response elements permits interplay between nuclear receptors based on specificity at the level of DNA binding. Alterations in the core sequence, spacing and orientation of TGACCT-like repeats and flanking sequences, coupled with the degree of DNA binding promiscuity of certain nuclear receptors, enable the differential regulation of target genes by distinct nuclear receptors via their respective response elements (Glass, 1994; Naar et al., 1991).

#### *1.4.2 Regulation of Genes Encoding peroxisomal $\beta$ -Oxidation Enzymes*

Peroxisome proliferators were shown to induce the levels of peroxisomal  $\beta$ -oxidation enzymes and subsequently found to increase transcription of a number of genes, including those encoding enzymes of the peroxisomal  $\beta$ -oxidation pathway (Dreyer et al., 1993; Lazarow and De Duve, 1976; McQuaid et al., 1987; Osumi and Hashimoto, 1978; Reddy et al., 1986a). Closer examination resulted in the identification of PPREs in the promoter/enhancer regions of the AOX and HD genes (Osumi et al., 1991; Tugwood et al., 1992; Zhang et al., 1993; Zhang et al., 1992). There are two DR1 sites upstream of the AOX gene (between -580 and -516 nucleotides upstream of the site of transcription

initiation)(Osumi et al., 1991), however, only the most distal DR1 was found to be bound by PPAR/RXR and to potentiate transactivation in response to peroxisome proliferators (Tugwood et al., 1992). The HD-PPRE contains four tandem TGACCT-related repeats located between nucleotides -2956 to -2919 upstream of the site of transcription initiation of the HD gene (Zhang et al., 1992). This region is sufficient to confer responsiveness to peroxisome proliferators and for binding of PPAR/RXR heterodimers (Zhang et al., 1993). The HD-PPRE is more complex than the AOx-PPRE, as it consists of two DR1 sites separated by 2 nucleotides, yielding an internal DR2 site (Chu et al., 1995a; Zhang et al., 1993; Zhang et al., 1992). This arrangement increases the potential for competitive interactions between nuclear receptors with DR1 and DR2 core site requirements. The 3' region flanking the second TGACCT repeat of the DR1 is important for both the binding of, and transactivation by, PPAR/RXR, with a broad consensus of TGACCTAG(A/T)A being deduced by comparing numerous PPREs (Johnson et al., 1996; Osada et al., 1996). The A/T at position +3 is the most conserved nucleotide among the PPREs and appears to be essential for efficient binding of, and transactivation by, PPAR/RXR heterodimers (Johnson et al., 1996; Osada et al., 1996).

### **1.5 PPARs and the Nuclear Receptor Superfamily**

The nuclear receptor superfamily encompasses a large number of members, which are further classified into subfamilies based on their DNA-binding domains and their homology to other nuclear receptors (Keightley, 1998; Meier, 1997; Moras and Gronemeyer, 1998; Stunnenberg, 1993). The PPARs form a group of nuclear receptors within the vitamin




D3/thyroid hormone/retinoid receptor subfamily. PPARs have been identified in a number of species including human (Schmidt et al., 1992; Sher et al., 1993), rat (Gottlicher et al., 1992), mouse (Chen et al., 1993; Issemann and Green, 1990; Tontonoz et al., 1994b; Zhu et al., 1993), guinea pig (Bell et al., 1998), hamster (Aperlo et al., 1995), salmon (Ruyter et al., 1997) and *Xenopus laevis* (Dreyer et al., 1992). Three PPAR isoforms have been characterized:  $\alpha$ ,  $\beta/\delta$  and  $\gamma$ , and each of these isoforms has a specific tissue distribution and regulates transcription of both distinct and overlapping target genes (Latruffe and Vamecq, 1997; Schoonjans et al., 1996).

#### *1.5.1 General Structure of Nuclear Receptors*

Nuclear receptors contain distinct domains that are functionally interchangeable (Glass, 1994; Laudet et al., 1992; Stunnenberg, 1993). The A/B domain is localized near the amino terminus, shows the most variation among nuclear receptors and is necessary for transactivation (see Fig. 1-2). The C region is highly conserved, contains two zinc-finger motifs required for DNA-binding and also harbors a dimerization interface. The P-box sequence is located at the carboxy-terminal end of the first zinc finger and confers DNA-binding specificity. The P-box is highly conserved and serves as a means to subclassify nuclear receptors based on similarities within this region. Members of the thyroid/retinoid/peroxisome proliferator receptor family all contain an identical P-box motif of CEGCKG. The D region is less conserved and is characterized by a hinge domain. The ligand binding domain is found in the E/F region near the carboxy terminus. Within the ligand binding domain lies the activation function-2 (AF-2) region, which has been shown to be

**Figure 1-2. Nuclear receptor structure and comparison of PPAR isoforms.** The modular structure of members of the nuclear receptor superfamily is illustrated. The location of several common domains are indicated and include the NH<sub>2</sub>-terminal transactivation domain, DNA- and ligand-binding domains, both dimerization interfaces, the hinge region and the activation function-2 (AF-2) domain. PPAR isoforms from mouse (m), rat (r), human (h) and *Xenopus laevis* (x) are shown. The information contained in this figure is derived from papers by Forman et al., 1996, Dreyer et al., 1993, Forman et al., 1995b, Lazar, 1993 and Glass, 1994.



A/B

C

D

E/F

NH<sub>2</sub>

COOH

Transactivation

DNA Binding

Dimerization I  
Hinge Region

Ligand Binding  
Dimerization II

AF-2

essential for interactions between nuclear receptors and nuclear receptor co-regulatory factors. A second nuclear receptor dimerization interface exists within the ligand binding domain. Structural studies of the PPAR $\gamma$  ligand binding domain revealed a large ligand binding cavity, and this may explain how a variety of structurally diverse ligands can activate this receptor (Moras and Gronemeyer, 1998; Nolte et al., 1998).

### *1.5.2 Dimerization Requirements*

Nuclear receptors bind their response elements primarily as homo- or heterodimers; however, some receptors can bind as monomers (Carlberg et al., 1994; Keightley, 1998; Maruyama et al., 1998; Meier, 1997; Moras and Gronemeyer, 1998). PPARs require heterodimerization with the 9-*cis*-retinoic acid receptor (RXR $\alpha$ ) to enable association with PPREs (Gearing et al., 1993; Kliewer et al., 1992a). Nuclear receptors have two putative dimerization interfaces, one within the ligand binding domain and a second interface within the DNA binding domain (Lee et al., 1998; Perlmann et al., 1996). The recently characterized 40 amino acid ligand binding domain dimerization interface contains an 11 amino acid core region that is sufficient for directing dimerization preferences in a yeast two hybrid system (Lee et al., 1998). A two step method for nuclear receptor activation has been postulated, in which interaction between ligand binding domain-dimerization interfaces allows formation of dimers in solution capable of binding a range of DNA elements (Perlmann et al., 1996). This initial interaction is followed by contact between the DNA binding domain-dimerization interfaces which restricts DNA-binding specificity (Perlmann et al., 1996).

### 1.5.3 Orphan Receptors and the Role of Ligand Binding

The largest group of nuclear receptors are those for which no ligand has been identified, or may not exist (Keightley, 1998; Meier, 1997). These are referred to as orphan receptors. Nuclear receptors are activated by ligand binding; however, ligands are not required for binding of nuclear receptors to target DNA response elements (Keightley, 1998; Meier, 1997; Tenbaum and Baniahmad, 1997). The search for nuclear receptor ligands is being performed by numerous groups, some relying on the observation that *in vitro* binding affinity of nuclear receptors towards co-regulatory factors, termed co-activators, greatly increases in the presence of ligand (Desvergne et al., 1998; Krey et al., 1997). The ligand binding requirement for association of the steroid receptor co-activator-1 (SRC-1) with PPAR has allowed the development of a co-activator-dependent receptor ligand assay (Desvergne et al., 1998; Krey et al., 1997; Willson and Wahli, 1997). This assay has identified several natural fatty acids and hypolipidemic drugs as ligands for *Xenopus* PPAR $\alpha$ ,  $\beta/\delta$ , and  $\gamma$ . Additional assays based on conformational changes and using radiolabelled agonists have also been critical in determining PPAR ligands (Forman et al., 1997a; Kliewer et al., 1997). The identification of specific agonists and antagonists will prove fruitful in future experiments in order to determine the genes that are regulated by specific receptors and for characterization of novel co-regulatory interactions.

## 1.6 The Peroxisome Proliferator Activated Receptors

The PPAR family of nuclear receptors contains three isoforms,  $\alpha$ ,  $\gamma$  and  $\delta$ , that each exhibit a distinct pattern of expression. These nuclear receptors have demonstrated roles in

the regulation of lipid metabolism and homeostasis, the control of differentiation, oncogenesis and the progression of apoptosis (Cherkaoui Malki et al., 1990; Chinetti et al., 1998; Latruffe and Vamecq, 1997; Peters et al., 1997a; Tontonoz et al., 1998).

### *1.6.1 PPAR $\alpha$*

PPAR $\alpha$  is found primarily in liver, kidney, heart and brain tissues, and has a critical role in the regulation of genes involved in lipid metabolism and homeostasis (Forman et al., 1996; Issemann and Green, 1990; Latruffe and Vamecq, 1997; Schoonjans et al., 1996). Among the genes regulated by PPAR $\alpha$  are those responsible for peroxisomal  $\beta$ -oxidation: AOx (Osumi et al., 1991; Tugwood et al., 1992) and HD (Zhang et al., 1993; Zhang et al., 1992),  $\omega$ -hydroxylation: cytochrome P450 (CYP) 4A1 and 4A6 (Aldridge et al., 1995; Muerhoff et al., 1992), and fatty acid transport: fatty acid binding protein (Poirier et al., 1997), acyl CoA synthetase (Schoonjans et al., 1995), long chain fatty acid translocase and fatty acid transporter protein (Motojima et al., 1998), as well as apolipoproteins AI and CIII (Peters et al., 1997b; Vu-Dac et al., 1998), carnitine palmitoyl transferase (Mascaro et al., 1998), and mitochondrial HMG-CoA synthase (Rodriguez et al., 1994). In addition, PPAR $\alpha$  has been implicated in the regulation of Cu<sup>2+</sup>, Zn<sup>2+</sup> superoxide dismutase (Inoue et al., 1998), nitric oxide synthase (Colville-Nash et al., 1998), non-peroxisomal transthyretin and  $\alpha$ -2u-globulin (Motojima et al., 1997). Examination of transgenic PPAR $\alpha$ -deficient mice demonstrated prolonged inflammatory responses and increased levels of total serum cholesterol and apolipoprotein AI compared to those found in wild type animals (Peters et al., 1997b). These findings strengthen the proposed roles of PPAR $\alpha$  in lipid homeostasis and in

inflammation. PPAR $\alpha$  has also been linked to apoptosis, since activation of PPAR $\alpha$  by nafenopin prevents apoptosis in hepatocytes through inhibition of bcl-2 activity (Christensen et al., 1998). The suppression of apoptosis is thought to be a contributing factor in the tumorigenic effect of peroxisome proliferators. PPAR $\alpha$  is activated by certain fatty acids, fibrate hypolipidemic drugs and eicosanoids, with the hypolipidemic drug Wy-14,643 and leukotriene B<sub>4</sub> being shown to bind directly to and activate the PPAR $\alpha$  isoform (Forman et al., 1997a; Kliewer et al., 1997; Krey et al., 1997; Schoonjans et al., 1997).

### 1.6.2 PPAR $\gamma$

PPAR $\gamma$  is most abundant in fat tissue and has been named a master regulator of adipogenesis (Fajas et al., 1998; Hu et al., 1995; Tontonoz et al., 1994b; Tontonoz et al., 1994c). PPAR $\gamma$  is transiently expressed in cells of the nervous system during development and may co-operate with PPAR $\delta$  in neural differentiation (Braissant and Wahli, 1998; Cullingford et al., 1998; Granneman et al., 1998). Additionally, PPAR $\gamma$  and PPAR $\alpha$  appear to be involved in the differentiation and apoptotic programs of certain cells of the immune system, with possible roles in atherogenesis and arthritis (Chinetti et al., 1998; Christensen et al., 1998; Elstner et al., 1998; Jiang et al., 1998; Ricote et al., 1998; Tontonoz et al., 1998). Target genes for PPAR $\gamma$  include adipocyte lipid-binding protein 2 (Tontonoz et al., 1994a; Tontonoz et al., 1994b), phosphoenolpyruvate carboxykinase (Tontonoz et al., 1995), uncoupling proteins 1 and 2 (Aubert et al., 1997; Camirand et al., 1998; Digby et al., 1998), and long chain fatty acid transporter protein (Martin et al., 1997). The importance of PPAR $\gamma$  in differentiation first became apparent following the demonstration of Spiegelman and

colleagues that overexpression of PPAR $\gamma$  and CCAAT/enhancer binding protein  $\alpha$  was sufficient to induce the adipogenic program not only in preadipocytes, but also in normally non-adipogenic cells (Hu et al., 1995; Tontonoz et al., 1994c). A number of compounds that activate PPAR $\gamma$  have been discovered and include the thiazolidinediones class of antidiabetic compounds (Krey et al., 1997; Lehmann et al., 1995). It has further been shown that 15-deoxy- $\Delta^{12,14}$ -prostaglandin J<sub>2</sub> and thiazolidinedione BRL49653 are high-affinity ligands that activate PPAR $\gamma$  through direct binding interactions (Forman et al., 1995a; Kliewer et al., 1995; Kliewer et al., 1997; Nolte et al., 1998). Particular attention has focussed on the effects of the thiazolidinediones, since they are insulin sensitizers used in the treatment of type II diabetes (Ibrahimi et al., 1994; Lehmann et al., 1995). Recently, a clinical investigation has linked mutations at Ser114 in human PPAR $\gamma$  with obesity, and it was further demonstrated that this mutation accelerates adipocyte differentiation *in vitro* (Ristow et al., 1998).

### 1.6.3 PPAR $\delta$

Although PPAR $\delta$  has a very broad tissue distribution, little is known about the target genes and specific activators of this PPAR isoform. PPAR $\delta$  is highly expressed in the nervous system, and it appears that it plays a role in oligodendrocyte and astrocyte differentiation (Cullingford et al., 1998; Granneman et al., 1998). PPAR $\delta$  may co-operate with PPAR $\gamma$  in tissue-specific regulation of certain genes, including the uncoupling protein-2 gene (Aubert et al., 1997). PPAR $\delta$  is weakly activated by a number of peroxisome proliferators including Wy-14,643 and the thiazolidinediones BRL49653; however, L-631,033 cinnamic acid has been identified as a potent PPAR $\delta$ -specific activator (Johnson et al., 1997).



### 1.7 Hepatocyte Nuclear Factor-4

Hepatocyte nuclear factor-4 (HNF-4) is an orphan nuclear receptor that is present primarily in liver and kidney, and in lower amounts in brain, spleen and lung. This tissue distribution is similar to that observed for peroxisome proliferator-responsiveness (Dreyer et al., 1992; Issemann and Green, 1990). HNF-4 binds to DR1 response elements as a homodimer, and contains an extended consensus DNA binding site (Jiang et al., 1995; Sladek et al., 1990). The HNF-4 target DNA binding sequence is highly conserved within the AOX-PPRE and less conserved within the HD-PPRE (Osumi et al., 1991; Sladek et al., 1990; Tugwood et al., 1992; Zhang et al., 1992). Interestingly, it has been observed that fatty acyl-CoA thioesters with chain lengths greater than C12 act as ligands for HNF-4, further implicating HNF-4 in fatty acid regulation (Hertz et al., 1998).

Both PPAR and HNF families are key regulators of differentiation and metabolic pathways (Duncan et al., 1998; Latruffe and Vamecq, 1997; Schoonjans et al., 1996). A mutation in HNF-4 has been shown to be responsible for a form of early onset non-insulin dependent diabetes mellitus, indicating the importance of HNF-4 in glucose metabolism (Duncan et al., 1998; Stoffel and Duncan, 1997; Yamagata et al., 1996). Genes which are regulated by HNF-4 include the glycolytic enzymes aldolase B and glyceraldehyde-phosphate dehydrogenase, as well as liver pyruvate kinase and glucose transporter 2 (Gregori et al., 1998; Stoffel and Duncan, 1997). PPAR and HNF-4 co-regulate several genes responsible for fatty acid metabolism and homeostasis including apolipoprotein AI (Jiang et al., 1995; Peters et al., 1997b; Vu-Dac et al., 1998), fatty acid binding protein (Poirier et al., 1997; Stoffel and Duncan, 1997), mitochondrial HMG-CoA synthase (Rodriguez et al., 1994;

Rodriguez et al., 1998), liver transferrin (Hertz et al., 1996b) and apolipoprotein CIII (Fraser et al., 1998; Hertz et al., 1995). These nuclear receptors also affect the transcription of cytochrome P450 (CYP) genes, with HNF-4 modulating expression of CYP 2A4 (Yokomori et al., 1997) and CYP 3A23 (Huss and Kasper, 1998) and PPAR shown to influence transcription of CYP 4A1 and 4A6 (Aldridge et al., 1995; Johnson et al., 1996; Muerhoff et al., 1992). A list of processes which are governed by both HNFs and PPARs is rapidly emerging. In addition to directing the expression of genes involved in lipid homeostasis, these two receptor families have been shown to control genes encoding enzymes responsible for inflammation (Chinetti et al., 1998; Jiang et al., 1998; Peters et al., 1997b; Ricote et al., 1998; Ueda et al., 1998), and overlap in the control of certain viral genes including the pre-C and nucleocapsid promoters of hepatitis B virus (Raney et al., 1997; Yu and Mertz, 1997) and the composite element in the long terminal repeat of HIV-I isolates (Ladías, 1994).

### **1.8 Thyroid Hormone Receptors**

The thyroid hormone receptor (TR) family is a well characterized group of nuclear receptors which are activated 3, 3', 5-triiodothyronine (T3)(Evans, 1988; Lazar, 1991a; Ribeiro et al., 1995a; Shepard and Eberhardt, 1993) The actions of TR are widespread and this nuclear receptor family has been shown to be at the centre of regulating many cellular processes including lipid metabolism, cell growth and differentiation (Chatterjee and Tata, 1992; Desvergne, 1994; Goodridge, 1987; Silva and Rabelo, 1997; Xiong et al., 1998). The wide ranging effects of TR family members are due in part to the broad response element preference exhibited by these receptors. Thyroid hormone response elements may contain

direct, palindromic and inverted palindromic repeats with a range of spacing (Desvergne, 1994; Ribeiro et al., 1998; Ribeiro et al., 1995b; Tsai and O'Malley, 1994); however, the nucleotides flanking the TGACCT-related sites appear to influence TR binding (Olson and Koenig, 1997). In addition, TRs can function as monomers, homodimers and as heterodimers with RXR $\alpha$  (Desvergne, 1994; Ikeda et al., 1994; Lazar et al., 1991b; Tsai and O'Malley, 1994).

The TR family has been the focus of much of the pioneering work which has established general mechanisms of nuclear receptor function. In particular, the ability of TRs to function as transcriptional silencers has driven much of the early research and discovery of nuclear receptor accessory factors, and is discussed in more detail in a subsequent section (Baniahmad et al., 1992; Baniahmad et al., 1995; Damm et al., 1989; Graupner et al., 1989; Renkawitz, 1993; Sap et al., 1989). A member of the TR family was a natural choice as a candidate regulator for peroxisome proliferator responsive genes, based on overlapping tissue distributions and the observations that transcription of several thyroid hormone-responsive genes were shown to be modulated by PPAR $\alpha$  (Bogazzi et al., 1994; Castelein et al., 1994; Jow and Mukherjee, 1995). Genes which are co-regulated by PPAR $\alpha$  and TR $\alpha$  include those encoding malic enzyme, CYP 4A family proteins and uncoupling proteins (Aldridge et al., 1995; Dogra et al., 1998; IJpenberg et al., 1997; Johnson et al., 1996; Muerhoff et al., 1992; Silva and Rabelo, 1997; Webb et al., 1996).

## **1.9 Retinoid-like Z Receptor/Retinoid Orphan Receptors**

The retinoid-like Z receptor/retinoid orphan receptor (RZR/ROR) family consists of

three isoforms:  $\alpha$ ,  $\beta$ , and  $\gamma$ , which demonstrate broad, brain and skeletal muscle tissue distributions, respectively. The  $\alpha$  isoform includes four splice variants (ROR $\alpha$ 1, ROR $\alpha$ 2, ROR $\alpha$ 3 and RZR $\alpha$ ) (Becker-Andre et al., 1993; Carlberg et al., 1994; Giguere et al., 1994; Hirose et al., 1994). It has been reported that RZR $\alpha$ s are activated directly by melatonin, however this finding has been subject to discussion (Becker-Andre et al., 1994; Carlberg and Wiesenberg, 1995; Hazlerigg et al., 1996; Rafii-El-Idrissi et al., 1998; Steinhilber et al., 1995). Recently, it has also been reported that several thiazolidinedione-like agents function as RZR ligands (Missbach et al., 1996; Wiesenberg et al., 1998). Members of this family of nuclear receptors bind as monomers to target response elements and have been shown to regulate genes involved in inflammation, differentiation and lipid metabolism (Carlberg et al., 1994; Giguere et al., 1994; Hirose et al., 1994; Schrader et al., 1996; Vu-Dac et al., 1997). The  $\alpha$  and  $\gamma$  isoforms are induced early in adipogenesis suggesting involvement in pre-adipocyte differentiation and possible interplay with the adipogenic regulator, PPAR $\gamma$  (Austin et al., 1998; Spiegelman and Flier, 1996). Monomeric receptors from the NGFI-B and RZR/ROR families contain a carboxy-terminal extension of the zinc finger in their DNA binding domains (Wilson et al., 1992). This extended region is also found in the DNA-binding domain of PPAR, and is required for the recognition of extended 3' sequences that flank TGACCT-like core sites (Hsu et al., 1998; Wilson et al., 1992). The RZR consensus flanking sequence is similar to those which border a number of nuclear receptor response elements including several PPREs (Schrader et al., 1996). The HD-PPRE was identified as a possible RZR regulatory site in a large scale screen for candidate RZR response elements (Schrader et al., 1996).

### 1.10 Calreticulin

Calreticulin is a  $\text{Ca}^{2+}$ -binding protein which exhibits strong endoplasmic reticulum localization and is involved in a variety of cellular processes (Coppolino and Dedhar, 1998; Krause and Michalak, 1997). Calreticulin shares homology with the endoplasmic reticulum chaperone protein calnexin, and has demonstrated an important role as a chaperone involved in the proper folding and assembly of proteins within the endoplasmic reticulum (Vassilakos et al., 1998). Calreticulin can replace calnexin in the processing of mouse class I histocompatibility molecules (Krause and Michalak, 1997; Vassilakos et al., 1998). Calreticulin readily binds  $\text{Ca}^{2+}$  and is thought to serve as a key factor in the storage and regulation of intracellular  $\text{Ca}^{2+}$  (Mery et al., 1996).  $\text{Ca}^{2+}$  is an extremely important cell signalling molecule and as a consequence calreticulin may be a central player governing signalling events, both within the endoplasmic reticulum lumen and at other locations within the cell. It has been demonstrated that  $\text{Ca}^{2+}$ -binding by calreticulin leads to direct interaction with, and sequestration of, another endoplasmic reticulum luminal protein, protein disulfide isomerase, suggesting a regulatory role for calreticulin via  $\text{Ca}^{2+}$ -dependent protein-protein interactions (Baksh et al., 1995). Calreticulin has also been implicated in the regulation of cell adhesion to extracellular substrata (Leung-Hagesteijn et al., 1994; Opas et al., 1996). Although calreticulin can interact *in vitro* with KLGFFKR motifs in the cytoplasmic tail of  $\alpha$ -integrin, it appears that calreticulin is affecting adhesion indirectly by altering the level of vinculin (Opas et al., 1996; Rojiani et al., 1991). A role for calreticulin in regulation of gene expression has also been suggested since, *in vitro*, calreticulin binds to KXXFF(K/R)R motifs localized in the DNA-binding domains of nuclear receptors. Calreticulin interacts with these

motifs in the DNA-binding domains of glucocorticoid, vitamin D3, retinoic acid and androgen receptors *in vitro*, and inhibits transactivation by these receptors in transfection experiments (Burns et al., 1994; Dedhar et al., 1994; Desai et al., 1996; Michalak et al., 1996; St-Arnaud et al., 1995; Wheeler et al., 1995). Consequently it was proposed that calreticulin is a general regulator of nuclear receptor activity through physical interactions with the conserved KXXFF(K/R)R motif located in the DNA binding domains of nuclear receptors (Burns et al., 1994; Dedhar et al., 1994; Shago et al., 1997).

### **1.11 Nuclear Receptor Co-Regulatory Factors**

In the past three years the intense interest in nuclear receptors and the molecular means by which they influence transcription has resulted in the discovery of a novel class of nuclear receptor accessory factors (Horwitz et al., 1996; Nolte et al., 1998; Torchia et al., 1998). These co-regulatory proteins serve to direct the potentiation or inhibition of transcription through physical interactions with nuclear receptors, and are referred to as co-activators and co-repressors, respectively. By biochemical and structural examinations, a common mechanism of nuclear receptor regulation is emerging (Chen and Li, 1998; Lavinsky et al., 1998; Nolte et al., 1998; Westin et al., 1998). It appears that ligand binding causes a structural rearrangement in the nuclear receptor which results in the dissociation of a co-repressor complex and facilitates direct interaction between ligand-bound receptors and a co-activator complex. In addition to interacting with nuclear receptors, co-repressors and co-activators are directly involved in chromatin remodelling.

### *1.11.1 The Activation Function Region*

Nuclear receptors contain a discrete ligand binding domain within their carboxy-terminus. Through mutational studies with the estrogen receptor (ER), a region within the ligand binding domain was found to be absolutely necessary for transactivation (Danielian et al., 1992). This region was named the activation function-2 (AF-2) domain. Insights into structural alterations within the ligand binding domain following ligand association were obtained by performing X-ray crystallography of a number of nuclear receptors in the unbound, agonist- or antagonist-bound state (Bourguet et al., 1995; Brzozowski et al., 1997; Nolte et al., 1998; Renaud et al., 1995; Wagner et al., 1995). A high degree of structural conservation in the ligand binding domains of the nuclear receptors was observed. These regions consist of 12 alpha helices with a conserved beta-turn between helices 5 and 6 (Moras and Gronemeyer, 1998; Nolte et al., 1998; Torchia et al., 1998). Interestingly, the AF-2-containing helix 12 undergoes dramatic alterations in response to agonist or antagonist binding. In comparisons of the ligand binding domains of RAR, RXR, ER, PPAR $\gamma$  and TR, agonist binding appears to cause helix 12 to adopt a conformation in which it directly interacts with the ligand, suggesting that AF-2 switches between different conformations in the presence or absence of ligand (Bourguet et al., 1995; Brzozowski et al., 1997; Nolte et al., 1998; Renaud et al., 1995; Wagner et al., 1995). Binding of the ER antagonist, raloxifene, causes helix 12 to become partially buried and obscures residues known to be necessary for transcriptional activation (Brzozowski et al., 1997). PPAR $\alpha$  undergoes a conformational change following exposure to agonists, based on protease sensitivity assays (Dowell et al., 1997b). Structural data from studies with PPAR $\gamma$  and its ligand BRL49653 demonstrate the

importance of helix 12 structural rearrangements to facilitate interactions with the SRC-1 and transcriptional intermediary factor 2 (TIF2) co-activators (Leers et al., 1998; Nolte et al., 1998; Westin et al., 1998). Other aspects of nuclear receptor ligand binding domains are also modified following association with ligand, suggesting that agonists and antagonists directly alter the surface structure of nuclear receptor ligand binding domains (Bourguet et al., 1995; Moras and Gronemeyer, 1998; Renaud et al., 1995; Torchia et al., 1998; Wagner et al., 1995).

#### *1.11.2 Nuclear Receptor Co-Activators*

A number of co-activator proteins have been identified which mediate transcriptional activation in a ligand- and AF-2-dependent manner. Co-activators were primarily isolated by screens for factors that interacted with ligand binding domains, or with epitope-tagged or ligand-bound nuclear receptors (Glass et al., 1997; Hong et al., 1996; Monden et al., 1997; Zhu et al., 1997). These studies showed the involvement of accessory factors in transcriptional activation by nuclear receptors. A number of related nuclear receptor co-activators have been isolated and include glucocorticoid receptor-interacting protein 1 (GRIP1) (Hong et al., 1996), activator of the thyroid and retinoic acid receptor (ACTR) (Chen et al., 1997), receptor-associated co-activator 3 (RAC3) (Li et al., 1997a), p300/CBP co-integrator associate protein (p/CIP) (Torchia et al., 1997), and TR-activator molecule (TRAM) (Takeshita et al., 1997). The co-activators SRC-1 (DiRenzo et al., 1997), receptor-interacting protein 140 (RIP140) (Treuter et al., 1998), PPAR $\gamma$ -binding protein (PBP) (Zhu et al., 1997), TIF2 (Leers et al., 1998) and PPAR $\gamma$ -co-activator (PGC) (Puigserver et al.,



1998) have been observed to interact with PPARs. CREB binding protein (CBP) and the functionally related p300 exhibit a more general co-activator function for both nuclear receptors and other transcription factors such as cAMP response element-binding protein (CREB), activator protein-1, and signal transducer and activator of transcription (STAT) (Arany et al., 1994; Chakravarti et al., 1996; Horvai et al., 1997; Kamei et al., 1996). CBP/p300 not only interacts with transcription factors, but also possesses intrinsic histone acetyl transferase activity (Bannister and Kouzarides, 1996; Ogryzko et al., 1996). Chromatin remodelling through the acetylation of histones is thought to make DNA elements more accessible to transcriptional machinery and thereby increase transcription (Glass et al., 1997; Grunstein, 1997; Torchia et al., 1998; Wade et al., 1997; Wolffe, 1997; Wolffe, 1998). CBP/p300 interacts strongly with nuclear receptors, including PPAR $\gamma$ , in a ligand-dependent manner (Dowell et al., 1997a). The necessity of cooperation between nuclear receptors and CBP/p300 is evident since the majority of identified co-activators contain both nuclear receptor and CBP/p300-interaction domains (Ding et al., 1998; Schmidt et al., 1998; Takeshita et al., 1998; Voegel et al., 1998; Yao et al., 1996). Another factor which contains histone acetyl transferase activity, p300/CBP-associated factor (p/CAF), interacts with CBP/p300 and is a component of the proposed co-activator complex (Bannister and Kouzarides, 1996; Chen et al., 1997; Yang et al., 1996). It was recently observed that SRC-1 also functions as a histone acetyl transferase (Spencer et al., 1997). A common LXXLL motif (where L denotes leucine and X represents any amino acid) is present in co-activators and has been shown to be necessary for nuclear receptor/co-activator interactions (Ding et al., 1998; Heery et al., 1997; McInerney et al., 1998; Torchia et al., 1997; Voegel et al., 1998). It

appears that the LXXLL motifs are important for stabilizing interactions between ligand bound nuclear receptors and co-activators (Moras and Gronemeyer, 1998; Torchia et al., 1998; Westin et al., 1998). Addition of 25 amino acid fragments containing the LXXLL motif are sufficient to eliminate interactions between ligand-bound nuclear receptors and co-activators *in vitro*, and mutations in this motif eliminate co-activator mediated transcriptional activation by nuclear receptors in functional assays (Heery et al., 1997; Torchia et al., 1997; Westin et al., 1998). The co-activator complex is thought to consist of ligand bound nuclear receptors, CBP/p300, p/CAF, co-activators and as yet unidentified factors (Chen et al., 1997; Chen and Li, 1998; Glass et al., 1997; Westin et al., 1998). The association of ligand bound nuclear receptors with DNA results in structural changes that are thought to specify interactions with particular co-activators (Glass et al., 1997; Leers et al., 1998; Nolte et al., 1998; Torchia et al., 1998; Westin et al., 1998). Co-activator complexes may serve to increase transcription through several means including chromatin remodelling via histone acetylation, stabilization of nuclear receptor binding and facilitation of interactions with basal transcriptional machinery (Chen and Li, 1998; Glass et al., 1997; Horwitz et al., 1996; Torchia et al., 1998).

### *1.11.3 Nuclear Receptor Co-Repressors*

RAR and TR act as transcriptional silencers for many genes in the absence of ligand (Baniahmad et al., 1992; Burcin et al., 1994; Damm et al., 1989; Graupner et al., 1989; Nawaz et al., 1995; Renkawitz, 1993; Sap et al., 1989). Closer examination of this observation resulted in the discovery of silencing units within the RAR and TR ligand binding

domains, and of co-repressors which mediate the silencing effects of TR and RAR (Chen and Evans, 1995; Kurokawa et al., 1995; Sande and Privalsky, 1996). The silencing activity of TR and RAR has been localized to a site within the ligand binding domain, but distinct from the AF-2 region, termed the CoR box (Baniahmad et al., 1992; Baniahmad et al., 1995; Zamir et al., 1996). Co-repressors inhibit transcription of target genes by several nuclear receptors in the unbound or antagonist-bound states (Chen and Evans, 1995; Jackson et al., 1997; Lee et al., 1995; Shibata et al., 1997; Smith et al., 1997). The nuclear receptor co-repressor (NCoR) and silencing mediator for retinoid and thyroid receptors (SMRT) interact with a range of nuclear receptors (Chen and Evans, 1995; Heinzl et al., 1997; Lavinsky et al., 1998; Lee et al., 1995; Zamir et al., 1996; Zamir et al., 1997). Although NCoR and SMRT share homology, they show different preferences for specific nuclear receptors (Lavinsky et al., 1998). SMRT has been shown to repress transactivation by PPARs, whereas NCoR does not appear to interact with these receptors (Lavinsky et al., 1998; Yan et al., 1998; Zamir et al., 1997). Like co-activators, co-repressors also form complexes and these contain unliganded nuclear receptors, the co-repressor and factors responsible for chromatin remodelling (Heinzl et al., 1997; Laherty et al., 1997; Li et al., 1997b; Nagy et al., 1997; Torchia et al., 1998; Zhang et al., 1997). Recent investigation of core co-repressor complexes demonstrated that histone deacetylases and other components including Sin3 were present in the complex, and histone deacetylase activity was found to be required for re-establishment of the repressed chromatin state (Alland et al., 1997; Hassig et al., 1997; Heinzl et al., 1997; Laherty et al., 1998; Laherty et al., 1997; Nagy et al., 1997; Torchia et al., 1998; Zhang et al., 1997).

#### *1.11.4 Model of Co-Activator/Co-Repressor Interactions*

The current model of co-activator/co-repressor function proposes that DNA-bound nuclear receptors associate with a co-repressor complex, yielding histone deacetylation and transcriptional repression (see Fig. 1-3) (Chen and Li, 1998; Torchia et al., 1998). Upon ligand binding, the co-repressor complex is dissociated, allowing interaction of the ligand-bound nuclear receptor with a co-activator complex. This association enables histone acetylation and chromatin remodelling, revealing crucial DNA sites for transcriptional activation. In addition to being regulated by the availability and active state of nuclear receptors, it appears that additional signalling networks such as the mitogen-activated protein kinase and protein kinase A cascades, may affect the function of co-activator and co-repressor complexes (Lavinsky et al., 1998; Torchia et al., 1998).

#### **1.12 Nuclear Receptor Interplay**

Interactions between nuclear receptors allow the integration of signals from diverse signalling pathways to direct transcriptional regulation of target genes. In experiments that directly compared the DNA element requirements of several nuclear receptors that recognize DR1 sites (including HNF-4, PPAR $\alpha$  and RXR $\alpha$ ), it was observed that the important factors influencing DNA binding affinity are determinants within the core sequence, the identity of the spacing nucleotide and the sequence flanking the core TGACCT-related repeats (Jiang et al., 1995; Juge-Aubry et al., 1997; Nakshatri and Bhat-Nakshatri, 1998; Nishiyama et al., 1998). It was also observed that DNA binding affinity correlates with transactivation potential from a given element (Nakshatri and Bhat-Nakshatri, 1998).

**Figure 1-3. Model of nuclear receptor accessory factor interactions.** In the absence of ligand or presence of antagonist, a co-repressor/nuclear receptor complex forms and maintains chromatin in the deacetylated (repressed) state. Binding of ligand (L) by the nuclear receptor results in a conformational change which enables dissociation of the co-repressor complex and assembly of a co-activator/nuclear receptor complex. The co-activator complex contains histone acetyl-transferase activity which relieves chromatin repression and makes sites required for transcriptional initiation available, yielding transcriptional activation.

Co-Repressor Complex Interactions

Co-Activator Complex Interactions




Figure 1-3. Model of nuclear receptor co-regulatory factor interactions.

The coordinate transactivation of target genes by nuclear receptors is governed by several levels of specificity in addition to those inherent in the DNA binding/response element requirements. Nuclear receptor interplay can be dictated by dimerization requirements, ligand binding, and by competition for common binding sites and for co-activators or co-repressors. Nuclear receptors typically bind to target response elements as homo- or heterodimers, however some receptors such as RZR, NGFB-I and TR can bind to DNA as monomers (Carlberg et al., 1994; Desvergne, 1994; Giguere et al., 1994; Ikeda et al., 1994; Lazar et al., 1991b; Maruyama et al., 1998). RXR $\alpha$  is a common heterodimerization partner for many nuclear receptors including PPAR (Kliwer et al., 1992b), RAR, TR (Zhang et al., 1992), and vitamin D3 receptor (VDR) (Thompson et al., 1998), and therefore the availability of RXR $\alpha$  can serve to alter the transactivation capacity of specific nuclear receptors due to competition for this heterodimerization partner (Chu et al., 1995b; DiRenzo et al., 1997; Juge-Aubry et al., 1995; Miyata et al., 1996).

The classical model of nuclear receptor activation describes ligand binding as a requirement for receptor activation, however ligand binding is not necessary for binding of nuclear receptors to their target response elements (Carson-Jurica et al., 1990; Evans, 1988). Ligand binding by nuclear receptors results in a conformational change in the AF2 domain which facilitates interaction with co-activators and is thought to displace associated co-repressors (Bourguet et al., 1995; Brzozowski et al., 1997; Moras and Gronemeyer, 1998; Nolte et al., 1998; Renaud et al., 1995; Wagner et al., 1995). It has been reported that ligand association also increases DNA binding of PPAR $\alpha$ /RXR $\alpha$  heterodimers (Forman et al., 1997a). In the absence of ligand some nuclear receptors, such as TR, can repress

transcription. Therefore the availability of ligand critically affects transcriptional activation from target response elements.

Several different nuclear receptors can compete for binding to the same response element depending on their inherent promiscuity. COUP-TF is an orphan receptor that recognizes a wide variety of response elements and is a constitutive repressor of transcription. COUP-TF has been shown to compete for binding to response elements for PPAR, TR, RXR, RAR and VDR, and represses transactivation in part, through competition for shared response elements (Cooney et al., 1993; Cooney et al., 1992; Marcus et al., 1996; Miyata et al., 1993).

In recent years the discovery of accessory factors termed co-activators and co-repressors has added to our understanding of transcriptional regulation by nuclear receptors (Chen and Li, 1998; Glass et al., 1997; Horwitz et al., 1996; Nolte et al., 1998; Torchia et al., 1998; Westin et al., 1998). Nuclear receptors are dependent on interactions with accessory factors to potentiate or repress transactivation of target genes. In terms of nuclear receptor interplay, competition between nuclear receptors for accessory factors can affect the activity of nuclear receptors and thereby direct the activation or repression of a target response element.

### **1.13 Aims of This Project**

In order to better understand the mechanisms involved in the regulation of the AOX and HD genes in response to peroxisome proliferators, we have examined the interplay between several key nuclear receptors in modulating gene expression. These studies have



investigated the interactions of PPAR $\alpha$ , RXR $\alpha$ , HNF-4, TR $\alpha$  and RZR $\alpha$  in terms of recognition and binding of the AOX- and HD-PPREs and have been extended to examine the influences that these receptors exert in the transactivation of PPREs in a cellular context. I have also analyzed the role of the Ca<sup>2+</sup>-binding protein, calreticulin, in the modulation of peroxisome proliferator-directed gene expression. The results of my investigation have enabled the formation of a more complete model of peroxisome proliferator-directed gene expression and have provided additional insights into the general mechanisms of nuclear receptor signalling.

## CHAPTER 2

### **Transactivation of the Peroxisome Proliferator-Activated Receptor is Differentially Modulated by Hepatocyte Nuclear Factor-4<sup>†</sup>**

<sup>†</sup>A version of this chapter has been published. Winrow, C.J., Marcus S.L., Miyata, K.S., Zhang, B., Capone, J.P. and Rachubinski, R.A. 1994. *Gene Express.* 4:53-62. Used with permission from Cognizant Communications Corporation.

## 2.1 ABSTRACT

Peroxisome proliferator-activated receptors (PPARs) stimulate the expression of several genes involved in lipid metabolism by binding to specific *cis*-acting peroxisome proliferator-responsive elements (PPREs) via cooperativity with retinoid X receptors. We demonstrate here that hepatocyte nuclear factor-4 (HNF-4), another member of the nuclear hormone receptor superfamily, bound with differing affinities to the PPREs from the genes encoding rat acyl-CoA oxidase and hydratase-dehydrogenase, the first two enzymes of the peroxisomal  $\beta$ -oxidation pathway. In cotransfection assays, HNF-4 repressed rat PPAR-dependent activation of a reporter gene linked to the acyl-CoA oxidase PPRE, either in the absence or presence of the peroxisome proliferator, Wy-14,643. Rat PPAR-dependent activation of a reporter gene linked to the hydratase-dehydrogenase PPRE was less efficiently repressed by HNF-4 in the absence of Wy-14,643 than was activation from the acyl-CoA oxidase PPRE. However, in the presence of Wy-14,643, HNF-4 functioned cooperatively with PPAR to significantly enhance induction from the hydratase-dehydrogenase PPRE. These results suggest that the genes encoding the first two enzymes of the peroxisomal  $\beta$ -oxidation pathway are subject to differential regulation by the interplay of multiple members of the steroid/nuclear hormone receptor superfamily, mitigated in part by the structures of the PPREs and by the presence of activators of PPARs.

## 2.2 INTRODUCTION

Peroxisome proliferators constitute a large group of xenobiotic chemicals that include the fibrate family of hypolipidemic drugs, herbicides and phthalate ester plasticizers. Exposure to peroxisome proliferators increases both the number and metabolic capacity of peroxisomes (Lock et al., 1989). The mechanism of action of peroxisome proliferators is of considerable interest, since these agents have been shown to induce hepatomegaly, chromosomal aberrations and ultimately hepatocarcinogenesis in rodents (Rao and Reddy, 1991; Reddy, 1990; Reddy et al., 1980). Peroxisome proliferators do not mutate DNA directly and are therefore classified as non-genotoxic carcinogens.

The pleiotropic cellular responses to peroxisome proliferators are mediated in part by the transcriptional induction of a number of genes whose products are involved in lipid metabolism. These include the rat acyl-CoA oxidase (AOx) and hydratase-dehydrogenase (HD) genes, which encode the first two enzymes of the peroxisomal fatty acid  $\beta$ -oxidation system (Reddy et al., 1986a); the CYP4A6 gene, which encodes a member of the cytochrome P450 fatty acid  $\omega$ -hydroxylase family (Muerhoff et al., 1992); and the gene encoding the liver fatty acid-binding protein (Besnard et al., 1993; Issemann et al., 1992; Kaikaus et al., 1993). Transactivation of these peroxisome proliferator-responsive genes is mediated by peroxisome proliferator-activated receptors (Issemann and Green, 1990). PPARs are members of the steroid/thyroid hormone receptor superfamily that bind to specific *cis*-acting peroxisome proliferator-responsive elements (PPREs), resulting in an overall stimulation of transcription of responsive genes (Marcus et al., 1993; Osumi et al., 1991; Tugwood et al., 1992; Zhang et al., 1992). Human (Schmidt et al., 1992), rat (Gottlicher et al., 1992), mouse (Chen et al.,

1993; Issemann and Green, 1990; Zhu et al., 1993), and *Xenopus* (Dreyer et al., 1992) have been shown to harbor multiple PPAR-related genes, perhaps reflecting a requirement for different PPARs in regulating specific target genes or in mediating responsiveness to a variety of stimuli. PPARs can be activated by a wide spectrum of structurally diverse peroxisome proliferators, as well as by both synthetic and natural fatty acids (Dreyer et al., 1992; Keller et al., 1993). However, none of these agents binds directly to PPARs, and therefore the true ligands for these receptors remain to be identified.

The natural PPREs characterized so far consist of direct repeats of the core half-site motif TGACCT, which is also found in the cognate response elements of other nuclear hormone receptors of this class including the thyroid, retinoic acid, vitamin D3 and 9-*cis*-retinoic acid (RXR) receptors (Umesono et al., 1991). Target specificity is determined in part by the sequence of the half-sites, the number and relative spacing of their direct repeats, and the ability of some receptors to bind to cognate response elements as homodimers and/or heterodimers (Lucas and Granner, 1992; Mader et al., 1993). However, it is not uncommon for different nuclear hormone receptors to compete for binding to a particular DNA-response element. The potential for promiscuous binding, combinatorial interactions, and cross-talk among receptors serves to modulate transcription of hormone-responsive genes in multiple ways, thereby contributing to both the complexity and the diversity in signalling pathways. Indeed, we and others have demonstrated that PPARs bind to PPREs as heterodimers with 9-*cis*-retinoic acid receptors, implying that the peroxisome proliferator and retinoid signalling pathways converge (Gearing et al., 1993; Kliewer et al., 1992b; Marcus et al., 1993). Recently, we have shown that COUP-TF, an orphan member of the nuclear hormone receptor

superfamily, binds to the HD-PPRE and can antagonize rat (r) PPAR-mediated transactivation from this PPRE *in vivo* (Miyata et al., 1993). Therefore, PPAR function is dependent upon interactions with, and can be subject to modulation by, other members of the nuclear hormone receptor superfamily.

Hepatocyte nuclear factor-4 (HNF-4) is another orphan member of the nuclear hormone receptor family that binds to TGACCT-related elements (Sladek et al., 1990). HNF-4 is a liver-enriched factor that plays an important role in the transcriptional regulation of several genes whose products are involved in diverse metabolic pathways. These include the genes encoding apoAI, apoAII, apoB and apoCIII (Ladiaz et al., 1992; Mietus-Snyder et al., 1992),  $\alpha_1$ -antitrypsin and transthyretin (Costa et al., 1989; Costa et al., 1988), ornithine transcarbamylase (Nishiyori et al., 1994), and some members of the cytochrome P450 hydroxylase superfamily (Chen et al., 1994). HNF-4 mRNA is most abundant in liver and kidney and low in brain, spleen, and lung, a pattern of expression closely matching peroxisome proliferator tissue specificities and PPAR abundance (Dreyer et al., 1992; Issemann and Green, 1990). Since peroxisomes are responsible for the metabolism of long-chain fatty acids, and the PPREs of both the AOX and HD genes share significant homology with the consensus HNF-4 target site, we explored the possibility that HNF-4 might play a role in the regulation of these genes. We demonstrate here that HNF-4 present in extracts of peroxisome proliferator-responsive rat hepatoma cells or synthesized *in vitro* bound strongly to the AOX-PPRE and with lower affinity to the HD-PPRE. HNF-4 on its own had no effect on the transcription of reporter genes linked to the AOX- or HD-PPRE; however, it repressed both the peroxisome proliferator-independent and peroxisome proliferator-dependent transactivation from the AOX-PPRE by PPAR $\alpha$ . Surprisingly, while PPAR $\alpha$ -dependent

transactivation of a HD-PPRE-linked reporter gene was also repressed by HNF-4, its activity was potentiated in the presence of the peroxisome proliferator Wy-14,643. Our results suggest that HNF-4 plays an important role in the regulation of genes encoding enzymes of the peroxisomal  $\beta$ -oxidation pathway by differentially modulating transactivation by PPARs.

## 2.3 MATERIALS AND METHODS

### 2.3.1 Plasmids

The luciferase reporter plasmids pCPS*luc*, pHD(X3)*luc* and pAOX(X2)*luc* and the effector plasmids expressing the full-length cDNAs for rat PPAR and human RXR $\alpha$  have been described previously (Marcus et al., 1993; Zhang et al., 1993). pCPS*luc* is a luciferase reporter plasmid containing the minimal promoter (a 600 nucleotides region from upstream of the transcription start site) for the carbamoyl-phosphate synthetase gene. The pHD(X3)*luc* construct was made by inserting three tandem copies of the sequence 5'-gatCCTCTCCTTTGACCTATTGAACTATTACCTACATTTGA and its complement 5'-gatcTCAAATGTAGGTAATAGTTCAATAGGTCAAAGGAGAG corresponding to -2956 to -2919 of the HD gene into the unique *Bam*HI site of pCPS*luc*. The pAOx(X2)*luc* reporter plasmid contains two tandem copies of the sequence 5'-gatCCTTTCCCGAACGTGACCTT-TGTCCTGGTCCCCTTTTGCT and its complement 5'-gatctAGCAAAGGGGACC-AGGACAAAGGTCACGTTTCGGGAAG which encompasses the AOx-PPRE between -583 and -544 of the rat AOx gene. Nucleotides shown in lower case at the ends of the oligonucleotides were added to provide cohesive *Bam*HI-*Bgl*III ends at the 5' and 3' termini,

respectively. The rat PPAR cDNA in pBluescript II SK(+)(Stratagene) was provided by D. Noonan (Ligand Pharmaceuticals, San Diego) and was excised as a 2.6 Kbp *SpeI/EcoRV* fragment and inserted between the *XbaI/EcoRI* sites of pRC/CMV (Invitrogen). The human RXR $\alpha$  construct pSKXR3-1 was provided by R. Evans (The Salk Institute, La Jolla) and a 1.8 Kbp *EcoRI* fragment containing the cDNA was removed and inserted into the *EcoRI* site of the eukaryotic expression vector pSG5 (Stratagene). The  $\beta$ -galactosidase expression vector pCH110 (Amersham-Pharmacia Biotech) was included in transfections as an internal control for transfection efficiency. The plasmid for expression of rat HNF-4 was kindly provided by F.M. Sladek (University of California, Riverside) and contains the HNF-4 cDNA within a 3.0 Kbp fragment cloned into the *Bam*HI site of pSG5 (Sladek et al., 1990).

### 2.3.2 *Transfections and Measurement of Luciferase Activity*

All transfections were carried out in African green monkey kidney BSC40 monolayer cells using a modification of the calcium phosphate method (Ausubel, 1992; Zhang et al., 1993). Cells at 60-80% confluence were maintained in phenol-red free DMEM medium supplemented with 5% charcoal-stripped fetal bovine serum, 1% penicillin/streptomycin (Gibco BRL) and 4% L-glutamine (Gibco BRL), for 24 hours prior to and during transfection. Equal volumes of a 0.125 mM calcium chloride/DNA solution and a 2X HEPES-buffered saline solution (280 mM NaCl, 1.5 mM Na<sub>2</sub>HPO<sub>4</sub>·7H<sub>2</sub>O, 50 mM HEPES (pH 7.12)) were combined and incubated at room temperature for 30 minutes. During this time, fresh transfection media containing either 0.1 mM of the peroxisome proliferator Wy-14,643 (in dimethyl-sulfoxide) or an equivalent volume of dimethyl-sulfoxide was added to the cells.



The DNA/calcium phosphate co-precipitate was divided into 1 mL aliquots and added dropwise to 10 cm plates. Transfections contained 5  $\mu$ g of either the pHD(X3)*luc* or pAOx(X2)*luc* reporter gene construct, and where indicated, 2  $\mu$ g of PPAR $\alpha$ /RC/CMV, 2  $\mu$ g of RXR $\alpha$ /SG5 and varying amounts of rHNF-4/SG5. The  $\beta$ -galactosidase expression vector (2  $\mu$ g) was included in transfections and effector plasmid dosage was kept constant by addition of appropriate amounts of the corresponding empty vectors. Total DNA was kept at 20  $\mu$ g per plate by addition of sonicated salmon sperm DNA (Sigma). Fresh transfection media containing Wy-14,643 or dimethyl-sulfoxide was added 16 hours post-transfection, and cells were incubated for an additional 24 hours. Cell harvesting, lysate preparation and quantification of luciferase activity were carried out as described previously (Marcus et al., 1993; Miyata et al., 1993; Zhang et al., 1993). Briefly, the cells were washed with phosphate-buffered saline, scraped from plates in 1 mL of phosphate-buffered saline and pelleted by centrifugation for 30 seconds. The supernatant was aspirated and cells were resuspended in 100  $\mu$ L of 1X lysis buffer (2 mM CDTA, 1% (w/v) Triton X-100, 2 mM dithiothreitol, 10% (w/v) glycerol and 25 mM Tris-phosphate (pH7.8)). The lysates were prepared by intermittent vortexing for 10 minutes followed by brief centrifugation to remove cell debris. Aliquots of the supernatants were measured for luciferase activity by addition of reconstituted luciferase assay buffer and quantitation with a luminometer (Bio-Orbit, model 1523). The reconstituted luciferase assay buffer contains 0.53 mM ATP, 0.47 mM luciferin (Sigma), and 0.27 mM co-enzyme A (lithium salt, Sigma) in 10 mM dithiothreitol/25 mM glycylglycine (pH 7.8) added to basic luciferase assay buffer (20 mM Tricine, 1.07 mM (MgCO<sub>3</sub>)<sub>4</sub>Mg(OH)<sub>2</sub>•5H<sub>2</sub>O, 2.67 mM MgSO<sub>4</sub>, 0.1 mM EDTA, 33.3 mM dithiothreitol (pH

7.8)).

### 2.3.3 *In Vitro Transcription/Translation*

Transcription and translation of cDNAs encoding PPAR $\alpha$ , RXR $\alpha$  and HNF-4 were performed using the commercially available TNT T7 coupled rabbit reticulocyte lysate system according to manufacturer's protocol (Promega). Translation products were labelled with L-[ $^{35}$ S]methionine and were analyzed on 15% SDS-polyacrylamide gels. Synthesis of proteins used in electrophoretic mobility shift analysis (EMSA) was carried out in parallel with unlabelled methionine.

### 2.3.4 *Electrophoretic Mobility Shift Analysis*

H4IIEC3 nuclear extracts were prepared following Dignam's method (Dignam et al., 1983), except that following cell harvesting, cells were incubated on ice in buffer A for 10 minutes and subsequently lysed in buffer A supplemented with 0.5% NP-40. The nuclei were then collected by centrifugation at 2200 xg, and extracts were processed according to Dignam's method (Dignam et al., 1983). EMSA was performed with oligonucleotide probes corresponding to the HD-PPRE (5'-CCTCTCCTTTGACCTATTGAACTATTACCTACATTTGA and its complement), the AO $\alpha$ -PPRE (5'-CCTTTCCCGAACGTGACCTTTGT-CCTGGTCCCCTTTTGCT and its complement), and the M4-PPRE (5'-CCTCTCCTTattttaATTGAACTATTACCTACATTTGA and its complement (Miyata et al., 1993; Zhang et al., 1993). Underlined sequences indicate the locations of TGACCT-like motifs. M4-PPRE is a mutant version of the wild-type HD-PPRE

in which the first TGACCT-like repeat is mutated as indicated by the nucleotides in lower case. Complementary oligonucleotides were annealed and end-labeled with the Klenow fragment of DNA polymerase I and [ $\alpha$ - $^{32}$ P]dATP. 2 to 3  $\mu$ l of programmed lysate was incubated with 0.2 pmol of labeled probe at 25°C in a final reaction volume of 15  $\mu$ l containing 7 mM Hepes (pH 7.9), 140 mM NaCl, 1 mM EDTA, 7% (v/v) glycerol, 4  $\mu$ g bovine serum albumin, 4  $\mu$ g nonspecific competitor DNA (poly dI•dC and sonicated salmon sperm DNA, 1:1 weight ratio), 150  $\mu$ M phenylmethylsulfonyl fluoride and 0.2 mM dithiothreitol. The total amount of reticulocyte lysate in each reaction was kept constant by addition of unprogrammed lysate. Where indicated, 1  $\mu$ l of preimmune serum or 1  $\mu$ l of anti-rat HNF-4 serum was added to the binding reactions, which were then preincubated for 5 min prior to the addition of probe. Binding reactions were analyzed by electrophoresis at 4°C on prerun 3.5% polyacrylamide gels (30:1 acrylamide/N',N'-methylenebisacrylamide weight ratio) with 22 mM Tris base/22 mM boric acid/1 mM EDTA as running buffer, followed by autoradiography.


## 2.4 RESULTS

### 2.4.1 *HNF-4 Interacts with the AOx- and HD-PPREs*


Both the AOx-PPRE and the HD-PPRE contain two direct repeats of the TGACCT-related half site separated by a single nucleotide (denoted DR1) (Dreyer et al., 1992; Osumi et al., 1991; Zhang et al., 1993). The HD-PPRE contains an additional upstream direct repeat separated by a 2 nucleotide spacing (DR2) that is necessary for peroxisome proliferator-

responsiveness (Miyata et al., 1993). Fig. 2-1 compares the sequences of the AOx- and HD-PPREs with the HNF-4 consensus binding site (Sladek et al., 1990). The HNF-4 consensus binding sequence matches the AOx-PPRE in 11 of 12 positions, almost overlapping the DR1 repeat. The best match of the HNF-4 consensus binding sequence within the HD-PPRE is 8 of 12 nucleotides encompassing the entire first repeat and part of the second repeat of the HD-PPRE. This high degree of similarity led us to examine whether HNF-4 is among the rat liver nuclear proteins that bind to the AOx- and HD-PPREs.

As shown in Fig. 2-2, incubation of labelled AOx- (lane a) and HD-PPRE (lane d) probes with nuclear extracts prepared from H4IIEC3 cells, a rat hepatoma cell line responsive to peroxisome proliferators, resulted in the formation of a major protein/DNA complex in each case. We have previously demonstrated that PPAR, RXR, and COUP-TF are among the repertoire of DNA-binding proteins present in hepatoma cells that are capable of binding to both the AOx- and HD-PPREs (Miyata et al., 1993). We used antiserum to rat HNF-4 to determine whether HNF-4-like proteins were also present in these protein/DNA complexes. As shown in Fig. 2-2, inclusion of antiserum specific to HNF-4 (lane c) but not preimmune serum (lane b) led to the formation of a supershifted complex (arrowhead) and a corresponding decrease in the signal intensity of the major protein/DNA complex formed with the AOx-PPRE probe, indicating that HNF-4, or HNF-4-related proteins, present in this extract interact with the AOx-PPRE. A significant fraction of the AOx-PPRE/protein complex could be supershifted with the anti-HNF-4 serum, indicating that HNF-4 is a major component of the protein complexes formed on this element. In similar experiments carried out with the HD-PPRE, the generation of a supershifted complex with anti-HNF-4 serum was




**Figure 2-1. Comparison of the AOx-PPRE and the HD-PPRE to the consensus HNF-4 binding site.** The consensus HNF-4 binding site (Sladek et al., 1990) is shown at top and is compared to the AOx- and HD-PPREs. The arrows indicate the TGACCT-related repeat motifs.



**Figure 2-2. Endogenous HNF-4 related proteins of rat hepatoma H4IIEC3 cells bind to the AOx- and HD-PPREs.** Electrophoretic mobility shift assays were performed with  $^{32}\text{P}$ -labeled AOx-PPRE probe or HD-PPRE probe, as indicated at bottom. Probes were incubated with rat hepatoma H4IIEC3 nuclear extract alone (lanes a and d) or with nuclear extract supplemented with preimmune serum (+PI, lanes b and e) or anti-rat HNF-4 serum (+αHNF-4, lanes c and f). The arrowheads in lanes c and f indicate the supershifted complexes observed with anti-HNF-4 serum. This supershifted complex is readily evident with the AOx-PPRE probe (lane c) but is barely visible with the HD-PPRE probe (lane f). The band indicated by the open arrowhead in lane e is a nonspecific complex generated by a factor present in the preimmune serum that has a higher affinity for the HD-PPRE probe than for competitor DNA or for the AOx-PPRE probe.

at the limit of detection (arrowhead, lane f). The above results suggest that HNF-4 or HNF-4-related proteins present in extracts of rat liver hepatoma cells bind to the AOx-PPRE and, with much lower affinity, to the HD-PPRE. The differences in affinities correlate with the degree of relatedness of the respective PPREs to the HNF-4 consensus binding sequence.

To determine if HNF-4 could bind directly to the AOx- and HD-PPREs, we transcribed and translated rat HNF-4 *in vitro* from its cDNA and used the *in vitro* synthesized protein in mobility shift assays. As shown in Fig. 2-3, *in vitro* synthesized HNF-4 bound strongly to the AOx-PPRE probe (lane b) and more weakly to the HD-PPRE probe (lane d). The binding observed with both the AOx- and HD-PPREs was sequence specific, as determined by using mutant oligonucleotides as probes and competitors. For example, HNF-4 was unable to bind to a mutant HD-PPRE probe (M4) in which the second TGACCT repeat was mutated (Fig. 2-3, compare M4, lane f to HD, lane d), indicating that the first direct repeat in the HD-PPRE, which encompasses the HNF-4 consensus sequence, is required for binding to HNF-4. We have previously shown that luciferase reporter constructs containing the M4 mutant HD-PPRE have a reduced response to peroxisome proliferators *in vivo*, yet PPAR/RXR heterodimers could still efficiently bind to this element *in vitro* (Miyata et al., 1993). HNF-4 and PPAR/RXR may therefore have overlapping recognition sequences on the HD-PPRE, similar to what is observed with COUP-TF (Miyata et al., 1993). This finding would be consistent with the overlapping sequence identity between the middle repeat of the HD-PPRE and the HNF-4 consensus binding site (Fig. 2-1). It is not yet known whether all three receptors can occupy this site simultaneously.




**Figure 2-3.** HNF-4 synthesized *in vitro* binds to the AOx- and HD-PPREs. Electrophoretic mobility shift assays were carried out with the indicated labeled probes in the absence (-) or presence (+) of *in vitro* synthesized rat HNF-4. The arrowheads in lanes b and d indicate the HNF-4 protein/DNA complexes. The M4 probe used in lanes e and f is a mutant HD-PPRE in which the first TGACCT-like repeat is disrupted.



#### 2.4.2 *HNF-4 Represses PPAR $\alpha$ -dependent Induction from the AOx-PPRE both in the Presence and Absence of a Potent Peroxisome Proliferator*

To examine the consequences of ectopic expression of HNF-4 on transcription from the AOx- and HD-PPREs, we carried out transient transfection assays with luciferase reporter gene constructs containing the minimal carbamoyl phosphate synthetase promoter linked to either 2 tandem copies of the minimal AOx-PPRE (pAOx(X2)*luc*) or 3 tandem copies of the minimal HD-PPRE (pHD(X3)*luc*) (Zhang et al., 1993). The effector plasmids that were used expressed full-length cDNAs encoding rat PPAR $\alpha$ , human RXR $\alpha$ , and rat HNF-4. BSC40 cells were used in the transient transfection assays, since efficient induction by peroxisome proliferators in this cell line requires the cotransfection of both PPAR and RXR (Miyata et al., 1993).

Cotransfection of pAOx(X2)*luc* with PPAR $\alpha$  and RXR $\alpha$  expression vectors resulted in a 5- to 10-fold induction in activity of the reporter gene, as we have previously demonstrated (Fig. 2-4)(Marcus et al., 1993). This PPAR $\alpha$ -mediated induction is independent of exogenously added peroxisome proliferators and likely results from the presence of endogenous PPAR and/or RXR activators present in these cells. Addition of the potent peroxisome proliferator Wy-14,643 resulted in a 20-fold stimulation of activity over basal levels. This stimulation required the presence of both PPAR $\alpha$  and RXR $\alpha$ . Expression of HNF-4 during the transient transfection resulted in inhibition of both the PPAR $\alpha$ -dependent/Wy-14,643-independent induction and the PPAR $\alpha$ -dependent/Wy-14,643-dependent induction. Inhibition was dose-dependent under both circumstances. Inclusion of 4  $\mu$ g of the HNF-4 expression plasmid reduced PPAR $\alpha$ -dependent activation to the basal




**Figure 2-4. HNF-4 antagonizes rPPAR-mediated induction of an AOx-PPRE reporter gene.** pAOx(X2)*luc* was transfected into BSC40 cells in the presence or absence of the peroxisome proliferator Wy-14,643, along with a constant amount of rPPAR $\alpha$  and RXR $\alpha$  expression plasmids and increasing amounts of rat HNF-4 expression plasmid (in  $\mu$ g), as indicated. Transfections were carried out in duplicate and repeated a minimum of 3 times. The values shown ( $\pm$  standard error of the mean) are normalized to the value obtained from Wy-14,643-treated cells cotransfected with rPPAR and RXR $\alpha$  expression plasmids, which was taken as 100%.


levels observed with pAOx(X2)*luc* alone. Similarly, peroxisome proliferator-dependent induction was almost completely inhibited in the presence of 4  $\mu$ g of HNF-4 expression plasmid. HNF-4-dependent inhibition was specific for the PPAR $\alpha$ -mediated response, since the basal level expression of pAOx(X2)*luc* was not significantly affected by cotransfection with increasing amounts of HNF-4 expression plasmid (Fig. 2-6). These findings demonstrate that HNF-4 antagonizes induction from the AOx-PPRE by PPAR $\alpha$ , both in the absence and in the presence of an exogenously added peroxisome proliferator.

#### *2.4.3 HNF-4 Potentiates Activity of PPAR $\alpha$ on the HD-PPRE in the Presence of a Peroxisome Proliferator*

Transfection experiments like those carried out with the AOx-PPRE reporter gene construct were carried out with the HD-PPRE reporter gene construct. Cotransfection of pHD(X3)*luc* with both PPAR $\alpha$  and RXR $\alpha$  expression plasmids gave a 5- to 10-fold peroxisome proliferator-independent induction and a 50-fold peroxisome proliferator-dependent induction of reporter gene activity over basal levels (Fig. 2-5). As seen with the AOx-PPRE reporter gene construct, inclusion of HNF-4 during transient transfection resulted in inhibition of the peroxisome proliferator-independent activation mediated by PPAR $\alpha$ /RXR $\alpha$  from the HD-PPRE. However, HNF-4-mediated inhibition was less efficient with the HD-PPRE than with the AOx-PPRE, causing only a 50% inhibition of activation with 4  $\mu$ g of HNF-4 expression plasmid as compared to a control transfection in the absence of HNF-4. Surprisingly, when Wy-14,643 was included in the transfections, HNF-4 significantly stimulated the induction response of the HD-PPRE. Cotransfection with 4  $\mu$ g



**Figure 2-5. HNF-4 differentially modulates the function of rPPAR $\alpha$  on the HD-PPRE.** Cotransfection of pHD(X3)*luc* with various expression plasmids into BSC40 cells in the presence or absence of Wy-14,643 and luciferase activity measurements were performed as described in Fig. 2-4. The results demonstrate that HNF-4 antagonizes rPPAR $\alpha$  function in the absence of Wy-14,643 but potentiates rPPAR $\alpha$  activity in the presence of Wy-14,643.




**Figure 2-6. HNF-4 alters transactivation from the AOx- and HD-PPREs in the presence of rPPAR $\alpha$  and RXR $\alpha$ .** BSC40 cells were co-transfected as before with pAOx(X2)*luc* or pHD(X3)*luc* reporter plasmids and with rPPAR and RXR $\alpha$  (2  $\mu$ g each) or with HNF-4 (4  $\mu$ g) expression plasmids as indicated. Co-transfections were performed with or without the peroxisome proliferator Wy-14,643. Values are presented relative to the value obtained for cells co-transfected with rPPAR and RXR $\alpha$  in the presence of Wy-14,643 (taken as 100%). The values represent the average of three independent transfections carried out in duplicate.

of HNF-4 expression plasmid resulted in a 2- to 3-fold potentiation of the PPAR $\alpha$ -mediated response to the presence of this peroxisome proliferator. Both repression and stimulation by HNF-4 required the presence of PPAR $\alpha$  and RXR $\alpha$ , since HNF-4 alone had no significant effects on the basal level expression of the HD-PPRE reporter gene construct (Fig. 2-6). Therefore, in contrast to the results seen with the AOx-PPRE, HNF-4 cooperates with PPAR $\alpha$  to stimulate transcription of the HD-PPRE reporter gene construct, but only in the presence of a peroxisome proliferator. Because HNF-4 both repressed the PPAR $\alpha$ -mediated response in the absence of drug and increased the response in the presence of drug, the peroxisome proliferator-dependent induction ratio was effectively increased from 5-fold to approximately 35-fold.

#### 2.4.4 HNF-4 Does Not Cooperate with RXR $\alpha$ or PPAR $\alpha$ for DNA Binding

PPAR $\alpha$  heterodimerizes with RXR $\alpha$  and binds cooperatively to both the AOx- and HD-PPREs *in vitro* (Fig. 2-7, compare lanes b and c with lanes e)(Marcus et al., 1993). In contrast, HNF-4 has been shown to bind to cognate response elements exclusively as a homodimer, and there is no evidence that HNF-4 heterodimerizes with other members of the nuclear hormone receptor family (Sladek et al., 1990). Nevertheless, we explored the possibility that the effects of HNF-4 on PPAR $\alpha$ -dependent induction of transcription *in vivo* was mitigated by cooperative interactions with PPAR and/or RXR $\alpha$  by mobility shift analyses with *in vitro* translated receptor proteins. As shown in Fig. 2-7, the binding of HNF-4 to the AOx-PPRE or the HD-PPRE was not affected by the presence of RXR $\alpha$  or PPAR $\alpha$  and *vice versa* (compare lanes d to lanes f and g, respectively). Therefore, the ability of HNF-4 to



**Figure 2-7. HNF-4 does not cooperate with rPPAR $\alpha$  or RXR $\alpha$  for DNA binding.** *In vitro* translated human RXR $\alpha$ , rPPAR $\alpha$  and rat HNF-4 were incubated singly or in pairwise combinations, as indicated, with labeled AOx-PPRE probe (upper panel) or HD-PPRE probe (lower panel) and analyzed by electrophoretic mobility shift assay. Lane a contained probes incubated with unprogrammed reticulocyte lysate. Arrow, nonspecific complex formed between the HD-PPRE probe and reticulocyte lysate.

modulate PPAR function apparently is not the result of cooperative DNA binding by HNF-4 with either RXR $\alpha$  or PPAR $\alpha$ .

## 2.5 DISCUSSION

The results presented herein demonstrate that HNF-4 differentially modulates the functioning of the peroxisome proliferator-activated receptor on specific PPREs, suggesting that this nuclear hormone receptor plays a role in the regulation of expression of genes encoding peroxisomal  $\beta$ -oxidation enzymes. HNF-4 has been shown to act as a cell-restricted, positive regulator of genes whose products are involved in several metabolic pathways (Chen et al., 1994; Ladas et al., 1992; Sladek et al., 1990). Our findings are the first demonstration that HNF-4 can both activate and repress specific target genes involved in lipid metabolism through a common response element.

HNF-4 was a major component of the protein/DNA complex formed between the AOx-PPRE and rat liver hepatoma nuclear extracts, and HNF-4 synthesized *in vitro* interacted strongly with the AOx-PPRE. In transient transfection assays, HNF-4 competitively inhibited PPAR $\alpha$ -dependent transactivation of a reporter gene linked to the AOx-PPRE, both in the presence or the absence of the peroxisome proliferator, Wy-14,643. The mechanism by which HNF-4 antagonizes PPAR $\alpha$  functioning is not known at present, but one scenario is that HNF-4 can compete directly with PPAR $\alpha$ /RXR heterodimers for binding to the AOx-PPRE. This explanation would be consistent with the near perfect match between the consensus HNF-4 binding site and part of the AOx-PPRE and with the fact that the region of homology between the two sequences almost completely overlaps the two



TGACCT DR1 repeats that are essential for PPAR $\alpha$ /RXR $\alpha$  binding to the AOx-PPRE. HNF-4 was unable to bind cooperatively with either PPAR $\alpha$  or RXR $\alpha$ , implying that competition for heterodimerization partners is probably not involved in the observed inhibition of activation from the AOx-PPRE. However, it remains possible that under physiological conditions, non-DNA binding heteromers could form between HNF-4 and PPAR $\alpha$  or RXR $\alpha$ , thereby leading to inhibition of the PPAR $\alpha$ -mediated response via indirect mechanisms.

Inhibition of activation from the AOx-PPRE by HNF-4 could result from the absence of an appropriate ligand for HNF-4 in the BSC40 cells used in the transfection assays. It has been shown that, depending on the nature of the response element, the thyroid receptor can function as a repressor in the unliganded state but as an activator when associated with its cognate hormone (Damm et al., 1989; Graupner et al., 1989; Sap et al., 1989). The repression observed with the thyroid receptor in the absence of its ligand is thought to be mediated via an active intrinsic silencing domain that may directly inhibit the formation of the functional preinitiation complex (Banahmad et al., 1992; Fondell et al., 1993). There is no evidence as of yet that HNF-4 possesses an active transcriptional silencing function or even that a ligand exists for this receptor.

The results obtained with the HD-PPRE are even more intriguing than those obtained with the AOx-PPRE. HNF-4 had a repressive effect (50%) on the PPAR $\alpha$ -dependent activation of the HD-PPRE in the absence of exogenously added peroxisome proliferator. However, in the presence of Wy-14,643, HNF-4 cooperated with PPAR $\alpha$ /RXR $\alpha$  to provoke a significant stimulation of transcription. Therefore, under conditions of peroxisome proliferator administration, HNF-4 had diametric effects on the PPAR $\alpha$ -mediated, peroxisome

proliferator responsiveness of the AOx-PPRE *vis-à-vis* the HD-PPRE. It is unclear whether the effects of HNF-4 on activation from the HD-PPRE are mitigated directly through binding to the HD-PPRE or indirectly, for example, through interaction with, or modulation of, auxiliary coregulators involved in peroxisome proliferator signalling or transcriptional control.

HNF-4 synthesized *in vitro* or present endogenously in rat hepatoma cells had only a weak affinity for the HD-PPRE, although its binding to the HD-PPRE was specific for the sequence of this element. The structure of the HD-PPRE is more complex than that of the AOx-PPRE. The HD-PPRE consists of three TGACCT-related half-sites separated by 2 nucleotides and 1 nucleotide, respectively. The integrity of all three repeats, as well as the spacing between them, are necessary for efficient peroxisome proliferator responsiveness *in vivo*; however, only the DR1 repeats are necessary for PPAR $\alpha$ /RXR $\alpha$ -binding *in vitro* (Miyata et al., 1993). The region within the HD-PPRE that most closely matches the HNF-4 consensus sequence contains the first direct repeat. We have shown that this repeat is necessary for HNF-4 binding (Fig. 2-3). It is conceivable that binding of HNF-4 to the first repeat might interfere with binding of PPAR $\alpha$ /RXR to the HD-PPRE, leading to the inhibitory effects on activation observed in transient transfections. Activation of PPAR $\alpha$  by exogenously added peroxisome proliferators may serve to alter the binding affinity or conformation of the PPAR $\alpha$ /RXR $\alpha$  heterodimer specifically on the HD-PPRE, thereby allowing cooperative interactions with HNF-4 and consequently potentiation of induction. We have previously shown that the first repeat in the HD-PPRE is also required for binding to COUP-TF1, an orphan receptor that antagonizes peroxisome proliferator-responsiveness *in vivo* (Miyata et al., 1993). Therefore, the additional repeat element present in the HD-

PPRE compared to the AOx-PPRE may allow more complex strategies of regulation for the HD gene.

It is becoming increasingly apparent that PPARs play a central role in regulating lipid metabolic pathways. More recently, PPARs have been implicated in cellular differentiation and proliferation (Chawla and Lazar, 1994). Our findings suggest that HNF-4 can modulate PPAR function both positively and negatively, consequently resulting in differential regulation of at least two distinct peroxisome proliferator-responsive genes. These differential effects are conditional upon the structure of a particular target PPRE and the presence or absence of PPAR activators. This finding is interesting in light of the recent observation that HNF-4 activates transcription of the medium-chain acyl-CoA dehydrogenase gene (Carter et al., 1993). Medium-chain acyl-CoA dehydrogenase is the initial and rate-limiting enzyme of the mitochondrial fatty acid  $\beta$ -oxidation pathway, which metabolizes short- and medium-chain length fatty acids in response to energy demands of the cell. In contrast, the peroxisomal  $\beta$ -oxidation pathway preferentially uses long-chain fatty acids as substrates, and acyl-CoA oxidase is the initial and rate-limiting step of this pathway. Our finding that HNF-4 represses PPAR-mediated induction of the AOx gene suggests that HNF-4 may play a key role in coordinating and integrating the mitochondrial and peroxisomal  $\beta$ -oxidation pathways through the bidirectional regulation of the respective rate-limiting enzymes. Similarly, the HNF-4-mediated, peroxisome proliferator-dependent upregulation of the activity of the HD gene, whose product catalyzes the second step in the peroxisomal  $\beta$ -oxidation pathway, may represent an adaptive cellular response that primes the peroxisomal  $\beta$ -oxidation pathway to respond rapidly to cellular oxidative demands under physiological conditions where repression

of the AOX gene is alleviated. HNF-4/PPAR cooperativity on the HD-PPRE may be relevant to the observation that HD accumulates to higher levels in the liver compared to the other two enzymes of the peroxisomal  $\beta$ -oxidation pathway following administration of peroxisome proliferators (Reddy et al., 1986a). Our findings add an additional layer of complexity to peroxisome proliferator-signalling pathways and may begin to provide insight into how combinatorial interactions among distinct nuclear receptors and convergence of multiple signalling pathways integrate complex metabolic regulatory networks.

## CHAPTER 3

### **Calreticulin Modulates the In Vitro DNA Binding But Not the In Vivo Transcriptional Activation by Peroxisome Proliferator-Activated Receptor/Retinoid X Receptor Heterodimers<sup>†</sup>**

<sup>†</sup>A version of this chapter has been published. Winrow, C.J., Miyata, K.S., Marcus, S.L., Burns, K., Michalak, M., Capone, J.P. and Rachubinski. 1995. Mol. and Cell. Endocrinol. **111**:175-179. Used with permission from Elsevier Science.

### 3.1 ABSTRACT

Calreticulin is a ubiquitous calcium binding/storage protein found primarily in the endoplasmic reticulum. Calreticulin has been shown to inhibit DNA binding and transcriptional activation by glucocorticoid and androgen hormone receptors by binding to the conserved sequence KXFF(K/R)R, present in the DNA-binding domains of all known members of the steroid/nuclear hormone receptor superfamily. To determine whether calreticulin might be a general regulator of hormone-responsive pathways, we examined its effect on DNA binding *in vitro* and transcriptional activation *in vivo* by heterodimers of the peroxisome proliferator-activated receptor (PPAR) and the 9-*cis* retinoic acid receptor (RXR $\alpha$ ). We show here that purified calreticulin inhibits the binding of PPAR $\alpha$ /RXR $\alpha$  heterodimers and of other nuclear hormone receptors, to peroxisome proliferator-responsive DNA elements *in vitro*. However, overexpression of calreticulin in transiently transfected cultured cells had little or no effect on transactivation mediated by PPAR $\alpha$ /RXR $\alpha$ . Therefore, while calreticulin inhibits the binding of both nuclear and steroid hormone receptors to cognate response elements *in vitro*, our findings suggest that calreticulin does not necessarily play an important role in the regulation of all classes of hormone receptors *in vivo*.

### 3.2 INTRODUCTION

Members of the steroid/nuclear hormone receptor superfamily play crucial roles in organismal development, differentiation and homeostasis. These receptors mediate physiological and cellular responses to extracellular and intracellular stimuli through the regulation of diverse gene networks. Steroid/nuclear hormone receptors are structurally related ligand-activated transcription factors that bind to cognate response elements upstream of target genes, conferring responsiveness to steroid hormones such as glucocorticoids, androgens and estrogen and to non-steroid hormones such as retinoids, thyroid hormone, and vitamin D3. These transcription factors are subject to a variety of post-transcriptional regulatory strategies that serve to modulate their subcellular distribution, abundance, DNA-binding capacity, and function. Therefore, steroid receptors like the glucocorticoid receptor (GR), estrogen receptor, progesterone receptor, and androgen receptor (AR) are found in multi-component cytoplasmic complexes with non-receptor cellular factors, including the heat shock proteins hsp90, hsp70 and hsp56. Ligand binding facilitates the dissociation of the receptor from this complex, with subsequent nuclear translocation and binding of the receptor to cognate response sites (Evans, 1988; Truss and Beato, 1993; Tsai and O'Malley, 1994).

Recently, calreticulin has been shown to modulate DNA binding and transcriptional activation by AR and GR (Burns et al., 1994; Dedhar et al., 1994). Calreticulin is a multifunctional Ca<sup>2+</sup>-binding/storage, luminal protein of the endoplasmic reticulum (Michalak et al., 1992) that has been shown to interact with the conserved amino acid sequence KLGFFKR of integrins (Rojiani et al., 1991). This sequence is similar to the sequence KXFF(K/R)R found in the conserved DNA-binding domain of members of the steroid/nuclear

hormone receptor superfamily (Burns et al., 1994; Laudet et al., 1992). Calreticulin was shown to bind to this sequence present in GR and AR, resulting in the inhibition of these receptors to bind to DNA *in vitro* and to activate transcription *in vivo* (Burns et al., 1994; Dedhar et al., 1994). These findings are surprising since calreticulin is localized predominantly to the endoplasmic reticulum, and therefore would not be expected to have access to nuclear-resident transcription factors. The exact mechanism responsible for calreticulin-mediated modulation of steroid receptor function *in vivo* remains unclear.

Since the consensus calreticulin binding site is conserved in all members of the steroid/nuclear hormone receptor superfamily, calreticulin might be considered to be a general regulator of all nuclear hormone-responsive pathways. GR and AR are found both in the cytoplasm and the nucleus and bind to their response elements as homodimers. Receptors for retinoic acid (RAR), 9-*cis* retinoic acid (RXR $\alpha$ ), vitamin D3, thyroid hormone, and peroxisome proliferators (PPAR) constitute a functionally distinct subclass within the steroid/nuclear hormone receptor superfamily. These receptors are constitutively localized to the nucleus, do not form complexes with cytoplasmic factors, and can bind to DNA in the absence of ligand. Nuclear hormone receptors can bind to DNA as homodimers, although high-affinity DNA binding is generally manifested through heterodimerization with a common auxiliary cofactor such as RXR.

We explored the role of calreticulin on gene regulation by this class of nuclear hormone receptors by investigating its effects on the function of human RXR $\alpha$  and rat PPAR $\alpha$ . RXR and PPAR bind as heterodimers to peroxisome proliferator-response elements (PPREs) and mediate transcriptional activation by peroxisome proliferators, a broad class of



xenobiotic compounds, and by selected fatty acids (Gearing et al., 1993; Gottlicher et al., 1992; Issemann and Green, 1990; Kliewer et al., 1992a; Marcus et al., 1993). We show that calreticulin inhibits the *in vitro* binding of PPAR/RXR heterodimers to the PPRE of the gene encoding fatty acyl-CoA oxidase (AOx), the first enzyme of the peroxisomal  $\beta$ -oxidation cascade. However, in transiently transfected cultured cells, overexpression of calreticulin results in only marginal effects on the transcriptional activation of the AOx PPRE by PPAR/RXR. We conclude that while calreticulin can inhibit the DNA binding of both steroid and nuclear hormone receptors *in vitro*, it may not be physiologically important in the *in vivo* regulation of all members of this receptor superfamily.

### 3.3 MATERIALS AND METHODS

#### 3.3.1 Plasmids, Cell Culture, and Transfection

The luciferase reporter plasmids pCPS*luc*, containing the minimal promoter of the rat carbamoyl phosphate synthetase gene, and pAOx(X2)*luc*, containing 2 copies of the minimal PPRE of the rat AOx gene (5'-CCTTTCCGAACGTGACCTTTGTCCTG-GTCCCCTTTTGCT), have been described in Chapter 2 (Zhang et al., 1993). The glucocorticoid response element (GRE) reporter plasmid MMTV-GRE*luc* (pJA358) has been reported (Burns et al., 1994). Plasmid vectors for *in vivo* expression and *in vitro* transcription/translation of rat PPAR $\alpha$ , human RXR $\alpha$ , and human HNF-4 have been described (Marcus et al., 1993; Winrow et al., 1994). The expression plasmid for human COUP-TF1 contains the COUP-TF1 cDNA inserted as an *HindIII/XbaI* fragment into pRC/CMV (Miyata

et al., 1993). The expression vector pSVL-CRT was constructed by insertion of the cDNA encoding full-length rabbit skeletal muscle calreticulin into the plasmid pSVL (Pharmacia)(Burns et al., 1994). The plasmid VARO was used to express GR and was provided by K.R. Yamamoto (University of California, San Diego)(Pearce and Yamamoto, 1993). Transient transfection assays were carried out in BSC40 cells (Miyata et al., 1993) using various expression vectors as indicated in the figure legends. Luciferase activity was measured as described in Chapter 2. Where indicated, the peroxisome proliferator Wy-14,643 or dexamethasone was added to transfected cells at a final concentration of 0.1 mM and 1  $\mu$ M, respectively (Marcus et al., 1993). The plasmid pCH110 (Pharmacia) containing the *lacZ* gene was included in transfections to normalize for the efficiency of transfection.

### 3.3.2 Electrophoretic Mobility Shift Assays

Electrophoretic mobility shift assays were done with  $^{32}$ P-labelled AOx-PPRE probe and receptors transcribed and translated *in vitro* from their respective cDNAs, as previously described in Chapter 2 (Miyata et al., 1993). Where indicated, 1  $\mu$ g of purified calreticulin (Baksh et al., 1992) was added to the reactions prior to the addition of probe DNA.

## 3.4 RESULTS

### 3.4.1 Calreticulin Inhibits PPAR/RXR Binding to DNA

Calreticulin has been reported to interact with the DNA-binding domains of GR (Burns et al., 1994) and AR (Dedhar et al., 1994) via the conserved amino acid sequence


KXFF(R/K)R located between the two "zinc fingers" in these and other steroid receptors (Laudet et al., 1992). In this study we examined whether calreticulin could inhibit the cooperative DNA binding of PPAR/RXR heterodimers to the AOx-PPRE. Rat PPAR $\alpha$  (hereafter called PPAR) and human RXR $\alpha$  (hereafter called RXR) contain the sequences KGFFRR and KGFFKR, respectively, in their DNA-binding domains (Fig. 3-1). An electrophoretic mobility shift assay was done with *in vitro* translated PPAR and RXR and a radiolabelled synthetic oligonucleotide probe corresponding to the AOx-PPRE. *In vitro* translated PPAR and RXR formed a specific protein/DNA complex on the AOx-PPRE (Fig. 3-2, *lane c*). PPAR and RXR do not bind individually to the AOx-PPRE (Gearing et al., 1993; Kliwer et al., 1992a; Marcus et al., 1993). Addition of purified calreticulin abolished the binding of the PPAR/RXR heterodimer to the AOx-PPRE (*lane f*). Calreticulin alone did not interact with the AOx-PPRE (*lane b*). COUP-TF1 and HNF-4, two additional nuclear hormone receptors, also form complexes as homodimers on the AOx-PPRE (Miyata et al., 1993; Winrow et al., 1994)(Fig. 3-2, *lanes d* and *e*, respectively). Complex formation by either receptor was completely abolished by addition of calreticulin (compare *lane d* to *lane g* and *lane e* to *lane h*). Therefore, calreticulin interacts *in vitro* with PPAR/RXR heterodimers and COUP-TF1 and HNF-4 homodimers, preventing their binding to the AOx-PPRE.

#### 3.4.2 Calreticulin Shows Little or No Effect on Transactivation by PPAR/RXR In Vivo

Transient transfection assays were carried out to investigate the effects of calreticulin expression on transactivation mediated by PPAR/RXR *in vivo*. These experiments used the

<b>PPAR<math>\alpha</math></b>	VHACEGC	<b>KGFFRR</b>	T I R L K L A
<b>RXR<math>\alpha</math></b>	VYSCEGC	<b>KGFFKR</b>	T V R K D L T
<b>COUP-TF1</b>	VSSSV I E	<b>QLFFVR</b>	L V G K T P I
<b>HNF-4</b>	ASSCDGC	<b>KGFFRR</b>	S V R K N H M
<b>GR</b>	VLTCGSC	<b>KVFFKR</b>	A V E G Q H N
<b>CRT binding site</b>		<b>KXFF(K/R)R</b>	

**Figure 3-1. The DNA-binding domains of nuclear hormone and steroid receptors contain sequences homologous to the calreticulin binding site.** Portions of the DNA-binding regions of the nuclear hormone receptors rat PPAR $\alpha$ , human RXR $\alpha$ , human COUP-TF1, and rat HNF-4 and of the steroid receptor human GR are shown, along with the consensus calreticulin (CRT) binding site. The amino acids highlighted in bold conform to the consensus calreticulin binding site.





**Figure 3-2. Calreticulin inhibits PPAR/RXR heterodimer binding to the AOx-PPRE *in vitro*.** An electrophoretic mobility shift assay was performed with <sup>32</sup>P-labeled AOx-PPRE probe and *in vitro* translated PPAR and RXR (*lane c*). The AOx-PPRE probe was also incubated with *in vitro* translated COUP-TF1 (*lane d*) and HNF-4 (*lane e*). Incubation of the *in vitro* translated receptors with the AOx-PPRE probe was for 25 min at 28°C. Where indicated, 1 µg of purified rabbit calreticulin (CRT) was preincubated with *in vitro* translated PPAR/RXR (*lane f*), COUP-TF1 (*lane g*), and HNF-4 (*lane h*) for 15 min at 28°C, followed by incubation with the AOx-PPRE probe as described above. No complex was formed by the addition of unprogrammed reticulocyte lysate (*lane a*).


luciferase reporter gene plasmid construct pAOx(X2)*luc* containing 2 copies of the AOx-PPRE, which has been shown to be responsive to transactivation by PPAR/RXR (Marcus et al., 1993). In agreement with our earlier report (Marcus et al., 1993; Winrow et al., 1994), cotransfection of BSC40 cells with pAOx(X2)*luc* and plasmids expressing PPAR and RXR led to 6-fold and 20-fold increases in luciferase reporter gene activity over control transfections containing empty vectors in the absence and presence of the added peroxisome proliferator Wy-14,643, respectively (Fig. 3-3A). Addition of increasing amounts of the calreticulin expression plasmid pSVL-CRT to the transfections showed no significant effects on either the peroxisome proliferator-dependent or peroxisome proliferator-independent activation of the reporter gene construct by PPAR/RXR. The slight inhibition of transactivation (10 to 15%) observed with increasing amounts of pSVL-CRT is apparently nonspecific, as similar levels of inhibition were seen in control transfections carried out with pCPS*luc*, the parental reporter plasmid used to construct pAOx(X2)*luc* (Fig. 3-3B).

Given the dramatic inhibitory effect of calreticulin on the *in vitro* binding of PPAR/RXR to the AOx-PPRE (Fig. 3-2), it was surprising that calreticulin expression had very little or no effect on the PPAR/RXR-mediated *in vivo* activation of the pAOx(X2)*luc* reporter construct. Burns et al. (Burns et al., 1994) have shown that overexpression of calreticulin in mouse L cells results in a 50 to 80% inhibition in the GR-mediated expression of a luciferase reporter gene under the control of a MMTV-GRE promoter. Therefore, to insure that calreticulin was functionally expressed under the experimental conditions used in this study, we transfected BSC40 cells with the MMTV-GRE*luc* reporter plasmid pJA358, the VARO plasmid expressing GR, and the calreticulin expression vector pSVL-CRT. The

**Figure 3-3. Calreticulin fails to repress the *in vivo* transactivation from the AO $\alpha$ -PPRE by PPAR/RXR.** (A). pAO $\alpha$ (2X)*luc* was transfected into BSC40 cells along with effector plasmids expressing PPAR, RXR, and calreticulin (CRT). Transfections contained 5  $\mu$ g of the reporter plasmid pAO $\alpha$ (X2)*luc* and, where indicated, 2  $\mu$ g of hRXR $\alpha$ -expression plasmid, 2  $\mu$ g of rPPAR $\alpha$ -expression plasmid, and varying amounts (in  $\mu$ g) of the calreticulin-expression plasmid pSVL-CRT. Control transfections contained the corresponding empty vectors. Cells were grown in media that did or did not contain 0.1 mM of the peroxisome proliferator, WY-14,643. Cells were harvested 48-h post-transfection, and luciferase activity was determined. The values shown are relative activities from three independent transfections done in duplicate and were normalized to the value obtained for WY-14,643-treated cells cotransfected with rPPAR and hRXR $\alpha$  expression plasmids, which was taken as 100%. (B). Transfection was done as in (A), except that the reporter plasmid was pCPS*luc*. The values shown are relative activities from three independent transfections done in duplicate and were normalized as in (A).







**Figure 3-4. Calreticulin represses transactivation from the GRE by GR.** The GRE reporter plasmid pJA358 was transfected into BSC40 cells along with effector plasmids expressing GR and calreticulin. Transfections contained 5  $\mu$ g of the reporter plasmid pJA358 and, where indicated, 2  $\mu$ g of the GR-expression plasmid VARO and varying amounts (in  $\mu$ g) of the calreticulin-expression plasmid pSVL-CRT. Control transfections contained the corresponding empty vectors. Cells were grown in media that did or did not contain 1  $\mu$ M dexamethasone. Cells were harvested 48h post-transfection, and luciferase activity was determined. The values shown are relative activities from three independent transfections done in duplicate and were normalized to the value obtained for dexamethasone-treated cells transfected with the GR-expression plasmid VARO, which was taken as 100%.

presence of the VARO plasmid increased luciferase activity from pJA358 70-fold upon addition of dexamethasone (Fig. 3-4). This dexamethasone-dependent activation of the reporter gene was inhibited 50 to 90% when the calreticulin expression vector pSVL-CRT was included in the transfection (Fig. 3-4). Therefore, calreticulin inhibits GR-mediated stimulation of gene expression in BSC40 cells, as previously demonstrated in mouse L cells (Burns et al., 1994). This finding suggests that transfection of BSC40 cells with pSVL-CRT led to expression of functional calreticulin which, while capable of inhibiting the transcriptional activation function by GR, had little or no effect on transcriptional activation by PPAR/RXR.

### 3.5 DISCUSSION

We have shown that calreticulin inhibits the *in vitro* DNA binding of the nuclear hormone receptors PPAR and RXR, as well as COUP-TF1 and HNF-4, to a cognate response element, as was previously observed with GR and AR. However, in contrast to the results observed with GR and AR (Burns et al., 1994; Dedhar et al., 1994), overexpression of calreticulin in transiently transfected cells had little or no effect on PPAR/RXR-mediated activation of gene expression.

For calreticulin to play a physiologically relevant role in the regulation of steroid and nuclear hormone receptors, sufficient calreticulin must gain access to the cytoplasm and/or the nucleus (Bleackley et al., 1995). However, calreticulin is localized primarily to the lumen of the endoplasmic reticulum, since it is synthesized with a signal sequence and contains a KDEL retention sequence (Bleackley et al., 1995; Michalak et al., 1992).

Immunofluorescence studies have suggested that some calreticulin can be found in the nuclear envelope and the cytoplasm in some cell types, perhaps in amounts sufficient to affect steroid receptor function (Michalak et al., 1992; Nash et al., 1994; Opas et al., 1991). However, calreticulin did not inhibit PPAR/RXR function *in vivo*. The lack of a pronounced effect of calreticulin expression on PPAR/RXR function *vis-à-vis* GR and AR function may be due to differences in their pathways of activation. Calreticulin may inhibit the function of GR and AR at some step unique to the function of steroid receptors, such as dissociation from cytoplasmic proteins, nuclear compartmentalization, or ligand-dependent DNA binding. The exact mechanism of calreticulin inhibition awaits further study.

In contrast to our findings with PPAR and RXR, Dedhar et al. (1994) have postulated that calreticulin directly inhibits the function *in vivo* of the nuclear hormone receptor RAR. Thus, in a P19 embryonic carcinoma cell line selected for calreticulin overexpression, retinoic acid-induced expression of specific differentiation-associated markers was suppressed, whereas in a cell line in which calreticulin expression was inhibited by anti-sense RNA, retinoic acid-induced expression of these markers was elevated (Dedhar et al., 1994). It is possible that the observed effects were due to an unusual intracellular trafficking of calreticulin in P19 cells that allows for more efficient nuclear accumulation. However, there was no evidence in this system that calreticulin was modulating the transactivating function or DNA binding capacity of RAR *in vivo* by directly binding to RAR. Since calreticulin is thought to play a multifunctional role in the cell, large changes in the cellular levels of calreticulin may have pleiotropic and indirect effects on gene expression patterns.

In summary, we have shown that calreticulin inhibits the DNA binding of both steroid

and nuclear hormone receptors to their response elements *in vitro*. We have also shown that calreticulin does not have a general role in the *in vivo* regulation of all nuclear hormone receptors.

## CHAPTER 4

### **Interplay of the Peroxisome Proliferator-Activated Receptor and Thyroid Hormone Receptor Signalling Pathways in Regulating Peroxisome Proliferator-Responsive Genes<sup>†</sup>**

<sup>†</sup> A version of this chapter has been published. Winrow, C.J., Kassam, A., Miyata, K.S., Marcus, S.M., Hunter, J., Capone, J.P. and Rachubinski, R.A. 1996. *Ann. N.Y. Acad. Sci.* **804**:214-230. Used with permission from The Annals of the New York Academy of Sciences.

## 4.1 INTRODUCTION

Peroxisome proliferator-activated receptors (PPARs) are members of the nuclear hormone receptor superfamily. PPARs are ligand-modulated transcription factors that are activated by structurally diverse xenobiotic chemicals called peroxisome proliferators, as well as by naturally occurring and synthetic fatty acids (Green and Wahli, 1994; Issemann et al., 1993). PPARs modulate transcription of a wide spectrum of genes involved in lipid and metabolic homeostasis, including those encoding the peroxisomal  $\beta$ -oxidation enzymes fatty acyl-CoA oxidase (AOx) and 3-enoyl-CoA hydratase/3-hydroxyacyl-CoA dehydrogenase (Marcus et al., 1993; Tugwood et al., 1992), mitochondrial medium-chain acyl-CoA dehydrogenase (Gulick et al., 1994), malic enzyme (Castelein et al., 1994), phosphoenolpyruvate carboxykinase (Tontonoz et al., 1995), 3-hydroxy-3-methylglutaryl-CoA synthase (Rodriguez et al., 1994), fatty acid binding protein (Kaikaus et al., 1993), cytochrome P4504A6 (Muerhoff et al., 1992), and apolipoprotein AI (Vu-Dac et al., 1994). PPARs exist in a variety of types, some of which are expressed in cell-, tissue-, and developmentally specific manners and respond differentially to activators. This implies that PPARs play distinct roles in the cell. One example is the demonstration that adipocyte-specific PPAR $\gamma$ 2 is a central determinant in regulating the process of adipogenesis (Tontonoz et al., 1994c). PPARs therefore perform essential and diverse functions both in controlling lipid metabolic cascades and in modulating cell differentiation and proliferation.

Regulation of gene expression by PPARs involves the binding of PPARs to specific peroxisome proliferator-response elements (PPREs) through heterodimerization with the 9-*cis* retinoic acid receptor RXR $\alpha$ . Therefore, the retinoid and peroxisome proliferator signalling

pathways converge through the direct interaction of their respective nuclear hormone receptors (Gearing et al., 1993; Issemann et al., 1993; Keller et al., 1993; Kliewer et al., 1992a; Miyata et al., 1994; Tugwood et al., 1992). Moreover, the orphan nuclear hormone receptors COUP-TF1 and HNF-4 bind to specific PPREs and differentially modulate PPAR function (Miyata et al., 1993; Winrow et al., 1994). Gene regulation by PPARs thus appears to be an overall transcriptional response achieved through a dynamic balance among several nuclear hormone receptors functioning as both positive and negative regulators of PPAR activity.

Thyroid hormone receptors (TRs) might be relevant to PPAR function under various physiological conditions. TRs mediate cellular responses to thyroid hormone which, like peroxisome proliferators, affects the expression of many genes important for lipid metabolism, cellular growth, and development. Peroxisome proliferators and thyroid hormones have overlapping metabolic effects, and there is apparent cross-talk between the thyroid hormone and peroxisome proliferator signalling pathways, as thyroid hormone has been shown to attenuate peroxisome proliferator-mediated transcriptional induction of genes encoding peroxisomal  $\beta$ -oxidation enzymes (Pacot et al., 1993; Takeda et al., 1992). Moreover, a connection between thyroid hormones and peroxisome proliferators is suggested by the finding that PPAR activators such as the fibrate hypolipidemic drugs promote adipose differentiation via a process modulated by thyroid hormone (Gharbi-Chihi et al., 1993).

Recent studies provide evidence that peroxisome proliferator and thyroid hormone signalling pathways converge at the level of their respective nuclear hormone receptors (Bogazzi et al., 1994; Jow and Mukherjee, 1995; Meier-Heusler et al., 1995). Thus, rat PPAR $\alpha$  has been shown to negatively regulate the expression of certain genes classically

considered to be thyroid hormone-dependent, either by heterodimerizing directly with particular TR subtypes to generate non-DNA-binding PPAR/TR heterodimers or by forming DNA-binding heterodimers that interact with specific thyroid hormone response elements (TREs). PPARs can also indirectly inhibit TR function by competing with TR for the common auxiliary co-regulator RXR. The converse effects of TR expression on PPAR-mediated activation of peroxisome proliferator-responsive genes have not been addressed. We have examined the effect of rat TR $\alpha$  on the transcriptional activation and DNA-binding properties of rat PPAR $\alpha$  (hereafter called TR $\alpha$  and PPAR, respectively). We demonstrate that TR $\alpha$ , on its own as well as cooperatively with RXR $\alpha$ , modulates the transactivation function of PPAR *in vivo* and binds to the AOX-PPRE *in vitro*. Our findings indicate that PPAR and TR signalling pathways converge in the regulation of peroxisome proliferator-responsive genes.

## 4.2 MATERIALS AND METHODS

### 4.2.1 Plasmids

The reporter plasmids pCPS $luc$  and pAOx(X2) $luc$  and the *in vivo* and *in vitro* expression plasmids for rat PPAR and human RXR $\alpha$  have been described in Chapter 2 (Marcus et al., 1993; Miyata et al., 1993; Zhang et al., 1993; Zhang et al., 1992). The mammalian expression vector for rat TR $\alpha$  (pRSV-TR $\alpha$ ) was obtained from V. Nikodem (National Institutes of Health, Bethesda) and contains the rTR $\alpha$  cDNA inserted as an *Hind*III/*Hpa*I fragment into pRSV (Bogazzi et al., 1994). The reporter plasmid pTREp $alluc$ ,



containing a high-affinity palindromic TRE, was obtained from C. Glass (University of California, San Diego)(Glass et al., 1988). The TR $\alpha$  *in vitro* transcription vector was constructed by excising the full-length TR $\alpha$  cDNA from pRSV-TR $\alpha$  and cloning it into pGEM-7Zf(+).

#### 4.2.2 *In Vitro Transcription/Translation*

Transcription/translation was performed using a coupled TNT rabbit reticulocyte lysate system (Promega). Labelled translation products were analyzed on 15% SDS-polyacrylamide gels. Synthesis of proteins for use in gel retardation analysis was carried out in parallel with unlabeled methionine.

#### 4.2.3 *Gel Retardation Analysis*

Gel retardation analysis was carried out with *in vitro* translated proteins (Miyata et al., 1993). Oligonucleotides corresponding to the AOx-PPRE (5'-CCTTTCCCGAACGTTGACCTTTGTCCTGGTCCCCTTTTGCT) and the DR4 element 5'-GATCTTCTTGACCTCCTGTGACCTGG were annealed to their respective complementary strands and end-labelled with the Klenow fragment of DNA polymerase I and [ $\alpha$ -<sup>32</sup>P]dATP. The underlined sequences highlight TGACCT-like direct repeats. One to 2  $\mu$ l of programmed lysate was incubated for 30 min at 30°C with 0.2 pmol of labelled probe in a final reaction volume of 15  $\mu$ l containing 7 mM HEPES (pH 7.9), 120 mM NaCl, 1 mM EDTA, 7% (vol/vol) glycerol, 4  $\mu$ g bovine serum albumin, 8  $\mu$ g of nonspecific competitor DNA (a 1:1 mixture of poly(dI-dC)·poly(dI-dC) and sonicated salmon sperm DNA), 150  $\mu$ M

phenylmethylsulfonyl fluoride, and 0.2 mM dithiothreitol. The total amount of lysate in each reaction was kept constant with the addition of unprogrammed lysate. For competition assays, unlabeled annealed oligonucleotides were added to the binding reactions along with labelled probe. For supershift analysis, 1  $\mu$ l of monoclonal antibodies specific for TR $\alpha$  or polyclonal anti-RXR $\alpha$  antibodies was added to the binding reactions prior to the addition of labelled probe. Protein/DNA complexes were analyzed by electrophoresis at 4°C on prerun 4% polyacrylamide gels (30:1 acrylamide:*N,N'*-methylenebisacrylamide weight ratio) with 22 mM Tris base/22 mM boric acid/1 mM EDTA as running buffer.

#### 4.2.4 *Transfection and Luciferase Assay*


All transfections were carried out with BSC40 monolayer cells by the calcium phosphate method (Miyata et al., 1993). BSC40 cells were chosen for transfection since they contain low endogenous PPAR $\alpha$  and RXR $\alpha$  activity. Cells were maintained in medium without phenol red and containing 5% charcoal-stripped fetal bovine serum for 24 h prior to and during transfection. Transfections (using 10 cm dishes of subconfluent cells) contained 5  $\mu$ g of reporter gene (pCPS*luc*, pAOx(X2)*luc*, or pTREp*alluc*), 0.5  $\mu$ g to 4  $\mu$ g of various combinations of receptor expression plasmids (as indicated in the figure legends) and 2  $\mu$ g of pCHI10 (Pharmacia), a  $\beta$ -galactosidase expression plasmid used as an internal reference. Effector plasmid dosage was kept constant by addition of appropriate amounts of the corresponding empty vectors. Total DNA was normalized to 20  $\mu$ g by the addition of salmon sperm DNA. Where indicated, the peroxisome proliferator Wy-14,643 or the TR ligand 3,3',5-triiodothyronine (T3; Sigma) was added to final concentrations of 0.1 mM and  $10^{-7}$  M,

respectively (from 100X stocks in dimethyl sulfoxide). Control cells received the equivalent volume of dimethyl sulfoxide. Cell extracts were prepared 48 h after transfection, and luciferase activity was measured. Unless indicated otherwise, values represent the average from at least 3 independent experiments carried out in duplicate and normalized to the internal  $\beta$ -galactosidase transfection control.

### 4.3 RESULTS

#### 4.3.1 *TR $\alpha$ and TR $\alpha$ /RXR $\alpha$ Heterodimers Bind to the AOx-PPRE*

TRs and PPARs bind to cognate response elements composed of direct repeats of the core consensus half site TGACCT, preferentially through heterodimerization with RXR (Tsai and O'Malley, 1994). TR $\alpha$ /RXR $\alpha$  heterodimers bind with high affinity to TGACCT direct repeats spaced by 4 nucleotides (referred to as DR4), whereas PPAR/RXR $\alpha$  heterodimers bind preferentially to direct repeat response elements configured as DR1, as found in both the AOx- and hydratase-dehydrogenase PPREs (Tugwood et al., 1992; Zhang et al., 1993). However, natural TREs are configured in various manners and orientations and can be recognized by TR monomers, homodimers, or heterodimers with RXR and other partners (Desvergne, 1994). To explore whether TR $\alpha$  recognizes the AOx-PPRE and/or modulates PPAR/RXR $\alpha$  protein/DNA interactions on this element, we carried out DNA binding assays with *in vitro* transcribed and translated receptors. TR $\alpha$ /RXR $\alpha$  heterodimers could bind cooperatively to a synthetic DR4 element (Fig. 4-1, compare lanes b and c to lane d). TR $\alpha$  could also bind to the DR4 element weakly as a monomer (Fig. 4-1, lane b). PPAR alone or




**Figure 4-1. Interaction of TR $\alpha$  with a synthetic DR4 element.** *In vitro* synthesized TR $\alpha$ , PPAR $\alpha$ , and RXR $\alpha$  (1  $\mu$ l each) were used individually or in pairwise combinations (as indicated at top) in gel retardation assays with a radiolabeled synthetic DR4 probe. Reactions were normalized with unprogrammed lysate as appropriate. The positions of the TR $\alpha$  monomeric complex and the TR $\alpha$ /RXR $\alpha$  heterodimeric complex are indicated. Lane a is a control in which the DR4 probe was incubated with unprogrammed lysate.

with RXR $\alpha$  or TR $\alpha$  did not form a complex on the DR4 element (Fig. 4-1, lanes e, f, and g, respectively).

Mobility shift experiments carried out with the AOx-PPRE are shown in Fig. 4-2. As previously demonstrated (Marcus et al., 1993), PPAR and RXR $\alpha$  did not bind to the AOx-PPRE individually but did bind cooperatively (Fig. 4-2, compare lanes b and d with lane e). Interestingly, TR $\alpha$  alone formed a complex with the AOx-PPRE (Fig. 4-2, lane c). The mobility of this complex was consistent with TR $\alpha$  binding to the AOx-PPRE as a monomer. Additional complexes were not generated when TR $\alpha$  and PPAR were coincubated, indicating that PPAR did not heterodimerize with TR $\alpha$  to bind to the AOx-PPRE (Fig. 4-2, lane g). However, in the presence of RXR $\alpha$ , TR $\alpha$  formed an additional complex on the AOx-PPRE that had a slightly faster electrophoretic mobility than that of the PPAR/RXR $\alpha$  complex (Fig. 4-2, lane f). The TR $\alpha$ /RXR $\alpha$ -dependent complex was generated at an approximately 5-fold lower efficiency *vis-à-vis* the PPAR/RXR $\alpha$  complex when equivalent amounts of the respective receptors were used in the binding assays (Fig. 4-2, compare lane f to lane e). Coincubation of equal amounts of PPAR, RXR $\alpha$  and TR $\alpha$  led to the formation of three distinct complexes, corresponding to PPAR/RXR $\alpha$  heterodimer, TR $\alpha$ /RXR $\alpha$  heterodimer, and TR $\alpha$  monomer complexes (lane h). Under these conditions, the TR $\alpha$ /RXR $\alpha$  and PPAR/RXR $\alpha$  complexes formed with approximately the same efficiency, indicating that TR $\alpha$  and PPAR compete for limiting amounts of RXR $\alpha$  (see below).

To determine if TR $\alpha$  was a component of these protein/DNA complexes, supershift analysis with antibodies to TR $\alpha$  and RXR $\alpha$  were carried out. Addition of anti-TR $\alpha$  antibodies to the binding reaction disrupted the formation of both the faster migrating TR $\alpha$ -dependent




**Figure 4-2. TR $\alpha$  binds to the AOx-PPRE on its own and in association with RXR $\alpha$ .** *In vitro* synthesized receptors (1  $\mu$ l each) were incubated individually or in various combinations (as indicated at top) with radiolabeled AOx-PPRE probe and analyzed by gel retardation as in Fig. 4-1. Reactions were normalized with unprogrammed lysate as appropriate. The positions of the monomeric TR $\alpha$  complex and the heterodimeric TR $\alpha$ /RXR $\alpha$  and PPAR $\alpha$ /RXR $\alpha$  complexes are indicated. The TR $\alpha$ /RXR $\alpha$  complex has a slightly faster electrophoretic mobility than that of the PPAR $\alpha$ /RXR $\alpha$  complex.

complex and the more slowly migrating TR $\alpha$ /RXR $\alpha$ -dependent complex (Fig. 4-3, compare lane c to lane d, and lane e to lane h). In contrast, anti-RXR $\alpha$  antibodies disrupted the formation of only the more slowly migrating complex (Fig. 4-3, compare lane e to lane g). Preimmune serum had no effect on complex formation (Fig. 4-3, lane f). These results show that TR $\alpha$  was present in both the faster migrating TR $\alpha$ -dependent complex and the more slowly migrating TR $\alpha$ /RXR $\alpha$ -dependent complex, whereas RXR $\alpha$  was only a component of the more slowly migrating complex.


Addition of the TR ligand T3 to a binding reaction containing TR $\alpha$  and RXR $\alpha$  resulted in a slightly increased mobility of the TR $\alpha$ /RXR $\alpha$  complex but had little effect on its intensity (Fig. 4-3, compare lane j to lane i). This altered mobility may be related to previous observations which demonstrated that T3 induces a ligand-dependent conformational change in TR/RXR heterodimers (Leng et al., 1993).

The TR $\alpha$ /RXR $\alpha$ -dependent complex was sequence-specific, since it could be competed with unlabeled AO $\alpha$ -PPRE probe DNA but not with a nonspecific competitor oligonucleotide (Fig. 4-4; compare lane d to lanes e-g and to lanes i-k, respectively). The TR $\alpha$  monomeric complex could also be competed with specific, but not with nonspecific, competitor DNA; however, this competition required at least 100- fold molar excess of competitor DNA compared to the 10- to 50-fold molar excess that was sufficient to disrupt the PPAR/RXR $\alpha$  and TR $\alpha$ /RXR $\alpha$  complexes. These results suggest that the TR $\alpha$  monomeric complex is more stable than the TR $\alpha$ /RXR $\alpha$  complex. Our findings demonstrate that the TR $\alpha$  binds in a sequence-specific manner to the AO $\alpha$ -PPRE on its own and as a heterodimer with RXR $\alpha$ .



**Figure 4-3. TR $\alpha$  is present in complexes formed on the AOx-PPRE.** Gel retardation assays using radiolabeled AOx-PPRE were carried out with *in vitro* synthesized receptors (1  $\mu$ l each) in the presence of preimmune serum (lane f), monoclonal anti-TR $\alpha$  antibodies (lanes d and h), or polyclonal anti-RXR $\alpha$  antibodies (lane g), as indicated. 1  $\mu$ l of preimmune serum or antibodies was added along with the indicated receptors prior to the addition of the AOx-PPRE probe. Reactions were normalized with unprogrammed lysate as appropriate. Inclusion of anti-TR $\alpha$  antibodies resulted in the loss of both the fast migrating and slowly migrating complexes (lanes d and h), while inclusion of anti-RXR $\alpha$  antibodies resulted in the loss of only the slowly migrating complex (lane g). Specific supershifted complexes produced by addition of anti-TR $\alpha$  and anti-RXR $\alpha$  antibodies were apparent after longer exposure. Lane a is a control in which the AOx-PPRE probe was incubated with unprogrammed lysate. Lanes i and j are identical in composition to lane e, except that lane j contained T3 ( $10^{-7}$  M final concentration from a stock in dimethyl sulfoxide), while lane i contained an equal volume of dimethyl sulfoxide alone. T3 led to a slight and reproducible increase in the electrophoretic mobility of the TR $\alpha$ /RXR $\alpha$  complex, as indicated by the arrowhead.







**Figure 4-4. TR $\alpha$  binds to the AOx-PPRE in a sequence-specific manner.** *In vitro* synthesized RXR $\alpha$ , TR $\alpha$ , and PPAR $\alpha$  (1  $\mu$ l each) were coincubated with radiolabeled AOx-PPRE alone (lanes d and h) and in the presence of 10-, 50- and 100-molar excess of unlabeled, double-stranded nonspecific competitor DNA (5'GATCCCTACCCATACGACGTCCCAGACTACGCTTGAGCT and its complement; lanes e-g, respectively) or 10-, 50- and 100-molar excess of unlabeled AOx-PPRE (lanes i-k, respectively) and analyzed by gel retardation. Lanes a and b are incubations carried out with combinations of *in vitro* translated PPAR $\alpha$  + RXR $\alpha$  and of TR $\alpha$  + RXR $\alpha$  (1  $\mu$ l of each), respectively, to indicate the positions of the heterodimeric complexes. Lane c is a control in which the AOx-PPRE probe was incubated with unprogrammed lysate.

#### 4.3.2 *TR $\alpha$ Competes with PPAR for Binding to the AOx-PPRE*

As indicated above (Fig.4-2, lane h), coincubation of the AOx-PPRE with equal amounts of TR $\alpha$ , PPAR, and RXR $\alpha$  resulted in the formation of three species of complex corresponding to the TR $\alpha$  monomeric complex, the PPAR/RXR $\alpha$  complex, and the TR $\alpha$ /RXR $\alpha$  complex. When constant amounts of PPAR and RXR $\alpha$  were coincubated with increasing amounts of TR $\alpha$ , the PPAR/RXR $\alpha$  complex gradually disappeared concomitantly with an increase in the intensity of the TR $\alpha$ /RXR $\alpha$  complex and the TR $\alpha$  monomeric complex (Fig. 4-5, lanes b-d). This result could be due to a competition of TR $\alpha$  for limiting amounts of RXR $\alpha$ , a competition of TR $\alpha$ /RXR $\alpha$  heterodimers for the AOx-PPRE binding site, and/or the formation of non-DNA-binding TR $\alpha$ /PPAR heterodimers. These possibilities are addressed in Fig. 4-6 in which binding reactions were carried out with increasing amounts of RXR $\alpha$  and/or PPAR while keeping the amount of TR $\alpha$  constant. Increasing the RXR $\alpha$  concentration led to an increase in the amount of TR $\alpha$ /RXR $\alpha$  complex concomitantly with a decrease in the amount of TR $\alpha$  monomeric complex. This result would suggest that RXR $\alpha$  was limiting for complex formation under these conditions (Fig. 4-6, compare lane a to lanes b and c). Incubation of TR $\alpha$  with increasing PPAR led to a decrease in the formation of the TR $\alpha$  monomeric complex, suggesting that PPAR can sequester TR $\alpha$  (Fig. 4-6, compare lane a to lanes d and e). Coincubation of TR $\alpha$  and PPAR with increasing amounts of RXR $\alpha$  led to a reduction in the amount of TR $\alpha$  monomeric complex concomitant with increased amounts of the PPAR/RXR $\alpha$  and TR $\alpha$ /RXR $\alpha$  complexes (Fig. 4-6, compare lane a to lanes f and g). These results suggest that TR $\alpha$  can inhibit binding of PPAR to the AOx-PPRE by competing for the common heterodimerization partner RXR $\alpha$ .



**Figure 4-5. TR $\alpha$  competes with PPAR $\alpha$ /RXR $\alpha$  for binding to the AO $\alpha$ -PPRE.** *In vitro* translated PPAR $\alpha$  and RXR $\alpha$  (2  $\mu$ l of each) were incubated in the absence (lane a) or presence of 2, 4, and 6  $\mu$ l of *in vitro* translated TR $\alpha$  (lanes b, c and d, respectively) and analyzed by gel retardation. Volumes of lysate were kept constant by the addition of unprogrammed lysate as appropriate. TR $\alpha$  inhibits the formation of the PPAR $\alpha$ /RXR $\alpha$  complex concomitant with the formation of the TR $\alpha$ /RXR $\alpha$  complex and the TR $\alpha$  complex.



**Figure 4-6.** Effects of increasing amounts of RXR $\alpha$  and PPAR $\alpha$  on the formation of TR $\alpha$ -dependent protein/DNA complexes. *In vitro* translated TR $\alpha$  (2  $\mu$ l) was mixed with RXR $\alpha$  (2  $\mu$ l, lane b; 5  $\mu$ l, lane c), PPAR $\alpha$  (2  $\mu$ l, lane d; 5  $\mu$ l, lane e), PPAR $\alpha$  (1  $\mu$ l) + RXR $\alpha$  (1  $\mu$ l) (lane f) or PPAR $\alpha$  (1  $\mu$ l) + RXR $\alpha$  (4  $\mu$ l) (lane g) in the presence of radiolabeled AOX-PPRE and analyzed by gel retardation. Lysate volumes were kept constant by the addition of unprogrammed lysate as appropriate.

and by binding to the AOx-PPRE on its own or cooperatively with RXR $\alpha$ . Moreover, the availability of PPAR/RXR $\alpha$  heterodimers capable of binding to the AOx-PPRE may also be restricted through the formation of non-PPRE-binding TR $\alpha$ /PPAR complexes.


#### 4.3.3 *Unliganded TR $\alpha$ Stimulates Transactivation by PPAR/RXR $\alpha$ Heterodimers*

Transient transfection assays were carried out in BSC40 cells to examine the effects of expression of TR $\alpha$  on PPAR/RXR $\alpha$ -dependent transactivation of a reporter gene plasmid containing two copies of the AOx-PPRE (pAOx(X2)*luc*). Control transfections showed the T3-dependent activation of a luciferase reporter gene plasmid containing a palindromic TRE (pTRE*pal**luc*) following cotransfection with the TR $\alpha$  expression vector (Fig. 4-7A). TR $\alpha$  functions as a ligand-dependent transactivator in BSC40 cells, resulting in a 10- to 12-fold T3-dependent induction of the TRE reporter gene construct. Expression from the parental reporter gene plasmid not containing a TRE (pSVL $\Delta$ 5'*luc*) was not affected by expression of TR $\alpha$  or by the presence of T3, as reported previously (Glass et al., 1988).

Fig. 4-7B shows the effects of transfecting increasing amounts of the TR $\alpha$  expression plasmid on transactivation of the AOx-PPRE reporter gene plasmid by PPAR and RXR $\alpha$ . Cotransfection of PPAR and RXR $\alpha$  expression plasmids resulted in an approximately 7- to 10-fold activation of expression of the AOx-PPRE reporter gene plasmid over basal levels in the absence of exogenously added PPAR activators, as previously demonstrated (Marcus et al., 1993). This activator-independent induction is presumably due to the presence of endogenous PPAR activators. Addition of the potent peroxisome proliferator Wy-14,643 to the transfections resulted in a 50-fold stimulation of activity of the AOx-PPRE reporter


**Figure 4-7. TR $\alpha$  enhances transactivation by PPAR $\alpha$ /RXR $\alpha$  *in vivo*.** (A) BSC40 cells were cotransfected with the reporter plasmids pTREpalluc or pSVL $\Delta$ 5' luc and the TR $\alpha$  expression plasmid pRSV-TR $\alpha$  in the presence or absence of T3, as indicated. Luciferase assays were performed as indicated in **Materials and Methods**. The values reported are the averages ( $\pm$  SEM) of 3 independent transfections carried out in duplicate and normalized to the value obtained with pTREpalluc alone (taken as 1). (B) BSC40 cells were transfected with pAO $\alpha$ (X2) luc in the presence or absence of the peroxisome proliferator Wy-14,643, along with a constant amount of PPAR $\alpha$  and RXR $\alpha$  expression plasmids (2  $\mu$ g each) and increasing amounts of the TR $\alpha$  expression plasmid (in  $\mu$ g), as indicated. Plasmid dosage was normalized in each case with appropriate amounts of the corresponding empty expression vectors. The values reported are the averages ( $\pm$  SEM) of a minimum of 3 independent transfections carried out in duplicate and normalized to the value obtained from Wy-14,643-treated cells cotransfected with PPAR $\alpha$  and RXR $\alpha$  expression plasmids (taken as 100%). The results show that TR $\alpha$  potentiates Wy-14,643-mediated transactivation by PPAR $\alpha$ /RXR $\alpha$ .

**A**



pTREpalluc  
pSVLΔ5luc  
TRα

**B**




PPARα + RXRα  
TRα

gene plasmid over basal levels. Surprisingly, cotransfection with increasing amounts of the TR $\alpha$  expression plasmid resulted in progressively enhanced stimulation by PPAR/RXR $\alpha$ . Cotransfection of up to 4  $\mu$ g of the TR $\alpha$  expression plasmid (representing a 2-fold molar excess over PPAR/RXR $\alpha$ ) resulted in a 100- to 200-fold stimulation of Wy-14,643-dependent, PPAR/RXR $\alpha$ -mediated transactivation of the AOx-PPRE reporter gene plasmid over basal levels. TR $\alpha$  also stimulated the Wy-14,643-independent induction by PPAR/RXR $\alpha$ , although to a lesser extent (40- to 50-fold stimulation over basal levels compared to 10- to 20-fold stimulation in the absence of TR $\alpha$ ).

Transfection of the PPAR expression vector alone resulted in a modest (10-fold) stimulation of AOx-PPRE reporter gene activity in the presence of Wy-14,643, likely because of interaction between PPAR and endogenous RXR $\alpha$  and/or other cofactors (Fig. 4-8). However, TR $\alpha$  did not further stimulate PPAR-mediated activation under these conditions. Indeed, activation was reduced by approximately 50% when a 2-fold excess of TR $\alpha$  expression plasmid was transfected. Similarly, the slight activation of AOx-PPRE reporter gene expression mediated by RXR $\alpha$  alone (2- to 4-fold) was not appreciably affected by cotransfection of the TR $\alpha$  expression vector. Interestingly, TR $\alpha$  on its own stimulated reporter gene expression approximately 10-fold, but only when transfected at relatively high amounts. However, this TR $\alpha$ -mediated induction was not further stimulated by Wy-14,643, as compared to control transfections.

To examine the requirements of a PPRE for TR $\alpha$ -dependent stimulation, cotransfection experiments similar to those reported above were carried out with pCPS*luc*, the parental vector used to construct pAOx(X2)*luc*. Transfection of PPAR/RXR $\alpha$  alone, or







**Figure 4-8. TR $\alpha$ -mediated stimulation of Wy-14,643-dependent activation requires the presence of PPAR $\alpha$  and RXR $\alpha$ .** The TR $\alpha$  expression plasmid pRSV-TR $\alpha$  (in  $\mu$ g) was cotransfected with pAOx(X2)*luc* alone or in combination with plasmids expressing PPAR $\alpha$  (2  $\mu$ g) or RXR $\alpha$  (2  $\mu$ g) in the presence or absence of Wy-14,643, as indicated. The values reported are the averages ( $\pm$  SEM) of three independent transfections carried out in duplicate and normalized to the value obtained for cells transfected with the pAOx(X2)*luc* reporter plasmid alone (taken as 1).


in combination with TR $\alpha$ , had little effect on the expression of this reporter gene construct, either in the absence or presence of Wy-14,643 (Fig. 4-9A). Therefore, TR $\alpha$ -mediated stimulation of Wy-14,643-dependent transcriptional induction requires both exogenously added PPAR and RXR $\alpha$  and a functional PPRE.

In contrast to the stimulation of transactivation of the AOx-PPRE by PPAR/RXR $\alpha$  observed with unliganded TR $\alpha$ , addition of T3 to the transfections resulted in an almost complete repression of both Wy-14,643-independent transcriptional induction (Fig. 4-9B) and Wy-14,643-dependent transcriptional induction (Fig. 4-10). Repression was manifested through TR $\alpha$ , since T3 alone had no effect on basal level expression of the AOx-PPRE reporter gene plasmid or on transactivation by PPAR/RXR $\alpha$  (Fig. 4-9B). However, T3-mediated, TR $\alpha$ -dependent repression was independent of PPAR/RXR $\alpha$  or on the presence of a PPRE, since inhibition of basal level activity was also observed with the parental reporter plasmid pCPS $luc$  (Fig. 4-9A). These results suggest that liganded TR $\alpha$  has a generalized repressive effect on transcription, perhaps because it sequesters common co-activators required for basal level and/or activated transcription of the minimal promoters used (see **DISCUSSION**). This would be consistent with the observation that liganded TR $\alpha$  reduced activity below the basal levels obtained from cells transfected with pAOx(X2) $luc$  or pCPS $luc$  alone.

Overall, our findings suggest that TR $\alpha$  in the unliganded state is a positive modulator of PPAR/RXR $\alpha$ -mediated transactivation of the AOx-PPRE, despite the fact that *in vitro*, TR $\alpha$  appears to inhibit the binding of PPAR/RXR $\alpha$  heterodimers to this response element.

**Figure 4-9. TR $\alpha$  inhibits transactivation in the presence of T3.** (A) BSC40 cells were cotransfected with the reporter gene pCPS*luc* and plasmids (2  $\mu$ g each) expressing the indicated receptors in the presence or absence of Wy-14,643 and T3, as indicated. The values reported are the averages of duplicate transfections and are normalized to the value obtained with pCPS*luc* cotransfected with PPAR $\alpha$  and RXR $\alpha$  expression plasmids in the presence of Wy-14,643 (taken as 100%). Readings of duplicate samples did not vary by more than 15%. (B) BSC40 cells were cotransfected with pAOx(X2)*luc*, PPAR $\alpha$  and RXR $\alpha$  expression plasmids (2  $\mu$ g each), and increasing amounts of the TR $\alpha$  expression plasmid pRSV-TR $\alpha$  in the presence or absence of T3, as indicated. The values reported are the averages ( $\pm$  SEM) from 3 independent transfections carried out in duplicate and are normalized to the value obtained for cells transfected with the reporter gene pAOx(X2)*luc* alone (taken as 1).

**A****B**



**Figure 4-10. Liganded TR $\alpha$  also inhibits Wy-14,643-mediated transactivation by PPAR $\alpha$ /RXR $\alpha$ .** Transient co-transfections were performed with 5  $\mu$ g of pAOx(X2)luc, 2  $\mu$ g each of PPAR $\alpha$  and RXR $\alpha$ , and 2  $\mu$ g of TR $\alpha$  expression plasmids, as indicated. Transfection media was supplemented with Wy-14,643 alone or with both Wy-14,643 and T3, as shown in the figure legend. The values presented represent the average of duplicate transfections, and are normalized to the value obtained with PPAR $\alpha$ /RXR $\alpha$  in the presence of Wy-14,643 (taken as 100%).

#### 4.4 DISCUSSION

Recent findings from several groups have indicated that PPAR can differentially influence TR-mediated activation of thyroid-hormone responsive genes (Bogazzi et al., 1994; Jow and Mukherjee, 1995; Meier-Heusler et al., 1995). Conversely, our findings show that TR $\alpha$  modulates PPAR-mediated transactivation of peroxisome proliferator-responsive genes *in vivo* and can disrupt the binding of PPAR/RXR $\alpha$  heterodimers to PPREs. Thus, cross-talk and coupling between the thyroid hormone and peroxisome proliferator signalling pathways are important to the reciprocal regulation of both thyroid hormone- and peroxisome proliferator-responsive genes.

We have shown that TR $\alpha$  is capable of binding to the AOx-PPRE in a sequence-specific manner on its own as well as through association with RXR $\alpha$ . This is not necessarily unexpected, since thyroid receptors bind promiscuously as monomers or as homo- and heterodimers to a wide variety of structurally diverse response elements configured in different manners (Desvergne, 1994). Therefore, the AOx-PPRE defines a novel target for TR $\alpha$ . This finding, coupled with our previous observations that COUP-TF1 (Miyata et al., 1993) and HNF-4 (Chapter 2) also bind to the AOx-PPRE, provides further evidence that the AOx-PPRE is a composite response element that is a target for multiple members of the nuclear hormone receptor superfamily.

The binding of TR $\alpha$  and of TR $\alpha$ /RXR $\alpha$  to the AOx-PPRE led to a decrease in the binding of PPAR/RXR $\alpha$  to this element. Our findings suggest that this decrease was the net result of TR $\alpha$  competing for the common heterodimerization partner RXR $\alpha$  and for binding to the AOx-PPRE. Moreover, as shown by others, PPAR and TR $\alpha$  can heterodimerize in

solution, indicating that TR $\alpha$  can sequester PPAR as well as RXR $\alpha$  (Bogazzi et al., 1994). However, the TR $\alpha$ /PPAR heterodimer was unable to bind to the AOx-PPRE, and a natural DNA binding site for the TR $\alpha$ /PPAR heterodimer pair, if it exists, remains to be identified. Recently, the TR $\beta$ /PPAR heterodimer has been shown to be capable of binding to a DR2 element present in the promoter region of the gene encoding myelin proteolipid protein (Bogazzi et al., 1994).

Paradoxically, cotransfection of PPAR/RXR $\alpha$  with the TR $\alpha$  expression vector led to a significant Wy-14,643-mediated stimulation of transcription of the AOx-PPRE reporter gene over that seen with PPAR/RXR $\alpha$  alone. This stimulation required the presence of both RXR $\alpha$  and PPAR and a functioning PPRE in the reporter gene plasmid. In the absence of exogenously added RXR $\alpha$ , TR $\alpha$  led to a reduction in PPAR/Wy-14,643-dependent stimulation. TR $\alpha$  on its own gave a slight stimulation of AOx-PPRE reporter gene expression; however, this was negated by exogenously added RXR $\alpha$  or PPAR. Neither TR $\alpha$  alone nor in combination with RXR $\alpha$  conferred responsiveness of the AOx-PPRE to Wy-14,643 *in vivo*. These findings suggest that TR $\alpha$  cooperates with RXR $\alpha$  and PPAR to stimulate Wy-14,643-dependent transactivation of the AOx-PPRE.

The modest transcriptional activation of the AOx-PPRE observed with TR $\alpha$  on its own is inconsistent with the reports that activation by TRs is normally hormone dependent. Indeed, unliganded TRs typically repress the basal activity of promoters that contain TREs through active silencing or inhibition of formation of the preinitiation complex (Fondell et al., 1993). Nevertheless, examples of ligand-independent transactivation by TR $\alpha$  have been reported (Forman et al., 1988; Saatcioglu et al., 1993), and therefore it remains possible that

DNA-bound TR $\alpha$  directly contributes to the overall enhanced stimulation by PPAR/RXR $\alpha$ .

The results showing that unliganded TR $\alpha$  augments PPAR/RXR $\alpha$ -mediated transactivation of the AOx-PPRE *in vivo* appear to be at odds with the *in vitro* results showing that TR $\alpha$  inhibits the binding of PPAR/RXR $\alpha$  to the AOx-PPRE. It is possible that TR $\alpha$  may not bind to the AOx-PPRE *in vivo*, perhaps because PPAR/RXR $\alpha$  binds more avidly to the AOx-PPRE or because there exist additional cellular factors that stabilize PPAR/RXR $\alpha$  interactions on the AOx-PPRE. TR $\alpha$ -dependent stimulation of PPAR/RXR $\alpha$  activity *in vivo* may thus be manifested through mechanisms that do not involve the binding of TR $\alpha$  to the AOx-PPRE. For instance, it is possible that TR $\alpha$  titrates, and therefore counteracts, the effects of a repressor that attenuates transcriptional activation by PPAR/RXR $\alpha$  heterodimers, thereby relieving repression. Consistent with this possibility, unliganded TR $\alpha$  has been shown to bind to an inhibitory factor that also interacts with the retinoic acid receptor, RAR $\alpha$  (Casanova et al., 1994). Therefore, overexpression of TR $\alpha$  may lead to an alteration in the repertoire of RXR heterodimers present in the cell, which could effectively increase the amount of RXR $\alpha$  available for interaction with PPAR.

Interestingly, the addition of T3 led to strong TR $\alpha$ -dependent inhibition of transcription of the AOx-PPRE both in the presence and absence of Wy-14,643. This finding may be relevant to the fact that administration of thyroid hormone can antagonize activation of the AOx gene by peroxisome proliferators (Yamada et al., 1994). However, transcriptional repression by liganded TR $\alpha$  appeared to be indirect, since repression was also observed in the absence of PPAR/RXR $\alpha$  or with the reporter gene construct pCPS $luc$ , which does not contain a PPRE. Transcriptional repression by liganded TR $\alpha$  may be due to its ability, but not that



of unliganded TR $\alpha$ , to squelch transcription by sequestering general auxiliary transcription factors.

Negative transcriptional effects of liganded TR $\alpha$  resulting from squelching do not necessarily exclude liganded TR $\alpha$  from having a more direct negative effect on PPAR/RXR $\alpha$  activation. TRs have complex and diverse effects on transcriptional control, since they can act as both positive and negative regulators of gene expression depending on the particular response element, the heterodimerization partner, and the absence or presence of ligand. TR $\alpha$ /RXR $\alpha$  heterodimers undergo a ligand-dependent conformational change (Bendik and Pfahl, 1995; Leng et al., 1993). It is interesting to note that the addition of T3 led to a slight, but reproducible, altered electrophoretic mobility of the TR $\alpha$ /RXR $\alpha$  complex on the AOX-PPRE, suggesting a ligand-dependent conformational change on this element. Therefore, it remains possible that TR $\alpha$ /RXR $\alpha$  heterodimers bound to the AOX-PPRE *in vivo* can actively repress transcription in the presence of ligand.

In summary, we have established a link between PPARs and TRs that may be physiologically important in the regulation of lipid homeostasis and in mediating the thyromimetic effects of peroxisome proliferators. The mechanisms by which TR $\alpha$  differentially modulates transactivation by PPARs, and *vice versa*, are complex and appear to be dependent upon a dynamic balance between RXR $\alpha$ , PPAR, TR $\alpha$ , and possibly other cellular factors and nuclear hormone receptors. Elucidation of these pathways will provide a clearer understanding of the mechanism underlying the pleiotropic cellular effects mediated by PPARs.

## CHAPTER 5

### **Cross-talk Between Orphan Nuclear Hormone Receptor RZR $\alpha$ and Peroxisome Proliferator-Activated Receptor $\alpha$ in Regulation of the Peroxisomal Hydratase-Dehydrogenase Gene <sup>†</sup>**

<sup>†</sup> A version of this chapter has been published. 1998. Winrow, C.J., Capone, J.P. and Rachubinski, R.A. *J. Biol. Chem.* **273**:31442-31448. Used with permission from The American Society for Biochemistry and Molecular Biology.

## 5.1 ABSTRACT

The genes encoding the peroxisomal  $\beta$ -oxidation enzymes enoyl-CoA hydratase/3-hydroxyacyl-CoA dehydrogenase (HD) and fatty acyl-CoA oxidase (AOx) are coordinately regulated by peroxisome proliferator-activated receptor  $\alpha$  (PPAR $\alpha$ )/9-*cis* retinoic acid receptor (RXR $\alpha$ ) heterodimers that transactivate these genes in a ligand-dependent manner via upstream peroxisome proliferator-response elements (PPRE). Here we demonstrate that the monomeric orphan nuclear hormone receptor, RZR $\alpha$ , modulates PPAR $\alpha$ /RXR $\alpha$ -dependent transactivation in a response-element dependent manner. Electrophoretic mobility shift analysis showed that RZR $\alpha$  bound specifically as a monomer to the HD-PPRE but not the AOx-PPRE. Determinants in the HD-PPRE for binding of RZR $\alpha$  were distinct from those required for interaction with PPAR $\alpha$ /RXR $\alpha$  heterodimers. In transient transfections, RZR $\alpha$  stimulated ligand-mediated transactivation by PPAR $\alpha$  from an HD-PPRE luciferase reporter in the absence of exogenously added RXR $\alpha$ , but did not affect PPAR $\alpha$ -dependent transactivation of an AOx-PPRE reporter gene. These data illustrate cross-talk between the RZR $\alpha$  and PPAR $\alpha$  signaling pathways at the level of the HD-PPRE in the regulation of the HD gene and characterize additional factors governing the regulation of peroxisomal  $\beta$ -oxidation.

## 5.2 INTRODUCTION

Nuclear hormone receptors are a diverse group of structurally related ligand-activated transcription factors that direct the expression of target genes in response to physiological and environmental stimuli (Mangelsdorf and Evans, 1995; Tsai and O'Malley, 1994). Peroxisome proliferator-activated receptors (PPAR) are members of the steroid/nuclear hormone receptor superfamily that act to regulate a large number of genes involved in differentiation and lipid metabolism (Juge-Aubry et al., 1997; Latruffe and Vamecq, 1997; Spiegelman and Flier, 1996; Tontonoz et al., 1995; Tontonoz et al., 1994b; Tugwood et al., 1992; Zhang et al., 1992) in response to a variety of compounds collectively called peroxisome proliferators. Peroxisome proliferators include the fibrate family of hypolipidemic drugs, phthalate ester plasticizers, herbicides, pesticides, antidiabetic thiazolidinediones, as well as natural and synthetic fatty acids (Forman et al., 1997b; Kliewer et al., 1995; Kliewer et al., 1997; Lambe and Tugwood, 1996; Lehmann et al., 1995; Lock et al., 1989). Transactivation of target genes by PPARs is mediated through binding to *cis*-acting regulatory sequences called peroxisome proliferator-response elements (PPRE) that consist of direct repeats of the hexameric TGACCT/C core motif. PPARs heterodimerize with the 9-*cis* retinoic acid receptor, RXR $\alpha$ , and bind with preference to response elements with spacing of one nucleotide between hexameric repeats (DR1) (Chu et al., 1995a; Keller et al., 1993; Kliewer et al., 1992b; Marcus et al., 1993). PPREs have been identified in the regulatory regions of a number of genes, including those encoding the first two enzymes of the peroxisomal  $\beta$ -oxidation pathway: fatty acyl-CoA oxidase (AOx) (Osumi et al., 1991; Tugwood et al., 1992) and enoyl-CoA hydratase/3-hydroxyacyl-CoA dehydrogenase (HD) (Bardot et al., 1993;

Zhang et al., 1992).

In addition to PPARs, several other nuclear hormone receptors have been shown to bind to PPREs and differentially modulate PPAR function. These include chicken ovalbumin upstream promoter transcription factor (COUP-TF) (Miyata et al., 1993), hepatocyte nuclear factor-4 (HNF-4) (Chapter 2) and thyroid hormone receptor (TR) (Chapter 4). Transcriptional regulation via PPREs is thus a net aggregate response manifested in part by the availability of PPARs and other factors that bind to PPREs, the complexity of response elements, and the interplay of PPARs with other nuclear hormone receptors and cofactors. Overlying this network is a series of co-repressors and co-activators that serve to mediate receptor signaling. The overall intricacy of the system enables the integration of information from multiple signaling pathways to ensure appropriate transcriptional responses of target genes to various stimuli.

The retinoid Z receptor (RZR) family, also known as the retinoid orphan receptor (ROR) family, is a recently described group of nuclear hormone receptors shown to be involved in regulating genes responsible for cellular differentiation, the inflammatory response, and lipid metabolism (Carlberg et al., 1994; Giguere et al., 1994; Hirose et al., 1994; Vu-Dac et al., 1997). RZR/ROR target response elements have been identified in the promoters of genes for chicken  $\gamma$ F-crystallin, human and rat bone sialo protein, human 5-lipoxygenase, N-myc proto-oncogene, and apolipoprotein AI (Dussault and Giguere, 1997; Schrader et al., 1996; Steinhilber et al., 1995; Tini et al., 1995; Vu-Dac et al., 1997). Three RZR/ROR isoforms have been characterized ( $\alpha$ ,  $\beta$ ,  $\gamma$ ) that show differential tissue localization. The  $\alpha$  isoform includes four splice variants (ROR $\alpha$ 1, ROR $\alpha$ 2, ROR $\alpha$ 3 and RZR $\alpha$ ) and is

widely expressed, RZR $\beta$  is found primarily in brain tissues, and ROR $\gamma$  is localized to skeletal muscle (Becker-Andre et al., 1993; Carlberg et al., 1994; Hirose et al., 1994; Missbach et al., 1996). While nuclear hormone receptors typically bind response elements as hetero- or homodimers, RZR/ROR family members are distinct in that they are able to bind to, and transactivate from, target response elements as monomers (Carlberg et al., 1994; Giguere et al., 1994). RZR/ROR receptors recognize a TGACCT/C consensus half-site and have a demonstrated preference for sites flanked by a 6 base A/T-rich region (Carlberg et al., 1994; Giguere et al., 1995; Giguere et al., 1994; Schrader et al., 1996). This consensus half site is embedded in the HD-PPRE, and recently RZR $\alpha$  was shown to bind to this element, suggesting that RZR $\alpha$  might be a candidate regulator of peroxisomal  $\beta$ -oxidation (Schrader et al., 1996). To explore this possibility, I undertook to determine how RZR $\alpha$  might interact with PPAR $\alpha$  and RXR $\alpha$  in regulating the transcription of genes encoding the first two enzymes of the peroxisomal  $\beta$ -oxidation pathway, AOX and HD. The results presented here indicate that RZR $\alpha$  binds to the HD-PPRE but not the AOX-PPRE, and selectively potentiates transactivation from the HD-PPRE in a manner dependent on the relative availability of RXR $\alpha$ .

### 5.3 MATERIALS AND METHODS

#### 5.3.1 Plasmids

The *in vitro* and *in vivo* expression plasmids for rat PPAR $\alpha$  and human RXR $\alpha$ , and the luciferase reporter constructs pCPS $luc$  containing the minimal promoter for the gene

encoding carbamoyl phosphate synthetase, pHD(X3)*luc* containing two copies of the minimal HD-PPRE, and pAOx(X2)*luc* containing two copies of the minimal AOx-PPRE, have been described previously in Chapter 2 (Marcus et al., 1993; Zhang et al., 1993; Zhang et al., 1992). The plasmid pM2(X3)*luc*, constructed by cloning the mutant HD-PPRE synthetic oligonucleotide M2 (5'-GATCCTCTCCTTTAAAATATTGAACTATTACCTACATTTGA) and its complement into the *Bam*HI site of pCPS*luc*, contains three direct tandem copies of the M2 element. The expression plasmid for human RZR $\alpha$  (RZR $\alpha$ /SG5) (Schrader et al., 1996) was a kind gift of Carsten Carlberg (Hôpital Cantonal Universitaire de Genève, Geneva, Switzerland).

### 5.3.2 *In Vitro* Transcription/Translation

Transcription/translation of cDNAs encoding PPAR $\alpha$ , RXR $\alpha$  and RZR $\alpha$  was performed using the TNT T7 coupled rabbit reticulocyte lysate system according to the manufacturer's protocol (Promega). Translation products labeled with *L*-[<sup>35</sup>S]methionine were analyzed on 15% SDS-polyacrylamide gels. Synthesis of proteins for use in electrophoretic mobility shift analysis (EMSA) was carried out in parallel with unlabeled methionine.

### 5.3.3 *Electrophoretic Mobility Shift Analysis*

EMSA was carried out as previously described in Chapter 2 (Miyata et al., 1993; Zhang et al., 1993) with oligonucleotide probes corresponding to the HD-PPRE (5'-CCTCTCCTTTGACCTATTGAACTATTACCTACATTTGA and its complement) and the

AOx-PPRE (5'-CCTTTCCCGAACGTGACCTTTGTCCTGGTCCCCTTTTGCT and its complement). Underlined sequences indicate TGACCT-like direct repeats. Complementary oligonucleotides were annealed and end-labeled with the Klenow fragment of DNA polymerase I and [ $\alpha$ - $^{32}$ P]dATP. 1 to 2  $\mu$ l of programmed lysate was incubated with 0.2 pmol of labeled probe at 25°C in a final volume of 15  $\mu$ l containing 6 mM Hepes (pH 7.9), 120 mM NaCl, 0.4 mM MgCl<sub>2</sub>, 0.1 mM EDTA, 7% (v/v) glycerol, 4  $\mu$ g bovine serum albumin, 4  $\mu$ g nonspecific competitor DNA (poly dI•dC and sonicated salmon sperm DNA, 1:1 weight ratio), 150  $\mu$ M phenylmethylsulfonyl fluoride and 0.2 mM dithiothreitol. The total amount of reticulocyte lysate in each reaction was kept constant by addition of unprogrammed lysate. Binding reactions were analyzed by electrophoresis at 4°C on prerun 3.5% polyacrylamide gels (30:1 acrylamide/*N,N'*-methylenebisacrylamide weight ratio) with 22 mM Tris base/22 mM boric acid/1 mM EDTA as running buffer, followed by autoradiography.

#### 5.3.4 *Transient Transfections and Measurement of Luciferase Activity*

Transfections were performed in African green monkey kidney BSC40 monolayer cells by the calcium phosphate method (Zhang et al., 1992). Cells at 60-80% confluence were transfected in medium without phenol red and supplemented with 5% (v/v) charcoal-stripped fetal bovine serum. Transfections were carried out with 5  $\mu$ g of luciferase reporter construct (pCPS*luc*, pHD(X3)*luc*, pM2(X3)*luc* or pAOx(X2)*luc*), 2  $\mu$ g of each of RXR $\alpha$  and PPAR $\alpha$  expression plasmids, and 0.5-4  $\mu$ g of RZR $\alpha$  expression plasmid per 10 cm plate, as indicated. 2  $\mu$ g of the  $\beta$ -galactosidase expression plasmid pCH110 (Pharmacia) was included in transfections as an internal reference. Effector plasmid dosage was kept constant by addition



of corresponding amounts of empty expression plasmids, and total DNA was maintained at 20  $\mu$ g per 10 cm plate by addition of sonicated salmon sperm DNA. The peroxisome proliferator Wy-14,643 or the putative RZR $\alpha$  ligand melatonin (both in dimethyl sulfoxide) was added to fresh medium to a final concentration of 0.1 mM and 1  $\mu$ M, respectively, where indicated. An equivalent volume of dimethyl sulfoxide was added to control medium. Cells were harvested and lysates prepared 48 h post-transfection, and luciferase activity was quantitated as previously described in Chapter 2 (Zhang et al., 1992).

## 5.4 RESULTS

### 5.4.1 RZR $\alpha$ binds as a monomer to the HD-PPRE but not the AOx-PPRE

Several potential RZR $\alpha$  consensus binding sites exist within the HD- and AOx-PPREs (Fig. 5-1), and interaction of RZR $\alpha$  with the HD-PPRE has recently been observed (Schrader et al., 1996). To explore how RZR $\alpha$  might cooperate with PPAR $\alpha$  and RXR $\alpha$  to regulate transactivation from PPRES, we first examined the DNA-binding characteristics of these nuclear hormone receptors to the HD- and AOx-PPREs. EMSA was performed with radiolabelled HD- and AOx-PPREs and *in vitro* translated receptors. PPAR $\alpha$  and RXR $\alpha$  formed a characteristic heterodimer that bound strongly to the HD-PPRE (Fig. 5-2A). RZR $\alpha$  also bound to the HD-PPRE, apparently as a monomer based on a comparison of the mobility of the RZR $\alpha$  complex with that of the heterodimeric PPAR $\alpha$ /RXR $\alpha$  complex. Coincubation of RZR $\alpha$  with either PPAR $\alpha$  or RXR $\alpha$  alone did not yield heterodimeric complexes. When all three receptors were present in the binding reaction, only heterodimeric PPAR $\alpha$ /RXR $\alpha$  and

**AO<sub>x</sub>-PPRE**


  
 CCGAACGTGACCTTTGTCCTGGTCCC

**HD-PPRE**



  
 CCTCTCCTTTGACCTATTGAACTATTACCTACATT

**RZR $\alpha$  Consensus Site**


TGACC(T/C)(A/T)<sub>6</sub>

**Figure 5-1.** Comparison of the sequence of the RZR $\alpha$  consensus binding site to sequences within the AO<sub>x</sub>- and HD-PPREs. Arrows and roman numerals indicate the locations and directions of TGACCT-like motifs.

A



B




**Figure 5-2. RZR $\alpha$  binds as a monomer to the HD-PPRE but not the AOx-PPRE.** *In vitro* synthesized PPAR $\alpha$ , RXR $\alpha$  and RZR $\alpha$  (1  $\mu$ l each) were incubated alone or in combination with radiolabeled HD-PPRE (A) or AOx-PPRE (B) probe, as indicated. Lysate volumes were kept constant by addition of unprogrammed lysate. EMSA was performed as described under "Materials and Methods". The positions of the heterodimeric PPAR $\alpha$ /RXR $\alpha$  and monomeric RZR $\alpha$  complexes are indicated by *arrows*.

monomeric RZR $\alpha$  complexes were observed, suggesting that these receptors do not co-occupy the HD-PPRE under these conditions. Therefore, RZR $\alpha$  binds as a monomer to the HD-PPRE and does not form complexes with PPAR $\alpha$  or RXR $\alpha$  on the HD-PPRE *in vitro*. PPAR $\alpha$ /RXR $\alpha$  heterodimers readily formed a complex on the AOX-PPRE, as expected; however, RZR $\alpha$  was unable to bind to this element either alone or in combination with PPAR $\alpha$  or RXR $\alpha$  (Fig. 5-2B).

Competition analysis demonstrated that the interaction of RZR $\alpha$  with the HD-PPRE was specific. As shown in Fig. 5-3, both the RZR $\alpha$  and the PPAR $\alpha$ /RXR $\alpha$  complexes were refractory to competition by nonspecific unlabeled oligonucleotide (compare *lanes d-f* to *lanes a-c, g*), whereas increasing amounts of unlabeled HD-PPRE oligonucleotide effectively competed for binding of both RZR $\alpha$  and PPAR $\alpha$ /RXR $\alpha$  (compare *lanes h-j* to *lanes a-c, g*). These results show that binding of RZR $\alpha$  is specific for the HD-PPRE.

We next examined whether PPAR $\alpha$  might modulate binding of RZR $\alpha$  to the HD-PPRE. EMSA was performed with radiolabelled HD-PPRE probe, constant amounts of RXR $\alpha$  and RZR $\alpha$ , and decreasing amounts of PPAR $\alpha$ . Dilution of PPAR $\alpha$  resulted in progressively decreased amounts of the PPAR $\alpha$ /RXR $\alpha$  complex, as expected; however, there was no effect on formation of the RZR $\alpha$  monomer complex (Fig. 5-4A), suggesting that RZR $\alpha$  binding to the HD-PPRE *in vitro* is not influenced by PPAR $\alpha$ . Similarly, increasing amounts of RZR $\alpha$  resulted in correspondingly greater amounts of the RZR $\alpha$  monomeric complex, but no change in the PPAR $\alpha$ /RXR $\alpha$  complex (Fig. 5-4B). Additional analyses carried out using different amounts of radiolabelled HD-PPRE probe and titration of either RZR $\alpha$  (Fig. 5-5A) or PPAR $\alpha$ /RXR $\alpha$  (Fig. 5-5B) showed that PPAR $\alpha$ /RXR $\alpha$  and RZR $\alpha$  did




**Figure 5-3. RZR $\alpha$  monomer binds to the HD-PPRE in a sequence-specific manner.** EMSA was performed with *in vitro* synthesized receptors (1  $\mu$ l each) and radiolabeled HD-PPRE probe, as shown. Volumes were kept constant by addition of unprogrammed lysate. Where indicated, unlabeled competitor DNA (5'-GATCCCGTGCATGCTAATGATATTCT and its complement) (*Nonspecific*) or unlabeled HD-PPRE (*Specific*) was present in 10-, 50- or 100-fold molar excess. The positions of the heterodimeric PPAR $\alpha$ /RXR $\alpha$  and monomeric RZR $\alpha$  complexes are indicated by *arrows*.

**Figure 5-4. PPAR $\alpha$ /RXR $\alpha$  and RZR $\alpha$  bind independently to the HD-PPRE.** (A) *In vitro* synthesized RXR $\alpha$  and RZR $\alpha$  (1  $\mu$ l each) were incubated with 1  $\mu$ l of PPAR $\alpha$  or increasing dilutions of PPAR $\alpha$  in unprogrammed lysate, as shown, prior to addition of radiolabeled HD-PPRE probe. Complexes were analyzed by EMSA. The positions of PPAR $\alpha$ /RXR $\alpha$  heterodimer and RZR $\alpha$  monomer complexes are indicated by *arrows*. (B) Radiolabeled HD-PPRE probe was incubated in the presence of constant amounts of *in vitro* synthesized PPAR $\alpha$  and RXR $\alpha$  (1  $\mu$ l each) and in the absence or presence of 1, 2 and 5  $\mu$ l of *in vitro* synthesized RZR $\alpha$ . Lysate volumes were kept constant by addition of unprogrammed lysate. The positions of PPAR $\alpha$ /RXR $\alpha$  heterodimer and RZR $\alpha$  monomer complexes are indicated by *arrows*.



**A**

PPAR $\alpha$	-	+	-	+	1:10	1:50	1:100	1:250	1:500	1:1000
RXR $\alpha$	-	+	-	+	+	+	+	+	+	+
RZR $\alpha$	-	-	+	+	+	+	+	+	+	+




**B**

RZR $\alpha$		▲		
PPAR $\alpha$ /RXR $\alpha$	+	+	+	+

not affect each other's binding to the HD-PPRE. Therefore, RZR $\alpha$  and PPAR $\alpha$ /RXR $\alpha$  bind independently to the HD-PPRE.

#### 5.4.2 *RZR $\alpha$ and PPAR $\alpha$ /RXR $\alpha$ bind to different regions of the HD-PPRE in vitro*

The above results suggest that RZR $\alpha$  recognizes distinct determinants on the HD-PPRE compared to the requirements for PPAR $\alpha$ /RXR $\alpha$  interaction. The HD-PPRE is a complex response element that consists of 4 consensus TGACCT hexameric half sites (sites I-IV; Fig. 5-1). To determine which elements are responsible for RZR $\alpha$  binding specificity, oligonucleotide probes containing mutations in each of the four hexameric half sites were used in binding studies (Fig. 5-6). EMSA analysis showed that mutations in sites I, III, or IV did not adversely affect binding of RZR $\alpha$ , whereas disruption of site II eliminated binding (Fig. 5-6). This result is consistent with the fact that site II most closely matches the RZR $\alpha$  consensus sequence (TGACCT/C/(A/T)<sub>6</sub>). By comparison, the integrity of sites III and IV were essential for PPAR $\alpha$ /RXR $\alpha$  interaction, whereas sites I and II were dispensable (Fig. 5-6). This latter result is consistent with previous observations that the downstream DR1 element is necessary and sufficient for PPAR $\alpha$ /RXR $\alpha$  signaling in the cell (Zhang et al., 1993). Thus, RZR $\alpha$  and PPAR $\alpha$ /RXR $\alpha$  binding specificity is mediated by distinct and non-overlapping determinants in the HD-PPRE.


#### 5.4.3 *RZR $\alpha$ potentiates transactivation from the HD-PPRE by PPAR $\alpha$*

To define a cellular role for the observed *in vitro* interaction of RZR $\alpha$  with the HD-PPRE, a series of transient transfections was performed with the HD-PPRE luciferase




**Figure 5-5. RZR $\alpha$  and PPAR $\alpha$ /RXR $\alpha$  do not affect each other's binding to the HD-PPRE.** Binding reactions contained 2  $\mu$ l of *in vitro* synthesized PPAR $\alpha$ /RXR $\alpha$  and 0  $\mu$ l, 2  $\mu$ l or 5  $\mu$ l of *in vitro* synthesized RZR $\alpha$  (*Panel A*); or 1  $\mu$ l of *in vitro* synthesized RZR $\alpha$  and 0  $\mu$ l, 4  $\mu$ l or 10  $\mu$ l of *in vitro* synthesized PPAR $\alpha$ /RXR $\alpha$  (*Panel B*). Radiolabeled HD-PPRE probe was added to 200 fmol, 2 fmol or 0.2 fmol, as indicated. Lysate volumes were kept constant by addition of unprogrammed lysate. EMSA was performed as described in "Materials and Methods". Autoradiography was for 24 h (200 fmol) or 17 d (2 and 0.2 fmol).

A




B



**Figure 5-6. RZR $\alpha$  monomer and PPAR $\alpha$ /RXR $\alpha$  heterodimer bind to different sites within the HD-PPRE *in vitro*.** (A) Sequences of the wild-type HD-PPRE and of the oligonucleotides *M1*, *M2*, *M3* and *M4* containing mutations in consensus hexameric binding sites I, II, III and IV (*bold lettering*) of the HD-PPRE. *Open lettering*, sequences of mutant sites I, II, III and IV. (B) *In vitro* synthesized PPAR $\alpha$ , RXR $\alpha$  and RZR $\alpha$  (1  $\mu$ l each) were incubated with radiolabeled double-stranded oligonucleotides corresponding to mutations M1-M4. Lysate volumes were kept constant by addition of unprogrammed lysate. EMSA was performed as described under "Materials and Methods". *Arrows* show the positions of bound complexes.

**A**

HD-PPRE 5' GATCCTCTCCTTTGACCTATTGA<sup>I</sup>ACTATTACCTACATTG<sup>II</sup>A  
 M1 5' GATCCTATAATTTGACCTATTGA<sup>III</sup>ACTATTACCTACATTG<sup>IV</sup>A  
 M2 5' GATCCTCTCCTTTAAAAATTGA<sup>I</sup>ACTATTACCTACATTG<sup>II</sup>A  
 M3 5' GATCCTCTCCTTTGACCTATTGA<sup>III</sup>AGTATTACCTACATTG<sup>IV</sup>A  
 M4 5' GATCCTCTCCTTTGACCTATTGAACTAATCTTCACATTG<sup>I</sup>A




**B**


reporter plasmid, pHD(X3)*luc*, in BSC40 African monkey kidney cells. This cell line was chosen because it contains low levels of endogenous PPAR $\alpha$  and RXR $\alpha$  activity (Miyata et al., 1993; Zhang et al., 1992). Transfection of a PPAR $\alpha$  expression plasmid resulted in a 2-fold increase in luciferase activity over control transfections containing the reporter plasmid alone (Fig. 5-7A). Treatment of PPAR $\alpha$ -expressing cells with the peroxisome proliferator and PPAR $\alpha$  ligand, Wy-14,643, increased luciferase activity a further 4-fold. Transfection of increasing amounts of RXR $\alpha$  expression plasmid in the presence of fixed amounts of PPAR $\alpha$  expression plasmid led to a dose-dependent increase in the level of luciferase activity in cells treated with Wy-14,643, but did not affect the ligand-independent response. At the maximal level of RXR $\alpha$  plasmid used (4  $\mu$ g), ligand-dependent PPAR $\alpha$  activity was 23 times the basal level of activity of control cells transfected with reporter plasmid alone (and compared to the 8-fold PPAR $\alpha$ , ligand-dependent induction observed in the absence of RXR $\alpha$  expression plasmid). RXR $\alpha$  expression plasmid alone showed at most a 1.5-fold increase in luciferase activity over basal levels of the HD-PPRE luciferase reporter (Fig. 5-8), in agreement with previous studies that demonstrated that RXR $\alpha$  only weakly activates transcription via this element (Schrader et al., 1996). The modest increase observed reflects cooperation of RXR $\alpha$  with the HD-PPRE, as RXR $\alpha$  had no effect on basal expression levels of the parental pCPS*luc* reporter gene, which lacks a PPRE (see below). Importantly, functional cooperativity between PPAR $\alpha$  and RXR $\alpha$  in transactivation from the HD-PPRE is abrogated in the reporter construct pM2(X3)*luc* (Fig. 5-7B), in which site II of the HD-PPRE required for binding of RXR $\alpha$  has been mutated. These results show that binding of RXR $\alpha$  to the HD-PPRE is apparently required to achieve transcriptional

**Figure 5-7. RZR $\alpha$  potentiates transactivation by PPAR $\alpha$  from the HD-PPRE but not the AOx-PPRE.** BSC40 monolayer cells were transfected with 5  $\mu$ g of the luciferase reporter pHD(X3)*luc* (Panel A), pM2(X3)*luc* (Panel B) or pAOx(X2)*luc* (Panel C), and expression plasmids for PPAR $\alpha$  (2  $\mu$ g) and RZR $\alpha$  (0.5-4  $\mu$ g), in the absence or presence of 0.1 mM Wy-14,643. Plasmid dosage was normalized by addition of the appropriate empty vectors, where required. Cells were harvested 48 h post-transfection, and luciferase activity was quantitated, as described in Chapter 2. All values represent the average of three independent experiments carried out in duplicate. Values from individual transfections did not vary by more than 15%. The values are presented relative to the value obtained for cells transfected with PPAR $\alpha$  in the presence of Wy-14,643 (taken as 100%).

**A****HD-PPRE**

132

**B****HD(M2)-PPRE****C****AOX-PPRE**



**Figure 5-8. RZRα alone weakly activates transcription from the HD-PPRE.** Transient co-transfections were performed in BSC40 monolayer cells with 5 μg of pHD(X3)*luc* reporter plasmid, and expression plasmids for PPARα and RXRα (2 μg each) or RZRα (4 μg). Transfection media contained 0.1 mM of the peroxisome proliferator Wy-14,643 (in DMSO) or an equivalent volume of vehicle (DMSO). The data presented represent the average of three independent transfections performed in duplicate and values for individual transfections did not vary by more than 15%. The values are shown relative to the value obtained for cells transfected with PPARα/RXRα in the presence of Wy-14,643 (taken as 100%).




stimulation by PPAR $\alpha$ .

The above experiments demonstrate that RZR $\alpha$  cooperates with PPAR $\alpha$  to potentiate transactivation from the HD-PPRE in response to treatment with a peroxisome proliferator. In contrast, RZR $\alpha$  did not significantly affect PPAR $\alpha$ -dependent transactivation from the AO $\alpha$ -PPRE, in either the absence or presence of ligand (Fig. 5-7C). These data are in keeping with the *in vitro* binding data presented above, and together the results demonstrate that RZR $\alpha$  binds specifically to the HD-PPRE and modulates transactivation by PPAR $\alpha$  from this response element.

We next investigated whether melatonin, which has been suggested to be a putative ligand for RZR $\alpha$  (Carlberg and Wiesenberg, 1995; Steinhilber et al., 1995), could influence the functional cooperation between PPAR $\alpha$  and RZR $\alpha$  in stimulating transcription from the HD-PPRE. Under our experimental conditions, we were unable to observe any effect of melatonin on the functional cooperativity of these two receptors in stimulating transcription from the HD-PPRE (Fig. 5-9).




#### *5.4.4 Exogenous RXR $\alpha$ abrogates the potentiation effect of RZR $\alpha$ on transactivation by PPAR $\alpha$ from the HD-PPRE*

RXR $\alpha$  is an obligate heterodimerization partner for PPAR $\alpha$ , and the PPAR $\alpha$ -dependent activation presented above presumably arises from cooperation with low levels of endogenous RXR $\alpha$  present in BSC40 cells. We therefore examined the effects of RZR $\alpha$  under conditions where RXR $\alpha$  expression plasmid was included in the transfections. Coexpression of exogenous RXR $\alpha$  and PPAR $\alpha$  increased luciferase activity 14- and 10-fold



**Figure 5-9. Melatonin does not stimulate transactivation by PPAR $\alpha$ /RZR $\alpha$  from the HD-PPRE.** BSC40 cells were transfected with 5  $\mu$ g of the luciferase reporter pHD(X3)*luc* and the expression plasmids for PPAR $\alpha$  (2  $\mu$ g) and RZR $\alpha$  (4  $\mu$ g), with addition of vehicle (dimethyl sulfoxide) (*open bars*), 0.1 mM Wy-14,643 (*hatched bars*) or 1  $\mu$ M melatonin (*dashed bars*). Plasmid dosage was normalized by the addition of the appropriate empty vectors, where required. Cells were harvested 48 h post-transfection, and luciferase activity was quantitated. All values represent the average of three independent experiments carried out in duplicate. Values from individual transfections did not vary by more than 15%. The values are presented relative to the value obtained for cells transfected with PPAR $\alpha$  in the presence of Wy-14,643 (taken as 100%).

**Figure 5-10. RZR $\alpha$  does not potentiate transactivation by PPAR $\alpha$  from the HD-PPRE in the presence of exogenous RXR $\alpha$ .** BSC40 cells were transfected with plasmids expressing PPAR $\alpha$  (2  $\mu$ g), RXR $\alpha$  (2  $\mu$ g) and RZR $\alpha$  (0.5-4  $\mu$ g), and 5  $\mu$ g of luciferase reporter construct. Values are shown for cells transfected with either (A) pHD(X3)*luc* or (B) pAOx(X2)*luc* reporter plasmids. The values are the average of three independent transfections performed in duplicate. Values from individual transfections did not vary more than 15%. The values presented were relative to the value obtained for cells transfected with PPAR $\alpha$  and RXR $\alpha$  in the presence of Wy-14,643 (taken as 100%). (C) Transfection of BSC40 cells with the reporter plasmid pCPS*luc* in the presence or absence of PPAR $\alpha$  (2  $\mu$ g), RXR $\alpha$  (2  $\mu$ g) and RZR $\alpha$  (0.5-4  $\mu$ g), as indicated. Values represent the average of duplicate transfections.



over basal levels for the HD-PPRE (Fig. 5-10A) and AOx-PPRE (Fig. 5-10B) luciferase reporter plasmids, respectively. In the presence of Wy-14,643, activity was increased 60-fold for the HD-PPRE reporter (Fig. 5-10A) and 20-fold (Fig. 5-10B) for the AOx-PPRE reporter. The robust activity obtained with coexpressed RXR $\alpha$  is consistent with the fact that the endogenous level of this receptor is limiting for PPAR $\alpha$ -mediated transactivation. Interestingly, under these conditions, transactivation from the HD- and AOx-PPRE reporter plasmids was not significantly affected by addition of RZR $\alpha$  (Figs. 5-10A and B, respectively). Control transfections with the parental construct pCPS*luc*, which lacks a PPRE, showed that the presence of PPAR $\alpha$ , RXR $\alpha$ , and RZR $\alpha$  did not influence the levels of relative luciferase activity observed (Fig. 5-10C), indicating that these receptors did not alter the basal activity of the parental reporter construct. Together, these data suggest that the stimulatory effect observed with RZR $\alpha$  on transactivation by PPAR $\alpha$  is attenuated by increasing the level of RXR $\alpha$ .

## 5.5 DISCUSSION

Several observations have suggested a degree of interplay between the RZR/ROR and PPAR nuclear hormone receptor families in the regulation of genes, particularly those encoding proteins involved in lipid metabolism. For instance, both PPAR $\alpha$  and RZR $\alpha$  regulate transcription of the apolipoprotein AI gene (Vu-Dac et al., 1997; Vu-Dac et al., 1994). PPAR $\gamma$  has been shown to be a critical regulator of the adipogenic program, and ROR $\alpha$  and ROR $\gamma$  mRNAs are induced early in adipogenesis (Austin et al., 1998). Finally, the antidiabetic thiazolidinediones, which are potent activators of PPAR $\gamma$ , have recently been

shown to also be specific ligands for RZR $\alpha$  (Lehmann et al., 1995; Missbach et al., 1996).

Both the HD and AOx genes are regulated by a number of nuclear hormone receptors from several different signaling networks that converge on the respective PPREs or that directly modulate PPAR $\alpha$  activity (Miyata et al., 1996; Miyata et al., 1993; Winrow et al., 1996; Winrow et al., 1994). Consistent with these observations, Schröder *et al.* reported that RZR $\alpha$  weakly interacted with the HD-PPRE and was able to minimally activate transcription via this element (Schröder et al., 1996). Our results extend this finding to show that RZR $\alpha$  can strongly potentiate transactivation by PPAR $\alpha$  from the HD-PPRE when RXR $\alpha$  levels are limiting. We found that the monomeric binding of RZR $\alpha$  to the HD-PPRE was specific and required for potentiation of transactivation by PPAR $\alpha$  from the HD-PPRE, and that RZR $\alpha$  did not bind to, or stimulate PPAR $\alpha$ -dependent transactivation from, the AOx-PPRE.

The mechanism by which RZR $\alpha$  stimulates PPAR $\alpha$  activity on the HD-PPRE is not known. EMSA analysis did not show any obvious antagonizing or stabilizing effects between RZR $\alpha$  monomers and PPAR $\alpha$ /RXR $\alpha$  heterodimers on the HD-PPRE *in vitro*. This result agrees with our findings that the RZR $\alpha$  monomer and PPAR $\alpha$ /RXR $\alpha$  heterodimer target distinct and non-overlapping hexameric determinants for binding to the HD-PPRE. Moreover, we did not observe a higher order ternary complex containing PPAR $\alpha$ /RXR $\alpha$ /RZR $\alpha$  on the HD-PPRE. However, this does not preclude the possibility of such a complex forming *in vivo*, which may be dependent on cooperativity or interaction with auxiliary cofactors. The involvement of auxiliary factors such as SRC-1, p300 and N-COR in transcriptional regulation by PPARs and other nuclear hormone receptors is well established (DiRenzo et al., 1997; Dowell et al., 1997b; Zhu et al., 1996). Additionally,

PPAR $\alpha$ , RXR $\alpha$  and/or RZR $\alpha$  might require phosphorylation or other modification not provided in the *in vitro* transcription/translation system. We are currently investigating whether auxiliary factors and receptor post-translational modification affect cooperative transcriptional regulation by PPAR $\alpha$ , RXR $\alpha$  and RZR $\alpha$ .

The HD-PPRE is a complex response element and among a select few that contain 4 hexameric direct repeats. In the HD-PPRE, these hexameric half sites are organized in two tandem DR1 arrays that are separated by 2 nucleotides, an arrangement that is thought to facilitate diverse receptor interactions and thereby permit multiple levels of control. A model has been proposed in which either one or two PPAR $\alpha$ /RXR $\alpha$  heterodimers bind to the HD-PPRE to determine the state of transcriptional activation (Chu et al., 1995a). As we have shown previously, the 3' DR1 (sites III/IV) array is essential and sufficient for PPAR $\alpha$ /RXR $\alpha$  binding and activity (Zhang et al., 1993); however, the arrangement that has been suggested to yield the highest level of transactivation has PPAR $\alpha$ /RXR $\alpha$  heterodimers bound to both DR1 sites. Based on our observations *in vitro*, it is likely that under certain *in vivo* conditions, such as limited availability of RXR $\alpha$ , a single PPAR $\alpha$ /RXR $\alpha$  heterodimer may bind to the second DR1 element (sites III/IV), yielding a lower overall level of transactivation compared with the situation in which two PPAR $\alpha$ /RXR $\alpha$  bind to this PPRE. Since RZR $\alpha$  occupies site II within the HD-PPRE, this may preclude binding of PPAR $\alpha$ /RXR $\alpha$  to the first DR1 element and thereby favor binding of the heterodimer to the transcriptionally competent 3' DR1 array. RZR $\alpha$  may also contribute directly to transcriptional responses by stabilizing the PPAR $\alpha$ /RXR $\alpha$  complex *in vivo* or facilitating interactions with an auxiliary factor(s). This pathway of stimulation may be operative when RXR $\alpha$  is present in limiting amounts, for

example in transfections carried out with PPAR $\alpha$  alone, as RZR $\alpha$  had no effect when transfections were carried out in the presence of excess exogenous RXR $\alpha$ . Cotransfection of RXR $\alpha$  significantly increased the overall level of transactivation by PPAR $\alpha$ , as one would expect if endogenous RXR $\alpha$  is limiting. It is possible that this level of activity is beyond a threshold level at which RZR $\alpha$  might be expected to have a stimulatory effect. Moreover, when RXR $\alpha$  is present in excess *in vivo*, PPAR $\alpha$ /RXR $\alpha$  heterodimers may occupy both DR I s, thereby resulting in a maximal transcriptional response and preventing RZR $\alpha$  from accessing site II. This model suggests ligand-mediated regulation of the HD-PPRE is dependent on input and dynamic interplay among the PPAR $\alpha$ , RXR $\alpha$  and RZR $\alpha$  signaling pathways.

In summary, these findings demonstrate that the orphan receptor RZR $\alpha$  can work in cooperation with PPAR $\alpha$  to regulate expression of the gene encoding HD, the second enzyme of the peroxisomal  $\beta$ -oxidation pathway. Transcriptional regulation of peroxisomal  $\beta$ -oxidation has proven to be a dynamic process, integrating cues from a host of signaling pathways and rapidly responding to variations in levels of key components. An understanding of the principal regulators of peroxisomal  $\beta$ -oxidation, including transcription factors controlling and modulating the expression of the genes encoding the enzymes of this pathway, may provide for the development of pharmacologic agents that specifically target the peroxisomal  $\beta$ -oxidation system as a means to influence overall lipid metabolism and homeostasis.



## **CHAPTER 6**

### **GENERAL DISCUSSION**

## 6.1 Discussion

Currently the most prevalent diseases in our society are directly related to the control of lipid metabolism and homeostasis. Fatty acids serve not only as energy stores, but also as critical signalling molecules, altering gene expression, and eliciting widespread effects on metabolism, cell growth and differentiation (for reviews see Clarke and Jump, 1994; Jump et al., 1997). The influences of dietary fat have been examined in the initiation and progression of numerous chronic diseases including cancer, insulin resistance, hyperlipidemia, cardiovascular disease, obesity and hypertension (Hallaq et al., 1992; Pan et al., 1994; Phillipson et al., 1985; Storlien et al., 1991; Welsch, 1992). A better understanding of the core regulatory mechanisms which control fatty acid metabolism will provide additional targets for genetic screening and will direct the discovery of pharmacological agents which can be employed to ameliorate a variety of disease states. Nuclear receptors are critical regulators of fatty acid metabolism and provide excellent targets for pharmacological intervention since the majority are ligand-modulated and since they serve as specific activators and/or repressors of target genes. Members of the PPAR family of nuclear receptors were initially identified as regulators of peroxisomal  $\beta$ -oxidation and time has shown that these factors play an exceedingly important part in mounting the necessary responses to fluctuating fatty acid levels and maintaining the balance between lipid storage and metabolism. My research has focussed on the identification of factors which modify the activity of PPARs in response to peroxisome proliferators, with a particular eye towards the transcriptional regulation of enzymes of the peroxisomal fatty acid  $\beta$ -oxidation pathway. The results I have presented clearly demonstrate the role of several factors in the PPAR-mediated transactivation

of the rat AOX and HD genes which can be extended to other species, and add to the general understanding of nuclear receptor signalling and the part played by additional signalling molecules.

#### *6.1.1 Influence of HNF-4 on the AOX- and HD-PPREs*

HNF-4 regulates genes involved in lipid metabolism, exhibits a similar tissue distribution as PPAR $\alpha$  and recognizes response elements with DR1 spacing (Dreyer et al., 1992; Fraser et al., 1998; Gregori et al., 1998; Hall et al., 1995; Jiang et al., 1995; Murao et al., 1997; Sladek et al., 1990; Stoffel and Duncan, 1997). In our work with HNF-4, we observed that HNF-4 interacted readily with the AOX-PPRE and to a lesser extent with the HD-PPRE. We found that HNF-4 repressed PPAR $\alpha$  activation of the AOX- and HD-PPRE in the absence of a peroxisome proliferator. Interestingly, when a peroxisome proliferator was included in transient co-transfections, HNF-4 repressed the AOX-PPRE and potentiated the activation of the HD-PPRE by PPAR $\alpha$ . A recent study has confirmed our findings regarding the inhibition of PPAR $\alpha$ -mediated transactivation of the AOX gene by HNF-4 through direct competition for binding to the AOX-PPRE (Nishiyama et al., 1998). Our demonstration that HNF-4 differentially regulates the PPAR $\alpha$ -dependent transactivation of the AOX- and HD-PPRE was the first example that HNF-4 can function as both a transcriptional activator and repressor of lipid metabolism through a common response element. This research identifies HNF-4 as an important transcriptional regulator of peroxisomal  $\beta$ -oxidation and shows the interplay between the HNF and PPAR signalling networks.

The observation that HNF-4 represses Wy-14,643- independent transactivation of the AOx- and HD-PPREs and Wy-14,643-dependent transactivation of the AOx-PPRE can be accounted for by competition for DNA binding site occupancy by HNF-4 homodimers. This repression may also involve the formation of non-DNA binding HNF-4 heterodimers and may occur through interactions between the ligand binding domain-dimerization interfaces in the absence of DNA binding domain-dimerization interface interactions. Although this is possible, it may not be the case, since HNF-4 has only been reported to function as a homodimer (Jiang et al., 1995).

Since our initial examination of the interactions between HNF-4 and PPAR $\alpha$ /RXR $\alpha$  signalling, a number of studies have been performed which have investigated the cross-talk between these nuclear receptor families. HNF-4 and PPAR interact as both activators and as repressors of co-regulated genes (Hertz et al., 1995; Hertz et al., 1996b; Murao et al., 1997; Nishiyama et al., 1998; Peters et al., 1997b; Rodriguez et al., 1994; Rodriguez et al., 1998). Response elements in the promoters of the long terminal repeat of HIV-1, the hepatitis B virus nucleocapsid gene, phosphoenolpyruvate carboxykinase, fatty acid binding protein and cellular retinol binding protein genes, are all synergistically activated through the co-operation of HNF-4 and PPARs (Hall et al., 1995; Kliewer et al., 1992b; Ladas, 1994; Poirier et al., 1997; Raney et al., 1997; Stoffel and Duncan, 1997; Tontonoz et al., 1995). In addition to AOx and HD, HNF-4 and PPAR $\alpha$  have been subsequently found to differentially regulate expression of the liver transferrin and apolipoprotein CIII genes and the hepatitis B virus pre-C promoter (Fraser et al., 1998; Hertz et al., 1995; Hertz et al., 1996b; Yu and Mertz, 1997).

Early studies of the regulation of the apolipoprotein CIII gene showed that although

HNF-4 served as a transcriptional activator of this gene, fibrate peroxisome proliferators suppressed apolipoprotein CIII gene transcription (Fraser et al., 1998; Hertz et al., 1996a; Hertz et al., 1995). It was determined that PPAR $\alpha$  competes with HNF-4 for binding to apolipoprotein CIII response elements and displaces the transactivating HNF-4 (Hertz et al., 1996a; Hertz et al., 1995). In addition, peroxisome proliferators repress expression of the HNF-4 gene (Hertz et al., 1996b). The interplay between HNF-4 and PPAR $\alpha$ /RXR $\alpha$  in the regulation of the apolipoprotein CIII gene is thought to partially account for the hypolipidemic effects of peroxisome proliferators (Hertz et al., 1995; Hertz et al., 1996b). Apolipoprotein CIII inhibits the catabolism of triglyceride-rich lipoproteins by the inhibition of lipoprotein lipase and by blocking liver receptor mediated uptake (Hertz et al., 1996a; Hertz et al., 1995). Therefore, peroxisome proliferator inhibition of apolipoprotein CIII expression coupled with increases in  $\beta$ -oxidation might account for an overall decrease in plasma lipid levels (Hertz et al., 1996a; Hertz et al., 1995).

HNF-4 represses PPAR $\alpha$  transactivation of the mitochondrial HMG-CoA synthase gene, illustrating another example of differential control by these nuclear receptors (Rodriguez et al., 1994; Rodriguez et al., 1998). These observations reiterate the importance of the interplay between the HNF-4 and PPAR signalling pathways which we first described in the differential regulation of the AOx- and HD-PPREs. Prior to our investigation, HNF-4 had been shown to stimulate expression of mitochondrial medium chain acyl-CoA dehydrogenase, which is a key enzyme of the mitochondrial  $\beta$ -oxidation pathway (Carter et al., 1993). Our results indicate that HNF-4 represses the transcription of the peroxisomal  $\beta$ -oxidation AOx and HD genes in the absence of a peroxisome proliferator. Taken together,

these findings suggest that HNF-4 may serve as a molecular switch to direct activation of mitochondrial  $\beta$ -oxidation, while down regulating peroxisomal  $\beta$ -oxidation. It appears, however, that these interactions may be more complex than first thought, since PPAR $\alpha$  has also been shown to activate the mitochondrial medium chain acyl-CoA dehydrogenase gene in response to peroxisome proliferators (Gulick et al., 1994).

Recent evidence supports the role of cross-talk between PPAR and HNF-4 receptors on a number of fronts. Hypolipidemic peroxisome proliferators reduce the expression of the HNF-4 gene, and this may prove effective in modulating cross-talk between the PPAR and HNF-4 pathways (Hertz et al., 1996b). Interplay between these receptors is not unique to the AOx and HD genes, since HNF-4 and PPAR $\alpha$  differentially regulate expression of the apolipoprotein CIII, HMG-CoA synthase and liver transferrin genes and the hepatitis B virus pre-C promoter (Hertz et al., 1996a; Hertz et al., 1996b; Rodriguez et al., 1998; Yu and Mertz, 1997).

Perhaps the most striking indication of the convergence of the PPAR and HNF-4 signalling networks is that fatty acyl-CoA thioesters with chain lengths greater than C12 act as ligands for HNF-4 (Hertz et al., 1998). Saturated C14 or C16 fatty acyl-CoAs serve as agonists, potentiating HNF-4 homodimerization, DNA binding and gene activation. C18 and polyunsaturated fatty acyl-CoAs act as antagonists by increasing the formation of non-DNA binding HNF-4 homodimers, resulting in a significant decrease in transcription of target genes. The fatty acid chain length and degree of saturation appear to determine whether the transcriptional effects will be activating or repressive (Hertz et al., 1998). PPARs are activated by mono- and polyunsaturated long chain free fatty acids, including those with chain

lengths greater than C16 (Bentley et al., 1993; Dreyer et al., 1993; Forman et al., 1997a; Kliewer et al., 1997; Krey et al., 1997; Willson and Wahli, 1997). These findings imply a mechanism for synergistic and antagonistic cross-talk between PPARs and HNF-4 nuclear receptors at the level of ligand binding and availability (Hertz et al., 1998). In addition, peroxisomal  $\beta$ -oxidation preferentially metabolizes very long chain fatty acids, with chain lengths between C18-C30 (Hiltunen et al., 1996; Mannaerts and van Veldhoven, 1996; Mannaerts and Van Veldhoven, 1993; Reddy and Lalwai, 1983).

Coupled with our results, these findings implicate HNF-4 as an extremely important mediator of fatty acid metabolism, interacting with the peroxisomal proliferator signalling pathway to direct either mitochondrial or peroxisomal  $\beta$ -oxidation of fatty acids. This holds implications for HNF-4 and PPAR $\alpha$  interplay in the monitoring of physiological lipid levels and stimulation of fatty acid metabolism via  $\beta$ -oxidation. It will be interesting to determine more specifically how the PPAR and HNF-4 signalling pathways converge, and to characterize these interactions within the context of a whole animal. Given the discovery of an increasing number of genes that are co-regulated by HNF-4 and PPAR $\alpha$ , the interplay between these nuclear receptors will likely become more apparent in other critical processes such as mitochondrial  $\omega$ -hydroxylation, the inflammatory response, and the activation of viral genes (Aldridge et al., 1995; Huss and Kasper, 1998; Ladas, 1994; Muerhoff et al., 1992; Peters et al., 1997b; Raney et al., 1997; Ueda et al., 1998; Yokomori et al., 1997; Yu and Mertz, 1997).

### 6.1.2 Calreticulin and PPAR $\alpha$ /RXR $\alpha$ activity

Calreticulin is a Ca<sup>2+</sup>-binding protein which exerts effects on cell adhesion, gene expression, and signal transduction, and functions as an endoplasmic reticulum chaperone responsible for protein folding and processing (Coppolino and Dedhar, 1998; Krause and Michalak, 1997; Mery et al., 1996; Opas et al., 1996; Vassilakos et al., 1998). Based on initial observations with glucocorticoid, retinoic acid, vitamin D3 and androgen receptors, a number of groups had proposed that calreticulin might generally regulate nuclear receptor activity through interactions with the conserved KXXFF(K/R)R motifs in their DNA binding domains (Burns et al., 1994; Dedhar et al., 1994; St-Arnaud et al., 1995; Wheeler et al., 1995). The research that we have performed demonstrated that although calreticulin could reduce the *in vitro* DNA binding of COUP-TF, HNF-4 and PPAR $\alpha$ /RXR $\alpha$ , the effects of calreticulin overexpression on PPAR $\alpha$ /RXR $\alpha$ -mediated gene activation were negligible in a cellular context. We also observed repression of GR-mediated gene expression as has been previously reported. (Burns et al., 1994) Our experiments were the first demonstration that calreticulin is not a global regulator of nuclear receptor transactivation, and these findings have been strengthened in light of more recent findings.

In order to directly affect nuclear receptor function through physical interactions with the KXXFF(K/R)R sites within the nuclear receptor DNA-binding domains, calreticulin would need to be present outside of the endoplasmic reticulum. It had been originally reported that calreticulin is found in several locations including the cell surface, within the secretory pathway, in the cytoplasm and within the nucleus (Dedhar et al., 1994; Dupuis et al., 1993; Michalak et al., 1992; White et al., 1995). Since that time, the localization of



calreticulin outside of the endoplasmic reticulum has been questioned (Krause and Michalak, 1997; Michalak et al., 1996). Using a combination of separately derived antibodies towards native calreticulin as well as recombinant dystrophin- and green fluorescent protein-epitope tagged calreticulin, Opas concluded that calreticulin is localized exclusively to the endoplasmic reticulum network and is not a constitutive nuclear protein (reviewed in Krause and Michalak, 1997). Extensive immunohistological work in several cell types and with a number of calreticulin antibodies, and with both intact and fractionated nuclei has been conducted (Michalak et al., 1996). The conclusion of these studies was that calreticulin is not present as a nuclear protein. Use of a cytoplasmically targeted calreticulin mutant showed that only overexpression of the native endoplasmic reticulum-resident calreticulin was able to affect GR-mediated gene expression (Michalak et al., 1996). In addition, when GR is bound to DNA, calreticulin is unable to compete for GR binding, suggesting that the DNA-bound nuclear form of GR does not interact with calreticulin (Michalak et al., 1996). These findings imply that calreticulin is not interacting directly with the KXXFF(K/R)R motifs within the DNA-binding sites of nuclear receptors in the nucleus and thereby disrupting gene activation (Krause and Michalak, 1997; Michalak et al., 1996). Due to the nature of these experiments, however, one cannot completely exclude the possibility that calreticulin is present transiently in the nucleus. The weight of this evidence suggests that the effects of calreticulin on gene expression are occurring through other means, possibly by endoplasmic reticulum signalling or through alteration of  $\text{Ca}^{2+}$ -signalling (Krause and Michalak, 1997; Opas et al., 1996).

It is known that calreticulin can interact with factors within the endoplasmic reticulum and may be modulating nuclear receptor activity through associations with endoplasmic

reticulum-resident signalling molecules. Protein disulfide isomerase is localized to the endoplasmic reticulum and differentially interacts with calreticulin in response to  $\text{Ca}^{2+}$ -binding (Baksh et al., 1995). Calreticulin is a critical regulator of  $\text{Ca}^{2+}$  stores within the endoplasmic reticulum lumen and also appears to govern  $\text{Ca}^{2+}$  levels in the nuclear envelope and cytoplasm (Krause and Michalak, 1997; Mery et al., 1996; Zhu et al., 1998). As a key regulator of  $\text{Ca}^{2+}$  levels, calreticulin may affect nuclear receptor activity by altering transport of molecules across the nuclear membrane and could have upstream effects on  $\text{Ca}^{2+}$ -mediated nuclear receptor signalling and/or receptor localization. Regulatory interactions between calreticulin and nuclear receptors are very plausible when considering the role of calreticulin in regulating  $\text{Ca}^{2+}$  stores (Mery et al., 1996).

There is precedent for the involvement of nuclear receptors in the regulation of, and response to changes in intracellular  $\text{Ca}^{2+}$  levels, and in the mediation of cross-talk between hormonal and  $\text{Ca}^{2+}$ -signalling networks. Among the nuclear receptors with demonstrated roles in  $\text{Ca}^{2+}$  homeostasis and/or  $\text{Ca}^{2+}$ -signalling are the vitamin D3, estrogen and glucocorticoid receptors (Carlberg and Polly, 1998; Dennis et al., 1987; Yamaguchi and Oishi, 1995). Our laboratory and others have demonstrated that the transcriptional activities of VDR and GR are dramatically repressed by overexpression of calreticulin (Burns et al., 1994; Dedhar et al., 1994; Krause and Michalak, 1997; St-Arnaud et al., 1995; Wheeler et al., 1995; Winrow et al., 1995). Interestingly, the peroxisome proliferators DHEA-S and 5-androstene-3- $\beta$ -17  $\beta$ -diol, and the PPAR $\alpha$  ligand leukotriene B<sub>4</sub> are linked to changes in intracellular  $\text{Ca}^{2+}$  levels (Waxman, 1996; Yokomizo et al., 1997). The interplay between calreticulin and nuclear receptor-directed gene expression appears to involve aspects of  $\text{Ca}^{2+}$ -

signalling and not direct physical interactions between calreticulin and conserved motifs in the DNA binding domains of nuclear receptors. An understanding of what role calreticulin is playing in gene expression is currently an area of intense investigation, however the precise molecular mechanisms involved remain to be determined.

The findings that we have reported provide opportunities for future studies to determine the mechanisms by which calreticulin is affecting the activity of certain nuclear receptors. It will be interesting to ascertain which other nuclear receptors are unaffected or possibly activated by increased calreticulin levels and to compare these receptors with PPAR $\alpha$  and RXR $\alpha$  in order to better understand how calreticulin is affecting gene expression. An additional interesting area to investigate is whether calreticulin is capable of modulating the activity of PPAR $\delta$  or PPAR $\gamma$ . The differences in nuclear receptor susceptibility to calreticulin control could be due to the Ca<sup>2+</sup>-signalling requirements of each particular nuclear receptor. The mechanisms by which endoplasmic reticulum signalling occur are not well understood, but as they become more apparent they will yield greater insight into new means by which signal transduction by the nuclear receptor and other signalling pathways are regulated. This will undoubtedly result in the identification of novel signalling factors and illustrate the co-integration of distinct signalling pathways responsible for directing gene expression.

### *6.1.3 The Effects of TR $\alpha$ on Peroxisome Proliferator Signalling*

The thyroid hormone receptor (TR) family is a well studied group of nuclear receptors whose members are responsible for directing a wide variety of cellular processes, including lipid metabolism and homeostasis (Goodridge, 1987; Shepard and Eberhardt, 1993; Silva and

Rabelo, 1997; Xiong et al., 1998). Most recently, examinations of TR functions have illustrated general concepts for other nuclear receptors in terms of interactions with co-activators and co-repressors (Chen and Evans, 1995; Fondell et al., 1996; Lee et al., 1995; Wagner et al., 1995; Yuan et al., 1998). We observed that TR $\alpha$  binds to the AOx-PPRE as both a monomer and as a TR $\alpha$ /RXR $\alpha$  heterodimer. We further demonstrated that unliganded TR $\alpha$  augmented transactivation by PPAR $\alpha$  from the AOx-PPRE in the presence of a peroxisome proliferator. When activated by its ligand, T3, TR $\alpha$  potently repressed transactivation of the AOx-PPRE by PPAR $\alpha$ .

Following our illustration that TR $\alpha$  modulates peroxisome proliferator-directed expression of the AOx-PPRE, two other groups showed interplay between TR, PPAR and RXR in mediating expression from this element (Chu et al., 1995b; Miyamoto et al., 1997). There have also been a number of additional findings which illustrate cross-talk between thyroid hormone and peroxisome proliferator signalling networks in modulating expression of target genes. These include the co-regulation of the malic enzyme gene, microsomal NADPH:cytochrome P450 oxidoreductase, CYP 4A1, 4A2 and 4A3, and uncoupling protein genes (Ijpenberg et al., 1997; Silva and Rabelo, 1997; Webb et al., 1996). The interplay between the TR and PPAR families that is evident in the regulation of genes involved in lipid metabolism and differentiation, may also overlap in governing other key processes such as apoptosis (Chinetti et al., 1998; Christensen et al., 1998; Llanos et al., 1998; Nagasawa et al., 1997; Tata, 1994). An understanding of the roles of co-activators/co-repressors, dimerization requirements, chromatin remodelling, DNA core site recognition and receptor agonists and antagonists in co-regulation by TR, RXR and PPAR will evolve through examinations of

common target response elements, similar to those we have conducted with the AOX-PPRE.

Subsequent to our demonstration of PPAR $\alpha$ , RXR $\alpha$  and TR $\alpha$  interplay, it was reported that TR $\alpha$  inhibits peroxisome proliferator-induced transactivation of the AOX-PPRE in CV-1, H4IIEC3 and COS-1 cells both in the presence and absence of T3 (Chu et al., 1995b; Miyamoto et al., 1997). The differences between our findings and those presented by these groups may be due to several cell specific factors. We employed the BSC40 African green monkey kidney cell line derived from BS-C-1 cells for transient transfections (Hopps et al., 1963). It is possible that differences in the endogenous levels of nuclear receptors and/or accessory factors may contribute to the differences observed. It would be interesting to examine to what degree known PPAR $\alpha$ /RXR $\alpha$  interacting co-repressors, such as SMRT, are present in BSC40 cells and to compare the expression levels of these factors in CV-1, H4IIEC3 and COS-1 cells. Some nuclear receptor accessory factors demonstrate a restricted tissue distribution and this may account for one of the means by which tissue specific gene expression is realized (Jain et al., 1998; Misiti et al., 1998; Smith et al., 1997). Additionally, the differences in the observed effects of unliganded TR $\alpha$  on the AOX-PPRE may be due to differences in the expression or endogenous levels of PPAR $\alpha$ , RXR $\alpha$  or TR $\alpha$  in the cell lines employed. One of the reasons for using the BSC40 cells is that this cell line contains low levels of intrinsic PPAR, RXR and TR activity, however some residual RXR activity is observed. Since competition for RXR $\alpha$  is proposed to be a major means by which TR affects PPAR-signalling (Chu et al., 1995b; Juge-Aubry et al., 1995), the relative abundance of RXR $\alpha$  in the cell lines examined must be taken into consideration.

TR $\alpha$  can function as a monomer, homodimer and as a heterodimer (Desvergne, 1994;

Ikeda et al., 1994; Lazar et al., 1991b; Tsai and O'Malley, 1994). Although we observe binding of TR $\alpha$  monomers and TR $\alpha$ /RXR $\alpha$  heterodimers to the AOx-PPRE *in vitro*, it may be that non-DNA binding TR $\alpha$  dimers are also contributing to the overall transactivation from this response element *in vivo*. TR $\alpha$  heterodimerizes with RXR $\alpha$  and it is reported that TR $\alpha$ /RXR $\alpha$  heterodimers have less restrictive DNA-binding requirements than TR $\alpha$  homodimers (Olson and Koenig, 1997). TR $\alpha$  was observed to decrease the binding of PPAR $\alpha$ /RXR $\alpha$  to the AOx-PPRE and this is likely due in part to the sequestration of RXR $\alpha$  and competition of TR $\alpha$ /RXR $\alpha$  heterodimers for the binding to the AOx-PPRE. These mechanisms have been suggested to explain TR $\alpha$  actions with the AOx-PPRE since our initial investigation (Chu et al., 1995b; Miyamoto et al., 1997). TR $\alpha$  can also sequester PPAR $\alpha$  in solution, and this may also account for the reduced PPAR $\alpha$ /RXR $\alpha$  binding to the AOx-PPRE which we observed (Bogazzi et al., 1994). It appears that, contrary to what is reported for TR $\alpha$ /RXR $\alpha$  heterodimers, TR $\alpha$ /PPAR $\alpha$  heterodimers have a much more restricted DNA-binding preference, possibly as a result of the associations between their heterodimerization interfaces.

One explanation for the TR-mediated potentiation of the AOx-PPRE by PPAR $\alpha$ /RXR $\alpha$  could be the sequestration of a shared co-repressor, such as SMRT, by TR $\alpha$ . In the absence of ligand, TRs act as potent transcriptional silencers (Banahmad et al., 1992; Burcin et al., 1994; Nawaz et al., 1995; Renkawitz, 1993). Analysis of this observation showed that the transcriptional silencing by TRs is dependent on interactions with co-repressors (Banahmad et al., 1995; Chen and Evans, 1995; Kurokawa et al., 1995; Lee et al., 1995; Sande and Privalsky, 1996). It has been clearly demonstrated that unliganded TR $\alpha$

associates with co-repressors and it is possible that TR $\alpha$  which is not bound to DNA may be sequestering co-regulatory factors which would otherwise interact with PPAR $\alpha$  or RXR $\alpha$  (Chen and Evans, 1995; Kurokawa et al., 1995; Lee et al., 1995). Such interactions would facilitate the association of liganded PPAR $\alpha$ /RXR $\alpha$  heterodimers with positive co-activators, with the net result being an increase in the transactivation by PPAR $\alpha$ /RXR $\alpha$ .

In the presence of the T3, transactivation of the AO $\alpha$ -PPRE is potently inhibited when TR $\alpha$  is included in transfections. There may be several reasons for this, including the titration of common co-activators from PPAR/RXR, formation of ligand-directed non-DNA binding TR $\alpha$ /PPAR $\alpha$  or TR $\alpha$ /RXR $\alpha$  heterodimers, or binding of liganded TR $\alpha$  monomers or TR $\alpha$ /RXR $\alpha$  heterodimers to the AO $\alpha$ -PPRE in a conformation which represses transcription. TR $\alpha$  is functional in our system, since it is able to activate transcription of a thyroid response element in response to T3. Although it is more commonly observed that ligand binding increases transactivation, liganded nuclear receptors can also function as transcriptional repressors depending on the promoter context. This phenomenon has been observed in PPAR $\alpha$ /RXR $\alpha$  interactions with response elements of the apolipoprotein CIII gene (Hertz et al., 1995). Therefore, it remains possible that ligand-bound TR $\alpha$  is directly repressing transactivation from the AO $\alpha$ -PPRE.

We have shown that the efficient potentiation of the AO $\alpha$ -PPRE by TR $\alpha$  requires the transfection of both PPAR $\alpha$  and RXR $\alpha$ . Interestingly, in the absence of transfected RXR $\alpha$ , TR $\alpha$  slightly represses the peroxisome proliferator-dependent transactivation of the AO $\alpha$ -PPRE reporter gene in a dose dependent manner. Therefore, it appears that the availability of RXR $\alpha$  is also a critical determinant in the TR $\alpha$  regulation of the AO $\alpha$ -PPRE. Some groups

have suggested that TR $\alpha$ /RXR $\alpha$  heterodimers are more promiscuous in their DNA binding site preference compared to TR $\alpha$  homodimers (Olson and Koenig, 1997). Although heterodimerization between TR $\alpha$  and PPAR $\alpha$  has been reported, however no natural target element for TR $\alpha$ /PPAR $\alpha$  heterodimers has been characterized (Bogazzi et al., 1994). TR $\alpha$  associates with the AOx-PPRE as both a monomer and as a TR $\alpha$ /RXR $\alpha$  heterodimer *in vitro*; however, it remains possible that TR $\alpha$ /PPAR $\alpha$  heterodimers are influencing the *in vivo* response to peroxisome proliferators. It may be that the monomeric form of TR $\alpha$  is responsible for maintaining a repressive state, whereas unliganded TR $\alpha$ /RXR $\alpha$  augments transactivation by PPAR $\alpha$ /RXR $\alpha$ , either through direct interactions with the AOx-PPRE or by other means.

In order to obtain a more precise picture of TR $\alpha$ , PPAR $\alpha$  and RXR $\alpha$  interactions with the AOx-PPRE several further experiments would be beneficial. An examination of the roles of shared nuclear receptor accessory factors, such as SMRT and SRC-1 will help to identify the requirements of PPAR $\alpha$ , TR $\alpha$  and RXR $\alpha$  for co-repressors and co-activators in the regulation of the AOx-PPRE. Studies which employ mutant TR $\alpha$  will assist in determining a more specific role for this receptor in peroxisome proliferator-signalling. Several TR $\alpha$  mutants have been characterized which exhibit altered capacities for dimerization, ligand binding or association with distinct response elements (Kitajima et al., 1995; Liu et al., 1996; Nagaya et al., 1992). In addition, specific mutations in the ligand binding domains of TRs have been identified which permit differential interactions with co-regulatory factors independent of ligand association (Liu et al., 1998; Nagaya et al., 1998). The use of a yeast system for analysis of nuclear receptor interactions and transactivation has been employed




successfully (Marcus et al., 1996), and may provide a suitable environment in which to further study TR/RXR/PPAR regulation of the AOx-PPRE. A closer investigation of the possible roles of TR $\alpha$ /PPAR $\alpha$  heterodimers *in vivo*, the effect of their respective ligands on dimerization and the specific actions of co-regulatory factors will aid in developing a more complete model of TR $\alpha$ , PPAR $\alpha$  and RXR $\alpha$  interplay.

#### 6.1.4 RZR $\alpha$ , PPAR $\alpha$ and RXR $\alpha$ Interplay

RZR $\alpha$ s are one of the few identified nuclear receptors which bind to target response elements as monomers (Carlberg et al., 1994; Giguere et al., 1994). The RZR consensus binding site is conserved within the HD-PPRE and to a lesser extent in the AOx-PPRE. We observed binding of RZR $\alpha$  specifically to the HD-PPRE, and saw potentiation of PPAR $\alpha$  transactivation by RZR $\alpha$  from this response element when RXR $\alpha$  levels were limiting. The results of these experiments are summarized in Fig. 6-1. The RZR subfamily of receptors contain a carboxy terminal extension in their DNA-binding domains which is required for the recognition of an extended 3' sequence flanking the core TGACCT-like site (Giguere et al., 1995; McBroom et al., 1995; Wilson et al., 1992). PPAR $\alpha$  also contains a similar carboxy terminal extension of its zinc finger which may account for the requirement for 3' sequences flanking PPREs (Hsu et al., 1998). This may also explain why PPAR $\alpha$ /RXR $\alpha$  heterodimers exhibit the opposite polarity from most other RXR $\alpha$  heterodimers (Chu et al., 1995a; IJpenberg et al., 1997; Rastinejad et al., 1995). The most commonly observed orientation for RXR $\alpha$  heterodimers is with RXR $\alpha$  binding to the 3' TGACCT-like site (Kurokawa et al., 1994; Rastinejad et al., 1995).

Transcriptional Activity



**Figure 6-1. Summary of results from studies of RZR $\alpha$  effects on peroxisome proliferator-mediated transactivation from the HD-PPRE.**

The model I have provided in Figure 6-2 details the potential interactions by which RZR $\alpha$  may be influencing PPAR $\alpha$ -mediated gene expression, however the general means through which RZR $\alpha$  is functioning can be extended to explain how cross-talk may be occurring with other nuclear receptors. This model was derived from the observations that the RZR $\alpha$  monomer recognizes the second TGACCT-related repeat within the HD-PPRE (site II) and that peroxisome proliferator-dependent RZR $\alpha$  potentiation of PPAR $\alpha$  activity is only observed in conditions where levels of RXR $\alpha$  are limiting. The overall level of PPAR $\alpha$  transactivation is greatly increased when exogenous RXR $\alpha$  is included in co-transfections. It may be that under these conditions any potentiation by RZR $\alpha$  is concealed. It has been proposed that *in vivo*, PPAR $\alpha$ /RXR $\alpha$  heterodimers bind to each DR1 site within the HD-PPRE, yielding the highest level of transactivation (Chu et al., 1995a). In conditions of RXR $\alpha$  excess *in vivo*, this conformation may preclude association of RZR $\alpha$  with site II. Based on our *in vitro* DNA-binding studies with HD-PPRE mutants, it appears that PPAR $\alpha$ /RXR $\alpha$  heterodimer binding is much stronger at sites III and IV. PPAR $\alpha$ /RXR $\alpha$  heterodimers may require additional components that are not present in our *in vitro* system, to enable binding to sites I and II of the HD-PPRE.

The intermediate level of PPAR $\alpha$ -directed transactivation provided by RZR $\alpha$  in conditions of limiting RXR $\alpha$  may be accounted for by several non-mutually exclusive means (see Fig. 6-2). It may be that for the RZR $\alpha$  monomer, association at site II augments PPAR $\alpha$ /RXR $\alpha$  transactivation by stabilizing the activated PPAR $\alpha$ /RXR $\alpha$  heterodimer bound at sites III and IV, and may help to maintain PPAR $\alpha$ /RXR $\alpha$  in a ligand associated conformation. Since RZR $\alpha$  binds to the HD-PPRE at a site directly beside RXR $\alpha$ , it would

**Figure 6-2. Possible roles for RZR $\alpha$  in directing peroxisome proliferator-mediated transactivation from the HD-PPRE.** Arrows indicate putative sites of interaction for the RZR $\alpha$  monomer. RZR $\alpha$  may be altering transcription by (i) stabilizing activated PPAR $\alpha$ /RXR $\alpha$  heterodimers, (ii) facilitating assembly/association of a co-activator complex or (iii) assisting dissociation of a co-repressor complex.




Figure 6-2. Possible roles for RZRα in directing peroxisome proliferator-mediated transactivation from the HD-PPRE.

be interesting to determine if RZR $\alpha$  interacts weakly with aspects of RXR $\alpha$  and to examine what effect the RXR $\alpha$  ligand, 9-*cis*-retinoic acid may have on interactions with RZR $\alpha$ . It does not appear that RZR $\alpha$  is affecting transactivation indirectly, since mutation of the site II eliminates the potentiation of PPAR $\alpha$ /RXR $\alpha$  transactivation by RZR $\alpha$ .

The slight repressive effect of RZR $\alpha$  in the absence of a peroxisome proliferator indicates that it is specifically the interactions between ligand activated PPAR $\alpha$ /RXR $\alpha$  and RZR $\alpha$  which increased activation from the HD-PPRE, hinting at the importance of PPAR $\alpha$ /RXR $\alpha$ /RZR $\alpha$ -co-regulator interactions. Yeast two-hybrid analysis may be of use to identify RZR $\alpha$ -interacting factors and would help to determine what role this nuclear receptor is playing in the assembly of a co-activator complex. RZR $\alpha$  does contain an AF-2 region which is necessary for interaction with known co-activators, however the association of RZR $\alpha$  with co-activators has not yet been examined. Additionally, RZR $\alpha$  may be interacting with p300/CBP or p/CAF or other components of the co-activator complex. RZR $\alpha$  may also assist the displacement of a co-repressor and prevent association of PPAR $\alpha$ /RXR $\alpha$  heterodimers with components of a co-repressor complex. It will be interesting to determine if regions of the RZR $\alpha$  monomer interact directly with RXR and/or with PPAR/RXR co-activators or co-repressors.

The conclusive identification of an RZR $\alpha$  ligand will also open the doors for future studies. Although melatonin has been proposed to function as an RZR agonist, we saw no effects of melatonin on RZR $\alpha$  activity in our experiments. Recent evidence has indicated that several thiazolidenediones act as RZR ligands, and this suggests additional possibilities for examining the effects of RZR $\alpha$  on PPAR-directed gene expression (Missbach et al., 1996).

All transfections that we performed employed depleted serum, however it remains possible that an endogenous RXR $\alpha$  ligand is present in our transfection system and therefore RXR $\alpha$  may be already present in an active form.

## 6.2 General Model of Nuclear Receptor Interactions with the AOx- and HD-PPREs

The regulation of the AOx- and HD-PPREs by nuclear receptors is dynamic and the general modulation of target gene expression by nuclear receptors is by nature a fluid arrangement, allowing rapid and concerted transcriptional responses to appropriate physiological stimuli. A model of nuclear receptor interplay involved in the regulation of the AOx- and HD-PPREs is emerging based on studies in our laboratory and elsewhere (see Fig. 6-3).

The means by which nuclear receptors cross-talk are both direct and indirect (Katzenellenbogen et al., 1996). Nuclear receptors influence transcription directly through competition for binding sites, sequestration of common heterodimerization partners, such as RXR $\alpha$ , and by mediating interactions with co-activators or co-repressors. The effects of nuclear receptor cross-talk may be indirect, through changes in upstream signals, producing physical modifications in downstream receptors or altering ligand availability.

We observed that nuclear receptor competition for shared DNA binding sites occurs with both the AOx- and HD-PPREs. This competition was seen for the AOx-PPRE with PPAR $\alpha$ , RXR $\alpha$ , COUP-TF, HNF-4 and TR $\alpha$  through both *in vitro* binding studies and in transient transfections (Marcus et al., 1996; Miyata et al., 1993; Winrow et al., 1996; Winrow et al., 1994). HNF-4 and COUP-TF were found to compete with PPAR $\alpha$ /RXR $\alpha$




Figure 6-3. Overview of nuclear receptor crosstalk in the transactivation from the AOX- and HD-PPREs.



heterodimers for binding to the HD-PPRE *in vitro* (Marcus et al., 1996; Miyata et al., 1993; Winrow et al., 1994). By virtue of experiments with mutated HD-PPREs, it is becoming apparent that although sites III and IV are important for PPAR $\alpha$ /RXR $\alpha$  binding, site II is necessary for the co-regulatory effects of several nuclear receptors (Marcus et al., 1996; Miyata et al., 1993; Winrow et al., 1998; Winrow et al., 1994). Mutations of sites I or II have minimal effects on PPAR $\alpha$ /RXR $\alpha$  DNA binding *in vitro* (see Chapter 5, Fig. 5-6) (Miyata et al., 1993). The mutation of site II yields a reduced level of overall activity but is still transactivated by PPAR $\alpha$ /RXR $\alpha$  in a peroxisome proliferator-dependent manner in our transfection system (see Chapter 5, Fig. 5-7). Site II is required for the *in vitro* binding of HNF-4, COUP-TF and RZR $\alpha$  and was also shown to be necessary for the PPAR $\alpha$ -dependent potentiation of HD-PPRE by RZR $\alpha$  (Miyata et al., 1993; Winrow et al., 1998; Winrow et al., 1994). It is interesting to note that based on the reported polarity of PPAR $\alpha$ /RXR $\alpha$  heterodimers, RXR $\alpha$  binds upstream to site III and PPAR $\alpha$  binds downstream to site IV (see Fig.6-2) (Chu et al., 1995a; IJpenberg et al., 1997; Kurokawa et al., 1994). Given the importance of site II in the co-regulation of the HD-PPRE, it will be interesting to determine if, for HNF-4 and RZR $\alpha$ , the proximity to RXR $\alpha$  facilitates physical inter-receptor contact, which thereby modifies transactivation. Insight into which domains are necessary for these interactions and how association of 9-*cis*-retinoic acid may influence this interplay will be beneficial to our overall understanding of the co-regulatory mechanisms involved.

TAK1 is a novel orphan nuclear receptor found in hepatocytes, that represses PPAR $\alpha$  transactivation from both the AOX - and HD-PPREs (Yan et al., 1998). TAK1 functions as a homodimer and does not interact with either PPAR $\alpha$  or RXR $\alpha$ . It was determined that

TAK1 inhibits PPAR $\alpha$ /RXR $\alpha$  transactivation by both competing for binding to these PPRES and by sequestering the PPAR co-activator RIP140 (Yan et al., 1998). Similar mechanisms are likely involved in the interplay between PPAR, RXR and other co-regulatory nuclear receptors. In particular, our results suggest that the cross-talk between TR $\alpha$ , PPAR $\alpha$  and RXR $\alpha$  involves competition for DNA binding and titration of shared co-activators and co-repressors.

RZR $\alpha$  binds target response elements as a monomer and HNF-4 functions as a homodimer, however TR $\alpha$  can bind the AOx-PPRE as both a monomer and as a heterodimer with RXR $\alpha$ . Competition for a common heterodimeric partner such as RXR $\alpha$  is another means by which TR $\alpha$  may cross-talk with PPAR $\alpha$  in regulating the AOx-PPRE. Nuclear receptors may affect PPAR $\alpha$ /RXR $\alpha$  transactivation by sequestering either PPAR $\alpha$  or RXR $\alpha$  as a heterodimer which will not associate with either the AOx- or HD-PPRE (Miyata et al., 1996). This may be the case for TR $\alpha$ , where TR $\alpha$ /RXR $\alpha$  and TR $\alpha$ /PPAR $\alpha$  heterodimers may not recognize the AOx-PPRE *in vivo*. Liver X receptor (LXR) is an orphan nuclear receptor which was identified as a PPAR $\alpha$ -interacting factor by a yeast two-hybrid screen (Miyata et al., 1996). LXR $\alpha$  decreases PPAR $\alpha$ /RXR $\alpha$  transactivation from the AOx- and HD-PPRES but does not appear to bind to these response elements (Miyata et al., 1996). It was determined that LXR $\alpha$  may be affecting peroxisome proliferator transactivation of the AOx- and HD-PPRES by sequestering PPAR $\alpha$  or RXR $\alpha$ , through the formation of LXR $\alpha$  heterodimers which do not recognize these response elements (Miyata et al., 1996).

The presence of a peroxisome proliferator served to augment PPAR $\alpha$ /RXR $\alpha$ -dependent transactivation from the AOx-PPRE by TR $\alpha$  and from the HD-PPRE by RZR $\alpha$  and

HNF-4. This may be a result of direct stabilization of the active PPAR $\alpha$ /RXR $\alpha$  heterodimer, possibly by maintaining PPAR $\alpha$  in an agonist bound state. Additionally, this potentiation may be occurring through the titration of a co-repressor, thereby preventing assembly of a co-repressor complex and sustaining the chromatin in an acetylated conformation. The positive transcriptional effects of TR $\alpha$ , HNF-4 and RZR $\alpha$  may also be manifested by facilitating interactions of PPAR $\alpha$ /RXR $\alpha$  with a co-activator or assisting assembly of a co-activator complex. PPAR $\alpha$ , RXR $\alpha$  and TR $\alpha$  associate with common co-activators, and it is probable that HNF-4 and RZR $\alpha$  also share a number of identical co-activator molecules.

HNF-4, RZR $\alpha$  and TR $\alpha$  could affect peroxisome proliferator modulated transactivation indirectly by modifying the upstream signalling of the peroxisome proliferator pathway. It has been reported that PPAR $\gamma$  is a phosphoprotein that is phosphorylated on serine residues through the mitogen-activated protein kinase cascade (Adams et al., 1997; Camp and Tafuri, 1997). It is possible that HNF-4, RZR $\alpha$  and/or TR $\alpha$  might modulate the phosphorylation states of PPARs or RXR $\alpha$  by indirectly altering the activity of a required kinase. Alternatively, these nuclear receptors may affect some other step of peroxisome proliferator signal transduction, such as ligand availability, and thereby alter PPAR $\alpha$ /RXR $\alpha$ -mediated gene expression. This can be easily envisioned in the case of PPAR where ligands include prostaglandins, fatty acids and leukotrienes whose synthesis or modification may very well occur as a result of pathways directed by TR $\alpha$ , HNF-4 or RZR $\alpha$  or other nuclear receptors. The recent observation that HNF-4 activity is modulated by binding of fatty acyl-CoA thioesters is particularly suggestive of ligand directed interplay with PPAR $\alpha$ , given that many long chain fatty acids function as peroxisome proliferators (Dreyer et al., 1993; Forman

et al., 1997a; Hertz et al., 1998; Kliewer et al., 1997; Krey et al., 1997; Willson and Wahli, 1997).

We have demonstrated that calreticulin, which modulates cellular  $\text{Ca}^{2+}$ -storage and signalling, can dramatically alter the transactivation potential of the GR, while exerting no discernible influence on PPAR $\alpha$ /RXR $\alpha$  signalling. It is thought that although calreticulin recognizes a consensus amino acid motif found in the DNA binding domains of virtually all nuclear receptors, the effect of calreticulin on receptor activity may be indirect and possibly due to a combination of alterations of cellular  $\text{Ca}^{2+}$  levels and endoplasmic reticulum signalling (Baksh et al., 1995; Krause and Michalak, 1997; Mery et al., 1996; Michalak et al., 1996).

### 6.3 Summary

Peroxisomes are the site of multiple processes including bile acid and cholesterol biosynthesis, degradation of  $\text{H}_2\text{O}_2$  and  $\beta$ -oxidation of fatty acids. The regulation of gene expression by nuclear receptors is a dynamic process which provides for transcriptional responses to a multitude of signalling pathways. I have examined the interplay between nuclear receptors in the transcriptional regulation of genes involved in peroxisomal  $\beta$ -oxidation. In addition I have investigated the influence of a  $\text{Ca}^{2+}$ -regulatory protein, calreticulin, which has been shown to affect the activity of several nuclear receptors, in PPAR-mediated transcriptional activation.

HNF-4 binds target response elements as a homodimer and was shown to bind readily to the AOx-PPRE and to a lesser extent to the HD-PPRE. In the absence of a peroxisome

proliferator, HNF-4 inhibited transactivation by PPAR $\alpha$ /RXR $\alpha$  heterodimers from both the AOx- and HD-PPREs. HNF-4 differentially regulated transactivation by PPAR $\alpha$ /RXR $\alpha$  when a peroxisome proliferator was included in co-transfections. Under these conditions PPAR $\alpha$ /RXR $\alpha$  transactivation from the AOx-PPRE was inhibited by HNF-4, whereas activation of the HD-PPRE was increased by HNF-4. This differential effect of HNF-4 could be a reflection of differences in DNA response element affinity of the receptor and may involve additional interactions with common nuclear receptor accessory factors.

Calreticulin was shown to interact *in vitro* with a number of nuclear receptors via a conserved amino acid motif in their DNA binding domains. Although calreticulin inhibited transactivation by GR, no effect of calreticulin on the activity of PPAR $\alpha$ /RXR $\alpha$  heterodimers was observed in a whole cell setting. We demonstrated that calreticulin is not a general regulator of nuclear receptor activity, as had been proposed. It is most likely that the effect of calreticulin on gene expression is indirect, possibly due to alterations in intracellular Ca<sup>2+</sup> levels which might influence responsive nuclear receptors.

Unliganded TR $\alpha$  is able to bind to the AOx-PPRE *in vitro* and potentiates transactivation of this element by peroxisome proliferator-stimulated PPAR $\alpha$  in a cellular context. Addition of the TR ligand, T3, potently repressed PPAR $\alpha$ -mediated transcriptional activity for the AOx-PPRE when TR $\alpha$  was present. Although TR $\alpha$  is able to bind to the AOx-PPRE as both a monomer and as a TR $\alpha$ /RXR $\alpha$  heterodimer, it is possible that unliganded TR $\alpha$  potentiates PPAR $\alpha$  activity by sequestration of a common co-repressor.

RZR $\alpha$  binds specifically to the HD-PPRE as a monomer but does not bind to the AOx-PPRE. The binding of the RZR $\alpha$  monomer was localized to the second TGACCT-like

repeat within the HD-PPRE, and this site was shown to be essential for RXR $\alpha$  potentiation of PPAR $\alpha$  activity *in vivo*. In conditions where RXR $\alpha$  was limiting, RXR $\alpha$  augmented the activity of PPAR $\alpha$  from the HD-PPRE in an agonist-dependent manner. It appears that RXR $\alpha$  is mediating this effect by binding to the HD-PPRE at a site adjacent to RXR $\alpha$  and possibly stabilizing the ligand-bound form of the PPAR $\alpha$ /RXR $\alpha$  complex, inhibiting association with a co-repressor and/or facilitating assembly of a co-activator complex.

These findings show how the PPAR, RXR, HNF-4, TR and RXR signalling pathways are interconnected in the synergistic or differential expression of the AOx and HD genes by virtue of their PPREs. The intricate communication between diverse networks of nuclear receptors enables a high degree of specificity and confers the ability to rapidly respond to diverse physiological stimuli, be they transient or long-term. This research extends the current knowledge regarding peroxisome proliferator stimulation of gene expression and conveys the complexity of interactions involved in the coordinate regulation of the peroxisomal  $\beta$ -oxidation pathway at the level of transcription.

#### **6.4 Conclusions**

The research presented herein illustrates the integration of multiple signalling networks in the concerted regulation of genes encoding the first two enzymes of the peroxisomal  $\beta$ -oxidation cascade. Taken together these findings demonstrate the multilevel control of the AOx- and HD-PPRES through cross-talk between diverse nuclear receptor signalling pathways. The observations obtained, provide insight into the specific mechanisms of gene regulation by PPAR $\alpha$  and RXR $\alpha$ , and at the same time indicate several directions for

future studies which will prove fruitful in expanding our knowledge of the general mechanisms through which the effects of nuclear receptors are mediated. The implications of this research touch not only the transcriptional regulation of genes responsible for peroxisomal  $\beta$ -oxidation, but also underscore the overall importance of nuclear receptor interplay and afford firm indications as to the co-regulatory roles of distinct nuclear receptors.

## 6.5 BIBLIOGRAPHY

- Adams, M., Reginato, M.J., Shao, D., and Lazar, M.A., Chatterjee, V.K. (1997). Transcriptional activation by peroxisome proliferator-activated receptor gamma is inhibited by phosphorylation at a consensus mitogen-activated protein kinase site. *J. Biol. Chem.* 272, 5128-32.
- Aldridge, T. C., Tugwood, J. D., and Green, S. (1995). Identification and characterization of DNA elements implicated in the regulation of CYP4A1 transcription. *Biochem J* 306, 473-9.
- Alland, L., Muhle, R., Hou, H., Jr., Potes, J., Chin, L., Schreiber-Agus, N., and DePinho, R. A. (1997). Role for N-CoR and histone deacetylase in Sin3-mediated transcriptional repression. *Nature* 387, 49-55.
- Aperlo, C., Pognonec, P., Saladin, R., Auwerx, J., and Boulukos, K. E. (1995). cDNA cloning and characterization of the transcriptional activities of the hamster peroxisome proliferator-activated receptor haPPAR gamma. *Gene* 162, 297-302.
- Arany, Z., Sellers, W. R., Livingston, D. M., and Eckner, R. (1994). E1A-associated p300 and CREB-associated CBP belong to a conserved family of coactivators. *Cell* 77, 799-800.
- Aubert, J., Champigny, O., Saint-Marc, P., Negrel, R., Collins, S., Ricquier, D., and Ailhaud, G. (1997). Up-regulation of UCP-2 gene expression by PPAR agonists in preadipose and adipose cells. *Biochem Biophys Res Commun* 238, 606-11.
- Austin, S., Medvedev, A., Yan, Z. H., Adachi, H., Hirose, T., and Jetten, A. M. (1998). Induction of the nuclear orphan receptor ROR $\gamma$  during adipocyte differentiation of D1 and 3T3-L1 cells. *Cell Growth Differ* 9, 267-76.
- Ausubel, F. M. (1992). *Current protocols in molecular biology* (New York: Published by Greene Pub. Associates and Wiley-Interscience : J. Wiley), pp. Section 9.1.
- Baksh, S., Burns, K., Busaan, J., and Michalak, M. (1992). Expression and purification of recombinant and native calreticulin. *Protein Expr Purif* 3, 322-31.
- Baksh, S., Burns, K., Andrin, C., and Michalak, M. (1995). Interaction of calreticulin with protein disulfide isomerase. *J Biol Chem* 270, 31338-44.
- Baniahmad, A., Kohne, A. C., and Renkawitz, R. (1992). A transferable silencing domain is present in the thyroid hormone receptor, in the v-erbA oncogene product and in the retinoic acid receptor. *Embo J* 11, 1015-23.



- Baniahmad, A., Leng, X., Burris, T. P., Tsai, S. Y., Tsai, M. J., and O'Malley, B. W. (1995). The tau 4 activation domain of the thyroid hormone receptor is required for release of a putative corepressor(s) necessary for transcriptional silencing. *Mol Cell Biol* *15*, 76-86.
- Bannister, A. J., and Kouzarides, T. (1996). The CBP co-activator is a histone acetyltransferase. *Nature* *384*, 641-3.
- Bardot, O., Aldridge, T. C., Latruffe, N., and Green, S. (1993). PPAR-RXR heterodimer activates a peroxisome proliferator response element upstream of the bifunctional enzyme gene. *Biochem Biophys Res Commun* *192*, 37-45.
- Becker-Andre, M., Andre, E., and DeLamarter, J. F. (1993). Identification of nuclear receptor mRNAs by RT-PCR amplification of conserved zinc-finger motif sequences. *Biochem Biophys Res Commun* *194*, 1371-9.
- Becker-Andre, M., Wiesenberg, I., Schaeren-Wiemers, N., Andre, E., Missbach, M., Saurat, J. H., and Carlberg, C. (1994). Pineal gland hormone melatonin binds and activates an orphan of the nuclear receptor superfamily (published erratum appears in *J Biol Chem* 1997 Jun 27;272(26):16707). *J Biol Chem* *269*, 28531-4.
- Bell, A. R., Savory, R., Horley, N. J., Choudhury, A. I., Dickins, M., Gray, T. J., Salter, A. M., and Bell, D. R. (1998). Molecular basis of non-responsiveness to peroxisome proliferators: the guinea-pig PPAR $\alpha$  is functional and mediates peroxisome proliferator-induced hypolipidaemia. *Biochem J* *332*, 689-93.
- Bendik, I., and Pfahl, M. (1995). Similar ligand-induced conformational changes of thyroid hormone receptors regulate homo- and heterodimeric functions. *J Biol Chem* *270*, 3107-14.
- Bentley, P., Calder, I., Elcombe, C., Grasso, P., Stringer, D., and Wiegand, H. J. (1993). Hepatic peroxisome proliferation in rodents and its significance for humans. *Food Chem Toxicol* *31*, 857-907.
- Besnard, P., Mallordy, A., and Carlier, H. (1993). Transcriptional induction of the fatty acid binding protein gene in mouse liver by bezafibrate. *FEBS Lett* *327*, 219-23.
- Bieri, F., and Lhuguenot, J. C. (1993). Toxicity of peroxisome proliferators. *Biochimie* *75*, 263-8.
- Bleackley, R. C., Atkinson, E. A., Burns, K., and Michalak, M. (1995). Calreticulin: a granule-protein by default or design? *Curr Top Microbiol Immunol* *198*, 145-59.

- Bogazzi, F., Hudson, L. D., and Nikodem, V. M. (1994). A novel heterodimerization partner for thyroid hormone receptor. Peroxisome proliferator-activated receptor. *J Biol Chem* 269, 11683-6.
- Bourguet, W., Ruff, M., Chambon, P., Gronemeyer, H., and Moras, D. (1995). Crystal structure of the ligand-binding domain of the human nuclear receptor RXR-alpha. *Nature* 375, 377-82.
- Braissant, O., and Wahli, W. (1998). Differential expression of peroxisome proliferator-activated receptor- alpha, -beta, and -gamma during rat embryonic development. *Endocrinology* 139, 2748-54.
- Brzozowski, A. M., Pike, A. C., Dauter, Z., Hubbard, R. E., Bonn, T., Engstrom, O., Ohman, L., Greene, G. L., Gustafsson, J. A., and Carlquist, M. (1997). Molecular basis of agonism and antagonism in the oestrogen receptor. *Nature* 389, 753-8.
- Burcin, M., Kohne, A. C., Runge, D., Steiner, C., and Renkawitz, R. (1994). Factors influencing nuclear receptors in transcriptional repression. *Semin Cancer Biol* 5, 337-46.
- Burns, K., Duggan, B., Atkinson, E. A., Famulski, K. S., Nemer, M., Bleackley, R. C., and Michalak, M. (1994). Modulation of gene expression by calreticulin binding to the glucocorticoid receptor. *Nature* 367, 476-80.
- Camirand, A., Marie, V., Rabelo, R., and Silva, J. E. (1998). Thiazolidinediones stimulate uncoupling protein-2 expression in cell lines representing white and brown adipose tissues and skeletal muscle. *Endocrinology* 139, 428-31.
- Camp, H. S., and Tafuri, S. R. (1997). Regulation of peroxisome proliferator-activated receptor gamma activity by mitogen-activated protein kinase. *J Biol Chem* 272, 10811-6.
- Carlberg, C., Hooft van Huijsduijnen, R., Staple, J. K., DeLamarter, J. F., and Becker-Andre, M. (1994). RZR, a new family of retinoid-related orphan receptors that function as both monomers and homodimers. *Mol Endocrinol* 8, 757-70.
- Carlberg, C., and Wiesenberg, I. (1995). The orphan receptor family RZR/ROR, melatonin and 5-lipoxygenase: an unexpected relationship. *J Pineal Res* 18, 171-8.
- Carlberg, C., and Polly, P. (1998). Gene regulation by vitamin D3. *Crit Rev Eukaryot Gene Expr* 8, 19-42.
- Carson-Jurica, M. A., Schrader, W. T., and O'Malley, B. W. (1990). Steroid receptor family: structure and functions. *Endocr Rev* 11, 201-20.

Carter, M. E., Gulick, T., Raisher, B. D., Caira, T., Ladias, J. A., Moore, D. D., and Kelly, D. P. (1993). Hepatocyte nuclear factor-4 activates medium chain acyl-CoA dehydrogenase gene transcription by interacting with a complex regulatory element. *J Biol Chem* 268, 13805-10.

Casanova, J., Helmer, E., Selmi-Ruby, S., Qi, J. S., Au-Fliegner, M., Desai-Yajnik, V., Koudinova, N., Yarm, F., Raaka, B. M., and Samuels, H. H. (1994). Functional evidence for ligand-dependent dissociation of thyroid hormone and retinoic acid receptors from an inhibitory cellular factor. *Mol Cell Biol* 14, 5756-65.

Castelein, H., Gulick, T., Declercq, P. E., Mannaerts, G. P., Moore, D. D., and Baes, M. I. (1994). The peroxisome proliferator activated receptor regulates malic enzyme gene expression. *J Biol Chem* 269, 26754-8.

Castelein, H., Declercq, P. E., and Baes, M. (1997). DNA binding preferences of PPAR $\alpha$ /RXR $\alpha$  heterodimers. *Biochem Biophys Res Commun* 233, 91-5.

Cattley, R. C., DeLuca, J., Elcombe, C., Fenner-Crisp, P., Lake, B. G., Marsman, D. S., Pastoor, T. A., Popp, J. A., Robinson, D. E., Schwetz, B., Tugwood, J., and Wahli, W. (1998). Do peroxisome proliferating compounds pose a hepatocarcinogenic hazard to humans? *Regul Toxicol Pharmacol* 27, 47-60.

Chakravarti, D., LaMorte, V. J., Nelson, M. C., Nakajima, T., Schulman, I. G., Juguilon, H., Montminy, M., and Evans, R. M. (1996). Role of CBP/P300 in nuclear receptor signalling. *Nature* 383, 99-103.

Chatterjee, V. K., and Tata, J. R. (1992). Thyroid hormone receptors and their role in development. *Cancer Surv* 14, 147-67.

Chawla, A., and Lazar, M. A. (1994). Peroxisome proliferator and retinoid signaling pathways co-regulate preadipocyte phenotype and survival. *Proc Natl Acad Sci U S A* 91, 1786-90.

Chen, F., Law, S. W., and O'Malley, B. W. (1993). Identification of two mPPAR related receptors and evidence for the existence of five subfamily members. *Biochem Biophys Res Commun* 196, 671-7.

Chen, D., Lepar, G., and Kemper, B. (1994). A transcriptional regulatory element common to a large family of hepatic cytochrome P450 genes is a functional binding site of the orphan receptor HNF-4. *J Biol Chem* 269, 5420-7.

Chen, J. D., and Evans, R. M. (1995). A transcriptional co-repressor that interacts with nuclear hormone receptors. *Nature* 377, 454-7.

- Chen, H., Lin, R. J., Schiltz, R. L., Chakravarti, D., Nash, A., Nagy, L., Privalsky, M. L., Nakatani, Y., and Evans, R. M. (1997). Nuclear receptor coactivator ACTR is a novel histone acetyltransferase and forms a multimeric activation complex with P/CAF and CBP/p300. *Cell* 90, 569-80.
- Chen, J. D., and Li, H. (1998). Coactivation and corepression in transcriptional regulation by steroid/nuclear hormone receptors. *Crit Rev Eukaryot Gene Expr* 8, 169-90.
- Cherkaoui Malki, M., Lone, Y. C., Corral-Debrinski, M., and Latruffe, N. (1990). Differential proto-oncogene mRNA induction from rats treated with peroxisome proliferators. *Biochem Biophys Res Commun* 173, 855-61.
- Chinetti, G., Griglio, S., Antonucci, M., Torra, I. P., Delerive, P., Majd, Z., Fruchart, J. C., Chapman, J., Najib, J., and Staels, B. (1998). Activation of proliferator-activated receptors alpha and gamma induces apoptosis of human monocyte-derived macrophages. *J Biol Chem* 273, 25573-80.
- Christensen, J. G., Gonzales, A. J., Cattley, R. C., and Goldsworthy, T. L. (1998). Regulation of apoptosis in mouse hepatocytes and alteration of apoptosis by nongenotoxic carcinogens. *Cell Growth Differ* 9, 815-25.
- Chu, R., Lin, Y., Rao, M. S., and Reddy, J. K. (1995a). Cooperative formation of higher order peroxisome proliferator-activated receptor and retinoid X receptor complexes on the peroxisome proliferator responsive element of the rat hydratase-dehydrogenase gene. *J Biol Chem* 270, 29636-9.
- Chu, R., Madison, L. D., Lin, Y., Kopp, P., Rao, M. S., Jameson, J. L., and Reddy, J. K. (1995b). Thyroid hormone (T3) inhibits ciprofibrate-induced transcription of genes encoding beta-oxidation enzymes: cross talk between peroxisome proliferator and T3 signaling pathways. *Proc Natl Acad Sci U S A* 92, 11593-7.
- Clarke, S. D., and Jump, D. B. (1994). Dietary polyunsaturated fatty acid regulation of gene transcription. *Annu Rev Nutr* 14, 83-98.
- Colville-Nash, P. R., Qureshi, S. S., Willis, D., and Willoughby, D. A. (1998). Inhibition of inducible nitric oxide synthase by peroxisome proliferator-activated receptor agonists: correlation with induction of heme oxygenase 1. *J Immunol* 161, 978-84.
- Cooney, A. J., Tsai, S. Y., O'Malley, B. W., and Tsai, M. J. (1992). Chicken ovalbumin upstream promoter transcription factor (COUP-TF) dimers bind to different GGTC A response elements, allowing COUP-TF to repress hormonal induction of the vitamin D3, thyroid hormone, and retinoic acid receptors. *Mol Cell Biol* 12, 4153-63.

- Cooney, A. J., Leng, X., Tsai, S. Y., O'Malley, B. W., and Tsai, M. J. (1993). Multiple mechanisms of chicken ovalbumin upstream promoter transcription factor-dependent repression of transactivation by the vitamin D, thyroid hormone, and retinoic acid receptors. *J Biol Chem* *268*, 4152-60.
- Coppolino, M. G., and Dedhar, S. (1998). Calreticulin. *Int J Biochem Cell Biol* *30*, 553-8.
- Costa, R. H., Grayson, D. R., Xanthopoulos, K. G., and Darnell, J. E., Jr. (1988). A liver-specific DNA-binding protein recognizes multiple nucleotide sites in regulatory regions of transthyretin,  $\alpha$ 1-antitrypsin, albumin, and simian virus 40 genes. *Proc Natl Acad Sci U S A* *85*, 3840-4.
- Costa, R. H., Grayson, D. R., and Darnell, J. E., Jr. (1989). Multiple hepatocyte-enriched nuclear factors function in the regulation of transthyretin and  $\alpha$ 1-antitrypsin genes. *Mol Cell Biol* *9*, 1415-25.
- Cullingford, T. E., Bhakoo, K., Peuchen, S., Dolphin, C. T., Patel, R., and Clark, J. B. (1998). Distribution of mRNAs encoding the peroxisome proliferator-activated receptor alpha, beta, and gamma and the retinoid X receptor  $\alpha$ ,  $\beta$ , and  $\gamma$  in rat central nervous system. *J Neurochem* *70*, 1366-75.
- Damm, K., Thompson, C. C., and Evans, R. M. (1989). Protein encoded by v-erbA functions as a thyroid-hormone receptor antagonist. *Nature* *339*, 593-7.
- Danielian, P. S., White, R., Lees, J. A., and Parker, M. G. (1992). Identification of a conserved region required for hormone dependent transcriptional activation by steroid hormone receptors. *Embo J* *11*, 1025-33.
- Dedhar, S., Rennie, P. S., Shago, M., Hagesteijn, C. Y., Yang, H., Filmus, J., Hawley, R. G., Bruchofsky, N., Cheng, H., Matusik, R. J., and et al. (1994). Inhibition of nuclear hormone receptor activity by calreticulin. *Nature* *367*, 480-3.
- Dennis, G., June, C. H., Mizuguchi, J., Ohara, J., Witherspoon, K., Finkelman, F. D., McMillan, V., and Mond, J. J. (1987). Glucocorticoids suppress calcium mobilization and phospholipid hydrolysis in anti-Ig antibody-stimulated B cells. *J Immunol* *139*, 2516-23.
- Desai, D., Michalak, M., Singh, N. K., and Niles, R. M. (1996). Inhibition of retinoic acid receptor function and retinoic acid-regulated gene expression in mouse melanoma cells by calreticulin. A potential pathway for cyclic AMP regulation of retinoid action. *J Biol Chem* *271*, 15153-9.

- Desvergne, B. (1994). How do thyroid hormone receptors bind to structurally diverse response elements? *Mol Cell Endocrinol* *100*, 125-31.
- Desvergne, B., A. I. J., Devchand, P. R., and Wahli, W. (1998). The peroxisome proliferator-activated receptors at the cross-road of diet and hormonal signalling. *J Steroid Biochem Mol Biol* *65*, 65-74.
- Digby, J. E., Montague, C. T., Sewter, C. P., Sanders, L., Wilkison, W. O., O'Rahilly, S., and Prins, J. B. (1998). Thiazolidinedione exposure increases the expression of uncoupling protein 1 in cultured human preadipocytes. *Diabetes* *47*, 138-41.
- Dignam, J. D., Lebovitz, R. M., and Roeder, R. G. (1983). Accurate transcription initiation by RNA polymerase II in a soluble extract from isolated mammalian nuclei. *Nucleic Acids Res* *11*, 1475-89.
- Ding, X. F., Anderson, C. M., Ma, H., Hong, H., Uht, R. M., Kushner, P. J., and Stallcup, M. R. (1998). Nuclear receptor-binding sites of coactivators glucocorticoid receptor interacting protein 1 (GRIP1) and steroid receptor coactivator 1 (SRC-1): multiple motifs with different binding specificities. *Mol Endocrinol* *12*, 302-13.
- DiRenzo, J., Soderstrom, M., Kurokawa, R., Ogliastro, M. H., Ricote, M., Ingrey, S., Horlein, A., Rosenfeld, M. G., and Glass, C. K. (1997). Peroxisome proliferator-activated receptors and retinoic acid receptors differentially control the interactions of retinoid X receptor heterodimers with ligands, coactivators, and corepressors. *Mol Cell Biol* *17*, 2166-76.
- Dogra, S. C., Whitelaw, M. L., and May, B. K. (1998). Transcriptional activation of cytochrome P450 genes by different classes of chemical inducers. *Clin Exp Pharmacol Physiol* *25*, 1-9.
- Dowell, P., Ishmael, J. E., Avram, D., Peterson, V. J., Nevriy, D. J., and Leid, M. (1997a). p300 functions as a coactivator for the peroxisome proliferator-activated receptor alpha. *J Biol Chem* *272*, 33435-43.
- Dowell, P., Peterson, V. J., Zabriskie, T. M., and Leid, M. (1997b). Ligand-induced peroxisome proliferator-activated receptor alpha conformational change. *J Biol Chem* *272*, 2013-20.
- Dreyer, C., Krey, G., Keller, H., Givel, F., Helftenbein, G., and Wahli, W. (1992). Control of the peroxisomal beta-oxidation pathway by a novel family of nuclear hormone receptors. *Cell* *68*, 879-87.

- Dreyer, C., Keller, H., Mahfoudi, A., Laudet, V., Krey, G., and Wahli, W. (1993). Positive regulation of the peroxisomal beta-oxidation pathway by fatty acids through activation of peroxisome proliferator-activated receptors (PPAR). *Biol Cell* 77, 67-76.
- Duncan, S. A., Navas, M. A., Dufort, D., Rossant, J., and Stoffel, M. (1998). Regulation of a transcription factor network required for differentiation and metabolism. *Science* 281, 692-5.
- Dupuis, M., Schaerer, E., Krause, K. H., and Tschopp, J. (1993). The calcium-binding protein calreticulin is a major constituent of lytic granules in cytolytic T lymphocytes. *J Exp Med* 177, 1-7.
- Dussault, I., and Giguere, V. (1997). Differential regulation of the N-myc proto-oncogene by ROR alpha and RVR, two orphan members of the superfamily of nuclear hormone receptors. *Mol Cell Biol* 17, 1860-7.
- Eaton, S., Bartlett, K., and Pourfarzam, M. (1996). Mammalian mitochondrial beta-oxidation. *Biochem J* 320, 345-57.
- Elstner, E., Muller, C., Koshizuka, K., Williamson, E. A., Park, D., Asou, H., Shintaku, P., Said, J. W., Heber, D., and Koeffler, H. P. (1998). Ligands for peroxisome proliferator-activated receptor gamma and retinoic acid receptor inhibit growth and induce apoptosis of human breast cancer cells in vitro and in BNX mice. *Proc Natl Acad Sci U S A* 95, 8806-11.
- Evans, R. M. (1988). The steroid and thyroid hormone receptor superfamily. *Science* 240, 889-95.
- Fajas, L., Fruchart, J. C., and Auwerx, J. (1998). Transcriptional control of adipogenesis. *Curr Opin Cell Biol* 10, 165-73.
- Fondell, J. D., Roy, A. L., and Roeder, R. G. (1993). Unliganded thyroid hormone receptor inhibits formation of a functional preinitiation complex: implications for active repression. *Genes Dev* 7, 1400-10.
- Fondell, J. D., Ge, H., and Roeder, R. G. (1996). Ligand induction of a transcriptionally active thyroid hormone receptor coactivator complex. *Proc Natl Acad Sci U S A* 93, 8329-33.
- Forman, B. M., Yang, C. R., Stanley, F., Casanova, J., and Samuels, H. H. (1988). c-erbA protooncogenes mediate thyroid hormone-dependent and independent regulation of the rat growth hormone and prolactin genes. *Mol Endocrinol* 2, 902-11.

Forman, B. M., Tontonoz, P., Chen, J., Brun, R. P., Spiegelman, B. M., and Evans, R. M. (1995a). 15-Deoxy- $\Delta^{12,14}$ -prostaglandin J<sub>2</sub> is a ligand for the adipocyte determination factor PPAR gamma. *Cell* 83, 803-12.

Forman, B.M., Umesono, K., Chen, J., and Evans, R.M. (1995b). Unique response pathways are established by allosteric interactions among nuclear hormone receptors. *Cell* 81, 541-50.

Forman, B. M., Chen, J., and Evans, R. M. (1996). The peroxisome proliferator-activated receptors: ligands and activators. *Ann N Y Acad Sci* 804, 266-75.

Forman, B. M., Chen, J., and Evans, R. M. (1997a). Hypolipidemic drugs, polyunsaturated fatty acids, and eicosanoids are ligands for peroxisome proliferator-activated receptors alpha and delta. *Proc Natl Acad Sci U S A* 94, 4312-7.

Forman, B. M., Ruan, B., Chen, J., Schroepfer, G. J., Jr., and Evans, R. M. (1997b). The orphan nuclear receptor LXR $\alpha$  is positively and negatively regulated by distinct products of mevalonate metabolism. *Proc Natl Acad Sci U S A* 94, 10588-93.

Fraser, J. D., Martinez, V., Straney, R., and Briggs, M. R. (1998). DNA binding and transcription activation specificity of hepatocyte nuclear factor 4. *Nucleic Acids Res* 26, 2702-7.

Gearing, K. L., Gottlicher, M., Teboul, M., Widmark, E., and Gustafsson, J. A. (1993). Interaction of the peroxisome-proliferator-activated receptor and retinoid X receptor. *Proc Natl Acad Sci U S A* 90, 1440-4.

Gharbi-Chihi, J., Teboul, M., Bismuth, J., Bonne, J., and Torresani, J. (1993). Increase of adipose differentiation by hypolipidemic fibrates drugs in Ob 17 preadipocytes: requirement for thyroid hormones. *Biochim Biophys Acta* 1177, 8-14.

Giguere, V., Tini, M., Flock, G., Ong, E., Evans, R. M., and Otulakowski, G. (1994). Isoform-specific amino-terminal domains dictate DNA-binding properties of ROR alpha, a novel family of orphan hormone nuclear receptors. *Genes Dev* 8, 538-53.

Giguere, V., McBroom, L. D., and Flock, G. (1995). Determinants of target gene specificity for ROR alpha 1: monomeric DNA binding by an orphan nuclear receptor. *Mol Cell Biol* 15, 2517-26.

Glass, C. K., Holloway, J. M., Devary, O. V., and Rosenfeld, M. G. (1988). The thyroid hormone receptor binds with opposite transcriptional effects to a common sequence motif in thyroid hormone and estrogen response elements. *Cell* 54, 313-23.



- Glass, C. K. (1994). Differential recognition of target genes by nuclear receptor monomers, dimers, and heterodimers. *Endocr Rev* 15, 391-407.
- Glass, C. K., Rose, D. W., and Rosenfeld, M. G. (1997). Nuclear receptor coactivators. *Curr Opin Cell Biol* 9, 222-32.
- Goodridge, A. G. (1987). Dietary regulation of gene expression: enzymes involved in carbohydrate and lipid metabolism. *Annu Rev Nutr* 7, 157-85.
- Gottlicher, M., Widmark, E., Li, Q., and Gustafsson, J. A. (1992). Fatty acids activate a chimera of the clofibrilic acid-activated receptor and the glucocorticoid receptor. *Proc Natl Acad Sci U S A* 89, 4653-7.
- Granneman, J., Skoff, R., and Yang, X. (1998). Member of the peroxisome proliferator-activated receptor family of transcription factors is differentially expressed by oligodendrocytes. *J Neurosci Res* 51, 563-73.
- Graupner, G., Wills, K. N., Tzukerman, M., Zhang, X. K., and Pfahl, M. (1989). Dual regulatory role for thyroid-hormone receptors allows control of retinoic-acid receptor activity. *Nature* 340, 653-6.
- Green, S. (1993). Nuclear hormone receptors. Promiscuous liaisons. *Nature* 361, 590-1.
- Green, S., and Wahli, W. (1994). Peroxisome proliferator-activated receptors: finding the orphan a home. *Mol Cell Endocrinol* 100, 149-53.
- Gregori, C., Porteu, A., Lopez, S., Kahn, A., and Pichard, A. L. (1998). Characterization of the aldolase B intronic enhancer. *J Biol Chem* 273, 25237-43.
- Gronemeyer, H. (1992). Control of transcription activation by steroid hormone receptors. *Faseb J* 6, 2524-9.
- Grunstein, M. (1997). Histone acetylation in chromatin structure and transcription. *Nature* 389, 349-52.
- Gulick, T., Cresci, S., Caira, T., Moore, D. D., and Kelly, D. P. (1994). The peroxisome proliferator-activated receptor regulates mitochondrial fatty acid oxidative enzyme gene expression. *Proc Natl Acad Sci U S A* 91, 11012-6.
- Hall, R. K., Sladek, F. M., and Granner, D. K. (1995). The orphan receptors COUP-TF and HNF-4 serve as accessory factors required for induction of phosphoenolpyruvate carboxykinase gene transcription by glucocorticoids. *Proc Natl Acad Sci U S A* 92, 412-6.

Hallaq, H., Smith, T. W., and Leaf, A. (1992). Modulation of dihydropyridine-sensitive calcium channels in heart cells by fish oil fatty acids. *Proc Natl Acad Sci U S A* *89*, 1760-4.

Hashimoto, T. (1996). Peroxisomal beta-oxidation: enzymology and molecular biology. *Ann N Y Acad Sci* *804*, 86-98.

Hassig, C. A., Fleischer, T. C., Billin, A. N., Schreiber, S. L., and Ayer, D. E. (1997). Histone deacetylase activity is required for full transcriptional repression by mSin3A. *Cell* *89*, 341-7.

Hawkins, J. M., Jones, W. E., Bonner, F. W., and Gibson, G. G. (1987). The effect of peroxisome proliferators on microsomal, peroxisomal, and mitochondrial enzyme activities in the liver and kidney. *Drug Metab Rev* *18*, 441-515.

Hazlerigg, D. G., Barrett, P., Hastings, M. H., and Morgan, P. J. (1996). Are nuclear receptors involved in pituitary responsiveness to melatonin? *Mol Cell Endocrinol* *123*, 53-9.

Heery, D. M., Kalkhoven, E., Hoare, S., and Parker, M. G. (1997). A signature motif in transcriptional co-activators mediates binding to nuclear receptors. *Nature* *387*, 733-6.

Heinzel, T., Lavinsky, R. M., Mullen, T. M., Soderstrom, M., Laherty, C. D., Torchia, J., Yang, W. M., Brard, G., Ngo, S. D., Davie, J. R., Seto, E., Eisenman, R. N., Rose, D. W., Glass, C. K., and Rosenfeld, M. G. (1997). A complex containing N-CoR, mSin3 and histone deacetylase mediates transcriptional repression. *Nature* *387*, 43-8.

Hertz, R., Bishara-Shieban, J., and Bar-Tana, J. (1995). Mode of action of peroxisome proliferators as hypolipidemic drugs. Suppression of apolipoprotein C-III. *J Biol Chem* *270*, 13470-5.

Hertz, R., Bishara-Shieban, J., and Bar-Tana, J. (1996a). Hypolipidemic effect of peroxisome proliferators. Transcriptional suppression of apolipoprotein C-III. *Ann N Y Acad Sci* *804*, 731-3.

Hertz, R., Seckbach, M., Zakin, M. M., and Bar-Tana, J. (1996b). Transcriptional suppression of the transferrin gene by hypolipidemic peroxisome proliferators. *J Biol Chem* *271*, 218-24.

Hertz, R., Magenheim, J., Berman, I., and Bar-Tana, J. (1998). Fatty acyl-CoA thioesters are ligands of hepatic nuclear factor-4 $\alpha$ . *Nature* *392*, 512-6.

- Hess, R., Staubli, W., and Riess, W. (1965). Nature of the hepatomegalic effect produced by ethyl-chlorophenoxy- isobutyrate in the rat. *Nature* 208, 856-8.
- Hijikata, M., Ishii, N., Kagamiyama, H., Osumi, T., and Hashimoto, T. (1987). Structural analysis of cDNA for rat peroxisomal 3-ketoacyl-CoA thiolase. *J Biol Chem* 262, 8151-8.
- Hijikata, M., Wen, J. K., Osumi, T., and Hashimoto, T. (1990). Rat peroxisomal 3-ketoacyl-CoA thiolase gene. Occurrence of two closely related but differentially regulated genes. *J Biol Chem* 265, 4600-6.
- Hiltunen, J. K., Filppula, S. A., Koivuranta, K. T., Siivari, K., Qin, Y. M., and Hayrinen, H. M. (1996). Peroxisomal beta-oxidation and polyunsaturated fatty acids. *Ann N Y Acad Sci* 804, 116-28.
- Hirose, T., Smith, R. J., and Jetten, A. M. (1994). ROR gamma: the third member of ROR/RZR orphan receptor subfamily that is highly expressed in skeletal muscle. *Biochem Biophys Res Commun* 205, 1976-83.
- Hong, H., Kohli, K., Trivedi, A., Johnson, D. L., and Stallcup, M. R. (1996). GRIP1, a novel mouse protein that serves as a transcriptional coactivator in yeast for the hormone binding domains of steroid receptors. *Proc Natl Acad Sci U S A* 93, 4948-52.
- Hopps, H. E., Bernham, B. C., Nisalak, A., Tjio, J. H., and Smadel, J. E. (1963). Biologic characteristics of a continuous kidney cell line derived from the African green monkey. *J. Immunol.* 91, 416-424.
- Horvai, A. E., Xu, L., Korzus, E., Brard, G., Kalafus, D., Mullen, T. M., Rose, D. W., Rosenfeld, M. G., and Glass, C. K. (1997). Nuclear integration of JAK/STAT and Ras/AP-1 signaling by CBP and p300. *Proc Natl Acad Sci U S A* 94, 1074-9.
- Horwitz, K. B., Jackson, T. A., Bain, D. L., Richer, J. K., Takimoto, G. S., and Tung, L. (1996). Nuclear receptor coactivators and corepressors. *Mol Endocrinol* 10, 1167-77.
- Hsu, M. H., Palmer, C. N. A., Song, W., Griffin, K. J., and Johnson, E. F. (1998). A carboxyl-terminal extension of the zinc finger domain contributes to the specificity and polarity of peroxisome proliferator-activated receptor DNA binding. *J Biol Chem* 273, 27988-97.
- Hu, E., Tontonoz, P., and Spiegelman, B. M. (1995). Transdifferentiation of myoblasts by the adipogenic transcription factors PPAR gamma and C/EBP alpha. *Proc Natl Acad Sci U S A* 92, 9856-60.

- Huss, J. M., and Kasper, C. B. (1998). Nuclear receptor involvement in the regulation of rat cytochrome P450 3A23 expression. *J Biol Chem* 273, 16155-62.
- Ibrahimi, A., Teboul, L., Gaillard, D., Amri, E. Z., Ailhaud, G., Young, P., Cawthorne, M. A., and Grimaldi, P. A. (1994). Evidence for a common mechanism of action for fatty acids and thiazolidinedione antidiabetic agents on gene expression in preadipose cells. *Mol Pharmacol* 46, 1070-6.
- IJpenberg, A., Jeannin, E., Wahli, W., and Desvergne, B. (1997). Polarity and specific sequence requirements of peroxisome proliferator-activated receptor (PPAR)/retinoid X receptor heterodimer binding to DNA. A functional analysis of the malic enzyme gene PPAR response element. *J Biol Chem* 272, 20108-17.
- Ikeda, M., Rhee, M., and Chin, W. W. (1994). Thyroid hormone receptor monomer, homodimer, and heterodimer (with retinoid-X receptor) contact different nucleotide sequences in thyroid hormone response elements. *Endocrinology* 135, 1628-38.
- Inoue, I., Noji, S., Awata, T., Takahashi, K., Nakajima, T., Sonoda, M., Komoda, T., and Katayama, S. (1998). Bezafibrate has an antioxidant effect: peroxisome proliferator-activated receptor alpha is associated with Cu<sup>2+</sup>, Zn<sup>2+</sup>-superoxide dismutase in the liver. *Life Sci* 63, 135-44.
- Issemann, I., and Green, S. (1990). Activation of a member of the steroid hormone receptor superfamily by peroxisome proliferators. *Nature* 347, 645-50.
- Issemann, I., Prince, R., Tugwood, J., and Green, S. (1992). A role for fatty acids and liver fatty acid binding protein in peroxisome proliferation? *Biochem Soc Trans* 20, 824-7.
- Issemann, I., Prince, R. A., Tugwood, J. D., and Green, S. (1993). The peroxisome proliferator-activated receptor:retinoid X receptor heterodimer is activated by fatty acids and fibrate hypolipidaemic drugs. *J Mol Endocrinol* 11, 37-47.
- Jackson, T. A., Richer, J. K., Bain, D. L., Takimoto, G. S., Tung, L., and Horwitz, K. B. (1997). The partial agonist activity of antagonist-occupied steroid receptors is controlled by a novel hinge domain-binding coactivator L7/SPA and the corepressors N-CoR or SMRT. *Mol Endocrinol* 11, 693-705.
- Jain, S., Pulikuri, S., Zhu, Y., Qi, C., Kanwar, Y. S., Yeldandi, A. V., Rao, M. S., and Reddy, J. K. (1998). Differential expression of the peroxisome proliferator-activated receptor gamma (PPAR $\gamma$ ) and its coactivators steroid receptor coactivator-1 and PPAR-binding protein PBP in the brown fat, urinary bladder, colon, and breast of the mouse. *Am J Pathol* 153, 349-54.

- Jiang, G., Nepomuceno, L., Hopkins, K., and Sladek, F. M. (1995). Exclusive homodimerization of the orphan receptor hepatocyte nuclear factor 4 defines a new subclass of nuclear receptors. *Mol Cell Biol* 15, 5131-43.
- Jiang, C., Ting, A. T., and Seed, B. (1998). PPAR-gamma agonists inhibit production of monocyte inflammatory cytokines. *Nature* 391, 82-6.
- Johnson, E. F., Palmer, C. N., Griffin, K. J., and Hsu, M. H. (1996). Role of the peroxisome proliferator-activated receptor in cytochrome P450 4A gene regulation. *Faseb J* 10, 1241-8.
- Johnson, T. E., Holloway, M. K., Vogel, R., Rutledge, S. J., Perkins, J. J., Rodan, G. A., and Schmidt, A. (1997). Structural requirements and cell-type specificity for ligand activation of peroxisome proliferator-activated receptors. *J Steroid Biochem Mol Biol* 63, 1-8.
- Jow, L., and Mukherjee, R. (1995). The human peroxisome proliferator-activated receptor (PPAR) subtype NUC1 represses the activation of hPPAR alpha and thyroid hormone receptors. *J Biol Chem* 270, 3836-40.
- Juge-Aubry, C. E., Gorla-Bajszczak, A., Pernin, A., Lemberger, T., Wahli, W., Burger, A. G., and Meier, C. A. (1995). Peroxisome proliferator-activated receptor mediates cross-talk with thyroid hormone receptor by competition for retinoid X receptor. Possible role of a leucine zipper-like heptad repeat. *J Biol Chem* 270, 18117-22.
- Juge-Aubry, C., Pernin, A., Favez, T., Burger, A. G., Wahli, W., Meier, C. A., and Desvergne, B. (1997). DNA binding properties of peroxisome proliferator-activated receptor subtypes on various natural peroxisome proliferator response elements. Importance of the 5'-flanking region. *J Biol Chem* 272, 25252-9.
- Jump, D. B., Clarke, S. D., Thelen, A., Liimatta, M., Ren, B., and Badin, M. V. (1997). Dietary fat, genes, and human health. *Adv Exp Med Biol* 422, 167-76.
- Kaikaus, R. M., Chan, W. K., Lysenko, N., Ray, R., Ortiz de Montellano, P. R., and Bass, N. M. (1993). Induction of peroxisomal fatty acid  $\beta$ -oxidation and liver fatty acid-binding protein by peroxisome proliferators. Mediation via the cytochrome P-450IVA1 omega-hydroxylase pathway. *J Biol Chem* 268, 9593-603.
- Kamei, Y., Xu, L., Heinzl, T., Torchia, J., Kurokawa, R., Gloss, B., Lin, S. C., Heyman, R. A., Rose, D. W., Glass, C. K., and Rosenfeld, M. G. (1996). A CBP integrator complex mediates transcriptional activation and AP-1 inhibition by nuclear receptors. *Cell* 85, 403-14.

- Katzenellenbogen, J. A., O'Malley, B. W., and Katzenellenbogen, B. S. (1996). Tripartite steroid hormone receptor pharmacology: interaction with multiple effector sites as a basis for the cell- and promoter-specific action of these hormones. *Mol Endocrinol* 10, 119-31.
- Keightley, M. C. (1998). Steroid receptor isoforms: exception or rule? *Mol Cell Endocrinol* 137, 1-5.
- Keller, H., Dreyer, C., Medin, J., Mahfoudi, A., Ozato, K., and Wahli, W. (1993). Fatty acids and retinoids control lipid metabolism through activation of peroxisome proliferator-activated receptor-retinoid X receptor heterodimers. *Proc Natl Acad Sci U S A* 90, 2160-4.
- Kindl, H. (1993). Fatty acid degradation in plant peroxisomes: function and biosynthesis of the enzymes involved. *Biochimie* 75, 225-30.
- Kitajima, K., Nagaya, T., and Jameson, J. L. (1995). Dominant negative and DNA-binding properties of mutant thyroid hormone receptors that are defective in homodimerization but not heterodimerization. *Thyroid* 5, 343-53.
- Kliwer, S. A., Umesono, K., Mangelsdorf, D. J., and Evans, R. M. (1992a). Retinoid X receptor interacts with nuclear receptors in retinoic acid, thyroid hormone and vitamin D3 signalling. *Nature* 355, 446-9.
- Kliwer, S. A., Umesono, K., Noonan, D. J., Heyman, R. A., and Evans, R. M. (1992b). Convergence of 9-cis retinoic acid and peroxisome proliferator signalling pathways through heterodimer formation of their receptors. *Nature* 358, 771-4.
- Kliwer, S. A., Lenhard, J. M., Willson, T. M., Patel, I., Morris, D. C., and Lehmann, J. M. (1995). A prostaglandin J<sub>2</sub> metabolite binds peroxisome proliferator-activated receptor gamma and promotes adipocyte differentiation. *Cell* 83, 813-9.
- Kliwer, S. A., Sundseth, S. S., Jones, S. A., Brown, P. J., Wisely, G. B., Koble, C. S., Devchand, P., Wahli, W., Willson, T. M., Lenhard, J. M., and Lehmann, J. M. (1997). Fatty acids and eicosanoids regulate gene expression through direct interactions with peroxisome proliferator-activated receptors alpha and gamma. *Proc Natl Acad Sci U S A* 94, 4318-23.
- Krause, K. H., and Michalak, M. (1997). Calreticulin. *Cell* 88, 439-43.
- Krey, G., Braissant, O., L'Horset, F., Kalkhoven, E., Perroud, M., Parker, M. G., and Wahli, W. (1997). Fatty acids, eicosanoids, and hypolipidemic agents identified as ligands of peroxisome proliferator-activated receptors by coactivator-dependent receptor ligand assay. *Mol Endocrinol* 11, 779-91.

- Kurokawa, R., DiRenzo, J., Boehm, M., Sugarman, J., Gloss, B., Rosenfeld, M. G., Heyman, R. A., and Glass, C. K. (1994). Regulation of retinoid signalling by receptor polarity and allosteric control of ligand binding. *Nature* 371, 528-31.
- Kurokawa, R., Soderstrom, M., Horlein, A., Halachmi, S., Brown, M., Rosenfeld, M. G., and Glass, C. K. (1995). Polarity-specific activities of retinoic acid receptors determined by a co-repressor. *Nature* 377, 451-4.
- Ladiaz, J. A., Hadzopoulou-Cladaras, M., Kardassis, D., Cardot, P., Cheng, J., Zannis, V., and Cladaras, C. (1992). Transcriptional regulation of human apolipoprotein genes ApoB, ApoCIII, and ApoAII by members of the steroid hormone receptor superfamily HNF- 4, ARP-1, EAR-2, and EAR-3. *J Biol Chem* 267, 15849-60.
- Ladiaz, J. A. (1994). Convergence of multiple nuclear receptor signaling pathways onto the long terminal repeat of human immunodeficiency virus-1. *J Biol Chem* 269, 5944-51.
- Laherty, C. D., Yang, W. M., Sun, J. M., Davie, J. R., Seto, E., and Eisenman, R. N. (1997). Histone deacetylases associated with the mSin3 corepressor mediate mad transcriptional repression. *Cell* 89, 349-56.
- Laherty, C. D., Billin, A. N., Lavinsky, R. M., Yochum, G. S., Bush, A. C., Sun, J. M., Mullen, T. M., Davie, J. R., Rose, D. W., Glass, C. K., Rosenfeld, M. G., Ayer, D. E., and Eisenman, R. N. (1998). SAP30, a component of the mSin3 corepressor complex involved in N-CoR- mediated repression by specific transcription factors. *Mol Cell* 2, 33-42.
- Lake, B. G., and Gray, T. J. (1985). Species differences in hepatic peroxisome proliferation. *Biochem Soc Trans* 13, 859-61.
- Lambe, K. G., and Tugwood, J. D. (1996). A human peroxisome-proliferator-activated receptor-gamma is activated by inducers of adipogenesis, including thiazolidinedione drugs. *Eur J Biochem* 239, 1-7.
- Latruffe, N., and Vamecq, J. (1997). Peroxisome proliferators and peroxisome proliferator activated receptors (PPARs) as regulators of lipid metabolism. *Biochimie* 79, 81-94.
- Laudet, V., Hanni, C., Coll, J., Catzeflis, F., and Stehelin, D. (1992). Evolution of the nuclear receptor gene superfamily. *Embo J* 11, 1003-13.
- Lavinsky, R. M., Jepsen, K., Heinzl, T., Torchia, J., Mullen, T. M., Schiff, R., Del-Rio, A. L., Ricote, M., Ngo, S., Gemsch, J., Hilsenbeck, S. G., Osborne, C. K., Glass, C. K., Rosenfeld, M. G., and Rose, D. W. (1998). Diverse signaling pathways modulate nuclear receptor recruitment of N- CoR and SMRT complexes. *Proc Natl Acad Sci U S A* 95, 2920-5.

- Lazar, M. A. (1991a). Steroid and thyroid hormone receptors. *Endocrinol Metab Clin North Am* 20, 681-95.
- Lazar, M. A., Berrodin, T. J., and Harding, H. P. (1991b). Differential DNA binding by monomeric, homodimeric, and potentially heteromeric forms of the thyroid hormone receptor. *Mol Cell Biol* 11, 5005-15.
- Lazarow, P. B., and De Duve, C. (1976). A fatty acyl-CoA oxidizing system in rat liver peroxisomes; enhancement by clofibrate, a hypolipidemic drug. *Proc Natl Acad Sci U S A* 73, 2043-6.
- Lee, J. W., Choi, H. S., Gyuris, J., Brent, R., and Moore, D. D. (1995). Two classes of proteins dependent on either the presence or absence of thyroid hormone for interaction with the thyroid hormone receptor. *Mol Endocrinol* 9, 243-54.
- Lee, M. J., Gee, P., and Beard, S. E. (1997). Detection of peroxisome proliferators using a reporter construct derived from the rat acyl-CoA oxidase promoter in the rat liver cell line H-4-II-E. *Cancer Res* 57, 1575-9.
- Lee, S. K., Na, S. Y., Kim, H. J., Soh, J., Choi, H. S., and Lee, J. W. (1998). Identification of critical residues for heterodimerization within the ligand-binding domain of retinoid X receptor. *Mol Endocrinol* 12, 325-32.
- Leers, J., Treuter, E., and Gustafsson, J. A. (1998). Mechanistic principles in NR box-dependent interaction between nuclear hormone receptors and the coactivator TIF2. *Mol Cell Biol* 18, 6001-13.
- Lehmann, J. M., Moore, L. B., Smith-Oliver, T. A., Wilkison, W. O., Willson, T. M., and Kliewer, S. A. (1995). An antidiabetic thiazolidinedione is a high affinity ligand for peroxisome proliferator-activated receptor gamma (PPAR gamma). *J Biol Chem* 270, 12953-6.
- Lehmann, J. M., Lenhard, J. M., Oliver, B. B., Ringold, G. M., and Kliewer, S. A. (1997). Peroxisome proliferator-activated receptors alpha and gamma are activated by indomethacin and other non-steroidal anti-inflammatory drugs. *J Biol Chem* 272, 3406-10.
- Lehninger, A. L., Nelson, D. L., and Cox, M. M. (1993). *Principles of Biochemistry* (New York, NY: Worth Publishers), pp. 485-502.
- Leng, X., Tsai, S. Y., O'Malley, B. W., and Tsai, M. J. (1993). Ligand-dependent conformational changes in thyroid hormone and retinoic acid receptors are potentially enhanced by heterodimerization with retinoic X receptor. *J Steroid Biochem Mol Biol* 46, 643-61.



Leung-Hagesteijn, C. Y., Milankov, K., Michalak, M., Wilkins, J., and Dedhar, S. (1994). Cell attachment to extracellular matrix substrates is inhibited upon downregulation of expression of calreticulin, an intracellular integrin alpha-subunit-binding protein. *J Cell Sci* *107*, 589-600.

Li, H., Gomes, P. J., and Chen, J. D. (1997a). RAC3, a steroid/nuclear receptor-associated coactivator that is related to SRC-1 and TIF2. *Proc Natl Acad Sci U S A* *94*, 8479-84.

Li, H., Leo, C., Schroen, D. J., and Chen, J. D. (1997b). Characterization of receptor interaction and transcriptional repression by the corepressor SMRT. *Mol Endocrinol* *11*, 2025-37.

Liu, R. T., Suzuki, S., Takeda, T., and DeGroot, L. J. (1996). An artificial thyroid hormone receptor mutant without DNA binding can have dominant negative effect. *Mol Cell Endocrinol* *120*, 85-93.

Liu, Y., Takeshita, A., Misiti, S., Chin, W. W., and Yen, P. M. (1998). Lack of coactivator interaction can be a mechanism for dominant negative activity by mutant thyroid hormone receptors. *Endocrinology* *139*, 4197-204.

Llanos, S., Caelles, C., Azorin, I., Renau-Piqueras, J., Fernandez-Luna, J. L., Bosca, L., and Munoz, A. (1998). The c-erbA alpha protooncogene induces apoptosis in glial cells via a protein kinase C- and bcl-2-suppressible mechanism. *J Neurochem* *70*, 2315-26.

Lock, E. A., Mitchell, A. M., and Elcombe, C. R. (1989). Biochemical mechanisms of induction of hepatic peroxisome proliferation. *Annu Rev Pharmacol Toxicol* *29*, 145-63.

Lucas, P. C., and Granner, D. K. (1992). Hormone response domains in gene transcription. *Annu Rev Biochem* *61*, 1131-73.

Mader, S., Leroy, P., Chen, J. Y., and Chambon, P. (1993). Multiple parameters control the selectivity of nuclear receptors for their response elements. Selectivity and promiscuity in response element recognition by retinoic acid receptors and retinoid X receptors. *J Biol Chem* *268*, 591-600.

Mangelsdorf, D. J., and Evans, R. M. (1995). The RXR heterodimers and orphan receptors. *Cell* *83*, 841-50.

Mannaerts, G. P., and Van Veldhoven, P. P. (1993). Metabolic pathways in mammalian peroxisomes. *Biochimie* *75*, 147-58.

- Mannaerts, G. P., and van Veldhoven, P. P. (1996). Functions and organization of peroxisomal  $\beta$ -oxidation. *Ann N Y Acad Sci* 804, 99-115.
- Marcus, S. L., Miyata, K. S., Zhang, B., Subramani, S., Rachubinski, R. A., and Capone, J. P. (1993). Diverse peroxisome proliferator-activated receptors bind to the peroxisome proliferator-responsive elements of the rat hydratase/dehydrogenase and fatty acyl-CoA oxidase genes but differentially induce expression. *Proc Natl Acad Sci U S A* 90, 5723-7.
- Marcus, S. L., Capone, J. P., and Rachubinski, R. A. (1996). Identification of COUP-TFII as a peroxisome proliferator response element binding factor using genetic selection in yeast: COUP-TFII activates transcription in yeast but antagonizes PPAR signaling in mammalian cells. *Mol Cell Endocrinol* 120, 31-9.
- Martin, G., Schoonjans, K., Lefebvre, A. M., Staels, B., and Auwerx, J. (1997). Coordinate regulation of the expression of the fatty acid transport protein and acyl-CoA synthetase genes by PPAR $\alpha$  and PPAR $\gamma$  activators. *J Biol Chem* 272, 28210-7.
- Maruyama, K., Tsukada, T., Ohkura, N., Bandoh, S., Hosono, T., and Yamaguchi, K. (1998). The NGFI-B subfamily of the nuclear receptor superfamily (review). *Int J Oncol* 12, 1237-43.
- Mascaro, C., Acosta, E., Ortiz, J. A., Marrero, P. F., Hegardt, F. G., and Haro, D. (1998). Control of human muscle-type carnitine palmitoyltransferase I gene transcription by peroxisome proliferator-activated receptor. *J Biol Chem* 273, 8560-3.
- McBroom, L. D., Flock, G., and Giguere, V. (1995). The nonconserved hinge region and distinct amino-terminal domains of the ROR alpha orphan nuclear receptor isoforms are required for proper DNA bending and ROR alpha-DNA interactions. *Mol Cell Biol* 15, 796-808.
- McInerney, E. M., Rose, D. W., Flynn, S. E., Westin, S., Mullen, T. M., Krones, A., Inostroza, J., Torchia, J., Nolte, R. T., Assa-Munt, N., Milburn, M. V., Glass, C. K., and Rosenfeld, M. G. (1998). Determinants of coactivator LXXLL motif specificity in nuclear receptor transcriptional activation. *Genes Dev* 12, 3357-68.
- McQuaid, S., Russell, S. E., Withe, S. A., Pearson, C. M., Elcombe, C. R., and Humphries, P. (1987). Analysis of transcripts homologous to acyl-CoA oxidase and enoyl-CoA hydratase/3-hydroxyacyl-CoA dehydrogenase induced in rat liver by methylclofenapate. *Cancer Lett* 37, 115-21.
- Meier, C. A. (1997). Regulation of gene expression by nuclear hormone receptors. *J Recept Signal Transduct Res* 17, 319-35.

- Meier-Heusler, S. C., Zhu, X., Juge-Aubry, C., Pernin, A., Burger, A. G., Cheng, S. Y., and Meier, C. A. (1995). Modulation of thyroid hormone action by mutant thyroid hormone receptors, c-erbA alpha 2 and peroxisome proliferator-activated receptor: evidence for different mechanisms of inhibition. *Mol Cell Endocrinol* 107, 55-66.
- Mery, L., Mesaeli, N., Michalak, M., Opas, M., Lew, D. P., and Krause, K. H. (1996). Overexpression of calreticulin increases intracellular Ca<sup>2+</sup> storage and decreases store-operated Ca<sup>2+</sup> influx. *J Biol Chem* 271, 9332-9.
- Michalak, M., Milner, R. E., Burns, K., and Opas, M. (1992). Calreticulin. *Biochem J* 285, 681-92.
- Michalak, M., Burns, K., Andrin, C., Mesaeli, N., Jass, G. H., Busaan, J. L., and Opas, M. (1996). Endoplasmic reticulum form of calreticulin modulates glucocorticoid-sensitive gene expression. *J Biol Chem* 271, 29436-45.
- Mietus-Snyder, M., Sladek, F. M., Ginsburg, G. S., Kuo, C. F., Ladias, J. A., Darnell, J. E., Jr., and Karathanasis, S. K. (1992). Antagonism between apolipoprotein AI regulatory protein 1, Ear3/COUP-TF, and hepatocyte nuclear factor 4 modulates apolipoprotein CIII gene expression in liver and intestinal cells. *Mol Cell Biol* 12, 1708-18.
- Misiti, S., Schomburg, L., Yen, P. M., and Chin, W. W. (1998). Expression and hormonal regulation of coactivator and corepressor genes. *Endocrinology* 139, 2493-500.
- Missbach, M., Jagher, B., Sigg, I., Nayeri, S., Carlberg, C., and Wiesenberg, I. (1996). Thiazolidine diones, specific ligands of the nuclear receptor retinoid Z receptor/retinoid acid receptor-related orphan receptor alpha with potent antiarthritic activity. *J Biol Chem* 271, 13515-22.
- Miyamoto, T., Kaneko, A., Kakizawa, T., Yajima, H., Kamijo, K., Sekine, R., Hiramatsu, K., Nishii, Y., Hashimoto, T., and Hashizume, K. (1997). Inhibition of peroxisome proliferator signaling pathways by thyroid hormone receptor. Competitive binding to the response element. *J Biol Chem* 272, 7752-8.
- Miyata, K. S., Zhang, B., Marcus, S. L., Capone, J. P., and Rachubinski, R. A. (1993). Chicken ovalbumin upstream promoter transcription factor (COUP-TF) binds to a peroxisome proliferator-responsive element and antagonizes peroxisome proliferator-mediated signaling. *J Biol Chem* 268, 19169-72.
- Miyata, K. S., McCaw, S. E., Marcus, S. L., Rachubinski, R. A., and Capone, J. P. (1994). The peroxisome proliferator-activated receptor interacts with the retinoid X receptor in vivo. *Gene* 148, 327-30.

- Miyata, K. S., McCaw, S. E., Patel, H. V., Rachubinski, R. A., and Capone, J. P. (1996). The orphan nuclear hormone receptor LXR $\alpha$  interacts with the peroxisome proliferator-activated receptor and inhibits peroxisome proliferator signaling. *J Biol Chem* 271, 9189-92.
- Monden, T., Wondisford, F. E., and Hollenberg, A. N. (1997). Isolation and characterization of a novel ligand-dependent thyroid hormone receptor-coactivating protein. *J Biol Chem* 272, 29834-41.
- Moras, D., and Gronemeyer, H. (1998). The nuclear receptor ligand-binding domain: structure and function. *Curr Opin Cell Biol* 10, 384-91.
- Motojima, K., Peters, J. M., and Gonzalez, F. J. (1997). PPAR alpha mediates peroxisome proliferator-induced transcriptional repression of nonperoxisomal gene expression in mouse. *Biochem Biophys Res Commun* 230, 155-8.
- Motojima, K., Passilly, P., Peters, J. M., Gonzalez, F. J., and Latruffe, N. (1998). Expression of putative fatty acid transporter genes are regulated by peroxisome proliferator-activated receptor alpha and gamma activators in a tissue- and inducer-specific manner. *J Biol Chem* 273, 16710-4.
- Muerhoff, A. S., Griffin, K. J., and Johnson, E. F. (1992). The peroxisome proliferator-activated receptor mediates the induction of CYP4A6, a cytochrome P450 fatty acid  $\omega$ -hydroxylase, by clofibric acid. *J Biol Chem* 267, 19051-3.
- Murao, K., Bassyouni, H., Taylor, A. H., Wanke, I. E., and Wong, N. C. (1997). Hepatocyte nuclear factor 4 inhibits the activity of site A from the rat apolipoprotein AI gene. *Biochemistry* 36, 301-6.
- Naar, A. M., Boutin, J. M., Lipkin, S. M., Yu, V. C., Holloway, J. M., Glass, C. K., and Rosenfeld, M. G. (1991). The orientation and spacing of core DNA-binding motifs dictate selective transcriptional responses to three nuclear receptors. *Cell* 65, 1267-79.
- Nagasawa, T., Suzuki, S., Takeda, T., and DeGroot, L. J. (1997). Thyroid hormone receptor beta 1 expression in developing mouse limbs and face. *Endocrinology* 138, 1276-81.
- Nagaya, T., Madison, L. D., and Jameson, J. L. (1992). Thyroid hormone receptor mutants that cause resistance to thyroid hormone. Evidence for receptor competition for DNA sequences in target genes. *J Biol Chem* 267, 13014-9.

Nagaya, T., Fujieda, M., and Seo, H. (1998). Requirement of corepressor binding of thyroid hormone receptor mutants for dominant negative inhibition. *Biochem Biophys Res Commun* 247, 620-3.

Nagy, L., Kao, H. Y., Chakravarti, D., Lin, R. J., Hassig, C. A., Ayer, D. E., Schreiber, S. L., and Evans, R. M. (1997). Nuclear receptor repression mediated by a complex containing SMRT, mSin3A, and histone deacetylase. *Cell* 89, 373-80.

Nakshatri, H., and Bhat-Nakshatri, P. (1998). Multiple parameters determine the specificity of transcriptional response by nuclear receptors HNF-4, ARP-1, PPAR, RAR and RXR through common response elements. *Nucleic Acids Res* 26, 2491-9.

Nash, P. D., Opas, M., and Michalak, M. (1994). Calreticulin: not just another calcium-binding protein. *Mol Cell Biochem* 135, 71-8.

Nawaz, Z., Tsai, M. J., and O'Malley, B. W. (1995). Specific mutations in the ligand binding domain selectively abolish the silencing function of human thyroid hormone receptor beta. *Proc Natl Acad Sci U S A* 92, 11691-5.

Nishiyama, C., Hi, R., Osada, S., and Osumi, T. (1998). Functional interactions between nuclear receptors recognizing a common sequence element, the direct repeat motif spaced by one nucleotide (DR-1). *J Biochem (Tokyo)* 123, 1174-9.

Nishiyori, A., Tashiro, H., Kimura, A., Akagi, K., Yamamura, K., Mori, M., and Takiguchi, M. (1994). Determination of tissue specificity of the enhancer by combinatorial operation of tissue-enriched transcription factors. Both HNF-4 and C/EBP beta are required for liver-specific activity of the ornithine transcarbamylase enhancer. *J Biol Chem* 269, 1323-31.

Nolte, R. T., Wisely, G. B., Westin, S., Cobb, J. E., Lambert, M. H., Kurokawa, R., Rosenfeld, M. G., Willson, T. M., Glass, C. K., and Milburn, M. V. (1998). Ligand binding and co-activator assembly of the peroxisome proliferator-activated receptor-gamma. *Nature* 395, 137-43.

Ogryzko, V. V., Schiltz, R. L., Russanova, V., Howard, B. H., and Nakatani, Y. (1996). The transcriptional coactivators p300 and CBP are histone acetyltransferases. *Cell* 87, 953-9.

Olson, D. P., and Koenig, R. J. (1997). 5'-flanking sequences in thyroid hormone response element half-sites determine the requirement of retinoid X receptor for receptor-mediated gene expression. *J Biol Chem* 272, 9907-14.

- Opas, M., Dziak, E., Fliegel, L., and Michalak, M. (1991). Regulation of expression and intracellular distribution of calreticulin, a major calcium binding protein of nonmuscle cells. *J Cell Physiol* 149, 160-71.
- Opas, M., Szewczenko-Pawlikowski, M., Jass, G. K., Mesaeli, N., and Michalak, M. (1996). Calreticulin modulates cell adhesiveness via regulation of vinculin expression. *J Cell Biol* 135, 1913-23.
- Osada, S., Tsukamoto, T., and Osumi, T. (1996). Analysis of the essential sequence elements in the peroxisome proliferator-responsive enhancer of rat acyl-CoA oxidase gene. *Ann N Y Acad Sci* 804, 706-8.
- Osada, S., Tsukamoto, T., Takiguchi, M., Mori, M., and Osumi, T. (1997). Identification of an extended half-site motif required for the function of peroxisome proliferator-activated receptor alpha. *Genes Cells* 2, 315-27.
- Osumi, T., and Hashimoto, T. (1978). Enhancement of fatty acyl-CoA oxidizing activity in rat liver peroxisomes by di-(i-ethylhexyl)phthalate. *J Biochem (Tokyo)* 83, 1361-5.
- Osumi, T. (1993). Structure and expression of the genes encoding peroxisomal  $\beta$ -oxidation enzymes. *Biochimie* 75, 243-50.
- Osumi, T., Wen, J. K., and Hashimoto, T. (1991). Two cis-acting regulatory sequences in the peroxisome proliferator-responsive enhancer region of rat acyl-CoA oxidase gene. *Biochem Biophys Res Commun* 175, 866-71.
- Ozasa, H., Miyazawa, S., Furuta, S., Osumi, T., and Hashimoto, T. (1985). Induction of peroxisomal beta-oxidation enzymes in primary cultured rat hepatocytes by clofibric acid. *J Biochem (Tokyo)* 97, 1273-8.
- Pacot, C., Charmoillaux, M., Goudonnet, H., Truchot, R. C., and Latruffe, N. (1993). Role of thyroid state on induction by ciprofibrate of laurate hydroxylase and peroxisomal enzymes in rat liver microsomes. *Biochem Pharmacol* 45, 1437-46.
- Paget, G. E. (1963). Experimental studies of the toxicity of atomid with particular reference to fine structural changes in the livers of rodents. *J. Atheroscl. Res.* 3, 729-736.
- Palmer, C. N., Hsu, M. H., Griffin, H. J., and Johnson, E. F. (1995). Novel sequence determinants in peroxisome proliferator signaling. *J Biol Chem* 270, 16114-21.
- Palmer, C. N., Hsu, M. H., Griffin, K. J., Raucy, J. L., and Johnson, E. F. (1998). Peroxisome proliferator activated receptor-alpha expression in human liver. *Mol Pharmacol* 53, 14-22.

- Pan, D. A., Hulbert, A. J., and Storlien, L. H. (1994). Dietary fats, membrane phospholipids and obesity. *J Nutr* 124, 1555-65.
- Parker, M. G. (1993). Steroid and related receptors. *Curr Opin Cell Biol* 5, 499-504.
- Pearce, D., and Yamamoto, K. R. (1993). Mineralocorticoid and glucocorticoid receptor activities distinguished by nonreceptor factors at a composite response element. *Science* 259, 1161-5.
- Perlmann, T., Umesono, K., Rangarajan, P. N., Forman, B. M., and Evans, R. M. (1996). Two distinct dimerization interfaces differentially modulate target gene specificity of nuclear hormone receptors. *Mol Endocrinol* 10, 958-66.
- Peters, J. M., Cattley, R. C., and Gonzalez, F. J. (1997a). Role of PPAR alpha in the mechanism of action of the nongenotoxic carcinogen and peroxisome proliferator Wy-14,643. *Carcinogenesis* 18, 2029-33.
- Peters, J. M., Hennuyer, N., Staels, B., Fruchart, J. C., Fievet, C., Gonzalez, F. J., and Auwerx, J. (1997b). Alterations in lipoprotein metabolism in peroxisome proliferator-activated receptor alpha-deficient mice. *J Biol Chem* 272, 27307-12.
- Phillipson, B. E., Rothrock, D. W., Connor, W. E., Harris, W. S., and Illingworth, D. R. (1985). Reduction of plasma lipids, lipoproteins, and apoproteins by dietary fish oils in patients with hypertriglyceridemia. *N Engl J Med* 312, 1210-6.
- Plant, N. J., Horley, N. J., Savory, R. L., Elcombe, C. R., Gray, T. J., and Bell, D. R. (1998). The peroxisome proliferators are hepatocyte mitogens in chemically- defined media: glucocorticoid-induced PPAR alpha is linked to peroxisome proliferator mitogenesis. *Carcinogenesis* 19, 925-31.
- Poirier, H., Braissant, O., Niot, I., Wahli, W., and Besnard, P. (1997). 9-cis-retinoic acid enhances fatty acid-induced expression of the liver fatty acid-binding protein gene. *FEBS Lett* 412, 480-4.
- Puigserver, P., Wu, Z., Park, C. W., Graves, R., Wright, M., and Spiegelman, B. M. (1998). A cold-inducible coactivator of nuclear receptors linked to adaptive thermogenesis. *Cell* 92, 829-39.
- Rafii-El-Idrissi, M., Calvo, J. R., Harmouch, A., Garcia-Maurino, S., and Guerrero, J. M. (1998). Specific binding of melatonin by purified cell nuclei from spleen and thymus of the rat. *J Neuroimmunol* 86, 190-7.

- Raney, A. K., Johnson, J. L., Palmer, C. N., and McLachlan, A. (1997). Members of the nuclear receptor superfamily regulate transcription from the hepatitis B virus nucleocapsid promoter. *J Virol* 71, 1058-71.
- Rao, M. S., and Reddy, J. K. (1991). An overview of peroxisome proliferator-induced hepatocarcinogenesis. *Environ Health Perspect* 93, 205-9.
- Rastinejad, F., Perlmann, T., Evans, R. M., and Sigler, P. B. (1995). Structural determinants of nuclear receptor assembly on DNA direct repeats. *Nature* 375, 203-11.
- Reddy, J. K., Azarnoff, D. L., and Hignite, C. E. (1980). Hypolipidaemic hepatic peroxisome proliferators form a novel class of chemical carcinogens. *Nature* 283, 397-8.
- Reddy, J. K., and Lalwai, N. D. (1983). Carcinogenesis by hepatic peroxisome proliferators: evaluation of the risk of hypolipidemic drugs and industrial plasticizers to humans. *Crit Rev Toxicol* 12, 1-58.
- Reddy, J. K., Goel, S. K., Nemali, M. R., Carrino, J. J., Laffler, T. G., Reddy, M. K., Sperbeck, S. J., Osumi, T., Hashimoto, T., Lalwani, N. D., and et al. (1986a). Transcription regulation of peroxisomal fatty acyl-CoA oxidase and enoyl-CoA hydratase/3-hydroxyacyl-CoA dehydrogenase in rat liver by peroxisome proliferators. *Proc Natl Acad Sci U S A* 83, 1747-51.
- Reddy, J. K., Reddy, M. K., Usman, M. I., Lalwani, N. D., and Rao, M. S. (1986b). Comparison of hepatic peroxisome proliferative effect and its implication for hepatocarcinogenicity of phthalate esters, di(2-ethylhexyl) phthalate, and di(2-ethylhexyl) adipate with a hypolipidemic drug. *Environ Health Perspect* 65, 317-27.
- Reddy, J. K., and Rao, M. S. (1989). Oxidative DNA damage caused by persistent peroxisome proliferation: its role in hepatocarcinogenesis. *Mutat Res* 214, 63-8.
- Reddy, J. K. (1990). Carcinogenicity of peroxisome proliferators: evaluation and mechanisms. *Biochem Soc Trans* 18, 92-4.
- Reddy, J. K., and Chu, R. (1996). Peroxisome proliferator-induced pleiotropic responses: pursuit of a phenomenon. *Ann N Y Acad Sci* 804, 176-201.
- Renaud, J. P., Rochel, N., Ruff, M., Vivat, V., Chambon, P., Gronemeyer, H., and Moras, D. (1995). Crystal structure of the RAR- $\gamma$  ligand-binding domain bound to all- trans retinoic acid. *Nature* 378, 681-9.
- Renkawitz, R. (1993). Repression mechanisms of v-ERBA and other members of the steroid receptor superfamily. *Ann N Y Acad Sci* 684, 1-10.



Ribeiro, R. C., Apriletti, J. W., West, B. L., Wagner, R. L., Fletterick, R. J., Schaufele, F., and Baxter, J. D. (1995a). The molecular biology of thyroid hormone action. *Ann N Y Acad Sci* 758, 366-89.

Ribeiro, R. C., Kushner, P. J., and Baxter, J. D. (1995b). The nuclear hormone receptor gene superfamily. *Annu Rev Med* 46, 443-53.

Ribeiro, R. C., Apriletti, J. W., Wagner, R. L., West, B. L., Feng, W., Huber, R., Kushner, P. J., Nilsson, S., Scanlan, T., Fletterick, R. J., Schaufele, F., and Baxter, J. D. (1998). Mechanisms of thyroid hormone action: insights from X-ray crystallographic and functional studies. *Recent Prog Horm Res* 53, 351-92.

Ricote, M., Li, A. C., Willson, T. M., Kelly, C. J., and Glass, C. K. (1998). The peroxisome proliferator-activated receptor- $\gamma$  is a negative regulator of macrophage activation. *Nature* 391, 79-82.

Ristow, M., Muller-Wieland, D., Pfeiffer, A., Krone, W., and Kahn, C. R. (1998). Obesity associated with a mutation in a genetic regulator of adipocyte differentiation. *N Engl J Med* 339, 953-9.

Rodriguez, J. C., Gil-Gomez, G., Hegardt, F. G., and Haro, D. (1994). Peroxisome proliferator-activated receptor mediates induction of the mitochondrial 3-hydroxy-3-methylglutaryl-CoA synthase gene by fatty acids. *J Biol Chem* 269, 18767-72.

Rodriguez, J. C., Ortiz, J. A., Hegardt, F. G., and Haro, D. (1998). The hepatocyte nuclear factor 4 (HNF-4) represses the mitochondrial HMG-CoA synthase gene. *Biochem Biophys Res Commun* 242, 692-6.

Rojiani, M. V., Finlay, B. B., Gray, V., and Dedhar, S. (1991). In vitro interaction of a polypeptide homologous to human Ro/SS-A antigen (calreticulin) with a highly conserved amino acid sequence in the cytoplasmic domain of integrin alpha subunits. *Biochemistry* 30, 9859-66.

Ruyter, B., Andersen, O., Dehli, A., Ostlund Farrants, A. K., Gjoen, T., and Thomassen, M. S. (1997). Peroxisome proliferator activated receptors in Atlantic salmon (*Salmo salar*): effects on PPAR transcription and acyl-CoA oxidase activity in hepatocytes by peroxisome proliferators and fatty acids. *Biochim Biophys Acta* 1348, 331-8.

Saatcioglu, F., Deng, T., and Karin, M. (1993). A novel cis element mediating ligand-independent activation by c-ErbA: implications for hormonal regulation. *Cell* 75, 1095-105.

- Sande, S., and Privalsky, M. L. (1996). Identification of TRACs (T3 receptor-associating cofactors), a family of cofactors that associate with, and modulate the activity of, nuclear hormone receptors. *Mol Endocrinol* 10, 813-25.
- Sap, J., Munoz, A., Schmitt, J., Stunnenberg, H., and Vennstrom, B. (1989). Repression of transcription mediated at a thyroid hormone response element by the v-erb-A oncogene product. *Nature* 340, 242-4.
- Schmidt, A., Endo, N., Rutledge, S. J., Vogel, R., Shinar, D., and Rodan, G. A. (1992). Identification of a new member of the steroid hormone receptor superfamily that is activated by a peroxisome proliferator and fatty acids. *Mol Endocrinol* 6, 1634-41.
- Schmidt, S., Baniahmad, A., Eggert, M., Schneider, S., and Renkawitz (1998). Multiple receptor interaction domains of GRIP1 function in synergy. *Nucleic Acids Res* 26, 1191-7.
- Schoonjans, K., Watanabe, M., Suzuki, H., Mahfoudi, A., Krey, G., Wahli, W., Grimaldi, P., Staels, B., Yamamoto, T., and Auwerx, J. (1995). Induction of the acyl-coenzyme A synthetase gene by fibrates and fatty acids is mediated by a peroxisome proliferator response element in the C promoter. *J Biol Chem* 270, 19269-76.
- Schoonjans, K., Staels, B., and Auwerx, J. (1996). The peroxisome proliferator activated receptors (PPARs) and their effects on lipid metabolism and adipocyte differentiation. *Biochim Biophys Acta* 1302, 93-109.
- Schoonjans, K., Martin, G., Staels, B., and Auwerx, J. (1997). Peroxisome proliferator-activated receptors, orphans with ligands and functions. *Curr Opin Lipidol* 8, 159-66.
- Schrader, M., Danielsson, C., Wiesenberg, I., and Carlberg, C. (1996). Identification of natural monomeric response elements of the nuclear receptor RZR/ROR. They also bind COUP-TF homodimers. *J Biol Chem* 271, 19732-6.
- Schulte-Hermann, R., Bursch, W., and Parzefall, W. (1991). Mitogenesis and programmed cell death as determinants of carcinogenicity of nongenotoxic compounds. *Prog Clin Biol Res* 369, 237-44.
- Shago, M., Flock, G., Leung Hagesteijn, C. Y., Woodside, M., Grinstein, S., Giguere, V., and Dedhar, S. (1997). Modulation of the retinoic acid and retinoid X receptor signaling pathways in P19 embryonal carcinoma cells by calreticulin. *Exp Cell Res* 230, 50-60.
- Shaw, I. C., and Jones, H. B. (1994). Mechanisms of non-genotoxic carcinogenesis. *Trends Pharmacol Sci* 15, 89-93.

- Shepard, A. R., and Eberhardt, N. L. (1993). Molecular mechanisms of thyroid hormone action. *Clin Lab Med* 13, 531-41.
- Sher, T., Yi, H. F., McBride, O. W., and Gonzalez, F. J. (1993). cDNA cloning, chromosomal mapping, and functional characterization of the human peroxisome proliferator activated receptor. *Biochemistry* 32, 5598-604.
- Shibata, H., Nawaz, Z., Tsai, S. Y., O'Malley, B. W., and Tsai, M. J. (1997). Gene silencing by chicken ovalbumin upstream promoter-transcription factor I (COUP-TFI) is mediated by transcriptional corepressors, nuclear receptor-corepressor (N-CoR) and silencing mediator for retinoic acid receptor and thyroid hormone receptor (SMRT). *Mol Endocrinol* 11, 714-24.
- Silva, J. E., and Rabelo, R. (1997). Regulation of the uncoupling protein gene expression. *Eur J Endocrinol* 136, 251-64.
- Sladek, F. M., Zhong, W. M., Lai, E., and Darnell, J. E., Jr. (1990). Liver-enriched transcription factor HNF-4 is a novel member of the steroid hormone receptor superfamily. *Genes Dev* 4, 2353-65.
- Smith, C. L., Nawaz, Z., and O'Malley, B. W. (1997). Coactivator and corepressor regulation of the agonist/antagonist activity of the mixed antiestrogen, 4-hydroxytamoxifen. *Mol Endocrinol* 11, 657-66.
- Spencer, T. E., Jenster, G., Burcin, M. M., Allis, C. D., Zhou, J., Mizzen, C. A., McKenna, N. J., Onate, S. A., Tsai, S. Y., Tsai, M. J., and O'Malley, B. W. (1997). Steroid receptor coactivator-1 is a histone acetyltransferase. *Nature* 389, 194-8.
- Spiegelman, B. M., and Flier, J. S. (1996). Adipogenesis and obesity: rounding out the big picture. *Cell* 87, 377-89.
- St-Arnaud, R., Prud'homme, J., Leung-Hagesteijn, C., and Dedhar, S. (1995). Constitutive expression of calreticulin in osteoblasts inhibits mineralization. *J Cell Biol* 131, 1351-9.
- Steinhilber, D., Brungs, M., Werz, O., Wiesenberg, I., Danielsson, C., Kahlen, J. P., Nayeri, S., Schrader, M., and Carlberg, C. (1995). The nuclear receptor for melatonin represses 5-lipoxygenase gene expression in human B lymphocytes. *J Biol Chem* 270, 7037-40.
- Stoffel, M., and Duncan, S. A. (1997). The maturity-onset diabetes of the young (MODY1) transcription factor HNF4 $\alpha$  regulates expression of genes required for glucose transport and metabolism. *Proc Natl Acad Sci U S A* 94, 13209-14.

Storlien, L. H., Jenkins, A. B., Chisholm, D. J., Pascoe, W. S., Khouri, S., and Kraegen, E. W. (1991). Influence of dietary fat composition on development of insulin resistance in rats. Relationship to muscle triglyceride and omega-3 fatty acids in muscle phospholipid. *Diabetes* 40, 280-9.

Stunnenberg, H. G. (1993). Mechanisms of transactivation by retinoic acid receptors. *Bioessays* 15, 309-15.

Svoboda, D. J., and Azarnoff, D. L. (1966). Response of hepatic microbodies to a hypolipidemic agent, ethyl chlorophenoxyisobutyrate (CPIB). *J Cell Biol* 30, 442-50.

Takeda, T., Ichikawa, K., Miyamoto, T., Kobayashi, M., Nishii, Y., Suzuki, S., Sakurai, A., and Hashizume, K. (1992). Regulation of rat hepatic peroxisomal enoyl-CoA hydratase-3-hydroxyacyl-CoA dehydrogenase bifunctional enzyme by thyroid hormone. *Biochem Biophys Res Commun* 185, 211-6.

Takeshita, A., Cardona, G. R., Koibuchi, N., Suen, C. S., and Chin, W. W. (1997). TRAM-1, A novel 160-kDa thyroid hormone receptor activator molecule, exhibits distinct properties from steroid receptor coactivator-1. *J Biol Chem* 272, 27629-34.

Takeshita, A., Yen, P. M., Ikeda, M., Cardona, G. R., Liu, Y., Koibuchi, N., Norwitz, E. R., and Chin, W. W. (1998). Thyroid hormone response elements differentially modulate the interactions of thyroid hormone receptors with two receptor binding domains in the steroid receptor coactivator-1. *J Biol Chem* 273, 21554-62.

Tata, J. R. (1994). Hormonal regulation of programmed cell death during amphibian metamorphosis. *Biochem Cell Biol* 72, 581-8.

Tenbaum, S., and Baniahmad, A. (1997). Nuclear receptors: structure, function and involvement in disease. *Int J Biochem Cell Biol* 29, 1325-41.

Thangada, S., Alvares, K., Mangino, M., Usman, M. I., Rao, M. S., and Reddy, J. K. (1989). An in vitro demonstration of peroxisome proliferation and increase in peroxisomal  $\beta$ -oxidation system mRNAs in cultured rat hepatocytes treated with ciprofibrate. *FEBS Lett* 250, 205-10.

Thompson, P. D., Jurutka, P. W., Haussler, C. A., Whitfield, G. K., and Haussler, M. R. (1998). Heterodimeric DNA binding by the vitamin D receptor and retinoid X receptors is enhanced by 1,25-dihydroxyvitamin D<sub>3</sub> and inhibited by 9- cis-retinoic acid. Evidence for allosteric receptor interactions. *J Biol Chem* 273, 8483-91.

- Tini, M., Fraser, R. A., and Giguere, V. (1995). Functional interactions between retinoic acid receptor-related orphan nuclear receptor (ROR alpha) and the retinoic acid receptors in the regulation of the gamma F-crystallin promoter. *J Biol Chem* 270, 20156-61.
- Tolbert, N. E. (1981). Metabolic pathways in peroxisomes and glyoxysomes. *Annu Rev Biochem* 50, 133-57.
- Tontonoz, P., Graves, R. A., Budavari, A. I., Erdjument-Bromage, H., Lui, M., Hu, E., Tempst, P., and Spiegelman, B. M. (1994a). Adipocyte-specific transcription factor ARF6 is a heterodimeric complex of two nuclear hormone receptors, PPAR gamma and RXR alpha. *Nucleic Acids Res* 22, 5628-34.
- Tontonoz, P., Hu, E., Graves, R. A., Budavari, A. I., and Spiegelman, B. M. (1994b). mPPAR gamma 2: tissue-specific regulator of an adipocyte enhancer. *Genes Dev* 8, 1224-34.
- Tontonoz, P., Hu, E., and Spiegelman, B. M. (1994c). Stimulation of adipogenesis in fibroblasts by PPAR gamma 2, a lipid-activated transcription factor (published erratum appears in *Cell* 1995 Mar 24;80(6):following 957). *Cell* 79, 1147-56.
- Tontonoz, P., Hu, E., Devine, J., Beale, E. G., and Spiegelman, B. M. (1995). PPAR gamma 2 regulates adipose expression of the phosphoenolpyruvate carboxykinase gene. *Mol Cell Biol* 15, 351-7.
- Tontonoz, P., Nagy, L., Alvarez, J. G., Thomazy, V. A., and Evans, R. M. (1998). PPAR $\gamma$  promotes monocyte/macrophage differentiation and uptake of oxidized LDL. *Cell* 93, 241-52.
- Torchia, J., Rose, D. W., Inostroza, J., Kamei, Y., Westin, S., Glass, C. K., and Rosenfeld, M. G. (1997). The transcriptional co-activator p/CIP binds CBP and mediates nuclear-receptor function. *Nature* 387, 677-84.
- Torchia, J., Glass, C., and Rosenfeld, M. G. (1998). Co-activators and co-repressors in the integration of transcriptional responses. *Curr Opin Cell Biol* 10, 373-83.
- Treuter, E., Albrektsen, T., Johansson, L., Leers, J., and Gustafsson, J. A. (1998). A regulatory role for RIP140 in nuclear receptor activation. *Mol Endocrinol* 12, 864-81.
- Truss, M., and Beato, M. (1993). Steroid hormone receptors: interaction with deoxyribonucleic acid and transcription factors. *Endocr Rev* 14, 459-79.
- Tsai, M. J., and O'Malley, B. W. (1994). Molecular mechanisms of action of steroid/thyroid receptor superfamily members. *Annu Rev Biochem* 63, 451-86.

- Tugwood, J. D., Issemann, I., Anderson, R. G., Bundell, K. R., McPheat, W. L., and Green, S. (1992). The mouse peroxisome proliferator activated receptor recognizes a response element in the 5' flanking sequence of the rat acyl CoA oxidase gene. *Embo J* 11, 433-9.
- Tugwood, J. D., Holden, P. R., James, N. H., Prince, R. A., and Roberts, R. A. (1998). A peroxisome proliferator-activated receptor-alpha (PPAR $\alpha$ ) cDNA cloned from guinea-pig liver encodes a protein with similar properties to the mouse PPAR $\alpha$ : implications for species differences in responses to peroxisome proliferators. *Arch Toxicol* 72, 169-77.
- Ueda, A., Takeshita, F., Yamashiro, S., and Yoshimura, T. (1998). Positive regulation of the human macrophage stimulating protein gene transcription. Identification of a new hepatocyte nuclear factor-4 (HNF-4) binding element and evidence that indicates direct association between NF-Y and HNF-4. *J Biol Chem* 273, 19339-47.
- Umesono, K., and Evans, R. M. (1989). Determinants of target gene specificity for steroid/thyroid hormone receptors. *Cell* 57, 1139-46.
- Umesono, K., Murakami, K. K., Thompson, C. C., and Evans, R. M. (1991). Direct repeats as selective response elements for the thyroid hormone, retinoic acid, and vitamin D3 receptors. *Cell* 65, 1255-66.
- Vassilakos, A., Michalak, M., Lehrman, M. A., and Williams, D. B. (1998). Oligosaccharide binding characteristics of the molecular chaperones calnexin and calreticulin. *Biochemistry* 37, 3480-90.
- Voegel, J. J., Heine, M. J., Tini, M., Vivat, V., Chambon, P., and Gronemeyer, H. (1998). The coactivator TIF2 contains three nuclear receptor-binding motifs and mediates transactivation through CBP binding-dependent and -independent pathways. *Embo J* 17, 507-19.
- Vu-Dac, N., Schoonjans, K., Laine, B., Fruchart, J. C., Auwerx, J., and Staels, B. (1994). Negative regulation of the human apolipoprotein A-I promoter by fibrates can be attenuated by the interaction of the peroxisome proliferator-activated receptor with its response element. *J Biol Chem* 269, 31012-8.
- Vu-Dac, N., Gervois, P., Grotzinger, T., De Vos, P., Schoonjans, K., Fruchart, J. C., Auwerx, J., Mariani, J., Tedgui, A., and Staels, B. (1997). Transcriptional regulation of apolipoprotein A-I gene expression by the nuclear receptor ROR $\alpha$ . *J Biol Chem* 272, 22401-4.

- Vu-Dac, N., Chopin-Delannoy, S., Gervois, P., Bonnelye, E., Martin, G., Fruchart, J. C., Laudet, V., and Staels, B. (1998). The nuclear receptors peroxisome proliferator-activated receptor alpha and rev-erbalpha mediate the species-specific regulation of apolipoprotein A-I expression by fibrates. *J Biol Chem* 273, 25713-20.
- Wade, P. A., Pruss, D., and Wolffe, A. P. (1997). Histone acetylation: chromatin in action. *Trends Biochem Sci* 22, 128-32.
- Wagner, R. L., Apriletti, J. W., McGrath, M. E., West, B. L., Baxter, J. D., and Fletterick, R. J. (1995). A structural role for hormone in the thyroid hormone receptor. *Nature* 378, 690-7.
- Waxman, D. J. (1996). Role of metabolism in the activation of dehydroepiandrosterone as a peroxisome proliferator. *J Endocrinol* 150 *Suppl*, S129-47.
- Webb, S. J., Xiao, G. H., Geoghegan, T. E., and Prough, R. A. (1996). Regulation of CYP4A expression in rat by dehydroepiandrosterone and thyroid hormone. *Mol Pharmacol* 49, 276-87.
- Welsch, C. W. (1992). Relationship between dietary fat and experimental mammary tumorigenesis: a review and critique. *Cancer Res* 52, 2040s-2048s.
- Westin, S., Kurokawa, R., Nolte, R. T., Wisely, G. B., McInerney, E. M., Rose, D. W., Milburn, M. V., Rosenfeld, M. G., and Glass, C. K. (1998). Interactions controlling the assembly of nuclear-receptor heterodimers and co-activators. *Nature* 395, 199-202.
- Wheeler, D. G., Horsford, J., Michalak, M., White, J. H., and Hendy, G. N. (1995). Calreticulin inhibits vitamin D3 signal transduction. *Nucleic Acids Res* 23, 3268-74.
- White, T. K., Zhu, Q., and Tanzer, M. L. (1995). Cell surface calreticulin is a putative mannoside lectin which triggers mouse melanoma cell spreading. *J Biol Chem* 270, 15926-9.
- Wiesenberg, I., Chiesi, M., Missbach, M., Spanka, C., Pignat, W., and Carlberg, C. (1998). Specific activation of the nuclear receptors PPARgamma and RORA by the antidiabetic thiazolidinedione BRL 49653 and the antiarthritic thiazolidinedione derivative CGP 52608. *Mol Pharmacol* 53, 1131-8.
- Willson, T. M., and Wahli, W. (1997). Peroxisome proliferator-activated receptor agonists. *Curr Opin Chem Biol* 1, 235-41.

- Wilson, T. E., Paulsen, R. E., Padgett, K. A., and Milbrandt, J. (1992). Participation of non-zinc finger residues in DNA binding by two nuclear orphan receptors. *Science* 256, 107-110.
- Winrow, C. J., Marcus, S. L., Miyata, K. S., Zhang, B., Capone, J. P., and Rachubinski, R. A. (1994). Transactivation of the peroxisome proliferator-activated receptor is differentially modulated by hepatocyte nuclear factor-4. *Gene Expr* 4, 53-62.
- Winrow, C. J., Miyata, K. S., Marcus, S. L., Burns, K., Michalak, M., Capone, J. P., and Rachubinski, R. A. (1995). Calreticulin modulates the in vitro DNA binding but not the in vivo transcriptional activation by peroxisome proliferator-activated receptor/retinoid X receptor heterodimers. *Mol Cell Endocrinol* 111, 175-9.
- Winrow, C. J., Kassam, A., Miyata, K. S., Marcus, S. L., Hunter, J., Capone, J. P., and Rachubinski, R. A. (1996). Interplay of the peroxisome proliferator-activated receptor and the thyroid hormone receptor-signaling pathways in regulating peroxisome proliferator-responsive genes. *Ann N Y Acad Sci* 804, 214-30.
- Winrow, C. J., Capone, J. P., and Rachubinski, R. A. (1998). Cross-talk between Orphan Nuclear Hormone Receptor RZR $\alpha$  and Peroxisome Proliferator-activated Receptor  $\alpha$  in Regulation of the Peroxisomal Hydratase-Dehydrogenase Gene. *J Biol Chem* 273, 31442-31448.
- Wolffe, A. P. (1997). Chromatin remodeling regulated by steroid and nuclear receptors. *Cell Res* 7, 127-42.
- Wolffe, A. P. (1998). Packaging principle: how DNA methylation and histone acetylation control the transcriptional activity of chromatin. *J Exp Zool* 282, 239-44.
- Xiong, S., Chirala, S. S., Hsu, M. H., and Wakil, S. J. (1998). Identification of thyroid hormone response elements in the human fatty acid synthase promoter. *Proc Natl Acad Sci U S A* 95, 12260-5.
- Yamada, J., Sugiyama, H., Tamura, H., and Suga, T. (1994). Hormonal modulation of peroxisomal enzyme induction caused by peroxisome proliferators: suppression by growth and thyroid hormones in cultured rat hepatocytes. *Arch Biochem Biophys* 315, 555-7.
- Yamagata, K., Furuta, H., Oda, N., Kaisaki, P. J., Menzel, S., Cox, N. J., Fajans, S. S., Signorini, S., Stoffel, M., and Bell, G. I. (1996). Mutations in the hepatocyte nuclear factor-4 $\alpha$  gene in maturity-onset diabetes of the young (MODY1). *Nature* 384, 458-60.



- Yamaguchi, M., and Oishi, K. (1995). 17 beta-Estradiol stimulates the expression of hepatic calcium-binding protein regucalcin mRNA in rats. *Mol Cell Biochem* 143, 137-41.
- Yan, Z. H., Karam, W. G., Staudinger, J. L., Medvedev, A., Ghanayem, B. I., and Jetten, A. M. (1998). Regulation of peroxisome proliferator-activated receptor  $\alpha$ -induced transactivation by the nuclear orphan receptor TAK1/TR4. *J Biol Chem* 273, 10948-57.
- Yang, X. J., Ogryzko, V. V., Nishikawa, J., Howard, B. H., and Nakatani, Y. (1996). A p300/CBP-associated factor that competes with the adenoviral oncoprotein E1A. *Nature* 382, 319-24.
- Yao, T. P., Ku, G., Zhou, N., Scully, R., and Livingston, D. M. (1996). The nuclear hormone receptor coactivator SRC-1 is a specific target of p300. *Proc Natl Acad Sci U S A* 93, 10626-31.
- Yokomizo, T., Izumi, T., Chang, K., Takuwa, Y., and Shimizu, T. (1997). A G-protein-coupled receptor for leukotriene B4 that mediates chemotaxis. *Nature* 387, 620-4.
- Yokomori, N., Nishio, K., Aida, K., and Negishi, M. (1997). Transcriptional regulation by HNF-4 of the steroid 15 $\alpha$ -hydroxylase P450 (Cyp2a-4) gene in mouse liver. *J Steroid Biochem Mol Biol* 62, 307-14.
- Youssef, J., and Badr, M. (1998). Extraperoxisomal targets of peroxisome proliferators: mitochondrial, microsomal, and cytosolic effects. Implications for health and disease. *Crit Rev Toxicol* 28, 1-33.
- Yu, X., and Mertz, J. E. (1997). Differential regulation of the pre-C and pregenomic promoters of human hepatitis B virus by members of the nuclear receptor superfamily. *J Virol* 71, 9366-74.
- Yuan, C. X., Ito, M., Fondell, J. D., Fu, Z. Y., and Roeder, R. G. (1998). The TRAP220 component of a thyroid hormone receptor-associated protein (TRAP) coactivator complex interacts directly with nuclear receptors in a ligand-dependent fashion. *Proc Natl Acad Sci U S A* 95, 7939-44.
- Zamir, I., Harding, H. P., Atkins, G. B., Horlein, A., Glass, C. K., Rosenfeld, M. G., and Lazar, M. A. (1996). A nuclear hormone receptor corepressor mediates transcriptional silencing by receptors with distinct repression domains. *Mol Cell Biol* 16, 5458-65.
- Zamir, I., Zhang, J., and Lazar, M. A. (1997). Stoichiometric and steric principles governing repression by nuclear hormone receptors. *Genes Dev* 11, 835-46.

Zhang, B., Marcus, S. L., Sajjadi, F. G., Alvares, K., Reddy, J. K., Subramani, S., Rachubinski, R. A., and Capone, J. P. (1992). Identification of a peroxisome proliferator-responsive element upstream of the gene encoding rat peroxisomal enoyl-CoA hydratase/3-hydroxyacyl-CoA dehydrogenase. *Proc Natl Acad Sci U S A* 89, 7541-5.

Zhang, B., Marcus, S. L., Miyata, K. S., Subramani, S., Capone, J. P., and Rachubinski, R. A. (1993). Characterization of protein-DNA interactions within the peroxisome proliferator-responsive element of the rat hydratase-dehydrogenase gene. *J Biol Chem* 268, 12939-45.

Zhang, X. K., Hoffmann, B., Tran, P. B., Graupner, G., and Pfahl, M. (1992). Retinoid X receptor is an auxiliary protein for thyroid hormone and retinoic acid receptors. *Nature* 355, 441-6.

Zhang, Y., Iratni, R., Erdjument-Bromage, H., Tempst, P., and Reinberg, D. (1997). Histone deacetylases and SAP18, a novel polypeptide, are components of a human Sin3 complex. *Cell* 89, 357-64.

Zhou, Y. C., and Waxman, D. J. (1998). Activation of peroxisome proliferator-activated receptors by chlorinated hydrocarbons and endogenous steroids. *Environ Health Perspect* 106 Suppl 4, 983-8.

Zhu, N., Pewitt, E. B., Cai, X., Cohn, E. B., Lang, S., Chen, R., and Wang, Z. (1998). Calreticulin: an intracellular Ca<sup>++</sup>-binding protein abundantly expressed and regulated by androgen in prostatic epithelial cells. *Endocrinology* 139, 4337-44.

Zhu, Y., Alvares, K., Huang, Q., Rao, M. S., and Reddy, J. K. (1993). Cloning of a new member of the peroxisome proliferator-activated receptor gene family from mouse liver. *J Biol Chem* 268, 26817-20.

Zhu, Y., Qi, C., Calandra, C., Rao, M. S., and Reddy, J. K. (1996). Cloning and identification of mouse steroid receptor coactivator-1 (mSRC-1), as a coactivator of peroxisome proliferator-activated receptor gamma. *Gene Expr* 6, 185-95.

Zhu, Y., Qi, C., Jain, S., Rao, M. S., and Reddy, J. K. (1997). Isolation and characterization of PBP, a protein that interacts with peroxisome proliferator-activated receptor. *J Biol Chem* 272, 25500-6.