

CANADIAN THESES ON MICROFICHE

I.S.B.N.

THESES CANADIENNES SUR MICROFICHE



National Library of Canada
Collections Development Branch

Canadian Theses on
Microfiche Service

Ottawa, Canada
K1A 0N1

Bibliothèque nationale du Canada
Direction du développement des collections

Service des thèses canadiennes
sur microfiche

NOTICE

The quality of this microfiche is heavily dependent upon the quality of the original thesis submitted for microfilming. Every effort has been made to ensure the highest quality of reproduction possible.

If pages are missing, contact the university which granted the degree.

Some pages may have indistinct print especially if the original pages were typed with a poor typewriter ribbon or if the university sent us a poor photocopy.

Previously copyrighted material (journal articles and book reviews) is not microfilmed.

Reproduction in full or in part of this film is governed by the Canadian Copyright Act, R.S.C. 1970, c. C-30. Please read the authorization forms which accompany this thesis.

THIS DISSERTATION
HAS BEEN MICROFILMED
EXACTLY AS RECEIVED

AVIS

La qualité de cette microfiche dépend grandement de la qualité de la thèse soumise au microfilmage. Nous avons tout fait pour assurer une qualité supérieure de reproduction.

S'il manque des pages, veuillez communiquer avec l'université qui a conféré le grade.

La qualité d'impression de certaines pages peut laisser à désirer, surtout si les pages originales ont été dactylographiées à l'aide d'un ruban usé ou si l'université nous a fait parvenir une photocopie de mauvaise qualité.

Les documents qui font déjà l'objet d'un droit d'auteur (articles de revue, examens publiés, etc.) ne sont pas microfilmés.

La reproduction, même partielle, de ce microfilm est soumise à la Loi canadienne sur le droit d'auteur, SRC 1970, c. C-30. Veuillez prendre connaissance des formulaires d'autorisation qui accompagnent cette thèse.

LA THÈSE A ÉTÉ
MICROFILMÉE TELLE QUE
NOUS L'AVONS REÇUE

Canada



National Library
of Canada

Bibliothèque nationale
du Canada

Canadian Theses Division

Division des thèses canadiennes

Ottawa, Canada
K1A 0N4

67278

PERMISSION TO MICROFILM — AUTORISATION DE MICROFILMER

• Please print or type — Écrire en lettres moulées ou dactylographier

Full Name of Author — Nom complet de l'auteur

CHOI, DONG-KOO

Date of Birth — Date de naissance

Feb. 13, 1946

Country of Birth — Lieu de naissance

KOREA

Permanent Address — Résidence fixe

#4, 3720-105 st

Edmonton, Alberta

CANADA

Title of Thesis — Titre de la thèse

Electronic ...

University — Université

University of Alberta

Degree for which thesis was presented — Grade pour lequel

Ph.D.

Year of degree — Année de la thèse

1980

Supervisor — Directeur de thèse

...

Permission is hereby granted to the NATIONAL LIBRARY OF CANADA to microfilm this thesis and to lend or sell copies of the film.

The author reserves other publication rights, and neither the thesis nor extensive extracts therefrom may be printed or otherwise reproduced without the author's written permission.

L'autorisation est par la présente accordée à la BIBLIOTHÈQUE NATIONALE DU CANADA de microfilmer cette thèse et de prêter ou de vendre des exemplaires du film.

L'auteur se réserve les autres droits de publication; ni la thèse ni de longs extraits de celle-ci ne doivent être imprimés autrement que par la permission écrite de l'auteur.

Signature

...

THE UNIVERSITY OF ALBERTA

ALGEBRAIC COMPUTATIONS OF SCALED PADÉ FRACTIONS

by

© Dong-Koo Choi

A thesis

submitted to the Faculty of Graduate Studies and Research
in partial fulfillment of the requirements for the degree
of Doctor of Philosophy

Department of Computing Science

Edmonton, Alberta

Spring 1994

RELEASE FORM

Year this degree was granted: 1984

The author reserves other publication rights, and neither the thesis nor extensive extracts from it may be printed or otherwise reproduced without the author's written permission.


(Signed)

#4 3720-105 Street
Edmonton, Alberta
Canada


THE UNIVERSITY OF ALBERTA

FACULTY OF GRADUATE STUDIES AND RESEARCH


The undersigned certify that they have read, and recommend to the Faculty of Graduate Studies and Research, for acceptance, a thesis entitled "Algebraic Computations of Scaled Pade Fractions", submitted by Dong-Koo Choi in partial fulfillment of the requirements for the degree of Doctor of Philosophy.



Prof. S. Cabay
Supervisor




Prof. C. Addison



Prof. W. Armstrong



Prof. J. Tartar



Prof. V. Gourishankar



Prof. K. Geddes
External Examiner

Date

Dec 20/83

ABSTRACT

Two companion algorithms are developed for constructing Pade fractions along an off-diagonal path of the Pade table for a function $-A(z)/B(z)$, where $A(z)$ and $B(z)$ are formal power series over a field.

One of the algorithms computes the first n Pade fractions along the off-diagonal in time $O(n^2)$. When $A(z)$ and $B(z)$ are finite power series (i.e., polynomials), it is shown that the algorithm is equivalent to Euclid's extended algorithm for computing greatest common divisors.

The other algorithm, a generalization of the first, proceeds along the off-diagonal in quadratic steps, and is of complexity $O(n \log^2 n)$. When $A(z)$ and $B(z)$ are polynomials, the second algorithm becomes a fast Euclid's extended algorithm for computing greatest common divisors. The algorithm is of the same complexity as other fast greatest common divisor methods, but its iterative nature provides a practical advantage during implementation.

The algorithms may also be used for computing Pade fractions along an anti-diagonal path of the Pade table, as well. The fast algorithm is of the same complexity as other fast algorithms for anti-diagonal computations. However, it has the advantage of being able to determine easily any specific Pade fraction along the anti-diagonal.

Finally, it is shown that two successive Pade fractions can be used to obtain the inverse of Hankel and Toeplitz matrices in time $O(n \log^2 n)$.

ACKNOWLEDGEMENT

I am deeply indebted to my thesis supervisor, Dr. S. Cabay, for introducing me to the problems, guiding me at each stage of my research, and offering valuable suggestions for and criticisms of the dissertation.

I would like to thank my committee members, Dr. C. Addison, Dr. K. Geddes, Dr. V. Gourishankar, Dr. J. Tartar, and Dr. W. Armstrong for the valuable comments on many improvements in the dissertation. I also wish to thank Mrs. S. Lim for her sincere help in typing and producing the dissertation.

Both the financial and technical supports provided by the Department of Computing Science are gratefully acknowledged.

Most of all, I thank my family, who gave me encouragement when I needed it most, and endured five years.

TABLE OF CONTENTS

	Page
CHAPTER 1 INTRODUCTION	1
CHAPTER 2 PADE THEORY	4
2.1 Introduction	4
2.2 Pade Forms	4
2.2.1 Definitions	4
2.2.2 Existence and Non-uniqueness	6
2.2.3 General Structure of Pade Forms	10
2.3 Pade Fractions	15
2.3.1 Scaled Pade Fractions	15
2.3.2 Non-scaled Pade Fractions	16
CHAPTER 3 COMPUTATION OF OFF-DIAGONAL SCALED PADE FRACTIONS	18
3.1 Introduction	18
3.2 Preliminary Results	19
3.3 Fast Off-diagonal Algorithm for a Single Power Series	29
3.3.1 The Algorithm	29
3.3.2 Proof of algorithm validity	33
3.3.3 Cost Analysis	36
3.4 Fast Off-diagonal Algorithm for a Quotient Power Series	41
3.5 Classical Off diagonal Algorithm for a Quotient Power Series	45

CHAPTER 4 GREATEST COMMON DIVISOR COMPUTATIONS OF POLYNOMIALS	50
4.1 Introduction	50
4.2 Euclid's Extended Algorithm	51
4.3 Fast GCD Computations	55
4.4 Antidiagonal Computations	56
CHAPTER 5 THE INVERSE OF HANKEL AND TOEPLITZ MATRICES	60
5.1 Introduction	60
5.2 The Inverse Formula	60
CHAPTER 6 CONCLUSIONS AND SUGGESTIONS FOR FUTURE RESEARCH	71
BIBLIOGRAPHY	74

CHAPTER 1

INTRODUCTION

The Pade table of a formal power series

$$A(z) = \sum_{i=0}^{\infty} a_i z^i \quad (1.1)$$

is a doubly infinite array of rational functions

$$\frac{U_{mn}(z)}{V_{mn}(z)} = \frac{\sum_{i=0}^m u_i z^i}{\sum_{i=0}^n v_i z^i} \quad (1.2)$$

determined in such a manner that the Maclaurin expansion of $U_{mn}(z)/V_{mn}(z)$ agrees with $A(z)$ as far as possible. The power series $A(z)$ is said to be normal if, for each pair (m, n) , this agreement is exact through the power z^{m+n} . The foundation for the development of Pade theory was laid by Cauchy(1821) in his famous "Cour d'Analyse". Later, Frobenius(1881) developed the basic algorithmic aspects of the theory, and Pade(1892) treated in detail certain abnormal cases.

Since Pade's time, Pade tables have become a classical tool of analysis. Their analytical properties have been studied in great depth and are surveyed, for example, by Gagg [GAG74] and by Baker [BAK75]. Traditionally, it is assumed that the coefficients in (1.1) and (1.2) lie in the field of complex numbers, and that the power series and the rational functions are to be evaluated at certain points in the complex plane.

Although the results obtained in this thesis are likely to have an impact in an analytical (or numerical) setting, the effects of this impact are not examined. The issues addressed are strictly algebraic ones: that is, no consideration is given to the goodness of the approximation of (1.2) to (1.1). Instead, the objective is to provide an effective tool to algebraically manipulate rational functions.

tions as truncated power series (for which the cost of operations is relatively cheap), and to transform back to rational form on request. It is assumed that the coefficients lie in an arbitrary field.

The coefficients of the rational function (1.2) satisfy an under-determined system of linear equations, known as the Hankel system. Various properties of the family of solutions to the Hankel system are described in Chapter 2. All the results given are those of Pade, and are included for the sake of completeness and for reference ease. The Hankel system can be solved directly for the coefficients of $U_m(z)$ and $V_m(z)$. With coefficients over a field, Rissanen [RIS73], for example, provides an algorithm which requires $O(n^2)$ arithmetic operations over the field. On the other hand, with coefficients over an arbitrary integral domain, Geddes [GED79] gives a fraction-free algorithm which requires $O(n^3)$ arithmetic operations over the integral domain.

Various relationships are known to exist between neighboring elements in the Pade table. These relationships have been used to derive numerous $O(n^2)$ methods for computing a sequence of elements in the Pade table. A survey and comparison of these methods are given in Brezinski [BRZ76], Claessens [CLA75] and Wynn [WYN60]. All these methods have a major flaw; they may fail in the abnormal case. In Chapter 3, a new relationship between elements lying along an off-diagonal path in the Pade table is derived. This leads to yet another $O(n^2)$ method, however, the new method succeeds in the abnormal case. Furthermore, if the coefficient field has an appropriate n -th root of unity (which permits fast multiplication and division of polynomials), the asymptotic complexity of the algorithm becomes $O(n \log^2 n)$.

The new algorithm can be applied to the quotient of two power series. Then, in particular, it can be applied to the quotient of two finite power series (i.e. the quotient of two polynomials). In Chapter 4, it is shown that if all elements along a specific off-diagonal of the Pade table are computed, then the new algorithm is equivalent to Euclid's extended algorithm for computing greatest common divisors. Furthermore, if fast polynomial operations can be performed, the new

algorithm can compute the greatest common divisor of two polynomials in $O(n \log^2 n)$ arithmetic operations. The algorithm has three advantages over the other fast methods (Moënck [MOE73], Aho et al [AHO74], and Brent et al [BRE80]) for computing greatest common divisors. It is basically an iterative algorithm rather than a recursive one, and consequently, significant cost savings can result during implementation. Secondly, it produces intermediate polynomial remainder sequences as a by-product, which is a valuable feature for some applications. Finally, various details about the nature of its behavior are easier to comprehend.

The algorithm can be applied to the quotient of the reciprocals of two truncated power series (polynomials). It is shown in Section 4.4 that this yields successive elements along an anti-diagonal path of the Pade table.

In Chapter 5, it is shown that two successive elements along the diagonal path of the Pade table can be used to find the inverse of a Hankel matrix. Therefore, if fast polynomial operations are possible, the inverse of a Hankel matrix of order n can be determined in $O(n \log^2 n)$ arithmetic operations. The new inversion method handles abnormalities with greater ease, and less cost, than does Brent et al's algorithm [BRE80]. Furthermore, by proceeding along an anti-diagonal path, the algorithm can be used to compute the inverse of a Toeplitz matrix of order n in time $O(n \log^2 n)$.

CHAPTER 2

PADE THEORY

2.1. Introduction

This chapter examines the theoretical background of Pade theory. None of the results are new; however, some of the proofs may be original. All of the proofs are obtained directly from the properties of Hankel systems.

The highlight of this chapter is Corollary 2.6, due to Pade, which completely describes the family of solutions to the Hankel system. We define one specific member of this family to be a scaled Pade fraction. Scaled Pade fractions exist uniquely, and are fundamental to the development of subsequent algorithms.

2.2. Pade Forms

2.2.1. Definitions

The class P of formal power series over a field F consists of expressions of the form

$$A(z) = \sum_{i=0}^{\infty} a_i z^i$$

with coefficients $a_i \in F$. We denote the units of P by

$$U = \{A(z) = \sum_{i=0}^{\infty} a_i z^i \mid a_0 \neq 0, A(z) \in P\}.$$

Associated with each such unit is a set of rational functions defined as follows:

Definition: Let $A(z) \in U$, and let m and n be non-negative integers. The rational form

$$\frac{U_{mn}(z)}{V_{mn}(z)} = \frac{u_0 + u_1 z + \cdots + u_m z^m}{v_0 + v_1 z + \cdots + v_n z^n} \quad (2.1)$$

is called a **Pade form** of type (m, n) for $A(z)$ if

$$(a) \quad V_{mn}(z) \neq 0, \text{ and} \quad (2.2)$$

$$(b) \quad A(z) V_{mn}(z) - U_{mn}(z) = O(z^{m+n+1}) \quad (2.3)$$

The (algebraic) O -symbol indicates that the right side is a power series beginning with the power z^{m+n+1} , $0 \leq k < \infty$; $k = +\infty$ means that $A(z) V_{mn}(z) - U_{mn}(z) = 0$.

From an analytical point of view, this means that the rational form $U_{mn}(z)/V_{mn}(z)$ is determined so that its Maclaurin expansion agrees with $A(z)$ as far as possible. That is, equation (2.3) corresponds to a linear system of $m+n+1$ equations for the u_i and v_i on

which u_0, \dots, u_m and v_0, \dots, v_n are the unknowns.

$$\sum_{i=0}^n a_i (v_i z^i)^{(k)} = \begin{cases} u_k & k = 0, 1, \dots, m \\ 0 & k = m+1, \dots, m+n+1 \end{cases} \quad (2.4)$$

where $(v_i z^i)^{(k)}$ denotes the k th derivative of $v_i z^i$ with respect to z .

$$\begin{vmatrix} 1 & 1 & 1 \\ 1 & 2 & 4 \\ 1 & 3 & 9 \end{vmatrix} \quad \begin{vmatrix} 1 & 1 & 1 \\ 1 & 2 & 4 \\ 1 & 3 & 9 \end{vmatrix}$$

¹ By convention, $v_i = 0$ if $i < 0$.
² Through the use of the chain rule, the above system of equations can be written in the form

$$\begin{bmatrix} a_{m-n+1} & \cdots & a_{m+1} \\ \vdots & \ddots & \vdots \\ a_m & \cdots & a_{m+n} \end{bmatrix} \begin{bmatrix} v_n \\ \vdots \\ v_0 \end{bmatrix} = \begin{bmatrix} 0 \\ \vdots \\ 0 \end{bmatrix}. \quad (2.6)$$

The matrix

$$H_m = \begin{bmatrix} a_{m-n+1} & \cdots & a_m \\ \vdots & \ddots & \vdots \\ a_m & \cdots & a_{m+n-1} \end{bmatrix} \quad (2.7)$$

associated with the power series $A(z)$ is called a **Hankel matrix**. With this definition, the system (2.6) becomes

$$H_m \begin{bmatrix} v_n \\ \vdots \\ v_0 \end{bmatrix} = v_0 \begin{bmatrix} a_{m+1} \\ \vdots \\ a_{m+n} \end{bmatrix}. \quad (2.8)$$

Denote the determinant of H_m by $\det(H_m)$, and let $\det(H_0) = 1$.

2.2.2 Existence and Non uniqueness

Theorem 2.1 (Frobenius): There always exists a non trivial solution of the system (2.5) and (2.6). Equivalently, Padé forms of type (n, n) for $A(z) \in \mathbb{H}$ always exist.

Proof: The existence of $V^* = (v_n, \dots, v_0)$ satisfying (2.6) follows immediately from the fact that the $n+1$ columns of the matrix in (2.6) are vectors of length n and must therefore be linearly dependent. The vector V^* satisfying (2.5) is then obtained simply by

It is clear that if U_{mn} and V_{mn} is a solution of (2.5) and (2.6), then so is $\alpha \cdot U_{mn}$ and $\alpha \cdot V_{mn}$, where α is a non-zero constant. The non-uniqueness of the solution, however, is more profound than this, and the next few results identify the true nature of the family of solutions.

Lemma 2.2: Let $U_{mn}(z)/V_{mn}(z)$ be a Padé form of type (m, n) for $\Lambda(z) \in U$, and let $D(z)$ be a non-zero polynomial of degree $\partial(D) \leq \lambda$, where $\lambda = \min\{m - \partial(U_{mn}), n - \partial(V_{mn})\}$. Then $U'_{mn}(z)/V'_{mn}(z)$, where

$$U'_{mn}(z) = U_{mn}(z)D(z) \quad \text{and} \quad V'_{mn}(z) = V_{mn}(z)D(z)$$

is also a Padé form of type (m, n) for $\Lambda(z)$.

Proof: With the $U_{mn}(z)$ and $V_{mn}(z)$ defined above, equation (2.5) becomes

$$\begin{bmatrix} 0 \\ U_{mn} \\ \vdots \\ U_n \end{bmatrix} = \begin{bmatrix} a_{m-n} \\ \vdots \\ a_1 \\ a_0 \end{bmatrix} \begin{bmatrix} 0 \\ V_{mn} \\ \vdots \\ V_n \end{bmatrix} \quad (2.9)$$

where the zero vector of length n is the zero vector of length n .

$$\begin{bmatrix} 0 \\ U_{mn} \\ \vdots \\ U_n \end{bmatrix} = \begin{bmatrix} 0 \\ a_{m-n} \\ \vdots \\ a_1 \\ a_0 \end{bmatrix} \begin{bmatrix} 0 \\ V_{mn} \\ \vdots \\ V_n \end{bmatrix}$$

$$\begin{bmatrix} 0 \\ U_{mn} \\ \vdots \\ U_n \end{bmatrix} = \begin{bmatrix} 0 \\ a_{m-n} \\ \vdots \\ a_1 \\ a_0 \end{bmatrix} \begin{bmatrix} 0 \\ V_{mn} \\ \vdots \\ V_n \end{bmatrix}$$

$$= \begin{bmatrix} a_{m-n} & \cdots & a_m \\ \vdots & & \vdots \\ a_{-n} & \cdots & a_0 \end{bmatrix} \begin{bmatrix} d_0 & \cdots & d_\lambda \\ & \ddots & d_\lambda \\ & & 0 \\ & & & d_0 \end{bmatrix} \begin{bmatrix} 0_\lambda \\ v_{n-\lambda} \\ \vdots \\ v_0 \end{bmatrix}.$$

That is,

$$\begin{bmatrix} u'_m \\ \vdots \\ u'_0 \end{bmatrix} = \begin{bmatrix} a_{m-n} & \cdots & a_m \\ \vdots & & \vdots \\ a_{-n} & \cdots & a_0 \end{bmatrix} \begin{bmatrix} v'_n \\ \vdots \\ v'_0 \end{bmatrix}.$$

Therefore U'_m and V'_m satisfy equation (2.5). The fact that V'_m also satisfies (2.6) follows from

$$\begin{aligned} & \begin{bmatrix} a_{m-n+1} & \cdots & a_{m+1} \\ \vdots & & \vdots \\ a_{-n} & \cdots & a_m \end{bmatrix} \begin{bmatrix} v'_{n-1} \\ \vdots \\ v'_0 \end{bmatrix} \\ &= \begin{bmatrix} a_{m-n+1} & \cdots & a_{m+1} \\ \vdots & & \vdots \\ a_{-n} & \cdots & a_m \end{bmatrix} \begin{bmatrix} d_0 & \cdots & d_\lambda \\ & \ddots & d_\lambda \\ & & 0 \\ & & & d_0 \end{bmatrix} \begin{bmatrix} 0_\lambda \\ v_{n-\lambda} \\ \vdots \\ v_0 \end{bmatrix} \\ &= \begin{bmatrix} d_0 & \cdots & d_\lambda \\ & \ddots & d_\lambda \\ & & 0 \\ & & & d_0 \end{bmatrix} \begin{bmatrix} a_{m-n+1} & \cdots & a_{m+1} \\ \vdots & & \vdots \\ a_{-n} & \cdots & a_m \end{bmatrix} \begin{bmatrix} v_{n-\lambda} \\ \vdots \\ v_0 \end{bmatrix} \\ &= \begin{bmatrix} d_0 & \cdots & d_\lambda \\ & \ddots & d_\lambda \\ & & 0 \\ & & & d_0 \end{bmatrix} \begin{bmatrix} a_{m-n+1} & \cdots & a_{m+1} \\ \vdots & & \vdots \\ a_{m-n} & \cdots & a_m \\ \vdots & & \vdots \\ a_{-n} & \cdots & a_{m+1} \end{bmatrix} \begin{bmatrix} 0_\lambda \\ v_{n-\lambda} \\ \vdots \\ v_0 \end{bmatrix} \end{aligned}$$

$$= \begin{bmatrix} 0 \\ \vdots \\ 0 \end{bmatrix} \quad (2.10)$$

In the last equality of (2.10), use was made both of equation (2.9) and of equation (2.6) with

$$V_{mn} = (0, v_{n-1}, \dots, v_0) \quad \blacksquare$$

The converse of Lemma 2.2 is given by

Lemma 2.3. Let $U'_{mn}(z)/V'_{mn}(z)$ be a Padé form of type (m, n) for $A(z) \in \mathbb{U}$, and let $D(z)$, $D(0) \neq 0$, be a common divisor of U'_{mn} and V'_{mn} . Then $U_{mn}(z)/V_{mn}(z)$, where

$$U_{mn}(z) = U'_{mn}(z)/D(z) \quad \text{and} \quad V_{mn}(z) = V'_{mn}(z)/D(z),$$

is also a Padé form of type (m, n) for $A(z)$.

Proof: The arguments for the proof are similar to those of the proof of Lemma 2.2. \square

The condition that $D(0) \neq 0$ in Lemma 2.3 is necessary only for the case of the Padé form of type $(0, n)$. For the case of the Padé form of type $(m, 0)$, the condition is not necessary.

Example 2.1. Let $A(z) = 1/z^2 + 2z^2 + 3z^3 + 4z^4 + 5$.

Let $U'_{2,2}(z) = 1/z^2 + 2z^2 + 3z^3 + 4z^4 + 5$ and $V'_{2,2}(z) = 1/z^2 + 2z^2 + 3z^3 + 4z^4 + 5$ yield a Padé form of type $(2, 2)$ for $A(z)$.

$$U'_{2,2}(z)/V'_{2,2}(z) = 1/z^2 + 2z^2 + 3z^3 + 4z^4 + 5 = A(z)$$

A common divisor of $U'_{2,2}(z)$ and $V'_{2,2}(z)$ is $D(z) = 1/z^2 + 2z^2 + 3z^3 + 4z^4 + 5$. The rational function $U_{2,2}(z)/V_{2,2}(z)$ where

$$U_{2,2}(z) = U'_{2,2}(z)/D(z) = 1 \quad \text{and} \quad V_{2,2}(z) = V'_{2,2}(z)/D(z) = 1$$

is also a Padé form of type $(2, 2)$ for $A(z)$.

$$A(z) \cdot V(z) - U(z) = O(z^6). \quad \blacksquare$$

2.2.3. General Structure of Pade Forms

Let H_{mn} be given. If $\det(H_{mn}) \neq 0$, then system (2.8) exhibits a one-parameter family of solutions. Indeed, if $\det(H_{mn}) \neq 0$ and $v_0 = 1$ in (2.8), then the solution of (2.8) for $V_{mn}(z)$ and of (2.5) for $U_{mn}(z)$ is unique. Pade forms for which $V(0) = 1$, i.e., $v_0 = 1$ are said to be **normalized**. In addition, we have

Lemma 2.4. If $\det(H_{mn}) \neq 0$, then the solutions U_{mn} and V_{mn} of (2.5) and (2.6) are relatively prime.

Proof: Suppose the contrary, and let $D(z)$ be a non-trivial common divisor of $U_{mn}(z)$ and $V_{mn}(z)$.

If $D(0) = 0$, then $V_{mn}(0) = 0$, i.e., $v_0 = 0$. Consequently, (2.8) has only the trivial solution and this contradicts the definition of a Pade form.

If $D(0) \neq 0$, let $\bar{D}(z) = D(z)/D(0)$. Then, by Lemma 2.3, $U'(z) = U(z)/\bar{D}(z)$ and $V'(z) = V(z)/\bar{D}(z)$ are solutions to systems (2.5) and (2.6). But $V'(0) = V(0)$, and equation (2.8) together with the condition that $\det(H_{mn}) \neq 0$ then imply that $V'(z) = V(z)$. Thus, $D(z)$ must be a polynomial of degree zero, which contradicts the initial assumption that $D(z)$ is non-trivial. \blacksquare

The converse of Lemma 2.4, however, is not true, as can be seen in the following example.

Example 2.2: Consider again the power series $A(z)$ in example 2.1. It can be verified that a Pade form of type (4, 3) for $A(z)$ is $U_{4,3}(z)/V_{4,3}(z) = (1 + z + z^2)/(1 - 2z + z^2)$, where $U_{4,3}(\cdot)$ and $V_{4,3}(\cdot)$ are relatively prime. However, the Hankel matrix

$$H_{4,3} = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 4 \\ 3 & 4 & 5 \end{bmatrix}$$

associated with the Pade form of type (4, 3) is singular, so that $\det(H_{4,3}) = 0$. \blacksquare

Given the Hankel matrix $H_{m,n}$, let $H_{m-\lambda, n-\lambda}$ be its largest non-singular principal submatrix. That is, λ is the smallest non-negative integer for which $\det(H_{m-\lambda, n-\lambda}) \neq 0$. From Lemma 2.4, a Pade form

$$\frac{U_{m-\lambda, n-\lambda}^*(z)}{V_{m-\lambda, n-\lambda}^*(z)} = \frac{\sum_{l=0}^{m-\lambda} u_l^* z^l}{\sum_{l=0}^{n-\lambda} v_l^* z^l} \quad (2.11)$$

of type $(m-\lambda, n-\lambda)$ for $A(z)$ exists, where $U_{m-\lambda, n-\lambda}^*(z)$ and $V_{m-\lambda, n-\lambda}^*(z)$ are relatively prime and $v_0^* \neq 0$. The Pade form is unique up to a multiplicative constant and satisfies

$$\begin{bmatrix} u_{m-\lambda}^* \\ \vdots \\ u_0^* \end{bmatrix} = \begin{bmatrix} a_{m-n} & \cdots & a_{m-\lambda} \\ \vdots & & \vdots \\ a_{-n+\lambda} & \cdots & a_0 \end{bmatrix} \begin{bmatrix} v_{n-\lambda}^* \\ \vdots \\ v_0^* \end{bmatrix} \quad (2.12)$$

and

$$\begin{bmatrix} a_{m-n+1} & \cdots & a_{m-\lambda+1} \\ \vdots & & \vdots \\ a_{-n} & \cdots & a_{-n-2\lambda} \end{bmatrix} \begin{bmatrix} v_{n-\lambda}^* \\ \vdots \\ v_0^* \end{bmatrix} = \begin{bmatrix} 0 \\ \vdots \\ 0 \end{bmatrix} \quad (2.13)$$

In addition

$$A(z) - V_{m-\lambda, n-\lambda}^*(z) \cdot U_{m-\lambda, n-\lambda}^*(z)^{-1} = \sum_{l=0}^{\infty} r_l z^l, \quad (2.14)$$

where $r_0 \neq 0$, if $k < \infty$.

Theorem 2.5: Let λ be such that $H_{m-\lambda, n-\lambda}$ is the largest non-singular principal submatrix in $H_{m,n}$, and let k be determined by (2.14). Then $k \geq \lambda$. In addition, let $l = \min \{2\lambda, k\}$. Then a basis for the solution space of

$$\begin{bmatrix} a_{m-n+1} & \cdots & a_{m+1} \\ \vdots & & \vdots \\ a_{m-\lambda} & \cdots & a_{m+1} \end{bmatrix} \begin{bmatrix} v_n \\ \vdots \\ v_0 \end{bmatrix} = \begin{bmatrix} 0 \\ \vdots \\ 0 \end{bmatrix} \quad (2.15)$$

is given by the $l-\lambda+1$ vectors

$$\{(v_{n-\lambda}^*, \dots, v_0^*, 0_\lambda)^\top, \dots, (0_{l-\lambda}, v_{n-\lambda}^*, \dots, v_0^*, 0_{2\lambda-l})^\top\}.$$

Proof: If $\lambda=0$, then the theorem follows trivially from Lemma 2.4. If $\lambda \neq 0$, then the multiplication of (2.15) by

$$\begin{matrix} n-\lambda \\ \lambda \end{matrix} \left\{ \begin{bmatrix} 1 & & & & 0 \\ & \ddots & & & \\ & & 0 & & \\ & & & 1 & \\ & & & & \ddots \\ & & & & & 0 \end{bmatrix} \right.$$

yields

$$\begin{bmatrix} a_{m-n+1} & \cdots & a_{m+1} \\ \vdots & & \vdots \\ a_{m-\lambda} & \cdots & a_{m+1} \\ \vdots & & \vdots \\ 0 & r_0 & \cdots & r_{2\lambda-k-1} \end{bmatrix} \begin{bmatrix} v_n \\ \vdots \\ v_0 \end{bmatrix} = \begin{bmatrix} 0 \\ \vdots \\ 0 \end{bmatrix} \quad (2.16)$$

in the case that $k \leq \lambda$, or

$$\begin{bmatrix}
 a_{m-n+1} & \dots & a_m & a_{m+1} \\
 & & & \\
 a_{m-\lambda} & \dots & a_{m+n-\lambda} \\
 & & 0_{l-\lambda} \\
 & & r_0 \\
 0 & & r_0 & r_1 \\
 & & & \\
 & & r_0 & \dots & r_{2\lambda-l-1}
 \end{bmatrix}
 \begin{bmatrix}
 v_n \\
 \vdots \\
 v_0
 \end{bmatrix}
 =
 \begin{bmatrix}
 0 \\
 \vdots \\
 0
 \end{bmatrix}
 \quad (2.17)$$

in the case that $k \geq \lambda$. But the largest non-singular principal submatrix in (2.16) has order n , which violates the assumption that $\lambda \neq 0$. Thus, when $\lambda \neq 0$, only (2.17) is possible, which proves that $k \geq \lambda$.

From (2.17), it follows that the coefficient matrix of (2.17) has rank $n + \lambda - l$, and therefore the solution space of (2.15) is of dimension $l - \lambda + 1$. But, using (2.13) and (2.14), it is easy to see that all the vectors

$$\{(v_{n-\lambda}^*, \dots, v_0^*, 0_\lambda)', \dots, (0_{l-\lambda}, v_{n-\lambda}^*, \dots, v_0^*, 0_{2\lambda-l})'\}$$

satisfy (2.17): ■

Corollary 2.6 (Pade): The general solution of

$$\begin{bmatrix}
 u_m \\
 \vdots \\
 u_0
 \end{bmatrix}
 =
 \begin{bmatrix}
 a_{m-n} & \dots & a_m \\
 \vdots & & \vdots \\
 a_{m-\lambda} & \dots & a_0
 \end{bmatrix}
 \begin{bmatrix}
 v_n \\
 \vdots \\
 v_0
 \end{bmatrix}$$

and

$$\begin{bmatrix}
 a_{m-n} & \dots & a_{m-1} \\
 \vdots & & \vdots \\
 a_m & \dots & a_{m-n}
 \end{bmatrix}
 \begin{bmatrix}
 v_n \\
 \vdots \\
 v_0
 \end{bmatrix}
 =
 \begin{bmatrix}
 0 \\
 \vdots \\
 0
 \end{bmatrix}$$

is given by

$$U_{mn}(z) = z^{2\lambda-l} \left(\sum_{i=0}^{l-\lambda} \alpha_i z^i \right) \left(\sum_{i=0}^{n-\lambda} u_i^* z^i \right) \quad (2.18)$$

$$V_{mn}(z) = z^{2\lambda-l} \left(\sum_{i=0}^{l-\lambda} \alpha_i z^i \right) \left(\sum_{i=0}^{n-\lambda} v_i^* z^i \right), \quad (2.19)$$

where l and λ are defined in Theorem 2.5 and α_i , $0 \leq i \leq l-\lambda$, is arbitrary.

Proof: Linear combinations of the basis vectors for the solution space can be written as

$$V_{mn} = \alpha_{l-\lambda} \begin{bmatrix} v_{n-\lambda}^* \\ \vdots \\ v_0^* \\ 0_\lambda \end{bmatrix} + \dots + \alpha_0 \begin{bmatrix} 0_{l-\lambda} \\ v_{n-\lambda}^* \\ \vdots \\ v_0^* \\ 0_{2\lambda-l} \end{bmatrix}$$

$$= \begin{bmatrix} 0_\lambda & \alpha_0 & \dots & \alpha_{l-\lambda} & 0 \\ & & & \alpha_{l-\lambda} & \\ & 0 & & \alpha_0 & \\ & & & 0_{2\lambda-l} & \end{bmatrix} \begin{bmatrix} 0_\lambda \\ v_{n-\lambda}^* \\ \vdots \\ v_0^* \end{bmatrix},$$

which in polynomial form becomes (2.19). Furthermore, using equation (2.12) and the results of Theorem 2.5, it follows that

$$\begin{bmatrix} 0_l \\ u_{m-\lambda}^* \\ \vdots \\ u_0^* \\ 0_{\lambda-l} \end{bmatrix} = \begin{bmatrix} a_{m-n} & \dots & a_m \\ \vdots & & \vdots \\ a_{-n} & \dots & a_0 \end{bmatrix} \begin{bmatrix} 0_l \\ v_{n-\lambda}^* \\ \vdots \\ v_0^* \\ 0_{\lambda-l} \end{bmatrix}$$

for $0 \leq i \leq l-\lambda$. Thus, the polynomial form for $U_{mn}(z)$ corresponding to $V_{mn}(z)$ is given by (2.18). ■

2.3. Pade Fractions

2.3.1. Scaled Pade Fractions

Let H_{mn} for $A(z) \in U$ be given, and let $H_{m-\lambda, n-\lambda}$ be its largest non-singular principal submatrix. If the rational function

$$\frac{U_{m-\lambda, n-\lambda}^*(z)}{V_{m-\lambda, n-\lambda}^*(z)} = \frac{\sum_{i=0}^{m-\lambda} u_i^* z^i}{\sum_{i=0}^{n-\lambda} v_i^* z^i} \quad (2.20)$$

is the Pade form of type $(m-\lambda, n-\lambda)$ for $A(z)$, then

Definition: The scaled Pade fraction of type (m, n) for $A(z)$ is defined to be $\gamma_{mn}(z) = S_{mn}(z) / T_{mn}(z)$, where

$$S_{mn}(z) = z^\lambda \sum_{i=0}^{m-\lambda} u_i^* z^i \quad (2.21)$$

$$T_{mn}(z) = z^\lambda \sum_{i=0}^{n-\lambda} v_i^* z^i \quad (2.22)$$

Theorem 2.7: The scaled Pade fraction γ_{mn} of type (m, n) for $A(z)$ is a Pade form of type (m, n) for $A(z)$. Furthermore, γ_{mn} is unique up to a multiplicative constant.

Proof: The scaled Pade fraction is obtained simply by setting $\alpha_{i-\lambda} \neq 0$ and $\alpha_0 = \alpha_1 = \dots = \alpha_{i-\lambda-1} = 0$ in equations (2.18) and (2.19) of Corollary 2.6. ■

As an immediate consequence, we have

Corollary 2.8: $\det(H_{mn}) \neq 0$ if and only if $T_{mn}(0) \neq 0$.

An alternative definition of scaled Pade fractions is given by

Theorem 2.9. $S_{mn}(z)/T_{mn}(z)$, where $T_{mn}(z) \neq 0$ is the scaled Pade fraction of type (m,n) for $A(z)$

if

$$(a) \min \{ m - \partial(S_{mn}), n - \partial(T_{mn}) \} = 0,$$

$$(b) \text{GCD}(S_{mn}, T_{mn}) = z^{\lambda_{mn}}, \text{ for some } \lambda_{mn} \geq 0, \text{ and}$$

$$(c) A(z) \cdot T_{mn}(z) - S_{mn}(z) = O(z^{m+n+1}).$$

Proof: The result is an immediate consequence of Theorem 2.7. ■

Thus, when normalized, the scaled Pade fraction of type (m,n) for $A(z)$ exists uniquely, and in addition satisfies the systems (2.5) and (2.6). This is tremendous advantage for our purposes over other definitions of Pade fractions described below, since it simplifies the development of algorithms in subsequent chapters.

2.3.2. Non-scaled Pade Fractions

Let $U_{mn}(z)/V_{mn}(z)$ be a Pade form of type (m,n) for $A(z)$, and let

$$D(z) = \text{GCD}(U_{mn}, V_{mn}).$$

Gragg [GRA 72] defines the Pade fraction of type (m,n) for $A(z)$ to be $P_{mn}(z)/Q_{mn}(z)$, where

$$P_{mn}(z) = U_{mn}(z) / D(z)$$

$$Q_{mn}(z) = V_{mn}(z) / D(z),$$

and in addition $Q_{mn}(0) = 1$.

That is,

$$P_{mn}(z) = z^{-\lambda} S_{mn}(z) = U_{mn-\lambda, n-\lambda}^s(z)$$

$$Q_{mn}(z) = z^{-\lambda} T_{mn}(z) = V_{m-\lambda, n-\lambda}^*(z),$$

where $U_{m-\lambda, n-\lambda}^*(z)$ and $V_{m-\lambda, n-\lambda}^*(z)$ are defined in (2.20). Thus, the Pade fraction of type (m, n) always exists and is unique. However, $P_{mn}(z) / Q_{mn}(z)$ is a member of the family of Pade forms of type (m, n) for $A(z)$ given by (2.18) and (2.19) only if $l = 2\lambda$, or equivalently, if $k \geq 2\lambda$ in equation (2.14). Thus, in the case that $k \geq 2\lambda$, the Pade fraction is obtained from the Pade form by setting $\alpha_0 = 1, \alpha_1 = \dots = \alpha_{l-\lambda} = 0$ in equation (2.18) and (2.19). In the remaining case that $k < 2\lambda$, the Pade fraction $P_{mn}(z) / Q_{mn}(z)$ does not satisfy the systems (2.5) and (2.6).

Baker's [BAK75] perspective is different in a very subtle way. He requires that $P_{mn}(z) / Q_{mn}(z)$ satisfies (2.5) and (2.6) and in addition $Q_{mn}(0) = 1$ and $\text{GCD}(P_{mn}, Q_{mn}) = 1$. But once again, from Theorem 2.5, this is possible only if $k \geq 2\lambda$, in which case $P_{mn}(z) / Q_{mn}(z)$ is obtained by setting $\alpha_0 = 1, \alpha_1 = \dots = \alpha_{l-\lambda} = 0$ in the equations (2.18) and (2.19). Thus, from Baker's point of view, a Pade fraction may not exist but whenever it does, it is unique and satisfies systems (2.5) and (2.6).

CHAPTER 3

COMPUTATION OF OFF-DIAGONAL SCALED PADE FRACTIONS

3.1. Introduction

In this chapter, derived is a new relationship between three scaled Pade fractions lying along an off-diagonal path of the Pade table for a power series $A(z)$. This relationship is described in Section 3.2 and is used in Section 3.3 to develop an algorithm to iteratively compute a sequence of scaled Pade fractions along the off-diagonal path. However, the algorithm is not totally iterative, since the relationship recursively involves a scaled Pade fraction of a different power series $\bar{A}(z)$, computed from $A(z)$. By doubling the step size along the off-diagonal path at each iteration, it is shown that the complexity of the algorithm for computing the n -th scaled Pade fraction along the path is $(n \log^2 n)$, assuming fast polynomial methods are used.

The algorithm of Section 3.3 can be used to compute scaled Pade fractions for the quotient $A(z)/B(z)$ of two power series $A(z)$ and $B(z)$. This can be accomplished by formally computing the inverse of $B(z)$, multiplying it by $A(z)$, and then applying the algorithm to the result. The dominating cost of this procedure is the cost of computing the inverse of $B(z)$. This cost, however, can be eliminated by an obvious modification of the algorithm, which is given in Section 3.4.

If the step size along the off-diagonal path is shortened, all scaled Pade fractions along the path can be computed. The recursive call alluded to earlier then becomes trivial, and the resulting algorithm, given in Section 3.5, becomes truly iterative. In the case that the path is along the diagonal, the algorithm is identical to the one given by Calvey and Kao [CABP1]. This algorithm can be adapted to fast polynomial multiplication, giving a complexity $O(n^2)$.

3.2. Preliminary Results

The scaled Padé fractions can be arranged in a doubly infinite array as follows:

Let $Y_{m,n}$ denote the (m,n) -th scaled Padé fraction for $A(z)$. Then

$Y_{-1,-1}$	$Y_{-1,0}$	$Y_{-1,1}$	$Y_{-1,2}$
$Y_{0,-1}$	$Y_{0,0}$	$Y_{0,1}$	$Y_{0,2}$
$Y_{1,-1}$	$Y_{1,0}$	$Y_{1,1}$	$Y_{1,2}$
$Y_{2,-1}$	$Y_{2,0}$	$Y_{2,1}$	$Y_{2,2}$
$Y_{3,-1}$	$Y_{3,0}$	$Y_{3,1}$	$Y_{3,2}$
$Y_{4,-1}$	$Y_{4,0}$	$Y_{4,1}$	$Y_{4,2}$

is the (extended) scaled Padé table for $A(z)$. The modifying

operation used in the table is the following:

$$Y_{m,n} = \frac{P_m(z)Q_n(z)}{Q_m(z)P_n(z)}, \quad \text{where } P_m(z) = \sum_{k=0}^m p_k z^k, \quad Q_n(z) = \sum_{k=0}^n q_k z^k,$$

$$p_k = a_k, \quad q_k = a_{-k}, \quad \text{for } k = 0, 1, \dots, m, n.$$

Table 1. The scaled Padé table for $A(z)$.

$Y_{-1,-1}$	$Y_{-1,0}$	$Y_{-1,1}$	$Y_{-1,2}$
$Y_{0,-1}$	$Y_{0,0}$	$Y_{0,1}$	$Y_{0,2}$
$Y_{1,-1}$	$Y_{1,0}$	$Y_{1,1}$	$Y_{1,2}$
$Y_{2,-1}$	$Y_{2,0}$	$Y_{2,1}$	$Y_{2,2}$
$Y_{3,-1}$	$Y_{3,0}$	$Y_{3,1}$	$Y_{3,2}$
$Y_{4,-1}$	$Y_{4,0}$	$Y_{4,1}$	$Y_{4,2}$

Let m and n be non-negative integers such that $m+n \leq M$. The modified Padé fraction is

$$Y_{m,n} = \frac{P_m(z)Q_n(z)}{Q_m(z)P_n(z)},$$

Note that the (extended) scaled Padé table for $A(z)$ is a doubly infinite array of fractions. The

with the construction of the scaled Pade fraction $\gamma_m(z)$, given that $\Gamma_N(A)$ already exists.

Without loss of generality, assume that $m \geq n$ (otherwise, the same arguments can be applied to

$1/A(z) \in U$). Let

$$M = N + (m - n). \quad (3.3)$$

Then, (m, n) and (M, N) both lie along the $(m-n)$ th off-diagonal path of the scaled Pade table

$\Gamma(A)$ (see Figure 3.1)




Figure 3.1

Let $\gamma_m(z)$ be the scaled Pade fraction of type (M, N) for $A(z)$, equation (2.3) becomes

$$A(z) = \gamma_m(z) = \frac{P_m(z)}{Q_m(z)} = \frac{P_m(z)}{Q_m(z)} \quad (3.4)$$

where $\mu_{MN} \geq 0$, and $R_1(0) \neq 0$ if $\mu_{MN} < \infty$. Let

$$z^{\lambda_{MN}} = \text{GCD}(S_{MN}, T_{MN}). \quad (3.5)$$

To construct $\gamma_{mn}(z)$, two separate cases, $\mu_{MN} \geq (n-N)$ and $\mu_{MN} < (n-N)$, arise. These two cases are considered separately in Theorem 3.1 and Theorem 3.4.

Theorem 3.1: If $\mu_{MN} \geq (n-N)$ in (3.4), then the scaled Dole function $\gamma_{mn}(z) = S_{mn}(z)/T_{mn}(z)$ is given by

$$S_{mn}(z) = z^{n-N} S_{MN}(z) \quad (3.6)$$

$$T_{mn}(z) = z^{n-N} T_{MN}(z) \quad (3.7)$$

Proof: Clearly from relation (3.3), it follows that

$$\partial(S_{mn}) = n - N + \partial(S_{MN})$$

$$< n + M - N$$

$$= m$$

Similarly $\partial(T_{mn}) \leq n$. Furthermore, it is clear that $\min\{m - \partial(S_{mn}), n - \partial(T_{mn})\} = 0$ from the equality $\min\{M - \partial(S_{MN}), M - \partial(T_{MN})\} = 0$. Using the fact that $(n-N) \geq 0$, it follows that

$$S_{mn}(z) = z^{n-N} (A(z)T_{MN}(z) + S_{MN}(z))$$

$$= z^{n-N} (z^{M-\partial(S_{MN})} R_1(z) + S_{MN}(z))$$

$$= z^{n-N} z^{M-\partial(S_{MN})} R_1(z)$$

$$= z^{m-\partial(S_{mn})} R_1(z)$$

$$\text{Hence, } \text{GCD}(S_{mn}, T_{mn}) = z^{n-N} \text{GCD}(S_{MN}, T_{MN}) = z^{\lambda_{MN}}.$$

$$\text{Hence, } \gamma_{mn}(z) = \gamma_{MN}(z) = \frac{S_{MN}(z)}{T_{MN}(z)} = \frac{S_{mn}(z)}{T_{mn}(z)} = \gamma_{mn}(z).$$

For the case $\mu_{MN} < (n-N)$, let

$$M^* = M - \lambda_{MN} - 1, \quad (3.8)$$

$$N^* = N - \lambda_{MN} - 1 \quad (3.9)$$

(see Figure 3.1) Clearly, the scaled fraction $\gamma_{M^*N^*}(z)$ for $A(z)$ satisfies

$$A(z) \cdot T_{M^*N^*}(z) - S_{M^*N^*}(z) = z^{M^*+N^*+1} R_0(z), \quad (3.10)$$

where $R_0(0) \neq 0$. In addition, by Theorem 2.7 and by the definition of M^* and N^* , it follows that

$$S_{MN}(z) / T_{MN}(z) \neq S_{M^*N^*}(z) / T_{M^*N^*}(z). \quad (3.11)$$

From the two unit power series, $R_1(z)$ and $R_0(z)$ given by (3.4) and (3.10), respectively, construct the unique power series

$$\bar{A}(z) = R_0(z) / R_1(z). \quad (3.12)$$

Associated with $\bar{A}(z) \in U$, let

$$\bar{m} = n - N + \lambda_{MN}, \quad (3.13)$$

$$\bar{n} = n - N - \mu_{MN} - 1. \quad (3.14)$$

Now assume that the \bar{n} -truncated scaled Padé table, $\Gamma_{\bar{n}}(\bar{A})$, for $\bar{A}(z)$ has been constructed, as well.

That is, assume that the (\bar{m}, \bar{n}) scaled Padé fraction, $\bar{\gamma}_{\bar{m}\bar{n}}(z) = \bar{S}_{\bar{m}\bar{n}}(z) / \bar{T}_{\bar{m}\bar{n}}(z)$, such that

$$\bar{A}(z) - \bar{T}_{\bar{m}\bar{n}}(z) / \bar{S}_{\bar{m}\bar{n}}(z) = O(z^{\bar{m}+\bar{n}+1}) \quad (3.15)$$

is available.

The scaled Padé fractions $\gamma_{MN}(z)$, $\gamma_{M^*N^*}(z)$ and $\bar{\gamma}_{\bar{m}\bar{n}}(z)$ provide sufficient information to obtain directly a Padé form of type (m, n) for $A(z)$. This Padé form is constructed in Lemma 3.2 below. It is shown, later in Theorem 3.1, that this form is also the scaled Padé fraction of type (m, n) for $A(z)$.

Lemma 3.2: Let $\mu_{MN} < (n-N)$ in equation (3.4), and let

$$S_{mn}(z) = z^{-\lambda_{MN}} S_{MN}(z) \bar{S}_{\bar{m}\bar{n}}(z) - z^\alpha S_{M^*N^*}(z) \bar{T}_{\bar{m}\bar{n}}(z) \quad (3.16)$$

$$T_{mn}(z) = z^{-\lambda_{MN}} T_{MN}(z) \bar{S}_{\bar{m}\bar{n}}(z) - z^\alpha T_{M^*N^*}(z) \bar{T}_{\bar{m}\bar{n}}(z), \quad (3.17)$$

where $\alpha = \lambda_{MN} + \mu_{MN} + 2$. Then $S_{mn}(z) / T_{mn}(z)$ is a Padé form of type (m, n) for $A(z)$.

Proof:

$$\begin{aligned} \partial(S_{mn}) &= \max\{\partial(z^{-\lambda_{MN}} S_{MN} \bar{S}_{\bar{m}\bar{n}}), \partial(z^\alpha S_{M^*N^*} \bar{T}_{\bar{m}\bar{n}})\} \\ &< \max\{-\lambda_{MN} + M + \bar{m}, \alpha + M^* + \bar{n}\} \\ &= \max\{-\lambda_{MN} + M + (n - N + \lambda_{MN}), \\ &\quad (\lambda_{MN} + \mu_{MN} + 2) + (M - \lambda_{MN} - 1) + (n - N - \mu_{MN} - 1)\} \\ &= \max\{m, m\} \\ &= m. \end{aligned}$$

Similarly, $\partial(T_{mn}) \leq n$. Moreover, the fact that $S_{MN}(z) / T_{MN}(z)$, $S_{M^*N^*}(z) / T_{M^*N^*}(z)$ and $\bar{S}_{\bar{m}\bar{n}}(z) / \bar{T}_{\bar{m}\bar{n}}(z)$ are the scaled Padé fractions yields immediately that

$$\min\{m - \partial(S_{mn}), n - \partial(T_{mn})\} = 0$$

In addition, from (3.16), (3.17) and (3.15), it follows that

$$\begin{aligned} A(z) &= \frac{S_{mn}(z)}{T_{mn}(z)} = \frac{z^{-\lambda_{MN}} S_{MN}(z) \bar{S}_{\bar{m}\bar{n}}(z) - z^\alpha S_{M^*N^*}(z) \bar{T}_{\bar{m}\bar{n}}(z)}{z^{-\lambda_{MN}} T_{MN}(z) \bar{S}_{\bar{m}\bar{n}}(z) - z^\alpha T_{M^*N^*}(z) \bar{T}_{\bar{m}\bar{n}}(z)} \\ &= \frac{z^{-\lambda_{MN}} \bar{S}_{\bar{m}\bar{n}}(z) \{A(z) T_{MN}(z) - S_{M^*N^*}(z)\}}{z^{-\lambda_{MN}} \bar{T}_{\bar{m}\bar{n}}(z) \{A(z) T_{MN}(z) - S_{M^*N^*}(z)\}} \\ &= \frac{\bar{S}_{\bar{m}\bar{n}}(z) \{P_1(z) - S_{M^*N^*}(z)\}}{\bar{T}_{\bar{m}\bar{n}}(z) \{P_1(z) - S_{M^*N^*}(z)\}} \end{aligned}$$

$$\begin{aligned}
&= z^{M+N+\mu_{MN}-\lambda_{MN}+1} \{R_1(z) \bar{S}_{\bar{m}\bar{n}}(z) - R_0(z) \bar{T}_{\bar{m}\bar{n}}(z)\} \\
&= -R_1(z) z^{M+N+\mu_{MN}-\lambda_{MN}+1} \{ \bar{T}_{\bar{m}\bar{n}}(z) R_0(z) / R_1(z) - \bar{S}_{\bar{m}\bar{n}}(z) \} \\
&= -R_1(z) z^{M+N+\mu_{MN}-\lambda_{MN}+1} O(z^{\bar{m}+\bar{n}+1}) \\
&= O(z^{M+N+\mu_{MN}-\lambda_{MN}+1+\bar{m}+\bar{n}+1}) \\
&= O(z^{m+n+1}). \quad \blacksquare
\end{aligned}$$

In order that S_{mn}/T_{mn} in (3.16) and (3.17) be a scaled Pade fraction of type (m, n) , it remains to show that $GCD(S_{mn}, T_{mn}) = z^{\lambda_{mn}}$ for some λ_{mn} . With this intent, consider again the (\bar{m}, \bar{n}) -th entry of $\Gamma_{\bar{n}}(\bar{\lambda})$, and let

$$\bar{m}^* = \bar{m} - \bar{\lambda}_{\bar{m}\bar{n}} - 1 \quad (3.18)$$

$$\bar{n}^* = \bar{n} - \bar{\lambda}_{\bar{m}\bar{n}} - 1, \quad (3.19)$$

where

$$z^{\bar{\lambda}_{\bar{m}\bar{n}}} = GCD(\bar{S}_{\bar{m}\bar{n}}, \bar{T}_{\bar{m}\bar{n}}). \quad (3.20)$$

The scaled Pade fraction $\bar{\gamma}_{\bar{m}^*, \bar{n}^*}(z)$ for $\bar{A}(z)$ satisfies

$$\begin{aligned}
\bar{\gamma}_{\bar{m}^*, \bar{n}^*}(z) &= \bar{S}_{\bar{m}^*, \bar{n}^*}(z) / \bar{T}_{\bar{m}^*, \bar{n}^*}(z) = O_E(z^{\bar{m}^* + \bar{n}^* + 1}) \\
&= O_E(z^{\bar{m} + \bar{n} - 2\bar{\lambda}_{\bar{m}\bar{n}} - 1}). \quad (3.21)
\end{aligned}$$

and, in addition,

$$\frac{\bar{S}_{\bar{m}\bar{n}}(z)}{\bar{T}_{\bar{m}\bar{n}}(z)} \neq \frac{\bar{S}_{\bar{m}^*, \bar{n}^*}(z)}{\bar{T}_{\bar{m}^*, \bar{n}^*}(z)} \quad (3.22)$$

² $R(z) = O_E(z^k)$ means that $R(z)$ is a power series whose first non-zero coefficient is the coefficient of z^k , exactly.

Lemma 3.3: Let

$$S_{m^* n^*}(z) = z^{-\lambda_{MN}} S_{MN}(z) \bar{S}_{m^* n^*}(z) - z^\alpha S_{M^* N^*}(z) \bar{T}_{m^* n^*}(z) \quad (3.23)$$

$$T_{m^* n^*}(z) = z^{-\lambda_{MN}} T_{MN}(z) \bar{S}_{m^* n^*}(z) - z^\alpha T_{M^* N^*}(z) \bar{T}_{m^* n^*}(z) \quad (3.24)$$

where $\alpha = \lambda_{MN} + \mu_{MN} + 2$. Then $\gamma_{m^* n^*}(z) = S_{m^* n^*}(z) / T_{m^* n^*}(z)$ is a Pade form of type (m^*, n^*) for $A(z)$, where

$$m^* = m - \bar{\lambda}_{m^* n^*} - 1 \quad (3.25)$$

$$n^* = n - \bar{\lambda}_{m^* n^*} - 1. \quad (3.26)$$

Proof: The proof is identical to the proof of Lemma 3.2. ■

Theorem 3.4: Let $\mu_{MN} < (n - N)$ in equation (3.4), and let

$$\gamma_{mn}(z) = S_{mn}(z) / T_{mn}(z)$$

be defined by (3.16) and (3.17). Then $\gamma_{mn}(z)$ is a scaled Pade fraction of type (m, n) for $A(z)$.

Proof: Let

$$G_{mn}(z) = \text{GCD}(S_{mn}, T_{mn}).$$

We first show that $\partial(G_{mn}) < \bar{\lambda}_{m^* n^*}$, where $\bar{\lambda}_{m^* n^*}$ is given by (3.20). Suppose that $\partial(G_{mn}) > \bar{\lambda}_{m^* n^*}$, and proceed by contradiction. Let

$$U_{m^* n^*}(z) = z^{\partial(G_{mn}) - \bar{\lambda}_{m^* n^*} - 1} S_{mn}(z) / G_{mn}(z),$$

$$V_{m^* n^*}(z) = z^{\partial(G_{mn}) - \bar{\lambda}_{m^* n^*} - 1} T_{mn}(z) / G_{mn}(z),$$

where m^* and n^* are given by (3.25) and (3.26). Then $\partial(U_{m^* n^*}) \leq m^*$, $\partial(V_{m^* n^*}) \leq n^*$, and

$$A(z) = V_{m^* n^*}(z) - U_{m^* n^*}(z)$$

$$\begin{aligned}
&= z^{\partial(G_{mn}) - \bar{\lambda}_{mn} - 1} \{A(z)T_{mn}(z) - S_{mn}(z)\} / G_{mn}(z) \\
&= O(z^{(-\bar{\lambda}_{mn} - 1) + (m+n+1)}) \\
&= O(z^{m^* + n^* + \bar{\lambda}_{mn} + 1}).
\end{aligned}$$

Thus, $U_{m^*, n^*}(z) / V_{m^*, n^*}(z)$ is a Pade form of type (m^*, n^*) for $A(z)$.

But $S_{m^*, n^*}(z) / T_{m^*, n^*}(z)$, given by Lemma 3.3, is also a Pade form of type (m^*, n^*) . Then, from the general structure of Pade forms given in equations (2.18) and (2.19) of Corollary 2.6, it follows that

$$\begin{aligned}
S_{m^*, n^*}(z) / T_{m^*, n^*}(z) &= U_{m^*, n^*}(z) / V_{m^*, n^*}(z) \\
&= S_{mn}(z) / T_{mn}(z),
\end{aligned}$$

or, equivalently, that

$$S_{m^*, n^*}(z) T_{mn}(z) - S_{mn}(z) T_{m^*, n^*}(z) = 0. \quad (3.27)$$

Replacing $S_{mn}(z) / T_{mn}(z)$ and $S_{m^*, n^*}(z) / T_{m^*, n^*}(z)$ in equation (3.27) by the expanded forms (3.16), (3.17), (3.23) and (3.24), it follows that

$$z^n z^{-\lambda_{MN}} \{S_{MN}(z) T_{m^*, n^*}(z) - S_{m^*, n^*}(z) T_{MN}(z)\} \{\bar{S}_{m^*, n^*}(z) \bar{T}_{m^*, n^*}(z) - \bar{S}_{m^*, n^*}(z) \bar{T}_{m^*, n^*}(z)\} = 0.$$

Thus, either $S_{MN}(z) / T_{MN}(z) = S_{m^*, n^*}(z) / T_{m^*, n^*}(z)$, or $\bar{S}_{m^*, n^*}(z) / \bar{T}_{m^*, n^*}(z) = \bar{S}_{m^*, n^*}(z) / \bar{T}_{m^*, n^*}(z)$, which contradicts (3.11) and (3.22).

Thus, $\partial(G_{mn}) \leq \bar{\lambda}_{mn}$. But,

$$z^{\bar{\lambda}_{mn}} = \text{GCD}(\bar{S}_{m^*, n^*}, \bar{T}_{m^*, n^*}),$$

which implies that $z^{\bar{\lambda}_{mn}}$ divides both $S_{mn}(z)$ and $T_{mn}(z)$ in equations (3.16) and (3.17). That is, $z^{\bar{\lambda}_{mn}}$ divides $G_{mn}(z)$, and consequently

$$G_{mn}(z) = z^{\lambda_{mn}}$$

Thus, $S_{mn}(z) / T_{mn}(z)$ given in Lemma 3.2 is not only a Pade form of type (m, n) for $A(z)$, but also the scaled Pade fraction of type (m, n) for $A(z)$, where

$$GCD(S_{mn}, T_{mn}) = z^{\lambda_{mn}}$$

and $\lambda_{mn} = \bar{\lambda}_{mn}$. ■

Theorem 3.5: Let $\mu_{MN} < (n - N)$ in (3.4) and let $\gamma_{mn}(z) = S_{mn}(z) / T_{mn}(z)$ be the scaled Pade fraction of type (m, n) for $A(z)$. If

$$m^* = m - \lambda_{mn} - 1 \quad (3.28)$$

and

$$n^* = n - \lambda_{mn} - 1, \quad (3.29)$$

where

$$z^{\lambda_{mn}} = GCD(S_{mn}, T_{mn}), \quad (3.30)$$

then the scaled Pade fraction of type (m^*, n^*) for $A(z)$ is $\gamma_{m^*n^*}(z) = S_{m^*n^*}(z) / T_{m^*n^*}(z)$, where $S_{m^*n^*}(z)$ and $T_{m^*n^*}(z)$ are given by equation (3.23) and (3.24).

Proof: The theorem follows using arguments identical to those of proof of Theorem 3.4, and using the results of Lemma 3.3. ■

A simple example for the off-diagonal computation is presented.

Example: Let $A(z) = 1 + z^4 + z^5 + z^9 + z^{10} + 2z^{15} + \dots$. This example constructs the scaled Pade fraction $\gamma_{7,6}(z) = S_{7,6}(z) / T_{7,6}(z)$ of type $(7, 6)$ for $A(z)$. Since $m - n = 1$, the construction proceeds along the 1st off-diagonal path of the scaled Pade table $\Gamma(A)$.

Assume that $\Gamma_3(A)$ is already available, from which it can be determined that the scaled Pade fraction $\gamma_{4,3}(z)$ of type $(M, N) = (4, 3)$ is given by

$$\gamma_{4,3}(z) = \frac{S_{4,3}(z)}{T_{4,3}(z)} = \frac{1 - z + z^2 - z^3 + z^4}{1 - z + z^2 - z^3} \quad (3.31)$$

From (3.4), the residual for $S_{4,3}(z) / T_{4,3}(z)$ is given by

$$A(z) T_{4,3}(z) - S_{4,3}(z) = R_1(z) z^{4+3+1}, \quad (3.32)$$

where $R_1(z) = -1 + z - z^2 + 2z^7 + \dots$

Consequently, equation (3.32) yields that $\mu_{4,3} = 0$. Since $\mu_{4,3} < n - N$, Theorem 3.4 is applicable. Observe that

$$z^{\lambda_{MN}} = z^0 = \text{GCD}(S_{4,3}, T_{4,3}),$$

and consequently the predecessor of $\gamma_{4,3}(z)$ along the 1st off-diagonal path is $\gamma_{M \cdot N}(z) = \gamma_{3,2}(z)$.

Therefore, $\gamma_{3,2}(z)$ is contained in $\Gamma_3(A)$, and is found to be

$$\gamma_{3,2}(z) = \frac{S_{3,2}(z)}{T_{3,2}(z)} = \frac{z^2}{z^2}$$

The residual for $S_{3,2}(z) / T_{3,2}(z)$ is given from

$$A(z) T_{3,2}(z) - S_{3,2}(z) = R_0(z) z^{3+2+1}, \quad (3.33)$$

where $R_0(z) = 1 + z + z^5 + z^6 + \dots$

The two residuals $R_1(z)$ and $R_0(z)$ give the residual power series $\bar{A}(z) \in \mathcal{U}$, where

$$\begin{aligned} \bar{A}(z) &= R_0(z) / R_1(z) \\ &= -1 - 2z - 2z^2 - 2z^3 - \dots \end{aligned} \quad (3.34)$$

Using equation (3.13) and (3.14),

$$\bar{m} = n - N + \lambda_{MN} = 3$$

$$\bar{n} = n - N - \mu_{MN} - 1 = 2,$$

and in order to apply Theorem 3.4, it is therefore required to obtain the scaled Padé fraction

$\bar{\gamma}_{3,2}(z)$ of type (3,2) and its predecessor for $\bar{A}(z)$. Assuming that $\Gamma_3(\bar{A})$ is available, from it can be

determined that

$$\bar{\gamma}_{3,2}(z) = \frac{\bar{S}_{3,2}(z)}{\bar{T}_{3,2}(z)} = \frac{-z - z^2}{-z - z^2} \quad (3.35)$$

Similarly, the predecessor of $\bar{\gamma}_{3,2}(z)$ along the first off-diagonal path of $\Gamma_3(\bar{A})$ is given by

$$\bar{\gamma}_{1,0}(z) = \bar{S}_{1,0}(z) / \bar{T}_{1,0}(z) = (-1 - 2z)/1.$$

Now by applying the formulae (3.16) and (3.17), the (7,6) entry of $\Gamma(\bar{A})$ is computed by

$$S_{7,6}(z) = (1 - z + z^2 - z^3 + z^4) \cdot (-z - z^2) - z^\alpha (z^2) \cdot (z - z^2),$$

$$T_{7,6}(z) = (1 - z + z^2 - z^3) \cdot (-z - z^2) - z^\alpha (z^2) \cdot (z - z^2),$$

where $\alpha = \lambda_{4,3} + \mu_{4,3} + 2 = 2$. Thus,

$$\gamma_{7,6}(z) = S_{7,6}(z) / T_{7,6}(z) = z(-1 - z^4) / z(-1 + z^5). \quad (3.36)$$

Note that $\bar{\gamma}_{1,0}(z)$ is not used for computing $\gamma_{7,6}(z)$. It is used instead in formulae (3.23) and (3.24)

for computing the predecessor of $\gamma_{7,6}(z)$. Since

$$z^{\lambda_{mn}} = z^{\lambda_{m,n}} = z^1 = \text{GCD}(\bar{S}_{3,2}, \bar{T}_{3,2}),$$

it is known that the predecessor is $\gamma_{5,4}(z)$, which is determined by (3.23) and (3.24) to be

$$\gamma_{5,4}(z) = \frac{S_{5,4}(z)}{T_{5,4}(z)} = \frac{(-1 - z + z^2 - z^3 - 2z^5)}{(-1 - z + z^2 - z^3 + z^4)}$$

3.3. Fast Off-diagonal Algorithm for a Single Power Series

3.3.1. The Algorithm

The algorithm given in this section constructs the scaled Pade fraction $\gamma_{m,n}$ of type (m,n) for $A(z)$ in a quadratic fashion. The iteration assumes the existence of

$$\gamma_{MN}(z) = S_{MN}(z) / T_{MN}(z), \quad (3.37)$$

where $M = N + (m - n)$, and where for notational convenience we set

$$S_1 = S_{MN}(z) \quad (3.38)$$

and

$$T_1 = T_{MN}(z) \quad (3.39)$$

Also assumed to exist is the predecessor

$$\gamma_{M^*N^*}(z) = S_{M^*N^*}(z) / T_{M^*N^*}(z) \quad (3.40)$$

of $\gamma_{MN}(z)$ on the the $(m-n)$ -th off-diagonal path of $\Gamma(A)$, where

$$M^* = M - \lambda_{MN} - 1,$$

$$N^* = N - \lambda_{MN} - 1$$

and

$$z^{\lambda_{MN}} = \text{GCD}(S_{MN}, T_{MN}).$$

Again for notational convenience, we set

$$S_0 = S_{M^*N^*}(z) \quad (3.41)$$

and

$$T_0 = T_{M^*N^*}(z) \quad (3.42)$$

To advance the solution from N to $N+s$ (that is, to construct $\gamma_{M+N+s, N+s}(z)$), where s is the step size, the algorithm first computes μ_{MN} such that

$$A(z)T_1 - S_1 \bmod z^{M+N+2s+\lambda_{MN}+1} = z^{M+N+\mu_{MN}+1} R_1(z),$$

where

$$z^{\lambda_{MN}} = \text{GCD}(S_1, T_1)$$

and $R_1(0) \neq 0$ if $\mu_{MN} < 2s + \lambda_{MN}$.

If $\mu_{MN} \geq s$, then $\gamma_{M+N+s, N+s}(z)$ is constructed trivially by means of Theorem 3.1. Otherwise,

Theorems 3.4 and 3.5 are applied.

ALGORITHM 1: OFFDIAG

INPUT: A, m, n , where

- (1) m and n are non-negative integers with $m \geq n$, and
- (2) A is a unit power series. (Note that only $A \bmod z^{m+n+1}$ is required)

OUTPUT: $\begin{bmatrix} S_1 & S_0 \\ T_1 & T_0 \end{bmatrix}$, where

- (1) S_1 / T_1 is the scaled Padé fraction of type (m, n) for A , and
- (2) S_0 / T_0 is the scaled Padé fraction of type $(m - \lambda_{mn} - 1, n - \lambda_{mn} - 1)$ for A , given that

$$z^{\lambda_{mn}} = \text{GCD}(S_1, T_1)$$

Step 1: # Initialization #

$$\begin{aligned} i &= -1 \\ M &= (m - n) \\ N &= 0 \end{aligned}$$

$$\begin{bmatrix} S_1 & S_0 \\ T_1 & T_0 \end{bmatrix} = \begin{bmatrix} A(z) \bmod z^{M+1} & 1 \\ 1 & 0 \end{bmatrix}$$

Step 2: # Calculation of step size #

$$\begin{aligned} i &= i + 1 \\ s &= \min \{2^i, N, n - 1\} \end{aligned}$$

Step 3: # Termination condition #

If $s = 0$ then exit

Step 4: # Calculation of scaling factor for S_1 / T_1 #

Determine λ_{MN} such that $z^{\lambda_{MN}} = \text{GCD}(S_1, T_1)$

Step 5: # Computation of residual for $\gamma_{MN} = S_1 / T_1$ #

Compute μ_{MN} and R_1 such that

$$(A T_1 - S_1) \bmod z^{M+N+2s+\lambda_{MN}+1} = z^{M+N+\mu_{MN}+1} R_1,$$

where $R_1(0) \neq 0$ if $\mu_{MN} < 2s + \lambda_{MN}$

Step 6: # Identification of Cases #

if $\mu_{MN} \geq s$

then # Case of Theorem 3.1 #

$$\begin{bmatrix} S_1 & \hat{S}_0 \\ T_1 & \hat{T}_0 \end{bmatrix} = \begin{bmatrix} S_1 & \hat{S}_0 \\ T_1 & \hat{T}_0 \end{bmatrix} \begin{bmatrix} z^s & 0 \\ 0 & 1 \end{bmatrix}$$

go to step 1)

else # Case of Theorem 3.2 #

go to step 2)

Step 7: # Calculation of the residue γ_{MN} and the polynomial \bar{p} #

$$\bar{p} = \frac{z^{s+1} \lambda_{MN}}{z^{s+1} \lambda_{MN} + 1}$$

Step 8: # Compute Series of γ_{MN} and \bar{p} #

$$(A(z) \cdot T_0 - S_0) \bmod \frac{z^{N+M} - \bar{z}^{N+M} - 1}{z^{N+M} - \bar{z}^{N+M} - 1} \Gamma_{N+M}$$

where $R_0(0) \neq 0$.

Step 9: # Computation of residual power series //

$$\bar{A}(z) = R_0(z) - R_0(0) \pmod{z^{\bar{N}+1}}$$

Step 10: # Computation of residual power series //

$$\begin{bmatrix} \bar{S}_1 & \bar{S}_0 \\ \bar{T}_1 & \bar{T}_0 \end{bmatrix} = \text{COMPUTATION}$$

Step 11: # Computation of residual power series //

$$\begin{bmatrix} \bar{S}_1 & \bar{S}_0 \\ \bar{T}_1 & \bar{T}_0 \end{bmatrix} = \begin{bmatrix} \bar{S}_1 & \bar{S}_0 \\ \bar{T}_1 & \bar{T}_0 \end{bmatrix}$$

Step 12: #

g'

$$\{y_i^{(j)}\}_{i=1}^n$$

Step 13: #

Step 14: #

Step 15: #

Also, define

$$\Omega_N = \{\Gamma_N(A) \text{ for all } A \in U\} \quad (3.44)$$

for $N = 0, 1, \dots$

The proof proceeds by induction on Ω_N for $i = 0, 1, 2, \dots$. Clearly the algorithm is correct for Ω_0 .

It is shown that if the algorithm is correct for Ω_N , then it is also correct for Ω_{N+1} , where

$$\Omega_{N+1} = \begin{cases} \Omega_0 & \text{if } N = 0 \\ \Omega_N & \text{if } N \geq 1 \end{cases} \quad (3.45)$$

Let $A(z) \in U$ be an arbitrary unit power series. It is shown that the algorithm correctly computes $\gamma_n^{(N)}(z)$, where $m \geq n$ and $1 \leq n \leq N+1$, given that it is correct for Ω_N .

If $1 \leq n \leq N$, then $\gamma_n^{(N)}(z) \in U$, $A(z) \in \Omega_N$, and by the inductive hypothesis the algorithm computes it correctly.

If $1 \leq n \leq N+1$, let $M = L+1$ ($m = n$). By the inductive hypothesis, the algorithm (using eq. (3.45)) correctly computes $\gamma_n^{(N)}(z) = S_N^{(N)}(z) + T_N^{(N)}(z) \in \Omega_N$, and its predecessor

$\gamma_{n-1}^{(N)}(z) \in U$. The algorithm then computes $\gamma_n^{(N+1)}(z)$ and its predecessor

(3.46)

by eq. (3.45) and let $\gamma_n^{(N+1)}(z) \in \Omega_{N+1}$. The algorithm then computes

yielding $\gamma_n^{(N+1)}(z) \in U$. The algorithm then computes

$$\gamma_n^{(N+1)}(z) = \gamma_n^{(N)}(z) + \gamma_{n-1}^{(N+1)}(z) \in \Omega_{N+1} \quad (3.46)$$

If $\mu_{MN} \geq s$, then by Theorem 3.1, step 5 correctly yields $\gamma_{mn}^{(A)}(z)$. Furthermore, from equations (3.6) and (3.7),

$$\begin{aligned} z^{\lambda_{mn}} &= \text{GCD}(S_{mn}^{(A)}(z), \tau_{mn}^{(A)}(z)) \\ &= z^{\lambda_{mn}^{(A)}}. \end{aligned}$$

Consequently, the predecessor of $\gamma_{mn}^{(A)}(z)$ is given by

$$\gamma_{mn}^{(A)}(z) = \gamma_{mn}^{(1)}(z) \quad (3.15)$$

since

$$m' = m - \lambda_{mn} = m - \lambda_{mn}^{(A)} = m - (1 + M)$$

and

$$n' = n - \lambda_{mn} = n - \lambda_{mn}^{(A)} = n - (1 + M).$$

Thus, in step 8, the output of step 5 is

$$R_1(z) = \gamma_{mn}^{(A)}(z) = 1 + z^{\lambda_{mn}^{(A)}}.$$

and

$$\bar{m}' = m' + \lambda_{mn} \geq \bar{m}.$$

The polynomial $R_1(z)$, computed in step 5, therefore satisfies $R_1(0) \neq 0$ and is of degree $\bar{m}' + \bar{n}$. In addition, step 8 produces a polynomial $R_0(z)$ of degree $\bar{m}' + \bar{n}$ such that $R_0(0) \neq 0$. Consequently, enough terms are available in $R_1(z)$ and $R_0(z)$ to compute, in step 9, the residual unit power series $\bar{A}(z)$ mod $z^{\bar{m}' + \bar{n} + 1}$ defined by (3.14). By the induction hypothesis, step 10 therefore correctly computes $\gamma_{\bar{m}' + \bar{n}}^{(A)}(z) \in \Gamma_{\bar{m}' + \bar{n}}(\bar{A}) \in \Omega_{\bar{m}' + \bar{n}}$ and step 11 produces $\gamma_{\bar{m}' + \bar{n}}^{(A)}(z)$ as the correct predecessor of $\gamma_{mn}^{(A)}(z)$. This completes the proof of the correctness of the algorithm.

3.3.3. Cost Analysis

Let $C(m, n)$ be the cost of computing the scaled Pade fraction of type (m, n) and its predecessor for an arbitrary power series $A(z) \in U$ using Algorithm 1. For the sake of simplicity, assume $0 \leq n \leq m \leq 2n$. The case that $m > 2n$ is considered later. In this section, asymptotic estimates of $C(m, n)$ are derived by counting the number of operations (additions, subtractions, multiplications and divisions in F) performed by the algorithm. A detailed cost analysis and an implementation of the algorithm are described by Verheijen [VER83], who compares Algorithm 1 with other algorithms for calculating Pade fractions.

When obtaining the asymptotic cost estimates, it is assumed that the algorithm makes use of fast methods for polynomial and power series arithmetic. Using fast Fourier transforms, two polynomials with coefficients in the field F and of degree M and N , respectively, can be multiplied in $O((M + N) \log(M + N))$ operations in F . Using Newton's method and fast multiplication of polynomials, the first N terms of the quotient of two power series in U can be obtained in $O(N \log N)$ operations in F . These and other fast methods for polynomial and power series arithmetic are described, for example, in Aho, Hopcraft and Ullman [AHO74] and in Lipson [LIP91].

Let $k = \lceil \log n \rceil$. Then it is easy to verify that the algorithm terminates after k iterations and that the number of operations of step 2 of iteration i

$$= \begin{cases} 1 & , i = 0 \\ 2^{i-1} & , 0 < i < k \\ n - 2^i & , i = k \end{cases} \quad (3.48)$$

and the number of operations of step 3

$$N = \begin{cases} 0 & , i = 0 \\ 2^{i-1} & , \text{otherwise,} \end{cases} \quad (3.49)$$

$$M = N + (m-n). \quad (3.50)$$

Assume that during iteration i , the scaled Padé fraction S_1/T_1 of type (M, N) and its predecessor S_0/T_0 are available.

The first nontrivial step requires the computation of the residual $R_1(z)$ in step 5. Let

$$A_1(z) = \sum_{j=0}^M a_j z^j \quad (3.51)$$

and

$$A_2(z) = \sum_{j=0}^{N+2r-1} a_{M+j+1} z^j \quad (3.52)$$

With λ_{MN} given by step 4, it is known that

$$\begin{aligned} A(z) T_1 - S_1 &\text{ mod } z^{M+N+2r+\lambda_{MN}+1} \\ &= A_1(z) T_1 + z^{M+1} A_2(z) T_1 - S_1 \text{ mod } z^{M+N+2r+\lambda_{MN}+1} \\ &= O(z^{M+N+1}). \end{aligned} \quad (3.53)$$

Since $A_1 T_1$ and S_1 are both of degree at most $M+N$, then μ_{MN} and $R_1(z)$ can be obtained directly from $z^{M+1} A_2 T_1$. The product $A_2 T_1$ is a polynomial of at most degree $2N + 2r - 1 < 4N$. Thus, using fast polynomial multiplication, step 5 can be executed in $O(N \log N)$ operations.

If it is determined as a result of step 5 that $\mu_{MN} \geq r$, then step 6 is performed trivially and the iteration is complete. Otherwise, the algorithm continues in step 8 with the calculation of the residual R_0 of the predecessor scaled Padé fraction S_0/T_0 . Making the same observations as in step 4, it follows that R_0 can be obtained from the product of $A_0 T_0$ where

$$A_0 = \sum_{j=0}^{N+\bar{m}+\bar{n}-\lambda_{MN}} a_{M-\lambda_{MN}+j+1} z^j \quad (3.54)$$

and T_0 is a polynomial of degree at most $N - \lambda_{MN} - 1$. Since the degree of the product is

bounded by $2N + \bar{m} + \bar{n} - 2\lambda_{MN} - 1 < 4N$, step 8 can be executed in $O(N \log N)$ operations.

The computation of the residual power series $\bar{A}(z) \bmod x^{\bar{m} + \bar{n} + 1}$ in step 9 requires the computation of the first $\bar{m} + \bar{n} = 2s + \lambda_{MN} - \mu_{MN} + 1 \leq 3N$ terms of the quotient of R_0 / R_1 .

Again, this may be performed in $O(N \log N)$ operations.

In step 8, the recursive call of Algorithm 1 in order to compute the scaled Pade fraction of type (\bar{m}, \bar{n}) for $A(z)$, requires $C(\bar{m}, \bar{n})$ operations by assumption. For later purposes, it is important to observe that $0 \leq \bar{n} \leq N$ and that $\bar{n} \leq \bar{m} \leq 2N$.

The final non-trivial step requires eight polynomial multiplications to obtain the scaled Pade fraction of type $(M+s, N+s)$ and its predecessor for $A(z)$. Since $M \leq 2N$, each of the polynomial products are of degree at most $M+s < 3N$. Consequently, step 8 can be executed in $O(N \log N)$ operations.

It is an easy matter to show that

$$C(m_1, n_1) \leq C(m_2, n_2), \quad (3.55)$$

whenever $m_1 \leq m_2$ and $n_1 \leq n_2$. The total cost of the i -th iteration is then bounded by

$$\begin{aligned} & C(\bar{m}, \bar{n}) + c(2N) \log(2N) \\ & \leq C(2N, N) + c(2N) \log(2N) \\ & \leq C(2^i, 2^{i-1}) + ci2^i \end{aligned} \quad (3.56)$$

operations, for an appropriate constant c . Consequently, we have

Theorem 3.6. Given that $0 \leq n \leq m \leq 2n$, Algorithm 1 can compute the scaled Pade fraction of type (m, n) for $A(z) \in U$ in time $O(n \log^2 n)$.

Proof: Consider the recurrence relation

$$C(2^{k+1}, 2^k) = \sum_{i=1}^k [C(2^i, 2^{i-1}) + ci2^i], \quad k \geq 1,$$

where c is a positive constant. Then

$$\begin{aligned} C(2^{k+1}, 2^k) &= \sum_{i=1}^k [C(2^i, 2^{i-1}) + ci2^i] + C(2^k, 2^{k-1}) + ck2^k \\ &= 2C(2^k, 2^{k-1}) + ck2^k. \end{aligned}$$

With $n = 2^k$, results on recurrence relations (see Bentley, Hanken and Saxe [BEN80]) then yield

$$\begin{aligned} C(2n, n) &= n[C(2, 1) + c \sum_{i=1}^k i] \\ &= n[C(2, 1) + ck(k+1)/2] \\ &= O(n \log^2 n). \end{aligned}$$

The theorem now follows, since from equation (3.56) for m and n satisfying $0 \leq n \leq m \leq 2n$

$$C(m, n) \leq \sum_{i=1}^k [C(2^i, 2^{i-1}) + ci2^i]. \quad \blacksquare$$

Lemma 3.7: Let $m > n$, and let $A(z) \in U$. Determine an integer $\delta \geq 1$ such that

$$A(z) = A_1(z) + z^{m-n+\delta} A_2(z), \quad (3.57)$$

where

$$A_1(z) = A(z) \bmod z^{m-n+1} \quad (3.58)$$

and $A_2(0) \neq 0$ if $\delta < \infty$. If $\delta < n$, let $S_{n, n-\delta}(z) / T_{n, n-\delta}(z)$ be the scaled Padé fraction of type $(n, n-\delta)$ for $1/A_2(z)$. Then the scaled Padé fraction $S_{mn}(z) / T_{mn}(z)$ of type (m, n) for $A(z)$ is given by

$$S_{mn}(z) = \begin{cases} A_1(z)S_{n,n-\delta}(z) + z^{m-n+\delta}T_{n,n-\delta}(z), & \text{if } \delta \leq n \\ A_1(z)z^n, & \text{otherwise,} \end{cases} \quad (3.59)$$

$$T_{mn}(z) = \begin{cases} S_{n,n-\delta}(z), & \text{if } \delta \leq n \\ z^n, & \text{otherwise.} \end{cases} \quad (3.60)$$

Proof: If $\delta \leq n$, then,

$$\begin{aligned} \partial(S_{mn}) &= \partial(A_1S_{n,n-\delta} + z^{m-n+\delta}T_{n,n-\delta}) \\ &\leq \max\{(m-n) + n, (m-n+\delta) + (n-\delta)\} \\ &= m. \end{aligned}$$

Also, $\partial(T_{mn}) \leq n$. Furthermore, either $\partial(S_{mn}) = m$ and $\partial(T_{mn}) = n$ or both. Moreover,

$$\begin{aligned} A(z) \cdot T_{mn}(z) - S_{mn}(z) &= \{A_1(z) + z^{m-n+\delta}A_2(z)\}S_{n,n-\delta}(z) - \{A_1(z)S_{n,n-\delta}(z) + z^{m-n+\delta}T_{n,n-\delta}(z)\} \\ &= z^{m-n+\delta}\{A_2(z)S_{n,n-\delta}(z) - T_{n,n-\delta}(z)\} \\ &= z^{m-n+\delta}\{A_2(z)O(z^{2n-\delta+1})\} \\ &= O(z^{m+n+1}). \end{aligned}$$

Finally,

$$\begin{aligned} \text{GCD}(S_{mn}, T_{mn}) &= \text{GCD}(A_1S_{n,n-\delta} + z^{m-n+\delta}T_{n,n-\delta}, S_{n,n-\delta}) \\ &= \text{GCD}(z^{m-n+\delta}T_{n,n-\delta}, S_{n,n-\delta}) \\ &= \text{GCD}(T_{n,n-\delta}, S_{n,n-\delta}). \end{aligned}$$

If $\delta > n$, then clearly $\partial(S_{mn}) \leq m$ and $\partial(T_{mn}) = n$. Moreover,

$$\begin{aligned} A(z) \cdot T_{mn}(z) - S_{mn}(z) &= z^n\{A_1(z) + z^{m-n+\delta}A_2(z)\} - z^nA_1(z) \\ &= z^{m+\delta}A_2(z) \\ &= O(z^{m+n+1}). \end{aligned}$$

Finally, $\text{GCD}(S_{mn}, T_{mn}) = z^n$. ■

Theorem 3.8: For arbitrary $m \geq n$, Algorithm 1 can compute the scaled Pade fraction of type (m, n) for $A(z) \in U$ in time $O(m \log m) + O(n \log^2 n)$.

Proof: If $\delta > n$ in equation (3.57), the result is trivial.

If $\delta \leq n$, computation of the first $2n - \delta + 1$ terms of $1/A_2(z)$ requires $O(n \log n)$ operations. Using (3.55),

$$C(n, n-\delta) \leq C(n, n),$$

and consequently by Theorem 3.6, it follows that the cost of computing the scaled Pade fraction $S_{n, n-\delta}(z)/T_{n, n-\delta}(z)$ for $1/A_2(z)$ is bounded by $O(n \log^2 n)$ operations. Finally, the cost of computing $S_{mn}(z)$ in equation (3.59) is $O(m \log m)$ operations. ■

3.4. Fast Off-diagonal Algorithm for a Quotient Power Series

Let $A(z), B(z)$ be unit power series and let

$$C(z) = -A(z)/B(z) \quad (3.59)$$

be the quotient power series. Given the non-negative integers $m \geq n$, then Algorithm 1 OFF-DIAG can be used to compute the scaled Pade fractions $S_{mn}(z)/T_{mn}(z)$ of type (m, n) for $C(z)$. As a result,

$$A(z)T_{mn}(z) + B(z)S_{mn}(z) = O(z^{m+n+1}) \quad (3.60)$$

Before applying the algorithm, the quotient

$$C(z) \bmod z^{m+n+1} = -A(z)/B(z) \bmod z^{m+n+1} \quad (3.61)$$

must be calculated. However, this division need be computed modulo z^{m+n+1} , only, by modifying

Algorithm 1 as follows:

ALGORITHM 2: OFFDIAG

INPUT: $A(z), B(z), m, n$, where

- (1) m and n are non-negative integers with $m \geq n$, and
- (2) $A(z), B(z)$ are unit power series. (Note that only $A(z) \bmod z^{m+n+1}$ and $B(z) \bmod z^{m+n+1}$ are required).

OUTPUT: $\begin{bmatrix} S_1 & S_0 \\ T_1 & T_0 \end{bmatrix}$, where

- (1) S_1 / T_1 is the scaled Pade fraction of type (m, n) for $-A(z) / B(z)$, and
- (2) S_0 / T_0 is the predecessor scaled Pade fraction of type $(m - \lambda_{mn} - 1, n - \lambda_{mn} - 1)$ for $-A(z) / B(z)$, given that

$$z^{\lambda_{mn}} = \text{GCD}(S_1, T_1).$$

Step 1: # Initialization #

$$\begin{aligned} i &\leftarrow -1 \\ M &\leftarrow (m - n) \\ N &\leftarrow 0 \end{aligned}$$

$$\begin{bmatrix} S_1 & S_0 \\ T_1 & T_0 \end{bmatrix} \leftarrow \begin{bmatrix} -A(z) / B(z) \pmod{z^{M+1}} & z^{M+1} \\ 1 & 0 \end{bmatrix}$$

Step 2: # Calculation of step-size #

$$\begin{aligned} i &\leftarrow i + 1 \\ s &\leftarrow \min \{2^i - N, n - N\} \end{aligned}$$

Step 3: # Termination criterion #

If $s = 0$ then exit

Step 4: # Calculation of scaling factor for S_1 / T_1 #

Determine λ_{MN} such that

$$z^{\lambda_{MN}} = \text{GCD}(S_1, T_1)$$

Step 5: # Computation of residual for $\gamma_{MN}(z) = S_1 / T_1$ #

Compute μ_{MN} and R_1 such that

$$(AT_1 + BS_1) \bmod z^{M+N+2s+\lambda_{MN}-1} = z^{M+N+\mu_{MN}-1} R_1$$

where $R_1(0) \neq 0$ if $\mu_{MN} < 2s + \lambda_{MN}$.

Step 6: # Identification of Cases #

if $\mu_{MN} \geq s$

then # Case of Theorem 3.1. #

$$\begin{bmatrix} S_1 & S_0 \\ T_1 & T_0 \end{bmatrix} = \begin{bmatrix} S_1 & S_0 \\ T_1 & T_0 \end{bmatrix} \begin{bmatrix} z' & 0 \\ 0 & 1 \end{bmatrix}$$

go to step 11

else # Case of Theorem 3.4 and 3.5 #

go to step 7

Step 7: # Calculation of degrees for residual scaled Padé fractions #

$$\begin{aligned} \bar{m} &= s + \lambda_{MN} \\ \bar{n} &= s - \mu_{MN} - 1 \end{aligned}$$

Step 8: # Computation of residual for $\gamma_{M,N}(z) = S_0 / T_0$ #

Compute R_0 such that

$$(A \cdot T_0 + B \cdot S_0) \bmod z^{M+N+\bar{m}+\bar{n}-2\lambda_{MN}} = R_0 z^{M+N-2\lambda_{MN}-1}$$

where $R_0(0) \neq 0$.

Step 9: # Computation of residual scaled Pade fractions #

$$\begin{bmatrix} \bar{S}_1 & \bar{S}_0 \\ \bar{T}_1 & \bar{T}_0 \end{bmatrix} = \text{OFFDIAG} (R_0, R_1, \bar{m}, \bar{n})$$

Step 10: # Advancement of scaled Pade fraction computation #

$$\begin{bmatrix} \bar{S}_1 & \bar{S}_0 \\ \bar{T}_1 & \bar{T}_0 \end{bmatrix} = \begin{bmatrix} \bar{S}_1 & \bar{S}_0 \\ \bar{T}_1 & \bar{T}_0 \end{bmatrix} \begin{bmatrix} z^{-\lambda_{MN}} & 0 \\ 0 & z^{\lambda_{MN} + \lambda_{MN} + 2} \end{bmatrix} \begin{bmatrix} \bar{S}_1 & \bar{S}_0 \\ \bar{T}_1 & \bar{T}_0 \end{bmatrix}$$

Step 11: # Calculation of degrees of S_1 / T_1 #

$M \leftarrow M + s$
 $N \leftarrow N + s$
 goto step 2

Algorithm 2 is a generalization of Algorithm 1, since it can be used to produce scaled Pade fractions for a single power series $A(z)$ simply by setting $B(z) = -1$. It differs from Algorithm 1 in that the division

$$\bar{A}(z) = -R_0(z) / R_1(z), \bmod z^{\bar{m} + \bar{n} + 1} \quad (3.63)$$

in step 9 of Algorithm 1 is avoided. Instead, the division is delayed (i.e., immediately subsequent to, rather than prior to, the recursive call of OFFDIAG) until the initialization

$$S_1 = -A(z) / B(z) \bmod z^{n-m+1} \quad (3.64)$$

in step 1 of Algorithm 2.

There are also various sign changes introduced in steps 1, 5, 8 and 10. These account for the fact that the algorithm deals with the power series $-A(z) / B(z)$ rather than $A(z) / B(z)$. This notational change simplifies the development of subsequent results.

For implementation purposes, Algorithm 2 can result in considerable savings in cost. Practically, fast division is significantly slower than fast multiplication, by an asymptotic constant of approximately 7 (see Verheijen [VER83]). For example, if $-A(z) / B(z)$ is normal, then $\bar{m} = \bar{n} + 1 = 2$ and the division in step 1 of Algorithm 2 becomes trivial. However, the asymptotic cost of Algorithm 2 remains the same as the asymptotic cost of Algorithm 1 given in section 3.3.

The proof of the correctness of Algorithm 2 is nearly identical to the proof of correctness of Algorithm 1, and therefore it is not given.

3.5. Classical Off-diagonal Algorithm for a Quotient Power Series

Let $S_k = T_k$ be the k th Padé fraction of type (M, N) for $k = 0, 1, 2, \dots$

$$T_k = \frac{A_k(z)}{B_k(z)} \quad (3.65)$$

that is, $A_k(z) = 0$. Then

$$A_k(z) = T_k(z) - \frac{A_{k-1}(z)}{B_{k-1}(z)}$$

where $R_k(0) \neq 0$ if $\mu_{M,N} < k$. Thus, $(z^k - T_k) = (z^k - 1) \dots$ the k th order function of

$(M+k, N+k)$ for $A(z)/B(z)$ the series $0 \leq j \leq \mu_{M,N}$. If $\mu_{M,N} = \infty$, then the series

converges to $A(z)/B(z)$ as $k \rightarrow \infty$. The series $A_k(z)/B_k(z)$ converges to $A(z)/B(z)$ as $k \rightarrow \infty$.

$$s = \mu_{mn} + 1.$$

By so selecting the step-size s , Algorithm 2 may be used to compute all Pade fractions along the $(M+N)$ -th off-diagonal path. With this choice of s , step 7 in Algorithm 2 gives

$$\bar{m} = \mu_{MN} + 1$$

$$\bar{n} = 0$$

Consequently, the recursive call of OFFDIAG in step 10 reduces to

$$\begin{bmatrix} \bar{S}_1 & \bar{S}_0 \\ \bar{T}_1 & \bar{T}_0 \end{bmatrix} = \begin{bmatrix} P_n \bar{T}_0 - z^{\mu_{MN}+1} \bar{S}_0 & z^{\mu_{MN}} \bar{S}_1 \\ 1 & 0 \end{bmatrix},$$

and step 11 then yields

$$\begin{bmatrix} S_1 & S_0 \\ T_1 & T_0 \end{bmatrix} = \begin{bmatrix} S_1 \bar{S}_1 + z^{\mu_{MN}+1} S_0 & z^{\mu_{MN}} S_1 \\ T_1 \bar{S}_1 + z^{\mu_{MN}+1} T_0 & z^{\mu_{MN}} T_1 \end{bmatrix}$$

Since $\text{GCD}(\bar{S}_1, \bar{T}_1) = 1$, it follows from the proof of theorem 3.4 that

$$\text{GCD}(S_1 \bar{S}_1 + z^{\mu_{MN}+1} S_0, T_1 \bar{S}_1 + z^{\mu_{MN}+1} T_0) = 1.$$

Thus, the above computations can be repeated for the scaled Lode fraction of type $(M+s, N+s)$.

The full details are provided in Algorithm 3, below. To simplify the presentation of subsequent results, in the i -th iteration, the scaled Lode fraction S_i/T_i of type (M, N) is denoted by S_i and its denominator T_i by T_i .

ALGORITHM 3

(1) m and n are negative integers with $m > n$ and

(2) $4 \leq m+1 \leq n$ with m even and n odd, $m \neq n$ and $m \neq n-1$.

$$C_{m,n} = \begin{bmatrix} S & S \\ 0 & 0 \end{bmatrix} \quad \text{for}$$

(1) $S = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ for $m \neq n-1$ and $n \neq 1$ and

(2) $S = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$ for $n = 1$.

$$M = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

1

go to step 2 ■

In the case that $B(z) = -1$ and $m = n$, Algorithm 3 is precisely the algorithm given by Cabay and Kao [CAB83] for computing diagonal Pade fractions for a single power series. Their $O(n^2)$ algorithm (since steps 3 and 6 require only $\mu_{MN} + 2$ terms of a product polynomials, and since one of the multipliers in step 7 is a polynomial of degree $\mu_{MN} + 1$, for small μ_{MN} , no advantage can be gained from using fast methods for polynomial arithmetic) is shown to be faster than other $O(n^2)$ algorithms for computing diagonal Pade fractions, such as those of Trench [TRE65] and Rissanen [RIS73].

The more interesting observation about Algorithm 3 is made in the next chapter, however. It is shown that, when the given quotient power series is a rational function, Algorithm 3 along one specific off diagonal path corresponds exactly to Euclid's extended algorithm for computing the greatest common divisor of the numerator and denominator of the given rational function. In this sense, Algorithm 3 is a generalization of Euclid's extended algorithm. For this reason that we choose to call it "Algorithm 3" instead of "Algorithm 1".

CHAPTER 4

GREATEST COMMON DIVISOR COMPUTATIONS OF POLYNOMIALS

4.1. Introduction

In this chapter, Algorithm 2 and Algorithm 3 are examined when they are applied to $A(z)/B(z)$, where $A(z)$ and $B(z)$ are polynomials of degrees m and n respectively. In addition, it is assumed that $A(0) \neq 0$ and $B(0) \neq 0$.

To permit the analysis, we need the following

Definition: The reciprocal of a polynomial

$$P(z) = p_0 + p_1 z + \dots + p_n z^n \quad (4.1)$$

of at most degree n is defined to be

$$P^*(z) = p_0 z^n + p_1 z^{n-1} + \dots + p_n = z^n P(z^{-1}). \quad (4.2)$$

The name originates from the fact that the zeros of $P^*(z)$ are the reciprocals of those of $P(z)$ if $p_0 \neq 0$.

Clearly

$$[P^*(z)]^* = P(z); \quad (4.3)$$

that is, the operation of forming the reciprocal polynomial is involuntary. Also

$$[P(z) \cdot Q(z)]^* = P^*(z) \cdot Q^*(z), \quad (4.4)$$

and

$$[z^n P(z)]^* = P^*(z) \quad (4.5)$$

Let $A(z)$ be a unit power series. The first $n+1$ terms of $A(z)$ are denoted by

$$\begin{aligned} A_n(z) &= \sum_{i=0}^n a_i z^i \\ &= A(z) \bmod z^{n+1} \end{aligned} \quad (4.6)$$

The reciprocal $A_n^R(z)$ of $A_n(z)$ is then given by

$$A_n^R(z) = \sum_{i=0}^n a_{n-i} z^i. \quad (4.7)$$

4.2. Euclid's Extended Algorithm

Theorem 4.1. Algorithm 3 applied to $-A(z)/B(z)$, where $A(z)$ and $B(z)$ are polynomials of degree m and n respectively, is equivalent to Euclid's extended algorithm for computing the greatest common divisor of $A^R(z)$ and $B^R(z)$

Proof: For completeness in presentation, assume that modulo operations are not performed in steps 3 and 6 of Algorithm 3. Thus,

$$A \cdot T_i + B \cdot S_i = z^{M \cdot N \cdot i} R_i \quad (4.8)$$

$$A \cdot T_{i-1} + B \cdot S_{i-1} = z^{M \cdot N \cdot (i-1)} R_{i-1} \quad (4.9)$$

where $R_i(z)$ and $R_{i-1}(z)$ are polynomials of degrees $\leq N - i$ and $\leq N - (i-1)$ respectively. By taking reciprocals of (4.8) and (4.9), it follows that

$$A^R \cdot T_i^R + B^R \cdot S_i^R = R_i^R \quad (4.10)$$

and

$$A^R \cdot T_{i-1}^R + B^R \cdot S_{i-1}^R = R_{i-1}^R \quad (4.11)$$

¹By convention, a polynomial of negative degree is the zero polynomial.

Let

$$Q_{i+1}(z) = R_{i-1}(z) / R_i(z) \mod z^{\mu_{MN}+2} \quad (4.12)$$

be a polynomial of degree $\mu_{MN}+1$. Then

$$R_{i-1}(z) - Q_{i+1}(z) \cdot R_i(z) = z^{\mu_{MN}+2} \bar{R}(z), \quad (4.13)$$

for some polynomial $\bar{R}(z)$ of degree $m-N-\mu_{MN}-2$. Taking reciprocals of (4.13), it follows that

$$R_{i-1}^R(z) - Q_{i+1}^R(z) \cdot R_i^R(z) = \bar{R}^R(z). \quad (4.14)$$

That is, $Q_{i+1}^R(z)$ and $\bar{R}^R(z)$ are the quotient and remainder, respectively, on division of $R_{i-1}^R(z)$ by $R_i^R(z)$.

With $Q_{i+1}(z)$ defined by (4.12), step 7 of Algorithm 3 yields

$$\begin{bmatrix} S_{i+1} & S_i \\ T_{i+1} & T_i \end{bmatrix} = \begin{bmatrix} z^{\mu_{MN}+2} \cdot S_{i-1} - Q_{i+1} \cdot S_i & z^{\mu_{MN}} \cdot S_i \\ z^{\mu_{MN}+2} \cdot T_{i-1} - Q_{i+1} \cdot T_i & z^{\mu_{MN}} \cdot T_i \end{bmatrix}, \quad (4.15)$$

which on taking reciprocals becomes

$$\begin{bmatrix} S_{i+1}^R & S_i^R \\ T_{i+1}^R & T_i^R \end{bmatrix} = \begin{bmatrix} S_{i-1}^R - Q_{i+1}^R \cdot S_i^R & S_i^R \\ T_{i-1}^R - Q_{i+1}^R \cdot T_i^R & T_i^R \end{bmatrix}. \quad (4.16)$$

Looking ahead one iteration, we obtain

$$\begin{aligned} R_{i+1}^R &= A^R \cdot T_{i+1}^R + B^R \cdot S_{i+1}^R \\ &= A^R \cdot [T_{i-1}^R - Q_{i+1}^R \cdot T_i^R] + B^R \cdot [S_{i-1}^R - Q_{i+1}^R \cdot S_i^R] \\ &= R_{i-1}^R - Q_{i+1}^R \cdot R_i^R. \end{aligned} \quad (4.17)$$

Thus, $\bar{R}^R(z) = R_{i+1}^R(z)$ in equation (4.14).

Summarizing, let Q_{i+1}^R be the quotient on division of R_{i-1}^R by R_i^R . Then

$$\begin{aligned} R_{i+1}^R &= R_{i-1}^R - Q_{i+1}^R \cdot R_i^R \\ S_{i+1}^R &= S_{i-1}^R - Q_{i+1}^R \cdot S_i^R \end{aligned} \quad (4.18)$$

$$T_{i+1}^R = T_{i-1}^R - Q_{i+1}^R \cdot T_i^R,$$

which are the fundamental relations describing Euclid's extended algorithm.

To complete the proof, it remains to show that the initial conditions for Euclid's extended algorithm are satisfied. Initialization in step 1 of Algorithm 3 yields

$$\begin{bmatrix} S_0 & S_{-1} \\ T_0 & T_{-1} \end{bmatrix} = \begin{bmatrix} -Q_0 & z^{m-n-1} \\ 1 & 0 \end{bmatrix}, \quad (4.19)$$

where, as in (4.12), Q_0^R is the quotient on division of A^R by B^R . Steps 3 and 5 with $i = 0$ then become

$$A \cdot 1 + B \cdot (-Q_0) = z^{m-n+\mu_{m-n,0}+1} R_0 \quad (4.20)$$

and

$$A \cdot 0 + B \cdot z^{m-n-1} = z^{m-n-1} R_{-1} \quad (4.21)$$

Taking reciprocals of (4.19), (4.20) and (4.21), it follows that

$$\begin{array}{lll} R_{-1}^R = B^R & S_{-1}^R = 1 & T_{-1}^R = 0 \\ R_0^R = A^R - Q_0^R \cdot B^R & S_0^R = -Q_0^R & T_0^R = 1 \end{array} \quad (4.22)$$

Corollary 4.2. At the i -th iteration of Algorithm 3, let S_i / T_i be the scaled Pade fraction of type (M, N) for $-A / B$, such that

$$A \cdot T_i + B \cdot S_i = z^{M+N+\mu_{M,N}+1} R_i.$$

Then,

$$\begin{aligned} R_{i-1} \cdot S_i - z^{\mu_{M,N}+2} R_i \cdot S_{i-1} &= (-1)^i \cdot A \\ R_{i-1} \cdot T_i - z^{\mu_{M,N}+2} R_i \cdot T_{i-1} &= (-1)^{i+1} \cdot B \\ S_{i-1} \cdot T_i - S_i \cdot T_{i-1} &= (-1)^{i+1} z^{M+N-1}. \end{aligned} \quad (4.23)$$

Furthermore, the algorithm terminates for some $i = k$, where $k \leq n$. On termination, S_{k+1} / T_{k+1} is

the scaled Pade fraction of type (m, n) for $-A/B$ such that

$$A \cdot T_{k+1} + B \cdot S_{k+1} = 0. \quad (4.24)$$

In addition,

$$A \cdot T_k + B \cdot S_k = z^{m+n-2\lambda_m-1} R_k, \quad (4.25)$$

where

$$z^{\lambda_m} = \text{GCD}(S_{k+1}, T_{k+1})$$

and

$$R_k = \text{GCD}(A, B).$$

Proof: From Euclid's extended algorithm (4.18) with initial conditions (4.22), it follows that (see, for example, McEliece [McE78])

$$\begin{aligned} R_{i-1} \cdot S_i^R - R_i^R \cdot S_{i-1}^R &= (-1)^i \cdot A^R \\ R_{i-1}^R \cdot T_i^R - R_i^R \cdot T_{i-1}^R &= (-1)^{i+1} \cdot B^R \\ S_{i-1}^R \cdot T_i^R - S_i^R \cdot T_{i-1}^R &= (-1)^{i+1}. \end{aligned} \quad (4.26)$$

By using the correspondence established in theorem 4.1 and taking reciprocals, equations (4.26) result in equations (4.23).

Furthermore, Euclid's extended algorithm terminates for some $k < n$ when

$$A^R \cdot T_{k+1}^R + B^R \cdot S_{k+1}^R = R_{k+1}^R = 0 \quad (4.27)$$

and

$$A^R \cdot T_k^R + B^R \cdot S_k^R = R_k^R = \text{GCD}(A^R, B^R), \quad (4.28)$$

where $\lambda = \partial(R_k^R)$, $\partial(S_{k+1}^R) \leq m - \lambda$, $\partial(T_{k+1}^R) \leq n - \lambda$, $\partial(S_i^R) < m - \lambda$ and $\partial(T_i^R) < n - \lambda$.

Taking reciprocals of (4.28) with respect to $z^{m+n-\lambda-1}$ results in

$$A \cdot T_k + B \cdot S_k = z^{m+k-\lambda-1} \cdot R_k,$$

where $R_k(0) \neq 0$. Thus, S_k / T_k is the scaled Pade fraction of type $(m - \lambda - 1, n - \lambda - 1)$ for $-A / B$. Taking reciprocals of (4.27) with respect to z^{m+n} gives

$$A \cdot T_{k+1} + B \cdot S_{k+1} = 0, \quad (4.29)$$

where $\partial(S_{k+1}) \leq m$ and $\partial(T_{k+1}) \leq n$. Thus, S_{k+1} / T_{k+1} is the scaled Pade fraction of type (m, n) such that

$$\text{GCD}(S_{k+1}, T_{k+1}) = z^\lambda. \quad (4.30)$$

Thus $\lambda = \lambda_{m,n}$. ■

As a consequence of Corollary 4.2, Algorithm 3 computes co-multipliers S_k and T_k , only, such that

$$A^R \cdot T_k^R + B^R \cdot S_k^R = R_k^R = \text{GCD}(A^R, B^R). \quad (4.31)$$

The remainder R_k^R is available only if the multiplications in steps 3 and 6 are performed without the modulo operation.

4.3. Fast GCD Computations

Theorem 4.3. Algorithm 2 applied to $-A(z) / B(z)$, where $A(z)$ and $B(z)$ are polynomials of degree m and n , respectively, returns the scaled Pade fraction S_1 / T_1 of type (m, n) such that

$$A^R \cdot T_1^R + B^R \cdot S_1^R = 0 \quad (4.32)$$

and its predecessor S_0 / T_0 of type $(m - \lambda_{m,n} - 1, n - \lambda_{m,n} - 1)$ such that

$$A^R \cdot T_0^R + B^R \cdot S_0^R = R_0^R. \quad (4.33)$$

where

$$z^{\lambda_{mn}} = \text{GCD}(S_1, T_1)$$

and

$$R_0 = \text{GCD}(A, B).$$

Proof: Since scaled Pade fractions are unique, the result is an immediate consequence of Corollary 4.2. ■

The greatest common divisor R_0 is not explicitly computed by Algorithm 2. However, $\partial(R_0) = \lambda_{mn} \leq n$. Using fast multiplication, it can therefore be determined in $O(n \log n)$. As a consequence of this and Theorem 3.8, we have

Theorem 4.4. Algorithm 2 can compute the greatest common divisor, the cofactors and the comultipliers of two polynomials of degrees m and n , where $m \geq n$, in time $O(m \log m) + O(n \log^2 n)$. ■

Thus, Algorithm 2 for GCD computations is basically of the same asymptotic complexity as the fast algorithm of Moenck [MOE73], Aho, Hopcroft and Ullman [AHO74] and Brent, Gustavson and Yun [BRE80], which are of complexity $O((m+n) \log^2(m+n))$. However, Algorithm 2 has the advantage of being partly iterative (approximately half as many recursive calls are used in comparison with the other fast methods). This can result in significant cost savings in an implementation environment (Verheijen [VER83]).

4.4. Antidiagonal Computations

Let

$$A(z) = \sum_{i=0}^{\infty} a_i z^i \tag{4.34}$$

be a unit power series, and let

$$A_d^R(z) = \sum_{i=0}^d a_{d-i} z^i, \quad d \geq 0, \quad (4.35)$$

be the reciprocal of the first $d+1$ terms of $A(z)$. In this section, we examine the intermediate results obtained by Algorithm 2² while computing the scaled Pade fraction of type (n,n) , $2n \leq d$, for the two polynomials $1 + z A_d^R(z)$ and $B^R(z) = -1$.

At the i -th iteration of Algorithm 2, the scaled Pade fraction $S_{NN}(z) / T_{NN}(z)$ of type (N,N) for $\{1 + z A_d^R(z)\} / B^R(z)$ and its predecessor $S_{N^*N^*}(z) / T_{N^*N^*}(z)$ are determined such that

$$\{1 + z A_d^R(z)\} \cdot T_{NN}(z) - S_{NN}(z) = z^{2N^* + \lambda_{NN} + 1} R_1(z) \quad (4.36)$$

$$\{1 + z A_d^R(z)\} \cdot T_{N^*N^*}(z) - S_{N^*N^*}(z) = z^{2N^* + 1} R_0(z), \quad (4.37)$$

where

$$N^* = N - \lambda_{NN} - 1$$

and

$$z^{\lambda_{NN}} = \text{GCD}(S_{NN}, T_{NN}).$$

By taking reciprocals of (4.36) and (4.37), it follows that

$$A_d(z) \cdot T_{NN}^R(z) - R_1^R(z) = z^{d+1} \{S_{NN}^R(z) - T_{NN}^R(z)\} \quad (4.38)$$

$$A_d(z) \cdot T_{N^*N^*}^R(z) - R_0^R(z) = z^{d+1} \{S_{N^*N^*}^R(z) - T_{N^*N^*}^R(z)\} \quad (4.39)$$

where

$$d(R_1^R) = d - N - \lambda_{NN} \quad (4.40)$$

and

$$d(R_0^R) = d - N^* \quad (4.41)$$

² It is assumed that no truncation of polynomial operations takes place.

Theorem 4.5: Let $M = d - N$ and $M^* = d - N^*$. Then the scaled Pade fractions of type (M, N) and (M^*, N^*) for $A(z)$ are $R_1^R(z) / T_{NN}^R(z)$ and $R_0^R(z) / T_{N^*N^*}^R(z)$, respectively.

Proof: From (4.38), (4.39), (4.40) and (4.41), clearly the degree and order conditions for scaled Pade fractions are satisfied. To complete the proof, let

$$G(z) = \text{GCD}(T_{NN}^R, R_1^R).$$

Then, from (4.38), $G(z)$ must divide not only T_{NN}^R but $z^{d+1} S_{NN}^R(z)$, as well. Since $\text{GCD}(S_{NN}^R, T_{NN}^R) = 1$, it follows that

$$G(z) = z^\lambda$$

for some $\lambda \geq 0$. Thus, $R_1^R(z) / T_{NN}^R(z)$ is a scaled Pade fraction of type (M, N) . Similarly, $R_0^R(z) / T_{N^*N^*}^R(z)$ is a scaled Pade fraction of type (M^*, N^*) . ■

Observe that

$$M + N = d \quad (4.42)$$

and

$$M^* + N^* = d \quad (4.43)$$

The scaled Pade fractions $R_1^R(z) / T_{NN}^R(z)$ and $R_0^R(z) / T_{N^*N^*}^R(z)$ both lie along the d -th anti-diagonal path of the scaled Pade table $I(A)$, where

Definition (see, for example, Brent [BRE80]). The d -th anti-diagonal path of the scaled Pade table $I(A)$ for a unit power series $A(z)$ is defined to the set of all scaled Pade fractions of type (i, j) , where

$$i + j = d. \quad (4.44)$$

But, by Theorem 4.5, the scaled Pade fraction $R_1^R(z) / T_{NN}^R(z)$ and its predecessor $R_0^R(z) / T_{N^*N^*}^R(z)$, along the d -th anti-diagonal path, are obtained by applying Euclid's extended

algorithm to the reciprocals of $1 + z A_d^R(z)$ and $B^R(z) = -1$, that is, to the polynomials $A_d(z) + z^{d+1}$ and $-z^{d+1}$. However,

$$\text{GCD}(A_d(z) + z^{d+1}, -z^{d+1}) = \text{GCD}(A_d(z), -z^{d+1}). \quad (4.45)$$

This is precisely the result given by McFlice [McE78]; namely, Euclid's extended algorithm applied to $A_d(z)$ and $-z^{d+1}$ yields all the Pade fractions along the d -th anti-diagonal path of the Pade table for $A(z)$. Equivalently, by Theorem 4.5, the same Pade fractions can be obtained by applying Algorithm 3 to the two polynomials $1 + z A_d^R(z)$ and $B^R(z) = -1$, and then taking reciprocals of the results.

In addition, by Theorem 4.5, Algorithm 2 can be used to compute any specific scaled Pade fraction $R_1^R(z) / T_{NN}^R(z)$ of type (M, N) , where $M + N = d$, along the d -th anti-diagonal path for $A(z)$. This requires $O(N \log^2 N)$ arithmetic operations to compute, by Algorithm 2, the scaled Pade fraction $S_{NN}(z) / T_{NN}(z)$ of type (N, N) for $1 + z A_d^R(z)$, plus an additional $O((d-N) \log(d-N))$ operations to determine $R_1^R(z)$ which satisfies (4.36). This asymptotic cost is the same as the cost of the fast algorithm of Brent [BRE80] for computing the greatest common divisor of $A_d(z)$ and $-z^{d+1}$. However, their BRENT algorithm can compute all the Pade fractions along the d -th anti-diagonal only.

CHAPTER 5

THE INVERSE OF HANKEL AND TOEPLITZ MATRICES

5.1. Introduction

In this chapter, scaled Bode functions are used to construct the inverse of a nonsingular Hankel matrix

$$H = \begin{bmatrix} a_1 & a_n \\ \vdots & \vdots \\ \vdots & \vdots \\ \vdots & \vdots \end{bmatrix}$$

and the inverse of the Toeplitz matrix

$$T = \begin{bmatrix} a_n & \vdots & \vdots & \vdots & \vdots & \vdots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \end{bmatrix}$$

associated with the Hankel matrices

The formula derived for the inverse is similar to those given by Trench [TREN65], Zohar [ZOH74], and Knuth et al [KAT88] but it has the advantage of being successful in the degenerate case where a principal minor of H is zero. The formula is due to Ahlberg and [AHL79].

$$S(z)/T(z) = \sum_{j=0}^n s_j z^j + \sum_{j=0}^n t_j z^j \quad (5.1)$$

be the scaled L -th function $f(z) = f_n(z)$ for

$$f(z) = \sum_{j=0}^n f_j z^j \quad (5.2)$$

Assume that

$$f_n \neq 0$$

Let $\alpha_1, \alpha_2, \dots, \alpha_n$ be the zeros of $f(z)$, the above equation (5.1) can be written

$$\begin{bmatrix} 1 & \alpha_1 & \alpha_1^2 & \dots & \alpha_1^n \\ 1 & \alpha_2 & \alpha_2^2 & \dots & \alpha_2^n \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & \alpha_n & \alpha_n^2 & \dots & \alpha_n^n \end{bmatrix} \begin{bmatrix} s_0 \\ s_1 \\ s_2 \\ \vdots \\ s_n \end{bmatrix} = \begin{bmatrix} t_0 \\ t_1 \\ t_2 \\ \vdots \\ t_n \end{bmatrix}$$

First, we consider the case $n=1$, then the above equation (5.1) can be written

1

$$\begin{bmatrix} 1 & \alpha_1 \\ 1 & \alpha_2 \end{bmatrix} \begin{bmatrix} s_0 \\ s_1 \end{bmatrix} = \begin{bmatrix} t_0 \\ t_1 \end{bmatrix} \quad (5.3)$$

$$\begin{bmatrix} a_1 & \dots & a_n \\ \vdots & & \vdots \\ a_{n-1} & \dots & a_{2n-1} \end{bmatrix} \begin{bmatrix} v_{n-1} \\ \vdots \\ v_1 \\ v_0 \end{bmatrix} = \begin{bmatrix} 0 \\ \vdots \\ 0 \end{bmatrix} \quad (5.6)$$

Lemma 5.1 Let

$$y = (a_n, \dots, a_{n-1}) (v_{n-1}, \dots, v_0)^t. \quad (5.7)$$

Then $y \neq 0$.

Proof. Suppose $y = 0$. Then, from (5.6), $(v_{n-1}, \dots, v_0, 0)^t$ satisfies

$$\begin{bmatrix} a_1 & \dots & a_n & a_{n+1} \\ \vdots & & \vdots & \vdots \\ a_{n-1} & \dots & a_{2n-2} & a_{2n-1} \\ a_n & \dots & a_{2n-1} & 0 \end{bmatrix} \begin{bmatrix} v_{n-1} \\ \vdots \\ v_0 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ \vdots \\ 0 \\ 0 \end{bmatrix};$$

that is, $(v_{n-1}, \dots, v_0, 0)^t$ is also a solution of (5.4). This implies $\det(H_{n+1}) = 0$, which is a contradiction. \square

Lemma 5.2 Let $\lambda \in \mathbb{C}$. Denote the inverse of $H_{n+1}(\lambda)$

$$L(\lambda) = (v_{n+1}(\lambda), \dots, v_0(\lambda))^t \quad (5.8)$$

where $v_i(\lambda)$ is given by (5.1)

and $L(\lambda)$ is a solution of system (5.6) becomes

$$\begin{bmatrix} \lambda & \dots & \lambda \\ \vdots & & \vdots \\ a_{1n-1} & \dots & a_{2n-1} \end{bmatrix} \begin{bmatrix} v_{n-1} \\ \vdots \\ v_0 \end{bmatrix} = \begin{bmatrix} 0 \\ \vdots \\ 0 \end{bmatrix}$$

Equivalently,

$$\begin{bmatrix} v_{n-1} \\ \vdots \\ v_0 \end{bmatrix} = \begin{bmatrix} b_{11} & \dots & b_{1n} \\ \vdots & & \vdots \\ b_{n1} & \dots & b_{nn} \end{bmatrix} \begin{bmatrix} 0 \\ \vdots \\ 0 \\ \gamma \end{bmatrix}$$

and (5.8) now follows since B_{nn} is symmetric. ■

To obtain the remaining elements of B_{nn} , two cases, $t_n \neq 0$ and $t_n = 0$, of the solution of (5.1) are identified.

Case 1: Assume that $t_n \neq 0$. Then (5.4) becomes

$$\begin{bmatrix} a_1 \\ \vdots \\ a_n \\ \vdots \\ a_{n+1} \end{bmatrix} = \begin{bmatrix} a_2 & \dots & a_n & a_{n+1} \\ \vdots & & \vdots & \vdots \\ a_n & \dots & a_{2n-1} & a_{2n} \\ a_{n+1} & \dots & a_{2n-1} & 0 \end{bmatrix} \begin{bmatrix} t_n \\ \vdots \\ t_1 \end{bmatrix} = \begin{bmatrix} 0 \\ \vdots \\ \gamma \\ 0 \end{bmatrix} \quad (5.9)$$

$$H_{2n-1,n} = \begin{bmatrix} a_1 & \dots & a_{n+1} \\ \vdots & & \vdots \\ a_n & \dots & a_{2n-1} \\ a_{n+1} & \dots & a_{2n-1} & 0 \end{bmatrix}$$

If $\det(H_{2n-1,n}) \neq 0$, for otherwise (5.9) also has a solution with $t_n = 0$. Then t_n is a scalar

of the unique solution t which is a solution of the solution of (5.4).

$$t_n = \frac{1}{\det(H_{2n-1,n})} \det \begin{bmatrix} a_2 & \dots & a_n & a_{n+1} \\ \vdots & & \vdots & \vdots \\ a_n & \dots & a_{2n-1} & a_{2n} \end{bmatrix}$$

Thus

$$b_{ij} = \bar{b}_{i-1,j} - (t_{n+1-i}/t_0) \bar{b}_{n,j}, \quad i, j = 1, \dots, n, \quad (5.11)$$

where $\bar{b}_{0,j} = 0$, $j = 1, \dots, n$.

Proof: From (5.4)

$$\begin{bmatrix} t_n \\ \vdots \\ t_1 \end{bmatrix} [\bar{h}_{n1}, \dots, \bar{b}_{nn}] = -t_0 \begin{bmatrix} b_{11} & \dots & b_{1n} \\ \vdots & & \vdots \\ h_{n1} & \dots & b_{nn} \end{bmatrix} \begin{bmatrix} a_{n+1} \\ \vdots \\ a_{2n-1} \\ 0 \end{bmatrix} [\bar{b}_{n1}, \dots, \bar{b}_{nn}]. \quad (5.12)$$

However,

$$H_{nn} \begin{bmatrix} 0 & \dots & 0 \\ \bar{b}_{11} & & \bar{b}_{1n} \\ \vdots & & \vdots \\ \bar{b}_{n-1,1} & \dots & \bar{b}_{n-1,n} \end{bmatrix} = I_n - \begin{bmatrix} a_{n+1} \\ \vdots \\ a_{2n-1} \\ 0 \end{bmatrix} [\bar{b}_{n1}, \dots, \bar{b}_{nn}]. \quad (5.13)$$

Consequently, by substituting (5.13) into (5.12), it follows that

$$\begin{bmatrix} h_{11} & \dots & h_{1n} \\ \vdots & & \vdots \\ h_{n1} & \dots & h_{nn} \end{bmatrix} = \begin{bmatrix} 0 & \dots & 0 \\ \bar{h}_{11} & & \bar{h}_{1n} \\ \vdots & & \vdots \\ \bar{h}_{n-1,1} & \dots & \bar{h}_{n-1,n} \end{bmatrix} - t_0^{-1} \begin{bmatrix} t_n \bar{b}_{n1} & \dots & t_n \bar{b}_{nn} \\ \vdots & & \vdots \\ t_1 \bar{h}_{n1} & \dots & t_1 \bar{b}_{nn} \end{bmatrix} \quad (5.14)$$

Case 2. Assume $t_n = 0$. Consider the subsystem

$$\begin{bmatrix} a_2 & \cdots & a_n \\ \vdots & & \vdots \\ a_n & \cdots & a_{2n-2} \end{bmatrix} \begin{bmatrix} t_{n-1} \\ \vdots \\ t_1 \end{bmatrix} = -t_0 \begin{bmatrix} a_{n+1} \\ \vdots \\ a_{2n-1} \end{bmatrix} \quad (5.15)$$

of (5.4). Then the matrix on the left-hand-side of (5.15) is nonsingular. For, suppose otherwise.

Then there exists a nontrivial vector $[\alpha_{n-1}, \dots, \alpha_1]^T$ such that

$$\begin{bmatrix} a_2 & \cdots & a_n \\ \vdots & & \vdots \\ a_n & \cdots & a_{2n-2} \end{bmatrix} \begin{bmatrix} \alpha_{n-1} \\ \vdots \\ \alpha_1 \end{bmatrix} = \begin{bmatrix} 0 \\ \vdots \\ 0 \end{bmatrix} \quad (5.16)$$

However, by taking the transpose of (5.15), it follows from (5.16) that

$$\begin{bmatrix} \alpha_{n-1} \\ \vdots \\ \alpha_1 \end{bmatrix} = -t_0^{-1} \begin{bmatrix} a_{n+1} & \cdots & a_{2n-1} \end{bmatrix} \begin{bmatrix} a_2 & \cdots & a_n \\ \vdots & & \vdots \\ a_n & \cdots & a_{2n-2} \end{bmatrix} \begin{bmatrix} \alpha_{n-1} \\ \vdots \\ \alpha_1 \end{bmatrix} = \begin{bmatrix} 0 \\ \vdots \\ 0 \end{bmatrix} \quad (5.17)$$

Consequently,

$$\begin{bmatrix} a_1 & \cdots & a_n \\ \vdots & & \vdots \\ a_n & \cdots & a_{2n-1} \end{bmatrix} \begin{bmatrix} 0 \\ \vdots \\ \alpha_{n-1} \\ \vdots \\ \alpha_1 \end{bmatrix} = \begin{bmatrix} 0 \\ \vdots \\ 0 \end{bmatrix}$$

which implies that $\det(M) \neq 0$. This contradiction implies that

$$\det \begin{bmatrix} a_2 & \cdots & a_n \\ \vdots & & \vdots \\ a_n & \cdots & a_{2n-2} \end{bmatrix} \neq 0.$$

Now, since $t_n = 0$, it follows from (5.4) that $\det(H_{n+1,n}) = 0$. Let

$$H_{n+1,n}^* = \begin{bmatrix} a_2 & \cdots & a_n & a_{n+1} \\ \vdots & & \vdots & \vdots \\ a_n & & a_{2n-2} & a_{2n-1} \\ a_{n+1} & \cdots & a_{2n-1} & 1 \end{bmatrix} \quad (5.18)$$

Then, expansion with respect to the last column yields

$$\det(H_{n+1,n}^*) = \det \begin{bmatrix} a_2 & \cdots & a_n \\ \vdots & & \vdots \\ a_n & & a_{2n-2} \\ a_{n+1} & \cdots & a_{2n-1} \end{bmatrix} \neq 0. \quad (5.19)$$

Lemma 5.4. Assume $t_n = 0$, and let $B_{n+1,n}^* = (b_{ij}^*)_{i,j=1}^n$ denote the inverse of $H_{n+1,n}^*$. Then

$$b_{ij} = b_{i-1,j}^* + (b_{i,n}^* - t_{n+1-i} / t_0) b_{n,j}^*, \quad i, j = 1, \dots, n, \quad (5.20)$$

where $b_{0,j}^* = 0$, $j = 1, \dots, n$.

Proof: The result follows using arguments similar to those of the proof of Lemma 5.3. ■

Theorem 5.5 Let $(t_n, t_{n-1}, \dots, t_0)^t$ and $(v_{n-1}, v_{n-2}, \dots, v_0)^t$ be given by (5.1) and (5.5), respectively. If $t_0 \neq 0$, then the inverse of the H_n is given by the Christoffel-Darboux formula

$$\begin{aligned}
 H_{nn}^{-1} = & \frac{1}{\gamma t_0} \left\{ \begin{bmatrix} t_{n-1} & \cdots & t_0 \\ \vdots & \ddots & \vdots \\ t_0 & & 0 \end{bmatrix} \begin{bmatrix} v_{n-1} & \cdots & v_0 \\ \vdots & \ddots & \vdots \\ 0 & & v_{n-1} \end{bmatrix} \right. \\
 & \left. - \begin{bmatrix} v_{n-2} & \cdots & v_0 & 0 \\ \vdots & \ddots & \vdots & \vdots \\ v_0 & & 0 & \vdots \\ 0 & & & 0 \end{bmatrix} \begin{bmatrix} t_n & \cdots & t_1 \\ \vdots & \ddots & \vdots \\ 0 & & t_n \end{bmatrix} \right\} \quad (5.21)
 \end{aligned}$$

Proof: Consider the case $t_n \neq 0$. Then equation (5.11) of Lemma 5.3 yields

$$\bar{b}_{nn} = -(t_0 / t_n) b_{1n}$$

and

$$\bar{b}_{i+1,n} = \bar{b}_{i,n} - (t_{n-i} / t_0) \bar{b}_{nn}, \quad i = 1, 2, \dots, n.$$

Consequently,

$$b_{i+1,n} = \bar{b}_{i,n} + (t_{n-i} / t_n) b_{1n}, \quad i = 1, 2, \dots, n. \quad (5.22)$$

Since $\bar{b}_{i,n} = \bar{b}_{n-i}$, equation (5.22) inserted into (5.11) gives

$$b_{ij} = \bar{b}_{i+1,j} - (t_{n-i-1} / t_0) [b_{j+1,n} - (t_{n-j} / t_n) b_{1n}].$$

Thus,

$$b_{j+1,i+1} = \bar{b}_{j+1,i+1} - (t_{n-j} / t_0) [b_{i,n} - (t_{n-i-1} / t_n) b_{1n}].$$

Therefore,

$$b_{ij} = b_{j+1,i+1} + t_0^{-1} [t_{n-j} b_{i,n} - t_{n-i-1} b_{j+1,n}].$$

From (5.8) of Lemma 5.2, it now follows that

$$b_{ij} = b_{j+1,i+1} + (\gamma t_0)^{-1} [t_{n-j} v_{n-1} - t_{n-i-1} v_{n-1-j}],$$

$$i, j = 1, 2, \dots, n, \quad (5.23)$$

where all variables with subscripts outside the range $1, 2, \dots, n$ are assumed to be zero. But equation (5.23) corresponds exactly to (5.21).

The case $r_n = 0$ makes use of Lemma 5.4, and the proof proceeds in a similar manner. ■

Now consider the nonsingular Hankel system

$$H_{n,n} \mathbf{x} = \begin{bmatrix} a_1 & \dots & a_n \\ \vdots & & \vdots \\ a_n & \dots & a_{2n-1} \end{bmatrix} \mathbf{x} = \mathbf{b}, \quad (5.24)$$

for an arbitrary vector \mathbf{b} . From (5.21), the computation of $\mathbf{x} = H_{nn}^{-1} \cdot \mathbf{b}$ can be done by performing four polynomial multiplications. Using fast polynomial multiplication methods, this requires $O(n \log n)$ arithmetic operations. By theorem 3.6, the polynomials $T(z)$ and $V(z)$ can be computed in $O(n \log^2 n)$ operations, and therefore

Theorem 5.6. The system (5.24) can be solved in $O(n \log^2 n)$ arithmetic operations.

Example: Let

$$H_{3,3} = \begin{bmatrix} 2 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 4 \end{bmatrix},$$

and let

$$A(z) = a_0 + 2z + z^4 + 4z^5 + 0z^6 + \dots$$

be the associated unit power series. Then the scaled Pade fraction of type (3,3) for $A(z)$ is given by $S(z)/T(z)$, where

$$S(z) = 2a_0 + 4(1 - 2a_0)z - 16(1 - 2a_0)z^2 + (64 - a_0)z^3$$

and

$$T(z) = 2 - 8z + 32z^2 - z^3.$$

Thus, $H_{3,3}$ is nonsingular, since $T(0) = t_0 \neq 0$. The predecessor of $S(z)/T(z)$ is the scaled Pade fraction $U(z)/V(z)$ of type (2,2), where

$$U(z) = a_0 z + 2 z^2$$

and

$$V(z) = z.$$

Therefore, the solution of system (5.4),

$$\begin{bmatrix} 2 & 0 & 0 & 1 \\ 0 & 0 & 1 & 4 \\ 0 & 1 & 4 & 0 \end{bmatrix} \begin{bmatrix} t_3 \\ t_2 \\ t_1 \\ t_0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix},$$

is $(t_3, t_2, t_1, t_0)' = (-1, 32, -8, 2)'$; and the solution of system (5.6),

$$\begin{bmatrix} 2 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} v_2 \\ v_1 \\ v_0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix},$$

is $(v_2, v_1, v_0)' = (0, 1, 0)$.

The inverse $H_{3,3}$ is obtained by the Christoffel-Darboux formula (5.21) to be

$$H_{3,3}^{-1} = (1/2) \left\{ \begin{bmatrix} 32 & -8 & 2 \\ -8 & 2 & 0 \\ 2 & 0 & 0 \end{bmatrix} \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix} - \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} -1 & 32 & -8 \\ 0 & -1 & 32 \\ 0 & 0 & -1 \end{bmatrix} \right\}$$

$$= \begin{bmatrix} 1/2 & 0 & 0 \\ 0 & -4 & 1 \\ 0 & 1 & 1 \end{bmatrix} \quad \blacksquare$$

Since Toeplitz matrices are equivalent to Hankel matrices (simply by a change of notation), the results of theorem 5.6 apply to Toeplitz matrices, as well.

Consequently, by Theorem 5.6, the inverse of a Toeplitz matrix of order n can be obtained in $O(n \log^2 n)$ arithmetic operations. This is done with one call of Algorithm 1 for the computation of a specific scaled Pade fraction. The fast algorithm of Brent [BRE80] is also of complexity $O(n \log^2 n)$, but the computation of the Toeplitz inverse may require as many as four calls of their $O(n \log^2 n)$ PRSDC algorithm for computing a specific Pade fraction. In addition, each call is burdened with complications, alluded to earlier in Section 4.4.

CHAPTER 6

CONCLUSIONS AND SUGGESTIONS FOR FUTURE RESEARCH

Central to the classical theory of Padé approximants of power series are the concepts of the Padé form, which always exists but may not be unique, and the Padé fraction, which is unique but in a certain sense may not exist. The fundamental definition introduced in this thesis is the scaled Padé fraction which exists uniquely. It is shown that scaled Padé fractions satisfy a three-term relationship between elements along an off-diagonal path of the scaled Padé table. This relationship circumvents the problems of the degenerate case - i.e., the problems which plague other relationships such as those upon which the ϵ -algorithm and the qd-algorithm are based.

The three-term relationship is used to develop an $O(n^2)$ algorithm (Algorithm 3) which computes along an off-diagonal path, a finite sequence of successive scaled Padé fractions for the quotient of two power series. In the case that the power series is normal, Algorithm 3 is identical to the one described by Brezinski [BRZ76]. In the case that computations progress along the diagonal, Algorithm 3 becomes that of Cabay and Kao [CK83]. Furthermore, it is shown that if the two power series are finite (i.e., polynomials), then Algorithm 3 with computations along one specific off-diagonal path is exactly equivalent to Euclid's extended algorithm for computing the greatest common divisor of two polynomials. However, other off-diagonal paths can be used to compute greatest common divisors, and it remains a subject for future research to determine the optimal one.

By doubling the step at each iteration, the three term relationship gives rise to an $O(n \log^2 n)$ algorithm (Algorithm 1 or Algorithm 2). The cost complexity assumes the existence of fast methods for polynomial arithmetic. The decision to double the step size (rather than to triple it, for example) is not accidental. We believe that this choice is optimal (within the constraints placed by fast polynomial arithmetic methods), but a formal proof remains to be obtained.

When applied to polynomials, Algorithm 2 at each iteration routinely produces intermediate polynomial remainder sequences. For this reason, and because Algorithm 2 is simply a fast version of Algorithm 3, Algorithm 2 is truly a fast Euclid's extended algorithm. The PGCD, HGCD and EMGCD algorithms of Moenck [MOE73], of Aho, Hopcroft and Ullman [AHO74], and of Brent, Gustavson and Yun [BRE80], respectively, also compute greatest common divisors with the same cost complexity, but these methods are simply GCD methods. They do not produce intermediate polynomial remainder sequences. In addition, the artificial splitting of polynomials used by these methods to achieve their speed makes them difficult to understand. Finally, these methods are totally recursive, which makes them more costly to implement than Algorithm 2 which is semi-iterative.

For computing scaled Pade fractions for a power series, Brent et al [BRE80] have shown that the EMGCD algorithm can be modified, with substantial extra detail, to compute entries along an anti-diagonal path of the Pade table. We have shown that Algorithm 2 and Algorithm 3 can also be used to compute such entries, routinely. From a practical point of view, however, it seems to us that computations along an anti-diagonal path are not as natural as they are along an off-diagonal path. If the choice of the anti-diagonal path is incorrect, computations must be restarted. For off-diagonal computations, n need not be known a priori, and from an application point of view, this may be one of the most important contributions of this thesis.

It is not clear at this time whether or not the fast method, Algorithm 2, is useful in a practical environment. Initial experiments performed by Verheijen [VER83] indicate that the fast method, Algorithm 2, outperforms the classical one, Algorithm 3, only when n is greater than approximately 1600. However, as for the fast methods for polynomial multiplication, a hybrid of Algorithm 2 and Algorithm 3 should significantly lower this cross over point. This remains a subject for future research.

An effective implementation of an off-diagonal algorithm should prove to be a substantial tool in the design of a symbolic and algebraic system. A single routine set can

- (1) to obtain rational approximants of power series,
- (2) to convert rational functions from their power series representation
- (3) to compute greatest common divisors of polynomials, and
- (4) to solve Hankel and Toeplitz systems of equations.

The scope of this thesis includes those power series with coefficients over a field only. This restriction is relevant whenever division is required by the off diagonal algorithms. By choosing to perform pseudo-division, rather than division, the algorithms can be modified so that they are applicable to power series over a Euclidean domain rather than a field. As for Euclid's extended algorithm, the modified algorithms shall experience exponential growth of coefficients. It is a subject for future research to determine if methods similar to those of Collins [COL67] can be used to keep the growth linear. Other plausible methods for extending the results to Euclidean domains which need to be investigated, include the Chinese Remainder and Hensel algorithms.

The extension of results to Euclidean domains includes as a special case the multivariate power series. It is of interest to determine how this extension would compare with the methods for solving block Hankel systems, and for computing the least common multiple of power series. This remains a subject for future research.

As a final suggestion for future research, the validity of the results for the case of power series over a Euclidean domain should be investigated.

BIBLIOGRAPHY

- [AHU74] Aho, A.V., J.E. Hopcroft and J.D. Ullman, *The Design and Analysis of Computer Algorithms*, Addison-Wesley Mass. 1974.
- [AK75] Akaith, H. Block Toeplitz matrix inversion. *SIAM J. Appl. Math.* 24 (1975), pp. 234-241.
- [RAK75] Raker, G. In *Essentials of Padé Approximants*, Academic Press N.Y. 1975.
- [BH80] Bentley J., D. Hanken and J. Saxe, "A General Method for Solving Linear Recurrence Sequences", *ACM SIGACT NEWS* 21(1980), pp. 33-44.
- [BER68] Berlekamp E. R., *Algebraic Coding Theory*, McGraw Hill Co. N.Y. 1968.
- [BOC22] Bocher, M., *Introduction to High Algebra*, Macmillan Co. N.Y. 1922.
- [DB79] Dene, H.V. and S. Dasu, "2-D Matrix Existence Theorems and Decidability of Computation", *IEEE Trans. AC* (1979), pp. 502-512.
- [BRE75] Brent, D. J. C. Christyann, and D. V. V. Yun, "Fast Solution of Toeplitz Systems of Equations", *Mathematics of Computer*, *J. of Algebra* 36 (1975), pp. 259-295.
- [BR76] Brerinski, C., "Computation of Padé Approximants", *Comp. Appl. Math.* 2 (1976), pp. 113-127.
- [BR77] Brown, R. S., "On Euclid's Algorithm and the Computation of Padé Approximants", *Commun. IN. Pures. & Appl. Math.* 30 (1977), pp. 478-501.
- [BUT77] Butcher, A., "Recursive Algorithms for the Computation of Padé Approximants", *Math. Comp.* 31 (1977), pp. 113-127.

- [Bul79] Bultheel, A., "Recursive Algorithms for the Pade Tables: Two Approaches", in *Lecture Notes in Mathematics* 765, Springer Verlag Berlin, 1979, pp. 210-230.
- [CAB70] Cabay, S., CS566: Symbolic and Algebraic Manipulation Class Note, 1970
- [CAB77] Cabay, S. and T. P. Lam, "Congruential Technique for the Exact Solution of Integral Systems of Linear Equations", *ACM Trans. on Math. Systems*, 7(1977), no. 1, pp. 386-397.
- [CAB83] Cabay, S. and L. J. C. Kao, "The Diagonal Pade Table and the Triangular Decomposition of Hankel Matrices", TR83-10, Spring 1983.
- [Cla75] Claessens, G., "A new look at the Pade table and the different methods for computing its elements", *J. of Comp. and Appl. Math.*, 1(1975), pp. 141-152.
- [Col66] Collins, G.E., "Polynomial Determinants and their Minors", *Amer. Math. Mon.*, 73(1966), no. 7, pp. 577-581.
- [Col67] Collins, G.E., "Subresultant and Resultant Algorithms for Polynomials", *ACM*, 14(1967), no. 1, pp. 129-139.
- [Col71] Collins, G.E., "Computer Algebra", *Math. Mon.*, 80(1971), pp. 1-10.
- [Col72] Collins, G.E., "Symbolic Algebraic Manipulation", *Math. Mon.*, 81(1972), pp. 1-10.
- [Gag73] Gagg, W.B., "The Pade Table and the Padé Approximation", *Math. Mon.*, 82(1973), pp. 1-10.
- [Gag74] Gagg, W.B., "Padé Interpolations and the Padé Approximation", *Math. Mon.*, 83(1974), pp. 1-10.
- [Gag75] Gagg, W.B., "Fast Computation of Padé Approximations", *Math. Mon.*, 84(1975), pp. 1-10.

- [GUS79] Gustavson, F.G. and D.Y.Y. Yun, "Fast algorithms for rational Hermite approximation and solution of Toeplitz systems", *IEEE Trans. CS, CAS-26* (1979), No.9, pp. 750-755.
- [HOU69] Householder, A.S. and G.W. Stewart, "Eigengradients, Hankel Determinants and the Pade Table", in B. Dejon and P. Henrici (eds.), *Constructive Aspects of the Fundamental Theorem of Algebra*, Academic Press, New York, pp. 131-150.
- [KAI78] Kailath, T., A. Vieira and M. Morf, "Inverses of Toeplitz Operators, Innovations and Orthogonal Polynomials", *SIAM Review* 20, pp. 106-119.
- [KNU69] Knuth, D.F., *Seminumeric Algorithms*, Addison-Wesley, 1969.
- [LIP76] Lipson, J.D., "Newton's Method: A Greatest Algebraic Algorithm", *Proc. 1976 ACM Symp. on Symbolic and Algebraic Comp.* pp. 260-270.
- [LIP81] Lipson, J.D., *Elements of Algebra and Algebraic Computing*, Addison-Wesley, 1981.
- [McE78] McEliece, R.J. and J.B. Shearer, "A Property of Euclid's Algorithm and an Application to Pade Approximation", *SIAM J. Appl. Math.* 34(1978), No.4, pp. 611-617.
- [MOS73] Moses, I. and D.Y.Y. Yun, "The EMGCD Algorithm", *Proceeding of ACM annual conf.* (1973), pp. 159-166.
- [MOE73] Moenck, R., *Studies in Fast Algebraic Algorithms*, TR-57 University of Toronto, 1973.
- [RIS74] Rissanen, J., "Solution of Linear Equations with Hankel and Toeplitz Matrices", *Numer. Math.* 22(1974), pp. 361-366.
- [TRE61] Trench, W.F., "An Algorithm for the Inversion of Finite Toeplitz Matrices", *SIAM J.* 12(1964), No. 3, pp. 515-522.
- [TRE61] Trench, W.F., "An Algorithm for the Inversion of Finite Hankel Matrices", *SIAM J. Appl. Math.* 13, pp. 1102-1107.

- [VER83] Verheijen, A. *Evaluation of Diagonal Pade Fractions*, M.Sc. Thesis, University of Alberta, 1983, In preparation.
- [WYN60] Wynn, P., "The rational approximation of function which are formally defined by a power series expansion", *Math. Comp.*, 14(1960), pp. 147-186.
- [YUN79] Yun, D.Y.Y., "Uniform Bounds for a Class of Algebraic Mappings", *SIAM J. Comp.* 8(1979), No.3, pp. 348-356.
- [YUN74] Yun, D.Y.Y., "The Hensel Lemmma in Algebraic Manipulation", *Mac TR-138*, 1974.
- [ZOH69] Zohar, S., "Toeplitz Matrix Inversion: The Algorithm of W.F. Trench", *J. Assoc. Comp. Mach.* 16(1969), No.4, pp. 792-801.
- [ZOH74] Zohar, S., "The solution of a Toeplitz set of linear equations", *J. Assoc. Comp. Mach.* 21(1974), pp. 272-276.