

1 **Mechanisms and consequences of flight polyphenisms in an outbreaking bark beetle species**

2 **Authors:** Kelsey L. Jones¹, Rahmatollah Rajabzadeh², Guncha Ishangulyyeva², Nadir Erbilgin²,
3 and Maya L. Evenden¹

4 1. Department of Biological Sciences, University of Alberta

5 2. Department of Renewable Resources, University of Alberta

6 Corresponding Author: Kelsey Jones, kljones1@ualberta.ca

7 Summary Statement: This article explores the relationship between energy-use during flight and
8 host colonisation in the mountain pine beetle. The resulting information suggests potential
9 selection mechanisms maintaining flight polyphenisms.

10 Key Words: mountain pine beetle, dispersal, pheromone, host colonisation, polyphenism,
11 Scolytinae

12

13 **Abstract**

14 Flight polyphenisms naturally occur as discrete or continuous traits in insects. Discrete flight
15 polyphenisms include winged and wingless morphs, whereas continuous flight polyphenisms can
16 take the form of short- or long-distance fliers. The mountain pine beetle (*Dendroctonus*
17 *ponderosae*) exhibits polyphenic variation in flight distance but the consequences of this flight
18 variation on life history strategies of beetles is unknown. . This study assessed the effect of flight
19 on two particular aspects of beetle biology: (1) an energetic trade-off between flight distance and
20 host colonisation capacity; and (2) the relationship between flight distance and pheromone
21 production. A 23-h flight treatment was applied to a subset of beetles using computer. After
22 flight treatment, both flown and unflown (control) beetles were given the opportunity to colonise
23 bolts of host trees, and beetles that entered hosts were aerated to collect pheromone. A trade-off
24 occurred between initiation of host colonisation and percent body weight lost during flight,
25 which indicates energy-use during flight affects host acceptance in female mountain pine beetles.
26 Furthermore, production of the aggregation pheromone *trans*-verbenol by female beetles was
27 influenced by both percent weight lost during flight and flight distance. Male production of *exo*-
28 brevicomin was affected by beetle condition following flight but not by the energy used during
29 flight. These novel results give new insight into the polyphenic flight behaviour of mountain pine
30 beetles. Flight variation is adaptive by acting to maintain population levels through safe and
31 risky host colonisation strategies. These findings suggest mechanisms that facilitate the
32 extremities of the continuous flight polyphenism spectrum. These opposing mechanisms appear
33 to maintain the high variation in flight exhibited by this species.

34 **Introduction**

35 Polyphenisms are traits that exhibit two or more distinct phenotypes from a single genotype in
36 response to environmental conditions. The link between phenotypes and environmental factors
37 promotes individual success under changing environmental conditions (Simpson et al., 2011).
38 Although these distinct phenotypes may be advantageous for certain functions under different
39 conditions, they may develop at a cost to other life history traits (Kopp & Tollrain, 2003;
40 Karlsson et al., 2008).

41 Flight is costly, and trade-offs between resource allocation to flight and other life history
42 traits (Karlsson & Johansson, 2008), such as host (Latty & Reid, 2009; 2010) and reproduction
43 (Roff & Fairbairn, 1991; Lin et al., 2018) are common. The most notable flight polyphenism in
44 insects is the occurrence of winged and flightless morphs within the same species. Although
45 many polyphenisms are discrete, continuous flight polyphenisms can also exist as short- vs. long-
46 distance fliers (Karlsson & Johansson, 2008; Simpson et al., 2011). Most studies focus on
47 understanding the effects of discrete flight polyphenisms on subsequent life history strategies of
48 adult insects (Cisper et al., 2000), but the effects of continuous flight polyphenisms remain less
49 studied.

50 Continuous flight polyphenisms occur in aggressive tree-killing bark beetle species in the
51 genera *Dendroctonus* and *Ips* (Coleoptera: Curculionidae, Scolytinae) (Jones et al. 2019), which
52 influences obligatory dispersal for host colonisation and reproduction (Raffa et al., 2005).
53 Successful attack of a host tree requires the production of aggregation pheromones to attract
54 conspecifics for mass attack (Safranyik et al., 2010). The pioneering beetle (females in
55 *Dendroctonus* and males in *Ips*) releases aggregation pheromone that triggers the mass attack by
56 both sexes (Raffa et al., 2015). Beetles of the same sex as the pioneer initiate new attacks along
57 the tree bole, while beetles of the opposite sex enter existing galleries to mate (Gitau et al.,
58 2013). Bark beetles synthesize pheromone components *de novo* or through the activity of
59 microbial symbionts (Cale et al. 2019), but also require monoterpene precursors from the host
60 tree for pheromone synthesis (Blomquist et al., 2010).

61 Differences in pheromone production by beetles, however, have some fitness
62 consequences (Raffa, 2001). If production is low, beetle aggregation on the host tree will fail due
63 to adult beetle mortality as a result of exposure to toxic host secondary compounds (Raffa &
64 Berryman, 1982).

65 The host colonisation process is costly and depends on the physiological condition of the
66 adult bark beetles arriving at the host after dispersal (Reid et al., 2017). Several hypotheses have
67 been put forth to explain the relationships between dispersal behavior, host choice, and host
68 colonisation in bark beetles (Latty & Reid, 2010). The “desperation” hypothesis states that
69 beetles with low energy reserves enter a tree independent of host quality decisions because low
70 energy reserves prohibit further flight (Byers, 1999). The “safe site” hypothesis posits that

71 beetles enter high quality hosts to promote mate attraction and successful attack (Latty & Reid,
72 2010). The “condition matching” hypothesis suggests that host colonisation by the beetle should
73 interact with the quality of the host tree; as a result, beetles in good energetic condition can enter
74 well-defended trees (Chubaty et al., 2014).

75 The mountain pine beetle, (*Dendroctonus ponderosae*), is native to Western North
76 America, and has expanded its range eastward and northward (Cullingham et al., 2011)
77 following the most recent population outbreak that started in the early 2000s, and killed millions
78 of pine trees (Safranyik et al. 2010). Dispersal by flight dictates the spread of this species and it
79 is arguably the least understood aspect of mountain pine beetle ecology (Chen & Walton, 2011).

80 After emerging from the natal host, mountain pine beetles exhibit two patterns of
81 dispersal within the stand – spot growth and spot proliferation (Robertson et al., 2007). Spot
82 growth involves short distance movements from the natal host to a reproductive host located only
83 a few metres away. Spot proliferation results from beetle flight past suitable hosts followed by
84 host selection much further away from the natal host. Understanding the mechanism underlying
85 these flight polyphenisms in the mountain pine beetle and the cascading effects of flight
86 polyphenisms on subsequent host selection and colonisation are essential for understanding
87 population dynamics of the beetle (Robertson et al., 2007). Although some variation in flight
88 distance is explained by lipid content (Evenden et al., 2014), energy reserves alone do not
89 account for the large degree of flight variation exhibited by the mountain pine beetle (Shegelski
90 et al., 2019). One explanation of the varied flight behaviour in mountain pine beetle populations
91 is that beetles may require a flight period before becoming responsive to semiochemicals (Gray
92 et al., 1972), similar to other bark beetle species (Thompson & Bennett, 1971). Beetles with high
93 lipid levels need to spend energy before settling on a host, which could explain flight variation
94 over geographic and temporal scales (Robertson et al., 2007).

95 While beetle body condition (high lipid to body volume ratio) affects host colonisation
96 behaviour in mountain pine beetle (Elkin & Reid, 2005), it is unknown if the same lipid
97 resources consumed during flight (Evenden et al., 2014) are also allocated to host colonisation.
98 Although metabolic costs associated with pheromone production may be insignificant
99 (Pureswaran et al., 2006), mountain pine beetle aggregation pheromones are produced and/or
100 stored in the fat body (Song et al., 2014; Chiu et al., 2018). It is unknown whether lipid-use

101 during flight influences the production of the male-produced aggregation pheromone *exo-*
102 *brevicomini*, or the storage and use of *exo-brevicomini* and the female-produced aggregation
103 pheromone, *trans-verbenol*. Mountain pine beetle reproduction is also linked to body condition;
104 beetles in poor condition produce smaller eggs (Elkin & Reid, 2005), and there is a trade-off
105 between energy-use during flight and offspring production (Wijerathna et al., 2019).

106 In this study, we test the influence of flight polyphenisms on (1) female beetle host
107 acceptance; and (2) male and female production of aggregation pheromones. The outcome of this
108 study will reveal the relationship between energy-use during the obligatory dispersal phase of
109 mountain pine beetle and the subsequent host colonisation process.

110 **Materials & Methods**

111 *Collection of beetles*

112 Beetle-infested lodgepole pine, *Pinus contorta* var. *latifolia* Douglas, was collected as 50-cm
113 long cylindrical cross sections of a tree bole, hereafter referred to as “bolts”. Bolts were collected
114 from three trees at each of three sites in Hinton, Alberta (53° 20.530 117° 35.208, 53° 22.825
115 117° 32.561 and 53° 16.527 117° 39.916) in June 2018, and from two trees at each of two sites
116 in Slave Lake, Alberta (54° 51.751 115° 09.751 and 54° 53.842 115° 08.708) in November 2017.
117 The localities were chosen to ensure that beetles collected were in the epidemic population range
118 of Alberta. Only mass attacked trees (>40 attacks per m²) that were larger than 27 cm diameter at
119 breast height were felled. Two, 50-cm bolts from each tree, removed from 1-2 m above the
120 ground were transported to the University of Alberta. Cut ends of the bolts were sealed with
121 paraffin wax (parowax®) to minimize desiccation, and bolts were stored at 5°C until July 2018
122 when bioassays were conducted.

123 When beetles were needed for bioassays, bolts were removed from cold storage and
124 placed in 121 L emergence bins fitted with a glass jar. Mountain pine beetles are positively
125 phototactic and when they emerge from bolts they follow the light towards the glass jar where
126 they are collected. Bins were housed at 21°C under a 16:8 h light:dark cycle. Emerging beetles
127 caught in the glass jars were collected daily, separated by sex, labelled, and placed in 1.5 mL
128 microcentrifuge tubes with a small strip of paper to hold onto (Evenden et al., 2014). Beetles
129 were stored at 4°C before use in the bioassay at 3-5 days post emergence from the bolt.

130 ***Flight mills***

131 Flight on flight mills was used as an experimental treatment to assess the impact of flight on
132 subsequent host colonization and pheromone production (Fig. 1). Beetles (3-5 days post
133 emergence) were weighed to the nearest 0.01 mg (Mettler Toledo XPE205 Microbalance,
134 Columbus, OH, USA). Beetles were assigned randomly to one of two treatments: 23 h flight
135 period (flown), or 23 h without the opportunity to fly (control). Beetles in the flown treatment
136 were tethered using a 2 cm long, 30 gauge aluminum wire (0.02 mm diam.) with a small loop at
137 the end. The loop was attached to the pronotum of each beetle using Press-Tite Contact Cement
138 (LePage, Mississauga, ON, CAN) so that elytra movement was not restricted. Twenty-two
139 tethered beetles were positioned on flight mills on each of 13 days, and given the opportunity to
140 fly during the 23 h treatment period. Control beetles were housed with a piece of paper in
141 perforated 1.5 mL microcentrifuge tubes in the flight mill room during the treatment period. The
142 flight mill room was kept at 23°C with a 16:8 h light to dark cycle. The distal end of each tether
143 was attached to the flight mill arm at a ~100° angle using a small piece of wire insulation. Light
144 (550 lux) was provided by high flicker frequency fluorescent bulbs (Evenden, Whitehouse &
145 Sykes, 2014).

146 A small magnetic transmitter positioned on the flight mill arm detected the arm rotation
147 propelled by beetle flight. The transmitter directed the signal to the attached computer. LabView
148 software (National Instruments Corporation, Austin, TX, USA) measured each revolution of the
149 flight mill arm (94.4 cm in circumference). Output included the duration and number of
150 revolutions for each flight burst initiated by the beetle. Total flight distance and duration, as well
151 as flight velocity and number of flight bursts were calculated from this output.

152 After the 23 h treatment period, the tether was removed from each flown beetle, and both
153 flown and control beetles were weighed to the nearest 0.01 mg. Beetles that died or became
154 detached from tethers during flight treatment were not included in the subsequent bioassays or
155 statistical analyses.

156 ***Inoculation material***

157 In July 2018, three uninfested lodgepole pine trees were felled at each of three sites (53° 20.530
158 117° 35.208, 53° 22.825 117° 32.561 and 53° 16.527 117° 39.916) in Hinton, Alberta. Trees

159 were chosen based on size and overall appearance; only those that were healthy looking (ie.
160 green needles and no large wounds) and larger than 27 cm diameter at breast height were felled.
161 From each tree, three 50-cm bolts were harvested between 1-2.5 m above the ground. Bolts were
162 transported to the University of Alberta, where the cut ends of each bolt were sealed with
163 paraffin wax and stored at 5°C until Aug 2018 when needed for bioassays.

164 *Host colonisation experiment*

165 The first experiment tested the hypothesis that flight treatment influences subsequent host
166 colonisation behaviour by female mountain pine beetle (Fig. 1). Host colonisation was measured
167 as capacity to enter lodgepole pine bolts and the time taken for successful host entry. Uninfested
168 bolts were removed from cold storage 24 h prior to beetle inoculation. Ten clear plastic cups (30
169 mL) were positioned 10 cm from the bottom of the bolt and secured with flagging tape. A
170 charcoal filter (Paasche Charcoal Filter, WY, USA) skirt was placed between the bolt and the
171 cup to fill any gaps.

172 Immediately following flight treatment and measurement of post-treatment weight, each
173 female beetle was introduced to one of ten individual cups positioned on a lodgepole pine bolt.
174 Flown and control beetles were placed in alternating order on each bolt. Beetle activity was
175 monitored for 72 h following the initial placement in the cup or until host entry or death. Boring
176 dust within the cup indicated host entry. Data for beetles that escaped from the cups (34 flown
177 beetles and 33 control beetles escaped) were removed.

178 *Pheromone production experiment*

179 A second experiment tested the hypothesis that flight treatment affects pheromone production by
180 mountain pine beetles following successful host entry (Fig. 1). A subset of female beetles, from
181 both treatment groups flown (n=12) and control (n=9), that entered host material within 24 h of
182 inoculation were used in aeration bioassays to measure semiochemicals released by the beetles.

183 A single flown (n=11) and control (n=7) male was introduced into galleries of individual
184 females 24 h after females were introduced to cups. Males were flown the day after females and
185 introduced to the bolts in a different manner. The bark was peeled back slightly around the
186 female entrance hole and boring dust was blown away to reveal the exact point of entrance.

187 Males were gently pushed into the female entrance hole. Once the male was firmly positioned
188 within the entrance hole, the set-up described below was assembled for aeration.

189 Aeration was conducted using the methods described in Erbilgin et al. (2014). Once
190 female beetles entered the bolt, the clear plastic cup was removed, and replaced with a glass
191 funnel (DWK Life Sciences Kimble K2895045, 45 mm diameter, 50 mm stem). The glass funnel
192 was positioned over a charcoal filter skirt pressed tightly against the bolt and secured with
193 flagging tape. The stem of the glass funnel was connected to a small, 10 cm portion of PTFE
194 tubing (Cole-Parmer, 3/16" x 1/4", RK-06605-32). A second piece of PTFE tubing was attached
195 to PVC tubing (Fisherbrand, 3/16" inner diameter, 1/16" wall) that was subsequently connected
196 to a laboratory bench vacuum. To collect the semiochemicals released by beetles, Porapak Q
197 tubes (6 x 110-mm, 2 sections: 75/150 mg sorbent, 20/40 mesh) were inserted between the two
198 portions of PTFE tubing. Over a 4 h duration, the vacuum pulled air ($100 \text{ mL} \cdot \text{min}^{-1}$) over the site
199 of beetle entry to trap semiochemicals produced by the beetle pair into the attached Porapak Q
200 tube. After the 4 h aeration, Porapak Q tubes were removed from the PTFE tubing and were
201 capped, wrapped in tinfoil, and stored at -80°C until extraction. Repeated aerations measured
202 pheromone production at 12, 24, 36, 48, 72, 96 and 120 h after introduction of females to cups.
203 Males were introduced 24 h after females, so the 12 h time point contained emissions from
204 females only; the subsequent collections were conducted on beetle pairs.

205 *Chemical extraction & analyses*

206 Each Porapak Q tube from each aeration sample was scored with a glass cutter to remove the
207 adsorbent beads from the tube into a 2 mL Axygen microtube that was placed onto dry ice. The
208 stock solution of the extraction solvent contained 500 mL DCM (dichloromethane, HPLC Grade,
209 Fisher Scientific, USA) with 5 μl of heptyl acetate (purity >98%, Sigma-Aldrich, USA) to act as
210 an internal standard. One mL of the stock solution was dispensed (0.5-5 mL dispenser,
211 Dispensette Organic, Eppendorf, GER) into each 2 mL microtube containing adsorbent material
212 from each sample. Microtubes containing adsorbent material and stock solution were vortexed
213 for 30 sec at maximum speed (3000) (VWR Pulsing Vortex Mixer) and were then placed into a
214 sonicator (Symphony) for 10 min. Microtubes were centrifuged for 15 min at 0°C at 16100 rcf
215 (Eppendorf AG 2231, GER) to create two phases. Dichloromethane with the extract was
216 collected from the lower phase.

217 To filter the extract, the solvent solution was pipetted into a modified pipette (Fisher
218 Scientific, borosilicate glass, 13-67-20A) containing a small amount of glass wool to act as a
219 filter. Filtered extract was collected in 2 mL Autosampler vials (Fisher Scientific, 9 mm/Amber-
220 ID, 03-391-9) that were capped (Autosampler caps, 9 mm screw thread/PTFE/Silicone, 03-391-
221 14) and stored at -40°C until chemical analyses.

222 Chemical analyses were performed using a Gas Chromatograph/Mass Spectrometer
223 (GC/MS, Agilent 7890A/5975C, Agilent Technologies, CA, USA) with a HP-CHIRAL-20 β
224 column (I.D. 0.25 mm, length 30 m, Agilent Tech.). Helium was the carrier gas with a flow rate
225 of 1 mL·min⁻¹. Two μ l samples from each extract were injected in a Pulsed Splitless mode. The
226 oven temperature started at 45°C for 2 min, increased to 70°C by 20°C·min⁻¹, increased to 90°C
227 by 10°C·min⁻¹, increased to 120°C by 2°C·min⁻¹, increased to 150°C by 3°C·min⁻¹, and then
228 increased up to 230°C by 30°C·min⁻¹ and held for 1 min. The data was acquired using both SIM
229 and SCAN modes. SCAN mode was conducted to identify the compounds of interest, whereas
230 SIM mode was used to quantify the collected data. The quantified compounds included (1) *trans*-
231 verbenol; and (2) *exo*-brevicommin. Compounds were quantified by comparison with
232 commercially available standards with a chemical purity > 99% (Contech Enterprises Inc., BC,
233 CAN).

234 ***Statistical Analyses***

235 All data analyses were performed in R version 3.4.1 (R Core Team, 2018). The explanatory
236 variable, percent weight lost during the flight treatment, was calculated by dividing the
237 difference between pre- and post-treatment weight by pre-treatment weight, and multiplying this
238 value by 100. Data were tested for normality and heteroscedasticity using visual techniques and
239 Shapiro-Wilks test. Due to the confounding nature of the variables (percent weight lost, pre-
240 treatment weight and distance flown) the effects of these independent factors were analyzed in
241 separate models to avoid spurious associations.

242 The effect of flight treatment on female beetle host acceptance was analysed using a
243 contingency table. Dichotomous entry data in the host colonisation experiment was analysed
244 using a binomial distribution in a generalized linear mixed effects model with natal bolt and
245 reproductive bolt defined as random factors in each model. The response variable, host entry,
246 was assessed in three separate models, (1) host entry explained by percent weight lost by both

247 flown and control female beetles, during the flight period; (2) host entry explained by distance
248 flown by female beetles during the flight period; and (3) host entry explained by pre-treatment
249 weight of both flown and control female beetles. For model 1, percent weight lost was square-
250 root transformed to meet the assumption of normality; for model 2, distance flown was
251 transformed to the fourth root to meet the assumption of normality. Cox proportional models are
252 regression models used to determine the relationship between survival time and predictor
253 variables. In the case of this study, the “survival” term was defined by entry success and time
254 until entry. Thus, instead of the “survival” term representing the length of time until death, it
255 represented the length of time until host entry. Four cox proportional models were used to
256 analyze entry success and time until host entry in relation to (1) square-root transformed percent
257 weight lost for all beetles; (2) percent weight lost for flown beetles; (3) fourth-root transformed
258 distance flown; and (4) pre-treatment weight for all beetles. For the beetles that entered, the
259 relationship between time until entry and percent weight lost was analysed using a mixed effects
260 linear model separately for flown and control beetles. Since multiple models were used to
261 analyse these groups separately, a Bonferroni correction of $\alpha = 0.025$ was applied to determine
262 significance. Both natal and reproductive hosts were included as random factors in both analyses.

263 Pheromone production data was analysed using linear mixed effects models with natal
264 bolt and reproductive bolt defined as random factors in each model. The response variable, total
265 *trans*-verbenol production across all aeration timepoints, was assessed in three separate models
266 (1) *trans*-verbenol production as explained by percent weight lost during treatment, by both
267 flown and control female beetles; (2) *trans*-verbenol production as explained by distance flown
268 by female beetles during the flight period; and (3) *trans*-verbenol production as explained by pre-
269 treatment weight of both flown and control female beetles. For models 1 and 3, total *trans*-
270 verbenol production was cube-root transformed to meet the assumption of normality. The
271 response variable, total *exo*-brevicomin production across all aeration periods, was assessed in
272 three separate models (1) *exo*-brevicomin production as explained by percent weight lost during
273 treatment, by both flown and control male beetles; (2) *exo*-brevicomin production as explained
274 by distance flown by male beetles during the flight period, and (3) *exo*-brevicomin production as
275 explained by pre-treatment weight of both flown and control male beetles.

276 **Results**

277 ***Host colonisation experiment***

278 Beetles placed on flight mills flew an average of 4.02 ± 0.54 km over the 23 h period (Fig. 2).
279 The minimum flight distance was 0.002 km and the maximum flight distance was 22.26 km. Of
280 the 267 flown and control female beetles used in the host colonisation study, 40% entered the
281 host material within 72 h. Initiation of host colonisation was influenced by flight treatment.
282 Beetles that flew on flight mills were 13% less likely to initiate host colonisation compared to
283 unflown control beetles ($\chi^2=5.2722$, $p=0.0216$).

284 Generalized linear models indicated a negative relationship between host entry and the
285 percent weight lost during the flight treatment ($\chi^2=31.774$, $p=1.732 \times 10^{-8}$). Female beetles that
286 lost less weight during flight treatment were more likely to enter a host (Fig. 3). No relationships
287 between host entry and distance flown ($\chi^2=0.0763$, $p=0.7824$) or pre-treatment weight
288 ($\chi^2=0.5286$, $p=0.4672$) were found.

289 Cox proportional models showed that percent weight lost affected host entry and entry
290 time for all beetles ($Z=6.264$, $p=3.74 \times 10^{-10}$) and flown beetles alone ($Z=2.184$, $p=0.029$, Fig. 3).
291 There was no relationship, however, between distance flown ($\chi^2=0.408$, $p=0.683$) or pre-
292 treatment weight ($\chi^2=0.704$, $p=0.4820$) and host entry. Of the beetles that entered the bolts, the
293 time until entry was negatively influenced by the percent weight lost during the treatment in
294 flown ($\chi^2=7.0248$, $p=0.0080$, Fig. 4) but not unflown control ($\chi^2=0.0093$, $p=0.923$) beetles.

295 ***Pheromone production experiment***

296 The production of *trans*-verbenol by female beetles was influenced by the percent weight lost
297 during flight treatment ($\chi^2= 3.8706$, $p=0.04914$) and the distance flown ($\chi^2=5.1584$, $p=0.0231$),
298 but not by pre-flight weight ($\chi^2=1.1417$, $p=0.2853$). Females that lost more weight and flew
299 further distances produced more *trans*-verbenol (Fig. 5).

300 The production of *exo*-brevicomin by male beetles was influenced by pre-flight treatment
301 weight ($\chi^2=5.6937$, $p= 0.0170$) and distance flown ($\chi^2=9.5932$, $p=0.0020$), but not by percent
302 weight lost during flight ($\chi^2=0.9912$, $p=0.3195$). Males that weighed more prior to flight
303 treatment produced more *exo*-brevicomin; males that flew further produced less *exo*-brevicomin
304 (Fig. 6).

305 Discussion

306 The important life history traits of adult mountain pine beetle include dispersal from the natal
307 host, host colonisation, aggregation triggered by pheromone production, and reproduction after
308 overcoming host defenses. The current study uncovers mechanisms by which energy-use during
309 flight influences host entry and pheromone production by beetles. The amount of lipids retained
310 by females following flight dictates the outcome of host colonisation success (Chubatý et al.,
311 2014). In the current study, female beetles that lost less than 10% of their body weight during
312 flight were more likely to enter hosts compared to those that lost more than 10%. In mountain
313 pine beetle, weight loss is linked to lipid metabolism during flight (Evenden et al., 2014). Our
314 findings are in agreement with the results of earlier studies on male pine engraver beetles (*Ips*
315 *pini*) in which beetles that enter host material have 21% more lipid compared to those that do not
316 enter (Wallin & Raffa, 2000). Certain silvicultural treatments, like stand thinning techniques, can
317 increase flight distance before host colonisation in managed stands. Mountain pine beetle was
318 detected in high numbers in thinned stands (Schmitz et al., 1988) as well as in clear cut stands
319 (Reid, 2008). In these stands, beetles are forced to fly further distances before host colonisation,
320 which could impact the number of successful attacks on trees.

321 The speed of host colonisation is also dependent on energy reserves remaining in female
322 beetles after dispersal. We found that the fastest beetles to enter the host had lost the most weight
323 during the flight treatment. This result indicates a trade-off between energy-use during flight and
324 host acceptance in female mountain pine beetle, which likely intensifies the flight-reproduction
325 trade-off previously suggested for this species (Wijerathna et al., 2019). These results lend
326 further support to the “desperation hypothesis” (Latty & Reid, 2010). In contrast to our findings,
327 time to host entry by pine engraver beetles declined with beetle starvation (Wallin & Raffa,
328 2002), suggesting that energy-use trade-offs may not be consistent across bark beetle species.

329 Distant dispersal away from the natal tree may increase the need for effective signalling
330 to attract conspecifics to mount a mass attack. We show that female flight distance and energy-
331 use is linked to a subsequent increase in *trans*-verbenol production by females following host
332 entry. Release of high concentrations of *trans*-verbenol should increase the success of pioneer
333 beetles that initiate attack on distant hosts to increase the aggregation of conspecifics (Erbilgin et
334 al., 2014). Similarly, attraction of a sister species *Dendroctonus frontalis* increases positively

335 with *trans*-verbenol dose (Shepherd & Sullivan, 2019). Beetles that disperse only a short
336 distance would benefit less from the production of strong pheromone signals.

337 Female mountain pine beetle release *trans*-verbenol upon initiation of gallery
338 construction and feeding (Pureswaran et al., 2000). *trans*-Verbenol production requires the
339 oxidation of the precursor, α -pinene (Hughes, 1975), obtained from the natal host (Chiu et al.,
340 2018). Additionally, *trans*-verbenol production varies with the concentration of α -pinene present
341 in the reproductive host (Taft et al., 2015), which suggests that the α -pinene necessary for
342 pheromone synthesis could be obtained from both sources. Female mountain pine beetles
343 accumulate α -pinene in the form of monoterpenyl esters which are fatty acid esters stored in the
344 fat body (Chiu et al., 2018). As we have shown that flight increases *trans*-verbenol production in
345 female mountain pine beetles, the biochemical mechanism dictating this increase may be the
346 result of lipid-use during flight through impact on the stored monoterpenyl esters. High
347 variability in pheromone production, including *trans*-verbenol, occurs in other bark beetles
348 (Pureswaran et al., 2008). Variation in pheromone production can also be linked to body size
349 (Pureswaran & Borden, 2003) and genetics (Domingue & Teale, 2008), but causes of variation
350 differ with pheromone component identity.

351 Flight distance negatively affected *exo*-brevicommin production by males. This difference
352 in pheromone production in response to flight between sexes could be due to the timing of
353 pheromone production. Males can begin to produce *exo*-brevicommin immediately upon
354 emergence from the natal host (Song et al., 2014). The complete biosynthetic pathway behind the
355 production of *exo*-brevicommin remains unknown; however, it is synthesized *de novo* from fatty
356 acyl-CoA precursors and stored in the fat body (Song et al., 2014). Energy-use during flight
357 could influence *exo*-brevicommin storage in the fat body, with more pheromone released during
358 periods of flight than rest. This may explain why males produce low levels of *exo*-brevicommin
359 when they enter the female nuptial galleries to reproduce (Song et al., 2014). These low levels of
360 *exo*-brevicommin are likely adaptive in mediation of aggregation behaviour as low concentrations
361 of *exo*-brevicommin are more attractive to conspecifics than higher concentrations (Rudinsky et
362 al., 1974). Flight may promote the release of low, attractive quantities of *exo*-brevicommin. Males
363 potentially have a finite amount of pheromone to release based on the condition of the beetle at
364 the time of pupation. Our finding that heavier males produce more *exo*-brevicommin than lighter

365 males supports this idea. The quality of the natal host likely has a large influence on the amount
366 of *exo*-brevicomins males can produce in a lifetime, as good quality hosts produce larger, more
367 robust offspring (Graf et al., 2012). This supports previous findings indicating a marginal link
368 between mountain pine beetle body weight and length to *exo*-brevicomins production
369 (Pureswaran & Borden, 2003). Heavier males fly further than lighter males (Evenden et al.,
370 2014), this difference in flight behaviour could promote optimal levels of *exo*-brevicomins release
371 at the reproductive host.

372 Interestingly, weight loss during flight influences pheromone production in females but
373 not males. This is potentially due to differential energy-use during flight between the sexes.
374 Females rely heavily upon lipids during long distance flight, while males use both lipids and
375 proteins to power flight (Wijerathna & Evenden, 2019). This is likely driven by variation in the
376 energy needed for host colonisation, as females require proteins for reproduction (Pitt et al.,
377 2014). The reliance on lipids by female beetles for flight likely has a direct impact on weight loss
378 during flight (Evenden et al., 2014), whereas weight loss by male beetles is a combination of the
379 depletion of multiple energy sources (Wijerathna & Evenden 2019). If lipid-use is responsible
380 for differing pheromone titres, the link between weight loss and pheromone production in males
381 would be lost. In the fat body, female beetles store monoterpenyl esters used in the production of
382 *trans*-verbenol (Chiu et al., 2018), while male beetles store *exo*-brevicomins in its final form in
383 the fat body (Song et al., 2014). Lipid-use during flight may allow for the release of pheromone
384 from storage in males and reduce the subsequent pheromone titre available to males calling at the
385 new host. In females, since the entire biosynthetic pathway of *trans*-verbenol remains unknown,
386 all that can be concluded is that flight could aid or promote biosynthesis of this compound.

387 Here we provide evidence for possible mechanisms that drive flight polyphenisms in bark
388 beetles. The trade-off between energy-use during flight and host colonisation could select for
389 short distance dispersal so that beetles have enough energy to successfully colonise their
390 reproductive host. Alternatively, long distance dispersal might be adaptive for outbreeding and
391 access to high quality and abundant hosts (Raffa et al. 1993). Energy-use during flight positively
392 impacts subsequent pheromone production in the pioneering female beetles. Increased *trans*-
393 verbenol production will aid beetles in mediating mass attacks at distant hosts, this in
394 combination with other benefits at these distant locations will select for long-distance dispersers.

395 These results provide evidence for mechanisms that promote contrasting selection on flight in
396 bark beetles. Selection at both ends of the polyphenism spectrum maintains high dispersal
397 variability within populations. This intraspecific variation in dispersal strategies promotes an
398 evolutionarily stable strategy for bark beetle populations (Kautz et al., 2016). Understanding
399 variation in spatial movement of bark beetles across landscapes will help to predict future
400 population spread of these aggressive tree pests.

401 **Acknowledgements**

402 We thank Caroline Whitehouse, Andrea Sharpe and Jenn McCormick from Alberta
403 Agriculture and Forestry for their assistance obtaining both mountain pine beetle-infested
404 material and clean host material for experimental use. We would like to thank Victor Shegelski
405 for his time and effort felling trees for this project and Ronald Batallas for his advice on
406 statistical analyses. This research was supported by a grant to from the Natural Science and
407 Engineering Research Council of Canada (grant no. NET GP 434810-12) to the TRIA Network,
408 with contributions from Alberta Agriculture and Forestry, fRI Research, Manitoba Conservation
409 and Water Stewardship, Natural Resources Canada - Canadian Forest Service, Northwest
410 Territories Environment and Natural Resources, Ontario Ministry of Natural Resources and
411 Forestry, Saskatchewan Ministry of Environment, West Fraser and Weyerhaeuser (Maya
412 Evenden and Nadir Erbilgin, co-PIs). Additionally, the Natural Sciences and Engineering
413 Research Council of Canada–Discovery to NE. Research stipends to support Kelsey Jones were
414 provided by teaching assistantships at the University of Alberta, and various scholarships
415 (NSERC CGS-M Alexander Graham Bell, Queen Elizabeth II Graduate Scholarship, Walter H.
416 John’s Graduate Fellowship, Julia O. Hrapko Scholarship).

417 **References**

- 418 **Blomquist, G.J., Figueroa-Teran, R., Aw, M., Song, M., Gorzalski, A., Abbott, N.L., Chang,**
419 **E., and Tittiger, C.** (2010). Pheromone production in bark beetles. *Insect Biochem. Molec.*
420 **40**(10), 699-712.
- 421 **Byers, J.A.** (1999). Effects of attraction radius and flight paths on catch of scolytid beetles
422 dispersing outward through rings of pheromone traps. *J. Chem. Ecol.* **25**(5), 985-1005.

- 423 **Chen H., and Walton A.** (2011). Mountain pine beetle dispersal: spatiotemporal patterns and
424 role in the spread and expansion of the present outbreak. *Ecosphere*. **2**(6), Article 66.
425 <https://doi.org/10.1890/ES10-00172.1>
- 426 **Chiu, C.C., Keeling, C.I., and Bohlmann, J.** (2018). Monoterpenyl esters in juvenile mountain
427 pine beetles and sex-specific release of the aggregation pheromone *trans*-verbenol. *P.*
428 *Natal. Acad. Sci. USA*. **115**(14), 3652-3657.
- 429 **Chubaty, A.M., Hart, M., and Rotiberg, D.** (2014). ‘To tree or not to tree’: the role of energy
430 limitation on host tree acceptance in a bark beetle. *Evol. Ecol. Res.* **16**, 337-349.
- 431 **Cisper, G., Zera, A.J., and Borst, D.W.** (2000). Juvenile hormone titer and morph-specific
432 reproduction in the wing-polymorphic cricket, *Gryllus firmus*. *J. Insect Physiol.* **46**, 585-
433 596.
- 434 **Cullingham, C.I., Cooke, J.E.K., Dang, S., Davis, C.S., Cooke, B.J., and Coltman, D.W.**
435 (2011). Mountain pine beetle host-range expansion threatens the boreal forest. *Mol. Ecol.*
436 **20**, 2157-2171.
- 437 **Domingue, M. J., and Teale, S.A.** (2008). The genetic architecture of pheromone production
438 between populations distant from the hybrid zone of the pine engraver, *Ips pini*.
439 *Chemoecology*. **17**, 255-262.
- 440 **Elkin, M., and Reid, M.L.** (2005). Low energy reserves and energy allocation decisions affect
441 reproduction by mountain pine beetles, *Dendroctonus ponderosae*. *Funct. Ecol.* **19**(1), 102-
442 109.
- 443 **Erbilgin, N., Ma, C., Whitehouse, C., Shan, B., Najjar, A., and Evenden, M.** (2014).
444 Chemical similarity between historical and novel host plants promotes range and host
445 expansion of the mountain pine beetle in naïve host ecosystem. *New Phytol.* **201**, 940-950.
- 446 **Evenden, M.L., Whitehouse, C.M., and Sykes, J.** (2014). Factors influencing flight capacity of
447 the mountain pine beetle (Coleoptera: Curculionidae: Scolytinae). *Environ. Entomol.* **43**(1),
448 187-196.
- 449 **Gitau, C.W., Bashford, R., Carnegie, A.J., and Gurr, G.M.** (2013). A review of
450 semiochemicals associated with bark beetle (Coleoptera: Curculionidae: Scolytinae) pests

451 of coniferous trees: a focus on beetle interactions with other pests and their associates.
452 *Forest Ecol. Manag.* **297**, 1-14.

453 **Graf, M., Reid, M.L., Aukema, B.H., and Lindgren, B.S.** (2012). Association of tree diameter
454 with body size and lipid content of mountain pine beetles. *Can. Entomol.* **144**, 467-477.

455 **Gray, B., Billings, R.F., Gara, R.I., and Johnsey, R.L.** (1972). On the emergence and initial
456 flight behaviour of the mountain pine beetle, *Dendroctonus ponderosae*, in Eastern
457 Washington. *Z. Angew. Entomol.* **71**, 250-259.

458 **Hughes, P.R.** (1975). Pheromones of *Dendroctonus*: Origin of α -pinene oxidation products
459 present in emergent adults. *J. Insect Physiol.* **21**(3), 687-691.

460 **Karlsson, B., and Johansson, A.** (2008). Seasonal polyphenism and developmental trade-offs
461 between flight ability and egg laying in a pierid butterfly. *Proc. R. Entomol. Soc. A.* **275**,
462 2131-2136.

463 **Karlsson, B., Stjernolm, F., and Wiklund, C.** (2008). Test of a developmental trade-off in a
464 polyphenic butterfly: direct development favours reproductive output. *Funct. Ecol.* **22**,
465 121-126. <https://doi.org/10.1111/j.1365-2435.2007.01334.x>

466 **Kautz, M., Imron, M.A., Dworschak, K., and Schopf, R.** (2016). Dispersal variability and
467 associated population-level consequences in tree-killing bark beetles. *Movement Ecol.* **4**, 9.
468 <https://doi.org/10.1186/s40462-016-0074-9>

469 **Kopp, M., and Tollrain, R.** (2003). Trophic size polyphenism in *Lembadion bullnium*: costs
470 and benefits of an inducible offense. *Ecology.* **84**(3), 641-651.

471 **Latty, T.M., and Reid, M.L.** (2009). First in line or first in time? Effects of settlement order and
472 arrival date on reproduction in a group-living beetle *Dendroctonus ponderosae*. *J. Anim.*
473 *Ecol.* **78**, 549-555.

474 **Latty, T.M., and Reid, M.L.** (2010). Who goes first? Condition and danger dependent
475 pioneering in a group-killing bark beetle (*Dendroctonus ponderosae*). *Behav. Ecol.*
476 *Sociobiol.* **64**, 639-646.

477 **Pitt, C., Robert, J.A., Bonnett, T.R., Keeling, C.I., Bohlmann, J., and Huber, D.P.W.**
478 (2014). Proteomics indicators of the rapidly shifting physiology from whole mountain pine
479 beetle, *Dendroctonus ponderosae* (Coleoptera: Curculioidea), adults during early host
480 colonization. *PLOS One*. **9**(10), e110673.

481 **Pureswaran, D.S., and Borden, J.H.** (2003). Is bigger better? Size and pheromone production
482 in the mountain pine beetle, *Dendroctonus ponderosae* Hopkins (Coleoptera: Scolytidae).
483 *J. Insect Behav.* **16**(6), 765-782.

484 **Pureswaran, D.S., Gries, R., Boren, J.H., and Pierce Jr., H.D.** (2000). Dynamics of
485 pheromone production and communication in the mountain pine beetle, *Dendroctonus*
486 *ponderosae* Hopkins, and the pine engraver, *Ips pini* (Say) (Coleoptera: Scolytidae).
487 *Chemoecology*. **10**, 153-168.

488 **Pureswaran, D.S., Sullivan, B.T., and Ayres, M.P.** (2006). Fitness consequences of
489 pheromone production and host selection in a tree-killing bark beetle (Coleoptera:
490 Curculionidae: Scolytinae). *Oecologia*. **148**, 720-728.

491 **Pureswaran, D.S., Sullivan, B.T., and Ayres, M.P.** (2008). High individual variation in
492 pheromone production by tree-killing bark beetles (Coleoptera: Curculionidae: Scolytinae).
493 *Naturwissenschaften*. **95**, 33-44.

494 **R Core Team.** (2018). R: A Language and Environment for Statistical Computing. R
495 Foundation for Statistical Computing, Vienna. <https://www.R-project.org>

496 **Raffa, K.** (2001). Mixed messages across multiple trophic levels: the ecology of bark beetle
497 chemical communication systems. *Chemoecology*. **11**, 49-65.

498 **Raffa, K.F., and Berryman, A.A.** (1982). Physiological differences between lodgepole pines
499 resistant and susceptible to the mountain pine beetle and associated microorganisms.
500 *Environ. Entomol.* **11**(2), 486-492.

501 **Raffa, K. F., Phillips, T. W., and Salom, S. M.** (1993). Strategies and mechanisms of host
502 colonization by bark beetles. In Schowalter, T.D. and Filip, G.M. (Eds.) *Beetle Pathogen*
503 *Interactions in Conifer Forests* (pp. 103-128). Academic Press, London.

- 504 **Raffa, K.F., Aukema, B.H., Erbilgin, N., Klepzig, K.D., and Wallin, K.F.** (2005). Interactions
505 among conifer terpenoids and bark beetles across multiple levels of scale: an attempt to
506 understand links between population patterns and physiological processes. *Recent Adv.*
507 *Phytochem.* **39**, 79-118.
- 508 **Raffa, K.F., Aukema, B.H., Bentz, B.J., Carroll, A.L., Hicke, J.A., Turner, M.G., and**
509 **Romme, W.H.** (2008). Cross-scale drivers of natural disturbances prone to anthropogenic
510 amplification: the dynamics of bark beetle eruptions. *BioScience.* **58**(6), 501-517.
- 511 **Raffa, K.F., Grégoire, J.C., and Lindgren, B.S.** (2015). Natural history and ecology of bark
512 beetles. In Vega, F.E., and Hofstetter, R.W. (Eds.) *Bark beetles: Biology and ecology of*
513 *native and invasive species* (pp. 1-40). Academic Press, Elsevier.
- 514 **Reid, M.L., Sekhon, J.K., and LaFramboise, L.M.** (2017). Toxicity of monoterpene structure,
515 diversity and concentration to mountain pine beetles, *Dendroctonus ponderosae*: beetle
516 traits matter more. *J. Chem. Ecol.* **43**, 351-361.
- 517 **Robertson, C., Nelson, T.A., and Boots, B.** (2007). Mountain pine beetle dispersal: The spatial-
518 temporal interaction of infestations. *Forest Sci.* **53**(3), 395-405.
- 519 **Roff, D.A., and Fairbairn, D.J.** (1991). Wing dimorphisms and the evolution of migratory
520 polymorphisms among the Insecta. *Am. Zool.* **31**, 243-251.
- 521 **Rudinsky, J.A., Morgan, M.E., Libbey, L.M., and Putnam, T.B.** (1974). Antiaggregative-
522 rivalry pheromone of the mountain pine beetle, and a new arrestant of the southern pine
523 beetle. *Environ. Entomol.* **3**(1), 90-98.
- 524 **Safranyik, L., and Carroll, A.L.** (2006). The biology and epidemiology of the mountain pine
525 beetle in lodgepole pine forests. In L. Safranyik and B. Wilson (Eds.) *The mountain pine*
526 *beetle a synthesis of biology, management, and impacts on lodgepole pine* (pp. 3-66).
527 Pacific Forestry Centre, Victoria.
- 528 **Safranyik, L., Carroll, A.L., Régnière, J., Langor, D.W., Riel, W.G., Shore, T.L., Peter, B.,**
529 **Cooke, B.J., Nealis, V.G., and Taylor, S.W.** (2010). Potential for range expansion of the
530 mountain pine beetle into the boreal forest of North America. *Can. Entomol.* **142**, 415-442.

531 **Shegelski, V.A., Evenden, M.L., and Sperling, F.A.H.** (2019). Morphological variation
532 associated with dispersal capacity in tree-killing bark beetle *Dendroctonus ponderosae*
533 Hopkins. *Agric. For. Entomol.* **21**(1), 79-87.

534 **Shepherd, W.P., and Sullivan, B.T.** (2019). Southern pine beetle (Coleoptera: Curculionidae:
535 Scolytinae) pheromone component *trans*-verbenol: enantiomeric specificity and potential
536 as a lure adjuvant. *Environ. Entomol.* **48**(1), 193-201. <https://doi.org/10.1093/ee/nvy169>

537 **Simpson, S.J., Sword, G.A., and Lo, N.** (2011). Polyphenism in insects. *Curr. Biol.* **21**(18),
538 R738-R749. <https://doi.org/10.1016/j.cub.2011.06.006>

539 **Song, M., Gorzalski, A., Nguyen, T.T., Liu, X., Jeffery, C., Blomquist, G.J., and Tittiger, C.**
540 (2014). *exo*-Brevicommin biosynthesis in the fat body of the mountain pine beetle,
541 *Dendroctonus ponderosae*. *J. Chem. Ecol.* **40**, 181-189.

542 **Taft, S., Najar, A., and Erbilgin, N.** (2015). Pheromone production by an invasive bark beetle
543 varies with monoterpene composition of its naïve host. *J. Chem. Ecol.* **41**, 540-549.

544 **Thompson, S.S., and Bennett, R.B.** (1971). Oxidation of fat during flight of male Douglas-fir
545 beetles, *Dendroctonus pseudotsugae*. *J. Insect Physiol.* **17**, 1555-1563.

546 **Wallin, K.F., and Raffa, K.F.** (2000). Influences of host chemicals and internal physiology on
547 the multiple steps of postlanding host acceptance behaviour of *Ips pini* (Coleoptera:
548 Scolytidae). *Physiol. Chem. Ecol.* **29**, 442-453.

549 **Wallin, K.F., and Raffa, K.F.** (2002). Prior encounters modulate subsequent choices in host
550 acceptance behaviour by the bark beetle *Ips pini*. *Entomol. Exp. Appl.* **103**, 205-218.

551 **Wijerathna, A.N., and Evenden, M.L.** (2019). Energy use by the mountain pine beetle
552 (Coleoptera: Curculionidae: Scolytinae) for dispersal by flight. *Physiol. Entomol. In Press.*

553 **Wijerathna, A.N., Whitehouse, C.M., Proctor, H., and Evenden, M.L.** (2019). Testing for
554 trade-offs between flight and reproduction in the mountain pine beetle (Coleoptera:
555 Curculionidae) on two pine hosts. *Can. Entomol.* **151**, 298-310.

556 **Figures**

557 **Fig 1. Experimental set-up for host colonisation and aeration experiments.** For host
558 colonisation, beetles flown on flight mills for 23h were subsequently placed into small cups tied
559 to the side of a healthy bolt. Females that entered within the first 24h were paired with a flown
560 male and subject to the collection of pheromones in the aeration part of the bioassay.

561 **Fig 2. Histogram of flight distribution exhibited by females flown on flight mills for 23 h.**
562 The average flight distance was 4.02 ± 0.54 km, with a minimum flight distance of 0.002 km and
563 a maximum flight distance of 22.26 km (N=83).

564 **Fig 3. Histograms of percent weight lost during flight for female mountain pine beetle**
565 **(*Dendroctonus ponderosae*) that entered hosts (A) and failed to enter hosts (B).** Female
566 beetles that failed to enter hosts lost more weight on average than those that entered ($\chi^2=31.774$,
567 $p=1.732 \times 10^{-8}$) (N=83).

568 **Fig 4. Box plots of percent weight lost during the assay for flown female mountain pine**
569 **(*Dendroctonus ponderosae*) beetle that entered lodgepole pine hosts at different times post**
570 **inoculation.** The midline indicates the median and the bottom and top of the box represent the
571 25th and 75th percentiles, respectively. Vertical lines extending from the box (whiskers) represent
572 the maximum and minimum values. Beetles that entered host material (green bars, n=49) lost
573 less weight during the flight treatment compared to those that subsequently failed to enter hosts
574 (yellow bar, n=34) ($Z=2.184$, $p=0.029$). Weight lost after flight influenced the length of time it
575 took beetles to initiate colonisation after flight ($\chi^2=7.0248$, $p=0.0080$) (green bars).

576 **Fig 5. The relationship between percent weight lost during (A), flight distance (B) and**
577 **subsequent *trans*-verbenol ($\mu\text{g/ml}$) production for both flown (n=12) and control (n=9)**
578 **female mountain pine beetle (*Dendroctonus ponderosae*) in lodgepole pine bolts.** Beetles that
579 lost more body weight during the assay produced higher amounts of *trans*-verbenol ($\chi^2= 3.8706$,
580 $p=0.0491$). Flight promoted *trans*-verbenol production in female beetles ($\chi^2=5.1584$, $p=0.0231$).

581 **Fig 6. The relationship between pre-bioassay weight (A), flight distance (B) and subsequent**
582 ***exo*-brevicomin production ($\mu\text{g/ml}$) for flown (n=11) and control (n=7) male mountain pine**
583 **beetle (*Dendroctonus ponderosae*) in lodgepole pine bolts.** Heavier beetles produced more *exo*-

584 brevicomin ($\chi^2=5.6937$, $p=0.0170$). Flight distance is negatively associated with *exo*-brevicomin
585 production in male beetles ($\chi^2=9.5932$, $p=0.002$).

586