

Changes in Alberta's Grasslands Soil pH by Adopting the Adaptive Multi-Paddock Grazing System

Leah Stachniak, Laio Silva Sobrinho, Sarah Breitreuz, Scott Chang

Department of Renewable Resources, Faculty of Agriculture, Life, and Environmental Sciences, University of Alberta

Introduction

- Natural grasslands correspond to ~50% of Earth's terrestrial area and store ~30% of the world's soil organic carbon (SOC). As the largest terrestrial carbon sink, soil plays a vital role in carbon sequestration and in mitigating climate change (Dermer & Schuman, 2007).
- Despite its global significance, ~70% of Canada's original grassland habitats have been degraded as a consequence of poor land management (Gauthier & Wiken, 2003). As a result of this, the temperate grasslands are the most endangered ecosystem on the planet (Dodds et al., 2004).
- Fortunately, ranchers can sustain productivity, decrease the negative effects on the environment, improve animal performance, and help restore the grasslands by adopting the Adaptive Multi-Paddocks (AMP) system, an improved, innovative grazing practice [Fig. 1] (Teague et al., 2013).
- Under the AMP system, it has been found that there are less unwanted plants and barren ground as well as an increase in plant regrowth and pasture biomass production (Williams, 2017). The AMP system has also been found to increase SOC levels (Teague et al., 2013). Other soil properties, such as soil pH, have the potential to be impacted by implementing the AMP system.
- Since SOC has been shown to be correlated with soil pH, it might be used to indicate whether or not the AMP system affects SOC sequestration in AMP study areas (Jin & Wang, 2018).

Figure 1: AMP grazing system where the land is divided into multiple small paddocks. Cattle graze for shorter periods, are not allowed to consume >50% of the plant biomass, and are rotated more often.





Figure 2: Non-AMP (i.e. conventional) grazing system where cattle graze freely in a single, large, open paddock for a prolonged period of time until most of the grass has been consumed.

Objective

To study the effect of the AMP grazing system on grasslands soil pH within Alberta's different ecoregions and across the soil profile.

Methods

- Samples were collected in 12 study sites within Alberta and each site consisted of a pair of neighbouring AMP and Non-AMP managed ranches [Fig. 3]. From each ranch, 15 soil cores (1 m x 5 cm) were taken using a hydraulic soil probe.
- Each soil core was sectioned into 3-5 layers at various depth increments.
- After being weighed and dried, each core section was filtered with a 2 mm sieve to remove all visible roots and gravel, after which only soil remained.
- 5 g of soil from each soil section was added to 25 mL of milli-Q water (Kalra & Maynard, 1991).
- Soil solutions were mixed in a shaker machine for 30 minutes [Fig. 4].
- Using a digital pH meter, the pH of samples were analyzed [Fig. 5].




Figure 4: Shaker machine



Figure 5: pH meter



Results




Figure 6: Relationship between soil depth and soil pH from AMP and Non-AMP systems within Alberta.



pH of Soil Layers from Various Grasslands Ecoregions




Figure 7: Average soil pH at each depth increment in AMP and Non-AMP grazing systems within the ecoregions of Alberta's grasslands (i.e., Aspen Parkland, Boreal Transition, Fescue Grassland, and Mixed Grassland).

Conclusions

- Areas where error could have occurred resulting in an alteration of data:
 - Faulty or imprecise pH meter.
 - Incorrect data entry or data analysis.
 - AMP practice not being utilized long enough in order to have a significant effect.
- Trends that can be seen throughout the data:
 - Soil pH increases with increasing soil depth.
 - Across all soil depths, the AMP system seems to have slightly lower pH than Non-AMP.
 - Average soil pH seems to differ between the four ecoregions.
 - In both AMP and Non-AMP systems, topsoil is acidic (possible higher SOC concentration) while subsoil is more alkaline (higher concentration of carbonates).
 - The type of grazing system does not appear to considerably impact Alberta's grassland soil pH regardless of soil depth or ecoregion.
- Moving forward:
 - Once SOC data is acquired, the actual correlation between pH and SOC will be established.
 - After all of our samples, which are representative of the entire Canadian grasslands ecosystem, have been analyzed, we will be able to know the potential of the AMP system in increasing carbon sequestration in grasslands soils and alleviating climate change.
 - Ultimately, ranchers who implement improved grazing practices such as the AMP grazing system might be financially compensated for doing so.

Literature Cited

- Dermer, J. D., & Schuman, G. E. (2007). Carbon sequestration and rangelands: a synthesis of land management and precipitation effects. *Journal of soil and water conservation*, 62(2), 77-85. Retrieved from <http://www.jswnonline.org/content/62/2/77.full.pdf+html>
- Dodds, W. K., Gido, K., Whiles, M. R., Fritz, K. M., & Matthews, W. J. (2004). Life on the edge: the ecology of Great Plains prairie streams. *BioScience*, 54(3), 205-216. [https://doi.org/10.1641/0006-3568\(2004\)054\[0205:LOTETE\]2.0.CO;2](https://doi.org/10.1641/0006-3568(2004)054[0205:LOTETE]2.0.CO;2)
- Gauthier, D. A., & Wiken, E. B. (2003). Monitoring the conservation of grassland habitats, Prairie Ecozone, Canada. *Environmental Monitoring and Assessment*, 88(1-3), 343-364. <https://doi.org/10.1023/A:1025585527169>
- Jin, S., & Wang, H. (2018). Relationships between soil pH and soil carbon in china's carbonate soils. *Fresenius environmental bulletin*, 27(1), 605-611. Retrieved from https://www.researchgate.net/publication/323846896_Relationships_between_soil_pH_and_soil_carbon_in_China's_carbonate_soils
- Kalra, Y. P., & Maynard, D. G. (1991). *Methods manual for forest soil and plant analysis* (Vol. 319). Retrieved from <http://cfs.nrcan.gc.ca/pubwarehouse/pdfs/11845.pdf>
- Teague, R., Provenza, F., Kreuter, U., Steffens, T., & Barnes, M. (2013). Multi-paddock grazing on rangelands: why the perceptual dichotomy between research results and rancher experience? *Journal of Environmental management*, 128, 699-717. Retrieved from <https://www.sciencedirect.com/science/article/pii/S0301479713004131>
- Williams, A. R. (2017). In Defense of Soil Health. Retrieved from https://uknowledge.uky.edu/cgi/viewcontent.cgi?article=1143&context=ky_grazing

Acknowledgements

Thank you to Laio Silva Sobrinho for all of his help and guidance. To Dr. Scott Chang and everyone in my lab, thank you for allowing me to be a part of your team. I also want to thank my lab partner Sarah, and WISEST for this incredible opportunity. Special thanks to Isomass and the Faculty of ALES for sponsoring my lab placement. This project is being funded by the Agricultural Greenhouse Gases Program (AGGP), a federal initiative by Agriculture and Agri-Food Canada (AAFC).