

University of Alberta

Dissecting Arabidopsis root responses to salt stress

by

Yuanqing Jiang



A thesis submitted to the Faculty of Graduate Studies and Research in partial
fulfillment of requirements for the degree of Doctor of Philosophy

in

Plant Biology

Department of Biological Sciences

Edmonton, Alberta

Spring 2008



Library and
Archives Canada

Published Heritage
Branch

395 Wellington Street
Ottawa ON K1A 0N4
Canada

Bibliothèque et
Archives Canada

Direction du
Patrimoine de l'édition

395, rue Wellington
Ottawa ON K1A 0N4
Canada

Your file *Votre référence*
ISBN: 978-0-494-45539-5
Our file *Notre référence*
ISBN: 978-0-494-45539-5

NOTICE:

The author has granted a non-exclusive license allowing Library and Archives Canada to reproduce, publish, archive, preserve, conserve, communicate to the public by telecommunication or on the Internet, loan, distribute and sell theses worldwide, for commercial or non-commercial purposes, in microform, paper, electronic and/or any other formats.

The author retains copyright ownership and moral rights in this thesis. Neither the thesis nor substantial extracts from it may be printed or otherwise reproduced without the author's permission.

AVIS:

L'auteur a accordé une licence non exclusive permettant à la Bibliothèque et Archives Canada de reproduire, publier, archiver, sauvegarder, conserver, transmettre au public par télécommunication ou par l'Internet, prêter, distribuer et vendre des thèses partout dans le monde, à des fins commerciales ou autres, sur support microforme, papier, électronique et/ou autres formats.

L'auteur conserve la propriété du droit d'auteur et des droits moraux qui protègent cette thèse. Ni la thèse ni des extraits substantiels de celle-ci ne doivent être imprimés ou autrement reproduits sans son autorisation.

In compliance with the Canadian Privacy Act some supporting forms may have been removed from this thesis.

Conformément à la loi canadienne sur la protection de la vie privée, quelques formulaires secondaires ont été enlevés de cette thèse.

While these forms may be included in the document page count, their removal does not represent any loss of content from the thesis.

Bien que ces formulaires aient inclus dans la pagination, il n'y aura aucun contenu manquant.

■*■
Canada

Abstract

Salt stress is one of the dominant abiotic stresses that influence plant growth and decrease the harvestable yield of all crop species. Roots are the primary site of salinity perception. To better understand salt stress responses in *Arabidopsis* roots, both the transcriptome and proteome changes in *Arabidopsis* roots treated by 150 mM NaCl for short and long terms were monitored using a full-genome oligonucleotide microarray and two-dimensional gel electrophoresis (2-DGE) based technologies, respectively. Statistical analysis of microarray results revealed that 5, 280 (22%) genes were salt-regulated with 2, 367 genes up-regulated and 2, 913 genes down-regulated by a cutoff of 2-fold. Many novel salt-responsive genes including transporters and transcription factors (TFs) were identified. A comparative proteomic analysis of salt-stressed *Arabidopsis* roots using 2-DGE and liquid-chromatography-tandem mass spectrometry (LC-MS/MS) allowed identification of 86 differentially expressed proteins, including many previously characterized stress-responsive proteins and proteins implicated in reactive oxygen species (ROS) scavenging, signal transduction, translation, cell wall biosynthesis as well as energy, amino acid, and hormone metabolisms. These data confirmed the modest correlation between mRNA and protein levels. Three novel salt-inducible transcription factor genes, *bHLH92*, *WRKY25* and *WRKY33* were chosen for further characterization through a combination of reverse genetic strategies including knock-out and overexpression. Quantitative RT-PCR demonstrated that the three TFs were responsive to multiple abiotic stresses. Promoter-*GUS* fusion assays revealed the spatial and developmental expression patterns and demonstrated that promoters are inducible by some abiotic stresses. Knock-out mutants for the three TFs were

identified, and stress tolerance assays showed that null mutants displayed minor differences from wild-type plants under some specific abiotic stresses, suggesting a functional redundancy exists between members of the same gene families. Constitutive overexpression of the three TFs in Arabidopsis enhanced its tolerance to salt and/or osmotic stress. Finally microarrays were employed to identify the target genes of the three TFs under saline condition, and many target genes were found to be differentially expressed between mutants and wild-type. This study not only provides a more comprehensive description of salt-responsive genes and proteins, but also contributes to elucidating the roles of bHLH and WRKY TFs in abiotic stress.

Acknowledgements

I am grateful to my supervisor Dr. Michael Deyholos for great supervising, instruction, encouragement and technical help throughout my graduate study and research. I also thank: Dr. Anthony Cornish and Troy Locke in MBSU for help in slide printing, microarray scanner use, quantitative PCR and mass spectrometry system use; Lisa Ostafichuk and Pat Murray in MBSU for primer synthesis and DNA sequencing service. Many thanks to my supervisory committee members: Drs. Nat Kav, Enrico Scarpella and Neil Adams for suggestion and help on the proposal and thesis. I would also like to thank Dr. Neil Harris and Matt Bryman for help in atomic absorption spectrometry and conductivity meter use. I gratefully acknowledge the technical help and kind cooperation provided by fellow graduate students, a technician, a post-doctoral fellow and a visiting scientist working in Mike's lab: Mary DePauw, Naomi Hotte, Ming-feng Yang, Manjeet Kumari, Disa Brownfield, Melissa Roach, Mohsen Mohammadi, Hameed Khadair, Abdur Rashid, Ryan McKenzie and Sue Koziel. I also thank the SALK Institute, Syngenta and the Arabidopsis Biological Resource Center (ABRC) for providing T-DNA insertion lines and mutants. Finally, I would like to thank my family members, especially my wife, Bo Yang, for their support and encouragement throughout my doctoral research. The project was funded by NSERC and Alberta Ingenuity awards to M.K.D.

Table of Contents

Chapter 1: Literature Review

Introduction.....	1
Effects of salt toxicity on plant physiology.....	2
Water deficit due to salt stress.....	3
Sodium entry and efflux of plant cells.....	4
Sodium compartmentation.....	5
Reactive oxygen species and antioxidants.....	6
Role of ABA in salt stress.....	8
Salt signaling.....	11
Specificity and cross talk in abiotic stresses.....	12
Strategies of developing salt tolerant plants.....	16
Conclusions and future perspectives.....	26
Objects.....	27
References.....	27

Chapter 2: Microarray analysis of Arabidopsis transcriptome change in response to salt stress

Introduction.....	47
Materials and methods.....	47
Plant growth and stress treatment.....	48
Physiological analyses.....	48
Microarray preparation, hybridization and data extraction.....	48
Microarray data analysis.....	49
Quantitative PCR.....	50
Results and Discussion.....	52
Whole-plant responses to salt treatment.....	52
General transcriptomic responses.....	54
Validation of microarray results.....	55

Classification of NaCl-responsive transcripts.....	56
Osmoprotectants.....	59
Reactive oxygen network.....	59
Transporters.....	62
Primary energy metabolism, carbohydrates and cell walls.....	62
Protein metabolism.....	65
Signal transduction components and hormones.....	68
Transcription Factors.....	72
Conclusion.....	77
References.....	78

Chapter 3: Comparative proteomic analysis of Arabidopsis root response to salt stress

Introduction.....	88
Materials and methods.....	89
Plant materials and stress treatment.....	89
Physiological analyses.....	89
Protein extraction and quantification.....	90
IEF and SDS-PAGE.....	90
Image and statistical analysis.....	91
In-gel digestion and LC-MS/MS analysis.....	92
Results	93
Plant growth response to NaCl stress.....	93
2-DGE analysis of NaCl-responsive proteins in Arabidopsis roots.....	96
LC-MS/MS identification and classification of NaCl-responsive proteins.....	96
Correlation analysis of mRNA and proteins levels	98
Discussion.....	101
Energy metabolism.....	101
ROS scavenging and detoxifying enzymes.....	102
Protein translation, processing and degradation.....	109
Cell wall-related proteins.....	111

Hormone-related proteins.....	112
Signal transduction network involved in NaCl stress responses.....	113
Amino acid metabolism.....	113
Cytoskeleton.....	114
Transcription-related proteins.....	115
Conclusion.....	115
References.....	116

Chapter 4: Functional characterization of salt-induced bHLH92 transcription factor gene

Introduction.....	126
Materials and methods.....	128
Plant growth and stress treatment.....	128
Quantitative RT-PCR	129
Identification of homozygous T-DNA insertion mutants.....	130
Plasmid constructions.....	131
Arabidopsis transformation and selection.....	132
GFP observation	133
Histochemical GUS staining.....	133
Stress tolerance tests and REL measurement.....	134
Microarray profiling of <i>bhlh92</i> mutant.....	134
Phylogenetic tree.....	135
Results.....	136
Sequence and phylogenetic analysis of <i>bHLH92</i> gene.....	136
Responses of <i>bHLH92</i> gene to various abiotic stresses.....	139
Salt-induced accumulation of <i>bHLH92</i> transcript in ABA and SOS mutants.....	140
Promoter-GUS fusion.....	142
Subcellular localization.....	145
Identification of T-DNA insertion lines and stress tolerance tests.....	145
Over-expression of bHLH92 gene in <i>Arabidopsis</i>	149
Microarray identification of bHLH92 target genes	151

Discussion.....	152
References.....	157

Chapter 5: Functional characterization of salt-induced WRKY25 and WRKY33 transcription factor genes

Introduction.....	166
Materials and methods.....	168
Plant growth, stress treatments and mutant isolation	168
Quantitative RT-PCR.....	169
Plasmid construction and plant transformation.....	170
GFP observation and GUS staining.....	171
Stress tolerance tests.....	171
Microarray profiling, data analysis and validation	172
Analysis of promoter motifs in co-regulated gene sets.	173
Sequence and phylogenetic analysis.....	173
Results.....	174
Identification and phylogenetic analysis of <i>WRKY25</i> and <i>WRKY33</i> genes.....	174
Expression analysis of <i>WRKY25</i> and <i>WRKY33</i> transcripts in <i>Arabidopsis</i> tissues.....	177
Subcellular localization.....	178
<i>WRKY25</i> and <i>WRKY33</i> are induced by multiple abiotic stresses.....	180
Expression analysis of <i>WRKY25</i> and <i>WRKY33</i> transcripts in ABA and SOS mutants	181
Analysis of promoter-GUS fusion lines.....	183
Mutant phenotypes of <i>WRKY25</i> and <i>WRKY33</i>	186
Overexpression of <i>WRKY25</i> and <i>WRKY33</i> genes in <i>Arabidopsis</i>	190
Microarray identification of <i>WRKY25</i> and <i>WRKY33</i> target genes.....	193
W-boxes are highly enriched in <i>WRKY25</i> and <i>WRKY33</i> regulon promoters....	205
Discussion.....	207
WRKY 25 and WRKY33 are involved in abiotic stress responses.....	208
WRKY25 and WRKY33 regulate salt stress response by targeting to diverse downstream genes.....	212

References.....214

Chapter 6: General discussion and conclusions.....226

References.....234

List of Tables

Table 1-1. Engineered salt stress tolerance through genetic manipulation of plant-source genes.....	17
Table 2-1. Comparison of qRT-PCR and microarray results for selected genes.....	56
Table 2-2. NaCl-responsive transcripts related to osmoprotection, detoxification, and transport.....	60
Table 2-3. NaCl-responsive transcripts related to respiration and carbohydrate and cell wall metabolism.	63
Table 2-4. NaCl-responsive transcripts related to protein metabolism.....	67
Table 2-5. NaCl-responsive transcripts related to signal transduction and hormone biosynthesis.	69
Table 2-6. Classification of NaCl-regulated transcription factors.....	73
Table 3-1. Differentially expressed proteins identified by LC-MS/MS.	104
Table 4-1. Putative regulatory sequences identified in the 1 kb upstream region of <i>bHLH92</i>	144
Table 4-2. Genes that are regulated by 150 mM NaCl in <i>bhlh92</i> mutant.....	152
Table 5-1. Putative regulatory sequences found in the promoter regions of <i>WRKY25</i> and <i>WRKY33</i>	186
Table 5-2. Genes that are up-regulated by 150 mM NaCl in <i>wrky25-1</i> mutant.....	194
Table 5-3. Genes that are down-regulated by 150 mM NaCl in <i>wrky25-1</i> mutant.....	195
Table 5-4. Genes that are positively regulated by 150 mM NaCl in <i>wrky33-1</i> mutant.....	197
Table 5-5. Genes that are negatively regulated by 150 mM NaCl in <i>wrky33-1</i> mutant.....	200
Table 5-6. Occurrence of W-box and W-box-like motifs in the <i>WRKY25</i> and <i>WRKY33</i> regulons.....	206

List of Figures

Figure 1-1. Regulation of sodium, potassium and calcium homeostasis in plant cells...	7
Figure 1-2. Transcriptional cascades of salt, drought and cold signal transduction.....	13
Figure 2-1. Accumulation of anthocyanin and praline in Arabidopsis plants in responses to NaCl treatment.....	53
Figure 2-2. Venn diagrams of salt-responsive genes.....	54
Figure 2-3. NaCl-responsive transcripts grouped according to temporal expression profiles.....	58
Figure 3-1. Root elongation and relative electrolyte leakage assay.....	94
Figure 3-2. Changes in Na and K ion concentrations following NaCl exposure.....	95
Figure 3-3. A representative 2-DE gel of Arabidopsis root proteins.....	97
Figure 3-4. Functional classification of NaCl-responsive proteins.....	99
Figure 3-5. Comparison of NaCl-induced changes in mRNA and cognate protein abundance.....	100
Figure 4-1. Multiple alignment of bHLH92 and four other <i>Arabidopsis</i> bHLH TFs...	137
Figure 4-2. Phylogenetic tree of 162 AtbHLHs.....	139
Figure 4-3. qRT-PCR analysis of <i>bHLH92</i> transcript level in various tissues and organs.....	139
Figure 4-4. qRT-PCR profiling of <i>bHLH92</i> and <i>RD29A</i> transcript levels in Arabidopsis roots.	140
Figure 4-5. qRT-PCR analysis of <i>bHLH92</i> transcript levels in wild-type, ABA and SOS mutants.	141
Figure 4-6. Histochemical localization of GUS activity in representative transgenic Arabidopsis plants harboring PbHLH92-GUS construct.....	143
Figure 4-7. Subcellular localization of bHLH92 fused to GFP in Arabidopsis.	146
Figure 4-8. Identification of T-DNA insertion mutants and stress tolerance testing...	148
Figure 4-9. Identification and stress tolerance testing of <i>bHLH92</i> -overexpressing Arabidopsis.....	150
Figure 4-10. qRT-PCR validation of microarray results.....	151

Figure 5-1. Comparison of the deduced amino acid sequences of WRKY25 and WRKY33.....	175
Figure 5-2. Phylogenetic analysis of 72 members of the Arabidopsis WRKY gene family.....	176
Figure 5-3. Microarray results of <i>WRKY2</i> , <i>WRKY20</i> , <i>WRKY25</i> , <i>WRKY26</i> and <i>WRKY33</i> <i>WRKY33</i> transcript abundance changes in response to salt stress	177
Figure 5-4. qRT-PCR analysis of <i>WRKY25</i> and <i>WRKY33</i> transcript levels in various tissues and organs.....	178
Figure 5-5. Subcellular localization WRKY33 fused to GFP in Arabidopsis.....	179
Figure 5-6. qRT-PCR analysis of <i>WRKY25</i> and <i>WRKY33</i> transcript abundance in Arabidopsis roots subjected to various treatments.....	181
Figure 5-7. qRT-PCR analysis of <i>WRKY25</i> and <i>WRKY33</i> transcript levels in salt treated wild-type, ABA and SOS mutants.	182
Figure 5-8. Histochemical localization of GUS activity in representative transgenic Arabidopsis plants harboring promoter-GUS constructs.....	185
Figure 5-9. Identification of T-DNA insertion mutants and stress tolerance assay....	189
Figure 5-10. Identification and stress tolerance testing of <i>WRKY25</i> - and <i>WRKY33</i> -overexpressing Arabidopsis.	192
Figure 5-11. qRT-PCR validation of microarray results.	204
Figure 5-12. Proposed working model explaining role of WRKY25 and WRKY33 in salt stress responsive pathway.	213

List of Abbreviations

2-DGE, two-dimensional gel electrophoresis
4CL, 4-coumarate:CoA ligase
ABA, abscisic acid
ABC, ATP-binding cassette
ABF, ABA-responsive element binding factor
ABI, ABA insensitive
ABRC, Arabidopsis Biological Resource Center
ABRE, ABA-responsive element
ACS1, Arabidopsis cysteine synthase
AGI, Arabidopsis genome initiative
AKT, Arabidopsis K⁺ transporter
APX, ascorbate peroxidase
AREB, ABRE-binding protein
ARF, auxin Response Factor
At, *Arabidopsis thaliana*
AUX/IAA, auxin/indole-3-acetic acid.
BADH, betaine aldehyde dehydrogenase
bHLH, basic helix-loop-helix
bp, base pairs
BSA, bovine serum albumin
bZIP, basic leucine zipper
CaMV, cauliflower mosaic virus
CAX, Ca²⁺/H⁺ antiporter
CBF, C-repeat-binding factor
CBL, calcineurin B-like protein
CDPK, Ca²⁺-dependent protein kinase
CIPK, CBL-interacting protein kinase
CDS, coding region
CHAPS, 3-[(3-cholamidopropyl) dimethylammonio]-1-propanesulfonate;

COMT, caffeic acid O-methyltransferase
COR, cold regulated
CPK, calcium responsive kinases
CRT, C-repeat
DAS, days after stratification
DAF, days after flowering
DD, differential display
DMSO, dimethyl sulfoxide
dNTP, deoxyribonucleotides
DRE, dehydration-responsive element
DREB, DRE-binding protein
DSP, dual specific phosphatases
DTT, dithiothreitol
EDTA, ethylenediamine tetraacetic acid
ERD, early responsive to dehydration
ERF, ethylene-responsive element binding factor
EREBP, ethylene response element binding protein
ET, ethylene
FDR, false discovery rate
Fwd, forward
G6PDH, glucose-6-phosphate-1-dehydrogenase
GA, gibberellin
GCR, G-protein-coupled receptor
GFP, green fluorescent protein;
GH, glycoside hydrolase
GPX, glutathione peroxidase
GRP, glycine-rich protein
GSH, glutathione
GST, glutathione S-transferase
GT, glycosyltransferase
GUS, β -glucuronidase

HKT, high affinity K⁺ transporter
HPLC, high performance liquid chromatograph
HPTII, hygromycin B phosphotransferase
HSF, heat shock factor
ICAT, isotope-coded affinity tag
ICE1, inducer of CBF expression 1
IEF, isoelectric focusing
IPG, immobilized pH gradient
IPTG, isopropyl-beta-D-thiogalactopyranoside
JA, jasmonic acid
KIN, cold inducible
LB, left border
LEA, late-embryogenesis-abundant
LC-MS/MS, liquid chromatography coupled to tandem mass spectrometry;
LOB, lateral organ boundary
LOS, low expression of osmotically responsive gene
LRR, leucine-rich repeat
LTP, lipid transfer proteins
MAPK, mitogen-activated protein kinase
MAPKK, mitogen-activated protein kinase kinase
MAPKKK/MAP3K, mitogen-activated protein kinase kinase kinase
MATE, multidrug and toxic compound extrusion
MDAR, Monodehydroascorbate reductase
MeJA, methyl jasmonic acid
MKS1, MAP kinase substrate 1
MLO, mildew resistance locus O
MS, Murashige-Skoog
MudPIT, multidimensional protein identification technology
MV, methyl viologen (Paraquat)
MW, molecular weight
NHX, Na⁺/H⁺ antiporter

NLS, nuclear localization signal
NOS, nopaline synthase
NPK, *Nicotiana* protein kinase
NPR1, nonexpressor of PR genes 1
NSCC, non-selective cation channels
Os, *Oryza sativa*
P5CR, delta(1)-pyrroline-5-carboxylate reductase
P5CS, delta(1)-pyrroline-5-carboxylate synthetase
PDH, pea DNA helicase
PERK, proline-rich extensin-like kinases
PP2C, protein phosphatase 2C
PPI, peptidylprolyl isomerase
PPP, pentose phosphate pathway
PR, pathogenesis-related
PRX/PER, peroxidase
pI, Isoelectric point
PTM, post-translational modification
qRT-PCR, quantitative RT-PCR
QTL, quantitative trait locus
RAB, responsive to ABA
RB, right border
RCD1, radical-induced cell death 1
RD, response to dehydration
REL, relative electrolyte leakage
RING, really interesting new gene
RLCK, receptor like cytoplasmic kinase
RLK, receptor-like kinases
ROS, reactive oxygen species
RPS6K, ribosomal protein S6 kinases
RT-PCR, reverse transcription-polymerase chain reaction
Rvs, reverse

SA, salicylic acid
SAGE, serial analysis of gene expression
SAM, significance analysis of microarray
SCOF1, soybean C2H2-type zinc finger protein1
S.D., standard deviation
SDS, sodium dodecyl sulfate
SDS-PAGE, sodium dodecyl sulfate–polyacrylamide gel electrophoresis
SFR, sensitive to freezing
SGBF1, soybean G-box binding bZIP transcription factor 1
SOD, superoxide dismutase
SOS, salt overly sensitive
SSH, suppressive subtractive hybridization
STEM, short time-series expression miner
TAIR, the Arabidopsis Information Resource
TAP, tandem affinity purification
TCA, tricarboxylic or trichloroacetic acid
TF, transcription factor
TIGR, the Institute of Genome Research
TL, translational leader
TSAP, thermo-sensitive alkyline phosphatase
UBQ, ubiquitin
UTR, untranslated region
WAK, wall-associated kinase
WT, wild-type
X-gluc, 5-bromo-4-chloro-3-indolyl- β -D-glucuronide
XTH, xyloglucan endotransglucosylase
ZIM, Zinc-finger protein expressed in inflorescence meristem

Chapter 1: Literature Review

Introduction

Salinity is a severe and increasing constraint on the productivity and yield of agricultural crops and threatens food sustainability worldwide. Saline soil has toxic levels of NaCl and Na₂SO₄ and by definition, the electrical conductivity (EC) of saturation extracts of saline soil exceeds more than 4.0 dS/m (deciSiemen/metre, 1.0 dS/m \approx 10mM NaCl). However, many crops are also affected by soil with an EC less than 4 dS/m (Chinnusamy *et al.*, 2005).

It is estimated that 45 Mha of the current 230 Mha of irrigated land are salt-affected (20%) (Munns 2005). Primary salinity in soil mostly originates from previous exposure to sea water. Secondary salinization of arable land arises from low-quality irrigation water and improper drainage, which has led to the decline of arable land, and it is threatening the long-term sustainability of many present large-scale irrigation systems. Excess salts in the soil interfere with mineral nutrition and water uptake, and lead to undue accumulation of toxic ions in plants. Salinity imposes two stresses on plant tissue: an initial water-deficit that results from the relatively high solute concentration in the soil, which is caused by the osmotic effect of salt around the roots, and the subsequent ion-specific stress resulting from altered K⁺/Na⁺ ratio. Although sodium is an essential micronutrient for some C₄ plants, in which Na⁺/pyruvate co-transporters import pyruvate into mesophyll chloroplasts, most crop plants are natrophobic (Ohnishi *et al.*, 1990). With regards to salt sensitivity, plants can be divided into two groups: glycophytes, which are salt sensitive, and halophytes, which are salt tolerant that can grow in 200 to 500 mM NaCl. Halophytes can be further divided into two subcategories, euhalophytes or facultative halophytes, which require salt for optimum growth (e.g. *Suaeda maritima*, *Atriplex nummularia*), and miohalophytes, which are capable of growth under high levels of salt, although the growth rate is inhibited by salt, e.g. *Atriplex hastata*, *Spartina townsendii* and sugar beet (*Beta vulgaris* L.) (Glenn *et al.*, 1999). Nearly all salt tolerant plants are angiosperms, and these species are widely distributed amongst angiosperms, indicating a polyphyletic origin or that halophytes are primitive genetic remnants of different families (Flowers 2004).

Considerable efforts were taken in the past several decades to understand the underlying genetic bases of the physiology of salinity tolerance, but little progress was made until the introduction of *Arabidopsis* as a genetic model to study salinity tolerance (Zhu 2000), partly due to the difficulty of conducting genetic analysis in most of the well-characterized halophytes. As *Arabidopsis* is a glycophyte that does not exhibit salinity tolerance even approaching that of halophytes, its utilization in salinity tolerance studies was initially thought to be inappropriate. However, forward genetic studies using *Arabidopsis* have yielded essential knowledge about the genetic bases of salinity tolerance (Zhu 2002). In the following sections, I will review and discuss the toxicity of salt on plants, cellular dehydration caused by salinity, the molecular basis of ion homeostasis, the role of abscisic acid (ABA) in salt stress signaling, the cross-talk and specificity between abiotic stresses, as well as the utilization of various genes for engineering salt-tolerant plants.

Effects of salt toxicity on plant physiology

Most commonly, high Na^+ and Cl^- in the soil and water cause salt stress. Salt stress has three effects; i.e. it reduces water potential (osmotic stress) and causes ion imbalance or disturbances in ion homeostasis and at the same time leads to cellular toxicity (Hasegawa *et al.*, 2000). General symptoms of damage caused by salt stress are growth inhibition due to dehydration as a result of altered water status, accelerated development, senescence and death during prolonged exposure (Munns 2005). The immediate response to salt stress is a reduction in the rate of leaf surface expansion leading to cessation of expansion as salt concentration increases (Wang and Nil, 2000). Growth inhibition is the primary injury that leads to other symptoms although programmed cell death (PCD) may also occur under severe salinity shock. Since salt stress involves both osmotic and ionic stress (Munns 2002), growth suppression is directly related to the total concentration of soluble salts or osmotic potential of soil water. Water potential and osmotic potential of plants become more negative with an increase in salinity (Hernandez *et al.*, 1999; Romeroaranda *et al.*, 2001). Salt stress induces the synthesis of ABA (Jia *et al.*, 2001), which closes stomata when transported to guard cells (Zeevaart and Creelman, 1988). As a result of stomatal closure, photosynthesis declines

and photoinhibition and oxidative stress occur. Another reason for decreased photosynthetic rate is that chlorophyll and total carotenoid contents of leaves decrease in general under salt stress (Hernandez *et al.*, 1999). Oxidative stress occurs under salt stress because of the formation of reactive oxygen species (ROS) such as superoxides and hydroxy and peroxy radicals. The ROSs that are by-products of hyperosmotic and ionic stresses cause membrane dysfunction and cell death (Bohnert and Jensen, 1996).

Water deficit due to salinity stress

A major initial consequence of NaCl stress is a reduced ability to take up water from the soil, which leads to slower growth. The amount of water transport depends on the motive force generated by an osmotic gradient (difference in water potential between two compartments) and water permeability. When extracellular water potential is reduced by salinity, the osmotic gradient can be restored by decreasing the intracellular water potential with the accumulation of compatible solutes, also known as osmolytes or osmoprotectants (Chen and Murata, 2000). These solutes include proline, glycine betaine, raffinose, mannitol, sorbitol, sucrose and trehalose, which are all relatively small, non-toxic compounds that can stabilize proteins and cellular structures, and are proposed to increase the osmotic pressure of plant cells. Many genes involved in osmoprotectant biosynthesis are upregulated under salt and drought stresses (Chen and Murata, 2000). Over-expression of genes involved in the biosynthesis of compatible osmolytes, either from *E. coli* or plants, in transgenic *Arabidopsis*, rice, wheat, tobacco and *Brassica* shows enhanced germination, growth, survival, recovery, and production under salt and osmotic stresses (Bohnert and Jensen, 1996; Chinnusamy *et al.*, 2005). Cell-to-cell water transport is facilitated by membrane proteins called aquaporins, which form water channels (Maurel and Chrispeels, 2001). Some aquaporin genes are regulated by salt or drought stress. Over-expression of a barley aquaporin increased salt sensitivity in transgenic rice plants, indicating that the increase in water conductance of leaf cells accelerates water loss from transgenic leaves (Katsuhara *et al.*, 2003).

Another group of water stress induced proteins includes late-embryogenesis-abundant proteins (LEAs) and their close relatives, dehydrins. These proteins may help to stabilize protein structures when cells start to lose water. Xu *et al.* (1996) reported that

the expression of a barley LEA protein, HVA1, confers tolerance to water deficit and salt stress in transgenic rice. However, overexpression of a single LEA-type protein is not always sufficient to confer plant stress tolerance, suggesting that LEA-type proteins might function synergistically with other proteins (Wang *et al.*, 2003).

Sodium entry and efflux of plant cells

While Na^+ is deleterious for plant growth, K^+ is an essential element and is required by plants in large quantities. Movement of salt into roots and to shoots is driven by transpirational flux required to maintain the water status of plants. Na^+ entry into the cells of the root is passive and uniporter or ion channel type transporters are the candidates for sodium influx (Tester and Davenport, 2003). Many K^+ transport systems, such as Na^+/K^+ symporters, have some affinity for Na^+ , which can enter cells through the high-affinity K^+ transporter HKT1 and non-selective cation channels (NSCCs). AtHKT1 is a low affinity Na^+ transporter and mediates Na^+ entry into the root cells of *Arabidopsis* under salt stress, and mutation in *HKT1* suppresses Na^+ accumulation and hypersensitivity of the plants to Na^+ (Berthomieu *et al.*, 2003; Rus *et al.*, 2001). NSCCs catalyze passive fluxes of cations through plant membranes and do not show much selectivity between monovalent cations; several NSCCs are also permeable to divalent cations. NSCCs are directly involved in a multitude of stress responses, growth and development, uptake of nutrients and calcium signaling (Demidchik and Maathuis, 2007). Accumulated evidence has supported that NSCCs are the main pathway for Na^+ entry into the roots, at high soil NaCl concentrations (Tester and Davenport, 2003). In wheat, a low affinity cation transporter (LCT1) was isolated using a screen for complementation of K^+ uptake mutant of yeast (Schachtman *et al.*, 1997). In yeast, LCT1 catalyzed uptake of many cations, including Na^+ , Rb^+ , and Ca^{2+} . Introduction of wheat LCT1 into yeast causes hypersensitivity to sodium in yeast, which was mitigated by the addition of high (20 mM), but not low (2 mM) Ca^{2+} to the medium (Amtmann *et al.*, 2001). It was therefore suggested that LCT1 may be a component of the Ca^{2+} -insensitive Na^+ influx seen in wheat roots, as it was not saturated at low external Na^+ (7 mM), a property attributed to NSCCs (Amtmann *et al.*, 2001). Similar transporters in rice or

Arabidopsis have not been identified, suggesting LCT1 may be unique to wheat (Aspe and Blumwald, 2007).

Sodium extrusion from plant cells is mediated by the plasma membrane Na^+/H^+ antiporter SOS (Salt Overly Sensitive)1, which is composed of 12 trans-membrane domains (Shi *et al.*, 2000). Activation of SOS1 is controlled by SOS3 and SOS2. The *SOS3* gene encodes a myristoylated Ca^{2+} binding protein, also known as CBL4 (calcineurin B-like protein 4). In response to Ca^{2+} perturbation, SOS3 changes its conformation and transduces a signal downstream by interacting with an effector kinase, SOS2, also known as CIPK24 (CBL-interacting protein kinase 24). *SOS2* encodes a novel serine/threonine protein kinase with an N-terminal catalytic and C-terminal regulatory domain. SOS3 interacts with SOS2 via a FISL motif and relieves the protein from autoinhibition, thereby making the kinase active (Fig. 1-1, Zhu, 2002). Na^+ extrusion is powered by a plasma membrane H^+ -ATPase that generates an electrochemical H^+ gradient and allows Na^+/H^+ antiporters to couple the passive movement of H^+ inside the cells with the active extrusion of Na^+ . Overexpression of *SOS1* in *Arabidopsis* shows improved salt tolerance, demonstrating that limiting Na^+ accumulation in cytoplasm is an effective way to attain salt tolerance (Shi *et al.*, 2003). The discovery of SOS genes paved the way for elucidation of a novel pathway linking the Ca^{2+} signaling to salt stress (Zhu, 2002 & 2003).

Sodium compartmentation

Compartmentation of Na^+ in the vacuole allows plants to maintain a lower Na^+ concentration at the sites of biochemical reactions in the cytosol, and yet maintain a lower overall osmotic potential. Influx of Na^+ into the vacuole occurs through Na^+/H^+ antiporters, which use the proton gradient generated by V-type H^+ -ATPase and H^+ -pyrophosphatase (PPase) (Apse *et al.* 1999). Salinity up-regulates the expression of a V-type H^+ -ATPase gene (Gollidack and Dietz 2001) and a vacuolar Na^+/H^+ antiporter gene (Gaxiola *et al.* 1999; Shi and Zhu 2002).

In *Arabidopsis*, the *AtNHX1* gene encodes a tonoplast Na^+/H^+ antiporter. *AtNHX1* expression under salt stress is partially dependent on ABA biosynthesis and ABA signaling through the protein phosphatase 2C-ABI1 (ABA insensitive 1). Salt-stress

induced up-regulation of AtNHX1 expression is lower in ABA deficient mutants (*aba2-1* and *aba3-1*) and in the ABA insensitive mutant, *abil-1* (Shi and Zhu, 2002). It was also shown that SOS2 regulates the AtNHX1 activity (Qiu *et al.*, 2004). More recently, in a yeast two-hybrid analysis, it was shown that SOS2 interacts directly with V-ATPase regulatory subunits B1 and B2, suggesting that regulation of V-ATPase activity is an additional key function of SOS2 in coordinating changes in ion transport during salt stress and in promoting salt tolerance (Batelli *et al.*, 2007). Overexpression of AtNHX1 in *Arabidopsis*, canola and tomato resulted in significantly higher salt tolerance (Apse *et al.*, 1999; Zhang *et al.*, 2001; Zhang and Blumwald, 2001). Overexpression of AVP1 in *Arabidopsis* plants, a vacuolar H⁺-PPase, showed enhanced salt tolerance as well as drought tolerance (Gaxiola *et al.*, 2001). These studies confirm that sequestration of Na⁺ into the vacuole is an important trait of salt tolerance in plants (Fig. 1-1).

Reactive oxygen species and antioxidants

A secondary aspect of salinity stress in plants is the increased abundance of reactive oxygen species (ROS). In plants, ROSs, including superoxide radicals, hydrogen peroxide and hydroxyl radicals, are continuously produced predominantly in chloroplasts, mitochondria, and peroxisomes (Apel and Hirt, 2004). In other circumstances, plants appear to purposefully generate ROS as signaling molecules to control various processes including pathogen defense, programmed cell death, and stomatal behavior (Apel and Hirt, 2004). However, ROSs can also cause oxidative damage to membrane lipids, proteins and nucleic acids (Smirnoff, 1993). Therefore, production and removal of ROS must be strictly controlled. However, the equilibrium between production and scavenging of ROS may be perturbed by a number of adverse abiotic stress factors such as high salinity, high light, drought, low temperature, high temperature, and mechanical stress (Mittler *et al.*, 2004).

Studies on the role of ROS in signaling in plants revealed that ROS might be sensed by two-component histidine kinases (HKs), and activate mitogen-activated protein kinase (MAPK) signaling pathways (Jonak *et al.*, 2002). ROS might change gene expression by targeting and modifying the activity of transcription factors. In




Fig. 1-1. Regulation of sodium, potassium and calcium homeostasis in plant cells. Higher concentration of extracellular NaCl is perceived by an unknown receptor at the plasma membrane (PM), which induces a cytosolic calcium perturbation. SOS3 senses the Ca²⁺ increase and interacts with SOS2. This interaction relieves SOS2 of its auto-inhibition and activates it, which forms a complex with SOS3 to phosphorylate PM-localized Na⁺/H⁺ antiporter SOS1, resulting in efflux of excess Na⁺ ions. SOS3–SOS2 complex inhibits the low affinity Na⁺ transporter HKT1 activity thus restricting Na⁺ entry into the cytosol. SOS2 interacts and activates the vacuolar Na⁺/H⁺ exchanger NHX1 resulting in sequestration of excess Na⁺ ions into the vacuole. SOS2 also activates the H⁺/Ca²⁺ antiporter CAX1 on the vacuolar membrane reinstating cytosolic Ca²⁺ homeostasis. High salinity also causes osmotic stress, which induces the synthesis of ABA. A protein phosphatase 2C (ABI1) can regulate NHX1 expression. The protein phosphatase ABI2 physically interacts with SOS2 and inactivates SOS2. High intracellular Na⁺ also inhibits the K⁺ transporter-AKT1 on PM. Dotted lines indicate possible regulations (Adapted from Zhu 2003 and Chinnusamy *et al.*, 2005).

Arabidopsis, H₂O₂ activates the MAPKs, MPK3 and MPK6 via MAPKKK ANP1, and overexpression of a tobacco ANP1 orthologue, NPK1, in transgenic tobacco plants

resulted in increased tolerance to heat shock, freezing, and salt stress (Kovtun *et al.*, 2000). H₂O₂ also increases expression of the *Arabidopsis* nucleotide diphosphate kinase (AtNDPK) 2 (Moon *et al.*, 2003). Overexpression of AtNDPK2 reduced accumulation of H₂O₂ and enhanced tolerance to multiple stresses including cold, salt, and oxidative stress. The effect of NDPK2 might be mediated by the MAPKs, MPK3, and MPK6 because NDPK2 can interact and activate the MAPKs (Moon *et al.*, 2003).

Plants use two different mechanisms for ROS scavenging. Nonenzymatic scavenging of ROS uses cellular ascorbic acid, glutathione (GSH), tocopherol, flavonoids, alkaloids, and carotenoids. Enzymatic scavenging employs detoxifying enzymes including superoxide dismutases (SODs), catalases (CATs), ascorbate peroxidases (APXs), and glutathione peroxidases (GPXs) (Apel and Hirt, 2004). SODs act as the first line of defense against ROS, dismutating superoxide to H₂O₂. APX, GPX, and CAT subsequently detoxify H₂O₂. ROS detoxification could provide enhanced plant resistance to salt and other abiotic stresses. Transgenic plants overexpressing some of these enzymes demonstrated increased tolerance to osmotic, temperature and oxidative stresses (Alscher *et al.*, 2002; Roxas *et al.*, 1997; Wang *et al.*, 1999).

A recent report indicated that the plasma membrane Na⁺/H⁺ antiporter SOS1 is also involved in oxidative stress tolerance (Katiyar-Agarwal *et al.*, 2006). It was found that under salt or oxidative stress, SOS1 interacts through its predicted cytoplasmic tail with RCD1 (radical-induced cell death), a regulator of oxidative-stress responses. Without stress treatment, RCD1 is localized in the nucleus. Under high salt or oxidative stress, RCD1 is found not only in the nucleus but also in the cytoplasm. Like *rcd1* mutants, *sos1* mutant plants show an altered sensitivity to oxidative stresses. The *rcd1* mutation causes a decrease in salt tolerance and enhances the salt-stress sensitivity of *sos1* mutant plants. Several genes related to oxidative-stress tolerance were found to be regulated by both RCD1 and SOS1. These results shed light on the cross-talk between the ion-homeostasis and oxidative-stress detoxification pathways involved in plant salt tolerance.

The role of ABA in salt stress

ABA plays an important role in many aspects of plant growth and development and

also in abiotic stress response. In fact, the role of ABA in response to abiotic stresses has long been recognized and extensively studied utilizing mutants deficient in ABA synthesis and sensitivity (Finkelstein *et al.*, 2002; Nambara and Marion-Poll, 2005). Salt, drought, and to a lesser extent, cold stresses, increase the biosynthesis and accumulation of ABA, which then is rapidly catabolised after stress relief (Koornneef *et al.*, 1998; Cutler and Krochko, 1999; Schwartz *et al.*, 2003). Salt stress induced ABA accumulation appears to be due to both ionic and osmotic stresses in roots, while that in the leaf is mainly due to osmotic stress (Jia *et al.* 2002).

The plethora of ABA responses is initiated by ABA perception. Three different ABA receptors, the nuclear RNA binding protein flowering time control protein (FCA), the chloroplast protein Mg chelatase H subunit, and the plasma membrane-localized GCR2 (G-protein-coupled receptor 2) have been identified (Razem *et al.*, 2006; Shen *et al.*, 2006; Liu *et al.*, 2007); however, a more recent report points out that GCR2 is not an ABA receptor in seed germination and early seedling development. Instead, GCR2 is most likely a plant homolog of bacterial lanthionine synthetases (Gao *et al.*, 2007). Genetic disruption of the single $G\alpha$ prototype of heterotrimeric G protein GPA1 present in *Arabidopsis* caused insensitivity of stomatal opening to inhibition by ABA (Wang *et al.*, 2001). GPA1 presumably represents a downstream effector of GCR1, a putative G protein-coupled receptor which acts as a negative regulator of GPA1-mediated ABA responses in guard cells (Pandey and Assmann, 2004). Evidence suggests that G-proteins modulate ABA responses (Wang *et al.*, 2001; Lemichez *et al.*, 2001; Zheng *et al.*, 2002). The Rho-like small G-protein, ROP10, negatively regulates ABA-mediated stomatal closure, germination, and growth inhibition (Zheng *et al.*, 2002). Recruitment of ROP10 to the plasma membrane required a functional farnesylation site, which supports the observed role of a farnesyltransferase β subunit ERA1 as a negative regulator of ABA responses upstream of cytosolic Ca^{2+} (Allen *et al.*, 2002). ROP2 and ROP6/AtRac1 work as pleiotropic negative modulators of ABA responses linked to actin skeleton reorganization and vesicle transport and required for stomatal closure or tip growth (Lemichez *et al.*, 2001; Jones *et al.*, 2002). In line with this conclusion, impairment of vesicle transport or fusion by deficiency in the syntaxin OSM1 (Osmotic Stress-sensitive Mutant 1), resulted in ABA-insensitive stomatal regulation (Zhu *et al.*,

2002). Thus, OSM1 plays a critical role in root growth and in ABA regulation of stomatal responses under osmotic stresses (Zhu *et al.*, 2002).

ABA signaling for stomatal closure and gene expression is transduced through Ca^{2+} (Schroeder *et al.*, 2001). A growing number of protein kinases, such as calcium-dependent protein kinases (CDPKs) as well as protein phosphatases such as PP2Cs, have been implicated in ABA-induced stomatal closure and gene expression (Christmann *et al.*, 2006). ABA transcriptionally upregulates a number of PP2Cs in *Arabidopsis* which act as negative regulators of ABA responses (Saez *et al.*, 2004; Kuhn *et al.*, 2006). Protein phosphatases ABI1 and the highly homologous ABI2 are partly redundant key regulators of ABA-invoked seed dormancy, stomatal closure, and growth inhibition (Finkelstein and Rock, 2002). Previous studies showed that salt-stress induced ABA accumulation regulates the SOS pathway through the ABI2 protein phosphatase 2C (Ohta *et al.*, 2003). ABI2 interacts with the protein phosphatase interaction (PPI) motif of SOS2. This interaction is abolished by the *abi2-1* mutation, which enhances tolerance of seedlings to 150 mM NaCl and causes ABA insensitivity (Ohta *et al.*, 2003). Therefore, ABI2 may negatively regulate salt tolerance either by inactivating SOS2, or the SOS2 regulated Na^+/H^+ antiporters such as SOS1 or NHX1 (Fig. 1-1).

Osmotic stress imposed by high salt or drought is transmitted through at least two pathways: ABA-dependent and ABA-independent. ABA induced expression often relies on the presence of a *cis* acting element called ABA-responsive element (ABRE) (Uno *et al.*, 2000). Increased ABA level under stress conditions up-regulates many LEA-like stress responsive genes, including *RD* (responsive to dehydration), *ERD* (early responsive to dehydration), *KIN* (cold inducible), and *RAB* (responsive to ABA). Two *Arabidopsis* ABA-deficit mutants, *low expression of osmotically responsive gene mutant5 (los5)* and *los6* are impaired in the expression of stress responsive genes, such as *RD29A*, *COR15A*, *COR47*, *RD22*, and *P5CS*, under salt and osmotic stresses and they are also defective in ABA biosynthesis. *LOS5* encodes a molybdenum cofactor sulfuryase (MCSU) and is allelic to *ABA3*, while *LOS6* allelic to *ABAI*, encodes a zeaxanthin epoxidase (ZEP) (Xiong *et al.*, 2001, 2002a). These results demonstrate that stress responsive gene expression under salinity is mediated by ABA. Genetic analysis

indicates that there is no clear line of demarcation between ABA-dependent and ABA-independent pathways and the components involved may often cross talk or even converge in the signaling pathway (Chinnusamy *et al.*, 2004; Seki *et al.*, 2003; Xiong *et al.*, 2002b; Zhu, 2002). Calcium, which serves as a second messenger for various stresses, represents a strong candidate, which can mediate such cross-talk (Sanders *et al.*, 2002).

Salt stress signaling

Salt stress affects cellular ion homeostasis as well as osmotic homeostasis. Hence, excess ions (Na^+ and Cl^-) and osmotic stress-induced turgor change may act as inputs for salt stress signaling. Na^+ entry into cells causes membrane depolarization, which is known to activate Ca^{2+} channels (Zhu, 2002). Salinity induces the biosynthesis and accumulation of ABA and also induces accumulation of ROSs. Current evidence suggests that the primary salt stress signals (ionic and osmotic stress) are transduced through Ca^{2+} as well as receptor kinase pathways, while the secondary salt stress signals such as ABA and H_2O_2 also regulate plant salt tolerance (Chinnusamy *et al.*, 2005).

High salinity results in increased cytosolic Ca^{2+} that is transported from the apoplast as well as the intracellular compartments (Knight *et al.*, 1997). The ionic aspect of salt stress is signaled through the SOS pathway where a calcium-responsive SOS3-SOS2 protein kinase complex controls the expression and activity of ion transporters such as SOS1 (Liu and Zhu, 1998). SOS2 kinase also positively regulates the activities of AtNHX1 and a vacuolar $\text{Ca}^{2+}/\text{H}^+$ exchanger CAX1 and may negatively regulate AtHKT1 (Fig. 1-1, Zhu, 2003). More recently, genetic and biochemical analyses demonstrated that the calcium sensor CBL10 protein physically interacts with SOS2 (CIPK24), and the CBL10-SOS2 complex is associated with the vacuolar compartments that are responsible for salt storage and detoxification in plant cells (Kim *et al.*, 2007). These findings suggest that CBL10 and SOS2 constitute a novel salt-tolerance pathway that regulates the sequestration/compartimentalization of Na^+ in plant cells (Kim *et al.*, 2007). In another report, SOS2 is found to interact with subunits of the vacuolar H^+ -ATPase (V-ATPase) and upregulate its transport activity (Batelli *et al.*,

2007). Therefore, coordination exists between the transporters in the tonoplast and plasma membranes in part through the SOS2 kinase (Cheng *et al.*, 2004; Qiu *et al.*, 2004). Recently, the identification and characterization of the SOS pathway in rice supports the conception that this pathway is conserved across diverse taxa (Martinez-Atienza *et al.*, 2007).

Osmotic stress activates several protein kinases including mitogen-activated protein kinases (MAPKs), which mediate osmotic homeostasis and/or detoxification responses. Plant MAPKs are implicated in signaling development, cell division, hormones, biotic, and abiotic stresses. Salt stress quickly activates MAPKs in plants. Correlative evidence suggests that the ROS mediated signaling under salt stress occurs through MAPKs (Chinnusamy *et al.*, 2004). Transcript levels for a number of protein kinases including a two component histidine kinase MAPKKK, MAPKK and MAPK increase in response to osmotic stress (Mikolajczyk *et al.*, 2000). This ultimately results in the accumulation of osmoprotectants that helps reestablish the osmotic balance, protection from stress damage or repair mechanisms by induction of LEA/dehydrin-type stress genes.

Specificity and cross talk in abiotic stresses

Plants grow in a changing environment and experience various environmental stresses. To respond and adapt to these abiotic stresses, plants have evolved mechanisms for various physiological and metabolic changes that increase their tolerance. It is increasingly clear that signaling pathways are actually part of a more complex hierarchy of signaling networks and that common components are shared between different pathways. Many abiotic stresses induce the biosynthesis and accumulation of ABA, which is known to mediate the expression of many genes in response to osmotic, drought and cold stress. However, the fact the expression of some stress responsive genes is unaffected in ABA-deficient and ABA-insensitive mutants indicates the presence of ABA-independent pathways. It is proposed that at least four signaling pathways functions under drought stress: two are ABA-dependent and two are ABA-independent (Fig. 1-2, Seki *et al.*, 2003). ABA-dependent pathways involve basic leucine-zipper and MYB/MYC type transcription factors that recognize ABRE (Uno *et*

al., 2000) and MYB/MYC recognition sequences in the downstream LEA-like genes (Abe *et al.*, 2003), respectively (Fig. 1- 2). ABA-independent signaling pathways are mediated through transcription factors C-repeat binding factors (CBFs) or dehydration responsive element binding factors (*DREBs*) that activate DRE/CRT *cis*-elements of LEA-type protein encoding genes (Stockinger *et al.*, 1997; Liu *et al.*, 1998; Jaglo-Ottosen *et al.*, 1998).




Fig. 1-2. Transcriptional cascades of salt, drought and cold signal transduction. Salinity works through SOS pathway to reinstate cellular ionic equilibrium or through transcription factors MYB/MYC, bZIP, CBF4/DREB1D, DREB2 to activate downstream *LEA*-like genes. Drought stress works through both ABA-dependent and ABA-independent pathway and shares some components with salt stress. Cold stress works through ICE1 and trigger transcription factors CBF1/DREB1B, CBF2/DREB1C and CBF3/DREB1A to mediate their influence on stress genes mainly via ABA-independent pathway. Transcription factors such as CBF/DREB1, DREB2A, DREB2B, bZIP, MYC (bHLH) and MYB interacts with CRT/DRE, ABRE, MYBRS or E-box elements in the promoter of *LEA*-like genes such as *RD29A*, *RD29B*, *COR15a*, *KIN1*, *ERD10* as well as *RD22* to induce the expression of these genes. (adapted from Xiong *et al.*, 2002b; Chinnusamy *et al.*, 2005; Yamaguchi-Shinozaki and Shinozaki, 2006).

In *Arabidopsis*, the promoter regions of several cold- and drought-induced *COR* genes (*COR6.6*, *COR15a*, and *COR78*) contain a 9 bp consensus sequence named C-repeat (CRT)/drought responsive elements (DRE), which stimulates gene expression in response to cold, drought, and salt, but not in response to the exogenous application of ABA (Yamaguchi-Shinozaki and Shinozaki, 1993). Using the yeast one-hybrid system, five different DRE/CRT binding genes, namely, *CBF1(DREB1B)*, *CBF2(DREB1C)*, *CBF3(DREB1A)*, *DREB2A* and *DREB2B* were isolated and characterized (Stockinger *et al.*, 1997; Gilmour *et al.*, 1998; Liu *et al.*, 1998; Medina *et al.*, 1999). The three *DREB1* genes are specifically induced by cold whereas the two *DREB2* genes are specifically induced by drought and salt stresses (Liu *et al.*, 1998; Nakashima *et al.*, 2000), and all of them are able to transcriptionally activate the expression of *COR* genes. Another CBF transcription factor, *CBF4* was later isolated from *Arabidopsis* (Haake *et al.*, 2002). *CBF4* gene expression is up-regulated by drought stress and ABA treatment, but not by low temperature, and it was also found that ABA biosynthesis is required for the proper drought-induced induction of *CBF4* expression (Haake *et al.*, 2002, Fig. 1-2). An example of cross talk between ABA-dependent and ABA-independent stress-responsive pathways is well documented by the transcription of the *COR* genes. As the *COR* gene promoters contains both DRE/CRT and ABRE consensus sequences, ABA-independent TFs (*DREB1* and *DREB2*) and ABA-dependent basic leucine zipper (bZIP) TFs exist (Fig. 1-2, Uno *et al.*, 2000). The discovery and characterization of bZIP type ABRE binding factors (ABF or AREB) demonstrated that ABA is required for their full activations (Choi *et al.*, 2000; Uno *et al.*, 2000). More direct evidence for the existence of cross talk between abiotic stresses comes from the microarray profiling of transcriptome changes under cold, salt, drought and osmotic stresses (Kreps *et al.*, 2002; Seki *et al.*, 2002). For instance, Seki *et al.* (2002) found that there is greater cross talk between drought and high salinity than between cold and salt stress signaling processes by microarray analysis.

A MAPK cascade, which has already been discussed previously in the context of ABA- and osmotic mediated signalling is believed to be the strongest evidence for cross-talk during abiotic stress signalling in plants based on the observation that diverse signals produced by different stress pathways converge at this cascade (Chinnusamy *et*

al., 2004). MAPK cascades are vital to fundamental physiological functions involved in hormonal responses, cell cycle regulation, abiotic stress signaling, and defense mechanisms (Tena *et al.*, 2001). The Arabidopsis genome encodes approximately 60 MAPKKKs, 10 MAPKKs and 20 MAPKs (Tena *et al.*, 2001). As signals transduced to the 60 MAPKKKs have to be transduced through 10 MAPKKs to 20 MAPKs, it provides a scope for cross-talk between different stress signals. Members of MAPK cascades are activated by H₂O₂, ABA and by both abiotic (wounding, drought, salt, cold), and biotic stresses in Arabidopsis, rice and tobacco (Mizoguchi *et al.*, 1996; Ichimura *et al.*, 1998, 2000; Kovtun *et al.*, 2000; Yuasa *et al.*, 2001; Xiong and Yang 2003), which suggests that MAPK cascades act as points of convergence in stress signaling. Ichimura *et al.* (1998) identified a putative Arabidopsis MAPK cascade consisting of AtMEKK1, AtMEK1/AtMKK2, and AtMPK4, in which salt stress induces the expression and activity of AtMEKK1 (Ichimura *et al.*, 2000), which activates AtMPK4 in vitro (Huang *et al.*, 2000). The Arabidopsis MAPK phosphatase 1 (*mkp1*) mutant is resistant to salinity but hypersensitive to genotoxic stress induced by UV-C. A yeast two-hybrid screen showed that MKP1 could interact with MPK3, MPK6 and MPK4, and the interaction was strongest with MPK4 (Ulm *et al.*, 2002). Previous studies suggested that MPK3, MPK4 and MPK6 are implicated in salt stress signaling (Mizoguchi *et al.*, 1996; Ichimura *et al.*, 2000). Taken together, these suggest that MKP1 is a negative regulator of salt stress signaling through MAPK, while it functions as positive regulator in genotoxic stress tolerance (Ulm *et al.*, 2002). Microarray analysis of *mkp1* identified that the transcript level of a putative vacuolar Na⁺/H⁺-antiporter (At4g23700, CHX17) was increased under salt stress (Ulm *et al.*, 2002), which indicates that this gene may be upregulated by a MAPK cascade that is negatively controlled by MPK1. Transient expression studies in Arabidopsis mesophyll protoplasts showed that MPK3 and MPK4 function downstream of the flagellin receptor FLS2 (Asai *et al.*, 2002). Studies from alfalfa and tobacco also showed that a MAPK cascade is a node of cross-talk in biotic and abiotic stress signaling (Baluska *et al.*, 2000; Cardinale *et al.*, 2000). Thus biotic and abiotic stress signals converge at the level of MAPK cascades (Cardinale *et al.*, 2002; Fujita *et al.*, 2006).

Strategies of developing salt tolerant plants

Plant growth under salt stress depends on the reestablishment of proper cellular ion homeostasis with other concomitant processes such as detoxification and growth regulation. Low cytosolic Na^+ content is maintained through the concerted interplay of regulated ion uptake, vacuolar compartmentation, and active extrusion to the extracellular milieu (Zhang *et al.*, 2004). Increased salt tolerance requires identification of genes and germplasm that perform these processes.

Unlike plant resistance to biotic stresses, which is mostly dependent on monogenic traits, the responses to abiotic stresses are multigenic and thus more difficult to control and engineer (Vinocur and Altman, 2005). There are two main strategies of producing salt-tolerant plants based on genetic information: marker-based breeding through QTL mapping and transgenic approaches either by introducing novel genes or altering the expression levels of existing genes to affect the degree of salt stress tolerance. QTLs and marker-assisted selection have advantages over direct phenotypic screening. Ren *et al.* (2005) reported that allelic variation in *OsHKT8*, which encodes a Na^+ transporter, contributes to enhanced capacity of a salt-tolerant variety of rice to maintain shoot K^+ homeostasis under NaCl stress. The drawbacks in using marker-assisted breeding are “linkage drag” of undesirable traits due to the large size of regions of chromosomes identified by QTLs and, the influence of environment and genetic background on the QTLs that are identified (Flowers, 2004).

Transgenic strategies for increasing salt stress tolerance rely on the expression of genes that are involved in signaling, ion homeostasis, or enzymes for the synthesis of functional and structural metabolites (Table 1-1), which are discussed in more details as below.

Ion transporters

Under salt stress condition, plants use three strategies to maintain a high K^+/Na^+ ratio in the cytosol including extrusion of Na^+ ions out of the cytoplasm, vacuolar compartmentation of Na^+ ions and restricting the Na^+ inflow. As expected, overexpression of *SOS1* and *NHX1* in transgenic Arabidopsis, cotton, tomato and canola plants shows a significant increase in salt tolerance (Table 1-1, Apse *et al.*, 1999; He *et al.*, 2005; Shi *et al.*, 2003; Zhang *et al.*, 2001; Zhang and Blumwald, 2001). Antisense inhibition of

Table 1-1. Engineered salt stress tolerance through genetic manipulation of plant-source genes.

Gene	Gene Source	Transgenic plant	Method	Traits	Reference
Ion Transporters					
AVP1 (vacuolar H ⁺ -ATPase)	Arabidopsis	Arabidopsis	OX	salt tolerance	Gaxiola <i>et al.</i> , 2001
H ⁺ -ATPase (plasma membrane proton pump ATPase)	<i>N. plumbaginifolia</i>	tobacco	OX without autoinhibit ory domain	increased salt tolerance	Gévaudant <i>et al.</i> , 2007
HKT1 (high affinity K ⁺ transporter)	wheat	wheat	AS	salt tolerance	Laurie <i>et al.</i> , 2002
NHX1 (vacuolar Na ⁺ /H ⁺ antiporter)	Arabidopsis	Arabidopsis, canola, tomato, cotton	OX	salt tolerance	Apse <i>et al.</i> ,1999; Zhang <i>et al.</i> ,2001; Zhang and Blumwald, 2001; He <i>et al.</i> , 2005
NHX1 (vacuolar Na ⁺ /H ⁺ antiporter)	<i>A. gmelini</i>	rice	OX	salt tolerance	Ohta <i>et al.</i> ,2002
SOS1 (plasma membrane Na ⁺ /H ⁺ antiporter)	Arabidopsis	Arabidopsis	OX	salt tolerance	Shi <i>et al.</i> ,2003
Transcription Factors					
Alfin1	alfalfa	alfalfa	OX	salt tolerance	Winicov, 2000
ABF2(bZIP)	Arabidopsis	Arabidopsis	OX	Enhanced tolerance of drought, salt, heat, oxidative stress, but root elongation inhibited by ABA	Kim <i>et al.</i> , 2004
CAP2(AP2)	chickpea	tobacco	OX	tolerance to dehydration and salt stress	Shukla <i>et al.</i> , 2006
CpMYB10	<i>C.plantagineum</i>	Arabidopsis	OX	drought and Salt tolerance	Villalobos <i>et al.</i> , 2004
DREB1A(CBF3)	Arabidopsis	Arabidopsis	OX	freezing, salt, and drought tolerance	Kasuga <i>et al.</i> ,1999

CBF3	Arabidopsis	rice	OX	increased tolerance to salinity and drought but not to cold	Oh <i>et al.</i> , 2005
GmDREB2(AP2/EREBP family)	soybean	Arabidopsis	OX	enhanced tolerance to drought and high-salt stresses	Chen <i>et al.</i> , 2007
HARDY (AP2/ERF family)	Arabidopsis	Arabidopsis/ rice	activation tagged mutant/OX	Drought and salt tolerance/Improves drought resistance and water use efficiency	Karaba <i>et al.</i> , 2007
OsMYB3R-2	rice	Arabidopsis	OX	increased tolerance to freezing, drought, and salt stress	Dai <i>et al.</i> , 2007
OsNAC6	rice	rice	OX	exhibited growth retardation and low reproductive yields. showed improved tolerance to dehydration and high-salt stresses, and increased tolerance to blast disease	Nakashima <i>et al.</i> , 2007
SNAC1(NAC TF)	rice	rice	OX	enhances drought resistance and salt tolerance	Hu <i>et al.</i> , 2006
StEREBP1(Ethylene responsive element binding protein 1)	potato	potato	OX	enhanced tolerance to cold and salt stress	Lee <i>et al.</i> , 2007
Ts11(EREBP/AP2 TF)	tobacco	tobacco	OX	Salt tolerance	Park <i>et al.</i> , 2001
Compatible osmolytes					
AtALDH3 (Aldehyde dehydrogenase)	Arabidopsis	Arabidopsis	OX	improved drought, salt and oxidative stress tolerance	Sunkar <i>et al.</i> , 2003
BADH(betaine aldehyde dehydrogenase)	barley	rice	OX	improved salt tolerance	Kishitani <i>et al.</i> , 2000
IMT1((myo-Inositol-Omethyltransferase)	ice plant	tobacco	OX	drought and salinity stress	Sheveleva <i>et al.</i> , 1997
P5CS(Pyroline-5-carboxylate synthase)	mothbean	tobacco, rice	OX	improved salt and drought tolerance	Kishor <i>et al.</i> , 1995; Zhu <i>et al.</i> , 1998
ProDH(Proline dehydrogenase)	Arabidopsis	Arabidopsis	AS	improved salt tolerance	Nanjo <i>et al.</i> , 1999
ROS related					

Ascorbate peroxidase	Arabidopsis	tobacco	OX	enhanced tolerance to salt, PEG and water stresses	Badawi <i>et al.</i> , 2004
FeSOD (superoxide dismutase)	Arabidopsis	tobacco	OX	improved salt and oxidative stress tolerance	Van Camp <i>et al.</i> , 1996
glutathione peroxidase (GPX)	<i>Chlamydomonas</i>	tobacco	OX	tolerant to salinity (up to 250 mM NaCl) as well as chilling stress	Yoshimura <i>et al.</i> , 2004
GST(Glutathione-S-transferase)	tobacco	tobacco	OX	stimulated seedling growth under chilling and salt stress	Roxas <i>et al.</i> , 1997
MDAR(Monodehydroascorbate reductase)	Arabidopsis	tobacco	OX	enhanced tolerance against ozone, salt and PEG stress	Eltayeb <i>et al.</i> , 2007
Kinases					
AtGSK1 (GSK3/shaggy-like gene)	Arabidopsis	Arabidopsis	OX	improved salt and drought tolerance	Piao <i>et al.</i> , 2001
AtNDPK2 (Nucleotide diphosphate kinase)	Arabidopsis	Arabidopsis	OX	enhanced tolerance to salt, cold and methyl viologen treatments	Moon <i>et al.</i> , 2003
CIPK15(calcineurin B-like protein-interacting protein kinases)	rice	rice	OX	improved salt tolerance	Xiang <i>et al.</i> , 2007
OsMAPK5	rice	rice	OX	increased tolerance to drought, salt, and cold stresses	Xiong and Yang, 2003
MKK2(MAPK kinase)	Arabidopsis	Arabidopsis	OX	increased freezing and salt tolerance	Teige <i>et al.</i> , 2004
NPK1(MAPKKK)	tobacco	tobacco	OX	increased tolerance to heat shock, freezing, and salt stress	Kovtun <i>et al.</i> , 2000
OsCDPK(Calcium-dependent protein kinase)	rice	rice	OX	improved drought and salt tolerance	Saijo <i>et al.</i> , 2000
LEA-like					
DHN-5(dehydrin)	wheat	Arabidopsis	OX	enhanced tolerance to salt and osmotic stress	Brini <i>et al.</i> , 2007
HVA1	barley	rice, oat, mulberry	OX	tolerance to drought and salt stress	Xu <i>et al.</i> , 1996; Rohila <i>et al.</i> , 2002 ; Oraby <i>et</i>

LEA	<i>B. napus</i>	Chinese cabbage	OX	enhanced ability to grow under salt and drought stress	al., 2005 ; Lal et al., 2007 Park et al., 2005
Miscellaneous					
ABA2/GIN1(short-chain dehydrogenase/reductase)	Arabidopsis	Arabidopsis	OX	exhibited seed germination delay and more tolerance to salinity	Lin et al., 2007
ABR17(pathogenesis-related protein)	pea	Arabidopsis	OX	germinated better in the presence of salt, cold temperature or both salt tolerance	Srivastava et al., 2006
AtCTL1(chitinase-like protein)	Arabidopsis	Arabidopsis	OX	improved tolerance to salt and osmotic stresses	Kwon et al., 2007
AtHAL3(flavoprotein)	Arabidopsis	Arabidopsis	OX	improved tolerance to salt and osmotic stresses	Espinosa-Ruiz et al.,1999
AtHsp17.6A (Small heat shock protein)	Arabidopsis	Arabidopsis	OX	improved drought and salt tolerance	Sun et al., 2001
AtOAT (Ornithine amino transferase)	Arabidopsis	tobacco	OX	improved NaCl or mannitol tolerance	Roosens et al., 2002
AtRab7 (Vesicle trafficking protein)	Arabidopsis	Arabidopsis	OX	improved salt and osmotic stress tolerance	Mazel et al., 2004
AtVAMP7C(vesicle soluble N-ethylmaleimide-sensitive factor attachment protein receptors)	Arabidopsis	Arabidopsis	AS and KO	improved salt tolerance	Leshem et al., 2006
CBL1(calcineurin B-like protein)	Arabidopsis	Arabidopsis	OX	enhanced tolerance to salt and drought but reduced tolerance to freezing	Cheong et al., 2003
CBL4	maize	Arabidopsis	OX	enhanced tolerance to salt at the germination and seedling stages	Wang et al., 2007
Glyoxylase I and Glyoxylase II	<i>Brassica juncea</i> and rice	tobacco	OX	improved salt tolerance	Singla-Pareek et al.,2003
GlyoxylaseI	<i>B.juncea</i>	<i>B. juncea</i>	OX	improved salt tolerance	Veena et al., 1999

GmTP55 (antiquitin-like protein)	soybean	Arabidopsis and tobacco	OX	tolerance to salinity during germination and to water deficit during plant growth	Rodrigues <i>et al.</i> , 2006
GS2 (Chloroplastic glutamine synthetase)	rice	rice		improved salt tolerance	Hoshida <i>et al.</i> , 2000
LTL1(GDSL-motif lipase)	Arabidopsis	Arabidopsis	OX	increases salt tolerance	Naranjo <i>et al.</i> , 2006
MBF1a(multiprotein bridging factor 1a)	maize	Arabidopsis	OX	elevated salt tolerance and resistance to fungal disease	Kim <i>et al.</i> , 2007
MIPS(1-myo-Inositol-1-phosphate synthase)	halophytic rice	tobacco	OX	capable of growth in 200-300 mm NaCl	Majee <i>et al.</i> , 2004
OsISAP1(Zinc-finger protein)	rice	tobacco	OX	tolerance to cold, dehydration, and salt stress	Mukhopadhyay <i>et al.</i> , 2004
PDH45 (DNA helicase 45)	pea	tobacco	OX	improved salt tolerance	Sanan-Mishra <i>et al.</i> , 2005
PR10(Pathogenesis-related)	pea	canola	OX	enhanced germination and growth in the presence of NaCl	Srivastava <i>et al.</i> , 2004
STO (CONSTANS-like protein)	Arabidopsis	Arabidopsis	OX	root growth was increased by 33-70% in the transgenic plants under salt stress	Nagaoka and Takano 2003
ZAT10(C ₂ H ₂ -EAR zinc finger protein)	Arabidopsis	Arabidopsis	OX, RNAi and KO	enhanced tolerance to salinity	Mittler <i>et al.</i> , 2006

OX, overexpression; KO, knock-out; AS, anti-sense

the wheat high affinity K^+ transporter HKT1 has also resulted in improved salt tolerance (Laurie *et al.*, 2002). It was shown earlier that transgenic plants overexpressing an Arabidopsis vacuolar H^+ - pyrophosphatase (AVP1) showed higher tolerance to salinity and drought conditions (Gaxiola *et al.*, 2001).

Transcription factors

Previous researches have identified many transcription factors that are involved in the relaying of stress signals downstream of the sensors to bring about a specific effect.

Classically, two pathways have been implicated in the induction of these genes under stress conditions- an ABA independent and an ABA dependent pathway (Fig. 1-2, Xiong *et al.*, 2002b; Yamaguchi-Shinozaki and Shinozaki, 2006; Zhu, 2002). CBFs /DREBs are the transcription factors that bind to CRT/DRE cis elements in the stress induce promoters in an ABA-independent pathway and, ABFs bind to ABRE cis elements in an ABA-dependent pathway (Yamaguchi-Shinozaki and Shinozaki, 2006). Overexpression of *CBF3/DREB1A* confers drought, salt and freezing stress tolerance, however, enhanced expression under the constitutive CaMV35S promoter resulted in severe growth retardation in Arabidopsis (Kasuga *et al.*, 1999). This dwarfing phenotype was not found when the *CBF3* expression is regulated via a stress-inducible *RD29A* promoter (Kasuga *et al.*, 2004). Another group overexpressed Arabidopsis *CBF3* and *ABF3* in rice and observed increased tolerance to salinity and drought without any penalty on rice growth (Oh *et al.*, 2005), which may be partly explained by the differences in the fine regulation of transcript accumulation in the monocot rice and dicot Arabidopsis, although the inner mechanism is unclear. Overexpression of another Arabidopsis ABRE-binding bZIP gene-*ABF2* in Arabidopsis showed tolerance to multiple stresses and altered sensitivity to ABA (Kim *et al.*, 2004). *ABF2* overexpression also promoted glucose mediated inhibition of seedling development (Kim *et al.*, 2004). Dai *et al.* (2007) found that overexpression of a rice R1R2R3 MYB gene, *OsMYB3R-2*, increases tolerance to freezing, drought, and salt stress in transgenic Arabidopsis (Table 1-1).

NAC (NAM, ATAF1,2, CUC2) transcription factors are involved in many diverse plant functions (Olsen *et al.*, 2005). A rice salt-inducible NAC (*SNAC1*) gene, when overexpressed in rice, enhanced its tolerance to both drought and salt stress (Hu *et al.*,

2006). Transgenic rice plants overexpressing another rice NAC gene-*OsNAC6*, exhibited growth retardation and low reproductive yields with improved tolerance to dehydration and high-salt stresses, and also increased tolerance to blast disease (Nakashima *et al.*, 2007).

Wang *et al.* (2004) identified a tomato jasmonate (JA) and ethylene responsive factor, *JERF3*, which is inducible by ethylene, JA, cold, salt and ABA in tomato (Wang *et al.*, 2004). Overexpression of *JERF3* in tobacco results in the induction of pathogenesis related genes and enhanced tolerance towards salinity (Wang *et al.*, 2004).

Signaling components

Calcium is known to be an important signal transducer for many stress responsive genes. To date, three major classes of Ca^{2+} sensors have been characterized in plants and they are calmodulin (CaM), CDPKs and CBLs (Yang and Poovaiah, 2003). Several lines of evidence suggest that all these three classes of Ca^{2+} sensors are involved in stress signal transduction (Luan *et al.*, 2002). One CaM was found to bind to a transcription factor MYB2 and was reported to enhance its DNA binding activity. Overexpression of this CaM in *Arabidopsis* upregulated the transcription of MYB2-regulated genes including *P5CS1* which is known to confer salt tolerance by overproducing proline (Yoo *et al.*, 2005). The mechanism by which calcium regulates sodium homeostasis via SOS pathway has been well elucidated (Zhu, 2002). Xiang *et al.* (2007) systemically characterized 30 rice *CIPK* genes (*OsCIPK01-OsCIPK30*) and studied their responses to various environmental stresses. Three *CIPK* genes, *OsCIPK03*, *OsCIPK12*, and *OsCIPK15* were overexpressed in rice and transgenic plants showed significantly improved tolerance to cold, drought, and salt stress, respectively. In another study, an *OsCDPK* was overexpressed in rice which showed tolerance to salt and drought stress (Saijo *et al.*, 2000). In addition, MAPK-mediated signaling cascades have also been suggested to play an important role in stress responses by sensing ROS and phosphorylation of regulatory proteins. Transgenic tobacco plants that constitutively expressed an active *Arabidopsis* MAPKKK, *NPK1*, displayed better tolerance to salt and temperature shocks than wild-type plants (Kovtun *et al.*, 2000). In rice, overexpression of an ABA-inducible *OsMAPK5* increased tolerance to drought, salt, and cold stresses, whereas suppression of *OsMAPK5*

expression by RNAi significantly enhanced resistance to fungal (*Magnaporthe grisea*) and bacterial (*Burkholderia glumae*) pathogens (Xiong and Yang 2003), suggesting a diverse role of some MAPK in abiotic and biotic stress. Similarly, transgenic *Arabidopsis* that overexpressed a MAPK kinase, *MKK2*, showed tolerance to salt and freezing stresses (Teige *et al.*, 2004).

Osmolytes/compatible solutes

Osmolytes/compatible solutes such as sugars, polyols, amino acids are synthesized and accumulated intracellularly to maintain cell turgor. Over-production of compatible osmolytes such as proline, glycine betaine, trehalose, and mannitol also improve plants' resistance to drought and salt stresses, though the degree to which transformed plants' ability to tolerate salinity stress is not necessarily correlative with the amounts of osmoprotectants attained (Chen and Murata, 2000). In most of these cases, the source of such genes has been *E.coli* and yeast (Chen and Murata, 2000), although a few examples of plant-source genes involved in betaine and proline synthesis have also been reported (Kishor *et al.*, 1995; Kishitani *et al.*, 2000; Nanjo *et al.*, 1999; Zhu *et al.*, 1998).

A problem with the use of compatible solutes to increase salt tolerance is that there are metabolic limitations on the absolute levels of target osmolyte that can be accumulated and polyol over-accumulating transgenic plants show sterility (Apse and Blumwald, 2002). Further studies may be directed towards dissecting the signaling cascades regulating the compatible solute biosynthesis in higher plants.

ROS scavenging

Removal or degradation of ROSs accumulated in cells by a variety of enzymes can also contribute to stress tolerance. For instance, Roxas *et al.* (1997) found that overexpression of glutathione S-transferase (GST) /glutathione peroxidase (GPX) enhances the growth of transgenic tobacco seedlings during chilling and salt stress. Ascorbate peroxidase (APX) is a critical component of the ROS scavenging system in plant cells (Apel and Hirt, 2004). An APX cDNA from *Arabidopsis* was fused to the chloroplast transit peptide of glutathione reductase (GR) and overexpressed in the chloroplasts. The transgenic plants showed enhanced tolerance to salt and water stress (Badawi *et al.*, 2004). In another report, a *Chlamydomonas* GPX was overexpressed in tobacco in the

cytosol or chloroplasts. The transgenic plants showed increased tolerance to oxidative, chilling and salt stress, indicating a high level of GPX-like protein in tobacco plants functions to remove unsaturated fatty acid hydroperoxides generated under stress and thus, leading to the maintenance of membrane integrity (Yoshimura *et al.*, 2004).

Apart from the various genes discussed above which have been engineered to increase plants tolerance to salt stress as well as to other types of abiotic stresses, there are many other genes which have also been tested in enhancing stress tolerance including salt stress (Table 1-1), although their roles in stress responses are less understood.

However, it should be noted that overexpression of stress responsive genes may not always impart stress tolerance. A stress responsive gene may be involved in stress tolerance or just an effect of stress. Thus, it is not surprising to find that overexpression of a stress responsive gene does not bring any expected trait or even makes the transgenic plants more vulnerable to some types of stress. For instance, it was found that overexpression of a tobacco GST in cotton did not show improved tolerance to salinity, chilling or even to herbicides (Light *et al.*, 2005).

A common problem with supposedly salt-tolerant plants produced by biotechnology is that the assessment of stress tolerance in the lab often has little correlation to that in the field, since saline soil is usually associated with alkaline pH, presence of other sodic salts, low humidity as well as transpiration in the field also influences the assessment. Another problem with those supposedly salt-tolerant plants produced in labs is that salt stress tolerance phenotypes were only observed at specific stages such as seed germination or seedling growth. Therefore, those assessed abiotic stress tolerant plants may not be able to tolerate the true stress conditions in the fields at all. More importantly, salt tolerance needs to be evaluated from an aspect of yield, not just survival and biomass (Flowers, 2004). So far, very few genes have been tested in the fields. Transgenic wheat expressing AtNHX1 was field evaluated. Xue *et al.* (2004) overexpressed *AtNHX1* in wheat and found that the transgenic lines showed improved biomass, increased germination rates and higher grain yields in severe saline conditions.

Although much progress has been made in the past two decades in understanding

salt stress signal transduction mechanisms, there is still a lot unknown or unclear. Therefore, it is necessary to understand how plants respond to stress in a systematic way before we apply the results gained in the lab to crop plants to engineer stress tolerance. Arabidopsis will continue to play a critical role in the foreseeable future, not only in the understanding of biological mechanisms of abiotic stress tolerance, but also in providing a facile means for testing in biotechnology.

Conclusions and future perspectives

Salt stress presents an increasing threat to agriculture in many parts of the world. A major challenge faced by plant scientists is to increase crop productivity while maintaining the sustainability of agricultural practices. It is clear that improving crop tolerance to abiotic stresses will play an important role for fully exploiting the productivity potential of crops. In the past few decades, worldwide researchers have produced an enormous amount of information regarding to the different components of salt signaling pathway, the cross talk between salt and other abiotic stresses as well as the utilization of various genes to engineer salt-tolerant plants. So far, many identified key molecular factors that can be used for genetic engineering of salt-tolerant plants include overexpression of specific transcription factors, characterization of dehydrin proteins, overproduction of osmoprotectants, expression of water channel proteins and ion transporters. In recent years, small RNAs such as microRNAs (miRNAs) and small interfering RNAs (siRNAs), which are small noncoding RNAs working as important regulators of mRNA degradation, translational repression, and chromatin modification, have been shown to play important roles in plant growth and development and stress tolerance (Sunkar and Zhu, 2004; Borsani *et al.*, 2005; Fujii *et al.*, 2005; Phillips *et al.*, 2007 ; Sunkar *et al.*, 2007). Overexpression of these miRNAs and characterization of their phenotypes would shed light on their biological functions. Recently, researches on the new model plant, salt cress (*Thellungiella halophila*) have provided new insights into salt and cold tolerance mechanisms in this small winter annual crucifer (Inan *et al.*, 2004; Volkov *et al.*, 2004; Vera-Estrella *et al.*, 2005; Kant *et al.*, 2006; Volkov and Amtmann, 2006; Wang *et al.*, 2006). Salt cress has a short life cycle and a small genome (~230 Mb) with high sequence identity (average 92%) with Arabidopsis. Salt

ress is an extremophile native to harsh environments and can reproduce after exposure to extreme salinity (500 mM NaCl) or cold to -15 °C (Inan *et al.*, 2004). Future priorities can be directed towards the identification of molecules connecting pathways and of key components in each pathway. Past researches in this field suggest that our understanding of plant stress tolerance can be greatly refined by thorough characterization of individual genes and assessing their contribution to stress tolerance. Such experiments indicated that many individual genes appear to have some positive impact on stress tolerance; mainly master switches such as transcription factors or their upstream signaling molecules are promising candidate genes for biotechnology-ical approaches. Therefore, a deeper understanding of the transcription factors and their regulated target genes, the products of the major stress responsive genes and cross talk between divergent signaling components should remain an area of intense research activity in the near future. The knowledge generated through these studies can be applied to engineer transgenic plants that can tolerate stress conditions without showing any growth and yield penalty.

Objectives

The short term objectives of my research are to identify a complete set of salt responsive genes through full-genome microarray profiling and 2D-gel electrophoresis of NaCl-treated *Arabidopsis* roots; and to characterize a few novel salt responsive transcription factor genes through characterizing their expression and function using a variety of techniques including reverse genetics. The long-term goal is to develop salt-tolerant plants which may also be tolerant of other abiotic stresses like drought.

References

- Abe H, Urao T, Ito T, Seki M, Shinozaki K and Yamaguchi-Shinozaki K.** (2003) *Arabidopsis* AtMYC2 (bHLH) and AtMYB2 (MYB) function as transcriptional activators in abscisic acid signaling. *Plant Cell* 15: 63–78
- Allen G J, Murata Y, Chu S P, Nafisi M and Schroeder J I.** (2002) Hypersensitivity of abscisic acid-induced cytosolic calcium increases in the *Arabidopsis* farnesyltransferase mutant *era1-2*. *Plant Cell* 14, 1649–1662

- Alscher RG, Alscher RG, Erturk N and Heath LS.** (2002) Role of superoxide dismutases (SODs) in controlling oxidative stress in plants. *J Exp Bot* 53:1331-1341.
- Amtmann A, Fischer M, Marsh EL, Stefanovic A, Sanders D and Schachtman DP.** (2001) The wheat cDNA LCT1 generates hypersensitivity to sodium in a salt-sensitive yeast strain. *Plant Physiol* 126, 1061–1071
- Apel K and Hirt H.** (2004) Reactive oxygen species: metabolism, oxidative stress, and signal transduction. *Annu Rev Plant Biol* 55: 373–399
- Apse MP, Aharon GS, Snedden WA and Blumwald E.** (1999) Salt tolerance conferred by overexpression of a vacuolar Na⁺/H⁺ antiporter in *Arabidopsis*. *Science* 285:1256-8.
- Apse MP and Blumwald E.** (2002) Engineering salt tolerance in plants. *Curr Opin Biotechnol* 13:146-150
- Apse MP and Blumwald E.** (2007) Na⁺ transport in plants. *FEBS Lett* 581: 2247–2254.
- Asai T, Tena G, Plotnikova J, Willmann MR, Chiu W-L, Gomez-Gomez L, Boller T, Ausubel FM and Sheen J.** (2002) MAP kinase signalling cascade in *Arabidopsis* innate immunity. *Nature* 415, 977–983
- Badawi G H, Kawano N, Yamauchi Y, Shimada E, Sasaki R, Kubo A and Tanaka K** (2004) Over-expression of ascorbate peroxidase in tobacco chloroplasts enhances the tolerance to salt stress and water deficit. *Physiol Plant* 121: 231–237
- Baluska F, Ovecka M and Hirt H.** (2000) Salt stress induces changes in amounts and localization of the mitogen-activated protein kinase SIMK in alfalfa roots. *Protoplasma* 212, 262–267
- Batelli G, Verslues PE, Agius F, Qiu Q, Fujii H, Pan S, Schumaker KS, Grillo S and Zhu JK.** (2007) SOS2 promotes salt tolerance in part by interacting with the vacuolar H⁺-ATPase and upregulating its transport activity. *Mol Cell Biol* 27:7781-7790
- Berthomieu P, Conejero G, Nublat A, Brackenbury WJ, Lambert C, Savio C, Uozumi N, Oiki S, Yamada K, Cellier F, Gosti F, Simonneau T, Essah PA, Tester M, Very AA, Sentenac H and Casse F.** (2003) Functional analysis of AtHKT1 in *Arabidopsis* shows that Na⁺ recirculation by the phloem is crucial for salt tolerance.

EMBO J 22: 2004-2014

Bohnert HJ and Jensen RG. (1996) Strategies for engineering water stress tolerance in plants. *Trends Biotechnol* 14, 89–97

Borsani O, Zhu J, Verslues PE, Sunkar R and Zhu J-K. (2005) Endogenous siRNAs derived from a pair of natural cis-antisense transcripts regulate salt tolerance in *Arabidopsis*. *Cell* 123: 1279–1291

Brini F, Hanin M, Lumbreras V, Amara I, Khoudi H, Hassairi A, Pagès M and Masmoudi K. (2007) Overexpression of wheat dehydrin DHN-5 enhances tolerance to salt and osmotic stress in *Arabidopsis thaliana*. *Plant Cell Rep* 26:2017-2026

Cardinale F, Jonak C, Ligterink W, Niehaus K, Boller T and Hirt H. (2000) Differential activation of four specific MAPK pathways by distinct elicitors. *J Biol Chem* 275, 36734–36740

Cardinale H, Meskiene I, Ouaked F and Hirt H. (2002). Convergence and divergence of stress-induced mitogen-activated protein kinase signalling pathways at the level of two distinct mitogen-activated protein kinase kinases. *Plant Cell* 14, 703–711

Chen M, Wang QY, Cheng XG, Xu ZS, Li LC, Ye XG, Xia LQ and Ma YZ. (2007) GmDREB2, a soybean DRE-binding transcription factor, conferred drought and high-salt tolerance in transgenic plants. *Biochem Biophys Res Commun* 353: 299-305

Chen THH and Murata N. (2000) Enhancement of tolerance of abiotic stress by metabolic engineering of betaines and other compatible solutes. *Curr Opin Plant Biol* 5, 250-257

Cheng Y and Long M. (2007) A cytosolic NADP-malic enzyme gene from rice (*Oryza sativa* L.) confers salt tolerance in transgenic *Arabidopsis*. *Biotechnol Lett.* 29:1129-1134

Cheng NH, Pittman JK, Zhu J-K and Hirschi KD. (2004) The protein kinase SOS2 activates the *Arabidopsis* H^+/Ca^{2+} antiporter CAX1 to integrate calcium transport and salt tolerance. *J Biol Chem* 279: 2922–2926

Cheong YH, Kim KN, Pandey GK, Gupta R, Grant JJ and Luan S. (2003) CBL1, a calcium sensor that differentially regulates salt, drought, and cold responses in *Arabidopsis*. *Plant Cell* 15:1833-1845

- Chinnusamy V, Jagendorf A and Zhu J-K.** (2005). Understanding and improving salt tolerance in plants. *Crop Sci* 45: 437-438
- Chinnusamy V, Schumaker K and Zhu J-K.** (2004) Molecular genetic perspectives on cross-talk and specificity in abiotic stress signalling in plants. *J Exp Bot* 55: 225-236
- Choi H, Hong J, Ha J, Kang J and Kim SY.** (2000) ABFs, a family of ABA-responsive element binding factors. *J Biol Chem* 275:1723-1730
- Christmann A, Moes D, Himmelbach A, Yang Y, Tang Y and Grill E.** (2006) Integration of abscisic acid signalling into plant responses. *Plant Biol (Stuttg)*. 8:314-325.
- Cutler AJ and Krochko JE.** (1999) Formation and breakdown of ABA. *Trends Plant Sci.* 4: 472-478
- Dai X, Xu Y, Ma Q, Xu W, Wang T, Xue Y and Chong K.** (2007) Overexpression of an R1R2R3 MYB gene, OsMYB3R-2, increases tolerance to freezing, drought, and salt stress in transgenic *Arabidopsis*. *Plant Physiol* 143:1739-1751
- Demidchik V and Maathuis FJ.** (2007) Physiological roles of nonselective cation channels in plants: from salt stress to signalling and development. *New Phytol* 175:387-404
- Dubouzet JG, Sakuma Y, Ito Y, Kasuga M, Dubouzet EG, Miura S, Seki M, Shinozaki K and Yamaguchi-Shinozaki K.** (2003) OsDREB genes in rice, *Oryza sativa* L., encode transcription activators that function in drought-, high-salt- and cold-responsive gene expression. *Plant J* 33: 751–763
- Eltayeb AE, Kawano N, Badawi GH, Kaminaka H, Sanekata T, Shibahara T, Inanaga S and Tanaka K.** (2007) Overexpression of monodehydroascorbate reductase in transgenic tobacco confers enhanced tolerance to ozone, salt and polyethylene glycol stresses. *Planta* 225:1255-1264
- Espinosa-Ruiz A, Belles J M, Serrano R and Culianez-Macia F A.** (1999) *Arabidopsis thaliana* AtHAL3: A flavoprotein related to salt and osmotic tolerance and plant growth. *Plant J* 20: 529–539
- Finkelstein RR, Gampala SS and Rock CD.** (2002) Abscisic acid signaling in seeds and seedlings. *Plant Cell* 14:S15-45
- Finkelstein RR and Rock CD.** (2002) Abscisic acid biosynthesis and response. In

Somerville CR, Meyerowitz EM, eds, *The Arabidopsis Book*. American Society of Plant Biologists, Rockville, MD

Flowers TJ. (2004) Improving crop salt tolerance. *J Exp Bot* 55: 307-319

Fujita M, Fujita Y, Noutoshi Y, Takahashi F, Narusaka Y, Yamaguchi-Shinozaki K and Shinozaki K. (2006) Crosstalk between abiotic and biotic stress responses: a current view from the points of convergence in the stress signaling networks. *Curr Opin Plant Biol* 9: 436-442

Fujii H, Chiou TJ, Lin SI, Aung K and Zhu J-K. (2005) A miRNA involved in phosphate-starvation response in Arabidopsis. *Curr Biol* 15: 2038-2043

Gao Y, Zeng Q, Guo J, Cheng J, Ellis BE and Chen JG. (2007) Genetic characterization reveals no role for the reported ABA receptor, GCR2, in ABA control of seed germination and early seedling development in Arabidopsis. *Plant J*, Sep 26; [Epub ahead of print].

Gaxiola RA, Li J, Undurraga S, Dang LM, Allen GJ, Alper SL and Fink GR. (2001) Drought- and salt-tolerant plants result from overexpression of the AVP1 H⁺-pump. *Proc Natl Acad Sci USA*. 98:11444-11449

Gaxiola RA, Rao R, Sherman A, Grisafi P, Alper SL and Fink GR. (1999) The Arabidopsis thaliana proton transporters, AtNHX1 and Avp1, can function in cation detoxification in yeast. *Proc Natl Acad Sci USA* 96:1480-1485

Gévaudant F, Duby G, von Stedingk E, Zhao R, Morsomme P and Boutry M. (2007) Expression of a constitutively activated plasma membrane H⁺-ATPase alters plant development and increases salt tolerance. *Plant Physiol* 144:1763-1776

Gilmour SJ, Zarka DG, Stockinger EJ, Salazar MP, Houghton JM and Thomashow MF. (1998) Low temperature regulation of the Arabidopsis CBF family of AP2 transcriptional activators as an early step in cold-induced COR gene expression. *Plant J* 16:433-442

Glenn EP, Brown JJ and Blumwald E. (1999) Salt tolerance and crop potential of halophytes. *Crit Rev Plant Sci* 18: 227–255

Golldack D and Dietz KJ. (2001) Salt-induced expression of the vacuolar H⁺-ATPase in the common ice plant is developmentally controlled and tissue specific. *Plant Physiol* 125:1643-1654

- Haake V, Cook D, Riechmann JL, Pineda O, Thomashow MF and Zhang JZ.** (2002) Transcription factor CBF4 is a regulator of drought adaptation in Arabidopsis. *Plant Physiol* 130:639-648
- Hasegawa PM, Bressan RA, Zhu J-K and Bohnert HJ.** (2000) Plant cellular and molecular responses to high salinity. *Annu Rev Plant Physiol Plant Mol Biol* 51, 463–499
- He C, Yan J, Shen G, Fu L, Holaday AS, Auld D, Blumwald E and Zhang H.** (2005) Expression of an Arabidopsis vacuolar sodium/proton antiporter gene in cotton improves photosynthetic performance under salt conditions and increases fiber yield in the field. *Plant Cell Physiol* 46:1848-1854
- Hernandez JA, Campillo A, Jimenez A, Alacon JJ and Sevilla F.** (1999) Response of antioxidant systems and leaf water relations to NaCl stress in pea plants. *New Phytol* 141, 241–251
- Hoshida H, Tanaka Y, Hibino T, Hayashi Y, Tanaka A, Takabe T and Takabe T.** (2000) Enhanced tolerance to salt stress in transgenic rice that overexpresses chloroplast glutamine synthetase. *Plant Mol Biol* 43: 103–111
- Hu H, Dai M, Yao J, Xiao B, Li X, Zhang Q and Xiong L.** (2006) Overexpressing a NAM, ATAF, and CUC (NAC) transcription factor enhances drought resistance and salt tolerance in rice. *Proc Natl Acad Sci U S A.* 103:12987-12992
- Huang Y, Li H, Gupta R, Morris PC, Luan S and Kieber JJ.** (2000) ATMPK4, an Arabidopsis homolog of mitogen-activated protein kinase, is activated in vitro by AtMEK1 through threonine phosphorylation. *Plant Physiol* 122, 1301-1310
- Ichimura K, Mizoguchi T, Irie K, Morris P, Giraudat J, Matsumoto K and Shinozaki K.** (1998) Isolation of ATMEKK1 (a MAP kinase kinase kinase)-interacting proteins and analysis of a MAP kinase cascade in Arabidopsis. *Biochem Biophys Res Comm* 253, 532-543
- Ichimura K, Mizoguchi T, Yoshida R, Yuasa T and Shinozaki K.** (2000) Various abiotic stresses rapidly activate Arabidopsis MAP kinases ATMPK4 and ATMPK6. *Plant J* 24, 655-665
- Inan G, Zhang Q, Li P, Wang Z, Cao Z, Zhang H, Zhang C, Quist TM, Goodwin SM, Zhu J, Shi H, Damsz B, Charbaji T, Gong Q, Ma S, Fredricksen M,**

- Galbraith DW, Jenks MA, Rhodes D, Hasegawa PM, Bohnert HJ, Joly RJ, Bressan RA and Zhu JK.** (2004). Salt cress. A halophyte and cryophyte *Arabidopsis* relative model system and its applicability to molecular genetic analyses of growth and development of extremophiles. *Plant Physiol* 135:1718-1737
- Jaglo-Ottosen KR, Gilmour SJ, Zarka DG, Schabenberger O and Thomashow MF.** (1998) *Arabidopsis* CBF1 overexpression induces *cor* genes and enhances freezing tolerance. *Science* 280:104–106
- Jia W, Wang Y, Zhang S and Zhang J.** (2002) Salt-stress-induced ABA accumulation is more sensitively triggered in roots than in shoots. *J Exp Bot* 53:2201-2206
- Jia W, Zhang J and Liang J.** (2001) Initiation and regulation of water deficit-induced abscisic acid accumulation in maize leaves and roots: cellular volume and water relations. *J Exp Bot* 52, 295–300
- Jonak C, Okresz L, Bogre L and Hirt H.** (2002) Complexity, cross talk and integration of plant MAP kinase signaling. *Curr Opin Plant Biol* 5:415–424
- Jones M A, Shen J J, Fu Y, Li H, Yang Z and Grierson CS.** (2002) The *Arabidopsis* Rop2 GTPase is a positive regulator of both root hair initiation and tip growth. *Plant Cell* 14, 763–776
- Karaba A, Dixit S, Greco R, Aharoni A, Trijatmiko KR, Marsch-Martinez N, Krishnan A, Nataraja KN, Udayakumar M and Pereira A.** (2007) Improvement of water use efficiency in rice by expression of HARDY, an *Arabidopsis* drought and salt tolerance gene. *Proc Natl Acad Sci U S A*.104:15270-15275
- Kant S, Kant P, Raveh E and Barak S.** (2006) Evidence that differential gene expression between the halophyte, *Thellungiella halophila*, and *Arabidopsis thaliana* is responsible for higher levels of the compatible osmolyte proline and tight control of Na⁺ uptake in *T. halophila*. *Plant Cell Environ* 29, 1220-1234
- Kasuga M, Liu Q, Miura S, Yamaguchi-Shinozaki K and Shinozaki K.** (1999) Improving plant drought, salt, and freezing tolerance by gene transfer of a single stress-inducible transcription factor. *Nat Biotechnol* 17: 287–291
- Kasuga M, Miura S, Shinozaki K and Yamaguchi-Shinozaki K.** (2004) A combination of the *Arabidopsis* *DREB1A* gene and stress-inducible *rd29A* promoter improved drought and low-temperature stress tolerance in tobacco by gene transfer.

Plant Cell Physiol 45:346–350

Katiyar-Agarwal S, Zhu J, Kim K, Agarwal M, Fu X, Huang A and Zhu J-K. (2006) The plasma membrane Na^+/H^+ antiporter SOS1 interacts with RCD1 and functions in oxidative stress tolerance in Arabidopsis. Proc Natl Acad Sci U S A. 103:18816-18821

Katsuhara M, Koshio K, Shibasaka M, Hayashi Y, Hayakawa T and Kasamo K. (2003) Over-expression of a barley aquaporin increased the shoot/root ratio and raised salt sensitivity in transgenic rice plants. Plant Cell Physiol 44:1378-1383

Kim S, Kang JY, Cho DI, Park JH and Kim SY. (2004) ABF2, an ABRE-binding bZIP factor, is an essential component of glucose signaling and its overexpression affects multiple stress tolerance. Plant J 40:75–87

Kim MJ, Lim GH, Kim ES, Ko CB, Yang KY, Jeong JA, Lee MC and Kim CS. (2007) Abiotic and biotic stress tolerance in Arabidopsis overexpressing the multiprotein bridging factor 1a (MBF1a) transcriptional coactivator gene. Biochem Biophys Res Commun 354: 440-446

Kim BG, Waadt R, Cheong YH, Pandey GK, Dominguez-Solis JR, Schültke S, Lee SC, Kudla J and Luan S. (2007) The calcium sensor CBL10 mediates salt tolerance by regulating ion homeostasis in Arabidopsis. Plant J. 52: 473-484

Kishitani S, Takanami T, Suzuki M, Oikawa M, Yokoi S, Ishitani M, Alvarez-Nakase AM, Takabe T and Takabe T. (2000) Compatibility of glycinebetaine in rice plants: Evaluation using transgenic rice plants with a gene for peroxisomal betaine aldehyde dehydrogenase from barley. Plant Cell Environ. 23:107–114

Kishor PBK, Hong Z, Miao GH, Hu CAA and Verma DPS. (1995) Overexpression of [δ]-pyrroline-5-carboxylate synthetase increases proline production and confers osmotolerance in transgenic plants. Plant Physiol 108:1387–1394

Knight H, Trewavas AJ and Knight MR. (1997) Calcium signalling in *Arabidopsis thaliana* responding to drought and salinity. Plant J 12:1067–1078

Koornneef M, Léon-Kloosterziel KM, Schwartz SH and Zeevaart JA. (1998) The genetic and molecular dissection of abscisic acid biosynthesis and signal transduction in Arabidopsis. Plant Physiol Biochem 36: 83-89

Kovtun Y, Chiu W-L, Tena G and Sheen J. (2000) Functional analysis of oxidative

stress-activated mitogen-activated protein kinase cascade in plants. Proc Natl Acad Sci. USA 97:2940–2945

Kreps JA, Wu Y, Chang HS, Zhu T, Wang X and Harper JF. (2002) Transcriptome changes for *Arabidopsis* in response to salt, osmotic, and cold stress. Plant Physiol 130, 2129-2141

Kuhn J M, Boisson-Dernier A, Dizon M B, Maktabi M H and Schroeder JI. (2006) The protein phosphatase AtPP2CA negatively regulates abscisic acid signal transduction in *Arabidopsis*, and effects of *abh1* on AtPP2CA mRNA. Plant Physiol 140, 127–139

Kwon Y, Kim SH, Jung MS, Kim MS, Oh JE, Ju HW, Kim KI, Vierling E, Lee H and Hong SW. (2007) *Arabidopsis hot2* encodes an endochitinase-like protein that is essential for tolerance to heat, salt and drought stresses. Plant J 49:184-193

Lal S, Gulyani V and Khurana P. (2007) Overexpression of *HVA1* gene from barley generates tolerance to salinity and water stress in transgenic mulberry (*Morus indica*) Transgenic Res. DOI 10.1007/s11248-007-9145-4

Laurie S, Feeney KA, Maathuis FJ, Heard PJ, Brown SJ and Leigh RA. (2002) A role for HKT1 in sodium uptake by wheat roots. Plant J 32: 139–149

Lee HE, Shin D, Park SR, Han SE, Jeong MJ, Kwon TR, Lee SK, Park SC, Yi BY, Kwon HB and Byun MO. (2007) Ethylene responsive element binding protein 1 (StEREBP1) from *Solanum tuberosum* increases tolerance to abiotic stress in transgenic potato plants. Biochem Biophys Res Commun 353:863-868

Lemichez E, Wu Y, Sanchez J P, Mettouchi A, Mathur J and Chua N H. (2001) Inactivation of AtRac1 by abscisic acid is essential for stomatal closure. Genes Dev 15, 1808–1816

Leshem Y, Melamed-Book N, Cagnac O, Ronen G, Nishri Y, Solomon M, Cohen G and Levine A. (2006) Suppression of *Arabidopsis* vesicle-SNARE expression inhibited fusion of H₂O₂-containing vesicles with tonoplast and increased salt tolerance. Proc Natl Acad Sci U S A. 103:18008-18013

Light GG, Mahan JR, Roxas VP and Allen RD. (2005) Transgenic cotton (*Gossypium hirsutum* L.) seedlings expressing a tobacco glutathione S-transferase fail to provide improved stress tolerance. Planta 222: 346–354

- Lin PC, Hwang SG, Endo A, Okamoto M, Koshiha T and Cheng WH.** (2007) Ectopic expression of ABSCISIC ACID 2/GLUCOSE INSENSITIVE 1 in *Arabidopsis* promotes seed dormancy and stress tolerance. *Plant Physiol* 143:745-758
- Liu J and Zhu J-K.** (1998) A calcium sensor homolog required for plant salt tolerance. *Science* 280:1943-1945
- Liu Q, Kasuga M, Sakuma Y, Abe H, Miura S, Yamaguchi-Shinozaki K and Shinozaki K.** (1998) Two transcription factors, DREB1 and DREB2, with an EREBP/AP2 DNA binding domain separate two cellular signal transduction pathways in drought- and low-temperature-responsive gene expression, respectively, in *Arabidopsis*. *Plant Cell* 10: 1391–1406
- Liu X, Yue Y, Li B, Nie Y, Li W, Wu WH and Ma L.** (2007) A G protein-coupled receptor is a plasma membrane receptor for the plant hormone abscisic acid. *Science* 315:1712-1716
- Luan S, Kudla J, Rodriguez-Concepcion M, Yalovsky S, and Gruissem W.** (2002) Calmodulins and calcineurin-B like proteins: Calcium sensors for specific signal response coupling in plants. *Plant Cell* 14: S389–S400
- Majee M, Maitra S, Dastidar KG, Pattnaik S, Chatterjee A, Hait NC, Das KP and Majumder AL.** (2004) A novel salt-tolerant L-myo-inositol-1-phosphate synthase from *Porteresia coarctata* (Roxb.) *Tateoka*, a halophytic wild rice: molecular cloning, bacterial overexpression, characterization, and functional introgression into tobacco-conferring salt tolerance phenotype. *J Biol Chem* 279:28539–28552
- Martinez-Atienza J, Jiang X, Garcideblas B, Mendoza I, Zhu JK, Pardo JM and Quintero FJ.** (2007) Conservation of the salt overly sensitive pathway in rice. *Plant Physiol* 143:1001-1012
- Maurel C and Chrispeels MJ.** (2001) Aquaporins. A molecular entry into plant water relations. *Plant Physiol* 125:135–138
- Mazel A, Leshem Y, Tiwari B S and Levin A.** (2004) Induction of salt and osmotic stress tolerance by overexpression of an intracellular vesicle trafficking protein AtRab7 (AtRabG3e). *Plant Physiol* 134: 118–128
- Medina J, BARGUES M, Terol J, Pérez-Alonso M and Salinas J.** (1999) The *Arabidopsis* CBF gene family is composed of three genes encoding AP2 domain-

containing proteins whose expression is regulated by low temperature but not by abscisic acid or dehydration. *Plant Physiol* 119:463-470

Mikolajczyk M, Awotunde OS, Muszyńska G, Klessig DF and Dobrowolska G. (2000) Osmotic stress induces rapid activation of a salicylic acid-induced protein kinase and a homolog of protein kinase ASK1 in tobacco cells. *Plant Cell* 12:165–178

Mittler R, Kim Y, Song L, Coutu J, Coutu A, Ciftci-Yilmaz S, Lee H, Stevenson B and Zhu JK. (2006) Gain- and loss-of-function mutations in *Zat10* enhance the tolerance of plants to abiotic stress. *FEBS Lett* 580: 6537-6542

Mittler R, Vanderauwera S, Gollery M and Breusegem FV. (2004) Reactive oxygen gene network of plants. *Trends Plant Sci* 9: 490-498

Mizoguchi T, Irie K, Hirayama T, Hayashida N, Yamaguchi-Shinozaki K, Matsumoto K and Shinozaki K. (1996) A gene encoding a mitogen-activated protein kinase kinase kinase is induced simultaneously with genes for a mitogen-activated protein kinase and an S6 ribosomal protein kinase by touch, cold, and water stress in *Arabidopsis thaliana*. *Proc Natl Acad Sci USA* 93, 765-769

Moon H, Lee B, Choi G, Shin D, Prasad DT, Lee O, Kwak SS, Kim DH, Nam J, Bahk J, Hong JC, Lee SY, Cho MJ, Lim CO and Yun DJ. (2003) NDP kinase 2 interacts with two oxidative stress-activated MAPKs to regulate cellular redox state and enhances multiple stress tolerance in transgenic plants. *Proc Natl Acad Sci USA* 100: 358–363

Mukhopadhyay A, Vij S and Tyagi AK. (2004) Overexpression of a zinc-finger protein gene from rice confers tolerance to cold, dehydration, and salt stress in transgenic tobacco. *Proc Natl Acad Sci USA* 101: 6309–6314

Munns R. (2002) Comparative physiology of salt and water stress. *Plant Cell Environ* 25: 239-250

Munns R. (2005) Genes and salt tolerance: bringing them together. *New Phytol* 167: 645-663

Nagaoka S and Takano T. (2003) Salt tolerance-related protein *STO* binds to a Myb transcription factor homologue and confers salt tolerance in *Arabidopsis*. *J Exp Bot* 54:2231-2237

Nakashima K, Shinwari ZK, Sakuma Y, Seki M, Miura S, Shinozaki K and

- Yamaguchi-Shinozaki K.** (2000) Organization and expression of two *Arabidopsis DREB2* genes encoding DRE-binding proteins involved in dehydration- and high-salinity-responsive gene expression. *Plant Mol Biol* 42:657-665
- Nakashima K, Tran LS, Van Nguyen D, Fujita M, Maruyama K, Todaka D, Ito Y, Hayashi N, Shinozaki K and Yamaguchi-Shinozaki K.** (2007) Functional analysis of a NAC-type transcription factor OsNAC6 involved in abiotic and biotic stress-responsive gene expression in rice. *Plant J* 51:617-630
- Nambara E and Marion-Poll A.** (2005) Abscisic acid biosynthesis and catabolism. *Annu Rev Plant Biol* 56:165–185
- Nanjo T, Kobayashi M, Yoshida Y, Kakubari Y, Yamaguchi-Shinozaki K and Shinozaki K.** (1999) Antisense suppression of proline degradation improves tolerance to freezing and salinity in *Arabidopsis thaliana*. *FEBS Lett* 461: 205–210
- Naranjo MA, Forment J, Roldán M, Serrano R and Vicente O.** (2006) Overexpression of *Arabidopsis thaliana* LTL1, a salt-induced gene encoding a GDSL-motif lipase, increases salt tolerance in yeast and transgenic plants. *Plant Cell Environ* 29:1890-1900
- Oh SJ, Song SI, Kim YS, Jang HJ, Kim SY, Kim M, Kim YK, Nahm BH and Kim JK.**(2005) *Arabidopsis* CBF3/DREB1A and ABF3 in transgenic rice increased tolerance to abiotic stress without stunting growth. *Plant Physiol* 138:341–351
- Ohnishi J, Flugge UI, Heldt HW and Kanai R.** (1990) Involvement of Na⁺ in active uptake of pyruvate in mesophyll chloroplasts of some C4 plants: Na⁺/pyruvate cotransport. *Plant Physiol* 94: 950-959
- Ohta M, Guo Y, Halfter U and Zhu J-K.** (2003) A novel domain in the protein kinase SOS2 mediates interaction with the protein phosphatase 2C ABI2. *Proc Natl Acad Sci USA* 100:11771–11776
- Ohta M, Hayashi Y, Nakashima A, Hamada A, Tanaka A, Nakamura T and Hayakawa T.** (2002) Introduction of a Na⁺/H⁺ antiporter gene from *Atriplex gmelini* confers salt tolerance to rice. *FEBS Lett* 532: 279–282
- Olsen AN, Ernst HA, Leggio LL and Skriver K.** (2005) NAC transcription factors: structurally distinct, functionally diverse. *Trends Plant Sci* 10:79–87

- Oraby HF, Ransom CB, Kravchenko AN and Sticklen MB.** (2005) Barley *HVA1* gene confers salt tolerance in R3 transgenic oat. *Crop Sci.* 45:2218–2227
- Pandey S and Assmann S M.** (2004) The *Arabidopsis* putative G protein-coupled receptor GCR1 interacts with the G protein alpha subunit GPA1 and regulates abscisic acid signaling. *Plant Cell* 16, 1616–1632
- Park J, Park C, Lee S, Ham B, Shin R and Paek K.** (2001) Overexpression of the tobacco Tsi1 gene encoding an EREBP/AP2-type transcription factor enhances resistance against pathogen attack and osmotic stress in tobacco. *Plant Cell* 13:1035–1046
- Park BJ, Liu Z, Kanno A and Kameya T.** (2005) Transformation of radish (*Raphanus sativus* L.) via sonication and vacuum infiltration of germinated seeds with *Agrobacterium* harboring a group 3 *LEA* gene from *B. napus*. *Plant Cell Rep.* 24:494–500
- Phillips JR, Dalmay T and Bartels D.** (2007) The role of small RNAs in abiotic stress. *FEBS Lett* 581:3592-3597
- Piao H L, Lim J H, Kim S J, Cheong GW and Hwang I.** (2001) Constitutive overexpression of AtGSK1 induces NaCl stress responses in the absence of NaCl stress and results in enhanced NaCl tolerance in *Arabidopsis*. *Plant J* 27: 305–314
- Qiu QS, Guo Y, Quintero FJ, Pardo JM, Schumaker KS and Zhu JK.** (2004) Regulation of vacuolar Na⁺/H⁺ exchange in *Arabidopsis thaliana* by the SOS pathway. *J Biol Chem* 279: 207–215
- Razem F A, El-Kereamy A, Abrams S R and Hill R D.** (2006) The RNA-binding protein FCA is an abscisic acid receptor. *Nature* 439: 290–294
- Ren ZH, Gao JP, Li LG, Cai XL, Huang W, Chao DY, Zhu MZ, Wang ZY, Luan S and Lin HX.** (2005) A rice quantitative trait locus for salt tolerance encodes a sodium transporter. *Nat Genet* 37: 1141-1146
- Rodrigues SM, Andrade MO, Gomes AP, Damatta FM, Baracat-Pereira MC and Fontes EP.** (2006) *Arabidopsis* and tobacco plants ectopically expressing the soybean antiquitin-like *ALDH7* gene display enhanced tolerance to drought, salinity, and oxidative stress. *J Exp Bot* 57:1909–1918
- Rohila JS, Jain RK and Wu R.** (2002) Genetic improvement of Basmati rice for salt

- and drought tolerance by regulated gene expression of a barley *HVA1* cDNA. *Plant Sci* 163:525–532
- Romeroaranda R, Soria T and Cuartero J.** (2001) Tomato plant—water uptake and plant–water relationships under saline growth conditions. *Plant Sci* 160, 265–272
- Roosens N, Hal Bitar F, Loenders K, Angenon G and Jacobs M.** (2002) Overexpression of ornithine- δ -aminotransferase increases proline biosynthesis and confers osmotolerance in transgenic plants. *Mol Breed* 9: 73–80
- Roxas VP, Smith RK, Allen ER and Allen RD.** (1997) Overexpression of glutathione S-transferase/glutathione peroxidase enhances the growth of transgenic tobacco seedlings during stress. *Nat Biotechnol* 15:988-991
- Rus A, Yokoi S, Sharkhuu A, Reddy M, Lee BH, Matsumoto TK, Koiwa H, Zhu JK, Bressan RA and Hasegawa PM.** (2001) AtHKT1 is a salt tolerance determinant that controls Na⁺ entry into plant roots. *Proc Natl Acad Sci U S A.* 98:14150-14155
- Saez A, Apostolova N, Gonzalez-Guzman M, Gonzalez-Garcia M P, Nicolas C, Lorenzo O and Rodriguez P L.** (2004) Gain-of-function and loss-of-function phenotypes of the protein phosphatase 2C HAB1 reveal its role as a negative regulator of abscisic acid signalling. *Plant J* 37, 354–369
- Saijo Y, Hata S, Kyozuka J, Shimamoto K and Izui K.** (2000) Overexpression of a single Ca²⁺-dependent protein kinase confers both cold and salt/drought tolerance on rice plants. *Plant J* 23: 319–327
- Sanan-Mishra N, Pham XH, Sopory SK and Tuteja N.** (2005) Pea DNA helicase 45 overexpression in tobacco confers high salinity tolerance without affecting yield. *Proc Natl Acad Sci U S A* 102:509-514
- Sanders D, Pelloux J, Brownlee C and Harper JF.** (2002) Calcium at the crossroads of signaling. *Plant Cell* 14 (suppl.): S401–S417
- Schachtman DP, Kumar R, Schroeder JI and Marsh EL.** (1997) Molecular and functional characterization of a novel lowaffinity cation transporter (LCT1) in higher plants. *Proc Natl Acad Sci USA* 94, 11079–11084
- Schroeder JI, Allen GJ, Hugouvieux V, Kwak JM and Waner D.** (2001) Guard cell signal transduction. *Annu Rev Plant Physiol Plant Mol Biol* 52:627-658
- Schwartz SH, Qin X and Zeevaart JA.** (2003) Elucidation of the indirect pathway of

abscisic acid biosynthesis by mutants, genes, and enzymes. *Plant Physiol* 131:1591-1601

Seki M, Kamei A, Yamaguchi-Shinozaki K and Shinozaki K. (2003) Molecular responses to drought, salinity and frost: common and different paths for plant protection. *Curr Opin Biotechnol* 14:194-199

Seki M, Narusaka M, Ishida J, Nanjo T, Fujita M, Oono Y, Kamiya A, Nakajima M, Enju A, Sakurai T, Satou M, Akiyama K, Taji T, Yamaguchi-Shinozaki K, Carninci P, Kawai J, Hayashizaki Y and Shinozaki K. (2002) Monitoring the expression profiles of 7000 *Arabidopsis* genes under drought, cold and high-salinity stresses using a full-length cDNA microarray. *Plant J* 31: 279-292

Shen YY, Wang XF, Wu FQ, Du SY, Cao Z, Shang Y, Wang XL, Peng CC, Yu XC, Zhu SY, Fan RC, Xu YH and Zhang DP. (2006) The Mg-chelatase H subunit is an abscisic acid receptor. *Nature* 443:823-826

Sheveleva E, Chmara W, Bohnert H J and Jensen RG. (1997) Increased salt and drought tolerance by D-ononitol production in transgenic *Nicotiana tabacum* L. *Plant Physiol* 115:1211–1219

Shi H, Ishitani M, Kim C and Zhu JK. (2000) The *Arabidopsis thaliana* salt tolerance gene *SOS1* encodes a putative Na^+/H^+ antiporter. *Proc Natl Acad Sci U S A.* 97:6896-6901

Shi H, Lee BH, Wu SJ and Zhu JK. (2003) Overexpression of a plasma membrane Na^+/H^+ antiporter gene improves salt tolerance in *Arabidopsis thaliana*. *Nat Biotechnol* 21:81-85

Shi H and Zhu J-K. (2002) Regulation of expression of the vacuolar Na^+/H^+ antiporter gene *AtNHX1* by salt stress and ABA. *Plant Mol Biol* 50:543–550

Shukla RK, Raha S, Tripathi V and Chattopadhyay D. (2006) Expression of CAP2, an APETALA2-family transcription factor from chickpea, enhances growth and tolerance to dehydration and salt stress in transgenic tobacco. *Plant Physiol* 142:113-123

Singla-Pareek S L, Reddy MK and Sopory S K. (2003) Genetic engineering of the glyoxalase pathway in tobacco leads to enhanced salinity tolerance. *Proc Natl Acad Sci USA* 100: 14672–14677

- Smirnoff N.**(1993) The role of active oxygen in the response of plants to water deficit and desiccation. *New Phytol* 125: 27-58
- Srivastava S, Fristensky B and Kav NN.** (2004) Constitutive expression of a PR10 protein enhances the germination of *Brassica napus* under saline conditions. *Plant Cell Physiol* 45:1320-1324
- Srivastava S, Rahman MH, Shah S and Kav NN.** (2006) Constitutive expression of the pea ABA-responsive 17 (ABR17) cDNA confers multiple stress tolerance in *Arabidopsis thaliana*. *Plant Biotechnol J* 4:529-549
- Stockinger EJ, Gilmour SJ and Thomashow MF.** (1997) *Arabidopsis thaliana* CBF1 encodes an AP2 domain-containing transcriptional activator that binds to the C-repeat/DRE, a cis-acting DNA regulatory element that stimulates transcription in response to low temperature and water deficit. *Proc Natl Acad Sci U S A.* 94:1035-1040
- Sun W, Bernard C, van de Cotte, Van Montagu M and Verbruggen N.** (2001) *At-HSP17. 6A*, encoding a small heat-shock protein in *Arabidopsis*, can enhance osmotolerance upon overexpression. *Plant J* 27: 407–415
- Sunkar R, Bartels D and Kirch H-H.** (2003) Overexpression of a stress inducible aldehyde dehydrogenase gene from *Arabidopsis thaliana* in transgenic plants improves stress tolerance. *Plant J* 35: 452–464
- Sunkar R, Chinnusamy V, Zhu J and Zhu J-K.** (2007) Small RNAs as big players in plant abiotic stress responses and nutrient deprivation. *Trends Plant Sci* 12:301-309.
- Sunkar R and Zhu J-K.** (2004) Novel and stress-regulated microRNAs and other small RNAs from *Arabidopsis*. *Plant Cell* 16: 2001–2019
- Teige M, Scheikl E, Eulgem T, Dóczi R, Ichimura K, Shinozaki K, Dangl JL and Hirt H.** (2004). The MKK2 pathway mediates cold and salt stress signaling in *Arabidopsis*. *Mol Cell* 15:141-152
- Tena G, Asai T, Chiu WL and Sheen J.** (2001) Plant mitogen-activated protein kinase signaling cascades. *Curr Opin Plant Biol* 4:392-400
- Tester M and Davenport R.** (2003) Na⁺ tolerance and Na⁺ transport in higher plants. *Ann Bot* 91, 503–527
- Ulm R, Ichimura K, Mizoguchi T, Peck SC, Zhu T, Wang X, Shinozaki K and**

- Paszkowski J.** (2002) Distinct regulation of salinity and genotoxic stress responses by *Arabidopsis* MAP kinase phosphatase 1. *EMBO J* 21, 6483-6493
- Uno Y, Furihata T, Abe H, Yoshida R, Shinozaki K and Yamaguchi Shinozaki K.** (2000) *Arabidopsis* basic leucine zipper transcription factors involved in an abscisic acid-dependent signal transduction pathway under drought and high-salinity conditions. *Proc Natl Acad Sci USA* 97: 11632–11637
- Van Camp W, Capiou K, Van Montagu M, Inze D and Slooten L.** (1996) Enhancement of oxidative stress tolerance in transgenic tobacco plants overproducing Fe-superoxide dismutase in chloroplasts. *Plant Physiol* 112: 1703–1714
- Veena, Reddy V S, and Sopory SK.** (1999) Glyoxylase I from *Brassica juncea*: Molecular cloning, regulation and its overexpression confer tolerance in transgenic tobacco under stress. *Plant J* 17: 385–395
- Vera-Estrella R, Barkla BJ, Garcia-Ramirez L and Pantoja O.** (2005) Salt stress in salt cress activates Na transport mechanisms required for salinity tolerance. *Plant Physiol* 139, 1507-1517
- Villalobos M A, Bartels D and Iturriga G.** (2004) Stress tolerance and glucose insensitive phenotypes in *Arabidopsis* overexpressing the *CpMYB10* transcription factor gene. *Plant Physiol* 135: 309–324
- Vinocur B and Altman A.** (2005) Recent advances in engineering plant tolerance to abiotic stress: achievements and limitations. *Curr Opin Biotechnol* 16:123-132
- Volkov V and Amtmann A.** (2006) *Thellungiella halophila*, a salt-tolerant relative of *Arabidopsis thaliana*, has specific root ion channel features supporting K^+/Na^+ homeostasis under salinity stress. *Plant J* 48:342-353
- Volkov V, Wang B, Dominy PJ, Fricke W and Amtmann A.** (2004) *Thellungiella halophila*, a salt-tolerant relative of *Arabidopsis thaliana*, possesses effective mechanisms to discriminate between potassium and sodium *Plant Cell Environ* 27: 1-14.
- Wang B, Davenport R, Volkov V and Amtmann A.** (2006) Low unidirectional sodium influx into root cells restricts net sodium accumulation in *Thellungiella halophila*, a salt-tolerant relative of *Arabidopsis thaliana*. *J Exp Bot* 57, 1161-1170
- Wang J, Zhang H and Allen RD.**(1999) Overexpression of an *Arabidopsis* peroxisomal ascorbate peroxidase gene in tobacco increases protection against

oxidative stress. *Plant Cell Physiol* 40:725-732

Wang M, Gu D, Liu T, Wang Z, Guo X, Hou W, Bai Y, Chen X and Wang G. (2007) Overexpression of a putative maize calcineurin B-like protein in *Arabidopsis* confers salt tolerance. *Plant Mol Biol* Sep 20; [Epub ahead of print]

Wang X Q, Ullah H, Jones A M and Assmann S M. (2001) G protein regulation of ion channels and abscisic acid signaling in *Arabidopsis* guard cells. *Science* 292, 2070–2072

Wang W, Vinocur B and Altman A. (2003) Plant responses to drought, salinity and extreme temperatures: towards genetic engineering for stress tolerance. *Planta* 218: 1–14

Wang Y and Nil N. (2000) Changes in chlorophyll, ribulose biphosphate carboxylase–oxygenase, glycine betaine content, photosynthesis and transpiration in *Amaranthus tricolor* leaves during salt stress. *J Hortic Sci Biotechnol* 75, 623–627

Winicov I. (2000) Alfin1 transcription factor overexpression enhances plant root growth under normal and saline conditions and improves salt tolerance in alfalfa. *Planta* 210: 416–422

Xiang Y, Huang Y and Xiong L. (2007) Characterization of Stress-Responsive CIPK Genes in Rice for Stress Tolerance Improvement. *Plant Physiol* 144:1416-1428

Xiong L, Ishitani M, Lee H and Zhu J-K. (2001) The *Arabidopsis* *LOS5/ABA3* locus encodes a molybdenum cofactor sulfuryase and modulates cold stress- and osmotic stress- responsive gene expression. *Plant Cell* 13: 2063-2083

Xiong L, Lee H, Ishitani M and Zhu J-K. (2002a). Regulation of osmotic stress responsive gene expression by *LOS6/ABA1* locus in *Arabidopsis*. *J Biol Chem* 277: 8588-8596

Xiong L, Schumaker KS and Zhu J-K. (2002b) Cell signaling during cold, drought, and salt stresses. *Plant Cell* 14: S165-183

Xiong L and Yang Y. (2003) Disease resistance and abiotic stress tolerance in rice are inversely modulated by an abscisic acid-inducible mitogen-activated protein kinase. *Plant Cell* 15, 745-759

Xu D, Duan X, Wang B, Hong B, Ho T and Wu R. (1996) Expression of a late embryogenesis abundant protein gene, *HVA1*, from barley confers tolerance to water

deficit and salt stress in transgenic rice. *Plant Physiol* 110: 249-257

Xue ZY, Zhi DY, Xue GP, Zhang H, Zhao YX, and Xia GM (2004) Enhanced salt tolerance of transgenic wheat (*Triticum aestivum* L.) expressing a vacuolar Na⁺/H⁺ antiporter gene with improved grain yields in saline soils in the field and a reduced level of leaf Na⁺. *Plant Sci.* 167: 849–859

Yamaguchi-Shinozaki K and Shinozaki K. (1993) Characterization of the expression of a desiccation-responsive rd29 gene of *Arabidopsis thaliana* and analysis of its promoter in transgenic plants. *Mol Gen Genet* 236(2-3):331-340

Yamaguchi-Shinozaki K and Shinozaki K. (2006) Transcriptional regulatory networks in cellular responses and tolerance to dehydration and cold stresses. *Annu Rev Plant Biol* 57:781-803

Yang T and Poovaiah B W. (2003) Calcium/Calmodulin-mediated signal network in plants. *Trends Plant Sci.* 8: 505–512

Yoo JH, Park CY, Kim JC, Heo WD, Cheong MS, Park HC, Kim MC, Moon BC, Choi MS, Kang YH, Lee JH, Kim HS, Lee SM, Yoon HW, Lim CO, Yun DJ, Lee SY, Chung WS and Cho MJ. (2005) Direct interaction of a divergent CaM isoform and the transcription factor, MYB2, enhances salt tolerance in *Arabidopsis*. *J Biol Chem* 280:3697–3706

Yoshimura K, Miyao K, Gaber A, Takeda T, Kanaboshi H, Miyasaka H and Shigeoka S. (2004) Enhancement of stress tolerance in transgenic tobacco plants overexpressing *Chlamydomonas* glutathione peroxidase in chloroplasts or cytosol. *Plant J.* 37:21–33

Yuasa T, Ichimura K, Mizoguchi T and Shinozaki K.(2001) Oxidative stress activates ATMPK6, an *Arabidopsis* homologue of MAP kinase. *Plant Cell Physiol* 42, 1012-1016.

Zeevaart JAD and Creelman RA. (1988) Metabolism and Physiology of Abscisic Acid. *Annu Rev Plant Physiol Plant Mol Biol* 39: 439-473

Zhang HX and Blumwald E. (2001) Transgenic salt-tolerant tomato plants accumulate salt in foliage but not in fruit. *Nat Biotechnol* 19: 765–768

Zhang HX, Hodson JN, Williams JP and Blumwald E. (2001) Engineering salt tolerant Brassica plants: characterization of yield and seed oil quality in transgenic

plants with increased vacuolar sodium accumulation. Proc Natl Acad Sci USA 98: 12832–12836

Zhang JZ, Creelman RA and Zhu JK. (2004) From laboratory to field. Using information from *Arabidopsis* to engineer salt, cold, and drought tolerance in crops. Plant Physiol 135: 615-621

Zheng Z L, Nafisi M, Tam A, Li H, Crowell D N, Chary SN, Schroeder J I, Shen J and Yang Z. (2002) Plasma membrane-associated ROP10 small GTPase is a specific negative regulator of abscisic acid responses in *Arabidopsis*. Plant Cell 14, 2787–2797

Zhu B, Su J, Chang MC, Verma DPS, Fan YL and Wu R. (1998) Overexpression of a pyrroline-5-carboxylate synthetase gene and analysis of tolerance to water and salt stress in transgenic rice. Plant Sci 139:41–48

Zhu J, Gong Z, Zhang C, Song C P, Damsz B, Inan G, Koiwa H, Zhu J K, Hasegawa PM and Bressan R A. (2002) OSM1/SYP61: a syntaxin protein in *Arabidopsis* controls abscisic acid-mediated and non-abscisic acid-mediated responses to abiotic stress. Plant Cell 14, 3009–3028

Zhu J-K. (2000) Genetic analysis of plant salt tolerance using *Arabidopsis*. Plant Physiol 124: 941-948

Zhu J-K. (2002) Salt and drought stress signal transduction in plants. Annu Rev Plant Biol 53, 247-273

Zhu J-K. (2003) Regulation of ion homeostasis under salt stress. Curr Opin Plant Biol 6, 441-445

Chapter 2: Microarray analysis of Arabidopsis root transcriptome changes in response to salt stress

Introduction

Roots are the primary site of perception and injury for several types of water-limiting stress, including salinity and drought. In many circumstances, the stress-sensitivity of the root limits the productivity of the entire plant (Atkin *et al.*, 1973; Steppuhn and Raney, 2005). The physiological significance of roots is belied by their relative structural simplicity as compared to other organs: roots are largely lacking in some major metabolic pathways such as photosynthesis, and have a stereotypical morphology that is conserved across taxa and throughout the life cycle of individuals. This combination of physiological relevance and structural simplicity has made roots good targets for functional genomic studies. For example, detailed transcriptional profiles have now been resolved to single cell types within roots, and these are now being integrated into regulatory circuits and networks (Birnbaum and Benfey, 2004).

Salinity treatment of plants is also an attractive experimental system. NaCl treatments are easy to apply in laboratory settings, with dosage and timing controlled more precisely than with other major abiotic stresses such as chilling, freezing, and dehydration. So far, at least nine published reports used microarrays to analyze the response of Arabidopsis to NaCl. However, most of these studies have analyzed either cell cultures or whole plants, rather than specific tissues (Seki *et al.*, 2002a; Taji *et al.*, 2004; Takahashi *et al.*, 2004; Gong *et al.*, 2005; Kamei *et al.*, 2005; Ma *et al.*, 2006). Of the previous studies that analyzed roots specifically, none used microarray probe sets representing more than 8,100 of the originally predicted 25,498 genes in the Arabidopsis genome (Arabidopsis Genome, 2000; Chen *et al.*, 2002; Kreps *et al.*, 2002; Maathuis *et al.*, 2003). Although Affymetrix microarrays containing probes for at least 22,591 Arabidopsis genes have been used to profile NaCl responses specifically in

A version of this chapter was published: Jiang Y-Q, Deyholos MK. 2006. Comprehensive transcriptional profiling of NaCl-stressed Arabidopsis roots reveals novel classes of responsive genes. BMC Plant Biol 6, 25

roots, these data were generously deposited to public databases, but without detailed description or analysis in the primary literature. Thus, the absence of available, comprehensive transcriptomic data describing the response of *Arabidopsis* roots to NaCl treatment, in combination with the potential applications of these data in molecular physiology and systems biology, motivated us to conduct this research.

Materials and Methods

Plant growth and stress treatment

We sowed surface-sterilized wild-type seeds of *Arabidopsis thaliana* (ecotype Col-0) on (3.5 mm diameter) agar plugs arrayed on custom-made polypropylene rafts covered by 0.5 mm nylon mesh (Northwest Graphic Products Ltd, Edmonton) from beneath, and cold stratified them at 4°C for 2 d. Each plug consisted of ½ X Murashige and Skoog (MS, Sigma) basal salts, 1% sucrose, and 0.7% Phytagar (Invitrogen), and each raft held approximately 30 seeds, spaced ca. 15 mm apart. Rafts were transferred into opaque plastic tanks containing 1 L of autoclaved ¼ X MS solution (pH5.7) in a growth chamber at 22°C, 70% relative humidity, and light ($\sim 125 \mu\text{E m}^{-2} \text{s}^{-1}$) on a 16h light/8h dark cycle. The solution in the tanks was aerated and replaced with fresh, autoclaved ¼ X MS solution every 4 days to limit growth of microorganisms. At 20 days after stratification (DAS), seedlings were treated with ½ X MS medium (pH5.7) supplemented with 150 mM NaCl beginning at 8:30AM (1.5 h after the start of the light cycle), and roots were harvested at 6, 24 and 48 h post-treatment, and were frozen in liquid nitrogen and stored at -80°C. Untreated controls were harvested in parallel, at the same time points. Approximately 150 plants were harvested and pooled for each combination of time point and treatment, in each biological replicate. This process was repeated in three biologically and temporally independent replicates.

Physiological analyses

Proline concentrations were measured using the method of Bates (Bates *et al.*, 1973), with L-proline (Sigma) as a standard. Anthocyanin concentrations were determined according to Macinelli and Schwartz (1984).

Microarray preparation, hybridization and data extraction

QiagenOperon *Arabidopsis* 70-mer oligonucleotide probes (Array-Ready Oligo Set

Version 1.0) representing 26, 029 genes, plus additional probes for quality control were printed on superamine aminosilane-coated slides (Telechem) in a format of 12 X 4 subarrays with each containing 24 X 24 non-replicate spots. Probe spotting followed previously described methods (Wang *et al.*, 2003)

Total RNA was isolated from both control and stressed root tissues using the RNeasy Plant Midi kit (Qiagen), and was concentrated by Microcon YM-30 (Millipore) to ~ 1.2 µg/µL and quantified by NanoDrop ND-1000 Spectrophotometer. RNA integrity was checked on a formamide denaturing agarose gel. 5 µg total RNA was used to synthesize cDNAs using SuperscriptII (Invitrogen) with RT polyA-capture primers in 3D Array900 kit (Genisphere). Before hybridizations, microarray slides were rehydrated over a 60°C water bath for 5 seconds, snap-dried on a 65°C heating block for 5 seconds and allowed to cool for 1 min. This process was repeated for a total of five times. Then the slides were UV cross-linked by exposing the slides label-side up to 120 mJ for 5 min. Slides were then washed in 1% SDS for 5 minutes, dipped into 95% ethanol for 30 seconds before dried by centrifugation at 1,000 rpm for 1 min. A two-step hybridization was performed, i.e., cDNA from both control and treated tissue were hybridized together at 55°C overnight under a 24X 60 mm cover slip (Fisher Scientific) in a sealed humid 50 ml centrifuge tube. Secondary hybridization with Cy3 and Cy5 dendrimers were performed at 55°C for 4 h and washes were done according to the manufacturer's instruction. Slides were scanned immediately using ArrayWorxe (Applied Precision) and transformed into .tiff images. To avoid bias in the microarray evaluation as a consequence of dye-related differences in labelling efficiency and/or differences in recording fluorescence signals, dye labelling for each paired sample (stressed/control) was reversed in two individual hybridizations. Therefore, two hybridizations were performed for each of three independent biological replicates at each time point.

Microarray data analysis

Background-subtracted spot intensities were measured by Spotfinder v3.0, and normalized by the Lowess method in MIDAS v2.19 (Saeed *et al.*, 2003). Spots were defined as detectable above background if their signal intensity in at least one channel was greater than two standard-deviations above the mean signal intensities of all blank and

randomized negative control spots, and if they were detectable in at least two-thirds of arrays at any one time point. Next, MeV(v3.1) was used to find the differentially expressed genes between control and NaCl treatment. To do so, the integrated spot intensities were loaded into MeV and we used a two-channel spot intensity cut-off value of 1,000. Significance analysis of microarrays (SAM) contained in the was applied to find genes for which the fold change differed significantly from zero in a multiclass test (with inferred values at time point 0), using a false discovery rate (FDR) of $\leq 5\%$ (Tusher *et al.*, 2001). Fold changes (treated/control) were calculated in log₂ scale. Assignment of genes to temporal expression profiles and detection of statistically enriched gene families within each profile was conducted using STEM (Short Time Series Expression Miner) software, with maximum number of model profiles set to 16, and maximum unit change set to 3 (Ernst and Bar-Joseph, 2006). The input data for STEM analysis consisted of all normalized, intensity-filtered data points identified by SAM analysis as statistically distinct from 0. The p-values derived from STEM analysis were corrected for multiple hypothesis testing using a FDR of $< 5\%$ (Draghichi, 2003).

Quantitative PCR

Aliquots of the same RNA samples used for microarray analysis were also used for real-time PCR. Reverse transcription (RT) was performed with 2 μg of total RNA pretreated with DNase I (Ambion) to obtain cDNA with SuperScript II and Oligo(dT)₁₂₋₁₈ (Invitrogen) as the primer in a 20 μL reaction volume. Each cDNA sample was diluted 1:4 in sterile ddH₂O, and 1 μL of this dilution was used as template for qPCR. Primers for the PCR reactions were designed by PrimerSelect 5.00 of DNASTar (DNASTAR Inc) to have a T_m of $\sim 60^\circ\text{C}$ and an optimal annealing temperature of 53-55 $^\circ\text{C}$ with the length of the amplicons between 80 and 200 bp. Real-time PCR was performed with QuantiTech PCR SYBR Green I Kit (Qiagen) in 20 μL reactions using the LightCycler 1.5 System (Roche Diagnostics) according to the manufacturer's instruction. Each PCR reaction contained 1 μL of cDNA, 0.5 μM of each of the primers and 10 μL of master mix. The initial denaturing time was 15 min at 95 $^\circ\text{C}$, followed by 35 cycles consisting of 94 $^\circ\text{C}$ for 15 s, 53 $^\circ\text{C}$ for 30 s, and 72 $^\circ\text{C}$ for 20 s with a single fluorescence measurement. A melting curve (65 $^\circ\text{C}$ –95 $^\circ\text{C}$ with a

heating rate of 0.05°C s⁻¹ and a continuous fluorescence measurement) was run after the PCR cycles followed by a cooling step at 40°C.

For relative quantification, amplification efficiencies (*E*) for each gene were determined as follows: a portion of cDNAs transcribed from 5 µg of total root RNA was diluted with sterile ddH₂O to be 10⁻¹, 10⁻², 10⁻³ and 10⁻⁴. Standard curves for each gene was performed using 1 µL of the original and diluted cDNAs to cover the range of all template concentrations. Real-time PCR efficiencies (*E*) were calculated from the given slopes in the LightCycler software of the standard curves according to the equation: $E = 10^{-1/\text{slope}}$. Crossing points (CP), defined as the point at which the fluorescence rises above the background fluorescence was determined using the “Fit Point Method” in the LightCycler software 3.5.3 (Roche Diagnostics). Gene-specific PCR efficiency was used to calculate the expression of target genes relative to the expression of UBI10 reference gene according to the format: $\text{Ratio} = (E_{\text{target}})^{\Delta \text{CP}_{\text{target}}(\text{control} - \text{sample})} / (E_{\text{ref}})^{\Delta \text{CP}_{\text{ref}}(\text{control} - \text{sample})}$ (Paffl 2001).

The genes tested for expression levels by quantitative PCR were as follows:

At1g44830 (AP2/EREBP), forward primer (FP), 5'-AGTTGGGGTTCATGGGT TTCA-3', reverse primer (RP), 5'-ACATAGGAGTGCTGCGTCGTAGG-3'; *WRKY17* (At2g24570), FP, 5'-CAGCGAGGAAACACGTGGAAAGAGC -3', RP, 5'-CAAGC-CGAACCAAACACCAAACCAC -3'; *MYB15* (At3g23250), FP, 5'-AAATACTTGG-CAATAGATGGTCAG -3', RP, 5'-ACGGATTCAGATTTTGGTTTAGTA-3'; *XTH18* (At4g30280), FP, 5'-AGGCTCCTTTCACCGCTTCTA -3', RP, 5'-ACTC TGTACCCCTTTCATTCTTGTCTG -3'; At4g01550 (*NAC*), FP, 5'-AGAGAAGAA GAGCTGGTTTATTACA -3', RP, 5'-TTCATCGCCGGGTTC AAGT -3'; At2g04070 (*MATE*), FP, 5'-TCTTTCTGGTTTTTCGCTATGACACT -3', RP, 5'-ATTGCCGC TGATGG-GACTC -3'; At4g26410, FP, 5'-GAGCTGAAGTGGCTTCCATGAC-3', RP, 5'-GGT-CCGACATACCCATGATCC-3'; *CYP83B1/SUR2* (At4g31500), FP, 5'-ACTCTTGAC-CC TAACCGCCCTAAAC-3', RP, 5'-TGCAGCCGCGGTGTCAGT-3'; *bHLH92* (At5g43650), FP, 5'-CACGGAAGTGGATGATGTC-3', RP, 5'-GTC GCCGGCTT-CTTTCCTTCTCT-3'. *WRKY25* (At2g30250), FP, 5'-TGGTTCTTCC GGCGTTGACTGTTA-3', RP, 5'-GTGAAATCGGAAGAG-GTGGTGGTTG-3'; *WRKY33* (At2g38470), Fwd, 5'-TTACGCCACAAACAGAGCAC-3', Rvs, 5'-CCAA

AAGGCCCGGTATTAGT-3'; At3g17860 (*ZIM*), FP, 5'- GGCTCCAACAGTGGCAT TACCTCT -3', RP, 5'-TATGGGGATACGCTCGTGACCCTT -3'; *PER21* (At2g37130), FP, 5'- TGCGAGAGACGGTATTGTCA-3', RP, 5'- TCTCCCAAGTA GCTCCCTCT -3'; *PER27* (At3g01190), FP, 5'- CGCAATGGTTGCACTTGA-3', RP, 5'-TGAGCGAAA-ATCGCTGATAA -3'; *PER69* (At5g64100), FP, 5'- CTCTTG TTGGCGGACACA-3', RP, 5'- GTCGATTGATGGGTCAGGTT -3'.

The internal control gene was *UBQ10* (At4g05320), which is a suitable reference gene for abiotic stress analysis (Czechowski *et al.*, 2005). The primer sequences are 5'- TAC TCT TCA CTT GGT CCT GCG TCT TC-3' (FP) and 5'- GGC CCC AAA ACA CAA ACC ACC AA-3' (RP). qPCR for each gene was done on three biological replicates with duplicates for each biological replicate. The relative transcript level was determined for each sample, normalized using the *UBQ10* cDNA level, and averaged over the six replicates.

Results and Discussion

Whole-plant responses to salt treatment.

We applied a salt-shock treatment to 20 DAS (days after stratification) Arabidopsis plants by supplementing their hydroponic growth medium with 150 mM NaCl (Fig 2-1). This concentration of NaCl has been used in several previous gene expression studies, since it induces a moderate stress response and is not acutely lethal (He *et al.*, 2005; Ma *et al.*, 2006). Indeed, after application of 150 mM NaCl, we observed visible signs of stress including loss of turgor. However, even after 48h of exposure to media supplemented with 150 mM NaCl, nearly all of the treated plants recovered and resumed growth when transferred into NaCl-free hydroponic medium (data not shown). Obviously, these experimental conditions differ from those experienced by soil-grown plants, especially field-grown crops, where multiple stresses and nutrient limitations can occur simultaneously (Mittler, 2006). Nevertheless, we expect that many stress response pathways will be conserved across treatments. This assumption, along with the precise experimental control afforded by laboratory hydroponics, justifies the use of hydroponics as model system for salinity stress.

To further define the physiological state of the plants that we subjected to

transcriptional profiling, we monitored the accumulation of two stress-inducible metabolites, namely anthocyanin and proline (Yoshiba *et al.*, 1997). Recent evidence suggests that anthocyanins may mitigate photooxidative injury by efficiently scavenging free radicals and reactive oxygen species (Gould, 2004). Anthocyanin concentrations increased rapidly in leaves of stressed plants, with the absorbance at diagnostic wavelengths increasing nearly three-fold in the first 6h after treatment (Fig 2-1). These pigments continued to accumulate, with a five-fold increase observed at 24 h after treatment. Interestingly, the concentration of anthocyanin began to decrease after 24 h of stress, perhaps due to catabolism of the accumulated pigment and acclimation to the stress conditions, resulting in levels of anthocyanin at 48h post-stress




Fig 2-1. Accumulation of anthocyanin and proline in Arabidopsis plants in responses to NaCl treatment. a) Anthocyanin accumulation in Arabidopsis leaves. Anthocyanin concentration is reported as an absorbance ratio: $(A_{530-1/4} \cdot A_{657})/g$ fresh weight. b) Proline accumulation in roots and leaves of Arabidopsis plants. Arabidopsis plants were mock treated or exposed to 150mM NaCl for 6, 24 and 48 h. The means of six experiments \pm S.D. are shown.

that were slightly lower than levels observed at 6h. Proline concentration increased rapidly in both the leaves and roots of NaCl-stressed plants until at least 48h following NaCl treatment (Fig 2-1). We observed a 3-fold increase in proline concentration in roots after 6h, and an 18-fold increase after 48h. A similar pattern of increase, with

higher absolute concentrations of proline, was also observed in shoots. The accumulation of these metabolites indicates that these plants were actively mounting a stress response at the time of transcriptional profiling.

General transcriptomic responses

To characterize the effect of NaCl treatment on gene expression in Arabidopsis roots, we extracted RNA from control and stressed root samples at 6, 24, and 48h following treatment, and analyzed these in paired, dye-reversed hybridizations on spotted 70-mer oligonucleotide microarrays consisting of probes representing 23,687 unique genes identified by Arabidopsis Genome Initiative (AGI) locus identifiers. We identified probes for 7,217 unique genes whose treated:untreated \log_2 expression ratio differed significantly from 0 at one or more time points, according to the Significance Analysis of Microarrays (SAM) algorithm, with a false discovery rate (FDR) of less than 5% (Tusher *et al.*, 2001). Using a 2-fold cutoff, the number of salt up-regulated genes is 1540, 1444 and 1642 for 6, 24 and 48 h time points, respectively, whereas the number




Fig. 2-2. Venn diagrams of salt-regulated genes. Percentages in parentheses were calculated with the total numbers of genes assigned with AGIs (23,687). Figures in rectangles indicate salt treatment hours (h) and total number of salt-regulated genes at each time point. Only genes showing ≥ 2 -fold between treatment and control with a $FDR \leq 5\%$ in SAM were included here. Genes/probes that passed filterings (signal intensity above background on more 4 slides and $FDR \leq 5\%$) but have no AGI assigned are not listed here.

of salt down-regulated genes is 1354, 1924 and 2124 for 6, 24 and 48 h time points, respectively. In total, 5280 unique genes with AGI assigned were identified to be salt-regulated (Fig. 2-2). Among these statistically significant expression ratios, probes representing 2,367 unique genes showed a NaCl-induced increase in transcript abundance of at least 2.0 fold at one or more time points. Conversely, probes representing 2,913 unique genes showed a decrease in abundance of at least 2.0 fold or more. Thus, transcript accumulation for at least 10% of Arabidopsis genes was strongly (i.e. >2.0 fold) induced, and transcripts for at least 12% of genes were repressed at one or more time points, in response to NaCl-treatment. The proportion of NaCl-responsive genes we observed is somewhat higher than previously reported estimates based on cDNA microarrays, likely reflecting the higher specificity of the oligonucleotide-based arrays (Seki *et al.*, 2002a).

Validation of microarray results

We compared our results to publicly available microarray data from the AtGenExpress project, which used Affymetrix ATH1 arrays to describe the transcriptome of hydroponically grown Arabidopsis roots exposed to a 150 mM NaCl-shock for up to 24h (<http://www.arabidopsis.org/info/expression/ATGenExpress.jsp>). Qualitative trends in expression ratios are highly conserved between these datasets. Quantitatively, we observed a Pearson correlation co-efficient of 0.77 (6h) and 0.69 (24h) when comparing the ATH1 expression values with our signal-intensity filtered ratios. The transcriptomic responses we describe here are therefore generally reproducible across platforms and between laboratories. Moreover, probes for at least 3,277 genes were deposited on our spotted microarray that are not present on the ATH1 array, of which 730 were significantly responsive to NaCl treatment. For example, the gene with the highest-fold induction (mean ratio=7.4 (\log_2)) in our analysis is transcription factor *bHLH092* (At5g43650), but this gene is not represented on the ATH1 array. Thus, although the Affymetrix ATH1 array and the Operon 1.0 70-mer probe set are each commonly described as “whole-genome”, it is clear that either platform alone is deficient in the detection of hundreds of potentially significant genes, although in combination, they provide a nearly complete representation of the predicted transcriptome.

To provide further validation of our microarray data, we performed quantitative RT-PCR analysis on specific transcripts (Table 2-1). We selected 15 genes representing different functional categories, which according to our microarray analysis increased, decreased, or remained essentially unchanged in terms of transcript abundance following salt-shock. For all of these genes, the expression ratios measured by qRT-PCR and by microarray were highly correlated ($r= 0.91, 0.92$ and 0.88 for 6, 24 and 48h treatment, respectively). These results help to confirm the general accuracy of the microarray data we present here. The functional significance of the NaCl-responsive genes that were validated by qRT-PCR is discussed in more detail below.

Table 2-1 Comparison of qRT-PCR and microarray results for selected genes.

AGI#	annotation	microarray (log ₂ ratio)			qRT-PCR (log ₂ ratio)		
		6h	24h	48h	6h	24h	48h
At1g44830	AP2/ERF	4.37	5.42	4.89	6.05	6.37	6.47
At5g43650	bHLH92	7.21	7.58	7.45	11.05	11.93	11.87
At4g31500	CYP83B1/SUR2	-2	-1.89	-1.43	-1.84	-1.6	-1.22
At4g26410	expressed protein	0.01	0	0.04	-0.3	-0.23	-0.27
At2g04070	MATE	3.77	5.47	5.35	3.1	4.92	5.27
At3g23250	MYB15	4.05	3.25	1.31	5.39	5.67	5
At4g01550	NAC	1.93	3.35	3.21	2.46	4.87	5.65
At2g37130	PER21	-1.73	-1.56	-0.98	-1.32	-1.4	-0.69
At3g01190	PER27	-0.37	-0.02	0.14	-2.96	-1.21	-0.18
At5g64100	PER69	-1.78	-2.69	-1.8	-1.85	-2.4	-2.2
At2g24570	WRKY17	3.74	1.94	1.16	3.84	2.15	1.64
At2g30250	WRKY25	2.59	4.04	4.49	3.49	6	6.23
At2g38470	WRKY33	4.02	4.42	3.58	4.81	5.58	4.64
At4g30280	XTH18	2.54	1.3	1.57	2.65	1.58	2.5
At3g17860	ZIM	2.28	0.65	-0.84	2.42	1.02	-0.97

Genes representing a variety of expression patterns and functional categories were selected from the microarray dataset for further validation using qRT-PCR. Values shown are the mean of at least four independent measurements (microarray data), or three independent measurements (qRT-PCR).

Classification of NaCl-responsive transcripts

We used the STEM to summarize our filtered microarray data, by clustering it into 16 distinct temporal expression patterns (Fig 2-3). The algorithms implemented in STEM are designed specifically for microarray experiments that sample only a few time points, such as ours. STEM accordingly minimizes the potential for over-fitting that can occur when some other clustering methods are applied to time course data, and also

facilitates the identification of over-represented functional categories within each cluster, as described below.

The predominant temporal expression patterns detected by STEM analysis show that changes in transcript abundance can be detected within the first six hours following treatment for the majority of the 5,463 NaCl-responsive genes represented in the profiles (Fig 2-3.). At subsequent time points (i.e. 24h, 48h) the response generally either continued along the same trajectory (Fig 2-3 a,i), or remained close to the level of induction or repression observed at 6h (Fig 2-3 b,j). Another subgroup of genes had a peak response at 6h, which returned towards untreated levels of transcript abundance either gradually (Fig 2-3 c,k), or more rapidly (Fig 2-3 d,e,l,m). It is notable that a significant number of these genes show opposite patterns of induction and repression at 6h as compared to 48h (Fig 2-3 d,l). Still other genes that were responsive at 6h showed a peak change in expression at 24h (Fig 2-3 f, n), or, surprisingly, a moderate dampening of the response at 24h, and a renewed intensity of response at 48h (Fig 2-3 g,o). Finally, in contrast to all of the patterns discussed so far, which began with a marked response at 6h, a limited but significant number of genes showed relatively little response until the 48h time point (Fig 2-3 h,p). Thus, although the majority of genes are responsive by 6h of treatment, dynamic patterns of gene expression are also detected after 24h and 48h of treatment. These observations indicate that different regulatory networks and effectors are active at each of these different periods following the perception of stress.

STEM was used to identify categories of functionally and/or structurally related genes that are enriched (i.e. statistically over-represented at $FDR < 5\%$) in one or more of the clusters described above. Preliminary analysis using Gene Ontology consortium categories (data not shown) was used to guide the selection of a detailed list of gene families and superfamilies from primary literature sources, as well as from curated databases including KEGG (Kyoto Encyclopedia of Genes and Genomes), AraCyc, and TAIR (The Arabidopsis Information Resource) (Kanehisa and Goto, 2000; Rhee *et al.*, 2003; Zhang *et al.*, 2005). Most broadly, the functional themes represented by NaCl-responsive transcripts can be divided into effectors and regulators. According to this analysis, the most prominent effectors of the NaCl stress response are genes involved in




Fig. 2-3. NaCl-responsive transcripts grouped according to temporal expression profiles. Microarray expression data that had been previously filtered for significance based on signal intensity and SAM statistical analysis, was divided into 16 distinct temporal profiles, using STEM software. Each of the profiles (labelled a-o) is represented as a different plot, with mean expression ratios (\log_2 ; treated:control) for each of the assigned transcripts at each time point plotted in grey. The median of all assigned expression ratios in each profile is plotted in black. To emphasize that the y-axis range is different between plots, a dashed line corresponding to a \log_2 expression ratio of 0 is also shown in each plot.

detoxification, transport, protein turnover, energy metabolism, and production of osmoprotectants. The major classes of regulators detected are signal transduction

components, transcription factors, and hormone related genes. Each of these broad themes has been previously reported in association with abiotic stress responses (Sahi *et al.*, 2006). The analysis we present here highlights specific NaCl-responsive genes and groups of genes within these categories. Tables 2-2 to 2-6 show groups of functionally and/or phylogenetically related genes that, according to STEM analysis, were significantly over represented (FDR<5%) in one or more STEM profile (Fig 2-3), as well as a limited number of other groups of genes mentioned in the text, for comparison. The tables also include the number of genes within each category that were up- or down-regulated more than 2-fold, although this 2-fold threshold is not a criterion *per se* in STEM analysis.

Osmoprotectants

Osmoprotectants including proline and trehalose, and small hydrophilic proteins such as dehydrins help to maintain hydration of cellular components during osmotic stress (Alsheikh *et al.*, 2005). Genes regulating levels of osmoprotectants are highly stress responsive and were among the first stress-inducible transcripts reported in the literature (Gilmour *et al.*, 1992; Yamaguchi-shinozaki *et al.*, 1992; Verbruggen *et al.*, 1993; Welin *et al.*, 1994). As expected, we observed induction of many of these genes in our NaCl-treated tissues (Table 2-2). Two proline biosynthetic genes were induced (*P5CR*, At5g14800; *P5CS*, At2g39800), consistent with the observed increase in proline abundance (Fig 2-1). We also observed transcript-level induction of a large proportion of genes in the trehalose biosynthetic pathway including 7 of the 8 trehalose phosphatases detectable on our array, as well as some (but not all) dehydrins (Table 2-2). Although trehalose has been shown to act as an osmoprotectant in bacteria and fungi, recent studies have suggested that this disaccharide acts as a signalling molecule in plants, noting that its intracellular concentration is too low to be an effective osmoprotectant (Avonce *et al.*, 2004).

Reactive oxygen network

NaCl treatment, like many environmental stresses, generates reactive oxygen species (ROS) (Mittler *et al.*, 2004). These ROS and the products of their reactions with other cellular components (e.g. peroxidized lipids) are generally toxic. It is therefore not surprising that plants have evolved large gene families for detoxification of ROS and

Table 2-2. NaCl-responsive transcripts related to osmoprotection, detoxification, and transport.

family, group, or pathway	in genome	on array	signal above bkgnd.	number of genes		assignment to STEM profiles (Fig. 2-3) cluster:# of genes
				>2 fold up	>2 fold down	
osmoprotectants						
proline	5	5	5	2	1	a:1;b:1;j:1
trehalose biosynthesis*	22	21	15	8	1	a:4;b:4;c:1
trehalose phosphatase*	11	10	8	7	0	a:3;b:4
dehydrin*	10	10	6	4	1	b:1;c:2;h:1;i:1
ROS-response network						
alternative oxidase*	6	6	4	2	0	b:2
ascorbate peroxidase	9	8	6	1	1	b:1;h:1;o:1
blue copper protein	9	8	5	1	1	b:1;i:1;n:1
dehydroascorbate reductase	5	2	1	0	1	i:1
reductase						
ferritin*	4	4	3	2	0	a:2
glutaredoxin*	27	26	8	3	3	a:3;i:1;j:2
glutathione peroxidase	8	8	4	0	1	j:2
glutathione reductase*	5	4	4	0	0	m:1
monodehydroascorbate reductase	5	5	4	2	0	a:1;c:1
NADPH oxidase*	10	10	4	1	2	c:1;i:1;k:1
NADPH oxidase-like	9	8	4	0	3	i:2;j:1
peroxiredoxin	11	10	5	0	2	j:2
superoxide dismutase	8	8	3	0	3	i:1;j:2
thioredoxins	32	31	20	5	7	a:1;b:2;c:1;f:1;i:2;j:3;l:1;o:1
class III peroxidase*	73	71	45	10	27	a:6;b:3;c:1;d:1;i:3;j:17;k:4;l:1;o:3
glutathione-S-transferase*	53	49	37	19	7	a:4;b:9;c:1;d:3;e:1;f:1;h:2;i:3;j:2;l:2;n:1;o:1
GST-tau family*	28	26	20	13	3	a:1;b:9;c:1;d:3;i:1;j:1;n:1
transporters						
aquaporin*	35	33	24	1	18	a:1;i:4;j:12;n:1
V-Type ATPase*	22	18	14	0	4	j:4
antiporter	70	66	25	5	9	a:4;c:1;i:2;j:4;k:1;l:2
Na ⁺ /H ⁺ exchanger*	8	8	6	4	0	a:4
sugar transporter*	48	47	23	11	4	a:6;b:3;d:1;f:1;i:1;j:2;o:1
ABC transporter*	94	88	29	10	5	a:6;b:2;d:3;f:1;g:1;i:1;k:1;l:2
LeOPT1*	51	48	22	10	8	a:4;b:6;i:3;j:3;o:2
MATE*	56	51	26	13	1	a:4;b:6;c:1;d:1;f:1;i:1

Groups of genes that are significantly enriched in at least one STEM profile at a FDR of <5% are marked with an asterisk (*). The final column details the frequency of assignment of individual genes to specific STEM profiles identified by corresponding letters in Fig 2-3.

Functional categories in this table are defined according to the following sources: proline and trehalose biosynthesis (Zhang *et al.*, 2005); dehydrins (Alsheikh *et al.*, 2005); ROS-network (Mittler *et al.*, 2004); class III peroxidases (Tognolli *et al.*, 2002; Rhee *et al.*, 2003), GSTs (Rhee *et al.*, 2003); MATE transporters (Rogers and Guerinot, 2002); aquaporins (Rhee *et al.*, 2003; Boursiac *et al.*, 2005); all other transporters (Ward, 2001; Rhee *et al.*, 2003).

their by-products. At least 148 enzymes have been defined as part of the ROS scavenging network of Arabidopsis (Mittler *et al.*, 2004). However, of the 75 ROS network genes with sufficient hybridization signal intensity to be detected on our microarray, a majority of the responsive transcripts were down-regulated >2-fold by NaCl (Table 2-2). The predominance of non-responsive or down-regulated transcripts was evident within almost every category of ROS-scavenging enzymes, except for alternative oxidase, ferritin, and monodehydroascorbate reductase, in which all NaCl-responsive transcripts were induced. The limited proportion of genes with NaCl-inducible transcripts within the ROS-scavenging network points to a considerable amount of sub-functionalization at the regulatory and catalytic levels, as well as roles for these genes beyond detoxification of ROS (Noctor, 2006). A third family of peroxidases, namely the type III (heme-containing) peroxidases, is not usually considered part of the ROS scavenging network and may have diverse roles including generating ROS for cell wall modification (Passardi *et al.*, 2005). We arbitrarily selected three peroxidase genes PER21 (At2g37130), PER69 (At5g64100), and PER27 (At3g01190) for validation by qRT-PCR. The results confirmed that these genes were generally down-regulated by NaCl-treatment (Table 2-1). These observations are consistent with functions for type III peroxidases in biological processes that are largely distinct from ROS scavenging.

A large proportion of glutathione-S-transferases (GSTs) were induced by NaCl treatment (Table 2-2). We note that nearly all of the NaCl inducible GSTs are from the Tau family of GSTs, highlighting a distinct role for this sub-class of genes in oxidative stress responses, as has been previously inferred from proteomic studies (Kilili *et al.*, 2004).

Transporters

The regulated transport of molecules across the plasma and vacuolar membranes is a well characterized response to abiotic stress (Sahi *et al.*, 2006). Within the NaCl-responsive transcripts on our microarray, we observed enrichment of transporters for water, sugars, cations, and other molecules (Table 2-2). The NaCl-responsive aquaporins and vacuolar-type ATPases detected on our microarray were enriched almost exclusively in down-regulated transcripts. In contrast, almost all of the detectable NaCl-responsive Na^+/H^+ exchangers were induced by stress. The homogeneity of these responses indicates relatively little sub-functionalization within these groups of genes, as compared to the ROS scavenging network described above. Among the remaining antiporters, as well as sugar transporters, ABC (ATP-binding cassette), and LeOPT1-like transporters and MATE-like (multi-antimicrobial extrusion) efflux carriers, specific transcripts were induced and others were repressed by NaCl-treatment (Table 2-2) (Ward, 2001; Rogers and Guerinot, 2002). Within the MATE family in particular, the proportion of NaCl-inducible transcripts is relatively large, as transcripts representing approximately half of the 26 detectable genes were induced >2 fold by NaCl, while only one MATE gene was equivalently repressed. One of these transcripts was among those we selected for further validation by qRT-PCR (Table 2-1). Both qRT-PCR and microarray analysis showed an approximately 8-fold increase in transcript abundance after 6h of NaCl treatment, with further accumulation to a 32-fold increase by 48h. The large proportion of induced genes within the MATE family, plus the magnitude of their induction, suggests a previously underappreciated role for MATE-efflux carriers in the root response to NaCl. Very little is known about this family of carriers, although a previous meta-analysis of microarray data identified some MATE transporters among a list of stress-induced transporters (Maathuis, 2006).

Primary energy metabolism, carbohydrates and cell walls

Transcripts required for the main respiratory pathways of glycolysis, the tricarboxylic (TCA) cycle, and the pentose phosphate pathway (PPP) were generally non-responsive or were down-regulated by NaCl treatment (Table 2-3). Genes encoding components of the mitochondrial electron transport chain were especially enriched among down-regulated profiles (Table 2-3). The down-regulation of these energy-evolving

Table 2-3. NaCl-responsive transcripts related to respiration and carbohydrate and cell wall metabolism.

family, group, or pathway	in genome	on array	number of genes		assignment to STEM profiles (Fig. 2-3) cluster:# of genes	
			signal above bkgnd.	>2 fold up		>2 fold down
glycolysis*	70	64	45	6	10	a:2;b:1;d:1;f:1;h:2; i:3;j:5;k:3
tricarboxylic acid cycle	56	52	34	6	11	a:4;b:1;f:1;g:1; i:4;j:7
mitochondrial respiration*	95	66	55	0	26	g:1;i:2;j:23;o:1
non-oxidative pentose phosphate pathway	13	12	8	1	6	h:1;i:1;j:3;k:1;o:1
oxidative * pentose phosphate pathway	19	19	13	0	5	i:3;j:1;k:1
glycosyltransferase*	279	269	113	29	26	a:12;b:8;c:2;d:4;f:3; g:2;h:1;i:10;j:10;k:1; l:1;n:1;o:3
glycoside hydrolase (GH)*	306	289	121	42	27	a:17;b:16;c:1;d:4;e:3; h:2;i:11;j:6;k:2;l:2;o:3 p:1
GH family 16*	27	26	14	10	2	a:4;b:4;d:1;e:1;j:1; o:1
GH family 28*	52	46	10	4	1	a:3;b:1;d:1
expansins	35	32	21	7	4	a:1;b:3;c:1;d:2;i:3; m:1
lignification (exclusive of 4-CL like and COMT-like)	39	36	20	3	7	a:1;c:1;d:1;i:1;j:5; o:1
4CL-like*	9	8	3	2	1	b:1;d:1;k:1
COMT-like*	13	12	5	5	0	a:1;b:3
lipid transfer protein*	70	64	31	5	15	a:3;b:1;h:1;i:1;j:8; k:2;n:2;o:2

Groups of genes that are significantly enriched in at least one STEM profile at a FDR of <5% are marked with an asterisk (*). The final column details the frequency of assignment of individual genes to specific STEM profiles identified by corresponding letters in Fig 2-3. Functional categories in this table are defined according to the following sources: glycolysis, TCA, electron transport and PPP (Zhang *et al.*, 2005); all other groups: (Rhee *et al.*, 2003).

pathways is a common stress response that may serve to conserve energy and limit growth and further generation of ROS (Moller, 2001). The few glycolytic and TCA-related transcripts that were up-regulated were all isoforms of other genes that had been

down-regulated, although no correlation between cytoplasmic location and NaCl-response could be detected (data not shown). Transcript abundance for almost all components of the PPP was also decreased (Table 2-3), including D-ribulose-5-phosphate 3-epimerase (At3g01850, At1g63290), ribose 5-phosphate isomerase (At2g01290, At3g04790), 6-phosphogluconolactonase (At5g24410, At3g02360), and glucose-6-phosphate-1-dehydrogenase (G6PDH). The PPP is an important source of NADPH, especially in non-photosynthetic tissues, and therefore might be expected to have increased activity under oxidative stress (Rizhsky *et al.*, 2002; Chen *et al.*, 2004; Saher *et al.*, 2005). However, we noted that the detected down-regulated G6PDH transcripts were G6PD5 (At3g27300) and G6PD6 (At5g40760), which are the major cytosolic versions of G6PDH expressed in Arabidopsis roots and are distinct from the four genes encoding plastid-localized isozymes (Wakao and Benning, 2005). Previous analyses of G6PDH enzyme activity in potato and tobacco leaves has shown that isozymes located in plastids, but not the cytosol, have increased enzymatic activity following exposure to osmotic and oxidative stress (Hauschild and von Schaewen, 2003; Debnam *et al.*, 2004). Thus, G6PD5 and G6PD6 may function preferentially under conditions other than osmotic and oxidative stress. The sub-functionalization of isozymes into groups with contrasting responses to stress appears to be a common theme within the PPP, TCA, and glycolysis-related genes detected on our array.

Glycosyltransferases (GTs) and glycoside hydrolases (GHs) are two large superfamilies of carbohydrate- active enzymes (Coutinho and Henrissat, 1999). All GTs catalyze similar biochemical reactions, namely the transfer of sugar moieties to acceptor molecules. Conversely, GHs break bonds existing between sugar moieties and other molecules. Within each of these superfamilies, a significant and roughly equivalent number of genes were either up-regulated or down-regulated by NaCl at the transcript level, suggesting dynamic regulation of glycosylation in response to stress. These salinity responsive genes were distributed throughout the many constituent gene families of GTs and GHs, such that only two of the constituent families of GHs were significantly enriched in NaCl-responsive genes by our criteria: GH16, and GH28 (Table 2-3.). One member of the GH16 family, *xyloglucan endotransglucosylase /hydrolase18* (XTH18; At4g30280), was previously reported to be expressed in the

base of elongating roots, and is among the NaCl-induced genes we selected for validation by qRT-PCR (Table 2-1) (Vissenberg *et al.*, 2005). GH16 and GH28 are comprised largely of xyloglucanases and galacturonases, respectively, which are both often associated with cell wall metabolism. A limited number of other cell-wall related gene families were also enriched in NaCl-responsive transcripts, most notably the expansins (Table 2-3) (Lee *et al.*, 2001), for which at least one-third of the genes were transcriptionally up-regulated by NaCl. Increased activity of expansins and xyloglucanases is consistent with the increase in cell wall flexibility observed in response to osmotic stress in some species (Jones and McQueen-Mason, 2004).

A decrease in transcript abundance for many enzymes involved in lignification, as well as the potentially cell wall-related lipid transfer proteins (LTPs), were also observed from the microarray data (Table 2-3). However, we note that two families of enzymes that are sometimes grouped in the “lignification tool box”, namely the 4CL (4-coumarate:CoA ligase)-like and the COMT (Caffeic acid O-methyltransferase)-like families (Raes *et al.*, 2003), were almost entirely up-regulated in response to NaCl stress. It should be emphasized that these two families do not include their namesakes *4CL* and *COMT*, which are enzymes with clearly established roles in lignification. Thus the 4CL-like and COMT-like genes may have unique roles in the NaCl stress-response, and may not necessarily be related to lignification. LTPs are another group of proteins, some of which have been shown to affect cell wall extensibility or to be secreted in response to NaCl stress (Dani *et al.*, 2005; Nieuwland *et al.*, 2005). Fifteen LTP transcripts on our microarray were down-regulated by NaCl treatment (Table 2-3), although 5 LTPs were up-regulated. It is likely that these various LTPs represent a diversity of stress responses, some of which may also be related to cell-wall extensibility.

Protein metabolism

The impact of abiotic stress on protein metabolism is evident in our microarray results. A decrease in bulk protein translation following NaCl has been detected in proteomic studies, and is consistent with our observed down-regulation of the majority of transcripts for almost all cytosolic and plastidic ribosomal proteins (Table 2-4) (Ndimba *et al.*, 2005). Among proteins that promote the proper folding of proteins

and/or prevent the aggregation of nascent or damaged proteins, we detected enrichment of NaCl-responsive genes within several gene families. Within the peptidyl prolyl isomerase super-family, the 29 genes represented by detectable transcripts were largely down-regulated by NaCl (Table 2-4) (He *et al.*, 2004), although only the cyclophilins were significantly (<5% FDR) enriched in down-regulated transcripts. A second superfamily of genes related to protein folding and aggregation are the heat shock proteins (HSPs) (Lin *et al.*, 2001; Larkindale *et al.*, 2005). As expected, a majority of the 51 detectable HSP superfamily genes were transcriptionally induced by NaCl, and transcripts for only 5 genes were down-regulated by >2 fold (Table 2-4). Although the majority of HSPs are resident in the cytosol, most of the 5 down-regulated genes detected in the HSP superfamily were targeted to specific compartments, including two genes targeted to the endoplasmic reticulum (ER) (HSP70, At4g16660; HSP90-like *SHEPERD*, At4g24190), a chloroplast targeted HSP100/CLP (At5g50780). Consistent with these observations, some HSP family proteins that are targeted to chloroplasts or the ER have been previously shown to be repressed by particular abiotic stresses (Sung *et al.*, 2001).

Proteolytic activity has been attributed to over 600 peptidases within the plant genome, (Rawlings *et al.*, 2004). From our microarray data, both up- and down-regulated transcripts were detected for genes within each of the aspartic, cysteine, serine, and metallo-peptidase super-families (Table 2-4). Specific families and sub-families within the cysteine and serine peptidases were significantly enriched in NaCl responsive genes, as shown in Table 2-4. In contrast to the other peptidase superfamilies, the responsive threonine peptidases that comprise the 20S catalytic core of the 26S proteasome were exclusively down-regulated. A predominance of down-regulation was also observed among the transcripts comprising the other component of the 26S proteasome, namely the 19S regulatory particle (Table 2-4). However, many of the gene families putatively encoding functions upstream of the 26S proteasome in the ubiquitination pathway were up-regulated. Notably, SKP1 (S-phase-associated kinase protein) kinases, E3 U-box proteins, and E3 RING (Really Interesting New Gene) finger proteins were all significantly enriched in NaCl-induced genes (Table 2-4). Because the 26S proteasome works in conjunction with the ubiquitination pathway as

Table 2-4. NaCl-responsive transcripts related to protein metabolism.

family, group, or pathway	in genome	on array	signal above bkgnd.	number of genes		assignment to STEM profiles (Fig. 2-3) cluster:# of genes
				>2 fold up	>2 fold down	
ribosome large subunit*	121	100	83	0	59	i:19;j:36;n:4
ribosome small subunit*	94	87	74	0	53	i:24;j:24;n:1;o:1
ribosome plastidic	13	13	9	0	3	i:1;j:1;n:1
peptidylprolyl isomerase (PPI)						
PPI: cyclophilin*	28	28	19	1	9	b:1;d:1;i:1;j:8;l:1; m:1
PPI: FKB	23	22	7	1	2	a:1;j:2
PPI: parvulin	3	3	3	0	2	i:1;j:1
heat shock proteins*	70	68	51	36	5	a:31;b:3;c:1;f:1;i:2; j:2;o:1
peptidase*	557	524	265	43	73	a:20;b:12;c:1;d:4;f:6; g:4;h:1;i:16;j:46;k:3; l:2;n:2;o:6
aspartic peptidase	76	69	31	7	8	a:4;b:2;f:1;i:2;j:5;o:1
cysteine peptidase	102	102	55	11	15	a:5;b:2;d:3;f:3;i:4; j:7;k:2;l:1;n:1;o:1
family C26*	10	10	9	4	2	a:1;b:1;d:3;i:1;j:1
metallopeptidase	85	78	44	4	6	a:2;f:2;i:1;j:4
serine peptidase*	261	244	112	18	31	a:8;b:6;c:1;d:1;g:4; h:1;i:8;j:18;k:1;l:1; n:1;o:4
sub-family S10-004*	19	17	7	0	6	i:1;j:5
sub-family S10-005*	24	24	9	0	3	g:1;j:1;o:2
sub-family S33-UPW*	45	42	22	6	7	a:2;b:2;c:1;d:1;g:2; i:1;j:3;l:1;o:1
threonine peptidase *	28	28	22	3	13	a:1;b:2;i:1;j:12
20S proteasome subunit*	19	19	17	0	12	i:1;j:11
19S proteasome*	33	31	24	0	10	i:3;j:6
SKP1/ASK1*	17	16	9	5	3	a:2;b:1;h:2;j:3
E3 RING-type*	430	414	178	47	34	a:16;b:17;c:4;d:8;f:4; h:3;i:14;j:14;k:1;l:3;n:1 p:2
E3 Ubox-type*	61	53	30	16	3	a:6;b:8;c:1;d:1;j:2; o:1
Ubox class III*	12	11	9	7	2	b:5;c:1;d:1;j:1;o:1

Groups of genes that are significantly enriched in at least one STEM profile at a FDR of <5% are marked with an asterisk (*). The final column details the frequency of assignment of individual genes to specific STEM profiles identified by corresponding letters in Fig 2-3.

Functional categories in this table are defined according to the following sources: ribosomes (Barakat *et al.*, 2001) ; PPIs (He *et al.*, 2004); peptidases (Rawlings *et al.*, 2004); 19S

proteasome (Kanehisa and Goto, 2000); E3 RING (Stone *et al.*, 2005); HSPs, SKP1s, E3 Ubox (Rhee *et al.*, 2003).

the major proteolytic system of the nucleus and cytosol, it appears paradoxical that many ubiquitination-related proteins would be up-regulated, while the detectable 26S proteasome transcripts would be uniformly down-regulated by NaCl treatment (Smalle and Vierstra, 2004). This suggests a role for some SKP1, E3 U-box proteins, and E3 RING related enzymes that is independent of the 26S proteasome in the stress response, and may also hint at aspects of regulatory complexity not captured by microarray analysis (Kawaguchi *et al.*, 2004).

Signal transduction components and hormones

The predicted kinases of Arabidopsis form a very large superfamily of at least 979 genes that has been subdivided into 5 classes comprised in total of 81 families (Tchieu *et al.*, 2003; Shiu *et al.*, 2004). Nearly all of the NaCl-responsive kinases detected from our microarray were associated with the two largest classes in the PlantsP classification system (PPC:1, PPC:4) and twelve families therein (Table 2-5). These include eight families of receptor-like kinases (RLKs) and related kinases: legume lectin domain (LLD), leucine rich receptor (LRR) II & X, and LRR XI & XII, receptor like cytoplasmic kinase (RLCK) VII, wall-associated kinase (WAK)-like, proline-rich extensin-like kinases (PERK), S-Domain (type 1), and domain of unknown function (DUF) 26. Within each of these families, nearly all NaCl-responsive genes increased in transcript abundance following stress-treatment, and none of the detectable PERKs or S-Domain (type 1) kinases was repressed by NaCl (Table 2-5). Aside from the RLKs and related proteins in class PPC:1, we also detected enrichment of NaCl-responsive transcripts within several families of non-transmembrane kinases, including Mitogen Activated Protein Kinase Kinase Kinases (MAP3Ks), Calcium Responsive Kinases (CRKs), Ribosomal Protein S6 Kinases (RPS6K), and an unnamed family (PPC:4.4.1) of kinases. Among these, up-regulated transcripts were especially predominant in the RPS6K and PPC:4.4.1 families.

Protein phosphatases are divided into three major classes: Protein Serine /Threonine Phosphatases (STKs), Dual Specific Phosphatases (DSPs), and Protein

Table 2-5. NaCl-responsive transcripts related to signal transduction and hormone biosynthesis.

family, group, or pathway	in genome	on array	signal above bkgnd.	number of genes		assignment to STEM profiles (Fig. 2) cluster:# of genes
				>2 fold up	>2 fold down	
kinases	979	921	426	164	66	n,o:1;a:61;b:69;c:11;d:7;e:1;f:10;g:2;h:4;i:16;j:30;k:3;l:1;n:1;o:12;p:1
PPC:1 *	564	530	238	105	38	a:36;b:49;c:6;d:5;e:1;f:3;g:2;h:3;i:6;j:17;k:1;l:1;n:1;o:10
1.11.1 LLDK*	41	40	22	13	3	a:8;b:4;g:1;j:1;o:2
1.12.2 LRR II & X*	18	17	10	4	0	a:1;b:3
1.12.4 LRR XI & XII	45	44	27	8	5	a:2;b:2;c:1;d:2;f:1;h:1;i:1;j:3;n:1
1.2.2 RLCK VII*	45	44	25	19	0	a:6;b:9;c:1;d:2;e:1;f:1
1.5.1 WAK	27	23	11	6	5	b:3;c:1;j:3;l:1;o:1
1.6.2 PERK*	18	17	6	4	0	a:1;b:3
1.7.1 S-Domain (type 1)	14	14	10	7	0	a:3;b:3;c:1
1.7.2 DUF 26*	66	61	29	17	5	a:5;b:11;f:1;i:1;j:2;o:1
PPC:2	52	51	22	6	6	a:3;b:2;f:1;i:2;j:2;k:1;o:1
PPC:4 *	279	265	133	47	18	n,o:1;a:20;b:18;c:5;d:1;f:5;i:6;j:10;o:1
4.1 MAP3K*	51	48	20	10	2	a:6;b:3;d:1;i:1
4.2 CRK*	123	119	58	21	10	n,o:1;a:5;b:10;c:3;f:3;i:3;j:4;o:1
4.2.6 RPS6K *	40	40	16	7	1	n,o:1;b:6;c:1;f:1
4.4 unclassified*	27	23	8	7	1	a:1;b:4;c:1;f:1;j:1
phosphatase*	125	116	64	24	9	a:9;b:11;c:3;d:1;g:1;i:5;j:1;k:1
PPC:6.1 (STKs)	23	19	12	0	3	i:2
PPC:6.2 DSP	8	8	6	1	2	a:1;i:1;j:1
PPC:6.3 PP2C *	64	62	38	22	3	a:7;b:11;c:3;g:1;i:1;k:1
response regulator: A *	11	11	8	4	1	a:1;b:1;c:2;i:1
similar to MLO proteins*	14	14	5	3	2	b:3;i:1;j:1
14-3-3 proteins*	13	13	8	0	5	i:1;j:4
ethylene biosynthesis	28	27	16	6	5	a:2;b:3;f:1;i:2;j:2
jasmonic acid biosynthesis*	20	19	11	7	3	a:1;b:5;d:1;j:2;o:1
IAA biosynthesis*	10	9	7	2	2	a:1;f:1;h:1;i:1;n:1
auxin transport*	8	8	4	3	1	a:2;b:1;i:1
Lateral Organ Boundaries class I*	36	34	11	7	0	a:2;b:1;c:3;g:1;p:1

Groups of genes that are significantly enriched in at least one STEM profile at a FDR of <5% are marked with an asterisk (*). The final column details the frequency of assignment of individual genes to specific STEM profiles identified by corresponding letters in Fig 2-3.

Functional categories in this table are defined according to the following sources: kinases and phosphatases (Tchieu *et al.*, 2003); hormone biosynthesis (Zhang *et al.*, 2005); all others: (Ward, 2001; Rhee *et al.*, 2003).

Phosphatase 2C (PP2C) (Tchieu *et al.*, 2003). We observed that the STKs and DSPs differed greatly from the PP2Cs in their transcriptomic response to NaCl-shock (Table 2-5). Transcripts for only one of the 18 detected STKs or DSPs were strongly induced by NaCl treatment, whereas transcripts for more than half of the detectable PP2Cs were strongly induced by NaCl. Some PP2Cs have been previously identified as components of the ABA signalling pathway, and all of these are among the NaCl-induced PP2Cs detected from our arrays: *ABI1* (*ABA INSENSITIVE 1*; At4g26080), *ABI2* (At5g57050), *HABI* (*HOMOLOGY TO ABI1/ABI2*; (At1g72770), *HAB2* (At1g17550), and *AHG3* (*ABA-HYPERSENSITIVE GERMINATION3*, At3g11410) (Leung *et al.*, 1997; Merlot *et al.*, 2001; Saez *et al.*, 2004; Yoshida *et al.*, 2006). Our data indicate that nearly twenty other PP2Cs, including those outside of the ABI/HAB clade defined by Saez and colleagues, may play important roles in NaCl stress signalling.

The two-component signal transduction system, which consists of histidine protein kinases (HKs), His-containing phosphotransfer proteins (HPs) and response regulators (RRs), is involved in plant hormone, stress, and light signaling (Hwang *et al.*, 2002). The Arabidopsis genome contains 54 genes encoding putative HKs, HPs, RRs, and related proteins. In our data, none of the 16 HKs were induced by salt, while *AHK1*, a putative osmosensor (Urao *et al.*, 1999), was down-regulated at the 6h time-point. However, the A-type RRs were significantly enriched among clusters of NaCl-induced genes (Table 2-5). These include *ARR5*, *ARR6*, and *ARR7*, which act as negative regulators to repress the expression of the cytokinin-responsive genes (Hwang and Sheen, 2001), suggesting a potential link between NaCl stress and cytokinin signalling.

A handful of other families of putative regulatory proteins were statistically enriched among clusters of NaCl-responsive genes in our analysis. Transcripts for at least half of the six MLO (mildew resistance locus o) proteins detected on our microarray were induced by NaCl-treatment (Devoto *et al.*, 2003). These are seven-transmembrane domain proteins with features similar to G-protein coupled receptors,

and some members of this gene family have been reported to be abiotic stress responsive (Chen *et al.*, 2006). Finally, transcripts decreased for at least 5 of the 8 14-3-3 genes detectable in our root microarray, while no significant increase in transcript abundance for any 14-3-3 protein genes was observed (DeLille *et al.*, 2001).

Ethylene and jasmonic acid (JA) are hormones whose activity has been previously correlated with environmental stress (Devoto and Turner, 2005; He *et al.*, 2005; Ludwig *et al.*, 2005). The rate-limiting step of ethylene biosynthesis is the production of 1-aminocyclopropane-1-carboxylic acid (ACC) by ACC synthase (ACS), which is followed by the conversion of ACC to ethylene by ACC oxidase (ACO) (Chen *et al.*, 2005). ACS and ACO are each encoded by gene families. Four ACC synthase and one ACC oxidase genes were significantly up-regulated by NaCl-treatment, while another 3 ACO genes were down-regulated by salt. Differential regulation of members of the ACO gene family indicates that although ACS is the key regulatory point for ethylene biosynthesis, regulation of ACO expression also has functional significance (Chen *et al.*, 2005). Ethylene signalling transduction involves ethylene response factor (ERF)/ ethylene response element binding protein (EREBP), which have been shown to act as activators or repressors of additional downstream ethylene-responsive genes. JA and its derivatives are also known to be involved in a wide range of developmental processes, as well as stress and defence responses (Devoto and Turner, 2005). At least 7 genes implicated in the biosynthesis of JA were induced by salt stress (Table 2-5), including *LIPOXYGENASE3* (*LOX3*, At1g17420), *ALLENE OXIDE CYCLASE* (*AOC1*, At3g25760), *AOC2* (At3g25780), and *ALLENE OXIDE SYNTHASE* (*AOS*, At5g42650), indicating that biosynthesis of this hormone is also NaCl responsive.

Within the biosynthetic pathway for indole acetic acid (IAA), which is an auxin, we detected up-regulation of two genes in response to NaCl treatment: namely nitrilases *NIT1* (At3g44310) and *NIT2* (At3g44300) (Table 2-5) (Woodward and Bartel, 2005). Consistent with an increase in levels of bioactive auxin, we also detected down-regulation of transcripts for *SUPERROOT2* (*SUR2*, At4g31500), an auxin conjugating enzyme (Barlier *et al.*, 2000). We confirmed the repression of *SUR2* transcripts by qRT-PCR (Table 2-1). Furthermore, we observed that of 4 putative auxin efflux carriers detectable after hybridization, transcripts for *PIN1* (At1g73590), *PIN3*

(At1g70940), and *PIN7* (At1g23080) were induced by our NaCl treatment. Part of the increased auxin activity we inferred from our microarray data may be directed towards increased lateral root formation, which is an auxin-regulated, and PIN requiring process that has been previously described in NaCl-treated roots (He *et al.*, 2005; De Smet *et al.*, 2006). Lateral root formation also involves genes from the *LATERAL ORGAN BOUNDARIES (LOB)* family; consistent with this requirement, we observed that the class I sub-group of this family was exclusively enriched in up-regulated transcripts, with at least seven genes induced following NaCl-treatment (Table 2-5)(Shuai *et al.*, 2002). Thus, the initiation of lateral root primordia is a NaCl-stress response that appears to be distinctly represented within our microarray data, even though the process involves only a small number of cells within the entire root.

Transcription Factors

The Arabidopsis genome encodes about 1,922 predicted transcription factors (TFs), which have been classified into 64 families (Riechmann *et al.*, 2000; Guo *et al.*, 2005; Iida *et al.*, 2005). From the over 1,613 transcription factors represented as probes on our microarray, transcripts for 734 genes were detected at a signal intensity above background. Of these, 289 predicted TFs were up-regulated and 139 transcription factors were down-regulated at least 2 fold at one or more time points following NaCl treatment (Table 2-6). STEM analysis identified 17 families that were significantly enriched among clusters of NaCl responsive genes. These TF families range in size from 7 to 203 members, and are described below. Although these 17 families contain almost two-thirds of the predicted TFs in the Arabidopsis genome, it is worth noting that transcripts for several other relatively large TF families were detected on the array, but these were largely unresponsive to NaCl stress. Some of the notable TF families that were expressed in roots, but which were not significantly enriched in NaCl-responsive transcripts are: MADS (MCM-1, Agamous, Deficiens and Serum Response Factor), ARF (Auxin Response Factor), Nin-like (nodule inception), and TUB (Tubby).

At least four TF families have been reported to be enriched in stress-responsive genes: MYB (Myeloblastosis), HSF (Heat Shock Factor), AP2/EREBP (Apetala-2/EREBP), and WRKY (named after the WRKY amino acid motif). A recent analysis of MYB-transcription factor gene expression concluded that almost all MYB TFs are

Table 2-6. Classification of NaCl-regulated transcription factors.

family, group, or pathway	in genome	on array	signal above bkgnd.	number of genes		assignment to STEM profiles (Fig. 2) cluster:# of genes
				>2 fold up	>2 fold down	
ALFIN*	7	7	4	0	3	i:3
AP2/EREBP*	146	138	76	50	6	a:8;b:31;c:6;d:3;e:1; f:2;h:1;i:1;j:1;k:1;l:1; o:2
ARF*	23	20	12	4	0	a:3;d:1;f:1
AS2*	42	39	15	7	3	a:2;b:1;c:3;g:1;i:2;j:1; p:1
AUX/IAA	29	28	15	7	1	a:4;b:3;d:1;f:1;i:1;j:1
B3	39	36	8	2	2	a:1;b:1;i:2
bHLH	162	148	62	15	14	a:8;b:4;c:3;d:1;h:1;i:7; j:4;k:2;o:1;p:1
bZIP	75	68	42	11	11	a:5;b:5;c:2;i:3;j:4;l:1; o:2;p:1
C2C2-co-like*	31	31	13	5	2	b:1;d:1;e:1;f:2;i:1;j:1
C2C2-DOF*	36	31	15	8	1	b:5;c:1;d:2;g:1;k:1
C2C2-GATA*	29	27	14	3	3	b:3;d:2;h:1;i:1;j:1
C2H2*	130	122	57	22	14	a:5;b:15;c:1;d:1;f:1; i:5;j:6;o:2
C3H	33	31	13	5	2	a:1;b:3;c:1;j:1;n:1
CCAAT-HAP2	10	10	5	3	0	b:2;l:1
CCAAT-HAP3	11	11	7	2	2	a:2;i:1;j:1
CCAAT-HAP5	13	12	5	3	1	a:1;c:1;f:1;o:1
GARP-ARR-BARR19	12	12	4	1	1	b:1
GARP-G2-like*	43	43	26	8	8	b:4;c:1;d:3;f:1;i:3; j:4
GeBP	16	14	4	0	0	n/a
GRAS*	32	30	14	4	5	b:1;c:1;d:1;i:2;j:2;p:1
HB*	94	92	40	15	3	a:3;b:9;d:1;e:1;f:2;i:1; j:1;l:2;m:1
HMG	10	8	5	0	4	i:1;j:2;n:1
HSF*	24	23	13	10	1	a:6;b:4;d:1
JUMONJI	13	13	4	2	0	a:1;b:1;d:1
LIM	6	6	4	0	3	h:1;j:2;o:1
MADS	107	96	17	2	7	a:1;f:1;i:2;j:3;o:1
MYB*	203	188	84	33	12	a:9;b:18;c:1;d:5;f:2;i:3; j:5;n:1;o:1
NAC*	113	107	47	26	7	a:9;b:10;c:1;d:1;e:1;f:2; g:1;j:4;o:2
Nin-like	14	14	6	0	2	j:2
PHD	11	10	4	1	1	b:1;j:1
PLATZ*	9	8	5	2	0	b:2
SBP	16	15	8	3	1	a:1;b:2;i:1
TAZ	9	9	4	2	0	b:1;d:1;f:1
TCP	24	23	5	0	0	d:1

Trihelix	28	28	15	3	5	a:1;b:2;i:3;j:1;n:1
TUB	11	10	6	1	2	a:1;i:2
WRKY*	72	65	35	18	8	a:4;b:10;c:2;d:2;i:4; j:2;l:1
ZF-HD	15	15	6	1	2	d:1;j:2
ZIM*	15	12	10	8	1	b:3;c:3;d:1;f:1;o:1

Groups of genes that are significantly enriched in at least one STEM profile at a FDR of <5% are marked with an asterisk (*). The final column details the frequency of assignment of individual genes to specific STEM profiles identified by corresponding letters in Fig 2-3.

Functional categories in this table are defined according to the Database of Arabidopsis Transcription Factors (Guo *et al.*, 2005). The following small transcription factor families, which are each represented by probes for three or fewer genes on the microarray, are not listed in this table: ARID, BES1, C2C2-YABBY, CAMTA, CCAAT-DR1, CPP, E2F/DPE2FC, EIL, FHA, GIF, GRF, HRT-like, LFY, LUG, MBF1, NZZ, PBF-2-like(Whirly), PcG, S1Fa-like, SAP, SRS, ULT, VOZ.

responsive to stresses or hormones (Chen *et al.*, 2006). Consistent with this report, we observed that at least one third of the 84 detectable MYB TFs were induced by NaCl at the transcript abundance level, and only a small fraction were repressed (Table 2-6). *MYB15*(At3g23250), which is a R2R3-type MYB gene (Stracke *et al.*, 2001), was induced approximately 16-fold by NaCl at each of the three time points, and this was confirmed by qRT-PCR (Table 2-1). Similarly, most of the HSF TFs detected on our array were up-regulated by NaCl, as expected for this family of stress-inducible TFs (Busch *et al.*, 2005)(Table 2-6). The AP2/EREBP family includes some of the best characterized stress-responsive transcription factors, including the CBF (CRT/DRE binding factors) proteins (Ohmetakagi and Shinshi, 1995; Gilmour *et al.*, 1998). We observed NaCl-inducible transcript accumulation in more than half of the 76 AP2/EREBP TFs detectable on our array (Table 2-6); and these up-regulated transcripts were detected in each of the AP2/EREBP family's subgroups (i.e. AP2, RAV, ERF I-X) (Nakano *et al.*, 2006). We arbitrarily selected one AP2/EREBP TF (At1g44830, TINY-related) for confirmation by qRT-PCR. As shown in Table 2-1, both microarray and qRT-PCR analyses were in agreement about the strong inducibility of transcripts of At1g44830 following NaCl-treatment. The majority of WRKY family TF genes are known to be responsive to biotic and/or abiotic stress, although most researches have

focused on the roles of these genes in pathogen responses (Seki *et al.*, 2002a; Dong *et al.*, 2003; Narusaka *et al.*, 2004; Xu *et al.*, 2006). We observed that 18 transcripts out of the 35 WRKY TFs increased by at least 1.5 fold in response to NaCl treatment. *WRKY17* (At2g24570), *WRKY25* (At2g30250) and *WRKY33* (At2g38470) were chosen for further qRT-PCR validation. Both microarray and qRT-PCR confirmed that *WRKY17* and *WRKY33* transcripts increased at least 14-fold following NaCl treatment, although *WRKY17* transcript abundance peaked at 6h and gradually diminished, whereas *WRKY33* abundance remained very high throughout the time points sampled. The transcript level of *WRKY25* also increased and peaked at 48 h with an induction of at least 22-fold. These observations emphasize a potentially important role for WRKY transcription factors in abiotic stress responses.

Many TFs contain zinc-finger motifs including all members of the following families: C2H2 (Cys2-His2), C2C2 (Cys2-His2), C3H (Cys3-His) LIM (Lin-11, Isl-1 and Mec-3), PHD (Plant Homeodomain), WRKY, ZF-HD (zinc finger homeodomain), and ZIM (Zinc-finger protein expressed in Inflorescence Meristem). Of these, we observed the WRKY (as described above), and C2C2, C2H2, and ZIM families were particularly enriched in NaCl-responsive transcripts. The C2C2 and C2H2 families are relatively large, and had significant numbers both of up-regulated and down-regulated genes. The smaller ZIM family was represented by 10 genes with detectable transcripts in our analysis; 8 of these genes were up-regulated by NaCl treatment. Knowledge about the biological role of ZIM TFs is very limited. Previous studies have shown a putative relationship of some ZIM family members with GA-independent cell elongation in shoot tissues (Nishii *et al.*, 2000; Shikata *et al.*, 2004). The large proportion of ZIM family members with NaCl-inducible transcripts makes this group of TFs another attractive target for further functional characterization. We assayed the expression pattern of one ZIM transcription factor (At3g17860) by qRT-PCR and confirmed that it is induced after 6h of NaCl, although the transcript abundance falls off rapidly after 24h and 48h (Table 2-1).

Four other large families of TFs, each with distinct DNA binding motifs, are: bHLH (basic helix-loop-helix), bZIP (basic leucine zipper), NAC (NAM, ATAF1,2, CUC2), and HB (homeobox) genes. All of these families contained NaCl-induced

genes, although only the HB and NAC families were significantly enriched in NaCl-responsive transcripts. We selected one bHLH TF (*bHLH92*, At5g43650) and one NAC TF (At4g01550) for further confirmation by qRT-PCR. The NAC domain TF was chosen because its expression peaked at later time points than most TFs; *bHLH92* was selected because it was among the mostly highly induced transcripts of any gene on our microarray. Furthermore, this bHLH is not represented among probes on Affymetrix arrays, and therefore would not be detected in some previous transcriptomic studies, although it was among the NaCl-inducible transcripts identified in a long-oligomer microarray analysis of whole Arabidopsis plants (Kamei *et al.*, 2005). Our qRT-PCR analysis showed a NaCl-dependent increase of up to 50-fold for the NAC TF (At4g01550) transcripts, and an increase for *bHLH92* of over several magnitudes (Table 2-1). Clearly, these two TFs, and many of the other NaCl-responsive TFs shown in Table 2-6 are potentially important regulators of the NaCl-stress response in Arabidopsis roots.

An interesting property of the groups of transcription factors described in Table 2-6 is that the majority of the families (34/39) contain members that are assigned to different expression profiles. In many cases, related transcription factors show very divergent expression patterns, with some members of a single gene family induced early in the stress response, while other members are induced only at subsequent time points. For example, while the majority of AP2/ERF family members have a peak relative expression ratio at 6h (Table 2-6, and Fig 2-3, STEM profile b), some AP2/ERF transcripts had peak expression ratios at 24h (profile f), or increased in abundance throughout the experiment until at least 48h (profile a), while other genes did not show a significant relative change in expression until 48h (profile h), or showed a more complex pattern of regulation (profiles c, d, e), or were predominantly down-regulated (profiles i, j, k, l, o). Similarly divergent patterns of transcriptional responses have been described previously within families of oxidative stress inducible genes (Kim *et al.*, 2005). The large number of up- or down- regulated transcription factor genes we detected, as well as the diversity of their temporal expression patterns, is consistent with the existence of a highly complex regulatory network underlying the physiological response to NaCl treatment.

Conclusions

In this study, we identified 2,367 salt upregulated and 2,913 downregulated genes in *Arabidopsis* using the QiagenOperon microarray. Thus, microarray profiling of NaCl-treated *Arabidopsis* roots revealed dynamic changes in transcript abundance for ca. 22% of the genome for all the three time points examined, including hundreds of transcription factors, kinases/phosphatases, hormone-related genes, and effectors of homeostasis, all of which highlight the complexity of this stress response.

Many transcription factors were induced during salt stress. By contrast, fewer transcription factors were downregulated during salt stress. These results suggest that salt responses in plants are initiated mainly by transcriptional activation rather than repression of genes. The downregulation of other transcription factors later in salt stressed roots may be the result of early activation of transcription factors.

Many of the salt-regulated genes reported here are regulated not only by salt but also by other abiotic and biotic stresses and by ABA (Seki *et al.*, 2002a, Seki *et al.*, 2002b; Kreps *et al.* 2002; Fowler and Thomashow, 2002; Lee *et al.*, 2005). Different signal transduction pathways may crosstalk and converge in the activation of these stress genes.

The observations of the transcriptomic response to NaCl-treatment in *Arabidopsis* roots were consistent with the broad themes of abiotic stress responses previously reported in other systems (Sahi *et al.*, 2006). However, our functional enrichment analysis showed that a considerable amount of sub-functionalization has occurred within many gene families, demonstrating the complex relationship between stress and processes related to reactive oxygen species, cell wall growth, transport, signal transduction, hormone signalling, proteolysis and transcriptional regulation. Moreover, STEM enrichment analysis allowed us to detect previously underappreciated families within these broad categories that are strongly associated with stress response, including MATE transporters, LeOPT1-like transporters. PERK kinases, MLO-like receptors, carbohydrate active enzymes (e.g. XTH18), transcription factors, and other proteins (e.g. 4CL-like, COMT-like, LOB-Class 1). These data will facilitate the mapping of genetic regulatory networks in roots of *Arabidopsis* and a better understanding of stress phenomena in plants. The identified novel salt-responsive genes also

extend our ability to improve salt tolerance in plants. Further research could be directed towards exploring the network of gene regulation in the salt and determining the function of salt-responsive genes in salt tolerance through mutant analysis, overexpression, and other genetic and molecular biological approaches.

References

- Alsheikh MK, Svensson JT and Randall SK.** (2005) Phosphorylation regulated ion-binding is a property shared by the acidic subclass dehydrins. *Plant Cell Environ* 28: 1114-1122
- Arabidopsis Genome Initiative.** (2000) Analysis of the genome sequence of the flowering plant *Arabidopsis thaliana*. *Nature* 408: 796-815
- Atkin RK, Barton GE and Robinson DK.** (1973) Effect of root-growing temperature on growth-substances in xylem exudate of *Zea Mays*. *J Exp Bot* 24: 475-487
- Avonce N, Leyman B, Mascorro-Gallardo JO, Van Dijck P, Thevelein JM and Iturriaga G.** (2004) The Arabidopsis trehalose-6-P synthase AtTPS1 gene is a regulator of glucose, abscisic acid, and stress signaling. *Plant Physiol* 136: 3649-3659
- Barakat A, Szick-Miranda K, Chang IF, Guyot R, Blanc G, Cooke R, Delseny M and Bailey-Serres J.** (2001) The organization of cytoplasmic ribosomal protein genes in the arabidopsis genome. *Plant Physiol* 127: 398-415
- Barlier I, Kowalczyk M, Marchant A, Ljung K, Bhalerao R, Bennett M, Sandberg G and Bellini C.** (2000) The SUR2 gene of Arabidopsis thaliana encodes the cytochrome P450CYP83B1, a modulator of auxin homeostasis. *Proc Natl Acad Sci USA* 97: 14819-14824
- Bates LS, Waldren RP and Teare ID.** (1973) Rapid determination of free proline for water-stress studies. *Plant Soil* 39: 205-207
- Birnbaum K and Benfey PN.** (2004) Network building: transcriptional circuits in the root. *Curr Opin Plant Biol* 7: 582-588
- Boursiac Y, Chen S, Luu DT, Sorieul M, van den Dries N and Maurel C.** (2005) Early effects of salinity on water transport in Arabidopsis roots. Molecular and cellular features of aquaporin expression. *Plant Physiol* 139: 790-805
- Busch W, Wunderlich M and Schoffl F.** (2005) Identification of novel heat shock

factor-dependent genes and biochemical pathways in *Arabidopsis thaliana*. *Plant J* 41: 1-14

Chen KM, Gong HJ, Chen GC, Wang SM and Zhang CL. (2004) Gradual drought under field conditions influences glutathione metabolism, redox balance and energy supply in spring wheat. *J Plant Growth Regul* 23: 20-28

Chen W, Provart NJ, Glazebrook J, Katagiri F, Chang H-S, Eulgem T, Mauch F, Luan S, Zou G, Whitham SA, Budworth PR, Tao Y, Xie Z, Chen X, Lam S, Kreps JA, Harper JF, Si-Ammour A, Mauch-Mani B, Heinlein M, Kobayashi K, Hohn T, Dangl JL, Wang X and Zhu T. (2002) Expression profile matrix of *Arabidopsis* transcription factor genes suggests their putative functions in response to environmental stresses. *Plant Cell* 14: 559-574

Chen YF, Etheridge N and Schaller GE. (2005) Ethylene signal transduction. *Ann Bot* 95: 901-915

Chen YH, Yang XY, He K, Liu MH, Li JG, Gao ZF, Lin ZQ, Zhang YF, Wang XX, Qiu XM, Shen YP, Zhang L, Deng XH, Luo JC, Deng XW, Chen ZL, Gu HY and Qu LJ. (2006) The MYB transcription factor superfamily of *Arabidopsis*: Expression analysis and phylogenetic comparison with the rice MYB family. *Plant Mol Biol* 60: 107-124

Chen ZY, Hartmann HA, Wu MJ, Friedman EJ, Chen JG, Pulley M, Schulze-Lefert P, Panstruga R and Jones AM. (2006) Expression analysis of the AtMLO gene family encoding plant-specific seven-transmembrane domain proteins. *Plant Mol Biol* 60: 583-597

Coutinho PM and Henrissat B. (1999) Carbohydrate-active enzymes: an integrated database approach. *In* HJ Gilbert, G Davies, B Henrissat, B Svensson, eds, *Recent Advances in Carbohydrate Bioengineering*. The Royal Society of Chemistry, Cambridge, pp 3-12

Czechowski T, Stitt M, Altmann T, Udvardi MK, and Scheible W-R. (2005) Genome-wide identification and testing of superior reference genes for transcript normalization in *Arabidopsis*. *Plant Physiol* 2005 139: 5-17.

Dani V, Simon WJ, Duranti M and Croy RRD. (2005) Changes in the tobacco leaf apoplast proteome in response to salt stress. *Proteomics* 5: 737-745

- De Smet I, Vanneste S, Inze D and Beeckman T.** (2006) Lateral root initiation or the birth of a new meristem. *Plant Mol Biol* 60: 871-887
- Debnam PM, Fernie AR, Leisse A, Golding A, Bowsher CG, Grimshaw C, Knight JS and Emes MJ.** (2004) Altered activity of the P2 isoform of plastidic glucose 6-phosphate dehydrogenase in tobacco (*Nicotiana tabacum* cv. Samsun) causes changes in carbohydrate metabolism and response to oxidative stress in leaves. *Plant J* 38: 49-59
- DeLille JM, Sehnke PC and Ferl RJ.** (2001) The arabidopsis 14-3-3 family of signaling regulators. *Plant Physiol* 126: 35-38
- Devoto A, Hartmann HA, Piffanelli P, Elliott C, Simmons C, Taramino G, Goh CS, Cohen FE, Emerson BC, Schulze-Lefert P and Panstruga R.** (2003) Molecular phylogeny and evolution of the plant-specific seven-transmembrane MLO family. *J Mol Evol* 56: 77-88
- Devoto A and Turner JG.** (2005) Jasmonate-regulated Arabidopsis stress signalling network. *Physiol Plant* 123: 161-172
- Dong JX, Chen CH and Chen ZX.** (2003) Expression profiles of the Arabidopsis WRKY gene superfamily during plant defense response. *Plant Mol Biol* 51: 21-37
- Draghichi S.** (2003) *Data Analysis Tools for DNA Microarrays.* Chapman & Hall/CRC, Boca Raton
- Ernst J and Bar-Joseph Z.** (2006) STEM: a tool for the analysis of short time series gene expression data. *BMC Bioinformatics* 7
- Fowler S and Thomashow MF.** (2002). Arabidopsis transcriptome profiling indicates that multiple regulatory pathways are activated during cold acclimation in addition to the CBF cold response pathway. *Plant Cell* 14, 1675–1690
- Gilmour SJ, Artus NN and Thomashow MF.** (1992) cDNA sequence analysis and expression of 2 cold-regulated genes of *Arabidopsis thaliana*. *Plant Mol Biol* 18: 13-21
- Gilmour SJ, Zarka DG, Stockinger EJ, Salazar MP, Houghton JM and Thomashow MF.** (1998) Low temperature regulation of the Arabidopsis CBF family of AP2 transcriptional activators as an early step in cold-induced COR gene expression. *Plant J* 16: 433-442
- Gong QQ, Li PH, Ma SS, Rupassara SI and Bohnert HJ.** (2005) Salinity stress

adaptation competence in the extremophile *Thellungiella halophila* in comparison with its relative *Arabidopsis thaliana*. *Plant J* 44: 826-839

Gould KS. (2004) Nature's Swiss army knife: The diverse protective roles of anthocyanins in leaves. *J Biomed Biotech* 314-320

Hauschild R and von Schaewen A. (2003) Differential regulation of glucose-6-phosphate dehydrogenase isoenzyme activities in potato. *Plant Physiol* 133: 47-62

He XJ, Mu RL, Cao WH, Zhang ZG, Zhang JS and Chen SY. (2005) AtNAC2, a transcription factor downstream of ethylene and auxin signaling pathways, is involved in salt stress response and lateral root development. *Plant J* 44: 903-916

He ZY, Li LG and Luan S. (2004) Immunophilins and parvulins. Superfamily of peptidyl prolyl isomerases in *Arabidopsis*. *Plant Physiol* 134: 1248-1267

Hwang D, Chen HC and Sheen J. (2002) Two-component signal transduction pathways in *Arabidopsis*. *Plant Physiol* 129: 500-515

Hwang I and Sheen J. (2001) Two-component circuitry in *Arabidopsis* cytokinin signal transduction. *Nature* 413: 383-389

Iida K, Seki M, Sakurai T, Satou M, Akiyama K, Toyoda T and Konagaya A K S. (2005) RARTF: database and tools for complete sets of *Arabidopsis* transcription factors. *DNA Res* 12: 247-256

Jones L and McQueen-Mason S. (2004) A role for expansins in dehydration and rehydration of the resurrection plant *Craterostigma plantagineum*. *FEBS Lett* 559: 61-65

Kamei A, Seki M, Umezawa T, Ishida J, Satou M, Akiyama K, Zhu JK and Shinozaki K. (2005) Analysis of gene expression profiles in *Arabidopsis* salt overly sensitive mutants *sos2-1* and *sos3-1*. *Plant Cell Environ* 28: 1267-1275

Kanehisa M and Goto S. (2000) KEGG: Kyoto Encyclopedia of Genes and Genomes. *Nucleic Acids Res* 28: 27-30

Kawaguchi R, Girke T, Bray EA and Bailey-Serres J. (2004) Differential mRNA translation contributes to gene regulation under non-stress and dehydration stress conditions in *Arabidopsis thaliana*. *Plant J* 38: 823-829

Kilili KG, Atanassova N, Vardanyan A, Clatot N, Al-Sabarna K, Kanellopoulos PN, Makris AM and Kampranis SC. (2004) Differential roles of Tau class

- glutathione S-transferases in oxidative stress. *J Biol Chem* 279: 24540-24551
- Kim HS, Yu Y, Snesrud EC, Moy LP, Linford LD, Haas BJ, Nierman WC and Quackenbush J.** (2005) Transcriptional divergence of the duplicated oxidative stress-responsive genes in the *Arabidopsis* genome. *Plant J* 41: 212-220
- Kreps JA, Wu YJ, Chang HS, Zhu T, Wang X and Harper JF.** (2002) Transcriptome changes for *Arabidopsis* in response to salt, osmotic, and cold stress. *Plant Physiol* 130: 2129-2141
- Larkindale J, Mishkind M and Vierling E.** (2005) Plant responses to high temperature. *In* MA Jenks, PM Hasegawa, eds, *Plant Abiotic Stress*. Blackwell, Oxford, pp 100-144
- Lee Y, Choi D and Kende H.** (2001) Expansins: ever-expanding numbers and functions. *Curr Opin Plant Biol* 4: 527-532
- Lee BH, Henderson DA, and Zhu J-K.** (2005) The *Arabidopsis* cold-responsive transcriptome and its regulation by ICE1. *Plant Cell* 17:3155-3175
- Leung J, Merlot S and Giraudat J.** (1997) The *Arabidopsis* ABSCISIC ACID-INSENSITIVE2 (ABI2) and ABI1 genes encode homologous protein phosphatases 2C involved in abscisic acid signal transduction. *Plant Cell* 9: 759-771
- Lin BL, Wang JS, Liu HC, Chen RW, Meyer Y, Barakat A and Delseny M.** (2001) Genomic analysis of the Hsp70 superfamily in *Arabidopsis thaliana*. *Cell Stress Chaperones* 6: 201-208
- Ludwig AA, Saitoh H, Felix G, Freymark G, Miersch O, Wasternack C, Boller T, Jones JDG and Romeis T.** (2005) Ethylene-mediated cross-talk between calcium-dependent protein kinase and MAPK signaling controls stress responses in plants. *Proc Natl Acad Sci USA* 102: 10736-10741
- Ma SS, Gong QQ and Bohnert HJ.** (2006) Dissecting salt stress pathways. *J Exp Bot* 57: 1097-1107
- Maathuis FJM.** (2006) The role of monovalent cation transporters in plant responses to salinity. *J Exp Bot* 57:1137-1147
- Maathuis FJM, Filatov V, Herzyk P, Krijger GC, Axelsen KB, Chen SX, Green BJ, Li Y, Madagan KL, Sanchez-Fernandez R, Forde BG, Palmgren MG, Rea PA, Williams LE, Sanders D and Amtmann A.** (2003) Transcriptome analysis of

root transporters reveals participation of multiple gene families in the response to cation stress. *Plant J* 35: 675-692

Mancinelli AL and Schwartz OM. (1984) The photoregulation of anthocyanin synthesis IX. the photosensitivity of the response in dark and light-grown tomato seedlings. *Plant Cell Physiol* 25: 93-105

Merlot S, Gosti F, Guerrier D, Vavasseur A and Giraudat J. (2001) The ABI1 and ABI2 protein phosphatases 2C act in a negative feedback regulatory loop of the abscisic acid signalling pathway. *Plant J* 25: 295-303

Mittler R. (2006) Abiotic stress, the field environment and stress combination. *Trends Plant Sci* 11: 15-19

Mittler R, Vanderauwera S, Gollery M and Van Breusegem F. (2004) Reactive oxygen gene network of plants. *Trends Plant Sci* 9: 490-498

Moller IM. (2001) Plant mitochondria and oxidative stress: Electron transport, NADPH turnover, and metabolism of reactive oxygen species. *Annu Rev Plant Physiol Plant Mol Biol* 52: 561-591

Nakano T, Suzuki K, Fujimura T and Shinshi H. (2006) Genome-wide analysis of the ERF gene family in Arabidopsis and rice. *Plant Physiol* 140: 411-432

Narusaka Y, Narusaka M, Seki M, Umezawa T, Ishida J, Nakajima M, Enju A and Shinozaki K. (2004) Crosstalk in the responses to abiotic and biotic stresses in Arabidopsis: Analysis of gene expression in cytochrome P450 gene superfamily by cDNA microarray. *Plant Mol Biol* 55: 327-342

Ndimba BK, Chivasa S, Simon WJ and Slabas AR. (2005) Identification of Arabidopsis salt and osmotic stress responsive proteins using two-dimensional difference gel electrophoresis and mass spectrometry. *Proteomics* 5: 4185-4196

Nieuwland J, Feron R, Huisman BAH, Fasolino A, Hilbers CW, Derksen J and Mariani C. (2005) Lipid transfer proteins enhance cell wall extension in tobacco. *Plant Cell* 17: 2009-2019

Nishii A, Takemura M, Fujita H, Shikata M, Yokota A and Kohchi T. (2000) Characterization of a novel gene encoding a putative single zinc-finger protein, ZIM, expressed during the reproductive phase in Arabidopsis thaliana. *Biosci Biotech Biochem* 64: 1402-1409

- Noctor G.** (2006) Metabolic signalling in defence and stress: the central roles of soluble redox couples. *Plant Cell Environ* 29: 409-425
- Ohmetakagi M and Shinshi H.** (1995) Ethylene-inducible DNA-binding proteins that interact with an ethylene-responsive element. *Plant Cell* 7: 173-182
- Passardi F, Cosio C, Penel C and Dunand C.** (2005) Peroxidases have more functions than a Swiss army knife. *Plant Cell Rep* 24: 255-265
- Pfaffl MW.** (2001) A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Res.* 29:e45.
- Raes J, Rohde A, Christensen JH, Van de Peer Y and Boerjan W.** (2003) Genome-wide characterization of the lignification toolbox in Arabidopsis. *Plant Physiol* 133: 1051-1071
- Rawlings ND, Tolle DP and Barrett AJ.** (2004) MEROPS: the peptidase database. *Nucleic Acids Res* 32: D160-D164
- Rhee SY, Beavis W, Berardini TZ, Chen GH, Dixon D, Doyle A, Garcia-Hernandez M, Huala E, Lander G, Montoya M, Miller N, Mueller LA, Mundodi S, Reiser L, Tacklind J, Weems DC, Wu YH, Xu I, Yoo D, Yoon J and Zhang PF.** (2003) The Arabidopsis Information Resource (TAIR): a model organism database providing a centralized, curated gateway to Arabidopsis biology, research materials and community. *Nucleic Acids Res* 31: 224-228
- Rizhsky L, Liang HJ and Mittler R.** (2002) The combined effect of drought stress and heat shock on gene expression in tobacco. *Plant Physiol* 130: 1143-1151
- Rogers EE and Guerinot ML.** (2002) FRD3, a member of the multidrug and toxin efflux family, controls iron deficiency responses in Arabidopsis. *Plant Cell* 14: 1787-1799
- Saeed AI, Sharov V, White J, Li J, Liang W, Bhagabati N, Braisted J, Klapa M, Currier T, Thiagarajan M, Sturn A, Snuffin M, Rezantsev A, Popov D, Ryltsov A, Kostukovich E, Borisovsky I, Liu Z, Vinsavich A, Trush V and Quackenbush J.** (2003) TM4, A free, open-source system for microarray data management and analysis. *Biotechniques* 34: 374-378
- Saher S, Fernandez-Garcia N, Piqueras A, Hellin E and Olmos E.** (2005) Reducing properties, energy efficiency and carbohydrate metabolism in hyperhydric and normal

- carnation shoots cultured in vitro: a hypoxia stress? *Plant Physiol Biochem* 43: 573-582
- Sahi C, Singh A, Blumwald E and Grover A.** (2006) Beyond osmolytes and transporters: novel plant salt-stress tolerance-related genes from transcriptional profiling data. *Physiol Plant* 127: 1-9
- Seki M, Narusaka M, Ishida J, Nanjo T, Fujita M, Oono Y, Kamiya A, Nakajima M, Enju A, Sakurai T, Satou M, Akiyama K, Taji T, Yamaguchi-Shinozaki K, Carninci P, Kawai J, Hayashizaki Y and Shinozaki K.** (2002a) Monitoring the expression profiles of 7000 Arabidopsis genes under drought, cold and high-salinity stresses using a full-length cDNA microarray. *Plant J* 31: 279-292
- Seki M, Ishida J, Narusaka M, Fujita M, Nanjo T, Umezawa T, Kamiya A, Nakajima M, Enju A, Sakurai T, Satou M, Akiyama K, Yamaguchi-Shinozaki K, Carninci P, Kawai J, Hayashizaki Y and Shinozaki K.** (2002b) Monitoring the expression pattern of around 7,000 Arabidopsis genes under ABA treatments using a full-length cDNA microarray. *Funct Integr Genomics*. 2: 282-291.
- Shikata M, Matsuda Y, Ando K, Nishii A, Takemura M, Yokota A and Kohchi T.** (2004) Characterization of Arabidopsis ZIM, a member of a novel plant-specific GATA factor gene family. *J Exp Bot* 55: 631-639
- Shiu SH, Karlowski WM, Pan RS, Tzeng YH, Mayer KFX and Li WH.** (2004) Comparative analysis of the receptor-like kinase family in Arabidopsis and rice. *Plant Cell* 16: 1220-1234
- Shuai B, Reynaga-Pena CG and Springer PS.** (2002) The LATERAL ORGAN BOUNDARIES gene defines a novel, plant-specific gene family. *Plant Physiol* 129: 747-761
- Smalle J and Vierstra RD.** (2004) The ubiquitin 26S proteasome proteolytic pathway. *Annu Rev Plant Biol* 55: 555-590
- Steppuhn H and Raney JP.** (2005) Emergence, height, and yield of canola and barley grown in saline root zones. *Can J Plant Sci* 85: 815-827
- Stone SL, Hauksdottir H, Troy A, Herschleb J, Kraft E and Callis J.** (2005) Functional analysis of the RING-type ubiquitin ligase family of Arabidopsis. *Plant Physiol* 137: 13-30

- Stracke R, Werber M and Weisshaar B.** (2001) The R2R3-MYB gene family in *Arabidopsis thaliana*. *Curr Opin Plant Biol* 4: 447-456
- Sung DY, Vierling E and Guy CL.** (2001) Comprehensive expression profile analysis of the *Arabidopsis hsp70* gene family. *Plant Physiol* 126: 789-800
- Taji T, Seki M, Satou M, Sakurai T, Kobayashi M, Ishiyama K, Narusaka Y, Narusaka M, Zhu JK and Shinozaki K.** (2004) Comparative genomics in salt tolerance between *Arabidopsis* and *Arabidopsis*-related halophyte salt cress using *Arabidopsis* microarray. *Plant Physiol* 135: 1697-1709
- Takahashi S, Seki M, Ishida J, Satou M, Sakurai T, Narusaka M, Kamiya A, Nakajima M, Enju A, Akiyama K, Yamaguchi-Shinozaki K and Shinozaki K.** (2004) Monitoring the expression profiles of genes induced by hyperosmotic, high salinity, and oxidative stress and abscisic acid treatment in *Arabidopsis* cell culture using a full-length cDNA microarray. *Plant Mol Biol* 56: 29-55
- Tchieu JH, Fana F, Fink JL, Harper J, Nair TM, Niedner RH, Smith DW, Steube K, Tam TM, Veretnik S, Wang DG and Gribskov M.** (2003) The PlantsP and PlantsT functional Genomics Databases. *Nucleic Acids Res* 31: 342-344
- Tognolli M, Penel C, Greppin H and Simon P.** (2002) Analysis and expression of the class III peroxidase large gene family in *Arabidopsis thaliana*. *Gene* 288: 129-138
- Tusher VG, Tibshirani R and Chu G.** (2001) Significance analysis of microarrays applied to the ionizing radiation response. *Proc Natl Acad Sci USA* 98: 5116-5121
- Urao T, Yakubov B, Satoh R, Yamaguchi-Shinozaki K, Seki M, Hirayama T and Shinozaki K.** (1999) A transmembrane hybrid-type histidine kinase in *Arabidopsis* functions as an osmosensor. *Plant Cell* 11: 1743-1754
- Verbruggen N, Villarreal R and Vanmontagu M.** (1993) Osmoregulation of a Pyrroline-5-Carboxylate Reductase Gene in *Arabidopsis thaliana*. *Plant Physiol* 103: 771-781
- Vissenberg K, Oyama M, Osato V, Yokoyama R, Verbelen JP and Nishitani K.** (2005) Differential expression of AtXTH17, AtXTH18, AtXTH19 and AtXTH20 genes in *Arabidopsis* roots. Physiological roles in specification in cell wall construction. *Plant Cell Physiol* 46: 192-200
- Wakao S and Benning C.** (2005) Genome-wide analysis of glucose-6-phosphate

dehydrogenases in *Arabidopsis*. *Plant J* 41: 243-256

Wang H, Miyazaki S, Kawai K, Deyholos M, Galbraith DW and Bohnert HJ.

(2003) Temporal progression of gene expression responses to salt shock in maize roots. *Plant Mol Biol* 52: 873-891

Ward JM. (2001) Identification of novel families of membrane proteins from the model plant *Arabidopsis thaliana*. *Bioinformatics* 17: 560-563

Welin BV, Olson A, Nylander M and Palva ET. (1994) Characterization and differential expression of Dhn/Lea/Rab-like genes during cold-acclimation and drought stress in *Arabidopsis thaliana*. *Plant Mol Biol* 26: 131-144

Woodward AW and Bartel B. (2005) Auxin: Regulation, action, and interaction. *Ann Bot* 95: 707-735

Xu XP, Chen CH, Fan BF and Chen ZX. (2006) Physical and functional interactions between pathogen-induced *Arabidopsis* WRKY18, WRKY40, and WRKY60 transcription factors. *Plant Cell* 18: 1310-1326

Yamaguchishinozaki K, Koizumi M, Urao S and Shinozaki K. (1992) Molecular cloning and characterization of 9 cDNAs for genes that are responsive to desiccation in *Arabidopsis thaliana*: Sequence analysis of one cDNA clone that encodes a putative transmembrane channel protein. *Plant Cell Physiol* 33: 217-224

Yoshida Y, Kiyosue T, Nakashima K, Yamaguchi-Shinozaki K and Shinozaki K. (1997) Regulation of levels of proline as an osmolyte in plants under water stress. *Plant Cell Physiol* 38: 1095-1102

Yoshida T, Nishimura N, Kitahata N, Kuromori T, Ito T, Asami T, Shinozaki K and Hirayama T. (2006) ABA-Hypersensitive germination3 encodes a protein phosphatase 2C (AtPP2CA) that strongly regulates abscisic acid signaling during germination among *Arabidopsis* protein phosphatase 2Cs. *Plant Physiol* 140: 115-126

Zhang PF, Foerster H, Tissier CP, Mueller L, Paley S, Karp PD and Rhee SY.

(2005) MetaCyc and AraCyc. Metabolic pathway databases for plant research. *Plant Physiol* 138: 27-37

Chapter 3: Comparative proteomic analysis of Arabidopsis root response to salt stress

Introduction

Soil salinity is a prevalent abiotic stress that limits the productivity and geographical distribution of plants. Roots are the site of perception and injury for several types of stress, including salinity, nutrient deficiency and heavy metals. Under many circumstances, it is the stress-sensitivity of the root that limits the productivity of the entire plant (Atkin *et al.*, 1973; Steppuhn and Raney, 2005). An improved understanding of molecular mechanisms of responses in roots to NaCl treatment may therefore facilitate the development of crops with increased tolerance to NaCl and other stresses.

To build a useful description of the molecular mechanisms active in the response of roots to NaCl treatment, it is necessary to characterize the components of these mechanisms, including proteins. Proteomic profiles have been investigated for various stresses and species, including NaCl-treated roots of pea (*Pisum sativum* L.) (Kav *et al.*, 2004), rice (*Oryza sativa* L.) (Yan *et al.*, 2005) and wheat (*Triticum aestivum* L.) (Majoul *et al.*, 2000), drought treated rice (Salekdeh *et al.*, 2002), sugar beet (Hajheidari *et al.*, 2005), poplar (*Populus trichocarpa* Torr. & A.Gray) (Plomion *et al.*, 2006) and chickpea (*Cicer arietinum* L.) (Pandey *et al.*, 2007), and cold treated Arabidopsis (Bae *et al.*, 2003) and rice (Cui *et al.*, 2005), as well as heat treated rice (Lee *et al.*, 2007) and wheat (Majoul *et al.*, 2004). Recently, a 2-DGE based technology was employed to identify NaCl and osmotic responsive proteins in Arabidopsis cell suspension and root microsomal fraction (Ndimba *et al.*, 2005; Lee *et al.*, 2004). In the present study, we applied a moderate NaCl stress to hydroponic cultured Arabidopsis roots to identify proteins that are responsive to NaCl treatment.

Transcriptome profiling, a widely-used technique to identify NaCl-responsive genes, has contributed to our understanding of salinity stress in species including

A version of this chapter was published: Jiang Y-Q, Yang B, Harris NS, Deyholos MK. 2007. Comparative proteomic analysis of NaCl stress-responsive proteins in Arabidopsis roots. *J Exp Bot* 58(13):3591-3607. Adapted with permission from Oxford University Press.

Arabidopsis and rice (Kawasaki *et al.*, 2001; Kreps *et al.*, 2002; Seki *et al.*, 2002; Rabbani *et al.*, 2003; Chao *et al.*, 2005; Jiang and Deyholos, 2006). However, transcriptome profiling has some limitations because mRNA levels are not always correlated to those of corresponding proteins, due in part to post-transcriptional regulation. Only poor or moderate correlation between changes in the levels of specific mRNAs and their corresponding proteins has been previously reported in studies involving yeast (*Saccharomyces cerevisiae*), animals, or Arabidopsis (Gygi *et al.*, 1999; Tian *et al.*, 2004; Mooney *et al.*, 2006). Furthermore, post-translational modifications (PTM), such as phosphorylation and glycosylation, can result in a dramatic increase in proteome complexity without a concomitant increase in gene expression (Rose *et al.*, 2004; Jensen 2004). These biological realities motivated us to perform an analysis of NaCl stress responses at the proteome level, and to compare these results to our previous, microarray-based studies of similarly treated tissues (Jiang and Deyholos, 2006).

Materials and methods

Plant materials and stress treatment

Arabidopsis thaliana plants (wild-type, ecotype Col-0) were cultured hydroponically as described previously (Jiang and Deyholos, 2006). At 20 days after stratification (DAS), the hydroponic solution was changed to fresh, 1 x MS Murashige and Skoog (MS) medium, either with or without 150 mM NaCl, and was maintained for 6 or 48 h, with roots harvested from control or NaCl-treated plants in parallel at each time point and flash frozen in liquid nitrogen and stored at -80 °C. Three biologically independent replicates were prepared at separate times.

Physiological analyses

For root elongation assays, 5 DAS, wild-type Arabidopsis seedlings grown vertically on ½ x MS medium supplemented with 1% sucrose and 0.8% Phytoblend (Caisson Labs) were transferred onto ½ x MS plates supplemented with 1% sucrose and 0, 50, 100, 150, 200, or 250 mM NaCl in square petri dishes, with root lengths measured after 7 d. For ion concentration determination, hydroponic cultured Arabidopsis roots and shoots (20 DAS) were dried for 2 d in an oven at 65°C. 100-500 mg of dry root or leaf

samples was digested according to EPA 3050B method (<http://www.epa.gov/SW-846/pdfs/3050b.pdf>) with modifications. Na and K concentrations were determined by flame emission spectroscopy (AAAnalyst700, PerkinElmer). For the relative electrolyte leakage (REL) assay, ~ 150 mg of hydroponic cultured 18-DAS old seedlings were rinsed with ddH₂O and placed in test tubes containing 10 ml ddH₂O and incubated at room temperature for 2 h with the electrical conductivity of the solution (C₁) measured using conductivity meter (Orion 115Aplus, ThermoElectron). Then, the tubes were boiled for 15 min and cooled to room temperature, and the electrical conductivity (C₂) was measured again. The relative electrolyte leakage was calculated by the formula $C_1/C_2 \times 100\%$ (Cao *et al.*, 2007). Figures were drawn using SigmaPlot (v8.02, Systat Software Inc.).

Protein extraction and quantification

Total protein extracts were prepared essentially according to the method described by Tsugita and Kamo (1999) with modifications. In brief, approximately 1 g of control and NaCl-treated roots were homogenized separately to a fine powder in liquid nitrogen, and were transferred into three 2 ml tubes. 1 ml of 10% (w/v) trichloroacetic acid (TCA)/0.07% DTT in acetone was added to each tube and incubated at -20 °C for 1 h. Afterwards, tubes were centrifuged at 18,000x g for 15 min and the supernatants were discarded. The pellets were washed by resuspension in ice-cold acetone containing 0.07% DTT and centrifuged as described above. This wash was repeated three times, with the pellets dried at room temperature in a SpeedVac for 15 min and resuspended in 300 µL of lysis buffer (30 mM Tris-HCl, 7 M urea, 2 M thiourea, 4% CHAPS, pH 8.5). The samples were mixed vigorously, incubated overnight at 4 °C and centrifuged at room temperature for 15 min at 18,000 g, with the supernatants collected into fresh tubes. The protein extracts were cleaned up using 2-D Cleanup kit (GE Healthcare), dissolved in 200 µL of Destreak rehydration buffer containing 2% pH3-10 immobilized pH gradient (IPG) buffer (GE Healthcare), and quantified by using 2-D Quant kit (GE Healthcare) using bovine serum albumin (BSA) as the standard.

IEF and SDS-PAGE

Two-dimensional electrophoresis of protein extracts was performed using a GE Healthcare 2-DGE system according to the manufacturer's manuals. Briefly, IPG strips

(pH 3–10 NL, 24 cm) were rehydrated in 450 µl of Destreak rehydration buffer (containing 2% pH 3-10 IPG buffer) overnight (~15 h). Isoelectric focusing (IEF) was performed using IGPhor Isoelectric focusing unit with 400 µg of protein samples loaded by cup-loading. The voltage and duration used were as follows: 1st step and hold: 300 V, 3 hours; 2nd gradient: 1000 V, 6 hours; 3rd gradient: 8000 V, 3 hours; 4th step and hold: 8000 V, 4 hours 40 minutes. Prior to second dimension separation, the strips were incubated first in an equilibration buffer (6 M urea, 30% v/v glycerol, and 2% SDS in 0.05 M Tris-HCl, pH 8.8) containing 15 mM DTT for 15 min, then in an equilibration buffer containing 2.5% iodoacetamide (GE Healthcare) for another 15 min, followed by brief equilibration in 1 x SDS-Tris-glycine running buffer for 5 min. The 2nd dimension separation of proteins were performed on SDS-PAGE gel (12.5% acrylamide, Bio-Rad) using the Ettan Dalt Six apparatus (GE Healthcare) with protein markers (Cat#SM0661, Fermentas) loaded at the left most side. The electrophoresis was carried out at ~25 °C and 2.5 w/gel for 30 min and then 17 w/gel for ~5 h 40 min until the bromophenol blue dye front arrived at the bottom of the gels. Following SDS-PAGE, gels were washed in ddH₂O three times for 15 min and proteins were detected by a modified Colloidal Coomassie Brilliant Blue staining-blue silver protocol, which was assessed to have the sensitivity comparable to silver staining (Candiano *et al.*, 2004). After three time washes in ddH₂O, the 2-D gels were scanned immediately using a Fuji FLA-5000 scanner (Fujifilm) with a resolution of 100 µm and 16-bit grayscale pixel depth. A total of 12 gels were analyzed: i.e. three gels (biologically independent replicates) for each of two treatments (0mM or 150mM NaCl) at each of two time points (6 or 48 h). This experimental design was balanced with respect to all conditions, and the use of three gels per condition is consistent with recommendations based on previous statistical analyses of protein gel electrophoresis (Hunt *et al.* 2005). For pI and MW calibrations, we used the 2-D internal standards (Cat#161-0320, Bio-Rad) with IEF, SDS-PAGE, staining and scanning performed the same as above.

Image and statistical analysis

2-D gel images were analyzed using ImageMaster 2-D Platinum 6.0 (GE Healthcare). To verify the autodetected results, all spots were manually inspected and edited as necessary. After spot detection, quantification, and background subtraction, each

analyzed gel was matched individually to the reference gel, and matched spots were grouped into subclasses. To compensate for subtle differences in sample loading, gel staining, and destaining, the volume of each spot (i.e. spot abundance) was normalized as relative volume. This normalization method, provided by ImageMaster 2D Platinum 6.0 software, divides each spot volume value by the sum of total spot volume values to obtain individual relative spot volumes. Class reports were generated for spots of interest. The differences in expression between control and treatment were analyzed by the Student's *t*-test with $P \leq 0.05$ considered significant. The molecular masses of proteins on gels were determined by co-electrophoresis of standard protein markers (Fermentas) and internal 2-D internal standards (Bio-Rad) according to the software manual. For the figures shown in the manuscript, spot IDs were renumbered using the annotation tool in ImageMaster Platinum 6.0 and image brightness and contrast was adjusted and the protein marker sizes were added using Photoshop CS (Adobe).

In-gel digestion and LC-MS/MS analysis

Protein spots showing at least 1.5-fold difference in abundance between control and treatment at one or both time-points with $P < 0.05$ were selected and excised manually into 1.5 ml microtubes. The selection of 1.5-fold change as an arbitrary threshold allowed us to focus on the most responsive proteins for subsequent characterization, and is consistent with thresholds used previously by ourselves and others in microarray studies (Jiang and Deyholos, 2006 and references therein). Gel piece treatment and in-gel digestion of protein spots were performed following Jensen *et al.* (1999) with modifications. Briefly, gel pieces were first washed with 150 μL of HPLC grade water (Fisher Scientific), dehydrated with 50 μL of 100% acetonitrile (ACN), then destained with 100 μL of 50 mM NH_4HCO_3 /50% ACN for 2 h or longer until colorless. After dehydration with ACN again, gel pieces were reduced in 30 μL of 10 mM DTT/0.1 M NH_4HCO_3 at 56 °C for 30 min, dehydrated, and alkylated in 30 μL of 55 mM iodoacetamide/0.1 M NH_4HCO_3 at room temperature for 20 min in the dark, followed by rinsing with 200 μL of 0.1 M NH_4HCO_3 , dehydrated with ACN, dried in a SpeedVac for 10 min. Afterwards, 20 μL of trypsin solution (0.02 $\mu\text{g}/\mu\text{L}$ Trypsin Gold (Promega) in 40 mM NH_4HCO_3 /10% ACN) was added and incubated on ice for 1 h, then at 37 °C overnight (~14 h). Finally, 3 μL of 2% formic acid (FA) was added to

stop the digestion reactions, and the supernatant was collected into fresh tubes, followed by two extractions of peptides with 15 μ L of 50% ACN/0.1% FA with the collect (~50 μ L) mixed well and stored in -20 $^{\circ}$ C before use.

We performed LC-MS/MS analysis of digested peptide mixtures using an Agilent 1100 LC/MSD Trap XCT (Agilent Technologies). Briefly, an autosampler was used to inject 20 μ L of each tryptic digest onto the first of two C-18 columns. This short 5 μ m-enrichment column, Zorbax 300SB-C18 5 μ m 5 x 0.3 mm, served to trap and concentrate the samples. Next, the sample was eluted onto the next C-18 column (Zorbax 300SB-C18 5 μ m 150 x 0.3mm), which was used in conjunction with a solvent gradient to separate the peptides. The peptide-separation gradient started at 85% solvent A (0.1% FA in H₂O) and ended at 55% solvent B (0.1% FA, 5% H₂O in ACN) over a 42-minute span. This was followed by a 10-minute period of 90% solvent B to cleanse the columns before returning to 97% solvent A for the next sample. The ion trap mass spectrometer collected information by first running a MS 300-2200 m/z scan and followed that with a MS/MS analysis of the most intense ions. In addition to the most intense ion for each scan, the software was set to exclude this ion after two spectra and gather MS/MS information on the next most intense ion(s). Raw spectral data was processed into Mascot Generic File (.mgf) format using the default method in the ChemStation Data Analysis module. MS/MS ion search was performed using MASCOT (<http://www.matrixscience.com>) searching the NCBIInr database and taking *Arabidopsis* as the taxonomy. The parameters for searching were: a MS/MS tolerance of ± 0.8 Da, one missed cleavage site, enzyme of trypsin, fixed modifications of carbamidomethyl (C), peptide tolerance of ± 2 Da, peptide charge of 1+ 2+ 3+, monoisotopic and ESI-TRAP instrument. Only significant hits, as defined by the MASCOT probability analysis ($P < 0.05$), were accepted. In case multiple significant hits were found for a protein, only the highest scoring hit was listed in Table 3-1.

Results

Plant growth response to NaCl stress

Arabidopsis thaliana is a glycophyte and is sensitive to NaCl exposure. To find an appropriate concentration of NaCl to use for treatment of plants prior to proteomic

profiling, we performed a root elongation dose response assay. Our results demonstrated that 100 and 150 mM NaCl inhibited root elongation by 53% and 78% respectively (Fig. 3-1A). However, concentrations higher than 200 mM NaCl almost completely inhibited root growth and led to death of almost all seedlings. In a post-stress recovery assay, 20 DAS plants were treated by 150 mM NaCl for 6, 24 or 48 h, and were transferred into fresh media to recover for 1 week. Almost all of the NaCl-treated plants recovered and resumed normal growth (data not shown). We therefore selected 150 mM as our treatment concentration, because it induced visible signs of stress including retarded growth rate and loss of turgor. This concentration of NaCl has been used in several previous gene expression studies, because it induces a moderate stress response and is not acutely lethal (Jiang and Deyholos, 2006; Ma *et al.*, 2006). Higher concentrations of NaCl appear to cause plasmolysis and lethality (Munns 2005), although other previous Arabidopsis studies have used a NaCl treatment concentration of 250, 300 or even 600 mM (Ndimba *et al.*, 2005; Seki *et al.*, 2002).

Relative electrolyte leakage (REL) is an indicator of membrane damage caused by NaCl stress. We measured stress-induced changes in plants treated with 150 mM NaCl. As shown in Fig 3-1B, the REL of seedlings treated by 150 mM NaCl for 6h was 8.6%.




Fig. 3-1. Root elongation assay and relative electrolyte leakage. (A) Effect of increasing [NaCl] on root elongation of seedlings (n=9). (B) Relative electrolyte leakage from roots of control (filled circles) or NaCl-treated (closed circles) roots following 6, 24, or 48 h exposure to 150 mM NaCl (n=5). Vertical bars indicate standard deviation (S.D.).

After 24h treatment, the REL increased to 18% and further increased to be 26% after 48h of NaCl treatment. When compared to the control plants, the REL values of NaCl-treated seedlings were 1.7, 2.8 and 4.4 times higher at 6, 24 and 48 h, respectively, which indicated that stress-induced cellular damage accumulated throughout the duration of the experiment.

To further establish the physiological status of plants that we were to subject to proteome analysis, we next quantified the concentrations of K and Na in both leaves and roots of control and 150 mM-treated plants (Fig. 3-2). Na and K concentrations significantly increased and decreased, respectively, in both leaves and roots under 150 mM NaCl treatment. Moreover, the Na concentrations in roots treated by 150 mM NaCl for 6, 24, and 48h were 115, 167 and 132 fold higher than those of the control roots at comparable time points. The concentration of Na in roots reached a maximum after 24h (Fig. 3-2A), however, leaves continued to accumulate more Na after the 24h (Fig. 3-2B), presumably due to transpiration-driven flux. The K/Na ratios in both leaves and




Fig. 3-2. Changes in Na and K ion concentrations following NaCl exposure. Na (filled symbols) and K (open symbols) concentrations (mg/g dry weight) were determined by flame emission spectroscopy in roots (A) or rosette leaves (B) of hydroponically grown *Arabidopsis* plants, following 6, 24, or 48 h of exposure to media supplemented with either 0 mM NaCl (control, circles) or 150 mM NaCl (treated, triangles). Na concentrations in control samples are very low (< 0.4 mg/g). Data are from 3 biological replicates, and vertical bars indicate \pm S.D.

roots decreased as the NaCl treatment progressed. Part of the basis of Na⁺ toxicity in plants is that high Na⁺ concentration in the cells increases the Na/K ratio, which is adverse for most metabolic processes (Chinnusamy *et al.*, 2005). Thus, the quantitative ion analysis was consistent with the REL data in describing the progressive accumulation of stress symptoms throughout the 48 h duration of the treatments.

2-DGE analysis of NaCl-responsive proteins in Arabidopsis roots

To investigate the temporal changes of protein profiles during NaCl stress, we carried out 2-DGE analysis of the total proteins in Arabidopsis roots from three biologically independent replicate experiments. A representative gel is shown in Fig. 3-3. Approximately 1,000 protein spots were detected on Coomassie brilliant blue-stained gels and about 600 protein spots were matched between 6 control gels and 6 treatment gels. Quantitative image analysis revealed a total of 215 protein spots that changed their abundance (Vol%) significantly ($P \leq 0.05$) by more than 1.5 fold at one or two time points. We selected 1.5-fold as a threshold value in order to identify most responsive proteins which were consistent with our previous microarray experiments (Jiang and Deyholos, 2006). Essentially arbitrary threshold values ranging from 1.3- to 2.0-fold have been used in previous proteomics studies (Casati *et al.*, 2005; Amme *et al.*, 2006; Parker *et al.*, 2006; Yan *et al.*, 2006). We noted that some protein spots also demonstrated qualitative changes in intensity. For example, spots 21, 24, 28, and 75 were absent in the NaCl-treated gels at one or more time points while spots 26, 62, 82 and 85 were absent in control gels at one or more time points.

LC-MS/MS identification and classification of NaCl-responsive proteins

We arbitrarily selected and excised 89 of the 215 differentially expressed spots described above for tryptic digestion and analysis by LC-MS/MS. From 89 gel plugs excised, we successfully identified 86 proteins (Table 3-1), of which 81 were unique according to the protein identity. Some NaCl-responsive spots were not excised because of their low abundance. We found that at least 23 (26%) of these spots contained peptides that matched proteins from unrelated families (Table 3-1), indicating the presence of multiple proteins in some spots. Conversely, we found that four proteins were identified in more than one spot, although they were excised from the same gel (Table 3-1). For example, L-ascorbate peroxidase (APX1) was identified

from three spots (35, 36 and 61), glutathione S-transferase (AtGSTF2) was identified in two spots (41 and 42), glycosyl hydrolase family 17 protein was identified in two spots (26 and 62) and, glycosyl hydrolase family 1 protein was identified twice (65 and 86). Further examination of electrophoresis patterns indicated that the inferred mass or pI




Fig. 3-3. A representative 2-DE gel of Arabidopsis root proteins. 86 of the spots showing at least 1.5-fold change under NaCl treatment at least at one time-point with $p < 0.05$ were analyzed by LC-MS/MS.

values of these spots differed, due perhaps to post-translational modification (PTM) or degradation. PTMs such as glycosylation, phosphorylation, etc, can change the molecular weight and/or charge of proteins. Alternatively, proteins that were present in multiple spots could result from proteins being translated from alternatively spliced mRNAs (Ishikawa *et al.*, 1997). This phenomenon was also reported previously (Holmes-Davis *et al.*, 205; Ndimba *et al.*, 2005). Proteomic studies have also shown that some proteins may be degraded during abiotic stress. For example, rubisco large subunit (RcbL) was detected as 19 different fragments plus the intact protein in NaCl-

treated rice roots (Yan *et al.*, 2006). Similar phenomena have also been reported in pea mitochondrial proteome under chilling stress (Taylor *et al.*, 2005). It is possible that reactive oxygen species (ROS) may also contribute to the degradation of proteins under stress conditions (Desimone *et al.*, 1996; Kingston-Smith and Foyer, 2000).

We classified the identified proteins into 11 categories similar to the convention used by Ndimba *et al.* (2005) (Fig. 3-4). Proteins implicated in energy metabolism (e.g. glycolysis, citrate cycle, electron transport), ROS scavenging and defense, and protein metabolism (e.g. translation, processing and degradation) comprised 52% of the identified proteins. Further examination showed that after 6 h of stress, the abundance of most NaCl-responsive proteins had decreased. In contrast, after 48 h of treatment, we observed that the number of proteins that had increased in abundance was approximately equal to the number of proteins that had decreased in abundance (Fig. 3-4). This suggests that during the initial (6 h) phase of NaCl stress, the synthesis of many proteins was inhibited and/or their degradation increased, while after 48 h, plant roots began to adapt to water deficit and ionic accumulation by synthesizing selected stress-response proteins. This is generally consistent with our previous observations of NaCl-treated roots, in which abundance of transcripts for almost all ribosomal proteins decreased after 6 h, while transcripts for more than 30 peptidases increased at the same time point (Jiang and Deyholos, 2006). However, in our previous microarray analysis, transcripts for ribosomal proteins remained at decreased levels between 6 h and 48 h, suggesting that the increased expression of selected stress proteins we reported here may involve specialized mechanisms of translation that are not reflected in the bulk transcript level of all cellular ribosomal proteins.

Correlation analysis of mRNA and proteins levels

The relationship between gene expression measured at the mRNA level and the corresponding protein level has not been well characterized in plant roots under abiotic stresses. To evaluate the correlation between mRNA and the corresponding protein levels, we compared the differentially expressed protein levels to our previous oligonucleotide microarray data (Jiang and Deyholos, 2006). Tissue for both types of analyses were grown and treated under nearly identical conditions, to minimize the effects of experimental variation in our results. The oligonucleotide probes for two

proteins identified in the proteomic study were not represented in the microarray: ACO2 (At1g62380) and putative tubulin β -2/ β -3 chain (At5g62690). We filtered out those genes whose transcript levels as measured by their signal intensities on arrays did not pass the threshold background intensity and those with low signal in both channels of more than 1/3 of arrays at 6 or 48 h time points. Altogether, 70 and 69 data points were obtained for 6 and 48 h, respectively (Fig. 3-5).




Fig. 3-4. Functional classification of NaCl-responsive proteins. The number of proteins that either (A) increased or (B) decreased by at least 1.5-fold in eleven functional categories is shown for roots exposed to 150 mM NaCl for 6h (grey bars) or 48 h (black bars). Functional categories are named either according to the major metabolic substrate, except for the categories representing processes of signal transduction, transcriptional regulation (transcription), and unclassified or unknown (unclass. & unk.) proteins.

For the proteins whose abundance decreased at 6 h following NaCl treatment,

mRNA of 40 genes (57%, 40 out of 70) also decreased and, this percentage was calculated to be 52% for the 48 h time point. However, for the 22 and 29 proteins that increased in abundance after treated by NaCl for 6 or 48 h, respectively, only 4 (18%) and 6 (21%) genes also increased at the mRNA level (Fig. 3-5). The other spots showed different expression patterns between mRNA and protein at least at one time point (Fig. 3-5). These results support the conclusion of other authors that, in statistical terms, measurements of mRNA are not always correlated with protein abundance (Gygi *et al.*, 1999; Tian *et al.*, 2004; Mooney *et al.* 2006; Noir *et al.*, 2006; Yan *et al.*, 2006). Interestingly, a similar pattern was reported in a comparison of 2-DGE and qRT-PCR data for chilling stress in rice: 88% (15 out of 17) of proteins and their corresponding transcripts decreased in parallel following stress, while only 19% (5 of 27) of proteins that increased in abundance following stress had cognate transcripts that also increased in abundance (Yan *et al.*, 2006). Together, these studies suggest that




Fig. 3-5. Comparison of NaCl-induced changes in mRNA and cognate protein abundance. The relative change in abundance (treated/untreated) is shown in \log_2 scale for proteins extracted from roots treated for (A) 6 h or (B) 48 h, as compared to previously reported (Jiang and Deyholos, 2006) changes in mRNA abundance levels in tissues grown under identical conditions.

transcript abundance may be more directly relevant to expression of genes that are down-regulated following stress than those genes that are up-regulated. However, more study is required to demonstrate the generality of this pattern across a larger sample of proteins, treatments, and measurement techniques.

Discussion

In this study, 81 differentially expressed proteins in *Arabidopsis* roots in response to salt stress treatment were identified (Table 3-1). These proteins are found to be involved in a variety of plant processes (Fig. 3-4), suggesting an overall adaptation response in roots when challenged by osmotic and ionic stresses induced by high salinity.

Energy metabolism

Under NaCl stress, plants decrease energy metabolism rates to conserve energy and limit further generation of ROS (Moller 2001). We previously reported that the transcript abundance of components of the glycolytic, citrate cycle, mitochondrial respiration, and pentose phosphate pathways (PPP) was generally decreased in NaCl-treated *Arabidopsis* roots (Jiang and Deyholos, 2006). Therefore, it was not surprising to observe in the present study that the abundance of 11 proteins involved in glycolysis, citrate circle, PPP and electron transport decreased at one or both time points after NaCl treatment (Table 3-1). For example, aconitase (At2g05710), isocitrate dehydrogenase (At1g65930), fumarase (At2g47510) and malate dehydrogenase (At1g04410) are four enzymes of the citrate cycle that each also decreased in abundance following NaCl treatment. In contrast, the abundance of nucleoside diphosphate kinase 1 (NDPK1, At4g09320), an enzyme converting GTP to ATP, and a mitochondrial malate dehydrogenase (At1g53240) increased at the early phase (6 h) and late phase (48 h) following NaCl treatment, respectively. Differential regulation of structurally related transcripts and proteins has been previously reported in NaCl treated roots, and this may reflect sub-functionalization of related enzymes for optimal activity in different cells or cellular microenvironments (Jiang and Deyholos, 2006). The phosphopyruvate hydratase/enolase (LOS2, At2g36530), which catalyzes the formation of high-energy phosphoenol pyruvate (PEP) from 2-phosphoglycerate in the

glycolytic pathway, decreased after both 6 h and 48 h of stress. Previous reports have shown that mutation in *LOS2* locus results in repression of cold-responsive genes and therefore *LOS2* acts as a positive regulator of cold-responsive genes (Lee *et al.*, 2002). The decreased abundance of *LOS2* protein that we observed indicates that NaCl-induced perturbation of metabolic flux in glycolysis may be different from that of cold stress. Other energy related proteins, including three proton transporting ATPases, two vacuolar ATP synthases (At1g78900 and At4g11150) and a mitochondrial ATP synthase delta chain (At5g47030) were found to be responsive to NaCl stimulus. Vacuolar H⁺-ATPase can generate a proton electrochemical gradient, which is the driving force utilized by the tonoplast Na⁺/H⁺ antiporter, AtNHX1 to compartment Na⁺ into vacuole (Chinnusamy *et al.*, 2005). Vacuolar sequestration of Na⁺ is an important and cost-effective strategy for osmotic adjustment that also reduces the Na⁺ concentration in the cytosol in plants.

ROS scavenging and detoxifying enzymes

Abiotic stresses induce the production of ROS, which, on one hand can cause damage to cellular components, and on the other hand, can act as signaling molecules for stress responses (Apel and Hirt, 2004). Plants can regulate the ROS level through complex mechanisms such as scavenging them by ascorbate peroxidase (APX), glutathione peroxidase (GPX), glutathione S-transferase (GST), and superoxide dismutase (SOD), of which 9 proteins were identified to be differentially expressed in this study (Table 3-1). In general, the abundance of these 9 identified proteins was increased upon NaCl treatment. It is proposed that some members of APX, GPX, GST and SOD families are part of the antioxidant system employed by plants (Apel and Hirt, 2004) and, previous microarray results demonstrated that members of them are responsive to various stresses including NaCl, osmotica, drought and cold (Kreps *et al.*, 2002; Seki *et al.*, 2002; Jiang and Deyholos, 2006). The increased abundance of GST, PRX and SOD proteins following NaCl is consistent with the presence of oxidative stress in NaCl-stressed roots. The alleviation of oxidative damage and increased resistance to environmental stresses is correlated with an efficient antioxidative system (Smirnoff 1998). Overexpression of some SOD, APX and GST genes have been shown to improve oxidative stress tolerance in transgenic plants (Allen 1995; Roxas *et al.*, 1997;

Mittler 2002).

GSTs are abundant proteins encoded by a highly divergent, ancient gene family and have protective functions such as detoxification of herbicides, and the reduction of organic hydroperoxides formed during oxidative stress. Recent studies have also implicated GSTs as components of ultraviolet-inducible cell signaling pathways and as potential regulators of apoptosis (Dixon *et al.*, 2002). Our previous microarray analysis indicated that transcripts of at least 19 GST genes increased in abundance in NaCl-treated *Arabidopsis* roots (Jiang and Deyholos, 2006). Here, we identified proteins of four plant-specific GSTs, AtGST1/ GSTF6 (At1g02930), AtGSTF2/GST4 (At4g02520), AtGST6/GSTF8 (AT2g47730) and AtGST11/GSTF7 (AT1g02920), which all ultimately increased in abundance after 48 h of NaCl treatment, although several of them were slightly decreased in abundance at the initial time point sampled (6 h). Previous studies showed that AtGST1 was up-regulated by a variety of treatments while AtGSTF2 and AtGST6 each showed a selective spectrum of inducibility to different stresses, indicating that regulation of gene expression in this super-family is controlled by multiple mechanisms (Wagner *et al.*, 2002).

APX and GPX can directly detoxify H_2O_2 to H_2O , and previous studies showed that APX1 is a central component of the reactive oxygen gene network of *Arabidopsis* (Davletova *et al.*, 2005). In *Arabidopsis*, a family of seven related proteins named AtGPX1- AtGPX7 was identified and several AtGPX genes were up-regulated coordinately in response to abiotic stresses (Milla *et al.*, 2003). AtGPX6 possibly encodes mitochondrial and cytosolic isoforms by alternative initiation and, AtGPX6 transcript showed the strongest responses under most abiotic stresses tested, thus supporting an important role for it in protection against oxidative damage (Milla *et al.*, 2003).

In this study, we observed a NaCl-dependent increase in protein abundance for two class III plant peroxidases, PER22 (At2g38380) and PER23 (At2g38390). Class III peroxidases are plant-specific oxidoreductases that are implicated in various physiological processes such as H_2O_2 detoxification, auxin catabolism, lignification, suberization, stress response (wounding, pathogen attack, NaCl) and senescence (Hiraga *et al.*, 2001; Passardi *et al.*, 2005). This agrees with our previous results that

Table 3-1. Differentially expressed proteins identified by LC-MS/MS.

spot ID	AGI#	Putative identity	MW (kDa) thr/exp	pI thr/exp	score ^a	PM ^b	C(%) ^c	NCBI Acc#	fold change 6h	48h
energy metabolism										
18*	At3g52930	fructose-bisphosphate aldolase	38.9/37.9	6.05/5.95	711	27	54	NP_190861	0.91±0.12	0.65±0.07
11	At1g65930	isocitrate dehydrogenase (NADP+)-oxidoreductase	46.1/43.0	6.13/5.93	1433	42	62	NP_176768	0.4±0.10	0.69±0.13
19†	At3g04120	glyceraldehyde-3-phosphate dehydrogenase C subunit(GAPC)	37.0/37.0	6.62/6.62	852	41	52	NP_187062	0.55±0.08	0.46±0.11
43†	At3g55440	triose phosphate isomerase(TPI)	27.4/24.2	5.24/5.39	620	16	49	2009415A	0.62±0.15	0.75±0.09
12*	At2g47510	fumarate(FUM1)/fumarate hydratase	53.5/43.3	8.01/6.79	1004	27	47	NP_182273	0.63±0.20	0.88±0.16
22	At1g04410	malate	35.9/35.9	6.11/5.92	787	55	58	NP_171936	0.55±0.14	0.9±0.21
25†	At1g53240	dehydrogenase/oxidoreductase malate dehydrogenase (NAD), mitochondrial	37.2/33.4	8.54/5.95	726	31	47	A AF69549	1.08±0.12	1.52±0.23
47	At3g27890	NADPH:quinone oxidoreductase(NQR)	21.5/22.4	6.84/6.51	308	10	30	A AD37373	0.45±0.23	1.55±0.17
33†	At5g20080	cytochrome-b5 reductase/oxidoreductase	36.1/27.6	8.76/6.72	463	16	32	NP_568391	0.92±0.16	0.58±0.14
2*	At2g05710	cytoplasmic aconitate hydratase	98.7/103.2	5.79/5.83	791	28	28	A AD25640	0.96±0.18	0.59±0.11
24	At1g22450	Cytochrome c oxidase subunit ₁ (COX6B)	21.4/33.9	4.31/5.01	83	5	20	A AM63485	0.44±0.16	NDT
8	At2g36530	phosphopyruvate hydratase, enolase(LOS2)	48.0/48.0	5.54/5.54	1038	60	64	NP_181192	0.83±0.28	0.65±0.14
73*	At1g78900	vacuolar ATP synthase(VHA-A) subunit A	69.2/73.6	5.11/5.21	1483	37	52	NP_001031299	1.24±0.19	0.43±0.19
48	At5g47030	ATP synthase delta' chain, mitochondrial	21.5/21.4	6.2/5.2	127	6	16	NP_199514	1.53±0.26	1.39±0.30
31†	At4g11150	V-type proton-ATPase(TUF)	26.3/28.8	6.04/5.86	782	34	56	CAA63086	1.48±0.22	0.62±0.11
76*	At3g03250	UDP-glucose pyrophosphorylase (UGP)	51.9/47.0	5.8/5.83	1128	29	54	NP_186975	0.49±0.19	0.66±0.15
10*	At3g02360	phosphogluconate dehydrogenase	53.9/46.4	7.02/6.91	741	22	35	NP_850502	0.68±0.14	0.57±0.21

57	At4g09320	nucleoside-diphosphate kinase(NDPK1)	16.3/14.2	7.03/6.01	309	14	49	S31444	3.43±0.77	1.01±0.35
80 [†]	At1g47260	mitochondrial gamma carbonic anhydrase	30.2/29.0	6.71/6.79	602	17	46	NP_175159	0.63±0.12	0.7±0.10
ROS scavenging and defense										
67	At1g02930	glutathione S-transferase(ATGST1)	23.5/23.8	5.8/5.84	344	14	41	CAA72413	0.68±0.14	2.15±0.81
42	At4g02520	glutathione S transferase(ATGSTF2)	24.1/24.5	5.92/5.87	672	25	59	AAC78264	1.02±0.13	1.95±0.18
41*	At4g02520	Glutathione S-Transferase(ATGSTF2)	24.0/24.3	5.93/5.98	734	32	70	1BX9_A	0.79±0.25	1.71±0.20
44*	At2g47730	glutathione S-transferase (GST6)	24.1/23.2	6.09/5.93	582	18	55	AAC63629	0.78±0.10	1.5±0.17
58*	At1g02920	glutathione S-transferase(GST11)	23.6/24.0	6.31/6.19	468	17	48	CAA74639	0.61±0.11	2.06±0.50
36 [†]	At1g07890	L-ascorbate peroxidase(APX1)	27.8/26.4	5.72/5.82	681	50	54	CAA42168	0.62±0.13	1.51±0.07
61 [†]	At1g07890	L-ascorbate peroxidase(APX1)	27.8/26.8	5.72/5.65	387	13	51	NP_172267	1.08±0.25	1.58±0.10
35 [†]	At1g07890	L-ascorbate peroxidase(APX1)	27.8/26.1	5.72/5.71	473	16	54	CAA42168	0.64±0.15	0.64±0.11
21*	At2g38390	peroxidase23(PER23)	38.6/35.4	8.33/9.19	328	18	33	NP_181373	NDT	1.89±0.45
64*	At2g38380	peroxidase22(PER22)	38.7/39.6	5.95/5.87	352	16	20	AAA32842	1.05±0.31	2.15±0.58
52	At4g11600	glutathione peroxidase 6 (ATGPX6)	18.8/20.3	6.59/5.96	239	7	62	BAA24226	0.4±0.17	2.61±1.14
15 [†]	At1g77120	alcohol dehydrogenase(ADH1)	41.9/41.9	5.83/5.80	789	29	43	CAA54911	0.57±0.20	0.6±0.12
69	At3g10920	manganese superoxide dismutase(MSD1)	25.5/22.9	8.47/5.95	470	16	58	NP_187703	1.33±0.35	3.71±1.08
60	At1g51980	metalloendopeptidase	54.6/45.7	5.94/5.47	1126	33	44	NP_175610	0.35±0.18	0.91±0.23
49	At3g56240	copper homeostasis factor	13.1/21.3	4.91/5.17	374	12	53	CAB87423	0.56±0.17	1.41±0.25
13 [†]	At5g03630	monodehydroascorbate reductase, (ATMDAR2)	47.5/43.3	5.24/5.25	1231	43	70	NP_568125	0.88±0.22	0.58±0.13
46	At4g11650	osmotin-like protein (AtOSM34)	27.5/22.2	6.26/6.86	160	6	19	AAM61750	1.16±0.20	1.91±0.28
protein translation, processing and degradation										
23*	At3g09200	ribosomal protein L10	34.2/34.6	5/5.17	411	38	42	NP_187531	0.92±0.15	0.48±0.18
27*	At3g53870	ribosomal protein S3a	27.5/32.6	9.57/9.13	786	31	56	CAB88349	0.83±0.17	0.64±0.07
55	At1g15930	ribosomal protein S12 (RPS12A)	15.7/17.2	5.38/5.11	79	2	16	NP_173045	0.52±0.14	0.29±0.22
7 [†]	At4g34110	poly(A)-binding protein	68.8/67.3	8.19/7.56	901	24	30	CAB80128	0.71±0.10	0.48±0.09
32*	At3g12390	nascent polypeptide associated	22.0/29.0	4.3/5.0	405	17	59	NP_187845	1.65±0.21	0.79±0.11

75*	At1g77510	complex alpha chain protein disulfide isomerase(ATPDIL1-2)	56.6/67.3	4.9/5.16	1204	29	54	NP_177875	NDT	NDT	
63†	At1g21750	protein disulfide isomerase(ATPDIL1-1)	55.9/64.3	4.54/5.13	1480	40	53	NP_173594	1.5±0.37	0.33±0.19	
5*	At5g09590	mHSC70-2 (Heat shock protein 70)	73.2/73.6	5.63/5.34	1301	35	40	NP_196521	0.62±0.20	0.32±0.24	
40*	At4g31300	20S proteasome beta subunit A (PBA1)	25.2/24.5	5.31/5.58	669	16	49	CAA74028	0.67±0.30	1.5±0.21	
29	At5g23540	26S proteasome subunit RPN11	34.4/32.9	6.3/6.19	544	17	41	AAP86672	0.53±0.17	0.4±0.20	
17†	At5g43060	cysteine-type endopeptidase	52.4/39.6	5.86/5.11	528	18	24	NP_568620	0.53±0.10	0.85±0.11	
38†	At1g56450	endopeptidase	27.7/24.7	6.09/7.01	522	13	52	NP_176040	2.31±0.59	0.58±0.12	
78†	At2g46280	eukaryotic translation initiation factor 3 (eIF311/TRIP-1)	36.7/35.7	6.5/6.7	800	18	53	NP_182151	0.55±0.09	0.72±0.08	
cell wall-related											
6*	At1g47600	glycosyl hydrolase family 1 protein	58.1/58.8	8.34/7.96	976	42	38	NP_175191	1.92±0.40	0.88±0.12	
77*	At1g66280	glycosyl hydrolase family 1 protein	60.2/57.5	6.74/6.77	1177	35	37	NP_176802	0.56±0.10	0.51±0.13	
86*	At3g09260	glycosyl hydrolase family 1 protein	60.4/62.9	6.95/6.69	925	23	39	AAB38783	0.6±0.09	1.59±0.17	
65*	At3g09260	glycosyl hydrolase family 1 protein	60.3/62.9	6.95/6.56	629	23	24	AAB38783	0.54±0.11	3.04±1.01	
62	At4g16260	glycosyl hydrolase family 17 protein	37.7/32.4	6.43/8.33	100	4	22	AAL36038	NDC	1.63±0.12	
26	At4g16260	glycosyl hydrolases family 17 protein	37.7/32.6	6.43/6.95	793	54	67	AAL36038	NDC	1.59±0.23	
56*	At2g21660	glycine-rich RNA binding protein 7(GRP7)	16.9/16.0	5.85/5.44	642	27	76	AAM62447	1.51±0.07	0.85±0.09	
45	At4g14630	germin-like protein(GLP9)	23.2/23.2	5.82/5.84	113	8	11	AAD00509	0.92±0.10	1.85±0.21	
39	At5g38940	oxalate oxidase (germin protein)- like protein	23.8/24.2	8.62/7.76	146	13	13	BAB08650	1.83±0.19	1.58±0.15	
66*	At3g02230	reversibly glycosylated polypeptide (RGP1)	41.1/36.5	5.61/5.54	996	31	66	NP_186872	1.24±0.12	1.87±0.22	
hormone-related											
81†	At1g62380	ACC oxidase(ACO2)	36.4/38.3	4.97/5.1	490	12	31	AAC27484	1.31±0.21	0.66±0.09	
59†	At1g02500	S-adenosylmethionine synthetase (SAM1)	43.6/45.1	5.5/5.52	740	25	41	AAA32868	0.67±0.12	0.51±0.11	
85†	At3g25780	allene oxide cyclase 2 (AOC2)	28.5/22.4	9.19/7.6	141	6	27	NP_566777	NDC	NDC	

68†	At3g16470	jasmonate inducible protein(JR1)	48.6/45.1	5.12/5.24	898	27	32	BAB01146	0.83±0.13	1.66±0.15
signal transduction										
83	At1g56340	calreticulin 1 (CRT1)	48.7/61.5	4.46/5.04	807	20	40	NP_176030	0.57±0.10	1.53±0.10
84*	At1g09210	calreticulin 2 (CRT2)	48.4/53.7	4.37/5.02	956	38	43	NP_172392	0.61±0.17	1.62±0.19
28	At1g62480	vacuolar calcium-binding protein-related	16.6/32.6	4.05/4.95	57	2	30	NP_564795	0.79±0.20	NDT
34	At5g20010	small Ras-like GTP-binding protein (Ran-1)	25.6/27.2	6.38/6.45	237	7	35	AAA32851	0.43±0.15	2.69±1.10
amino acid metabolism										
16*	At5g07440	Glu dehydrogenase 2 (GDH2)	45.0/41.4	6.07/5.96	915	32	46	NP_196361	0.63±0.14	1.61±0.18
20*	At1g66200	Gln synthetase(GS)	47.5/39.6	5.97/5.11	499	14	36	1804333C	0.53±0.10	1.53±0.12
71*	At5g14200	3-isopropylmalate dehydrogenase (AtIMD1)	44.3/42.1	5.75/5.35	1051	24	47	AAU90074	1.02±0.35	0.57±0.07
3*	At5g17920	cobalamine-independent Met synthase	84.3/84.3	6.02/6.03	1319	41	35	IUIH_A	0.72±0.10	0.36±0.24
cytoskeleton										
14	At1g49240	actin 8	42.1/42.7	5.37/5.4	670	17	36	AAC49523	0.38±0.26	0.75±0.16
70*	At4g14960	tubulin alpha-6 chain	50.2/47.6	4.93/5.13	682	17	35	CAB10275	1.74±0.21	1.44±0.19
74*	At5g62690	putative tubulin beta-2/beta-3 chain	51.4/48.0	4.7/5.05	1064	39	52	BAC42096	0.54±0.11	0.63±0.04
transcription										
50	At1g73230	(NAC) domain-containing protein	18.0/20.8	5.91/5.66	193	3	30	AAM61406	1.68±0.23	1.89±0.31
53	At1g17880	(NAC) domain-containing protein	17.9/18.9	6.62/6.25	402	12	63	NP_173230	1.05±0.19	1.76±0.20
other metabolism										
79	At1g53580	hydroxyacylglutathione hydrolase, putative	27.0/28.8	5.45/5.26	348	11	34	2GCU_D	0.65±0.10	0.53±0.13
4	At3g60750	transketolase-like protein	81.9/84.3	5.8/5.46	698	20	35	CAB82679	0.83±0.12	0.54±0.11
30†	At5g09650	inorganic pyrophosphatase (AtPPA6)	33.7/31.4	5.71/5.17	592	15	33	AAS57950	0.65±0.10	0.61±0.05
unclassified and unknown										
37†	At2g43090	aconitase C-terminal domain-containing protein	27.1/27.6	6.33/5.19	558	15	54	NP_181837	1.31±0.19	2.46±0.71
82	At4g23670	major latex protein-related	17.6/15.4	5.91/5.86	277	17	42	NP_194098	1.55±0.10	NDC
1	At3g15950	unknown protein	85.2/115.0	4.61/5.09	1094	31	32	NP_188216	0.73±0.12	0.23±0.15
9*	At1g03220	unknown protein	46.4/45.4	8.97/8.79	837	42	47	NP_171821	0.57±0.17	0.59±0.10

51*	At3g52300	putative protein	19.6/20.1	5.09/5.07	888	23	67	CAC07921	1.25±0.18	0.61±0.12
54	At5g10860	unknown protein	22.8/18.2	9.1/7.01	396	9	37	NP_196647	0.75±0.16	1.52±0.11
72†	At2g20360	hypothetical protein	44.0/35.9	9.26/8.45	920	25	42	AA.T68351	1.77±0.22	0.81±0.14

The Arabidopsis Genomics Initiative (AGI) locus name retrieved from TAIR (www.arabidopsis.org) by running Blast tool is shown for each

protein. Where peptides from a single spot matched proteins from more than one AGI locus (score^a of >100), only the highest scoring match is shown, and the symbol (*) or (†) is appended to the spot ID for multiple matches within the same (*) or different (†) gene families. Fold change is expressed as a ratio of the Vol% between 150 mM NaCl treated/control roots, and each value represents the mean value ± S.D. of three biologically independent measurements. For some spots, fold-change cannot be accurately calculated because of complete absence of the spot in either treated or control samples; this is noted by the abbreviation NDC (not detected in control) or NDT (not detected in treatment).^a Probability based molecular weight search (Mowse) score; ^b Number of peptides matched; ^c sequence coverage percentage. * multiple protein matches in the same gene family as the best match shown. † multiple protein matches in different gene families from the best match shown.

the majority of class III peroxidases are responsive, at the transcript level, to NaCl treatment in *Arabidopsis* roots (Jiang and Deyholos, 2006). The diverse functions of class III peroxidases are in part due to two possible catalytic cycles, peroxidative and hydroxylic, involving the consumption or release of H₂O₂ and ROS (Passardi *et al.*, 2005). Although some functions of these peroxidases appear to be paradoxical, the whole process is probably regulated by a fine-tuning that has yet to be elucidated (Passardi *et al.*, 2005).

SODs catalyze the dismutation of superoxide into oxygen and H₂O₂ and constitute the first line of defence against ROS within a cell (Alscher *et al.*, 2002). Surprisingly, in our previous microarray analyses, transcripts for all three detectable SOD genes decreased in response to NaCl treatment (Jiang and Deyholos, 2006). Among these was manganese superoxide dismutase, MSD1 (At3g10920), which protein we here observed to increase in abundance by 3.7 fold at 48 h post NaCl-treatment (Table 3-1). The contrast in protein and transcript abundance in similarly treated tissues for MSD1 highlights the importance of an integrated proteomic and transcriptomic analysis of gene expression.

Ascorbate (AsA) is a major antioxidant and free-radical scavenger in plants. Monodehydroascorbate reductase (MDAR) is crucial for AsA regeneration and essential for maintaining a reduced pool of AsA. Surprisingly, we found that the abundance of one monodehydroascorbate reductase (AtMDAR2, At5g03630) was down-regulated by NaCl stress, suggesting that although plants require reduced AsA to remove free radicals, the fine tuning of the levels of various antioxidants is also an important consideration in stress responses (Lisenbee *et al.*, 2005).

Protein translation, processing and degradation

Regulation of gene expression is achieved at several levels, i.e. transcriptional, post-transcriptional, translational and post-translational. 13 proteins implicated in protein translation, processing and degradation were identified in our present study. A decrease in bulk *de novo* protein synthesis following NaCl treatment has been detected in *Arabidopsis* (Ndimba *et al.*, 2005), and our previous microarray data also demonstrated down-regulation of the majority of transcripts for almost all cytosolic and plastidic ribosomal proteins (Jiang and Deyholos, 2006). Similarly, microarray profiling of

Arabidopsis seedlings under hypoxia indicated a repression of bulk protein synthesis followed by selective translation of specific transcripts (Branco-Price *et al.*, 2005). Under dehydration condition, more than 90% of the Arabidopsis mRNAs showing a strong decrease in abundance as detected by microarray displayed reduced polysomal association, indicating a decreased translation of those transcripts (Kawaguchi *et al.*, 2004).

Consistent with these observations, we identified 3 ribosomal proteins (At1g15930, At3g09200 and At3g53870), whose abundance decreased following NaCl treatment, suggesting that short term NaCl stress represses protein synthesis *in vivo* (Table 3-1). The expression level of a eukaryotic translation initiation factor 3 subunit protein (eIF3I1, At2g46280), which is a homolog of mammalian TGF-beta receptor interacting protein (TRIP), was also found to be decreased in this study. Previous studies showed that eIF3I1/TRIP-1 is involved in brassinosteroid regulated plant growth and development, thereby revealing a putative link between a developmental signaling pathway and the control of protein translation (Jiang and Clouse 2001). We detected several proteins that promote the proper folding of proteins and/or prevent the aggregation of nascent or damaged proteins. Two protein disulfide isomerase-like (PDIL) proteins (PDIL1-1, At1g21750; PDIL1-2, At1g77510), putative nascent polypeptide associated complex alpha chain protein (At3g12390) and mitochondrial HSC70-2 (70-kDa heat-shock cognate, At5g09590) all ultimately decreased in protein abundance in NaCl treated tissues as compared to untreated controls after 48 h, although At3g12390 showed transient increases in abundance after 6 h of NaCl treatment. Members of HSC70 proteins are often involved in assisting the folding of *de novo* synthesized polypeptides and the import/translocation of precursor proteins (Wang *et al.*, 2004). It was proposed that HSC70 might be used as a motor for transporting the precursor protein through the membranes by interacting with the signal peptides (Zhang and Glaser, 2002). The decreased abundance of MtHSC70-2 after NaCl treatment suggests a decrease in the transportation of newly synthesized peptides into mitochondria, due partly to a decreased *de novo* protein synthesis under saline conditions.

Cell wall-related proteins

Moderate NaCl stress reduces water availability and leads to the inhibition of plant growth by increasing the threshold pressure for wall yielding in expanding cells or inducing hydraulic limitations to water uptake (Neumann *et al.*, 1994; Steudle 2000). In our current study, four glycosyl hydrolase (GH) family proteins, 3 of which belong to GH family 1 (At1g47600, At1g66280, At3g09260) and one GH family 17 (At4g16260) were identified (Table 3-1). GH1 and GH17 protein families include β -glucosidases and β -1,3-glucanases, respectively, which play important roles in many physiological processes in plants including cell wall remodeling (Bray 2004; Xu *et al.*, 2004). Each of the four GH proteins we identified had a distinctive temporal expression pattern: two of the GH1s decreased in abundance after 6 h, while the third GH1, At1g47600 increased at 6 h; however, after 48 h, At3g09260 (spots 65 and 86) showed increased abundance than controls. The identified two GH17 protein spots (spots 26 and 62) were more abundant in treated tissues than controls at both time points. The diversity of expression patterns suggested that several different physiological processes were represented by these results. Further experiments on the substrate specificity, localization of the enzymes with respect to potential substrates, and the activities of the substrates and hydrolysis products are required to determine the roles of these enzymes in NaCl root responses.

Glycine-rich proteins (GRPs) containing more than 60% glycine have been found in the cell walls of many higher plants and form a group of structural protein components of the wall in addition to extensins and proline-rich proteins (Ringli *et al.*, 2001). GRPs play roles in post-transcriptional regulation of gene expression in plants under various stress conditions, and in most cases, they are accumulated in the vascular tissues and that their synthesis is part of the plant's defense mechanism (Mousavi and Hotta, 2005). We observed that salinity caused a transient increase in the abundance of AtGRP7 (At2g21660) (Table 3-1). Previous studies showed that AtGRP7 transcript level was repressed by ABA, high NaCl and mannitol (Cao *et al.*, 2006). More recent studies suggest that AtGRP7 exhibits RNA chaperone activity and can promote the cold adaptation process in *E. coli* (Kim *et al.*, 2007).

We also identified two germin-like proteins (GLPs)- GLP9 (At4g14630) and

oxalate oxidase-like protein (At5g38940), whose abundance increased in Arabidopsis roots subjected to NaCl treatment (Table 3-1). GLPs exhibit sequence and structural similarity to cereal germins and may be associated with the cell wall (Membre *et al.*, 2000). Although GLPs mostly lack oxalate oxidase activity, some GLPs have SOD activity. GLPs are thought to play a significant role both during embryogenesis and in biotic and abiotic stress conditions. For example, GLP expression was detected in barley roots after exposure to NaCl (Hurkman *et al.*, 1994). A reversibly glycosylated polypeptide (RGP1, At3g02230) was also found to be induced by NaCl treatment. RGP1 is possibly involved in plant cell wall synthesis (Dhugga *et al.*, 1997). Cell wall rigidification, the formation of a physical barrier and a process of class III peroxidase-mediated cross-linking of several compounds (Passardi *et al.*, 2004), would protect plant roots from further dehydration under water deficit.

Hormone-related proteins

Ethylene and jasmonic acid (JA) are hormones whose activity has been previously correlated with environmental stress (Chen *et al.*, 2005; Devoto and Turner, 2005). The majority of ethylene and JA-related transcripts detected in our previous microarray experiments were responsive to NaCl treatment (Jiang and Deyholos, 2006). ACC oxidase (ACO) catalyzes the conversion of 1-aminocyclopropane-1-carboxylic acid (ACC) to ethylene. ACO are encoded by a small gene family. We found that ACO2 (At1g62380) increased in abundance at the early (6 h) time-point, later decreased in abundance (Table 3-1). S-adenosylmethionine synthetase catalyzes the production of S-adenosyl-L-methionine (SAM) from L-methionine and ATP. SAM serves as a methyl group donor in numerous transmethylation reactions and is the precursor for the biosynthesis of polyamines and ethylene among other metabolites. In plants, SAM synthetases are encoded by small gene families that contain members that are differentially regulated by NaCl stress (Espartero *et al.*, 1994). Here SAM1 (At1g02500) decreased in abundance following NaCl treatment, which is consistent with our previously reported microarray results (Jiang and Deyholos, 2006).

JA is involved in a wide range of stress, defence, and developmental processes (Devoto and Turner, 2005). We identified one enzyme implicated in JA biosynthesis, allene oxide cyclase2 (AOC2, At3g25780), whose abundance increased upon NaCl

challenge, indicating that increased JA biosynthesis may also be associated with NaCl-responses in Arabidopsis roots. One jacalin lectin family protein (JR1, At3g16470), which is similar to myrosinase binding protein was found to be positively regulated by NaCl. JR1 transcript was strongly induced by wounding and JA (Leon *et al.*, 1998), supporting the concept of cross-talk between various abiotic stresses.

Signal transduction network involved in NaCl stress responses

An increase in NaCl concentration in the extracellular space can be perceived by putative sensors in the cell membrane of Arabidopsis and transmitted to cellular machinery to regulate gene expression (Chinnusamy *et al.*, 2005). Some proteins involved in signal transduction were identified in this study (Table 3-1), i.e. two calcium ion binding proteins (CRT1, At1g56340 and CRT2, At1g09210), a vacuolar calcium-binding protein-related (At1g62480) and a small Ras-like GTP-binding protein (Ran-1, At5g20010). In plant cells, Ca²⁺ is a ubiquitous intracellular second messenger involved in numerous signaling pathways. Modulation of intracellular Ca²⁺ levels is partly regulated by calcium-binding proteins, which, after activated, induce specific kinases. Calreticulin (CRT) is a multifunctional protein mainly localized to the endoplasmic reticulum in eukaryotic cells. Plants have three CRT isoform groups (CRT1/CRT2 and CRT3) and Arabidopsis has 18 CRT proteins and members of the different isoform groups respond differently to applied external stimuli (Persson *et al.*, 2003). The identified CRT1 and CRT2 in this study support that they are the major isoforms, possibly due to an enhanced Ca²⁺-binding efficiency, and play important roles in Ca²⁺ homeostasis under osmotic stress. Ran is an evolutionarily conserved eukaryotic GTPase, which is likely involved in nuclear translocation of proteins and cell cycle progression (Yang 2002). However, little is known about the function of Ran in plant response to stresses. We found that the abundance of Ran-1 increased after 48 h NaCl treatment, suggesting that Ran could also play a specific role under saline condition.

Amino acid metabolism

The amount of Pro and certain other amino acids is reported to increase under NaCl treatment (Fougère *et al.* 1991; Di Martino *et al.* 2003). We observed that the abundance of four amino acid biosynthesis-related enzymes was influenced by NaCl

(Table 3-1). 3-isopropylmalate dehydrogenase (AtIMD1, At5g14200), which is involved in Leu biosynthesis, and cobalamine-independent methionine synthase (ATCIMS/AtMetE, At5g17920) decreased in abundance following NaCl treatment. However, glutamate dehydrogenase 2 (GDH2, At5g07440), and glutamine synthetase (GS, At1g66200) both decreased in abundance at the 6 h time-point, but increased at the 48 h time-point. GS functions as the major assimilatory enzyme for ammonia and GDH works as a link between carbon and nitrogen metabolism as it can aminate 2-oxoglutarate into glutamate (biosynthetic reaction) or deaminate glutamate into ammonium and 2-oxoglutarate (catabolic reaction). GS and GDH, together with a number of other enzymes play key roles in maintaining the balance of carbon and nitrogen (Mifflin and Habash 2002). Recent study showed that salinity-generated ROS signal induces α -GDH subunit expression, and the anionic iso-GDHs assimilate ammonia, acting as anti-stress enzymes in ammonia detoxification and production of Glu for Pro synthesis (Skopelitis *et al.*, 2006).

Cytoskeleton

Actin and tubulin dynamics have important functions in cellular homeostasis. The cytoskeleton is rapidly remodeled by various endogenous and external stimuli such as hormones, low temperature, aluminum, and NaCl (Abdrakhamanova *et al.*, 2003; Dhonukshe *et al.* 2003; Sivaguru *et al.*, 2003). For example, the transverse orientation of cortical microtubule arrays in tobacco BY-2 cells was remodeled to a more random arrangement after treatment with 150 mM NaCl for 15 min (Dhonukshe *et al.* 2003). More recent research suggests that NaCl stress compromises the organization of cortical microtubule arrays, in which SPR1 is involved, and inhibits anisotropic growth (Shoji *et al.*, 2006). We found that one actin protein, ACT8 (At1g49240) and one tubulin β -chain (At5g62690) decreased in abundance following NaCl treatment, while tubulin α -6 chain (TUA6) was induced by NaCl (Table 3-1). These observations are consistent with previously reported microarray results (Seki *et al.*, 2002; Jiang and Deyholos, 2006), and although their mechanistic significance is not fully clear, the stress-responsiveness of these common, cytoskeletal proteins calls into question their sometimes designation as house-keeping genes.

Transcription-related proteins

Transcriptional control of the expression of stress-responsive genes is a crucial part of the plant response to various abiotic and biotic stresses. Nascent-polypeptide-associated complex (NAC) is a heterodimeric complex (α - and β -NAC) that can reversibly bind to eukaryotic ribosomes. Rospert *et al.* (2002) suggested that NAC is a negative regulator of translocation into the endoplasmic reticulum and a positive regulator of translocation into the mitochondria. Previous studies found that α chain of NAC in osteoblasts functions as a transcriptional coactivator (Yotov *et al.*, 1998). Here we identified two NAC domain-containing proteins (At1g17880 and At1g73230), which are similar to human transcription factor BTF3 (RNA polymerase B transcription factor 3) and whose abundance increased following NaCl treatment (Table 3-1). Interestingly, transcripts for these proteins significantly decreased in abundance in our previous microarray study (Jiang and Deyholos, 2006). Furthermore, proteomic evidence showed that a rice α -NAC was down-regulated by NaCl and cold stresses (Yan *et al.*, 2005 & 2006). Based on previous microarray results produced by ourselves and others (Seki *et al.*, 2002; Jiang and Deyholos, 2006), it is likely that hundreds of other transcription factors changed in abundance in the tissues we examined, but that these proteins fell below the sensitivity threshold of analysis, and were therefore not detected in our protein gels.

Conclusion

In this study, we performed a proteomic analysis of Arabidopsis roots subjected to non-lethal NaCl treatment for 6 or 48 h, with physiologically defined responses (Fig. 3-1). Symptoms of stress, such as electrolyte leakage and Na concentration continued to increase until at least 48 h, even though considerable remodeling of the proteome had apparently occurred before this time point (Figs. 3-1& 3-2). We identified 81 different NaCl-responsive proteins by LC-MS/MS (Table 3-1). The identified proteins were implicated in a wide range of physiological processes, *i.e.* energy metabolism; ROS scavenging and detoxification; protein translation, processing and degradation; signal transduction; hormone and amine acid metabolisms; cell-wall modifications; as well as cytoskeleton remodeling, which might work cooperatively to re-establish cellular homeostasis under water deficient and ionic toxicity. Some of the proteins identified

here were also identified at the transcript level in previous microarray profiling of *Arabidopsis* response to NaCl stress (Seki *et al.*, 2002; Kreps *et al.*, 2002; Jiang and Deyholos, 2006) as well as in 2-DGE analysis of *Arabidopsis* cell suspension cultures (Ndimba *et al.*, 2005). The proteins identified in this study represent only a small part of *Arabidopsis* proteome responsive to NaCl treatment and many other NaCl-responsive proteins still need to be identified. Considering the limitations of proteomic study based on 2-D gel, i.e. inability to resolve membrane proteins and detect low-abundant proteins, complementary strategies at the transcript, protein and metabolite levels should be used to gain more insight into the intricate network of plant response to high salinity. To address some of these limitations with 2-DGE, alternative gel-free separation methods have been developed. One of them is the direct identification of proteins from complex protein mixtures based on peptide separation via liquid chromatography columns, followed by tandem mass spectrometry. This method has been applied in rice (Wienkoop and Weckwerth, 2006) and *Medicago truncatula* (Larrainzar *et al.*, 2007). Alternatively, sub-proteomics study (Lee *et al.*, 2004; Nohzadeh Malakshah *et al.*, 2007) and metabolic profiling (Kim *et al.*, 2007) have also provided new insights into stress responses. The identification of novel NaCl-responsive proteins provides not only new insights into NaCl stress responses but also a good starting point for further dissection of their functions using genetic and other approaches.

References

- Abdrakhamanova A, Wang QY, Khokhlova L and Nick P.** (2003) Is microtubule disassembly a trigger for cold acclimation? *Plant Cell Physiol* 44: 676–686
- Allen RD.** (1995) Dissection of oxidative stress tolerance using transgenic plants. *Plant Physiol* 107:1049-1054
- Amme S, Matros A, Schlesier B and Mock HP.** (2006) Proteome analysis of cold stress response in *Arabidopsis thaliana* using DIGE-technology. *J Exp Bot* 57:1537-1546
- Alscher RG, Erturk N and Heath LS.** (2002) Role of superoxide dismutases (SODs) in controlling oxidative stress in plants. *J Exp Bot* 53:1331-1341

- Apel K and Hirt H.** (2004) Reactive oxygen species: metabolism, oxidative stress, and signal transduction. *Annu Rev Plant Biol* 55: 373–399
- Atkin RK, Barton GE and Robinson DK.** (1973) Effect of root-growing temperature on growth-substances in xylem exudate of *Zea mays*. *J Exp Bot* 24: 475-487
- Bae MS, Cho EJ, Choi EY, and Park OK.** (2003) Analysis of the *Arabidopsis* nuclear proteome and its response to cold stress. *Plant J* 36:652-663
- Branco-Price C, Kawaguchi R, Ferriera R and Bailey-Serres J.** (2005) Genome-wide analysis of transcript abundance and translation in *Arabidopsis* seedlings subjected to oxygen deprivation. *Ann Bot* 96: 647-660.
- Bray EA.** (2004) Genes commonly regulated by water-deficit stress in *Arabidopsis thaliana*. *J Exp Bot* 55: 2331-2341
- Candiano G, Bruschi M, Musante L, Santucci L, Ghiggeri GM, Carnemolla B, Orecchia P, Zardi L and Righetti PG.** (2004) Blue silver: a very sensitive colloidal Comassie G-250 staining for proteome analysis. *Electrophoresis* 25: 1327-1333
- Cao S, Jiang L, Song S, Jing R and Xu G.** (2006) AtGRP7 is involved in the regulation of abscisic acid and stress responses in *Arabidopsis*. *Cell Mol Biol Lett* 11: 526-535
- Cao WH, Liu J, He XJ, Mu RL, Zhou HL, Chen SY and Zhang JS.** (2007) Modulation of ethylene responses affects plant salt-stress responses. *Plant Physiol* 143:707-719
- Casati P, Zhang X, Burlingame AL and Walbot V.** (2005) Analysis of leaf proteome after UV-B irradiation in maize lines differing in sensitivity. *Mol Cell Proteomics* 4:1673-1685
- Chao DY, Luo YH, Shi M, Luo D and Lin HX.** (2005). Salt-responsive genes in rice revealed by cDNA microarray analysis. *Cell Res* 15: 796-810
- Chen YF, Etheridge N and Schaller GE.** (2005) Ethylene signal transduction. *Ann Bot* 95: 901-915
- Chinnusamy V, Jagendorf A and Zhu J-K.** (2005) Understanding and improving salt tolerance in plants. *Crop Sci* 45: 437-448
- Cui S, Huang F, Wang J, Ma X, Cheng Y, and Liu J.** (2005) A proteomic analysis of cold stress responses in rice seedlings. *Proteomics* 5:3162-3172

- Davletova S, Rizhsky L, Liang H, Zhong S, Oliver DJ, Coutu J, Shulaev V, Schlauch K and Mittler R.** (2005) Cytosolic ascorbate peroxidase 1 is a central component of the reactive oxygen gene network of Arabidopsis. *Plant Cell* 17: 268-281
- Desimone M, Henke A and Wagner E.** (1996) Oxidative stress induces partial degradation of the large subunit of ribulose-1,5-bisphosphate carboxylase/oxygenase in isolated chloroplasts of barley. *Plant Physiol* 111: 789-796
- Devoto A and Turner JG.** (2005) Jasmonate-regulated Arabidopsis stress signalling network. *Physiol Plant* 123:161-172
- Dhonukshe P, Laxalt AM, Goedhart J, Gadella TWJ and Munnik T.** (2003) Phospholipase D activation correlates with microtubule reorganization in living plant cells. *Plant Cell* 15: 2666–2679
- Dhugga KS, Tiwari SC and Ray PM.** (1997) A reversibly glycosylated polypeptide (RGPI) possibly involved in plant cell wall synthesis: purification, gene cloning, and trans-Golgi localization. *Proc Nat Acad Sci USA* 94: 7679-7684
- Di Martino C, DelWne S, Pizzuto R, Loreto F and Fuggi A.** (2003) Free amino acids and glycine betaine in leaf osmoregulation of spinach responding to increasing salt stress. *New Phytol* 158: 455–463
- Dixon DP, Laphorn A and Edwards R.** (2002) Plant glutathione transferases. *Genome Biol* 3:REVIEWS3004
- Espartero J, Pintor-Toro JA and Pardo JM.** (1994) Differential accumulation of S-denosyl-methionine synthetase transcripts in response to salt stress. *Plant Mol Biol* 25:217-227
- Fougère F, Lerudulier D and Streeter JG.** (1991) Effects of salt stress on amino acid, organic acid, and carbohydrate composition of roots, bacteroids, and cytosol of alfalfa (*Medicago sativa*L). *Plant Physiol* 96:1228–1236
- Gygi SP, Rochon Y, Franza BR and Aebersold R.** (1999) Correlation between protein and mRNA abundance in yeast. *Mol Cell Biol* 19:1720-1730
- Hajheidari M, Abdollahian-Noghabi M, Askari H, Heidari M, Sadeghian SY, Ober ES, Salekdeh GH.** (2005) Proteome analysis of sugar beet leaves under drought stress. *Proteomics*. 5:950-960
- Hiraga S, Sasaki K, Ito H, Ohashi Y and Matsui H.** (2001) A large family of class

III plant peroxidases. *Plant Cell Physiol* 42: 462-468

Holmes-Davis R, Tanaka CK, Vensel WH, Hurkman WJ and McCormick S. (2005) Proteome mapping of mature pollen of *Arabidopsis thaliana*. *Proteomics* 5: 4864-4884

Hunt SMN, Thomas MR, Sebastian LT, Pedersen SK, Harcourt RL, Sloane AJ and Wilkins MR. (2005) Optimal replication and the importance of experimental design for gel-based quantitative proteomics. *J Proteome Res* 4, 809-819

Hurkman WJ, Lane BG and Tanaka CK. (1994) Nucleotide sequence of a transcript encoding a germin-like protein that is present in salt-stressed barley (*Hordeum vulgare* L.) roots. *Plant Physiol* 104: 803-804

Ishikawa T, Yoshimura K, Tamoi M, Takeda T and Shigeoka S. (1997) Alternative mRNA splicing of 3'-terminal exons generates ascorbate peroxidase isoenzymes in spinach (*Spinacia oleracea*) chloroplasts. *Biochem J* 328(Pt 3):795-800

Jensen ON, Wilm M, Shevchenko A and Mann M. (1999) Sample preparation methods for mass spectrometric mapping directly from 2-DE gels. In: Link AJ, ed. *2-D Proteome Analysis Protocols*. Humana Press, 112, 513-530

Jensen ON. (2004) Modification-specific proteomics: characterization of post-translational modifications by mass spectrometry. *Cur Opin Chem Biol* 8: 33-41

Jiang J and Clouse SD. (2001) Expression of a plant gene with sequence similarity to animal TGF-beta receptor interacting protein is regulated by brassinosteroids and required for normal plant development. *Plant J* 26: 35-45

Jiang Y-Q and Deyholos MK.(2006) Comprehensive transcriptional profiling of NaCl-stressed *Arabidopsis* roots reveals novel classes of responsive genes. *BMC Plant Biol* 6, 25

Kav NNV, Srivastava S, Goonewardene L and Blade SF. (2004) Proteome-level changes in the roots of *Pisum sativum* in response to salinity. *Ann Appl Biol* 145, 217-230

Kawaguchi R, Girke T, Bray EA and Bailey-Serres J. (2004) Differential mRNA translation contributes to gene regulation under non-stress and dehydration stress conditions in *Arabidopsis thaliana*. *Plant J* 38: 823-839

Kawasaki S, Borchert C, Deyholos MK, Wang H, Brazille S, Kawai K, Galbraith

- D and Bohnert HJ.** (2001) Gene expression profiles during the initial phase of salt stress in rice (*Oryza sativa* L.). *Plant Cell* 13: 889-905
- Kim JK, Bamba T, Harada K, Fukusaki E and Kobayashi A.** (2007) Time-course metabolic profiling in *Arabidopsis thaliana* cell cultures after salt stress treatment. *J Exp Bot* 58: 415-424
- Kim JS, Park SJ, Kwak KJ, Kim YO, Kim JY, Song J, Jang B, Jung CH and Kang H.** (2007) Cold shock domain proteins and glycine-rich RNA-binding proteins from *Arabidopsis thaliana* can promote the cold adaptation process in *Escherichia coli*. *Nucleic Acids Res* 35:506-516
- Kingston-Smith AH and Foyer CH.** (2000) Bundle sheath proteins are more sensitive to oxidative damage than those of the mesophyll in maize leaves exposed to paraquat or low temperatures. *J Exp Bot* 51: 123-130
- Kreps JA, Wu Y, Chang H, Zhu T, Wang X and Harper JF.** (2002) Transcriptome changes for *Arabidopsis* in response to salt, osmotic, and cold stress. *Plant Physiol* 130, 2129-2141
- Larrainzar E, Wienkoop S, Weckwerth W, Ladrera R, Arrese-Igor C, González EM.** (2007) *Medicago truncatula* root nodule proteome analysis reveals differential plant and bacteroid responses to drought stress. *Plant Physiol* 144:1495-1507
- Lee DG, Ahsan N, Lee SH, Kang KY, Bahk JD, Lee IJ, and Lee BH.** (2007) A proteomic approach in analyzing heat-responsive proteins in rice leaves. *Proteomics* 7:3369-3383
- Lee H, Guo Y, Ohta M, Xiong L, Stevenson B and Zhu JK.** (2002) LOS2, a genetic locus required for cold-responsive gene transcription encodes a bi-functional enolase. *EMBO J* 21:2692-2702
- Lee S, Lee EJ, Yang EJ, Lee JE, Park AR, Song WH and Park OK.** (2004) Proteomic identification of annexins, calcium-dependent membrane binding proteins that mediate osmotic stress and abscisic acid signal transduction in *Arabidopsis*. *Plant Cell* 16:1378-1391
- Leon J, Rojo E, Titarenko E and Sanchez-Serrano J J.** (1998) Jasmonic acid-dependent and -independent wound signal transduction pathways are differentially regulated by Ca²⁺/calmodulin in *Arabidopsis thaliana*. *Mol Gen Genet* 258: 412-419

- Lisenbee CS, Lingard MJ and Trelease RN.** (2005) Arabidopsis peroxisomes possess functionally redundant membrane and matrix isoforms of monodehydroascorbate reductase. *Planta* 43: 900-914
- Ma SS, Gong QQ and Bohnert HJ.** (2006) Dissecting salt stress pathways. *J Exp Bot* 57:1097-1107
- Majoul T, Chahed K, Zamiti E, Ouelhazi L and Ghrir R.** (2000) Analysis by two-dimensional electrophoresis of the effect of salt stress on the polypeptide patterns in roots of a salt-tolerant and a salt-sensitive cultivar of wheat. *Electrophoresis* 21: 2562-2565
- Majoul T, Bancel E, Triboï E, Ben Hamida J, and Branlard G.**(2004) Proteomic analysis of the effect of heat stress on hexaploid wheat grain: characterization of heat-responsive proteins from non-prolamins fraction. *Proteomics* 4:505-513
- Membre N, Bernier F, Staiger D and Berna A.** (2000) *Arabidopsis thaliana* germin-like proteins: common and specific features point to a variety of functions. *Planta* 211: 345-354
- Miflin BJ and Habash DZ.** (2002) The role of glutamine synthetase and glutamate dehydrogenase in nitrogen assimilation and possibilities for improvement in the nitrogen utilization of crops. *J Exp Bot* 53: 979-987
- Milla M A, Maurer A, Huete A R and Gustafson J P.** (2003) Glutathione peroxidase genes in Arabidopsis are ubiquitous and regulated by abiotic stresses through diverse signaling pathways. *Plant J* 36: 602-615
- Mittler R.** (2002) Oxidative stress, antioxidants and stress tolerance. *Trends Plant Sci* 7: 405-410
- Moller IM.** (2001) Plant mitochondria and oxidative stress: Electron transport, NADPH turnover, and metabolism of reactive oxygen species. *Annu Rev Plant Physiol Plant Mol Biol* 52:561-591
- Mooney BP, Miernyk JA, Greenlief CM and Thelen JJ.** (2006) Using quantitative proteomics of Arabidopsis roots and leaves to predict metabolic activity. *Physiol Plant* 128: 237-250
- Mousavi A and Hotta Y.** (2005) Glycine-rich proteins: a class of novel proteins. *Appl Biochem Biotech* 120:169-174

- Munns R.** (2005) Genes and salt tolerance: bringing them together. *New Phytol* **167**: 645-663
- Nohzadeh Malakshah S, Habibi Rezaei M, Heidari M and Hosseini Salekdeh G.** (2007) Proteomics reveals new salt responsive proteins associated with rice plasma membrane. *Biosci Biotechnol Biochem.* **71**:2144-2154
- Ndimba BK, Chivasa S, Simon WJ and Slabas AR.** (2005) Identification of *Arabidopsis* salt and osmotic stress responsive proteins using two-dimensional difference gel electrophoresis and mass spectrometry. *Proteomics* **5**: 4185-4196
- Neumann PM, Azaizeh H and Leon D.** (1994) Hardening of root cell walls: a growth inhibitory response to salinity stress. *Plant Cell Environ* **17**: 303–309
- Noir S, Brautigam A, Colby T, Schmidt J and Panstruga R.** (2005) A reference map of the *Arabidopsis thaliana* mature pollen proteome. *Biochem Biophys Res Comm* **337**:1257–1266
- Pandey A, Chakraborty S, Datta A, and Chakraborty N.** (2007) Proteomics approach to identify dehydration responsive nuclear proteins from chickpea (*Cicer arietinum* l.). *Mol Cell Proteomics.* Oct 6 [Epub ahead of print]
- Parker R, Flowers TJ, Moore AL and Harpham NV.** (2006) An accurate and reproducible method for proteome profiling of the effects of salt stress in the rice leaf lamina. *J Exp Bot* **57**:1109-1118
- Passardi F, Penel C and Dunand C.** (2004) Performing the paradoxical: how plant peroxidases modify the cell wall. *Trends Plant Sci* **9**:534-540
- Passardi F, Cosio C, Penel C and Dunand C.** (2005) Peroxidases have more functions than a Swiss army knife. *Plant Cell Rep* **24**:255-265
- Persson S, Rosenquist M, Svensson K, Galvao R, Boss WF and Sommarin M.** (2003) Phylogenetic analyses and expression studies reveal two distinct groups of calreticulin isoforms in higher plants. *Plant Physiol* **133**:1385-1396
- Plomion C, Lalanne C, Claverol S, et al.** (2006) Mapping the proteome of poplar and application to the discovery of drought-stress responsive proteins. *Proteomics* **6**, 6509-6527
- Rabbani MA, Maruyama K, Abe H, Khan MA, Katsura K, Ito Y, Yoshiwara K,**

- Seki M, Shinozaki K and Yamaguchi-Shinozaki K.** (2003) Monitoring expression profiles of rice genes under cold, drought, and high-salinity stresses and abscisic acid application using cDNA microarray and RNA gel-blot analyses. *Plant Physiol* 133:1755-1767
- Ringli C, Keller B, Ryser U.** 2001. Glycine-rich proteins as structural components of plant cell walls. *Cell Mol Life Sci* 58:1430-1441
- Rose JKC, Bashir S, Giovannoni JJ, Jahn MM and Saravanan RS.** (2004) Tackling the plant proteome: practical approaches, hurdles and experimental tools. *Plant J* 39: 715–733
- Rospert S, Dubaquitte Y and Gautschi M.** (2002) Nascent-polypeptide-associated complex. *Cell Mol Life Sci* 59:1632-1639
- Roxas VP, Smith RK Jr, Allen ER and Allen RD.** (1997) Overexpression of glutathione S-transferase/glutathione peroxidase enhances the growth of transgenic tobacco seedlings during stress. *Nat Biotechnol* 15:988-991
- Salekdeh GH, Siopongco J, Wade LJ, Ghareyazie B, and Bennett J** (2002) Proteomic analysis of rice leaves during drought stress and recovery. *Proteomics* 2:1131-1145.
- Seki M, Narusaka M, Ishida J, et al.** (2002) Monitoring the expression profiles of 7000 *Arabidopsis* genes under drought, cold and high-salinity stresses using a full-length cDNA microarray. *Plant J* 31: 279-292
- Shoji T, Suzuki K, Abe T, Kaneko Y, Shi H, Zhu JK, Rus A, Hasegawa PM and Hashimoto T.** (2006) Salt stress affects cortical microtubule organization and helical growth in *Arabidopsis*. *Plant Cell Physiol* 47:1158-1168
- Sivaguru M, Pike S, Gassmann W and Baskin TI.** (2003) Aluminum rapidly depolymerizes cortical microtubules and depolarizes the plasma membrane: evidence that these responses are mediated by a glutamate receptor. *Plant Cell Physiol* 44: 667–675
- Skopelitis DS, Paranychianakis NV, Paschalidis KA, Pliakonis ED, Delis ID, Yakoumakis DI, Kouvarakis A, Papadakis AK, Stephanou EG and Roubelakis-Angelakis KA.** (2006) Abiotic stress generates ROS that signal expression of anionic

- glutamate dehydrogenases to form glutamate for proline synthesis in tobacco and grapevine. *Plant Cell* 18:2767-2781
- Smirnoff N.** (1998) Plant resistance to environmental stress. *Curr Opin Plant Biol* 9: 214-219
- Steppuhn H and Raney JP.** (2005) Emergence, height, and yield of canola and barley grown in saline root zones. *Can J Plant Sci* 85: 815-827
- Steudle E.** (2000) Water uptake by roots: effects of water deficit. *J Exp Bot* 51: 1531-1542
- Taylor NL, Heazlewood JL, Day DA and Millar AH.** (2005) Differential impact of environmental stresses on the pea mitochondrial proteome. *Mol Cell Proteomics* 4:1122-1133
- Tian Q, Stepaniants SB, Mao M, Weng L, Feetham MC, Doyle MJ, Yi EC, Dai HY, Thorsson V, Eng J, Goodlett D, Berger JP, Gunter B, Linseley PS, Stoughton RB, Aebersold R, Collins SJ, Hanlon WA and Hood LE.** (2004) Integrated genomic and proteomic analyses of gene expression in mammalian cells. *Mol Cell Proteomics* 3:960-969
- Tsugita A and Kamo M.** (1999) 2-D electrophoresis of plant proteins. In: Link AJ, ed. *2-D Proteome Analysis Protocols*. Humana Press, 112, 95-97
- Wagner U, Edwards R, Dixon DP and Mauch F.** (2002) Probing the diversity of the *Arabidopsis* glutathione S-transferase gene family. *Plant Mol Biol* 49: 515-532
- Wang W, Vinocur B, Shoseyov O and Altman A.** (2004) Role of plant heat-shock proteins and molecular chaperones in the abiotic stress response. *Trends Plant Sci* 9: 244-252
- Wienkoop S and Weckwerth W.** (2006) Relative and absolute quantitative shotgun proteomics: targeting low-abundance proteins in *Arabidopsis thaliana*. *J Exp Bot* 57: 1529–1535
- Xu Z, Escamilla-Trevino L, Zeng L, Lalgondar M, Bevan D, Winkel B, Mohamed A, Cheng CL, Shih MC, Poulton J and Esen A.** (2004) Functional genomic analysis of *Arabidopsis thaliana* glycoside hydrolase family 1. *Plant Mol Biol* 55: 343-367
- Yan S, Tang Z, Su W and Sun W.** (2005) Proteomic analysis of salt stress-responsive proteins in rice root. *Proteomics* 5: 235-244

- Yan SP, Zhang QY, Tang ZC, Su WA and Sun WN.** (2006) Comparative proteomic analysis provides new insights into chilling stress responses in rice. *Mol Cell Proteomics* 5: 484-496
- Yang Z.** (2002) Small GTPases: versatile signaling switches in plants. *Plant Cell (Suppl)* 14: S375–S388
- Yotov WV, Moreau A and St-Arnaud R.** (1998) The alpha chain of the nascent polypeptide-associated complex functions as a transcriptional coactivator. *Mol Cell Biol* 18:1303-1311
- Zhang XP and Glaser E.** (2002) Interaction of plant mitochondrial and chloroplast signal peptides with the Hsp70 molecular chaperone. *Trends Plant Sci* 7: 14–21

Chapter 4: Functional characterization of a NaCl-inducible bHLH92 transcription factor

Introduction

Plant stress responses include changes in transcriptional regulation of gene expression, which depend on the interaction of transcription factors (TFs) with *cis*-regulatory sequences. Plants utilize a large number of TFs to regulate gene expression in response to developmental and environmental cues. Transcription factors belonging to different families have been shown to respond to abiotic stress (Chen *et al.*, 2002; Kreps *et al.*, 2002; Seki *et al.*, 2002; Jiang and Deyholos 2006). To activate or repress transcription, TFs must be localized to the nucleus and interact with the basal transcription apparatus. Therefore, environmental signals that regulate the activity of transcription factors may affect any one or a combination of these processes. Regulation of transcription factor activity is achieved by post-translational modifications such as phosphorylation and/or by *de novo* synthesis of TFs. The current analysis of identified TFs suggests that different stress signalling pathways may overlap or converge at specific points (Xiong *et al.*, 2002b; Yamaguchi-Shinozaki and Shinozaki, 2006; Zhu 2001 & 2002).

The basic/helix-loop-helix (bHLH) proteins are a superfamily of TFs in eukaryotes. This family is defined by the bHLH signature domain, which consists of ~60 amino acids with two functionally distinct regions: a basic region involved in DNA binding and an HLH region functioning as a dimerization domain (Murre *et al.*, 1989). bHLH proteins exhibit considerable sequence divergence, except for the conserved bHLH domain (Atchley *et al.*, 1999). According to their phylogenetic relationships, DNA-binding motifs, and functional properties, animal bHLH proteins were divided into six main groups (named as group A to F; Atchley and Fitch, 1997). The core motif recognized by the bHLH proteins is a consensus hexanucleotide sequence known as the E-box (5'-CANNTG-3'). There are different types of E-boxes, and one of the most common is the palindromic G-box (5'-CACGTG-3'). Certain conserved amino acids within the basic region of the protein provide recognition of the core consensus site, whereas other residues in the domain dictate specificity for a given type of E-box (Robinson *et al.*, 2000).

In *Arabidopsis*, 162 bHLH-encoding genes have been identified from the analysis of genome sequences (Bailey *et al.*, 2003; Heim *et al.*, 2003; Toledo-Ortiz *et al.*, 2003). The members of bHLH family of *Arabidopsis* were divided into 21 subfamilies by Toledo-Ortiz *et al.* (2003). Sequence analysis suggests that most of the plant bHLH proteins belong to Group B (Atchley and Fitch, 1997). In addition, recent reports have demonstrated that some plant bHLH proteins can interact with proteins that lack a bHLH domain. In particular, protein complexes with MYB, bHLH, and WD40 proteins were proposed to regulate guard cell and root hair differentiation (Ramsay and Glover, 2005). Compared to animals, only a small number of plant bHLH proteins have been characterized functionally, and in the model plant *Arabidopsis*, less than 40 bHLH genes have been characterized functionally so far. These studies have demonstrated that *Arabidopsis* bHLH genes (AtbHLH) are involved in controlling a variety of biological processes, including phytochrome-regulated light signaling pathways (Ni *et al.*, 1998; Huq and Quail, 2002), flavonoid biosynthesis (Nesi *et al.*, 2000), trichome development (Payne *et al.*, 2000), brassinosteroid response (Friedrichsen *et al.*, 2002), ABA signaling (Abe *et al.*, 1997, 2003), jasmonate (Anderson *et al.* 2004; Boter *et al.* 2004; Lorenzo *et al.* 2004), cold response (Chinnusamy *et al.*, 2003), iron acquisition (Jokaby *et al.*, 2004), auxin conjugate response (Rampey *et al.*, 2006) and stomatal development (Pillitteri and Torii, 2007). More recently, *Arabidopsis* AtNIG1 (NaCl-inducible gene) encoding a bHLH-type protein containing an EF-hand motif, was found to play a role in salt stress signaling (Kim and Kim 2006) and AtAIB (*Arabidopsis* ABA-inducible bHLH) was identified to be able to positively regulate ABA response (Li *et al.*, 2007).

In our previous microarray analysis of the *Arabidopsis* root response to salt stress, we identified 29 bHLH TFs that showed an at least 2-fold or higher changes in treated plants compared to controls (Jiang and Deyholos 2006). Among these 29 TFs, 15 increased in transcript abundance following salt treatment, whereas 14 decreased in abundance. Among the 15 up-regulated bHLH TFs, *bHLH92* (At5g43650) showed the highest fold-change (150 to 200-fold) by salt treatment across the three time-points examined. In the present study, we investigated further the function of *bHLH92* as a regulator of salt and osmotic stress responses through analyzing its mutant and

overexpression line phenotypes.

Materials and methods

Plant growth and stress treatment

Arabidopsis thaliana wild-type (WT, Col-0), two ABA mutants, *aba1-5* and *abi4-1* in Col genetic background, three *sos* mutants *sos1-1*, *sos2-1* and *sos3-1* in Col(*gll*) background were obtained from *Arabidopsis* Biological Resource Center (ABRC, Ohio, USA). Seeds were surface sterilized by 2.62% household bleach, sowed on ½ x Murashige-Skoog (MS) medium (Caisson Labs) supplemented with 1% sucrose and 0.3% Phytigel (Sigma-Aldrich), and stratified at 4 °C for 2 d. Seedlings were grown in 100 X 15 mm petri dishes for 7 d in a growth chamber and then were transplanted into soil (Sunshine Mix 1, Sunagro). The growth conditions were: photoperiod of 16 h light/8 h dark, relative humidity (RH) of 70%, 22 °C day and light, and the light intensity was ~125 $\mu\text{E m}^{-2} \text{s}^{-1}$. Rosetta leaves, stems, flower buds, young siliques (~7 DAF (days after flowering)) and mature seeds were harvested from WT plants for qRT-PCR analysis.

Hydroponics of *Arabidopsis* was performed as previously reported (Jiang and Deyholos, 2006). Twenty days after stratification (DAS), seedlings were treated with various stress treatments 1.5 h after the dark/light transition in the morning. Salt stress was performed by replacing the growth medium with fresh ½ X MS containing 150 mM NaCl; cold and heat stresses were applied by moving plants in fresh ½ X MS into a 4 °C or 37 °C growth cabinets (Danby); drought stress was applied by removing all liquid from the hydroponics tanks and leaving plants to dehydrate in the growth chamber; ABA and MeJA treatments were initiated by transferring plants into fresh ½ X MS containing 50 μM (\pm)-ABA(Sigma-Aldrich) or 10 μM MeJA (Sigma-Aldrich) and 0.05% ethanol; osmotic and oxidative stresses were applied by adding methyl viologen (Sigma-Aldrich) or mannitol (Sigma-Aldrich) into ½ X MS to a final concentration of 10 μM and 200 mM, respectively; AlCl_3 stress was applied by adding AlCl_3 (Sigma-Aldrich) to a final concentration of 25 μM in ½ x MS containing 200 μM CaCl_2 (Merck). Control plants were moved into fresh ½ X MS medium or ½ x MS supplemented with 0.05% ethanol at the same time. For cold and heat stresses, an

independent control was set up by moving plants into a growth cabinet with temperature set at 22 °C. All treatments lasted for 6 h or 24 h with roots and rosette leaves harvested separately from a pool of about 60 plants per treatment. Control samples were harvested at the same time. Samples were flash frozen in liquid nitrogen and kept at -80 °C before use.

Quantitative RT-PCR

Total RNA was extracted from 100 mg of tissue by RNeasy Plant Mini Kit (Qiagen). Dry seed RNA was extracted according to Suzuki *et al.*(2004). Formaldehyde agarose gel electrophoresis was run to examine the RNA quality (data not shown). Reverse transcription (RT) was performed with 2.5 µg of total RNA pretreated with DNase I by DNA-free RNA kit (Ambion) to obtain cDNA with SuperScript III (Invitrogen) or RevertAid H⁻ MMLV (Fermentas) and oligo(dT)₁₂₋₁₈ (Invitrogen) or oligo(dT)₁₈ (Fermentas) as the primer in a 20 µL reaction volume. Each cDNA sample was diluted 1:10 in sterile ddH₂O, and 2.5 µL of this dilution was used as template for qPCR. Primers for the PCR reactions were designed by PrimerSelect 5.00 of DNASTar (DNASTAR Inc) or PrimerExpress3.0 (Applied Biosystems) to have a T_m of 60 °C with the length of the amplicons between 80 and 180 bp. We first ran standard curves to check the amplification specificity and efficiency of each primer pair. cDNAs transcribed from 5 µg of total RNA was diluted with ddH₂O to be 1/4, 1/16, 1/64, 1/256, 1/1024 and 1/4096. Standard curves for each gene was performed using the dilution serials to cover the range of all template concentrations. Plots of Ct vs. log(input) were used to calculate the PCR efficiencies (*E*) according to the equation: $E = 10^{-1/\text{slope}}$. qRT-PCR was performed with a home-made 2 x SYBR GreenI master mix in 10 µL reactions using the ABI 7500 fast real-time PCR system (Applied Biosystems) according to the manufacturer's instruction. Each PCR reaction contains 2.5 µL of diluted cDNA, 0.4 µM of each of the primers and 5 µL of SYBR Green I mix (20 mM Tris (pH 8.3), 50 mM KCl, 3 mM MgCl₂, 0.8% glycerol 0.01% Tween 20, 2% dimethyl sulfoxide (DMSO), 0.2 mM dNTPs, 1×ROX, 0.25× SYBR Green I, 0.03 units/µL Platinum *Taq* DNA polymerase). The initial denaturing time was 2 min at 95°C, followed by 35 cycles consisting of 94°C for 15 s and 60°C for 1 min with a single fluorescence measurement. A dissociation curve was run after the PCR cycles to

examine that only single peak showed for each primer pair. The primers of genes tested for expression levels by quantitative PCR were as follows: *RD29A* (At5g52310), Fwd: 5'-AGGAAGTGAAAGGAGGAGGAGGAATG-3', Rvs: 5'-GTCGCACCACCACC AAACCAG-3'; *bHLH92* (At5g43650), Fwd, 5'-CACGGAAGTGGAGGATGATGTC-3', Rvs, 5'-GTCGCCGGCTTCTTTCCTTCTCT-3'; *UBQ10* (At4g05320), Fwd, 5'-GGCCTTGTATAATCCCTGATGAA -3', Rvs, 5'-AGAAGTTCGACTTGTCATTA GAAAGAAA-3'. Gene-specific PCR efficiency was used to calculate the expression of target genes relative to the expression of *UBQ10* reference gene (Pfaffl, 2001). qPCR for each gene was done on three biological replicates with duplicates for each biological replicate. The relative transcript level was determined for each sample, normalized using the *UBQ10* cDNA level, and averaged over the three replicates with triplicates for each biological replicate.

Identification of homozygous T-DNA insertion mutants

T-DNA insertion mutants (bulk T₄) developed by the SALK institute (Alonso *et al.*, 2003) and Syngenta (Sessions *et al.*, 2002) were obtained from ABRC (Ohio, USA). Genomic DNA was extracted from 3-week-old rosette leaves of about 14 independent T₄ lines for each T-DNA insertion mutant as well as from WT plants by 2% CTAB method (Chen and Ronald, 1999). PCRs were conducted to screen out putative homozygous mutant lines using two combinations of primers, LB/RP and LP/RP as per the protocol described at <http://signal.salk.edu/tdnaprimers.2.html>. The primer sequences are as follows: LBb1 for SALK line, 5'GCGTGGACCGCTTGCTGCA ACT3', SALK_033657, LP, 5'-AAATTTTGTGCAAGACACG-3'; RP, 5'-GATG GTCTTGAAACTTGTTCG-3'; SAIL_511_E04, LP, 5'-AAAACCGGATTCAAC CAATTC-3'; RP, 5'-GAAGAAAATACGGTGGCTCC-3'; LB3 for SAIL line, 5'-TAGCATCTGAATTCATAACCAATCTCGATACAC-3'.

To confirm the loss of gene expression in putative homozygous T-DNA insertion lines, total RNA was extracted from 4 week old rosette leaves according to Wadsworth *et al.*(1988) with modifications. The extracted RNA was treated with DNaseI (Ambion) before running RT using oligo(dT)₁₂₋₁₈ and SuperScriptII (Invitrogen) followed by PCR with gene-specific primers. The primers used are the same as for the overexpression construct (see below). The *ACTIN2* (At3g18780) gene primers used are 5'-GGTCGTA

CA ACCGGTATTGTG-3' and 5'-GATTCCTGGACCTGCCTCA-3'. PCR product was cloned into pBS SK+ based T vector constructed according to Hadjeb and Berkowitz (1996), sequenced using T7 and T3 promoter primers and ABI BigDye reagents (Applied Biosystems) and analyzed in DNASTar (DNASTAR Inc).

Plasmid constructions

General molecular biology experiments were performed according to Sambrook and Russel (2001). Total RNA was extracted from *Arabidopsis* leaves according to Wadsworth *et al.*(1988) with modifications as described previously and treated with DNaseI (Ambion). RT was performed with SuperScriptII (Invitrogen) and oligo(dT)₁₂₋₁₈ (Invitrogen) using 2 µg RNA as the template. For subcellular localization of TFs, binary vector-pCsGFPBT (GenBank Acc# DQ370426) was restricted with *Nco* I, dephosphorylated with TSAP (thermo-sensitive alkylphosphatase, Promega) and purified by a PCR purification kit (Qiagen). The coding region of *bHLH92* (744 bp) was PCR-amplified using primers 5'-CATG CCATGGATAACTTTTTCTAGGT-3' and 5'-CATGCCATGGCTCCACCTCCACCTCCGTAGTCTTTGTAAAAGGAAG-3''. The PCR products were purified, restricted by *Nco* I, purified again and cloned into *Nco* I restricted pCsGFPBT. For screening of positive clones, sequencing and PCR verification of T-DNA insertion into *Arabidopsis* genome, another three primers were designed: pCsGFPBT_TL_Fwd: 5'-CGAATCTCAAGCAATCAAGC-3', pCsGFPBT_sGFP_141Rvs: 5'-GAACTTCAGGGTCAGCTTGC-3', and pCsGFPBT-sGFP_729 Rvs: 5'-CTAGAGGATCCCCTTGACAG-3'.

For promoter-GUS fusion, a binary vector pCAMBIA1391Z (AF234312) containing a promoterless GUS gene interrupted by a catalase intron and a double cauliflower mosaic virus (CaMV) 35S promoter driving the hygromycin B phosphotransferase gene in the T-DNA region was modified. The double CaMV35S promoter was removed to eliminate its influence on the nearby promoter being analyzed. This was done by excision of the *Bst* XI-*Xho* I fragment (1881 bp) containing the 35S::hyg-35S terminator fragment from pCAMBIA1391Z, blunted by T4 DNA polymerase (Fermentas), and re-ligated to result in the modified vector p0381Z. The promoter region (PbHLH92, 1 kb upstream from the translation initiation codon) of *bHLH92* gene was PCR-amplified using WT genomic DNA as the template and

primers: 5'-CCC AAGCTT GAA ATG TGT GAG GTT TTG TGT -3' (Fwd) and 5'-CATG CCATGG CTTTCTCTCTTAGTGTGTTTG-3' (Rvs). The resulting fragments were digested with *Hind* III and *Nco* I and inserted into p0381Z.

For overexpression vectors, the coding region of *bHLH92* was cloned by RT-PCR. An *Nco* I site was introduced to the 5' end of forward primers and a *Bst*E II site was introduced to the 5' end of reverse primer. The primer sequences are as follows: Fwd, 5'-CATGCCATGGAT AAC TTT TTT CTA GGT TTG-3', Rvs, 5'-TCGG GGTCA CCTTAGTAGTCTTTGTAAAAGGA-3'. PCR products were purified by a Qiagen PCR purification kit, restricted with *Nco* I and *Bst* EII, and purified again using a Qiagen kit. Then it was ligated into *Nco* I and *Bst*E II digested of pCAMBIA1303 (www.cambia.org, AF234299) in place of *GUS-GFP*. For screening of positive clones, sequencing and PCR verification of T-DNA insertion into Arabidopsis genome, another two primers were designed: pCAM1303_SeqF, 5'CAACCACGTCTTCA AAGCAA3', pCAM1303_SeqR, 5'AAGACCGGCAACAGGATTC3'.

All PCR amplifications were conducted using high-fidelity *Pfx* DNA polymerase (Invitrogen). Five independent positive clones were sequenced using BigDye reagent in ABI3730 sequencer (Applied Biosystems) to confirm that no mismatch or frame-shift mutations occurred.

Arabidopsis transformation and selection

All the above constructs were introduced into agrobacterium GV3101 through the freeze-thaw method and transformed into WT Arabidopsis (Col-0) by floral dip (Clough and Bent, 1998). The PbHLH92-*GUS* construct was co-transformed into *Arabidopsis* with a GUS-free vector pCAMBIA1300 containing only 35S::hyg-Ter within T-DNA region. This was done by culturing agrobacterium harboring the PbHLH92-*GUS* construct and pCAMBIA1303 separately to an OD₆₀₀ of ~0.8 before combined in a ratio of (vol/vol, 2:1) just before transformation. pCsGFPBT and pCAMBIA1303 plasmids were also transformed independently into WT *Arabidopsis* as controls. T₁ seeds from ten independent lines were separately surface sterilized and selected on 10 separate ½ X MS medium plates containing 50 mg/L of hygromycin B (Sigma-Aldrich). After 7 d, putative transgenic seedlings were transplanted into pots containing potting medium and seeds were harvested (T₂), which were further sowed

on ½ X MS medium containing 25 mg/L of hygromycin B, and after 7 d, resistant lines were transplanted into soil pots. Genomic DNA was extracted from 28 d old rosette leaves using 2% CTAB buffer and PCR confirmation of transgene was performed using suitable primers. For co-transformed Arabidopsis, we ran PCR assay of hygromycin-resistant lines using the forward promoter primer (see above) and a *GUS*- specific reverse primer (5'-TGCCCAACCTTTCGGTATAA-3'). For overexpression lines, we first ran PCR confirmation of the integration of T-DNA into T₁ plant genome using the gene-specific forward primer and a pCAMBIA1303 vector-specific reverse primer (5'-AAGACCGGCAACAGGATT C-3'). qRT-PCR screening to detect high expression of transgenes in T₂ generation was performed as described above. Homozygous T₃ seeds were used for phenotypic scoring under various abiotic stresses and GUS staining, while T₂ lines were checked for GFP expression.

GFP observation

Five-day-old seedlings from ten independent T₂ lines were mounted on slides for GFP observation under confocal or epifluorescent microscopes. The roots or hypocotyls were observed and photographed with an Olympus BX51 microscope (Olympus Optical Co.) equipped with a Photometrix CoolSnap fx digital camera (Roper Scientific) and a MicroColor liquid crystal tunable RGB filter (Cambridge Research & Instrumentation, Inc.). For confocal laser scanning microscopy, samples were mounted in water and observed with a Zeiss Axiovert 100M microscope equipped with a Zeiss LSM510 laser module confocal unit (Carl Zeiss). GFP was visualized with the 488-nm line of an Argon laser at 55% of output and 5% transmission, and with either a 505–530-nm bandpass filter.

Histochemical GUS staining

GUS staining was conducted as described by Weigel and Glazebrook (2002) and Salinas and Sanchez-Serrano (2006). 5-d old T₃ seedlings from ten independent lines grown on MS agar plates supplemented with 25 mg/L hygromycin or hydroponically cultured, 20 DAS whole seedlings, flower buds and young siliques were harvested and incubated in a staining solution (0.1M sodium-phosphate buffer, pH 7, 2 mM K₄Fe(CN)₆, 2 mM K₃Fe(CN)₆, 0.2% Triton X-100 and, 10 mM EDTA, 2 mM X-Gluc (Rose Scientific)) in 50 ml Falcon tubes for 4-10 h at 37°C in the dark and then washed

2 times with 70% (v/v) ethanol at room temperature to remove chlorophyll. Samples were viewed and photographed with a Leica dissecting microscope (Leica Microsystems) equipped with a JVC digital camera. The representative and consistent results were reported. All images were assembled in Adobe Photoshop CS (Adobe Systems).

Stress tolerance tests and REL measurement

Seeds from wild-type (Col-0), homozygous T-DNA insertion lines, or three independent overexpression lines were harvested from plants grown under similar conditions and stress tolerance tests were performed essentially according to Verslues *et al.* (2006). In brief, seeds were surface sterilized and sowed on 1/2 x MS medium (pH5.7) containing 1% sucrose and 0.8% Phytoblend (Caisson Labs) contained in square petri dishes (Fisher Scientific). After stratification at 4 °C for 2-3 d, seed plates were placed vertically under the growth condition of 16 h light/8 h dark, RH~70%, light intensity ~130 $\mu\text{E}/\text{m}^2/\text{s}$. After 4-5 d, seedlings with similar growth vigor and root length were transferred using blunt forceps onto normal medium plates (1/2 x MS medium, pH5.7, 1% sucrose, 0.8% Phytoblend) or plates supplemented with 100 or 150 mM NaCl, 100, 150 or 200 mM mannitol, or 0.1 or 0.5 μM methyl viologen, or 0.5 μM (\pm)-ABA and continued to grow vertically for another one week before scored and photographed. Root length was measured either by using a ruler or the free ImageJ software available from <http://rsb.info.nih.gov/ij/index.html>. The relative electrolyte leakage (REL) assay was performed according to Jiang *et al.* (2007). Each assay was repeated two to three times and data were analyzed using two-tailed Student's *t*-test and significant differences ($p < 0.05$) are reported in the figures or figure legends.

Microarray profiling of *bhlh92* mutant

Qiagen Operon 26k *Arabidopsis* oligonucleotide microarrays were used to profile transcriptomes of *bhlh92* T-DNA insertion mutants (*bhlh92-1*) compared to WT control. Total RNA samples from hydroponically cultured roots with three independent biological replicates were extracted using the RNeasy Plant Mini kit (Qiagen). Potential DNA contamination was removed using a DNA-free kit (Ambion), and 5 μg of total RNA was reverse transcribed into cDNAs using Revert-Aid H⁻ MMLV (Fermentas) and primers contained in the Array900 kit (Genisphere). Microarray hybridizations

with dye-flip design were performed using an Array900 kit according to the manufacturer's manual and as described in Jiang and Deyholos (2006). Slides were scanned immediately using ArrayWox (Precision Scientific) and transformed into .tiff images. Data were analyzed using the TIGR TM4 software suite as described in Jiang and Deyholos (2006).

To validate microarray data, qRT-PCR was performed with six selected genes. The primers designed by PrimerExpress3.0 (Applied Biosystems) are as follows. At5g19890 (*PER59*), Fwd, 5'-TAACATCACCGACGTCGTAGCT-3'; Rvs, 5'-TCCCAAGCCGGTAAAGTTGA-3'; At3g02480, Fwd, 5'-GCAAAACGCGAGCTACCAA-3'; Rvs, 5'-TTGAGCTGAAGCAGCAGCAT-3'; At4g12550 (*AIR1*), Fwd, 5'-GGGACTTGTCCTAAAAATTCCAT-3'; Rvs, 5'-TTCCAAGAATGCTAGCCTTGA CT-3'; At1g20450 (*ERD10*), Fwd, 5'-CGCTGGTTGAAACAGCAACA-3'; Rvs, 5'-GTCTTGCGTGATAACCTGGAA-3'; At5g56270 (*WRKY2*), Fwd, 5'-ACCGGAGG TCAATACGAGTCA-3', Rvs, 5'-TTCCATGTGTCCCTCGATCA-3'; At5g05340 (*PER52*), Fwd, 5'-TCACCACGGCAGCTTCTTTC-3', Rvs, 5'-GTAGAG CCGCCGTTGAAGAG-3'. qRT-PCR was performed as described previously.

Phylogenetic tree

The protein sequences of 162 members of the *Arabidopsis* bHLH family were retrieved from DATF (<http://datf.cbi.pku.edu.cn/>) and RARTF (<http://rage.gsc.riken.jp/rartf/>). The bHLH DNA binding domain boundary was defined as per Atchley *et al* (1999) and Toledo-Ortiz *et al.*(2003). Only the ~ 60 amino acid residues in the bHLH domain were aligned using ClustalX (v1.83) (<http://bips.u-strasbg.fr/fr/Documentation/ClustalX/>) with a gap opening penalty of 35 and gap extension penalty of 0.75 in pairwise alignment, and a gap opening penalty of 15 and gap extension penalty of 0.30 in multiple alignment parameters parameter settings. Aligned sequences were further examined with gaps introduced manually for optimal alignment. The parsimony tree was drawn using PAUP*4.0 (Sinauer Associates, Inc.). For Fig. 4-1, Adobe Acrobat 7.0 professional was employed to output the alignment file and to display the phylogenetic tree in a PDF format.

Results

Sequence and phylogenetic analysis of *bHLH92* gene

The predicted bHLH92 protein encoded 247 amino acids with a molecular weight of 29.0 kD and an isoelectric point of 10.22 (Fig. 4-1). We further performed multiple alignment of bHLH92 protein to four other bHLH TFs: MYC2/JIN1/bHLH6, MYC3/bHLH5, MYC4/bHLH4 and ICE1/bHLH116, which had been shown to be involved in ABA (Abe *et al.* 2003), jasmonate (Anderson *et al.* 2004; Boter *et al.* 2004; Lorenzo *et al.* 2004) and cold signaling (Chinnusamy *et al.*, 2003). Using the simple modular architecture research tool (SMART, Letunic *et al.*, 2006, <http://smart.embl-heidelberg.de/index2.cgi>), we identified a bHLH domain for each of the five bHLH TFs we aligned (Fig. 4-1). The amino acid residues within the bHLH domain of bHLH92 are quite divergent from the those of MYC2, MYC3 and MYC4 (Fig.4-1).

Although the signature bHLH domain of the AtbHLH proteins is well conserved, the regions outside of this domain in the remainder of the protein generally are poorly conserved (data not shown). The length of AtbHLH proteins range from 129 aa to 1513 aa. Therefore, we used only the ~60 aa encoding the bHLH domain for phylogenetic analysis, as has been the case for previous studies of this kind (Atchley and Fitch, 1997; Atchley *et al.*, 1999; Morgenstern and Atchley, 1999; Ledent and Vervoort, 2001). Using the online SMART tool, we analyzed 162 putative bHLH protein sequences and found that 159 proteins had a typical bHLH domain whereas three, bHLH157, bHLH158, and bHLH159, did not contain a bHLH domain. InterPro (<http://www.ebi.ac.uk/interpro/>) was used to define the bHLH domains of the latter three bHLH proteins. As shown in the phylogenetic tree, bHLH92 is closely related to bHLH95 (At1g49770) and these two proteins form a relatively independent subclade (Fig 5-2). We also found that neither of these two bHLH gene probes is included in the Affymetrix ATH1 probe set.

In our previous microarray profiling, we found that the *bHLH95* mRNA level was not changed by salt treatment in any of the three time-points examined, whereas *bHLH92* transcript level was greatly up-regulated by salt treatment in *Arabidopsis* roots. Using qRT-PCR, we found that *bHLH92* mRNA abundance in WT control plant roots was very low. To analyze the expression patterns of *bHLH92* in different tissues,




Fig. 4-1. Multiple alignment of bHLH92 and four other *Arabidopsis* bHLH TFs. The symbol “-” in the sequence is the gap introduced for optimal alignment. The asterisks above the sequence indicate identical amino acids, while symbols “:” and “.” indicate conserved and less conserved amino acids, respectively. The bHLH domains at the C-termini are shown by four double-headed arrows under the sequence. The putative nuclear localization signal (NLS) is shown by a line above the sequence.




Fig. 4-2. Phylogenetic tree of 162 AtHLHs. The phylogenetic tree was based on the amino acid sequences from bHLH domains only. The numbers on the branches represent branch length. The arrow indicates bHLH92 and bHLH95.

qRT-PCR was used to measure RNA abundance in untreated roots, stems, leaves, flowers, siliques, and seeds. *bHLH92* transcripts were detected at low abundance in all tissues, except flowers and immature siliques, where they were considerably more abundant than in roots. (Fig. 4-3).




Fig. 4-3. qRT-PCR analysis of *bHLH92* transcript level in various tissues and organs. R, roots; L, rosette leaves; St, stems; F, flower buds; Si, young siliques; Se, dry seeds. The abundance of *bHLH92* in roots was normalized to be 1. Data were from 3 biological replicates with S.D. shown.

Responses of *bHLH92* gene to various abiotic stresses

To investigate the ability of *bHLH92* transcript abundance to be regulated by various environmental stresses, 20 DAS old hydroponic cultured wild-type *Arabidopsis* plants were treated various stresses for 6 or 24 h and qRT-PCR was used to compare the transcript levels in treated roots in relation to those in corresponding controls. The results showed that *bHLH92* was induced by salt, drought, mannitol, cold, MeJA and aluminum stresses at both 6 h and 24 h time points (2-fold cut-off), and it was also up-regulated by Paraquat at 6 h (Fig. 4-4A). For comparison, we also observed that a stress marker gene *RD29A* (Yamaguchi-Shinozaki and Shinozaki, 1993) was induced by 150 mM NaCl, cold, drought, mannitol, ABA and AlCl₃ treatments (Fig. 4-4B).




Fig. 4-4. qRT-PCR profiling of *bHLH92*(A) and *RD29A*(B) transcript levels in Arabidopsis roots treated by various abiotic stresses for 6 or 24 h as described in the Materials and Methods section. Values shown are the mean of three independent biological replicates with standard deviation.

Salt-induced accumulation of *bHLH92* transcript in ABA and SOS mutants

ABA regulates several aspects of plant development including seed development, desiccation tolerance of seeds and seed dormancy, and plays a crucial role in the plant's response to abiotic (drought, salinity, cold, and hypoxia) and biotic stresses. Stress-responsive genes have been proposed to be regulated by both ABA-dependent and

ABA-independent signalling pathways (Shinozaki and Yamaguchi-Shinozaki, 2000; Zhu, 2002; Chinnusamy *et al.*, 2005).

Regulation of cellular ion homeostasis during salinity stress is critical for plant salt tolerance. One of the responses of plant cells to salt stress is the generation of a cytosolic Ca^{2+} transient (Knight *et al.*, 1997) and the subsequent activation of Ca^{2+} sensor protein expression and/or activity. Molecular analysis of *sos* mutants of *Arabidopsis* led to the identification of components (SOS1, SOS2 and SOS3) of a pathway that transduce a salt stress-induced Ca^{2+} signal to reinstate cellular ion homeostasis (Zhu, 2002).

To examine whether the salt-induced expression of *bHLH92* is dependent on the ABA and SOS pathways, we used qRT-PCR analysis to investigate the changes of *bHLH92* in ABA biosynthesis and signaling mutants, *aba1-5*, *abi4-1* as well as *sos1-1*, *sos2-1* and *sos3-1* mutants. *aba1-5* is a mutant in which zeaxanthin epoxidase (ZEP), a key enzyme in ABA biosynthesis, is non-functional, while *abi4-1* is a mutation in a




Fig. 4-5. qRT-PCR analysis of *bHLH92* transcript levels in wild-type (WT), *aba1-5*, *abi4-1*, *sos1-1*, *sos2-1* and *sos3-1* mutants treated by 150 mM NaCl for 6 h. Data were the mean values from 3 biological replicates and error bars are the S.D. Asterisks indicate a significant difference ($p < 0.05$) between the mutant and WT.

AP2 domain transcription factor (Finkelstein and Rock, 2002). In the absence of functional *ABA1* and *ABI4*, the fold-induction of *bHLH92* decreased by 37% and 32%, respectively (Fig. 4-5), suggesting that salt-induced *bHLH92* expression is partially dependent on the ABA pathway. We found that the steady-state mRNA level of *bHLH92* in the *sos2-1* mutant was higher than that in WT ($p < 0.05$, Fig. 4-5), and in

sos1-1 and *sos3-1* mutants, the difference is insignificant ($p=0.05-0.1$). Kamei *et al.* (2005) found, through microarray profiling, that *bHLH92* was ~2-fold upregulated by 250 mM NaCl stress treatment for 2 h in *sos3* mutants compared to salt-treated WT plants. The difference between our data and Kamei *et al.* (2005) data may be due to the different concentrations of NaCl used.

Promoter-GUS fusion

To examine the role of a putative promoter region on the expression of *bHLH92* in *Arabidopsis* tissues, we fused the 1 kb region upstream of the translational start codon to the β -glucuronidase (*GUS*) reporter gene to determine the developmental, organ-specific and spatial expression patterns it conferred. Promoter analyses can be compromised by interactions between the test promoter and those driving the expression of other genes within the same construct. Information from the pCAMBIA vector series website (www.cambia.org/daisy/cambia/585) has acknowledged problems with some of the promoter-testing vectors. The constitutive CaMV 35S promoter in its various forms contains one or more enhancer sequence that can bidirectionally drive transcription and/or augment the ectopic expression of noncontiguous nearby genes (Xie *et al.*, 2001; Yoo *et al.*, 2005). It has been documented that transcriptional interference can occur between promoters within a construct, depending on their separation distance and relative orientation (Yoo *et al.*, 2005).

To avoid the problem with CaMV35S enhancers, we cloned the 1 kb promoter region (P**bHLH92**) into a modified binary vector-p0381Z (Fig 4-6A), which does not contain CaMV35S promoter and plant selectable marker gene. We therefore co-transformed our promoter-GUS constructs into WT *Arabidopsis* with another binary vector-pCAMBIA1300, which contains CaMV35S and plant selectable marker gene (hygromycin resistant gene) only within the T-DNA borders (Fig. 4-6A).

Ten independent T₃ transgenic *Arabidopsis* lines were examined to determine the GUS expression patterns conferred by the putative *bHLH92* promoter region. In 5 d old seedlings, a weak GUS expression was observed only in leaf veins (Fig. 4-6B). In 18 d old seedlings, GUS expression was evident in leaf veins (Fig. 4-6C). In inflorescences, GUS expression was visible at the base of young flower buds (Fig. 4-6D). In young siliques (7 d after anthesis, DAA), weak GUS expression was detected only in the seed




Fig. 4-6. Histochemical localization of GUS activity in representative transgenic *Arabidopsis* plants harboring PbHLH92-GUS construct or pCAMBIA1303 binary vector.

A. Map of T-DNA regions of promoter fusion transformation vector-p0381Z and pCAMBIA1300. LB, left border; RB, right border; Nos-ter, nopaline synthase terminator, CaMV 35S, cauliflower mosaic virus 35S promoter, HPTII, hygromycin phosphotransferase. GUS, β -glucuronidase. B-E, GUS staining patterns in 5-d old seedling (B), 18-d old seedling (C), inflorescence (D), and 7-DAF old siliques (E) harboring PbHLH92-GUS construct. F-I, GUS staining patterns in 5-d old seedling (F), 18-d old seedling (G), inflorescence (H), and 7-DAF old siliques (I) harboring pCAMBIA1303 vector. J-M, GUS staining patterns of 19-d old transgenic line harboring PbHLH92-GUS construct under normal condition (J) or treated by

salt (K), drought (L) and 10 μ M JA (M) for 6 h.

pod (Fig. 4-6E). In the control transgenic plants harboring 35S-driven *GUS*, almost all the tissue and organs examined showed strong *GUS* stainings (Fig. 4-6F-I).

We then examined the *GUS* expression in 19 d old seedlings treated by salt, drought, cold, heat, ABA or JA for 6 h and compared the staining pattern to the untreated seedling. Our results demonstrated that *GUS* expression under PbHLH92 was enhanced by salt, drought and JA treatments compared to the mock seedling (Fig. 4-6J-M) while the other abiotic stress treatments tested failed to affect the *GUS* staining (data not shown).

The putative *cis*-elements contained in the promoter region of *bHLH92* gene were analyzed using PlantCare (Lescot *et al.*, 2002, Table 4-1). We detected three ABRE (ABA-responsive elements, PyACGTGG/T, Yamaguchi-Shinozaki & Shinozaki, 1993) and one CE3 (coupling element). We also found four MeJA-responsive *cis*-

Table 4-1. Putative regulatory sequences identified in the 1 kb upstream region of *bHLH92*.

motif	Sequence	position	strand	Function of regulatory elements
CGTCA-motif	CGTCA	-990	-	MeJA-responsiveness
		-252	+	
		-286	+	
		-182	+	
HSE	AGAAAATTCG	-333	+	heat stress responsiveness
ABRE	TACGTG	-569	+	abscisic acid responsiveness
	CACGTG	-118	+	essential for the anaerobic induction
		-92	+	
ARE	TGGTTT	-198	+	
		-98	-	
TC-rich repeats	ATTTTCTCCA	-397	+	defense and stress responsiveness
TCA-element	CCATCTTTTT	-767	-	salicylic acid responsiveness
W box	TTGACC	-691	-	wounding and pathogen response
		-141	+	
		-628	-	
CE3	GACGCGTGTC	-985	+	ABA and VP1 responsiveness
WUN-motif	AAATTCCT	-593	-	wound-responsive element

Putative elements involved in light responses (found in all promoters) are not included in the table. PlantCare: <http://bioinformatics.psb.ugent.be/webtools/plantcare/html/>

elements, one heat shock element (HSE), one salicylic acid (SA)-responsive element and one wounding responsive element (Table 4-1), indicating that *bHLH92* may respond to these treatments. Interestingly, *bHLH92* promoter region also contains 3 W-box elements (Table 4-1), suggesting its activity might be regulated by some unknown upstream WRKY TF(s).

Sub-cellular localization

To examine the sub-cellular localization of bHLH92, its coding region was fused to the N-terminus of synthetic green fluorescent protein (sGFP) (Chiu *et al.*, 1996) and expressed constitutively under the control of cauliflower mosaic virus (CaMV) 35S promoter (Fig. 4-7A). The *sGFP* gene alone under the control of the CaMV 35S promoter served as a control. Stable transformation of WT *Arabidopsis* was performed and 5-7 day old roots from 10 independent T₂ seedlings were checked for GFP expression. The bHLH92-sGFP fusion protein was localized in the nucleus (Fig. 4-7B). With the control vector alone, sGFP signals were distributed in both cytoplasm and nucleus (Fig. 4-7C). This demonstrates that bHLH92 is a nuclear-localized protein, which is consistent with its predicted function as a transcription factor.

Identification of T-DNA insertion lines and stress tolerance tests

In order to test the sensitivity of mutants of *bHLH92* gene to abiotic stress, we obtained and screened out two homozygous T-DNA insertion mutants designated as *bhlh92-1* (SALK_033657) and *bhlh92-2* (SAIL_511E_04). We confirmed by qRT-PCR that the *bHLH92* transcript was undetectable in either of the two mutants, indicating presence of a null mutation (Fig. 4-9A) The T-DNA sites in both mutants are in the first intron (Fig. 4-8B).

Under normal growth conditions, we did not observe significant morphological differences between these null mutants and WT plants grown in soil. However, on normal medium plate, the root elongation of the mutant was significantly longer than that of WT (Fig. 4-8D), suggesting that bHLH92 might negatively regulate root elongation. Root elongation assay of the mutants (compared to wild-type) under NaCl, mannitol, and methyl viologen (Paraquat) stress conditions were performed. Under 100 or 150 mM NaCl treatments, *bhlh92-1* mutant roots were slightly shorter than WT roots, but the difference was not significant (Fig. 4-8C, D). Under 100 or 150 mM







Fig 4-7. Subcellular localization of bHLH92 fused to GFP in *Arabidopsis* root cells.

A. Map of T-DNA region of the GFP fusion vector-pCsGFPBT. HPTII, hygromycin phosphotransferase. GFP, green fluorescent protein. TL, translational leader.

B. Subcellular localization of bHLH92-sGFP in root cells of 5-d old transgenic seedlings under confocal microscope.

C. Vector control-pCsGFPBT. The fluorescence (left) and bright-field (middle) images are overlaid at right.

mannitol treatment, *bhlh92-1* mutant roots were significantly shorter than WT roots ($p < 0.04$) (Fig. 4-8C, D). No significant difference on root elongation was observed between mutants and WT under oxidative stress or ABA treatment (Fig. 4-8 C, D), although rosette leaves of *bhlh92-1* were a little yellowish on ABA medium (Fig. 4-8C). We further examined the relative electrolyte leakage (REL) of *bhlh92-1* seedlings under 150 mM NaCl treatment for 6 h, and found that REL of the *bhlh92* mutant was significantly higher than that of WT plants ($p < 0.05$, Fig. 4-8E). We did not observe any difference between the mutants and WT plants under other types of abiotic stress conditions (data not shown).







Fig. 4-8. Identification of T-DNA insertion mutants and stress tolerance testing.

A, RT-PCR verification of null mutants of SALK_033657 (*bhlh92-1*) and SAIL_511_E04 (*bhlh92-2*) using *ACTIN2* gene as a positive control. B, Schematic diagrams of *bHLH92* gene and T-DNA insertion sites. Closed boxes represent exons (black, coding regions; white, 5'UTR or 3'UTR), and lines between closed boxes represent introns. The positions of T-DNA insertions are indicated by triangles. C, Stress tolerance test of *bhlh92-1* mutant on $\frac{1}{2}$ x MS medium plates containing 100 or 150 mM NaCl or mannitol, 0.5 μ M MV or ABA. D, Quantitative assay of root elongation of *bhlh92-1* under stress treatments. Data are means \pm S.D., n=8-10. Asterisks indicate a significant difference ($p < 0.05$) between the mutant and WT. E, Relative electrolyte leakage (REL) from NaCl-treated WT and *bhlh92-1* seedlings following 6 h exposure to 150mM NaCl. Data are means \pm S.D., n=6. Asterisks indicate a significant difference ($p < 0.05$) between the mutant and WT.

Over-expression of *bHLH92* gene in *Arabidopsis*

The putative role of *bHLH92* in abiotic stress was also investigated by overexpressing the cDNA under the control of CaMV35S promoter in *Arabidopsis* (Fig. 4-9A). The expression level of the transgene was estimated for 10 independent T₂ lines by quantitative RT-PCR. Lines 2, 3 and 6 were found to have the highest abundance of transcript of the transgene (Fig. 4-9B) and were used for the root elongation assay.

None of the *35S:bHLH92* transgenic plants showed any phenotypic changes compared with control plants under normal conditions. We therefore conducted root elongation assays using homozygous T₃ progeny of these transgenic lines under normal and several types of abiotic stresses, i.e. NaCl, mannitol, Paraquat, ABA and, compared the responses of overexpressing lines to WT control seedlings. On normal medium plate, there is no significant difference between *bHLH92*-overexpressing lines and WT (Fig. 4-9 C, D). Under 100 mM NaCl treatment, *bHLH92*-overexpressing plants were 25% longer than WT roots and, we also detected more lateral roots on *bHLH92*-overexpressing plants than WT plants under salt treatment (Fig. 4-9 C, D). Under 0.1 μM Paraquat treatment, roots of overexpressing lines were also 15.5% longer than WT roots, and *bHLH92*-overexpression lines also had more lateral roots than WT control (Fig. 4-9 C, D). Under 200 mM mannitol treatment, overexpression plant roots were also 28.1% longer than WT control ($p < 0.05$, Fig 5-9 C, D). Therefore, overexpression of *bHLH92* increased plants tolerance to NaCl, mannitol, and Paraquat treatments, suggesting that overexpression of *bHLH92* in *Arabidopsis* can enhance plants' tolerance to high salinity, osmotic and oxidative stresses. We did not observe any significant difference between the *35S:bHLH92* lines and WT under cold or heat treatments (data not shown).






Fig. 4-9. Identification and stress tolerance testing of *bHLH92*-overexpressing *Arabidopsis*.
 A, Map of the T-DNA region of overexpression transformation vector-pCAMBIA1303.
 B, qRT-PCR analyses of transcript levels of *bHLH92* in *Arabidopsis* leaves of ten independent T_2 overexpression lines. The ratios were that of genes in overexpression lines compared to that of in wild-type and normalized to the level of endogenous gene UBQ10.
 C, Performance of *bHLH92* transgenic line and wild-type plants under salt, mannitol, paraquat or ABA treatments.
 D, Effect of 100 mM NaCl, 200 mM mannitol, 0.1 μ M Paraquat or 0.1 μ M (\pm)-ABA on root elongation of WT and *bHLH92*-overexpressing seedlings. Data are means \pm S.D. (n = 8–11). Asterisks indicate a significant difference ($p < 0.05$) between the *bHLH92*-overexpressing plants (OX) and WT exposed to the same treatment.

Microarray identification of bHLH92 target genes

To identify genes that are regulated, directly or indirectly, by the activity of bHLH92, we used a full-genome QiagenOperon 70-mer oligo microarray representing 26, 090 genes (Jiang and Deyholos 2006) to measure transcript abundance in hydroponically cultured *bhlh92* or wild-type plants treated with 150 mM NaCl for 6 h. With a 1.5-fold cutoff ($p < 0.05$), we detected ten and nine genes that decreased- and increased in abundance, respectively, in the *bhlh9-1* mutant compared to WT (Table 4-2). Among the up-regulated gene list, we detected two stress marker genes, *ERD10* and *KINI1*, which encode LEA-like proteins and were proposed to aid in damage repair (Zhu 2001). We also detected two transcription factor genes, one homeobox-leucine zipper family protein gene (*HB53*) and one WRKY gene, *WRKY2*. Interestingly, we detected three class III peroxidase genes (*PER10*, *PER52* and *PER59*) in the down-regulated gene list (Table 4-2), which suggests that there was an oxidative burst in plants under salinity condition. To validate the microarray data, we used qRT-PCR to measure transcript abundance of five selected genes. The transcript abundance data obtained by microarray analysis corresponded well with those obtained by qRT-PCR (Fig. 4-10).




Fig. 4-10. qRT-PCR validation of selected genes from microarray results. Data were from 3 biological replicates with the error bars representing S.D.

We further inspected 1 kb of sequence upstream of the predicted translational start codon of each gene (Table 4-2). We found that *cis*-element G-box (CACGTG) occurs in four of the nine up-regulated gene promoters and only one in ten of the down-regulated genes (Table 4-2). However, for E-box (CANNTG), these numbers increased

to be eight and seven for the up- and down-regulated datasets, respectively (Table 4-2). We propose that the other genes whose 1 kb upstream regions do not contain either G- or E-box may be indirect targets of bHLH92.

Table 4-2. Genes that are regulated by 150 mM NaCl in *bhlh92-1* mutant.

AGI	Putative identity	FC	S.D.	G-box	E-box
Up-regulated					
At1g20450	dehydrin (ERD10/LTI45)	1.70	0.35	+	+
At2g25625	auxin-regulated protein	1.68	0.26	-	+
At1g16850	expressed protein	1.66	0.30	+	+
At4g24411	Unknown protein	1.58	0.36	-	-
At5g66700	homeobox-leucine zipper family protein (HB53)	1.57	0.23	-	+
At4g33550	protease inhibitor/seed storage/lipid transfer protein (LTP) family	1.56	0.25	-	+
At5g15960	stress-induced protein KIN1	1.53	0.18	+	+
At5g56270	WRKY family transcription factor WRKY2	1.53	0.10	+	+
At3g62740	putative beta-glucosidase mRNA, complete cds	1.50	0.37	-	+
Down-regulated					
At4g18250	receptor serine/threonine kinase	0.67	0.22	-	-
At5g19890	peroxidase, putative(PER59)	0.67	0.08	-	+
At3g54120	reticulon family protein (RTNLB12)	0.66	0.22	+	+
At5g24540	glycosyl hydrolase family 1	0.65	0.20	-	+
At4g21920	hypothetical protein	0.65	0.17	-	-
At4g12545	protease inhibitor/seed storage/lipid transfer protein (LTP) family	0.64	0.08	-	+
At4g12550	protease inhibitor/seed storage/lipid transfer protein (LTP) family(AIR1)	0.64	0.08	-	-
At5g05340	peroxidase, putative(PER52)	0.64	0.22	-	+
At5g35940	jacalin lectin family	0.63	0.16	-	+
At1g49570	peroxidase, putative(PER10)	0.49	0.15	-	+

AGI, *Arabidopsis* genome initiative; fold change (FC) is expressed as the mRNA level of *bhlh92* mutant/WT with S.D. representing standard deviation; the presence or absence of G-box or E-box in the 1 kb upstream regions is represented by + or -, respectively.

Discussion

Transcriptional regulation of gene expression is one of the fundamental processes that occurs in response to a variety of signals. Although originally identified to be involved in light response, members of bHLH gene family have been found to play a role in quite distinct biological processes including abiotic stress. For instance, MYC2/JIN1 /bHLH6 has been found to be involved in the dehydration, ABA and JA response

pathway (Abe *et al.* 1997, 2003; Lorenzo *et al.*, 2004). ICE1/bHLH116 regulates the transcription of CBF/DREB1 genes in the cold stress and therefore is an upstream component of the CBF/DREB1 cold-signal transduction pathway (Chinnusamy *et al.*, 2003). We reported here the functional characterization of a salt-induced *bHLH92* gene in the context of abiotic stress response and tolerance.

The transcript of *bHLH92* predominantly accumulated in *Arabidopsis* flower buds and young siliques through qRT-PCR analysis (Fig. 4-3). Compared to the GUS staining pattern (Fig. 4-6B-E), there is difference between mRNA accumulation patterns and GUS expression pattern. This inconsistency between mRNA and reporter activity might be due to the fact that we fused only a 1 kb upstream fragment to the *GUS* reporter gene, and therefore, some *cis*-element(s) may exist beyond the 1 kb upstream region. However, using promoter-GUS fusion, we detected the inducibility of GUS expression by salt, drought and JA (Fig. 4-6J-M), which is consistent with the qRT-PCR profiling result (Fig. 4-4B), suggesting that the 1 kb upstream region contains the necessary *cis*-elements that confer stress response.

ABA is a stress hormone extensively involved in response to abiotic stresses such as salt and drought. At least four systems exist in the regulation of salt- and osmotic stress-inducible genes in *Arabidopsis thaliana* (Shinozaki & Yamaguchi-Shinozaki 1997; Shinozaki *et al.*, 2003). Although most salinity-inducible genes are regulated within an ABA-dependent pathway, some are not dependent on ABA for signalling. A conserved *cis*-acting element, referred to as DRE/CRT, functions in stress-inducible gene expression in an ABA-independent manner (Shinozaki *et al.*, 2003). In ABA-dependent pathways, the ABRE (ABA-responsive element) is a major *cis*-acting element in stress-responsive gene expression (Shinozaki *et al.*, 2003). Genetic analysis of two *Arabidopsis* ABA-deficit mutants, *aba1/los6* and *aba3/los5*, in which genes encoding molybdenum cofactor sulfurase (MCSU) and zeaxanthin epoxidase (ZEP) are dysfunctional, showed that ABA is necessary for the salt-stress induced expression of some late-embryogenesis-abundant (LEA) genes (Xiong *et al.*, 2001, 2002a). Our results showed that salt-induced *bHLH92* is partially dependent on the ABA biosynthesis (Fig. 4-5). ABA responsive *cis*-elements (ABREs) are found in the putative *bHLH92* promoter region, but *bHLH92* gene expression is not induced by

ABA treatment according to our criteria (Fig. 4-4B), suggesting a complex mechanism underlying regulating *bHLH92*. It has been shown that a single copy of ABRE is not sufficient for ABA-mediated gene expression, but multiple ABREs or the combination of an ABRE with CE could confer ABA responsiveness to a minimal promoter (Shen *et al.*, 1996; Hobo *et al.*, 1999). We observed that *bHLH92*-overexpressing plants were more sensitive to moderate ABA treatment (Fig. 4-9C), suggesting that ABA is included in the pathway that *bHLH92* works. However, we observed that the null mutant is not more tolerant of ABA (Fig. 4-8C, D). The fact that knock-out mutant and overexpressing did not always give opposite results implies the functional redundancy between *bHLH92* and other *bHLH* TF(s) in some respects or indirect activation of gene expression in the overexpressing plants.

Analysis of *salt overly sensitive (sos)* *Arabidopsis* mutants revealed a novel calcium-regulated protein kinase pathway for response to the ionic aspect of salt stress (Zhu 2003). Genetic and physiological data indicate that *SOS1*, *SOS2*, and *SOS3* function within the same pathway, in which the *SOS3*–*SOS2* complex regulates the expression level of *SOS1* and activates the transport activity of *SOS1*, which leads to Na^+ tolerance (Qiu *et al.* 2002; Zhu 2003). In addition, salt stress response was also shown to be mediated by *SOS*-independent signaling pathways with the identification of several protein kinases activated by salt or osmotic stress (Zhu 2001). In *Arabidopsis* seedlings, a 40-kDa kinase similar to *ASK1* (serine/threonine kinase 1) was identified as being rapidly activated by hyperosmotic stress in a calcium- and ABA-independent manner. Osmotic stress activation of the *Arabidopsis* 40-kDa protein kinase is not impaired by the *sos3* mutation, implying that its activation is independent of the *SOS* pathway (Hoyos and Zhang, 2000). Our results showed that salt-induced *bHLH92* transcript accumulation was not significantly different from between *sos2-1* and *sos3-1* mutants and WT (Fig. 4-5), but was higher in *sos2-1* mutant than WT, suggesting mutation in *SOS2* affect that salt-responsive *bHLH92* expression. Previous microarray profiling of *sos2-1* and *sos3-1* under 250 mM NaCl treatment revealed that *SOS3/SOS2* signalling pathway is independent of the *DRE/CRT*, *ABRE*, and *MYC/MYB* pathways and *SOS2* affects the regulation of stress-responsive genes much more strongly than *SOS3* (Kamei *et al.*, 2005).

In our experiment, we noted that *bHLH92* was induced by MeJA (Fig 5-4). Not surprisingly, we identified four MeJA-responsive cis-elements in the 1 kb upstream of *bHLH92* gene (Table 4-1). JAs modulate a number of vital physiological processes, including defense against pathogens and insects, wound responses, secondary metabolite biosynthesis, and flower development and fertility (Cheong and Choi, 2003). An *Arabidopsis* bHLH TF, MYC2/JIN1, has also been reported to play major roles in both ABA and JA signaling pathways (Abe *et al.*, 2003; Lorenzo and Solano, 2005; Dombrecht *et al.*, 2007).

Through utilizing T-DNA insertional mutants, we found that *bhlh92* mutant was more sensitive to mannitol treatment compared to WT control, while it did not show any sensitivity to salt stress (Fig. 4-8 C, D). Mannitol can cause osmotic stress in plant cells, making plants lose water to the extracellular environment, whereas high salinity can cause both ionic and osmotic stresses in plant cells (Munns 2005). Microarray profilings have found that many genes are responsive to both mannitol and salt treatment (Kreps *et al.*, 2002), suggesting that some signaling components are shared between them. It is widely believed that there is cross-talk between abiotic stresses (Zhu 2001, 2002; Seki *et al.*, 2003). The observation that *bhlh92* mutant was not significantly more sensitive to NaCl treatment suggests a functional redundancy exists between members of bHLH family. Through phylogenetic tree analysis, we propose that bHLH95 may replace bHLH92 in case bHLH92 is not functional. However, bHLH95 transcript abundance was not responsive to salt treatment.

Overexpression of TFs offers an alternative and complementary strategy to knockout analysis as it is less affected by functional redundancy. In plants, there are several examples in which knockouts of TF genes failed to generate informative phenotypes, but overexpression of these TFs revealed gene functions (Zhang 2003). Overexpression of *bHLH92* using the constitutive CaMV35S promoter slightly increased plant tolerance to salt, mannitol and oxidative stresses (Fig. 4-9), suggesting that bHLH92 is a positive regulator of salt and osmotic stress responses.

Using microarrays, we detected nine up- and ten down-regulated genes in the *bhlh92* mutant (Table 4-2). Among the nine up-regulated genes, *ERD10* and *KINI* are two well characterized genes involved in abiotic stress response (Kurkela and Franck,

1990; Gilmour *et al.*, 1992; Kiyosue *et al.*, 1994; Weilin *et al.*, 1995). *ERD10* and *KINI* encode proteins very similar to late embryogenesis abundant (LEA) proteins, and they may play a protecting role under desiccation condition (Wise and Tunnacliffe, 2004), although direct evidence for this is still lacking. As *bhlh92-1* mutant plants were only slightly more sensitive to salt treatment (Fig. 4-8 C, D), the increased abundance of *ERD10* and *KINI* in *bhlh92-1* mutant may indicate that bHLH92 is a negative regulator of some downstream LEA-like gene expression. The homeobox-leucine zipper family protein gene (*HB53*) is auxin-inducible and may be involved in root development (Son *et al.*, 2005). We identified one WRKY gene, *WRKY2*, which was up-regulated by salt in *bhlh92* (Table 4-2). In WT plant roots, *WRKY2* was not responsive to salt treatment (Jiang and Deyholos, 2006). Further examination showed that *WRKY2* promoter region contain one G-box *cis*-element (data not shown), suggesting *WRKY2* could be a target of bHLH92.

Among the ten genes that were down-regulated genes in the absence of *bHLH92*, three genes (*PER10*, *PER52* and *PER59*) were members of the class III peroxidase gene family were down-regulated (Table 4-2). Class III peroxidases are plant-specific oxidoreductases that are implicated in various physiological processes such as H₂O₂ detoxification, auxin catabolism, lignification, suberization, stress response (wounding, pathogen attack, NaCl) and senescence (Hiraga *et al.*, 2001). We previously reported that the majority of class III peroxidases are responsive, at the transcript level, to NaCl treatment in *Arabidopsis* roots (Jiang and Deyholos, 2006). However, the specific functions of these PER genes remain to be determined. Furthermore, three protease inhibitor/seed storage/lipid transfer protein (LTP) genes were found to be differentially expressed (Table 4-2). LTPs form a multigenic family whose gene expression in the different plant organs is temporally and spatially regulated (Marion *et al.*, 2007). LTPs can inhibit the growth of fungal pathogens *in vitro* and are also involved in the signalling of defence mechanism of plants against their pathogens (Blein *et al.*, 2002). However, the exact function of LTPs in abiotic stress has not been reported.

More recently, a tobacco (*Nicotiana tabacum*) plastid-resident basic helix-loop-helix protein-NtWIN4 (*N. tabacum* wound-induced clone 4), was found to be converted from a repressor of nuclear transcription into a plastid-resident regulatory factor

through replacement of the DNA-binding domain with a plastid transit sequence during evolution (Kodama and Sano, 2006, 2007). It was also found that bHLH92 protein can be relocated to the chloroplast when the first 30 amino acids are artificially removed (Kodama and Sano, 2006). Therefore, NtWIN4 uncovered a new evolutionary mechanism, although this mechanism is adopted in other cases and species remains unknown (Kleine and Leister, 2007).

In summary, this study shows that bHLH92 plays an important role in salt and osmotic stress responses. This strengthens our understanding of transcriptional regulation during salt and other types of abiotic stress responses in higher plants and provides a base for future work in this area. Future study can be directed to identify upstream gene(s) regulating bHLH92 activity and its interacting partners.

References

- Abe H, Yamaguchi-Shinozaki K, Urao T, Iwasaki T, Hosokawa D and Shinozaki K. (1997)** Role of *Arabidopsis* MYC and MYB homologs in drought and abscisic acid-regulated gene expression, *Plant Cell* 9, 1859–1868
- Abe H, Urao T, Ito T, Seki M, Shinozaki K and Yamaguchi-Shinozaki K. (2003)** *Arabidopsis* AtMYC2 (bHLH) and AtMYB2 (MYB) function as transcriptional activators in abscisic acid signaling. *Plant Cell* 15, 63–78
- Alonso JM, Stepanova AN, Lisse TJ, Kim CJ, Chen H, Shinn P, Stevenson DK et al. (2003)** Genome-wide insertional mutagenesis of *Arabidopsis thaliana*. *Science* 301: 653-657
- Anderson JP, Badruzsaufari E, Schenk PM, Manners JM, Desmond OJ, Ehlert C, Maclean DJ, Ebert PR and Kazan K. (2004)** Antagonistic interaction between abscisic acid and jasmonate-ethylene signaling pathways modulates defense gene expression and disease resistance in *Arabidopsis*. *Plant Cell* 16:3460–3479
- Atchley WR and Fitch WM. (1997)** A natural classification of the basic helix-loop-helix class of transcription factors. *Proc Natl Acad Sci U S A.* 94:5172-5176
- Atchley WR, Terhalle W and Dress A. (1999)** Positional dependence, cliques, and predictive motifs in the bHLH protein domain. *J Mol Evol* 48:501–516
- Bailey PC, Martin C, Toledo-Ortiz G, Quail PH, Huq E, Heim MA, Jakoby M,**

- Werber M and Weisshaar B.** (2003) Update on the basic helix-loop-helix transcription factor gene family in *Arabidopsis thaliana*. *Plant Cell* 15:2497-2502
- Blein JP, Coutos-Thévenot P, Marion D and Ponchet M.** (2002) From elicitors to lipid transfer proteins: a new insight in cell signaling involved in plant defence mechanism, *Trends Plant Sci* 7, 293–296
- Boter M, Ruiz-Rivero O, Abdeen A and Prat S.** (2004) Conserved MYC transcription factors play a key role in jasmonate signaling both in tomato and *Arabidopsis*. *Genes Dev* 18:1577–1591
- Buck MJ and Atchley WR.** (2003) Phylogenetic analysis of plant basic helix loop-helix proteins. *J Mol Evol* 56: 742–750
- Chen D-H and Ronald PC.** (1999) A rapid DNA miniprep method suitable for AFLP and other PCR applications. *Plant Mol Biol Rep* 17: 53-57
- Chen W, Provart NJ, Glazebrook J, Katagiri F, Chang H-S, Eulgem T, Mauch F, Luan S, Zou G, Whitham SA, Budworth PR, Tao Y, Xie Z, Chen X, Lam S, Kreps JA, Harper JF, Si-Ammour A, Mauch-Mani B, Heinlein M, Kobayashi K, Hohn T, Dangl JL, Wang X and Zhu T.** (2002) Expression profile matrix of *Arabidopsis* transcription factor genes suggests their putative functions in response to environmental stresses. *Plant Cell* 14: 559-574
- Cheong JJ and Choi YD.** (2003) Methyl jasmonate as a vital substance in plants. *Trends Genet* 19: 409–413
- Chinnusamy V, Ohta M, Kanrar S, Lee BH, Hong X, Agarwal M and Zhu JK.** (2003) ICE1: a regulator of cold-induced transcriptome and freezing tolerance in *Arabidopsis*. *Genes Dev* 17:1043-1054
- Chinnusamy V, Jagendorf A and Zhu J-K.** (2005) Understanding and improving salt tolerance in plants. *Crop Sci* 45: 437-448
- Chiu W, Niwa Y, Zeng W, Hirano T, Kobayashi H and Sheen J.** (1996) Engineered GFP as a vital reporter in plants. *Curr Biol* 6, 325-330
- Clough SJ and Bent AF.** (1998) Floral dip: A simplified method for *Agrobacterium*-mediated transformation of *Arabidopsis thaliana*. *Plant J* 16, 735–743
- Dombrecht B, Xue GP, Sprague SJ, Kirkegaard JA, Ross JJ, Reid JB, Fitt GP, Sewelam N, Schenk PM, Manners JM and Kazan K.** (2007) MYC2 differentially

modulates diverse jasmonate-dependent functions in *Arabidopsis*. *Plant Cell* 19:2225-2245

Duck PD and Fankhauser C. (2005) bHLH class transcription factors take centre stage in phytochrome signalling. *Trends Plant Sci* 10:51-54

Finkelstein RR and Rock CD. (2002) Abscisic acid biosynthesis and response. In Somerville CR, Meyerowitz EM, eds, *The Arabidopsis Book*. American Society of Plant Biologists, Rockville, MD

Friedrichsen DM, Nemhauser J, Muramitsu T, Maloof JN, Alonso J, Ecker JR, Furuya M, and Chory J. (2002) Three redundant brassinosteroid early response genes encode putative bHLH transcription factors required for normal growth. *Genetics* 162, 1445–1456

Gilmour SJ, Artus NN and Thomashow MF. (1992) cDNA sequence analysis and expression of two cold-regulated genes of *Arabidopsis thaliana*. *Plant Mol Biol* 18:13-21.

Hadjeb N, Berkowitz GA. 1996. Preparation of T-overhang vectors with high PCR product cloning efficiency. *BioTechniques* 20, 20-22

Heim MA, Jakoby M, Werber M, Martin C, Weisshaar B, and Bailey PC. (2003) The basic helix-loop-helix transcription factor family in plants: A genome-wide study of protein structure and functional diversity. *Mol Biol Evol* 20, 735–747

Hiraga S, Sasaki K, Ito H, Ohashi Y, and Matsui H. (2001) A large family of class III plant peroxidases. *Plant Cell Physiol* 42: 462-468

Hobo T, Asada M, Kowyama Y and Hattori T. (1999) ACGT-containing abscisic acid response element (ABRE) and coupling element 3 (CE3) are functionally equivalent. *Plant J* 19:679–689

Hoyos ME and Zhang S. (2000) Calcium-independent activation of salicylic acid-induced protein kinase and a 40-kilodalton protein kinase by hyperosmotic stress. *Plant Physiol* 122, 1355–1363

Huq E, and Quail PH. (2002) PIF4, a phytochrome-interacting bHLH factor, functions as a negative regulator of phytochrome B signaling in *Arabidopsis*. *EMBO J* 21, 2441–2450

Jakoby M, Wang HY, Reidt W, Weisshaar B, and Bauer P. (2004) FRU

(BHLH029) is required for induction of iron mobilization genes in *Arabidopsis thaliana*. FEBS Lett. 577:528-534

Jiang Y-Q and Deyholos MK. (2006) Comprehensive transcriptional profiling of NaCl-stressed *Arabidopsis* roots reveals novel classes of responsive genes. BMC Plant Biol 6, 25

Jiang Y-Q, Yang B, Harris NS and Deyholos MK. (2007) Comparative proteomic analysis of NaCl stress-responsive proteins in *Arabidopsis* roots. J Exp Bot 58:3591-3607

Kamei A, Seki M, Umezawa T, Ishida J, Satou M, Akiyama K, Zhu J-K and Shinozaki K. (2005) Analysis of gene expression profiles in *Arabidopsis* salt overly sensitive mutants *sos2-1* and *sos3-1*. Plant Cell Environ 28:1267-1275

Kim J and Kim H-Y. (2006) Functional analysis of a calcium-binding transcription factor involved in plant salt stress signaling. FEBS Lett 580:5251-5256

Kiyosue T, Yamaguchi-Shinozaki K and Shinozaki K. (1994) Characterization of two cDNAs (ERD10 and ERD14) corresponding to genes that respond rapidly to dehydration stress in *Arabidopsis thaliana*. Plant Cell Physiol 35:225-231

Kleine T and Leister D. (2007) Evolutionary tinkering: birth of a novel chloroplast protein. Biochem J. 403:e13-4

Knight H, Trewavas AJ and Knight MR. (1997) Calcium signalling in *Arabidopsis thaliana* responding to drought and salinity. Plant J 12:1067-1078

Kodama Y and Sano H. (2006) Evolution of a basic helix-loop-helix protein from a transcriptional repressor to a plastid-resident regulatory factor: involvement in hypersensitive cell death in tobacco plants. J Biol Chem. 281:35369-36380

Kodama Y and Sano H. (2007) Functional diversification of a basic helix-loop-helix protein due to alternative transcription during generation of amphidiploidy in tobacco plants. Biochem J 403: 493-499

Kreps JA, Wu Y, Chang HS, Zhu T, Wang X and Harper JF. (2002) Transcriptome changes for *Arabidopsis* in response to salt, osmotic, and cold stress. Plant Physiol 130, 2129-2141

Kurkela S and Franck M. (1990) Cloning and characterization of a cold- and ABA-inducible *Arabidopsis* gene. Plant Mol Biol 15:137-144

- Ledent V and Vervoort M.** (2001) The basic helix-loop-helix protein family: Comparative genomics and phylogenetic analysis. *Genome Res* 11, 754–770
- Letunic I, Copley RR, Pils B, Pinkert S, Schultz J and Bork P.** (2006) SMART 5: domains in the context of genomes and networks. *Nucleic Acids Res* 34, D257-D260
- Lescot M, Dehais P, Thijs G, Marchal K, Moreau Y, Van de Peer Y, Rouze P and Rombauts S.** (2002) PlantCARE, a database of plant *cis*-acting regulatory elements and a portal to tools for in silico analysis of promoter sequences. *Nucleic Acids Res* 30:325-327
- Li H, Sun J, Xu Y, Jiang H, Wu X and Li C.** (2007) The bHLH-type transcription factor AtAIB positively regulates ABA response in *Arabidopsis*. *Plant Mol Biol* 65: 655-65
- Lorenzo O, Chico JM, Sanchez-Serrano JJ and Solano R.** (2004) *JASMONATE-INSENSITIVE1* encodes a MYC transcription factor essential to discriminate between different jasmonate regulated defense responses in *Arabidopsis*. *Plant Cell* 16:1938–1950
- Lorenzo O and Solano R.** (2005) Molecular players regulating the jasmonate signalling network. *Curr Opin Plant Biol* 8: 532–540
- Marion D, Bakan B and Elmorjani K.** (2007) Plant lipid binding proteins: properties and applications. *Biotechnol Adv* 25:195-197
- Morgenstern B and Atchley WR.** (1999) Evolution of bHLH transcription factors: Modular evolution by domain shuffling? *Mol Biol Evol* 16, 1654–1663
- Munns R.** (2005) Genes and salt tolerance: bringing them together. *New Phytol* 167: 645-663
- Murre C, McCaw PS, and Baltimore D.** (1989) A new DNA binding and dimerization motif in immunoglobulin enhancer binding, daughterless, MyoD and myc proteins. *Cell* 56, 777–783
- Nesi N, Debeaujon I, Jond C, Pelletier G, Caboche M, and Lepiniec L.** (2000) The TT8 gene encodes a basic helix-loop-helix domain protein required for expression of DFR and BAN genes in *Arabidopsis* siliques. *Plant Cell* 12, 1863–1878
- Ni M, Tepperman JM, and Quail PH.** (1998) PIF3, a phytochrome-interacting factor necessary for normal photoinduced signal transduction, is a novel basic helix-loop-

helix protein. *Cell* 95, 657–667

Payne CT, Zhang F, and Lloyd AM. (2000) *GL3* encodes a bHLH protein that regulates trichome development in *Arabidopsis* through interaction with *GL1* and *TTG1*. *Genetics* 156, 1349–1362

Pfaffl MW. (2001) A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Res* 29: e45

Pillitteri LJ and Torii KU. (2007) Breaking the silence: three bHLH proteins direct cell-fate decisions during stomatal development. *Bioassays* 29:861-870

Qiu QS, Guo Y, Dietrich MA, Schumaker KS, and Zhu J-K. (2002) Regulation of *SOS1*, a plasma membrane Na^+/H^+ exchanger in *Arabidopsis thaliana*, by *SOS2* and *SOS3*. *Proc Natl Acad Sci USA* 99, 8436–8441

Ramsay NA and Glover BJ. (2005) MYB-bHLH-WD40 protein complex and the evolution of cellular diversity. *Trends Plant Sci* 10: 63–70

Rampey RA, Woodward AW, Hobbs BN, Tierney MP, Lahner B, Salt DE and Bartel B. (2006) An *Arabidopsis* basic helix-loop-helix leucine zipper protein modulates metal homeostasis and auxin conjugate responsiveness. *Genetics* 174:1841-1857

Robinson KA, Koepke JI, Kharodawala M, and Lopes JM. (2000) A network of yeast basic helix-loop-helix interactions. *Nucleic Acids Res.* 28, 4460–4466

Salinas J and Sanchez-Serrano JJ. (2006) *Arabidopsis* Protocols: 2nd Ed. Humana Press, NJ, USA

Sambrook J and Russel DW. (2001) *Molecular Cloning: a Laboratory Manual*, 3rd ed. Cold Spring Harbor: Cold Spring Harbor Laboratory Press. New York

Seki M, Kamei A, Yamaguchi-Shinozaki K and Shinozaki K. (2003) Molecular responses to drought, salinity and frost: common and different paths for plant protection. *Curr Opin Biotechnol* 14(2):194-199

Seki M, Narusaka M, Ishida J, Nanjo T, Fujita M, Oono Y, Kamiya A, Nakajima M, Enju A, Sakurai T, Satou M, Akiyama K, Taji T, Yamaguchi-Shinozaki K, Carninci P, Kawai J, Hayashizaki Y and Shinozaki K. (2002) Monitoring the expression profiles of 7000 *Arabidopsis* genes under drought, cold and high-salinity stresses using a full-length cDNA microarray. *Plant J* 31: 279-292

Sessions A, Burke E, Presting G, Aux G, McElver J, Patton D, Dietrich B, Ho P, Bacwaden J, Ko C, Clarke JD, Cotton D, Bullis D, Snell J, Miguel T, Hutchison D, Kimmerly B, Mitzel T, Katagiri F, Glazebrook J, Law M and Goff SA. (2002) A high-throughput *Arabidopsis* reverse genetics system. *Plant Cell* 14:2985-2994

Shen QJ, Zhang P and Ho TH. (1996) Modular nature of abscisic acid (ABA) response complexes: composite promoter units that are necessary and sufficient for ABA induction of gene expression in barley. *Plant Cell* 8:1107–1119

Shinozaki K and Yamaguchi-Shinozaki K. (2000) Molecular responses to dehydration and low temperature: differences and cross-talk between two stress signaling pathways. *Curr Opin Plant Biol* 3:217-223

Shinozaki K and Yamaguchi-Shinozaki K. (1997) Gene expression and signal transduction in water-stress response. *Plant Physiol* 115, 327-334

Shinozaki K, Yamaguchi-Shinozaki K and Seki M. (2003) Regulatory network of gene expression in the drought and cold stress responses. *Curr Opin Plant Biol* 6, 410–417

Singh K, Foley RC, Onate-Sanchez L. (2002) Transcription factors in plant defense and stress responses. *Curr Opin Plant Biol* 5, 430–436

Smolen GA, Pawlowski L, Wilensky SE, and Bender J. (2002) Dominant alleles of the basic helix-loop-helix transcription factor ATR2 activate stress-responsive genes in *Arabidopsis*. *Genetics* 161, 1235–1246

Son O, Cho HY, Kim MR, Lee H, Lee MS, Song E, Park JH, Nam KH, Chun JY, Kim HJ, Hong SK, Chung YY, Hur CG, Cho HT and Cheon CI. (2005) Induction of a homeodomain-leucine zipper gene by auxin is inhibited by cytokinin in *Arabidopsis* roots. *Biochem Biophys Res Commun* 326:203-209

Toledo-Ortiz G, Huq E and Quail PH. (2003) The *Arabidopsis* basic/helix-loop-helix transcription factor family. *Plant Cell* 15:1749-1770

Verslues PE, Agarwal M, Katiyar-Agarwal S, Zhu J and Zhu J-K. (2006) Methods and concepts in quantifying resistance to drought, salt and freezing, abiotic stresses that affect plant water status. *Plant J* 45: 523-539.

Wadsworth GJ, Redinbaugh MG and Scandalios JG. (1988) A procedure for the small-scale isolation of plant RNA suitable for RNA blot analysis. *Anal Biochem*

172:279-283

Weigel D and Glazebrook J. (2002) *Arabidopsis: A Laboratory Manual*. Cold Spring Harbor Laboratory Press, NY, USA

Welin BV, Olson A and Palva ET. (1995) Structure and organization of two closely related low-temperature-induced *dhn/lea/rab*-like genes in *Arabidopsis thaliana* L. Heynh. *Plant Mol Biol* 29:391-395

Wise MJ and Tunnacliffe A. (2004) POPP the question: what do LEA proteins do? *Trends Plant Sci* 9:13-17

Xie M, He Y, Gan S. (2001) Bidirectionalization of polar promoters in plants. *Nat Biotechnol* 19:677-679

Xiong L, Ishitani M, Lee H and Zhu J-K. (2001) The *Arabidopsis LOS5/ABA3* locus encodes a molybdenum cofactor sulfuryase and modulates cold stress- and osmotic stress- responsive gene expression. *Plant Cell* 13:2063-2083

Xiong L, Lee H, Ishitani M and Zhu J-K. (2002a) Regulation of osmotic stress responsive gene expression by *LOS6/ABA1* locus in *Arabidopsis*. *J Biol Chem* 277:8588-8596

Xiong L, Schumaker K and Zhu J-K. (2002b) Cell signaling during cold, drought, and salt stress. *Plant Cell* 14: S165-S183

Yoo SY, Bomblies K, Yoo SK, Yang JW, Choi MS, Lee JS, Weigel D and Ahn JH. (2005) The 35S promoter used in a selectable marker gene of a plant transformation vector affects the expression of the transgene. *Planta* 221:523-530

Yamaguchi-Shinozaki K and Shinozaki K. (1993) Characterization of the expression of a desiccation-responsive *rd29* gene of *Arabidopsis thaliana* and analysis of its promoter in transgenic plants. *Mol Gen Genet* 236(2-3):331-340

Yamaguchi-Shinozaki K and Shinozaki K. (2006) Transcriptional regulatory networks in cellular responses and tolerance to dehydration and cold stresses. *Annu Rev Plant Biol* 57:781-803

Zhang JZ.(2003) Overexpression analysis of plant transcription factors. *Curr Opin Plant Biol* 6:430-440

Zhu J-K.(2001) Cell signaling under salt, water and cold stresses. *Curr Opin Plant Biol* 4: 401–406

Zhu J-K.(2002) Salt and drought stress signal transduction in plants. *Annu Rev Plant Biol* 53, 247-273

Zhu J-K.(2003) Regulation of ion homeostasis under salt stress. *Curr Opin Plant Biol* 6: 441–445

Chapter 5: Functional characterization of NaCl-inducible *WRKY25* and *WRKY33* transcription factor genes

Introduction

Plants have evolved multiple signaling pathways and defence strategies in response to biotic or abiotic stress. Several classes of transcription factors (TFs) have been implicated in plant response to stress, including AP2/ERF, MYB, WRKY and bZIP (Singh *et al.*, 2002). WRKY TFs contain a WRKY zinc-finger motif, which binds to the W-box motif (C/TTGACC/T) (Eulgem *et al.*, 2000; Ülker and Somssich, 2004). The name of the WRKY family itself is derived from the most prominent feature of these proteins, the WRKY domain, constituted by about 60 amino acid residues. In this WRKY domain, a conserved WRKYGQK heptapeptide is followed by a C2H2- or C2HC-type of zinc finger motif (Eulgem *et al.*, 2000). WRKY transcription factor genes form large families in plants, with 72 members in Arabidopsis. These can be divided into three groups with several subgroups on the basis of their WRKY domains and their phylogenetic clades (Eulgem *et al.*, 2000). Although originally thought to be plant-specific, genes encoding WRKY proteins have been found in two non-photosynthetic eukaryotes: the slime mold *Dictyostelium discoideum* and the protist *Giardia lamblia*. These two lineages evolved prior to the divergence of plants from animals and fungi, indicating an ancient origin of WRKY transcription factors (Zhang and Wang 2005).

Abiotic and biotic stresses are major external factors influencing the expression of WRKY genes (Eulgem *et al.* 2000; Ülker and Somssich 2004). In plants, many WRKY proteins are involved in the defence against attack by phytopathogens such as bacteria, fungi and viruses (Asai *et al.*, 2002; Chen *et al.*, 2002; Dellagi *et al.*, 2000; Kalde *et al.*, 2003, Dong *et al.*, 2003; Yang *et al.*, 1999; Yoda *et al.*, 2002). Furthermore, WRKYs have been found to play essential roles in various physiological processes, including embryogenesis, seed coat and trichome development, senescence, regulation of biosynthetic pathways, and hormonal signaling (Ulker and Somssich, 2004; John *et al.*, 2002; Lagace and Matton, 2004; Xu *et al.*, 2004; Zhang *et al.*, 2004; Zou *et al.*, 2004; Xie *et al.*, 2005). WRKY genes have also been found to be responsive

to abiotic stress. For instance, barley WRKY38 is associated with cold- and drought-response (Mare *et al.*, 2004). Induction of *WRKY* genes has also been observed in *Retama raetam* by drought stress (Pnueli *et al.*, 2002) and in *Nicotiana tabacum* by wounding (Hara *et al.*, 2000). More recently, *Arabidopsis* WRKY75 was reported to be involved in regulating phosphate starvation response and root development (Devaiah *et al.*, 2007). The involvement of WRKY factors in these diverse processes is complicated. Characterization of the expression and the role of individual WRKY members is important and challenging considering the large number of WRKY TFs in plants (Ulker and Somssich, 2004). Recent studies have also revealed that there is cross-talk between abiotic and biotic stress responses and several molecules, including transcription factors (e.g. MYB, MYC) and kinases (e.g. MAPK), as promising candidates for common players that are involved in cross-talk between stress signaling pathways (Anderson *et al.*, 2004; Fujita *et al.*, 2006; Nakagami *et al.*, 2005). In *Arabidopsis*, WRKY29 has been identified as an important downstream component of a MAPK pathway that confers resistance to both bacterial and fungal pathogens (Asai *et al.*, 2002). Recently, *Arabidopsis* WRKY70 was also identified as a common regulatory component of salicylic acid (SA)- and JA- dependent defense response, mediating cross-talk between antagonistic pathways downstream of nonexpressor of pathogenesis-related genes 1 (NPR1) (Li *et al.*, 2004).

Arabidopsis MAP kinase 4 (MPK4) is an activator of jasmonic acid (JA) and ethylene (ET)-mediated defense, and is a repressor of SA-dependent resistance. In a previously reported study using yeast two-hybrid screening, MPK4 was found to interact with a substrate, MKS1, which in turn interacts with *Arabidopsis* WRKY25 and WRKY33 (Andreasson *et al.*, 2005). In addition, WRKY25 and WRKY33 were phosphorylated by MPK4 *in vitro*, and a *wrky33* knockout mutant expressed elevated levels of *PR1* (*PATHOGENESIS-RELATED1*) under a short-day growth conditions (Andreasson *et al.*, 2005). These results suggest that WRKY25 and WRKY33 may function as downstream components of the MPK4-mediated, SA-repressing and JA/ET-activating signaling pathways. Zheng *et al.* (2006) demonstrated that disruption of *WRKY33* results in enhanced susceptibility to necrotrophic fungal pathogens and impaired expression of JA/ET-regulated defense genes. No such phenotypes were

observed in the *wrky25* T-DNA insertion mutants. These results indicate that WRKY33 functions as a positive regulator of JA/ET-mediated pathways and plays an important role in disease resistance to necrotrophic fungal pathogens. More recently, Lippok *et al* (2007) found that rapid, pathogen-induced WRKY33 expression depends on PAD4 (PHYTOALEXIN-DEFICIENT 4), a key regulator upstream of SA, although it does not require salicylic acid (SA) signaling. In addition, WRKY25 functions as a negative regulator of SA-mediated defense responses to the bacterial pathogen *P. syringae* (Zheng *et al.*, 2007).

In our previous microarray study, we identified 18 WRKY TFs, including *WRKY25* (At2g30250) and *WRKY33* (At2g38470) that were induced by 150 mM NaCl treatment in *Arabidopsis* roots (Jiang and Deyholos 2006). This observation emphasizes a potentially important role for WRKY transcription factors in mediating salt stress responses. In this study, we investigated the function of *WRKY25* and *WRKY33* as regulators of salt stress responses. We analyzed their knock-out mutant and overexpression plant phenotypes, which suggest their role in salt, osmotic and ABA signalings.

Materials and methods

Plant growth, stress treatments and mutant isolation

Arabidopsis seeds, either wild-type (Col-0) or mutants, were sown and grown as described in Chapter 4. Leaves, stems, flower buds, young siliques and seeds were harvested from WT *Arabidopsis* (Col-0) plants for qRT-PCR analysis. Hydroponics of *Arabidopsis* was performed as previously reported (Jiang and Deyholos, 2006). At 20 DAS (days after stratification), seedlings were treated with various stress treatments as described in Chapter 4. All treatments lasted for 6 and 24 h with roots and rosette leaves harvested and pooled separately; control samples were harvested at the same time. Samples were flash frozen in liquid nitrogen and kept at -80°C until use.

Arabidopsis mutant seeds, *aba1-5*, *abi4-1*, *sos1-1*, *sos2-1* and *sos3-1* were obtained from *Arabidopsis* Biological Resource Center (ABRC, Ohio, USA) and hydroponically cultured as above for 20 d with roots harvested for qRT-PCR analysis. T-DNA insertion mutants (bulk T₄) developed at the Salk Institute (Alonso *et al.*, 2003)

and Syngenta (Sessions *et al.*, 2002) were obtained from ABRC. The homozygous T-DNA insertion lines for each gene were identified by PCR with primers designed according to <http://signal.salk.edu/tdnaprimers.2.html>. The primer sequences are as follows: LBb1 for SALK lines, 5'-GCGTGGACCGCTTGCTGCAACT-3'; 5'-GTCACAACAATC-CGGAAGAAC-3' (LP) and 5'-CAAACCTGGCATTGTACA CAGC-3' (RP) for SALK_006603; 5'-GTTGAGATTGTAGCTGCCAGG-3' (LP) and 5'-AGAATGA-TCGACGCAAAACC-3'(RP) for SALK_136966; LB3 for SAIL line, 5'-TAGCAT-CTGAATTTTCATAACCAATCTCGATACAC-3'; 5'-TGTTTGGTT GATCA-TCTTTCCTC-3' (LP) and 5'-GGAGGAACAACCTGAGATGAAG-3' (RP) for SAIL_529_B11.

Genomic DNA was extracted from 4-week-old rosette leaves of T₄ lines for each T-DNA insertion mutant as well as from WT plants by 2% CTAB method (Chen and Ronald, 1999). PCR was conducted using two combinations of primers (LBb1/RP and LP/RP) as per the protocol described in <http://signal.salk.edu/tdnaprimers.2.html>. To confirm the putative homozygous T-DNA insertion lines, total RNA was extracted from ~ 4 week old rosette leaves according to Wadsworth *et al.* (1988) with modifications, and dissolved in 20 µl RNase-free water. The extracted RNA was treated with DNaseI (Ambion), followed by reverse transcription using oligo(dT)₁₂₋₁₈ and SuperScriptII (Invitrogen), followed by PCR using gene-specific primers (see below). PCR products were cloned into a pBS SK+ (Stratagene)-based T vector and sequenced using T7 and T3 promoter primers (Invitrogen). General molecular biology experiments were performed according to Sambrook and Russel (2001).

Quantitative RT-PCR

Total RNA was extracted from ~100 mg of tissue using the RNeasy Plant Mini Kit (Qiagen). Dry seed RNA was extracted according to Suzuki *et al.*(2004). qRT-PCR was performed as described in Chapter 4. Data were analyzed in Sequencing Detection System (v1.4) and Microsoft Excel 2003.

The primers used for measuring transcript abundance were as follows: *RD29A* (At5g52310), Fwd, 5'-AGGAAGTGAAAGGAGGAGGAGGAATG-3', Rvs, 5'-GTCGCACCACCACCAAACCAG-3'; *WRKY25* (At2g30250), Fwd, 5'-TGGTTCTTC

CGGCGTTGACTGTTA-3', Rvs, 5'-GTGAAATCGGAAGAGGTGG-TGGTTG-3'; *WRKY33* (At2g38470), Fwd, 5'-TTACGCCACAAACAGAGCAC-3', Rvs, 5'-CCAAAAGGCCCGGTATTAGT-3'; *UBQ10* (At4g05320), Fwd, 5'-GGCCTTGTATAATCCCTGATGAA-3', Rvs, 5'-AGAAGTTCGACTTGTCATTA-GAAA GAAA-3'. qRT-PCR for each gene was assayed on three biological replicates with duplicates or triplicates for each biological replicate. The relative transcript abundance was determined for each sample, and was normalized using the UBQ10 cDNA level as described in Chapter 4.

Plasmid construction and plant transformation

Total RNA was extracted from *Arabidopsis* roots according to Wadsworth *et al.* (1988) with modifications. After treated with DNaseI (Ambion), 2 µg of RNA was reverse transcribed using SuperScriptII (Invitrogen) and Oligo(dT)₁₈ (Invitrogen). All primers were manually designed and restriction sites were analyzed using DNAMAN (v4.0, Lynnon Biosoft). For subcellular localization of TFs, the coding regions (CDS) of *WRKY25* (1182 bp) and *WRKY33* (1560 bp) were amplified by RT-PCR using the following primers: 5'-CATG CCATGGAT TCTTCCACTTCTTTCACCGA-3' (Fwd) and 5'-CATGCCATGGCAGCTCCACCTCCACCTCCCGAGCGACGTAGCGCGGT TGG-3' (Rvs) for *WRKY25*, 5'-CATG CCATGGCTGCTTCTTTTCTTACA-3' (Fwd) and 5'-TTAGGTACATGACAGCTCCACCTCCACCTCCGGGCATAAACGA ATCGAA AAA-3'(Rvs) for *WRKY33*, 5'-CATG CCATGGCGGCGACATCTTCC TCT-3'(Fwd) and 5'-TTAGGTACATGATAGCTCCACCTCCACCTCCAAGCTC GGGCTGCATTTGCTC-3' (Rvs) for *Arabidopsis ACS1*(At2g43750). The CDS regions were amplified from WT cDNAs using *Pfx* DNA polymerase (Invitrogen) and PCR products were purified using a Qiagen PCR purification kit, restricted by *Nco* I and/or *Bsp* HI, purified again and cloned into *Nco* I restricted pCsGFPBT with a Gly-Ala- rich peptide linker between CDSs and sGFP.

For promoter-GUS fusions, we started with binary vector pCAMBIA1391Z and modified it to be p0381Z as described in Chapter 4. The promoter regions of *WRKY25* and *WRKY33* genes were PCR-amplified using WT genomic DNA as the template with the following primer pairs: 5'-CCCAAGCTTCTACTTCAAATACATATATAC-3' (Fwd) and 5'-CATG CCATGG ACG AAA AAT GGA AGT TTG TTT-3' (Rvs) for

WRKY33 promoter, 5'-AGGGATCCCACTTGACGATTTATCAC-3' (Fwd) and 5'-CGAATTCGATGGTCTTTAATAAAGGAG-3' (Rvs) for WRKY25 promoter. The resulting fragments were digested with *Hind* III and *Nco* I and inserted into the p0381Z binary vector.

For overexpression, coding regions of *WRKY25* and *WRKY33* were cloned by RT-PCR using primers 5'-CATG CCATGGTAATGTCTTCCACTTCTTTTCAC-3' (Fwd) and 5'-TCGG GGTCACC TCACGAGCGACGTAGCGCGG-3' (Rvs) for *WRKY25*, 5'-CATG CCATGGCTGCTTCTTTTCTTACA-3' (Fwd) and 5'-TCGG GGTCACC TCAGGGCATAAACGAATCGAA-3' (Rvs) for *WRKY33*. PCR amplifications were conducted using high-fidelity *Pfx* DNA polymerase (Invitrogen) and, PCR products were purified by a Qiagen PCR purification kit, restricted with *Nco*I and *Bst* EII, and purified again using a Qiagen kit before ligated into *Nco* I and *Bst* E II digested of pCAMBIA1303 in place of *GUS-GFP*. All the inserts were sequenced using BigDye reagent (Applied BioSystems) and analyzed using DNASTar (DNASTAR Inc) to verify that no mismatches or frame-shifts had occurred.

All constructs plus controls were mobilized into *Agrobacterium* GV3101 through the freeze-thaw method and transformed into wild-type *Arabidopsis thaliana* (Col-0) by floral dip (Clough and Bent, 1998). Selection of hygromycin resistant lines was described in Chapter 4.

GFP observation and GUS staining

GFP signals in 3-5 d old transgenic seedlings from ten independent T₂ lines were observed under either an epifluorescent or confocal microscope as described in Chapter 4. GUS staining of T₃ lines was conducted as described in Chapter 4. Representative and consistent results were reported. All images were assembled in Adobe Photoshop CS (Adobe Systems).

Stress tolerance tests

Stress tolerance tests of mutants and 2-3 independent homozygous overexpression lines (T₃) compared to WT were performed according to Verslues *et al.* (2006) with modifications, which were reported in Chapter 4. The relative electrolyte leakage (REL) assay was performed according to Jiang *et al.* (2007). Each assay was repeated two to three times with similar results obtained. Data were analyzed using two-tailed

Student's *t*-test.

Microarray profiling, data analysis and validation

Qiagen Operon 26 k *Arabidopsis* oligonucleotide microarrays were used to profile transcriptomes of *wrky25-1* and *wrky33-1* T-DNA insertion mutants using WT as the reference. Total RNA samples from hydroponically cultured roots treated by 150 mM NaCl for 6 h with three independent biological replicates were extracted using the RNeasy Plant Mini kit (Qiagen). Possibly contaminating DNA was removed using the DNA-free kit (Ambion), and 5 µg of total RNA was reverse transcribed into cDNAs using RevertAid H- MMLV (Fermentas) and primers contained in the Array900 kit (Genisphere). Microarray hybridizations with dye-flip design were performed using the reagents contained in the Array900 kit according to the manufacturer's manual. Altogether six microarray slides were used with 12 image files analyzed. Data were analyzed as described in Jiang and Deyholos (2006).

To validate microarray data, qRT-PCR was performed with selected genes using the same batch of DNA-free RNA used for microarray hybridizations. The primers designed by PrimerExpress3.0 (Applied Biosystems) were: At5g05340 (*PER52*), Fwd, 5'-TCACCACGGCAGCTTCTTTC-3', Rvs, 5'-GTAGAGCCGCCGTTGAAGAG-3'; At5g24540 (glycosyl hydrolase family 1), Fwd, 5'-ACGGTTTGTGCTCACGAT-3', Rvs, 5'-GTGTCGCCGGTACGATTTGT-3'; At4g23170 (protein kinase family protein), Fwd, 5'-AACGTCACGGTGGGACAAAC-3', Rvs, 5'-GAGAAGGCAACG CAGTTA-3'; At1g12610 (*DDF1/DREB1F*), Fwd, 5'-CGGAGGAACGGTGACAAA TG-3', Rvs, 5'-CACGCGCTGCCATATCTG-3'; At1g76680 (*OPRI*), Fwd, 5'-GGAA TGAGGCTGTGTCAAAGG-3', Rvs, 5'-GGTGCATCCACTTGGAACCT-3'; At2g46830 (*CCA1*), Fwd, 5'-GTCATCCCCACACAGTTGCA-3', Rvs, 5'-GCGAAA GTTGCGGCAGTATAA-3'; At3g14225 (*GLIP4*), Fwd, 5'-TGAGCTTGGGTCCATT TGGT-3', Rvs, 5'-TGGAGATTGATCAGCTCTGTGACT-3'; At4g25470 (*CBF2 /DREB1C*), Fwd, 5'-CCGGTTTCCTCAGGCGGTGATTACA-3', Rvs, 5'-TCCTCTG TAAATTGGGTGACG-3'; At3g62720, Fwd, 5'-TCCGAGAAGAAGCGGGTAAA-3', Rvs, 5'-TCCGTCGCCAGCAGATAAAC-3'; At2g34100, Fwd, 5'-TGTCGCGCG AGCAGAGAA-3', Rvs, 5'-GCAATTGTCTCTCGCCTCACA-3'; At4g12480, Fwd, 5'-GCACTGCGCTTAGGGCTAAC-3', Rvs, 5'-AAGCCAGACGGAACCTTTCTG-

3'; At4g10270, Fwd, 5'-GTGAGCATCGGAGCCGTAGA-3', Rvs, 5'-TAACGTTG TTCCGGAGATGTTG-3'; At5g10040, Fwd, 5'-GGAGGTTGGGTGGTTGATGA-3', Rvs, 5'-AGAAACCGCCACAGCAAAC-3'. qRT-PCR assay was performed using SYBR Green I fluorescent dye in ABI7500 fast real-time PCR machine as described in Chapter 4.

Analysis of promoter motifs in co-regulated gene sets.

The sequence of a 1,000-bp region immediately upstream (5') of each gene's predicted translational start was scanned using the Motif Analysis program of TAIR (<http://www.arabidopsis.org/tools/bulk/motiffinder/index.jsp>). The frequency of all possible hexameric motifs in these 5' regions was queried and compared with the frequencies of the same motifs in the total genomic set of 1000-bp 5' regions. Motifs whose analysis yielded a *P* of the null hypothesis (no difference from the total genomic set) smaller than $1 \exp - 4$ were deemed significant.

Sequence and phylogenetic analysis

The predicted protein sequences of 72 members of the *Arabidopsis* WRKY family were retrieved from the Database of *Arabidopsis* Transcription Factors (DATF, (<http://datf.cbi.pku.edu.cn/>)) and Riken *Arabidopsis* Transcription Factors (RARTF, <http://rarge.gsc.riken.jp/rartf/>). The WRKY DNA binding domain boundary was defined as per Eulgem *et al* (2000) and only amino acid sequences from the single WRKY domain of group II and III members or the C-terminal WRKY domain of group I members was used, except WRKY19, for which, the N-terminal WRKY domain sequence was used because the C-terminal WRKY domain does not contain the typical WRKYGQK heptapeptide. Sequences were aligned according to Hall (2001) using ClustalX (v1.83) (Thompson *et al.*, 1997) with a gap opening penalty of 35 and gap extension penalty of 0.75 in pairwise alignment, and a gap opening penalty of 15 and gap extension penalty of 0.30 in multiple alignment parameter settings. Multiple alignment was further checked with gaps manually inserted for optimal alignment. The parsimony tree was drawn using PAUP*4.0 (Sinauer Associates, Inc.). For Fig. 5-1, a postscript file (.ps) generated from ClustalX (v1.83) was read by Adobe Acrobat 7.0 professional with the PDF format file output.

Results

Identification and phylogenetic analysis of *WRKY25* and *WRKY33* genes

The majority of WRKY family TF genes are known to be responsive to biotic and/or abiotic stress, although most research has focused on the role of these genes in pathogen responses (Eulgem *et al.*, 2000). In our previously reported microarray data, we identified 18 WRKY TFs, including *WRKY25* (At2g30250) and *WRKY33* (At2g38470) that were induced by 150 mM NaCl treatment in *Arabidopsis* roots (Jiang and Deyholos 2006). This observation emphasizes a potentially important role for WRKY transcription factors in salt stress response. We chose to characterize *WRKY25* and *WRKY33* genes as their physiological role in abiotic stress has not been reported.

WRKY proteins are divided into three main classes based on sequence similarity, and additional structural motifs allow the distinction of different subgroups (Eulgem *et al.*, 2000). *WRKY25* and *WRKY33* are group Ia proteins, which contain two WRKY domains followed by a Cys2/His2-type zinc-finger motif (Fig. 5-1). *WRKY25* (At2g30250) encodes a protein of 393 amino acids with a molecular weight of 44.1 kD and an isoelectric point of 6.43, while *WRKY33* (At2g38470) encodes a protein of 519 amino acids with a molecular weight of 57.2 kD and an isoelectric point of 7.89 (Fig. 5-1). *WRKY25* shares an overall identity of 28.90% at the amino acid level with the *WRKY33* protein.

Phylogenetic analysis of genes is important for the functional analysis of a gene family. Members of transcription factor gene families usually contain a highly conserved domain or more than one conserved domain involved in DNA binding. However, sequence similarity beyond the domains is often relatively low. Furthermore, a domain is considered as an evolutionary unit whose coding sequence can be duplicated and/or undergo recombination (Chothia *et al.*, 2003). Small proteins contain just one domain, whereas large proteins are formed by combinations of domains. As in the case of WRKY TFs, *WRKY43* (At2g46130) is the shortest with only 93 aa, whereas *WRKY19* (At4g12020) is the longest with 1798 aa. The main mechanisms of gene evolution are duplication, recombination, and sequence divergence (Ohno 1970; Chothia *et al.*, 2003). It was proposed that WRKY genes encoding a single WRKY domain may have been derived from the C-terminal WRKY domain of the genes




Fig. 5-1. Comparison of the deduced amino acid sequences of WRKY25 and WRKY33. The two WRKY motifs are underlined with the two cysteines and the two histidines of the zinc-finger motif indicated by arrows. The putative nuclear localization signal is shown by a double-headed arrow under the sequence. The asterisks above the sequence indicate identical amino acids, while symbols ':' and '.' indicate conserved and less conserved amino acids, respectively. PcWRKY1, parsley (*Petroselinum crispum*) WRKY1 (GenBank Acc# AAD55974).

harboring two WRKY domains (Wu *et al.*, 2005). We collected the protein sequences of all 72 *Arabidopsis* WRKY TFs and generated an inferred phylogenetic tree based on maximum parsimony analysis of the WRKY domain sequences (Fig. 5-2). *WRKY2*, *WRKY20*, *WRKY25*, *WRKY26* and *WRKY33* encode five structurally related WRKY proteins that belong to group Ia with each protein containing two highly conserved WRKY domains (Fig. 5-2).




Fig. 5-2. A parsimony tree of 72 members of the *Arabidopsis* WRKY gene family. Only the amino acid sequences from the single WRKY domain of group II and III members or the C-

terminal WRKY domain of group I members were used in the tree construction using PAUP*4.0. The numbers on the branches represent branch length.

Our previous microarray data indicated that *WRKY2* and *WRKY26* were not especially responsive to NaCl treatment, while *WRKY20* transcripts increased in abundance by over 2-fold following 24 h and 48 h of exposure (Fig. 5-3). The transcript levels of *WRKY25* and *WRKY33* in *Arabidopsis* roots were greatly increased following salt treatment (Fig.5-3). Of the more than 70 *Arabidopsis* WRKY factors, *WRKY33* is most similar to *WRKY25* and to parsley *WRKY1* involved in responses to pathogen elicitors (Andreasson *et al.*, 2005; Turck *et al.*, 2004). Therefore, we focused on *WRKY25* and *WRKY33* in this study.




Fig. 5-3. Microarray results of *WRKY2*, *WRKY20*, *WRKY25*, *WRKY26* and *WRKY33* transcript abundance changes in response to salt stress. 20 DAS roots were treated by fresh $\frac{1}{2}$ x MS or $\frac{1}{2}$ x MS supplemented with 150 mM NaCl for 6, 24 or 48 h (Jiang and Deyholos, 2006). Data were from three biological replicates with S.D. shown as error bars.

Expression analysis of *WRKY25* and *WRKY33* transcripts in *Arabidopsis* tissues

To examine the expression patterns of *WRKY25* and *WRKY33* in different *Arabidopsis* tissues, we used qRT-PCR to detect their transcript levels in rosette leaves, roots, stem, flower buds, young siliques and dry seeds compared to their levels in roots. *WRKY25* mRNA was more abundant in roots than in any other tissues sampled. *WRKY33* was more abundant in roots as compared to leaves, stems, siliques, and seeds, however the transcripts were slightly more abundant in flowers than roots (Fig. 5-4 A, B).




Fig. 5-4. qRT-PCR analysis of *WRKY25* (A) and *WRKY33* (B) transcript levels in various tissues and organs. R, roots; L, rosette leaves; St, stems; F, flower buds; Si, young siliques; Se, dry seeds. The abundance of mRNA in roots was assigned to be 1. Data were from three biological replicates with S.D. shown.

Subcellular localization

WRKY transcription factors are known to bind to cognate W-box *cis*-elements in the promoter regions of target genes to activate or repress their expression (Eulgem *et al.*, 2000; Rushton *et al.*, 1996). The deduced *WRKY25* protein contains one nuclear localization signal (NLS), PEMKRMK (289-295), whereas the deduced *WRKY33* protein contains two NLSs, PTKKKVE (215-221) and PEAKRWK (323-329) as predicted by PSORT (<http://wolfpsort.org/>, Horton *et al.*, 2006, Fig. 5-1). Additionally, *WRKY33* protein is predicted to contain a 7-amino acid signal peptide, which may target *WRKY33* to the chloroplast according to WoLF-PSORT and TargetP v1.1

(Emanuelsson *et al.*, 2000), as predicted by Schwacke *et al.* (2007). However, previous subcellular localization studies using GFP did not test if WRKY33 could be targeted to chloroplast, because either the 7 amino acid signal peptide was not included in the GFP fusion (Zheng *et al.*, 2006), or it simply was not investigated (Lippok *et al.*, 2007). To examine the sub-cellular localization of WRKY33, the coding regions was fused to the N-terminus of synthetic green fluorescent protein (sGFP) (Chiu *et al.*, 1996) and expressed constitutively under the control of cauliflower mosaic virus (CaMV) 35S promoter. The WRKY33-sGFP fusion protein was found to be localized in the nucleus in root cells (Fig. 5-5A) and in the hypocotyl cells of transgenic seedlings harboring




Fig. 5-5. Subcellular localization of WRKY33 fused to GFP in *Arabidopsis*.

A. Subcellular localization of WRKY33-sGFP in the root cells. The fluorescence (left) and bright-field (middle) images are overlaid at right.

B-C. Subcellular localization of WRKY33-sGFP (B) or ACS1-sGFP (C) in the hypocotyl cells. Left, chlorophyll autofluorescence; middle, GFP fluorescence; right, bright-field.

PWRKY33-sGFP, we failed to detect the fluorescence signal in the chloroplasts (Fig. 5-5B), which indicates that WRKY33 might not be targeted to chloroplasts except nuclei. As a positive control, subcellular of Arabidopsis cysteine synthase (*ACSI*, At2g43750, Hesse and Altmann, 1995) was also investigated. *ACSI* is a nuclear encoded gene with the protein being targeted to chloroplast. In hypocotyl cells, ACS1-GFP signal was observed in the chloroplasts only (Fig. 5-5C).

***WRKY25* and *WRKY33* are induced by multiple abiotic stresses**

Previous studies have shown that the expression of WRKY25 was elevated in cells in response to oxidative stress, heat shock, or wounding (Rizhsky *et al.*, 2004) and WRKY33 acts as a negative regulator of wound response/insect resistance genes and as a positive regulator of pathogen defense-related genes during JA signaling (Zheng *et al.*, 2006; Lippok *et al.*, 2007). To investigate the responses of *WRKY25* and *WRKY33* to various abiotic stresses or chemical treatments, qRT-PCR was used to compare the transcript levels in treated roots in relation to those in corresponding mocks or controls. Paraquat is a bipyridylium-type herbicide that generates superoxide anion inside cells.

The results showed that an increase in *WRKY25* transcript abundance was induced by salt at 6 or 24 h, which is consistent with our microarray results; and it was also induced by drought and Paraquat treatments at 6 h, or by mannitol, cold and heat at 24 h (Fig. 5-6A). *WRKY33* was induced by salt and cold at both time points, and it was also up-regulated by drought and ABA treatments at 6 h, and by osmotic stress at 24 h (Fig. 5-6B). We also monitored the changes of a stress marker gene *RD29A*, which was reported to be responsive to ABA, drought, cold, and salinity (Yamaguchi-Shinozaki and Shinozaki, 1993, 1994). Our qRT-PCR results confirmed that *RD29A* gene was induced by NaCl, cold, drought, ABA, and mannitol; and we also found that it was responsive to AlCl₃ treatment (Fig. 4-4B). On the whole, these results were consistent with the Affymetrix ATH1 results from the Genevestigator public microarray database (<https://www.genevestigator.ethz.ch/at/>) (data not shown).




Fig. 5-6. qRT-PCR analysis of *WRKY25* (A) and *WRKY33* (B) transcript abundance in *Arabidopsis* roots subjected to various treatments as described in Materials and Methods. Values shown are the mean of three biological replicates and S.D. shown as error bars.

Expression analysis of *WRKY25* and *WRKY33* transcripts in ABA and SOS mutants

ABA is a phytohormone regulating several aspects of plant development including seed development, desiccation tolerance of seeds and seed dormancy, and plays a crucial role in the plant's response to abiotic stress (Finkelstein *et al.*, 2002). Stress-responsive genes have been proposed to be regulated by both ABA-dependent and ABA-independent signalling pathways (Chinnusamy *et al.*, 2005; Shinozaki and Yamaguchi-Shinozaki, 2006; Zhu, 2002). For example, RD29A works in both ABA-dependent and

ABA-independent signalling pathways while NaCl-induced *RD29B* accumulation requires ABA (Shinozaki and Yamaguchi-Shinozaki, 1997; Uno *et al.*, 2000).

Regulation of cellular ion homeostasis during salinity stress is critical for plant salt tolerance. Sodium extrusion from plant cells is mediated by the plasma membrane Na^+/H^+ antiporter SOS (Salt Overly Sensitive)1. Activation of SOS1 is controlled by SOS3 and SOS2. The *SOS3* gene encodes a myristoylated Ca^{2+} binding protein, which can interact with and activate the effector serine/threonine kinase, SOS2 (Zhu, 2002).

To examine whether the NaCl-induced accumulation of *WRK25* and *WRKY33* transcripts is dependent on the ABA or SOS pathways, we investigated the expression of these two genes in in ABA biosynthesis and signaling mutants, *aba1-5* and *abi4-1*,




Fig. 5-7. qRT-PCR analysis of *WRKY25* (A) and *WRKY33* (B) transcript abundance in wild-type (WT), *aba1-5*, *abi4-1*, *sos1-1*, *sos2-1*, and *sos3-1* mutants treated by 150 mM NaCl for 6 h. Asterisks indicate a significant difference ($p < 0.05$) between the mutant and WT plants. Data shown were the mean values from three biological replicates and error bars are the S.D.

and three SOS mutants, *sos1-1*, *sos2-1* and *sos3-1* using qRT-PCR. *aba1-5* is a mutant in which the zeaxanthin epoxidase (ZEP), is non-functional, while *abi4-1* is a mutation in a AP2 domain transcription factor responsible for ABA induced transcription (Finkelstein *et al.*, 2002). qRT-PCR results showed that transcript levels of *WRKY25* in any of the five mutants were not significantly different from WT (Fig. 5-7A), suggesting that salt-induced *WRKY25* expression is independent of ABA and SOS pathways. In the absence of normal ABA1 or ABI4 function, accumulation of *WRKY33* was reduced by 38% and 36%, respectively (Fig. 5-7B), therefore, *WRKY33* expression is partially dependent on ABA biosynthesis and signaling pathway. However, salt-mediated *WRKY33* induction was not influenced by any of the three *sos* mutants examined, indicating it is implicated in a SOS-independent pathway. As the *WRKY33* upstream fragment contained ABA responsive elements (ABRE), it was not surprising that salt-induced *WRKY33* accumulation was dependent on ABA.

Analysis of promoter-GUS fusion lines

To examine the role of the upstream regions of *WRKY25* and *WRKY33* genes on their expression in various *Arabidopsis* tissues, we fused a genomic fragment from immediately upstream of the ATG translational start codon (315 bp for PWRKY25 and 1 kb for PWRKY33 where P designates a fragment containing a putative promoter region) to a β -glucuronidase (GUS) reporter gene to determine their developmental, organ-specific and spatial expression patterns. We noted that the length of the intergenic sequence between *WRKY25* (At2g30250) and At2g30260 (encodes U2 small nuclear ribonucleoprotein B) is only 203 bp, so we cloned this intergenic sequence plus the 5'UTR of At2g30250 into our reporter gene fusion.

To avoid potential problems with CaMV35s enhancers, we cloned the promoter regions into a modified binary vector, p0381Z, which did not contain the CaMV35S promoter or a plant selectable marker gene. We therefore co-transformed these promoter-GUS constructs into *Arabidopsis* along with another binary vector (pCAMBIA1300) which provided a selectable marker but not a reporter fusion, and which presumably integrated into the plant genome independently of our reporter fusions.

We examined ten independent T₃ transgenic *Arabidopsis* lines to determine the

GUS expression patterns conferred by the *WRKY25* and *WRKY33* upstream regions. For the PWRKY25-GUS fusion, in 5 d old seedlings, moderate GUS expression was visible in the hypocotyl, root, as well as at the tip of the cotyledon (Fig. 5-8a). In 18 d old seedlings, GUS expression was evident in the leaf veins and roots (Fig. 5-8b). In inflorescence, GUS expression is only visible in stamens, weak in petals (Fig. 5-8c). In young siliques, no GUS expression was observed (Fig. 5-8d).

For plants bearing PWRKY33-GUS fusions, in 5 d old, untreated seedlings, moderate GUS expression was visible in the leaf veins, and less intensely in roots (Fig. 5-8e). In 18 d old seedlings, GUS expression was also evident both in the leaf veins and in the roots (Fig. 5-8f). In inflorescences, GUS expression was only visible in petals and sepals (Fig. 5-8g). In young siliques, weak GUS expression was observed in the tip and restricted area of the silique (Fig. 5-8h).

We then examined the GUS expression in 19 d old seedlings treated by moderate salt, drought, ABA or JA for 6 h and compared the staining pattern to GUS expression in untreated leaves from the same transgenic lines. After 150 mM NaCl and 10 μ M MeJA treatments, we observed a stronger GUS staining in both roots and leaves of PWRKY25-GUS lines (Fig. 5-8j, k). We also observed stronger GUS staining in both roots and leaves of PWRKY33-GUS lines after 150 mM NaCl and dehydration treatments, (Fig. 5-8m, n). Our results demonstrated that the upstream sequence present in PWRKY25-GUS contains sufficient cis-elements to respond to salt or JA treatments, while the upstream fragment present in PWRKY33-GUS contains sufficient cis-elements to respond to salt or drought treatments.

We analyzed the cis-elements contained in the promoter regions of *WRKY25* and *WRKY33* genes using PlantCare (Lescot *et al.*, 2002; Table 5-1). Only one MeJA-, one GA- and one heat stress responsive elements were identified in the 315 bp upstream region of *WRKY25*. However, the 1 kb upstream region of *WRKY33* contained heat, ABA, wounding, SA and low temperature responsive elements, indicating that *WRKY33* may respond to these treatments. Interestingly, *WRKY33* promoter region also contained two W-box elements, suggesting its activity might be regulated by another WRKY TF.




Fig. 5-8. Histochemical localization of GUS activity in representative transgenic *Arabidopsis* plants harboring promoter-GUS constructs.

A-B, Histochemical localization of GUS activity in representative transgenic plants harboring PWRKY25-GUS (A) or PWRKY33-GUS (B). a & e, GUS staining pattern in 5 d old seedling; b&f, GUS staining pattern in 18 d old seedling; c&g, GUS staining pattern in inflorescence; d&h, GUS staining pattern in 7 DAF old siliques.

C, GUS staining pattern of transgenic lines harboring PWRKY25-GUS in 19 d old leaves untreated (i) or treated with 150 mM NaCl (j) or 10 μ M JA (k) for 6 h.

D, GUS staining pattern of transgenic lines harboring PWRKY33-GUS in 19 d old leaves untreated (l) or treated with 150 mM NaCl (m) or drought (n) for 6 h.

Table 5-1. Putative regulatory sequences found in the promoter regions of *WRKY25* and *WRKY33*.

motif	sequence	position	strand	function of regulatory elements
WRKY25 promoter				
GARE-motif	AAACAGA	-38	-	gibberellin-responsive element
HSE	AAAAAAT TTC	-61	+	heat stress responsiveness
TGACG-motif	TGACG	-308	+	MeJA-responsiveness
WRKY33 promoter				
HSE	AAAAAAT TTC	-509	-	heat stress responsiveness
ABRE	TACGTG	-412	+	abscisic acid responsiveness
ARE	TGGTTT	-694	-	essential for the anaerobic induction
AT-rich sequence	TAAAATACT	-624	+	maximal elicitor-mediated activation (2 copies)
W box	TTGACC	-868	-	wounding and pathogen response
		-270	-	
O2-site	GATGACATGA	-581	-	zein metabolism regulation
TC-rich repeats	ATTTTCTCA	-70	+	defense and stress responsiveness
TCA-element	GAGAAGAATA	-66	-	salicylic acid responsiveness
CCAAT-box	CAACGG	-274	+	MYBHv1 binding site
LTR	CCGAAA	-515	+	low-temperature responsiveness
MBS	CGGTCA	-271	+	MYB Binding Site

The 315 bp and 1 kb sequences upstream of ATG start codon of *WRKY25* and *WRKY33*, respectively were analyzed using PlantCare (<http://bioinformatics.psb.ugent.be/webtools/plantcare/html/>). Putative elements involved in light responses (found in all promoters) and the core promoter element (e.e. TATA-box) are not included in the table. The nucleotide immediately before ATG start codon is designated -1.


Mutant phenotypes of *WRKY25* and *WRKY33*





In order to investigate the function of *WRKY25* and *WRKY33* in abiotic stress, we analyzed their mutant phenotypes. We obtained two T-DNA insertion mutants for *WRKY25*, designated as *wrky25-1* (SAIL_529B11) and *wrky25-2* (SALK_136966), and one for *WRKY33*, designated as *wrky33-1* (SALK_006603) (Fig. 5-9A). We used PCR and RT-PCR to confirm homozygosity of these mutant alleles and absence of detectable wild-type transcripts for the mutated genes (Fig. 5-9B). In *wrky25-2*, we

detected a partly reduced *WRKY25* transcript abundance (Fig. 5-9B), so we used only *wrky25-1* for the following experiments. In SALK_006603, we detected a PCR product that was longer than the corresponding band from WT (Fig.5-9B), and after cloning and sequencing, concluded that this line contained a T-DNA induced chromosomal rearrangement. This allele contained no T-DNA insertion but had a 33 bp deletion followed by a 23 bp duplication at the 3'-end of the second intron of *WRKY33* that would prevent proper splicing; therefore, a larger transcript with an unspliced second intron would direct the translation of a truncated protein of 209 amino acids containing no WRKY zinc-finger motif, as reported by Zheng *et al.*(2006).

Under normal growth conditions, we did not observe any morphological differences between any of the confirmed null mutants and WT plants grown in soil. Root elongation and root bending assays of the mutants (compared to WT) were conducted under 100 or 150 mM NaCl or mannitol, 0.5 μ M methyl viologen (Paraquat), and 0.5 μ M ABA treatments. We found that *wrky25-1* mutants did not show significant difference compared to WT plants under all the stress conditions tested (Fig. 5-9C, F), while *wrky33* mutant was more sensitive to salt stress (Fig. 5-9D, G).

Considering the functional redundancy between members of WRKY gene family, especially between members belonging to the same group in the phylogenetic tree (Fig. 5-2), we crossed *wrky25-1* to *wrky33-1* and screened out homozygous double mutant *wrky25wrky33* by PCR and confirmed its homozygosity by RT-PCR. Similarly, we performed root elongation test on $\frac{1}{2}$ x MS medium plates containing 100 mM NaCl. We observed that *wrky25wrky33* mutant roots were significantly shorter than WT








Fig. 5-9. Identification of T-DNA insertion mutants of *WRKY25* and *WRKY33* genes and stress tolerance assay.

A. Schematic diagrams of *WRKY25* and *WRKY33* genes and T-DNA insertion sites. Closed

boxes represent exons (black, coding regions; empty, 5'UTR or 3'UTR), and lines between closed boxes represent introns.

B. RT-PCR validation of homozygous mutant lines of *wrky25-1(a)*, *wrky25-2(b)* and *wrky33-1(c)* using the coding region (CDS) primers and *ACTIN2* gene primers as a positive control. WT, wild-type (Col-0) cDNAs. For SALK_006603, the LP/RP primers were the same as those used for PCR screening (see "Materials and Methods" section).

C. Stress tolerance test of *wrky25-1* (SAIL_529_B11) mutants on ½ x MS containing 100 or 150 mM NaCl, 200 mM mannitol, 0.5 µM MV, or 0.5 µM ABA.

D. Stress tolerance test of *wrky33-1* (SALK_006603) mutants on ½ x MS containing 100 mM NaCl, 200 mM mannitol, 0.5 µM MV, or 0.5 µM ABA.

E. Stress tolerance test of *wrky25wrky33* double mutants on ½ x MS containing 100 mM NaCl.

F. Statistical analysis of the root elongation assay of *wrky25-1* mutants.

G. Statistical analysis of the root elongation assay of *wrky33-1* mutants.

H. Statistical analysis of the root elongation assay of *wrky25wrky33* mutants. Data are means ± S.D. Asterisks indicate a significant difference ($p < 0.05$) between the mutant and WT exposed to the same treatment on the same plate.


($p < 0.01$), suggesting that *wrky25wrky33* double mutant was more sensitive to salt (Fig. 5-9E, H). The performance of *wrky25wrky33* under other types of abiotic stress conditions has not been tested yet.

Overexpression of *WRKY25* and *WRKY33* genes in Arabidopsis

The putative roles of *WRKY25* and *WRKY33* in abiotic stress were further examined by overexpressing them in wild-type Arabidopsis. To do so, we cloned and fused their coding regions under the control of the CaMV35S promoter in Arabidopsis plants. None of the resulting transgenic plants showed detectable phenotypic changes compared with WT plants grown under the same, non-stressed conditions. The expression level of each transgene was evaluated for 10 independent T₂ lines by qRT-PCR. We chose 2-3 lines showing the highest transgene expression for each of the two *WRKY* genes for further characterization. For *WRKY25*, lines 3, 6 and 8 (Fig. 5-10A), and for *WRKY33*, lines 1 and 8 (Fig. 5-10B).

We subjected the selected transgenic lines to various abiotic stress treatments, NaCl, mannitol, Paraquat, ABA, cold and heat, and quantified the root elongation in

comparison to WT seedlings. Roots of *WRKY25*-overexpressing plants grew longer (8%) than that of WT under 100 mM NaCl (Fig 4-10C, D), and they were also more sensitive to Paraquat and ABA treatments as root elongation was inhibited by Paraquat and leaves showed senescence compared to those of WT (Fig 4-10C, D). The lateral roots of *WRKY25*-overexpressing plants were fewer than those of WT under the mannitol treatment, although root elongation was not influenced (Fig. 5-10C, D). Overexpression of *WRKY33* in *Arabidopsis* enhanced the tolerance of transgenic plants to salt stress, as 40% of WT seedling leaves were bleached by 150 mM NaCl, while almost no overexpressing line was bleached by salt treatment (Fig. 5-10E, F). However, similar to *WRKY25*-overexpressing lines, *WRKY33*-overexpressing plants were also more sensitive to Paraquat and ABA treatments than WT as shown by leaf senescence under ABA and significantly shorter roots under moderate ABA treatment ($p < 0.05$, Fig. 5-10E, F). Under Paraquat treatment, root elongation of *WRKY33*-overexpressing lines were significantly shorter than WT, and the leaves of *WRKY33* transgenic lines showed senescence with fewer lateral roots than those of WT plants (Fig. 5-10E, F).









Fig. 5-10. Identification and stress tolerance testing of *WRKY25*- and *WRKY33*-overexpressing

Arabidopsis plants.

A-B, qRT-PCR analyses of transcript levels of *WRKY25* and *WRKY33* in 10 independent transgenic lines (T₂). Rosette leaves harvested from 28 d old plants were used for the assay. The ratios were that of genes in overexpressed lines compared to that of in wild-type and normalized to the level of endogenous gene *UBQ10*.

C, E. Performance of *WRKY25*-(C) and *WRKY33*-overexpressing (E) transgenic plants and wild-type plants under salt, mannitol, paraquat or ABA treatments.

D-F. Effect of 100 or 150 mM NaCl, 200 mM mannitol, 0.1 μM Paraquat or 0.5 μM (±)-ABA on root elongation of WT and *WRKY25*- or *WRKY33*-overexpressing seedlings. Data are means ± S.D. (n = 8–11). Asterisks indicate a significant difference (p < 0.05) between the overexpressing plants (OX) and WT exposed to the same treatment.

Microarray identification of target genes of WRKY25 and WRKY33

To identify genes that had different expression patterns in the absence of WRKY25 or WRKY33 as compared to WT, we used full-genome oligonucleotide microarrays representing 26, 090 genes. RNA was extracted from *wrky25-1*, *wrky33-1* mutants or WT plants, where each of these three genotypes had been treated with 150 mM NaCl for 6 h. Differences in the expression of genes were analyzed by SAM with a FDR of ~5%. After applying a 1.5-fold cutoff, only 16 and 15 genes were down- and up-regulated, respectively, in *wrky25-1* mutant compared to WT (Tables 5-2 & 5-3). In contrast, we detected, using a 1.5-fold cutoff, 87 up-regulated genes and 121 down-regulated genes in *wrky33-1* mutants (Tables 5-4 & 5-5).

The genes that showed different profiles between the *wrky33* mutant and the wild type were classified into 10 categories (Tables 5-4 & 5-5). We observed that the down-regulated genes are quite distinct from the up-regulated genes. For instance, several types of transcription factor, such as AP2/ERF, AUX/IAA, MYB, and zinc-finger family members, were induced in the absence of *wrky33* (Table 5-4). In contrast, quite a few members of class III peroxidase family, glycosyl hydrolase family, and glutathione transferase (GST) family were down-regulated in *wrky33* (Table 5-5).

In the WRKY25 regulon, three protease inhibitor/seed storage/lipid transfer protein (LTP) genes were found to be up-regulated (Table 5-2). LTPs form a multigenic family and can inhibit the growth of fungal pathogens *in vitro* and are also

Table 5-2. Genes that are up-regulated by 150 mM NaCl in *wrky25-1* mutant.

AGI	Putative identity	F.C.	S.D.	W-box	W-box-like
At2g34100	unknown protein, similar to diacylglycerol kinase eta [Strongylocentrotus purpuratus]	37.2	15.00	+	+
At3g62720	Xyloglucan xylosyltransferase(ATXT1)	11.84	6.32	+	+
At4g12480	protease inhibitor/seed storage/lipid transfer protein (LTP) family	2.2	0.81	-	+
At4g12470	protease inhibitor/seed storage/lipid transfer protein (LTP) family	1.8	1.02	-	+
At4g22470	protease inhibitor/seed storage/lipid transfer protein (LTP) family	1.63	0.11	+	+
At2g33780	VQ motif-containing protein contains PF05678: VQ motif	1.78	0.86	+	-
At5g65140	trehalose-6-phosphate phosphatase, putative	1.77	0.61	-	+
At4g23590	aminotransferase family	1.71	0.67	+	+
At3g27360	histone H3	1.7	0.64	+	+
At1g56150	auxin-induced protein family	1.63	0.22	+	+
At5g57990	ubiquitin-specific protease 23 (UBP23), putative	1.55	0.65	+	-
At1g10880	Unknown protein	1.55	0.25	+	-
At5g26260	meprin and TRAF homology domain-containing protein	1.54	0.09	+	+
At5g47440	hypothetical protein	1.54	0.48	+	-
At2g36800	DON-Glucosyltransferase(DOGT1)	1.53	0.43	+	+

AGI, *Arabidopsis* genome initiative; fold change (F.C.) is expressed as the mean value the mRNA level of mutant/WT from three biological replicates; S.D. indicates the standard deviation. The presence or absence of W-box or W-box-like motif in the 1 kb upstream regions is represented by + or -, respectively.

involved in the signalling of defence mechanism of plants against their pathogens (Blein *et al.*, 2002). However, the exact function of LTPs in abiotic stress is not well understood. We also detected *ATXT1*, which is involved in plant cell wall modification (Cavalier and Keegstra, 2006) and *DOGTI* implicated in detoxification of the fusarium mycotoxin deoxynivalenol (Poppenberger *et al.*, 2003) in the up-regulated gene list (Table 5-2).

Among the down-regulated genes in absence of *WRKY25*, there are two pyruvate decarboxylase (PDC) genes working in fermentation, an acyl-(acyl-carrier-protein) desaturase gene for fatty acid biosynthesis, and one sucrose synthase gene (Table 5-3).

Table 5-3. Genes that are down-regulated by 150 mM NaCl in *wrky25-1* mutant.

AGI	Putative identity	F.C.	S.D.	W-box	W-box-like
At5g54960	pyruvate decarboxylase (PDC2)	0.67	0.17	-	+
At4g33070	pyruvate decarboxylase-1 (PDC1)	0.59	0.15	+	-
At5g58070	lipocalin, temperature stress-induced	0.66	0.19	+	+
At2g16060	class 1 non-symbiotic hemoglobin (AHB1/GLB1)	0.65	0.19	+	-
At4g30270	xyloglucan endotransglycosylase (MER15B)	0.65	0.24	+	-
At5g15120	expressed protein	0.64	0.19	+	+
At1g19530	expressed protein	0.62	0.26	+	+
At2g28550	AP2 domain transcription factor RAP2.7/TOE1	0.62	0.29	+	+
At1g43800	acyl-[acyl-carrier-protein] desaturase (stearoyl-ACP desaturase), putative	0.6	0.25	+	+
At5g39890	expressed protein	0.6	0.26	+	+
At3g03270	universal stress protein (USP) family protein / early nodulin ENOD18 family protein	0.58	0.25	-	+
At3g43190	sucrose synthase (UDP-glucose-fructose glucosyltransferase/sucrose-UDP glucosyltransferase), putative	0.57	0.23	+	+
At1g33055	expressed protein	0.52	0.19	+	-
At3g02550	lateral organ boundaries (LOB) domain protein 41 (LBD41)	0.52	0.20	+	-
At5g10040	expressed protein	0.51	0.10	+	+
At4g10270	probable wound-induced protein	0.47	0.13	-	+

In addition, one cell wall-modifying gene, *MER15B*, and, one AP2 TF (RAP2.7/TOE1) involved in miRNA-mediated flowering time control (Aukerman and Sakai, 2003) and, one hemoglobin gene shown to be able to protect plants from oxidative stress (Yang *et al.*, 2005).

In the WRKY33 regulon, we detected 208 differentially expressed genes belonging to 10 functional categories (Tables 5-4 and 5-5). Several members of the AP2/ERF family including two of the best characterized stress-responsive transcription factors, CBF1/DREB1B and CBF2/DREB1C (Stockinger EJ *et al.*, 1997; Gilmour SJ *et al.*, 1998; Medina J *et al.*, 1999) were up-regulated (Table 5-4). Another AP2/ERF family gene (*DDF1/DREB1F*) was previously found to be induced by high salinity and is involved in the regulation of GA biosynthesis and stress tolerance (Magome *et al.*, 2004). Three auxin induced genes (*IAA17*, *IAA13*, and *IAA3*) were up-regulated in *wrky33* mutant. These three IAAs were found to be involved in root development and

light response (Tian and Reed, 1999; Liscum and Reed, 2002). These results indicate that salt stress response and auxin signalling may also interact with each other. We also detected three MYB-related TFs, comprising a single MYB repeat instead of two or three as in most MYB proteins. Among these, CCA1 was found to regulate circadian rhythms (Wang and Tobin 1998). Furthermore, five zinc-finger family members were found to be up-regulated in the absence of *WRKY33*, but their function is largely unknown except COL2, which is implicated in flower development (Ledger *et al.*, 2001). Two genes (*OPR1* and *OPR2*) involved in jasmonic acid biosynthesis were also up-regulated under salt stress in *wrky33*. Both *OPR1* and *OPR2* were up-regulated by senescence, JA and salicylic acid, which confirmed their involvement in the modulation of JA- or SA- responsive gene expression and disease resistance (Eulgem and Somssich 2007). The discovery of up-regulation of *OPR1* and *OPR2* in *wrky33* suggests that *WRKY33* is a negative regulator of JA signaling.

Interestingly, seven members of class III peroxidase gene family were down-regulated in *wrky33* (Table 5-5). Class III peroxidases are plant-specific oxidoreductases that are implicated in various physiological processes such as H₂O₂ detoxification, auxin catabolism, lignification, suberization, stress response (wounding, pathogen attack, NaCl) and senescence (Hiraga *et al.*, 2001). We previously reported that the majority of class III peroxidases are responsive, at the transcript level, to NaCl treatment in *Arabidopsis* roots (Jiang and Deyholos, 2006). The specific functions of these genes remain to be determined. Three glutathione-S-transferases (GSTs) genes, all belonging to Tau sub-family, were also repressed in *wrky33* (Table 5-4). GSTs are proposed to play a role in oxidative stress and apoptosis (Kilili *et al.*, 2004; Apel and Hirt 2004). Although these genes were induced in salt-treated WT plants, they were down-regulated in the *wrky33* mutant, suggesting that *WRKY33* acts upstream of these three GST genes. Five glycosyl hydrolase family genes, were also identified to be down-regulated in *wrky33* mutant (Table 5-5). Glycosyl hydrolases play a role in cell wall remodeling (Xu *et al.*, 2004).

We identified one *WRKY* gene, *WRKY28*, that was down-regulated by salt in *wrky33* (Table 5-5). In WT plant roots, *WRKY28* was strongly induced by salt treatment (Jiang and Deyholos, 2006). Further examination showed that the *WRKY28*

Table 5-4. Genes that are positively regulated by 150 mM NaCl in *wrky33-1* mutant.

AGI	Putative identity	F.C.	S.D.	W-box	W-box-like
Cell growth and division					
At2g42070	MutT/nudix family protein	1.67	0.34	+	+
At3g53830	regulator of chromosome condensation (RCC1) family protein	1.5	0.25	+	-
Cell rescue, defense, death and aging					
At1g59860	heat shock protein, putative	1.52	0.34	+	+
At1g66090	disease resistance protein (TIR-NBS class), putative	1.51	0.4	+	-
Cellular communication/Signal transduction					
			0		
At3g62260	protein phosphatase 2C (PP2C), putative	1.72	0.36	+	+
At3g10190	calmodulin, putative	1.56	0.16	+	-
At1g73800	calmodulin-binding protein	1.62	0.46	+	-
Cellular structure, organization and biogenesis					
At2g28950	expansin, putative (EXP6)	1.6	0.26	-	+
At4g33930	glycine-rich protein	1.76	0.31	-	+
At2g36800	putative glucosyl transferase	1.8	0.72	+	+
At5g67230	glycosyltransferase family 43 protein	1.52	0.32	+	+
At1g23760	polygalacturonase isoenzyme 1 beta subunit - related	1.64	0.28	+	-
Cellular transport and transport facilitation					
At1g29100	copper-binding protein family	1.57	0.32	+	+
At3g44190	pyridine nucleotide-disulphide oxidoreductase family	1.67	0.34	-	-
At3g17690	cyclic nucleotide-binding transporter 2 (CNBT2/CNGC19)	1.71	0.44	-	-
At2g13790	leucine-rich repeat protein kinase family	1.51	0.39	+	+
At3g28740	cytochrome P450 family	1.51	0.17	-	+
At1g50590	pirin-like protein	1.65	0.25	+	+
At3g04620	nucleic acid binding	1.59	0.43	+	+
At1g04570	integral membrane transporter family protein	1.51	0.28	-	-
At5g51980	nucleic acid binding; similar to ZFWD1 (zinc finger WD40 repeat protein 1)	1.66	0.34	+	+
Metabolism					
At4g31780	1,2-diacylglycerol 3-beta-galactosyltransferase (UDP-galactose:diacylglycerol galactosyltransferase) (MGDG synthase) (MGD1), putative	1.51	0.32	+	-
At5g14740	CARBONIC ANHYDRASE 2	1.59	0.43	+	+
At5g22140	pyridine nucleotide-disulphide oxidoreductase family	1.51	0.18	-	+
At4g20320	CTP synthase (UTP--ammonia ligase), putative	1.64	0.37	+	-
At1g76680	12-oxophytodienoate reductase (OPR1)	2.12	0.89	+	-
At1g76690	12-oxophytodienoate reductase (OPR2)	1.87	0.54	+	+

Protein fate (folding, modification, and destination)

At5g56010	heat shock protein, putative	1.56	0.12	+	-
At2g22920	serine carboxypeptidase -related	1.64	0.17	+	-
At5g10540	peptidase M3 family protein / thimet oligopeptidase family protein	1.6	0.37	+	+
At4g23570	phosphatase – related(SGT1A)	1.57	0.21	+	+
At5g51450	RING-finger type ubiquitin ligases(RIN3)	1.92	0.82	+	-

Transcription

At1g12610	AP2/ERF transcription factor, putative(DREB subfamily)/dwarf and delayed-flowering1 (DDF1/DREB1F)	1.92	0.34	+	-
At4g25490	C-repeat/DRE binding factor 1 (CBF1/DREB1B)	1.68	0.18	+	+
At4g25470	C-repeat/DRE binding factor 2 (CBF2/DREB1C)	1.56	0.34	+	+
At1g74930	AP2/ERF transcription factor, putative(DREB subfamily)	1.66	0.28	+	-
At1g21910	AP2/ERF domain-containing transcription factor(DREB subfamily)	1.64	0.21	+	-
At4g34410	AP2/ERF domain transcription factor, putative(ERF subfamily)	1.54	0.4	+	+
At5g61600	AP2/ERF domain transcription factor, putative/ethylene responsive element binding factor 8(ERF subfamily)	1.62	0.21	+	-
At1g04250	auxin-responsive protein IAA17/AXR3	1.53	0.43	+	-
At2g33310	auxin-responsive protein IAA13	1.52	0.33	+	-
At1g04240	auxin-responsive protein IAA3/SHY2	1.73	0.69	+	+
At3g25710	bHLH protein family(BHLH32)	1.55	0.21	+	+
At3g60390	homeobox-leucine zipper protein HAT3	1.75	0.61	+	+
At2g46830	MYB-related transcription factor (CCA1)	1.73	0.38	+	+
At5g17300	myb-related family transcription factor	1.71	0.29	-	+
At1g74840	myb-related family transcription factor(CCA1-like)	1.61	0.25	+	-
At1g10170	zinc finger (NF-X1 type) family protein	1.57	0.41	+	+
At5g54470	zinc finger (B-box type) family protein	1.62	0.55	-	-
At3g02380	Zinc finger protein CONSTANS-LIKE 2 (COL2)	1.6	0.42	+	-
At1g66500	zinc finger (C2H2-type) family protein	1.55	0.25	-	-
At1g04500	zinc finger CONSTANS-related similar to Zinc finger protein constans-like 15 (SP:Q9FHH8)	1.55	0.49	+	-
At3g56790	RNA splicing factor-related contains weak similarity to U4/U6-associated RNA splicing factor [Homo sapiens] gi 2853287 gb AAC09069	1.65	0.43	+	-
At5g01270	Ser-5 specific C-terminal domain phosphatase(CPL2)	1.64	0.27	+	+

Unclassified

At1g34355	forkhead-associated domain-containing protein / FHA domain-containing protein	1.73	0.41	-	+
-----------	---	------	------	---	---

At5g43620	S-locus protein 4-related	1.57	0.3	+	-
At3g61620	exonuclease RRP41	1.54	0.34	+	-
At3g60260	phagocytosis and cell motility protein ELMO1-related	1.54	0.36	-	+
At1g11460	nodulin MtN21 family protein	1.62	0.53	+	+
At3g43660	nodulin - like protein	1.57	0.29	+	-
At1g21140	nodulin, putative similar to nodulin 21 (N-21) [Glycine max] SWISS-PROT:P16313	1.55	0.3	+	+
At3g43630	nodulin -related protein	1.53	0.22	+	+
At5g48850	male sterility MS5 family	1.52	0.25	+	-
At1g67880	N-acetylglucosaminyltransferase - related	1.51	0.28	-	+
At5g65630	bromodomain-containing protein	1.5	0.24	+	+
At3g07360	armadillo/beta-catenin repeat family protein / U-box domain-containing protein contains Pfam domain, PF00514: Armadillo/beta- catenin-like repeats and Pfam, PF04564: U- box domain	1.5	0.14	+	+
At1g69840	band 7 family protein strong similarity to hypersensitive-induced response protein [Zea mays] GI:7716466; contains Pfam profile PF01145: SPFH domain / Band 7 family	1.5	0.24	+	+
At2g03750	sulfotransferase family	1.63	0.54	+	-
At1g09800	tRNA pseudouridine synthase family protein, contains Pfam profile PF01416: tRNA pseudouridine synthaseβ	1.65	0.42	+	+
Unknown					
At5g61820	expressed protein MtN19, Medicago truncatula, EMBL:MTY15367	1.54	0.25	+	-
At2g37880	expressed protein contains Pfam profile PF04759: Protein of unknown function, DUF617	2.18	0.85	+	-
At1g73510	hypothetical protein	1.91	1.1	-	+
At5g17460	expressed protein	1.75	0.4	+	-
At5g41590	hypothetical protein	1.72	0.41	+	-
At3g12320	expressed protein	1.71	0.46	+	+
At1g72240	expressed protein	1.71	0.42	+	-
At3g22090	hypothetical protein	1.7	0.36	-	-
At2g21640	expressed protein	1.7	0.5	+	+
At5g06980	expressed protein	1.64	0.34	-	+
At4g22370	hypothetical protein	1.6	0.3	+	+
At1g62935	hypothetical protein	1.6	0.42	-	-
At1g48405	hypothetical protein	1.58	0.43	+	+
At5g02680	hypothetical protein	1.55	0.39	+	-
At1g73210	expressed protein	1.52	0.4	-	-
At1g05360	Unknown protein	1.51	0.35	+	-
At5g35110	hypothetical protein	1.5	0.43	+	+
At5g62170	Unknown protein	1.6	0.2	-	+

Table 5-5. Genes that are negatively regulated by 150 mM NaCl in *wrky33-1* mutant.

AGI	Putative identity	F.C.	S.D.	W-box	W-box-like
Cell growth and division					
At1g03870	fasciclin-like arabinogalactan-protein (FLA9)	0.65	0.19	+	-
At3g21070	ATP-NAD kinase family protein contains Pfam domain, PF01513: ATP-NAD kinase	0.62	0.11	+	+
At3g23450	glycine-rich protein	0.62	0.21	-	+
At2g05510	glycine-rich protein	0.66	0.08	+	-
At2g05520	glycine-rich protein (GRP)	0.52	0.17	+	+
At5g39190	germin-like protein (AtGER2)	0.48	0.14	+	+
At1g18980	germin-like protein, putative	0.58	0.1	+	-
At5g39110	germin-like protein (GLP6)	0.64	0.13	+	+
At5g38940	germin-like protein, putative	0.59	0.09	+	-
Cell rescue, defense, death and aging					
At1g74590	glutathione S-transferase, putative(AtGSTU10)	0.5	0.14	+	-
At1g69920	glutathione S-transferase, putative(AtGSTU12)	0.59	0.08	+	-
At1g69930	glutathione S-transferase, putative(AtGSTU11)	0.62	0.15	+	+
At1g55020	lipoxygenase (LOX1)	0.67	0.1	+	+
At5g05340	peroxidase, putative(PER52)	0.36	0.32	+	+
At4g08770	peroxidase, putative(PER37)	0.52	0.06	-	+
At4g08780	peroxidase, putative(PER38)	0.58	0.1	+	+
At5g06730	peroxidase, putative(PER54)	0.63	0.17	+	-
At4g37520	peroxidase, putative (PER50)	0.61	0.07	+	+
At1g68850	peroxidase, putative ATP23a(PER11)	0.46	0.13	+	+
At1g14550	anionic peroxidase, putative(PER5)	0.45	0.07	+	+
At2g43510	trypsin inhibitor -related	0.62	0.17	+	-
At2g15220	secretory protein, putative similar to NtPRp27	0.55	0.06	+	+
At4g11650	osmotin-like protein (OSM34)	0.62	0.16	+	+
At5g43570	serine protease inhibitor, potato inhibitor I-type family protein	0.62	0.11	+	+
At3g04720	hevein-related protein precursor (PR-4)	0.57	0.18	+	+
At1g77120	alcohol dehydrogenase (ADH)	0.57	0.09	-	-
At5g24760	alcohol dehydrogenase (ADH), putative	0.66	0.11	+	-
At1g19610	plant defensin protein, putative (PDF1.4)	0.63	0.18	+	+
At5g57625	pathogenesis-related protein PR1 precursor/allergen V5/Tpx-1-related family protein	0.52	0.26	+	+
At3g14225	GDSL-motif lipase/hydrolase protein(GLIP4)	0.35	0.29	+	+
At5g40990	GDSL-motif lipase/hydrolase protein(GLIP1)	0.6	0.03	-	+
At2g38870	protease inhibitor -related	0.67	0.16	-	+
Cellular communication/Signal transduction					
At3g51920	calmodulin 9	0.53	0.01	+	+

At4g33050	calmodulin-binding family protein/EDA39 (embryo sac development arrest 39)	0.64	0.17	+	+
At4g23170	protein kinase family protein, similar to receptor-like protein kinase 5	0.38	0.07	+	-
At3g46270	receptor protein kinase-related	0.44	0.08	+	+
At3g55150	exocyst subunit EXO70 family	0.56	0.15	+	+
At2g39380	exocyst subunit EXO70 family	0.44	0.07	+	-
At3g46280	protein kinase-related contains similarity to light repressible receptor protein kinase. gi 1321686 emb CAA66376	0.65	0.03	+	+
At5g15080	serine/threonine specific protein kinase - related	0.67	0.18	+	-
At2g40180	protein phosphatase 2C (PP2C), putative	0.61	0.06	+	-
At2g19190	light repressible receptor protein kinase, putative	0.37	0.28	+	+
At4g11530	serine/threonine kinase-related protein (fragment)	0.65	0.12	+	-
At4g04960	receptor lectin kinase, putative	0.66	0.18	+	+
At5g06740	lectin-related protein kinase	0.66	0.14	+	+
Cellular structure, organization and biogenesis					
At5g24540	glycosyl hydrolase family 1 protein	0.37	0.29	+	+
At5g24550	glycosyl hydrolase family 1 protein	0.4	0.28	+	-
At3g60120	glycosyl hydrolase family 1 protein	0.46	0.07	+	-
At4g16260	glycosyl hydrolase family 17(beta-1,3-glucanase class I precursor)	0.55	0.22	+	+
At5g24090	glycosyl hydrolase family 18 (acidic endochitinase)	0.55	0.11	+	-
At1g67980	S-adenosyl-L-methionine:trans-caffeoyl-Coenzyme A 3-O-methyltransferase homologue mRNA, complete cds	0.65	0.1	+	+
At4g34050	caffeoyl-CoA 3-O-methyltransferase	0.66	0.11	+	-
Cellular transport and transport facilitation					
At1g15520	ABC transporter family protein	0.62	0.12	-	-
At2g46430	cyclic nucleotide-regulated ion channel (CNGC3)	0.64	0.04	+	-
At1g44100	amino acid permease 5 (AAP5)	0.55	0.11	-	+
At1g77380	amino acid carrier, putative	0.62	0.14	+	-
At3g13610	oxidoreductase, 2OG-Fe(II) oxygenase family	0.51	0.15	+	-
At4g03540	hypothetical integral membrane protein common family	0.62	0.12	+	+
At2g35730	heavy-metal-associated domain-containing protein	0.61	0.06	+	+
At4g12550	protease inhibitor/seed storage/lipid transfer protein (LTP) family	0.42	0.14	+	-
At1g05450	protease inhibitor/seed storage/lipid transfer protein (LTP)-related similar to geranyl diphosphate synthase large subunit [Mentha x piperita] GI:6449052	0.65	0.16	+	+
At3g22600	protease inhibitor/seed storage/lipid transfer	0.63	0.11	+	-

	protein (LTP) family				
At2g44790	uclacyanin II	0.59	0.13	+	-
At2g02850	plastocyanin-like domain containing protein (plantacyanin, putative)	0.64	0.05	+	+
At2g30750	cytochrome p450 71A12	0.35	0.31	-	+
At4g31950	cytochrome P450 family	0.55	0.08	+	+
At2g45970	cytochrome P450, putative	0.66	0.12	+	+
At3g43890	CHP-rich zinc finger protein, putative	0.6	0.12	+	-
At2g34420	photosystem II type I chlorophyll a /b binding protein	0.53	0.19	-	-
At1g26400	FAD-linked oxidoreductase family	0.45	0.26	-	+
At1g26420	FAD-linked oxidoreductase family	0.53	0.23	+	+
At2g46750	FAD-linked oxidoreductase family	0.67	0.16	+	-
At1g54000	myrosinase-associated protein, putative	0.67	0.12	+	-
At4g32490	plastocyanin-like domain containing protein	0.67	0.06	+	-
Metabolism					
At4g25900	aldose 1-epimerase family	0.67	0.06	+	-
At2g02010	glutamate decarboxylase	0.55	0.09	+	+
At2g39400	hydrolase, alpha/beta fold family	0.56	0.18	+	+
At4g36610	hydrolase, alpha/beta fold family	0.63	0.2	+	-
At3g55190	esterase/lipase/thioesterase family	0.61	0.09	+	-
At4g26270	pyrophosphate-dependent phosphofructo-1-kinase	0.61	0.19	+	+
At5g14590	isocitrate dehydrogenase [NADP+], putative	0.54	0.15	+	-
At2g30140	glycosyltransferase family	0.66	0.15	+	-
At5g55120	similar to VTC2 (VITAMIN C DEFECTIVE 2)	0.48	0.04	+	+
At2g26560	lipid acyl hydrolase(PLP2)	0.51	0.09	+	+
At3g01420	feebly-related protein	0.67	0.16	+	-
Protein fate (folding, modification, and destination)					
At5g07460	peptide methionine sulfoxide reductase-related protein	0.63	0.04	+	+
At3g57810	OTU-like cysteine protease family protein	0.56	0.07	+	-
At1g53980	polyubiquitin -related	0.64	0.14	+	+
Transcription					
At3g12910	transcription factor; similar to ANAC042	0.36	0.09	+	+
At5g40590	CHP-rich zinc finger protein, putative	0.47	0.19	+	+
At4g18170	WRKY family transcription factor 28	0.59	0.08	+	+
At5g60710	zinc finger (C3HC4-type RING finger) protein family	0.66	0.08	+	+
At3g53820	zinc finger (C2H2 type) protein family	0.64	0.13	+	+
At1g56380	mitochondrial transcription termination factor family protein	0.45	0.17	+	+
Unclassified					
At2g27550	centroradialis protein, putative (CEN) strong similarity to SP Q41261 CENTRORADIALIS protein	0.62	0.16	+	+
At5g52390	photoassimilate-responsive protein PAR-related protein	0.47	0.23	+	+
At5g09480	proline-rich protein family	0.52	0.09	+	-

At1g53140	dynamain family protein low similarity to dynamain-like protein E	0.53	0.08	+	-
At5g53830	VQ motif-containing protein contains PF05678: VQ motif	0.57	0.12	-	-
At3g02550	lateral organ boundaries (LOB) domain protein 41 (LBD41)	0.58	0.15	+	-
At4g38080	proline-rich protein family	0.64	0.15	+	+
At5g41040	transferase family	0.66	0.09	+	-
At3g47380	invertase/pectin methylesterase inhibitor family protein	0.67	0.07	+	+
At2g33780	VQ motif-containing protein contains PF05678: VQ motif	0.45	0.1	+	-
Unknown					
At1g56320	expressed protein	0.56	0.23	+	+
At1g65500	expressed protein	0.56	0.04	+	+
At4g24130	expressed protein contains Pfam profile PF04398: Protein of unknown function, DUF538	0.58	0.16	+	+
At5g03390	expressed protein contains Pfam profile PF03478: Protein of unknown function (DUF295)	0.61	0.05	+	-
At1g28190	expressed protein	0.62	0.06	+	-
At5g59790	expressed protein	0.62	0.18	+	-
At5g12170	expressed protein similar to unknown protein (pir T09909)	0.62	0.08	-	+
At5g38310	hypothetical protein	0.62	0.09	+	+
At1g52550	expressed protein	0.63	0.07	-	-
At5g57000	expressed protein similar to unknown protein (gb AAF21159.1)	0.65	0.07	+	-
At3g55600	expressed protein	0.65	0.1	+	+
At5g22270	expressed protein similar to unknown protein (gb AAF02129.1)	0.52	0.29	+	+
At1g53970	hypothetical protein	0.36	0.05	+	+
At3g31390	hypothetical protein	0.54	0.06	+	-
At5g44990	hypothetical protein	0.64	0.12	+	+
At5g53820	expressed protein similar to ABA-inducible protein [Fagus sylvatica] GI:3901016, cold-induced protein kin1 [Brassica napus] GI:167146	0.67	0.12	+	-
At4g34120	CBS domain containing protein	0.67	0.05	+	+

promoter region contains three standard W-box *cis*-elements (data not shown), indicating *WRKY28* could be a target of *WRKY33*. Researches on *WRKY53* and *WRKY70* revealed that one *WRKY* gene can regulate the activity of several other *WRKY* genes (Miao *et al.*, 2004; Li *et al.*, 2004).

To validate the changes of some genes, we used qRT-PCR to examine the

changes of four differential genes in the *wrky25-1* dataset and, nine genes including five down-regulated genes and four up-regulated genes, for the *wrky33-1* dataset. These expression profile data obtained by microarray analysis corresponded well with those obtained by qRT-PCR analysis of (Fig. 5-11A, B).




Fig. 5-11. qRT-PCR validation of selected differentially expressed genes between *wrky25-1*(A) or *wrky33-1* (B) mutant and WT under saline condition. Data were the mean values of three biological replicates. Error bars represent S.D.

Only three genes, At2g33780 (VQ motif-containing protein), At2g36800 (glucosyl transferase -related), At3g02550 (lateral organ boundaries (LOB) domain protein 41 (LBD41) were commonly regulated in both *wrky25* and *wrky33*, indicating the WRKY25 and WRKY33 TFs may regulate the activity of quite different downstream genes.

W boxes are highly enriched in WRKY25 and WRKY33 regulon promoters

The regulatory effect of WRKY TFs is primarily through their binding with conserved W box elements on the promoters of specific genes (Ulker and Somssich, 2004). The coregulation of different genes may reflect transcriptional control mechanisms shared and may be based on conserved promoter elements. We inspected 1 kb upstream of the predicted translation start site (ATG). We found the binding site for WRKY transcription factors (W boxes; (C/T)TGAC(T/C); Eulgem *et al.*, 2000; Ulker and Somssich, 2004) was highly enriched in the promoters (Table 5-6). WRKY proteins bind to sequences with an invariant TGAC core, which is in most cases preceded and followed by a T or C. This (C/T)TGAC(T/C) hexamer appears 464 times on both strands in the 208 regulon promoters of WRKY33 regulon, an average of 2.2 copies per promoter (Table 5-6). In WRKY25 regulon, this (C/T)TGAC(T/C) hexamer appears 53 times on both strands in the 31 regulon promoters, an average of 1.7 copies per promoter (Table 5-6).

In contrast, using the most stringent definition for WRKY binding sites (a TTGACC/T hexamer), we still detected 162 genes (78%) whose promoters contain either a TTGACC or TTGACT motif or both. In total, we found a significant over-representation of 376 potential binding sites, an average of 1.8 copies per promoter, in these 57 *WRKY33*-regulated gene promoters (Table 5-6). We note that the *P* value for distribution of W-box hexamers is also significant. 46 of 208 *WRKY33* regulon promoters do not contain a copy of the TTGACC/T motif. However, we also observed that 116 promoters contain the W-box-like hexamer TTGACA. Previous report showed that this W-box-like motif is found 31 times within the 26 *PR-1* regulon promoters (Maleck *et al.*, 2000), which is twice the statistically expected frequency. Clustered occurrences of W boxes are common among promoters of WRKY-regulated genes. Association of these *cis*elements as palindromes or direct repeats with spacing of less than 20 bp results in binding cooperativity and synergistic transcriptional activation (Rushton *et al.*, 1996; Fukuda 1997; Eulgem *et al.*, 1999). We found both the core and stringent W-box elements (TTGAC and TTGACC/T) arranged in a palindromic or directly repeated manner on many promoters from the *WRKY33* regulon.

Table 5-6. Occurrence of W-box and W-box-like motifs in the *WRKY25* and *WRKY33* regulons.

oligomer	Absolute number of this oligomer in query set	Number of sequences in query set containing oligomer	average frequency	p-value from binomial distribution	data set
WRKY33 regulon					
TTGACT	169	90/121	1.9	9.09E-08	down
	77	47/87	1.6	7.39E-02	up
TTGACC	92	59/121	1.6	7.81E-04	down
	38	32/87	1.2	8.53E-02	up
CTGACC	20	20/121	1.0	7.49E-02	down
	20	17/87	1.2	4.22E-02	up
CTGACT	26	24/121	1.1	9.03E-02	down
	22	20/87	1.1	8.18E-02	up
TTGACA(W-box-like)	110	70/121	1.6	3.34E-02	down
	68	46/87	1.5	8.47E-02	up
GTGACT	39	36/121	1.1	5.90E-02	down
	21	16/87	1.3	2.14E-02	up
TTGACG	43	35/121	1.23	5.42E-02	down
	38	29/87	1.31	2.31E-02	up
WRKY25 regulon					
TTGACT	15	10/16.	1.5	1.33E-01	down
	18	9/15.	2.0	0.163	up
TTGACC	7	6/16.	1.2	1.99E-01	down
	6	5/15.	1.2	2.11E-01	up
CTGACT	3	3/16.	1.0	2.46E-01	down
	4	4/15.	1.0	1.89E-01	up
TTGACA(W-box-like)	19	11/16.	1.7	3.34E-02	down
	19	11/15.	1.7	8.52E-02	up
GTGACT	5	4/16.	1.3	2.23E-01	down
	6	5/15.	1.2	1.82E-01	up
TTGACG	3	3/16.	1.00	2.03E-01	down
	4	3/15.	1.33	2.22E-01	up

Transcription factors of the TGA subgroup of bZIP proteins (TGA-bZIPs) are positive regulators of INA-induced *PR-1* expression. They recognize a pentameric element, TGACG that potentially overlaps the W box consensus (Schindler *et al.*, 1992). The TGAC W box core, however, is followed by C or T in all available reports of WRKY/W box interactions. Thus, a distinction between binding sites for these two regulatory factor families can be predicted. We observed that 81 TTGACG motifs were found in 64 promoters.

Although WRKY factors show high binding affinity to a DNA sequence designated the W box, (C/T)TGAC(T/C), altered binding preferences have also been observed (Sun *et al.*, 2003). In barley, a WRKY transcription factor, SUSIBA2 implicated in sugar signaling was found to be able to bind the variant W-box motif GTGACT in the *isol* promoter. We counted 60 GTGACT motifs from 52 WRKY33 regulon promoters. We propose that the remaining 13 genes (6%) whose promoters do not contain W-box or W-box-like motifs may be the indirect targets of WRKY33.

As for WRKY25 regulon, we also detected 25 genes (81%) whose promoters contain W-box motif, and 22 genes (71%) contain W-box-like motif in their promoter regions. Furthermore, we found that 24 genes (77%) contain the stringent W-box motif (TTGACT/C) in their promoters. These data suggest that both WRKY25 and WRKY33 can positively and negatively control the expression of a distinct subset of genes, either directly or indirectly.

Discussion

Soil salinity has detrimental effects on plant growth and crop yields in agricultural production. Excess salts in the soil interfere with the plants' ability to take up nutrients and cause water deficit, followed by a decrease of the K^+/Na^+ ratio in cells. To minimize stress damage and to re-establish ion homeostasis, plants respond and adapt to salt stress by changing gene expression patterns, metabolic activity, and ion and water transport (Zhu, 2003). Many genes with various functions are induced by high salinity stress, and their gene products function in stress tolerance (Chinnusamy *et al.*, 2005). Numerous evidences indicate that specificity and cross-talk exist between various abiotic stresses as well as between biotic and abiotic stress (Seki *et al.*, 2003; Zhu *et al.*, 2001, 2002).

Proteins belonging to the WRKY superfamily of transcription factors are known to regulate a multiplicity of biological processes. Members of the "plant-specific" WRKY transcription factor family have been implicated in the regulation of genes involved in pathogen-induced stress (Dong *et al.*, 2003; Eulgem *et al.*, 2000; Eulgem and Somssich, 2007). Some WRKY genes were also found to be responsive to salt, as well as drought, and cold stresses (Seki *et al.*, 2002; Jiang and Deyholos, 2006). They

are also believed to play an important role in leaf senescence (Robatzek and Somssich, 2001; Miao *et al.*, 2004). Recent studies suggest that biotic and abiotic stress responses may share common components, indicating that there is cross-talk between abiotic and biotic stress signal transduction pathways (Park *et al.* 2001; Mengiste *et al.* 2003; Chini *et al.* 2004). This study shows that two members of Arabidopsis *WRKY* gene family, *WRKY25* and *WRKY33*, are involved in modulating abiotic stress responses.

WRKY 25 and WRKY33 are involved in abiotic stress responses

WRKY genes have been duplicated many times during plant evolution, resulting in a large gene family involved in regulating a large set of genes and thereby ensuring proper cellular responses to physiological processes, internal and external stimuli (Zhang and Wang, 2005). It is believed that functional redundancy might exist between members of *WRKY* gene family (Ulker and Somssich 2004). Therefore, a more practical approach to characterizing them is to study a group of phylogenetically closed *WRKY* genes together, as demonstrated by Xu *et al.*(2006). According to the phylogenetic tree of the 72 *WRKY* TFs in *Arabidopsis*, *WRKY2*, *WRKY20*, *WRKY25*, *WRKY26* and *WRKY33* genes belong to group Ia containing two *WRKY* domains. However, considering that *WRKY20* and *WRKY26* were not responsive to salt treatment, we studied *WRKY25* and *WRKY33* together. Our qRT-PCR results showed that the transcript abundance of *WRKY25* and *WRKY33* was increased by multiple abiotic stresses (Fig. 5-6A, B). Although these two genes were similarly induced by NaCl treatment, differences in their responses to other abiotic stresses were detected, for instance, *WRKY25* transcript abundance was increased by cold and heat treatments at 24 h time point (Fig. 5-6A), while *WRKY33* by cold treatment at both 6 and 24 h (Fig. 5-6B). In our experiment, *WRKY33* is not responsive to oxidative stress treatment induced by 10 μ M Paraquat (Fig. 5-6B), whereas Zheng *et al.*(2006) reported that *WRKY33* transcript level is increased by 100 μ M Paraquat treatment. This difference can be explained by the different concentrations of Paraquat used. Overall the different responses of *WRKY25* and *WRKY33* to abiotic stresses suggest that they may play different roles in Arabidopsis under abiotic stress.

WRKY proteins have been identified as transcription factors that bind DNA sequences containing W-boxes (C/TTGACC/T). The nuclear localization of *WRKY25*

and WRKY33 using GFP as a reporter gene suggests that they may possess transcription factor activity (Fig 5-5). Among the target genes identified by microarray (Tables 5-2 to 5-5), many contain W box or W box-like motif in their promoters. In order to test the WRKY protein–W-box-like motif interaction, we tried to express WRKY25 and WRKY33 in *E. coli* using pET32a(+) vector and BL21(DE3) or Rosetta gami strains (Novagen), which proved unsuccessful, most probably due to the high frequency of presence of rare codons encoding Arg, Lys and His amino acids (data not shown).

qRT-PCR assay demonstrated that transcripts of both *WRKY25* and *WRKY33* were most abundant in leaves, roots, and also in stems and flowers (Fig. 5-4). These mRNA accumulation patterns were generally consistent with the GUS expression patterns observed in promoter-GUS fusions of each gene (Fig. 5-8). This consistence between mRNA and GUS staining pattern can be partly explained by the fact that we used a co-transformation strategy to avoid the enhancer effect of CaMV35S promoter on the nearby reporter gene expression. Using promoter-GUS fusion, we detected the inducibility of GUS expression by salt for both PWRKY25 and PWRKY33 (Fig. 5-8), which were also induced by JA and drought, respectively (Fig. 5-8). However, we failed to observe the inducibility of PWRKY25 and PWRKY33 by other stress treatments tested, which is different from qRT-PCR results shown in Fig. 5-6. The minor difference on GUS expression between untreated and treated samples may only be detected by performing a quantitative GUS enzymatic assay. In addition, promoter fusion study using GUS may still not be a perfect way of studying gene expression patterns because regulatory elements may also exist within the coding region, introns, or downstream regions instead of the 5' upstream region (Taylor 1997).

ABA is an important stress hormone extensively involved in response to abiotic stresses such as salt, drought and osmotic stress. At least four regulatory systems exist for the regulation of salt- and osmotic stress-inducible genes in *Arabidopsis thaliana* (Shinozaki & Yamaguchi-Shinozaki 1997; Shinozaki *et al.*, 2003). Although most salinity-inducible genes are regulated by an ABA-dependent pathway, some are not dependent on ABA for signalling. A conserved *cis*-acting element, referred to as DRE/CRT, functions in stress-inducible gene expression in an ABA-independent

manner (Shinozaki *et al.*, 2003). In ABA-dependent pathways, the ABRE (ABA-responsive element) is a major *cis*-acting element in stress-responsive gene expression (Shinozaki *et al.*, 2003). Genetic analysis of two *Arabidopsis* ABA-deficit mutants, *aba1/los6* and *aba3/los5*, in which genes encoding molybdenum cofactor sulfurase (MCSU) and zeaxanthin epoxidase (ZEP) are dysfunctional, showed that ABA is necessary for the salt-stress induced expression of some late-embryogenesis-abundant (LEA) genes (Xiong *et al.*, 2001, 2002a, 2002b). qRT-PCR analysis of *Arabidopsis* roots treated by 50 μ M ABA for 6 h demonstrated that the transcript abundance of *WRKY25* and *WRKY33* was slightly increased compared to the control. The putative promoter region of *WRKY25* does not contain any ABRE element, whereas *WRKY33* contains one single ABRE element (Table 5-1). Skriver *et al.* (1991) reported that polymerized copies of ABREs can confer ABA responsiveness to a minimal promoter, whereas a single copy of ABRE is not sufficient for the full ABA response. Our results showed that salt-induced *WRKY25* transcript accumulation does not require ABA, while *WRKY33* is partially ABA-dependent, suggesting *WRKY25* and *WRKY33* may work in divergent pathways with respect to ABA. *wrky25* and *wrky33* mutants did not show significant difference from WT on the ABA medium plates (Fig. 5-9), however, *WRKY25*- and *WRKY33*- overexpressing plants were more sensitive to ABA treatment (Fig. 5-10), suggesting that *WRKY25* and *WRKY33* might work as positive regulators in mediating plant responses to ABA. Previous researches with rice *WRKY* genes have revealed positive and negative regulators of ABA signaling (Xie *et al.*, 2005).

Arabidopsis sos mutants were identified by genetic screening of seedlings that were hypersensitive to NaCl stress, characterized by retarded root growth (Liu & Zhu 1998; Liu *et al.* 2000; Shi *et al.* 2000). In particular, the *sos1*, *sos2*, and *sos3* mutants are specifically hypersensitive to Na⁺ and Li⁺ ions. Genetic and physiological data indicate that *SOS1*, *SOS2*, and *SOS3* function within the same pathway, in which the *SOS3*–*SOS2* complex regulates the expression level of *SOS1* and activates the transport activity of *SOS1*, which leads to Na⁺ tolerance (Qiu *et al.* 2002; Zhu 2003). In addition, salt stress responses were also shown to be mediated by *SOS*-independent signaling pathways with the identification of several protein kinases activated by salt or osmotic stress (Mikolajczyk *et al.*, 2000; Zhu 2001). In *Arabidopsis* seedlings, a 40-

kDa kinase similar to ASK1 (serine/threonine kinase 1) was identified as being rapidly activated by hyperosmotic stress in a calcium- and ABA-independent manner. Osmotic stress activation of the *Arabidopsis* 40-kDa protein kinase is not impaired by the *sos3* mutation, implying that its activation is independent of the SOS pathway (Hoyos and Zhang, 2000). Our results showed that salt-induced *WRKY25* or *WRKY33* transcript accumulation was not influenced in any of the three *sos* mutants, indicating both *WRKY25* and *WRKY33*-mediated salt response is independent of the SOS signaling pathway.

Knock-out mutation by T-DNA insertion and overexpression are two of the most frequently used techniques to characterize gene function. We found that *wrky25* mutant did not show any significant difference under stress condition compared to WT control, while *wrky33* is slightly more sensitive to salt treatment (Fig. 5-9). The observation suggests that functional redundancy exists between WRKY TFs within group Ia. Accordingly, the double mutant *wrky25wrky33* is more sensitive to salt treatment, although its sensitivity to other types of abiotic stress still needs to be tested. In the case of *Arabidopsis* WRKY proteins, structurally related WRKY18, WRKY40, and WRKY60 have been shown to have functional redundancy (Xu *et al.* 2006). On the other hand, AtWRKY44 and AtWRKY10 may act much less extent in functional redundancy (Johnson *et al.* 2002; Luo *et al.* 2005).

Overexpression of TFs offers an alternative and complementary strategy to knockout analysis as it is less affected by functional redundancy. In plants, there are several examples in which knockouts of TF genes failed to generate informative phenotypes, but overexpression of these TFs revealed gene functions (Baima *et al.*, 2001; van der Graaff *et al.*, 2002; Zhang 2003). Overexpression of *WRKY25* and *WRKY33* using the constitutive CaMV35S promoter demonstrated similar phenotypes, i.e. enhanced tolerance to salt stress, but more sensitive to oxidative and ABA treatments (Fig. 5-10). Our result on the sensitivity of *WRKY25*-overexpressing plants to oxidative stress is consistent with a previous report (Rizhsky *et al.*, 2004). These data suggest that both *WRKY25* and *WRKY33* contribute to salt tolerance in *Arabidopsis*.

WRKY25 and WRKY33 regulate salt stress response by targeting to diverse downstream genes

The role of WRKY25 and WRKY33 in salt stress response was further dissected by identifying their target genes through full-genome microarrays. The number of differentially expressed genes identified between *wrky25* and WT plants is 31, while the number increased to be 208 between *wrky25* and WT plants under a moderate salt treatment (Tables 5-2 to 5-5). The numbers of differentially expressed genes identified reflect the consequence of disrupting *WRKY25* and *WRKY33* expression, which is consistent with the mutant phenotypes under salt stress condition (Fig. 5-9). There are only three common genes between WRKY25 and WRKY33 regulons, suggesting that WRKY25 and WRKY33 have separate roles in salt stress response in addition to common ones. The identified target genes of WRKY33 were classified into ten categories, with many genes implicated in cell defense, cellular transport, signal transduction, metabolism and transcription, and still the identities of many genes are not clear yet (Tables 5-4 and 5-5), indicating a diverse roles of these target genes regulated directly or indirectly by WRKY33. We note that among the differentially expressed genes identified, many are those encoding transcription factors, especially in the WRKY33 regulon (Tables 5-4 and 5-5), which suggests that WRKY33 can relay the salt stress signal to downstream TFs by controlling their expression.

Inspection of 1 kb upstream regions of these target genes demonstrated that W boxes are highly enriched in WRKY25 and WRKY33 regulon promoters (Table 5-6); however, it is also possible that WRKY25 and WRKY33 interact with other factors directly or indirectly to regulate downstream gene expression. The flanking nucleotides around W boxes may affect the binding affinity of WRKY25 and/or WRKY33 transcription factors to their respective target sequences and can serve to discriminate among closely related factors.

Earlier research has demonstrated that the *Arabidopsis* MAPK kinase 2 (MKK2) with its upstream activator MAPK kinase kinase MEKK1 and the downstream MAPKs MPK4 and MPK6 mediate cold and salt stress signaling (Teige *et al.*, 2004). Salt stress induces the expression and activity of AtMEKK1 (Ichimura *et al.*, 2000). AtMPK4 is activated by cold, low humidity, osmotic stress, touch, and wounding (Ichimura *et al.*,

2000). MPK4 can phosphorylate WRKY25 and WRKY33 *in vitro*, and a substrate of MPK4, MKS1 can interact with WRKY25 and WRKY33 (Andreasson *et al.*, 2005), indicating that a MAPK cascade regulates WRKY25 and WRKY33 activity. A previous study with an Arabidopsis MAPK phosphatase 1 mutant (*mkp1*) demonstrated the complexity of the AtMPK4 signalling cascade. *mkp1* is resistant to salinity but hypersensitive to UV-C. A yeast two-hybrid screen showed that MKP1 could interact with AtMPK4 (Ulm *et al.*, 2002). Microarray analysis of *mkp1* indicated that the transcript level of a putative Na⁺/H⁺-antiporter (CHX17, At4g23700) is negatively regulated by AtMKP1 (Ulm *et al.*, 2002). More recent research on MEKK1 and MPK4 suggests that their functions are not limited to a single, linear signaling pathway




Fig. 5-12. Proposed working model explaining role of WRKY25 and WRKY33 in salt stress responsive pathway. High concentration of salt in the apoplast is sensed by an unknown sensor on the plasma membrane. This is transduced by a MAPK-mediated pathway to MPK4, which can phosphorylate MKS1. MKS1 activates WRKY25 and WRKY33, which can recognize and bind to the W-box in the downstream target genes and activate their transcription. Arrows indicate positive regulation, and open blocks indicate negative regulation.

pathway (Su *et al.*, 2007), which suggests the existence of other unknown component(s). These results, together with our analysis, suggest that two WRKY transcription factors function in the MPK4-mediated signaling pathways in plant responses to salt stress. Thus, these two transcription factors play a role in responses to both biotic as well as abiotic stresses and, biotic and abiotic stress signals converge at the level of MAPK cascades. On the basis of the results presented here as well as those reported previously (Ulm *et al.*, 2002; Teige *et al.*, 2004; Andreasson *et al.*, 2005), we present a model describing the interrelationships among MAPK signaling components, WRKY25 and WRKY33 in the salt signal transduction pathway (Fig. 5-12). This model, together with other recent studies of the role of MPK4 in salt and ROS signalings (Nakagami *et al.*, 2006; Su *et al.*, 2007), allows a better understanding of salt signal transduction pathway from sensing to the response in plants.

In summary, this study shows that WRKY25 and WRKY33 play important roles in abiotic stress responses. This strengthens our understanding of transcriptional regulation during salt and other types of abiotic stress responses in higher plants and provides a base for future work in this area.

References

- Alonso JM, Stepanova AN, Leisse TJ, Kim CJ, Chen H, Shinn P, Stevenson DK *et al.* (2003) Genome-wide insertional mutagenesis of *Arabidopsis thaliana*. *Science* 301: 653-657
- Anderson JP, Badruzsaufari E, Schenk PM, Manners JM, Desmond OJ, Ehlert C, Maclean DJ, Ebert PR and Kazan K. (2004) Antagonistic interaction between abscisic acid and jasmonate-ethylene signaling pathways modulates defense gene expression and disease resistance in *Arabidopsis*. *Plant Cell* 16:3460-3479
- Andreasson E, Jenkins T, Brodersen P, Thorgrimsen S, Petersen NH, Zhu S, Qiu JL, Micheelsen P, Rocher A, Petersen M, Newman MA, Bjorn Nielsen H, Hirt H, Somssich I, Mattsson O and Mundy J. (2005) The MAP kinase substrate MKS1 is a regulator of plant defense responses. *EMBO J* 24:2579-2589
- Apel K, and Hirt H. (2004) Reactive oxygen species: metabolism, oxidative stress, and signal transduction. *Annu Rev Plant Biol* 55:373-399

- Asai T, Tena G, Plotnikova J, Willmann MR, Chiu WL, Gomez-Gomez L, Boller T, Ausubel FM and Sheen J.** (2002) MAP kinase signalling cascade in *Arabidopsis* innate immunity. *Nature* 415, 977–983
- Aukerman MJ and Sakai H.** (2003) Regulation of flowering time and floral organ identity by a MicroRNA and its APETALA2-like target genes. *Plant Cell* 15:2730-2741
- Baima S, Possenti M, Matteucci A, Wisman E, Altamura MM, Ruberti I and Morelli G.** (2001) The *Arabidopsis* ATHB-8 HD-Zip protein acts as a differentiation-promoting transcription factor of the vascular meristems. *Plant Physiol* 126, 643–655
- Blein JP, Coutos-Thévenot P, Marion D and Ponchet M.** (2002) From elicitors to lipid transfer proteins: a new insight in cell signaling involved in plant defence mechanism, *Trends Plant Sci* 7, 293–296
- Cavalier DM and Keegstra K.** (2006) Two xyloglucan xylosyltransferases catalyze the addition of multiple xylosyl residues to cellohexaose. *J Biol Chem* 281:34197-34207
- Chen C and Chen Z.** (2002) Potentiation of developmentally regulated plant defense response by AtWRKY18, a pathogen-induced *Arabidopsis* transcription factor. *Plant Physiol* 129, 706–716
- Chen D-H and Ronald PC.** (1999) A rapid DNA minipreparation method suitable for AFLP and other PCR applications. *Plant Mol Biol Rep* 17: 53-57
- Chen WJ and Zhu T.** (2004) Networks of transcription factors with roles in environmental stress response. *Trends Plant Sci* 9:591-596.
- Chini A, Grant JJ, Seki M, Shinozaki K and Loake GJ.** (2004) Drought tolerance established by enhanced expression of the CC-NBS-LRR gene, *ADR1*, requires salicylic acid, EDS1 and ABI1. *Plant J* 38:810–822
- Chinnusamy V, Jagendorf A and Zhu J-K.** (2005) Understanding and improving salt tolerance in plants. *Crop Sci* 45: 437-448
- Chinnusamy V, Schumaker K, and Zhu J-K.** (2004) Molecular genetic perspectives on cross-talk and specificity in abiotic stress signalling in plants. *J Exp Bot* 55:225-236
- Chiu W, Niwa Y, Zeng W, Hirano T, Kobayashi H and Sheen J.** (1996) Engineered GFP as a vital reporter in plants. *Curr Biol* 6, 325-330
- Chothia C, Gough J, Vogel C and Teichmann S A.** (2003) Evolution of the protein

Repertoire. *Science*, 300, 701–1703

Clough SJ and Bent AF. (1998) Floral dip: A simplified method for *Agrobacterium*-mediated transformation of *Arabidopsis thaliana*. *Plant J* 16, 735–743

Dellagi A, Helibronn J, Avrova AO, Montesano M, Palva ET, Stewart HE, Toth IK, Cooke DE, Lyon GD and Birch PR. (2000) A potato gene encoding a WRKY-like transcription factor is induced in interactions with *Erwinia carotovora* subsp *atroseptica* and *Phytophthora infestans* and is coregulated with class I endochitinase expression. *Mol Plant–Microbe Interact* 13, 1092–1101

Devaiah BN, Karthikeyan AS and Raghothama KG. (2007) WRKY75 transcription factor is a modulator of phosphate acquisition and root development in *Arabidopsis*. *Plant Physiol* 143, 1789-1801

Dong J, Chen CH and Chen ZX. (2003) Expression profiles of the *Arabidopsis* WRKY gene superfamily during plant defense response. *Plant Mol Biol* 51, 21–37

Emanuelsson O, Nielsen H, Brunak S and von Heijne G. (2000) Predicting subcellular localization of proteins based on their N-terminal amino acid sequence. *J Mol Biol* 300: 1005-1016

Eulgem T, Rushton PJ, Robatzek S, and Somssich IE. (2000) The WRKY superfamily of plant transcription factors. *Trends Plant Sci* 5, 199–206

Eulgem T, Rushton PJ, Schmelzer E, Hahlbrock K and Somssich IE. (1999) Early nuclear events in plant defence signalling: rapid activation by WRKY transcription factors. *EMBO J* 18, 4689–4699

Eulgem T and Somssich IE. (2007) Networks of WRKY transcription factors in defense signaling. *Curr Opin Plant Biol* 10:366–371

Finkelstein RR, Gampala SS and Rock CD. (2002) Abscisic acid signaling in seeds and seedlings. *Plant Cell* 14:S15-45

Fujita M, Fujita Y, Noutoshi Y, Takahashi F, Narusaka Y, Yamaguchi-Shinozaki K and Shinozaki K. (2006) Crosstalk between abiotic and biotic stress responses: a current view from the points of convergence in the stress signaling networks. *Curr Opin Plant Biol* 9:436–442

Fukuda Y. (1997) Interaction of tobacco nuclear proteins with an elicitor responsive element in the promoter of a basic class I chitinase gene. *Plant Mol Biol* 34, 81–87

- Gilmour SJ, Zarka DG, Stockinger EJ, Salazar MP, Houghton JM and Thomashow MF.** (1998) Low temperature regulation of the *Arabidopsis* CBF family of AP2 transcriptional activators as an early step in cold-induced COR gene expression. *Plant J* 16:433-442
- Hall BG.** (2001) *Phylogentic trees made easy: A how-to manual for molecular biologists.* Sinauer Associates, Inc, Massachusetts, USA
- Hara K, Yagi M, Kusano T, and Sano H.** (2000) Rapid systemic accumulation of transcripts encoding a tobacco WRKY transcription factor upon wounding, *Mol Gen Genet* 263:30–37
- Hesse and Altmann.** (1995) Molecular cloning of a cysteine synthase cDNA from *Arabidopsis thaliana*. *Plant Physiol* 108: 851-852
- Hiraga S, Sasaki K, Ito H, Ohashi Y and Matsui H.**(2001) A large family of class III plant peroxidases. *Plant Cell Physiol* 42: 462-468
- Horton P, Park K-J, Obayashi T and Nakai K.** (2006) Protein subcellular localization prediction with WoLF PSORT. *Proceedings of the 4th Annual Asia Pacific Bioinformatics Conference APBC06, Taipei, Taiwan.* pp. 39-48
- Hoyos ME and Zhang S.** (2000) Calcium-independent activation of salicylic acid-induced protein kinase and a 40-kilodalton protein kinase by hyperosmotic stress. *Plant Physiol* 122, 1355–1363
- Ichimura K, Mizoguchi T, Yoshida R, Yuasa T and Shinozaki K.** (2000) Various abiotic stresses rapidly activate *Arabidopsis* MAP kinases ATMPK4 and ATMPK6. *Plant J* 24, 655-665
- Jiang Y-Q and Deyholos MK.** (2006) Comprehensive transcriptional profiling of NaCl-stressed *Arabidopsis* roots reveals novel classes of responsive genes. *BMC Plant Biol* 6, 25
- Jiang Y-Q, Yang B, Harris NS and Deyholos MK.** (2007) Comparative proteomic analysis of NaCl stress-responsive proteins in *Arabidopsis* roots. *J Exp Bot* 58:3591-3607
- Johnson CS, Kolevski B and Smyth DR.** (2002) *TRANSPARENT TESTA GLABRA2*, a trichome and seed coat development gene of *Arabidopsis*, encodes a WRKY transcription factor. *Plant Cell* 14, 1359–1375

- Kalde M, Barth M, Somssich IE and Lippok B.** (2003) Members of the *Arabidopsis* WRKY group III transcription factors are part of different plant defense signaling pathways. *Mol Plant–Microbe Interact* 16, 295–305
- Kilili KG, Atanassova N, Vardanyan A, Clatot N, Al-Sabarna K, Kanellopoulos PN, Makris AM and Kampranis SC.** (2004) Differential roles of Tau class glutathione S-transferases in oxidative stress. *J Biol Chem* 279: 24540-24551
- Lagace M and Matton DP.** (2004) Characterization of a WRKY transcription factor expressed in late torpedo-stage embryos of *Solanum chacoense*. *Planta* 219, 185–189
- Ledger S, Strayer C, Ashton F, Kay SA and Putterill J.** (2001) Analysis of the function of two circadian-regulated *CONSTANS-LIKE* genes. *Plant J* 26:15-22
- Lescot M, Dehais P, Thijs G, Marchal K, Moreau Y, Van de Peer Y, Rouze P and Rombauts S.** (2002) PlantCARE, a database of plant cis-acting regulatory elements and a portal to tools for *in silico* analysis of promoter sequences. *Nucleic Acids Res* 30:325-327
- Li J, Brader G and Palva ET.** (2004) The WRKY70 transcription factor: a node of convergence for jasmonate-mediated and salicylate-mediated signals in plant defense. *Plant Cell* 16:319-331
- Lippok B, Birkenbihl RP, Rivory G, Brümmer J, Schmelzer E, Logemann E and Somssich IE.** (2007) Expression of AtWRKY33 encoding a pathogen- or PAMP-responsive WRKY transcription factor is regulated by a composite DNA motif containing W box elements. *Mol Plant Microbe Interact* 20:420-429
- Liscum E and Reed JW.** (2002) Genetics of Aux/IAA and ARF action in plant growth and development. *Plant Mol Biol* 49:387-400
- Liu J, Ishitani M, Halfter U, Kim CS and Zhu J-K.** (2000) The *Arabidopsis thaliana* *SOS2* gene encodes a protein kinase that is required for salt tolerance. *Proc Natl Acad Sci USA* 97, 3730–3734
- Liu J and Zhu J-K.** (1998) A calcium sensor homolog required for plant salt tolerance. *Science* 280, 1943–1945
- Luo M, Dennis ES, Berger F, Peacock WJ and Chaudhury A.** (2005) *MINISEED3* (*MINI3*), a WRKY family gene, and *HAIKU2* (*IKU2*), a leucine-rich repeat (LRR) KINASE gene, are regulators of seed size in *Arabidopsis*. *Proc Natl Acad Sci U S A*.

102:17531-17536

- Magome H, Yamaguchi S, Hanada A, Kamiya Y and Oda K.** (2004) *dwarf and delayed-flowering 1*, a novel *Arabidopsis* mutant deficient in gibberellin biosynthesis because of overexpression of a putative AP2 transcription factor. *Plant J* 37: 720-729
- Maleck K, Levine A, Eulgem T, Morgen A, Schmid J, Lawton K, Dangl JL and Dietrich RA.** (2000) The transcriptome of *Arabidopsis thaliana* during systemic acquired resistance. *Nat Genet* 26:403-410
- Mare C, Mazzucotelli E, Crosatti C, Francia E, Stanca AM, and Cattivelli L.** (2004). Hv-WRKY38: a new transcription factor involved in cold- and drought-response in barley. *Plant Mol Biol* 55: 399-416
- Medina J, Bargues M, Terol J, Pérez-Alonso M and Salinas J.** (1999) The *Arabidopsis* CBF gene family is composed of three genes encoding AP2 domain-containing proteins whose expression is regulated by low temperature but not by abscisic acid or dehydration. *Plant Physiol* 119:463-470
- Mengiste T, Chen X, Salmeron J and Dietrich R.** (2003) The *BOTRYTIS SUSCEPTIBLE1* gene encodes an R2R3MYB transcription factor protein that is required for biotic and abiotic stress responses in *Arabidopsis*. *Plant Cell* 15:2551–2565
- Miao Y, Laun T, Zimmermann P and Zentgraf U.** (2004) Targets of the WRKY53 transcription factor and its role during leaf senescence in *Arabidopsis*. *Plant Mol Biol* 55:853-867
- Mikolajczyk M, Olubunmi SA, Muszynska G, Klessig DF and Dobrowolska G.** (2000) Osmotic stress induces rapid activation of a salicylic acid-induced protein kinase and a homolog of protein kinase ASK1 in tobacco cells. *Plant Cell* 12, 165–178
- Nakagami H, Pitzschke A and Hirt H.** (2005) Emerging MAP kinase pathways in plant stress signaling. *Trends Plant Sci*10:339-346
- Nakagami H, Soukupová H, Schikora A, Zárský V and Hirt H.** (2006) A mitogen-activated protein kinase kinase kinase mediates reactive oxygen species homeostasis in *Arabidopsis*. *J Biol Chem* 281:38697-38704
- Ohno S.** (1970) *Evolution by gene duplication*. Springer-Verlag, Berlin.
- Park JM, Park CJ, Lee SB, Ham BK, Shin R and Paek KH.** (2001) Overexpression of the tobacco Ts11 gene encoding an EREBP/AP2-type transcription factor enhances

resistance against pathogen attack and osmotic stress in tobacco. *Plant Cell* 13:1035-1046

Pfaffl MW. (2001) A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Res* 29: e45

Poppenberger B, Berthiller F, Lucyshyn D, Sieberer T, Schuhmacher R, Krska R, Kuchler K, Glössl J, Luschnig C and Adam G. (2003) Detoxification of the *Fusarium* mycotoxin deoxynivalenol by a UDP-glucosyltransferase from *Arabidopsis thaliana*. *J Biol Chem* 278: 47905-47914

Qiu QS, Guo Y, Dietrich MA, Schumaker KS and Zhu J-K. (2002) Regulation of SOS1, a plasma membrane Na⁺/H⁺ exchanger in *Arabidopsis thaliana*, by SOS2 and SOS3. *Proc Natl Acad Sci USA* 99, 8436–8441

Rizhsky L, Davletova S, Liang H and Mittler R. (2004) The zinc finger protein Zat12 is required for cytosolic ascorbate peroxidase 1 expression during oxidative stress in *Arabidopsis*. *J Biol Chem* 279:11736-11743

Robatzek S, and Somssich IE. (2001) A new member of the *Arabidopsis* WRKY transcription factor family, AtWRKY6, is associated with both senescence- and defence-related processes. *Plant J* 28, 123-133

Rushton PJ, Torres JT, Parniske M, Wernert P, Hahlbrock K and Somssich IE. (1996) Interaction of elicitor-induced DNA-binding proteins with elicitor response elements in the promoters of parsley PR1 genes. *EMBO J* 15, 5690–5700

Salinas J and Sanchez-Serrano JJ. (2006) *Arabidopsis* Protocols: 2nd Ed. Humana Press, NJ, USA

Sambrook J. and Russel DW. (2001) *Molecular Cloning: a Laboratory Manual*, 3rd edn. Cold Spring Harbor: Cold Spring Harbor Laboratory Press. New York

Schindler U, Beckmann H and Cashmore AR. (1992) TGA1 and G-box binding factors: two distinct classes of *Arabidopsis* leucine zipper proteins compete for the G-box-like element TGACGTGG. *Plant Cell* 4, 1309-1319

Schwacke R, Fischer K, Ketelsen B, Krupinska K and Krause K. (2007) Comparative survey of plastid and mitochondrial targeting properties of transcription factors in *Arabidopsis* and rice. *Mol Genet Genomics* 277:631-646

Seki M, Narusaka M, Ishida J, Nanjo T, Fujita M, Oono Y, Kamiya A, Nakajima

- M, Enju A, Sakurai T, Satou M, Akiyama K, Taji T, Yamaguchi-Shinozaki K, Carninci P, Kawai J, Hayashizaki Y and Shinozaki K.** (2002) Monitoring the expression profiles of 7000 *Arabidopsis* genes under drought, cold and high-salinity stresses using a full-length cDNA microarray. *Plant J* 31: 279-292
- Seki M, Kamei A, Yamaguchi-Shinozaki K and Shinozaki K.** (2003) Molecular responses to drought, salinity and frost: common and different paths for plant protection. *Curr Opin Biotechnol* 14:194-199
- Sessions A, Burke E, Presting G, Aux G, McElver J, Patton D, Dietrich B, Ho P, Bacwaden J, Ko C, Clarke JD, Cotton D, Bullis D, Snell J, Miguel T, Hutchison D, Kimmerly B, Mitzel T, Katagiri F, Glazebrook J, Law M and Goff SA.** (2002) A high-throughput *Arabidopsis* reverse genetics system. *Plant Cell* 14:2985-2994
- Shi H, Ishitani M, Kim C and Zhu J-K.** (2000) The *Arabidopsis thaliana* salt tolerance gene *SOS1* encodes a putative Na⁺/H⁺ antiporter. *Proc Natl Acad Sci USA* 97, 6896–6901
- Shinozaki K and Yamaguchi-Shinozaki K.** (1997) Gene expression and signal transduction in water-stress response. *Plant Physiol* 115, 327-334
- Shinozaki K, Yamaguchi-Shinozaki K and Seki M.** (2003) Regulatory network of gene expression in the drought and cold stress responses. *Curr Opin Plant Biol* 6, 410–417
- Singh K, Foley RC and Oñate-Sánchez L.** (2002) Transcription factors in plant defense and stress responses. *Curr Opin Plant Biol* 5, 430–436
- Skriver K, Olsen FL, Rogers JC and Mundy J.** (1991) cis-acting DNA elements responsive to gibberellin and its antagonist abscisic acid. *Proc Natl Acad Sci USA*. 88:7266-7270
- Stockinger EJ, Gilmour SJ and Thomashow MF.** (1997) *Arabidopsis thaliana* CBF1 encodes an AP2 domain-containing transcriptional activator that binds to the C-repeat/DRE, a cis-acting DNA regulatory element that stimulates transcription in response to low temperature and water deficit. *Proc Natl Acad Sci U S A* 94:1035-1040
- Su SH, Suarez-Rodriguez MC and Krysan P.** (2007) Genetic interaction and phenotypic analysis of the *Arabidopsis* MAP kinase pathway mutations *mekk1* and *mpk4* suggests signaling pathway complexity. *FEBS Lett* 581:3171-3177

- Sun C, Palmqvist S, Olsson H, Bore' n M, Ahlandsberg S and Jansson C.** (2003) A novel WRKY transcription factor, SUSIBA2, participates in sugar signaling in barley by binding to the sugar-responsive elements of the iso1 promoter. *Plant Cell* 15:2076-2092
- Suzuki Y, Kawazu T and Koyama H.** (2004) RNA isolation from siliques, dry seeds, and other tissues of *Arabidopsis thaliana*. *Biotechniques* 37: 542, 544
- Taylor CB.** (1997) Promoter fusion analysis: an insufficient measure of gene expression. *Plant Cell* 9, 273–275
- Teige M, Scheikl E, Eulgem T, Dóczy R, Ichimura K, Shinozaki K, Dangl JL and Hirt H.** (2004) The MKK2 pathway mediates cold and salt stress signaling in *Arabidopsis*. *Mol Cell* 15:141-152
- Thompson JD, Gibson TJ, Plewniak F, Jeanmougin F and Higgins DG.** (1997) The CLUSTAL_X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res* 25:4876-4882
- Tian Q and Reed JW.** (1999) Control of auxin-regulated root development by the *Arabidopsis thaliana* *SHY2/LAA3* gene. *Development* 126:711-721
- Turck F, Zhou A and Somssich IE.** (2004) Stimulus-dependent, promoter-specific binding of transcription factor WRKY1 to its native promoter and the defense-related gene *PcPRI-1* in parsley. *Plant Cell* 16: 2573–2585
- Ulker B and Somssich IE.** (2004) WRKY transcription factors: from DNA binding towards biological function. *Curr Opin Plant Biol* 7, 491-498
- Ulm R, Ichimura K, Mizoguchi T, Peck SC, Zhu T, Wang X, Shinozaki K and Paszkowski J.** (2002) Distinct regulation of salinity and genotoxic stress responses by *Arabidopsis* MAP kinase phosphatase 1. *EMBO J* 21, 6483-6493
- Uno Y, Furihata T, Abe H, Yoshida R, Shinozaki K and Yamaguchi Shinozaki K.** (2000) *Arabidopsis* basic leucine zipper transcription factors involved in an abscisic acid-dependent signal transduction pathway under drought and high-salinity conditions, *Proc Natl Acad Sci USA* 97: 11632–11637
- van der Graaff E, Hooykaas PJJ and Keller B.** (2002) Activation tagging of the two closely linked genes *LEP* and *VAS* independently affects vascular cell number. *Plant J* 32, 819-830

- Verslues PE, Agarwal M, Katiyar-Agarwal S, Zhu J and Zhu JK.** (2006) Methods and concepts in quantifying resistance to drought, salt and freezing, abiotic stresses that affect plant water status. *Plant J* 45:523-539
- Wadsworth GJ, Redinbaugh MG and Scandalios JG.** (1988) A procedure for the small-scale isolation of plant RNA suitable for RNA blot analysis. *Anal Biochem* 172:279-283
- Wang ZY and Tobin EM.** (1998) Constitutive expression of the *CIRCADIAN CLOCK ASSOCIATED 1 (CCA1)* gene disrupts circadian rhythms and suppresses its own expression. *Cell* 93:1207-1217
- Weigel D and Glazebrook J.** (2002) *Arabidopsis: A Laboratory Manual*. Cold Spring Harbor Laboratory Press, NY, USA
- Wu KL, Guo ZJ, Wang HH and Li J.** (2005) The WRKY family of transcription factors in rice and *Arabidopsis* and their origins. *DNA Res* 12:9-26
- Xie Z, Zhang ZL, Zou XL, Huang J, Ruas P, Thompson D, and Shen QJ.** (2005) Annotations and functional analyses of the rice WRKY gene superfamily reveal positive and negative regulators of abscisic acid signaling in aleurone cells. *Plant Physiol* 137, 176–189
- Xiong L, Ishitani M, Lee H and Zhu J-K.** (2001) The *Arabidopsis* *LOS5/ABA3* locus encodes a molybdenum cofactor sulfuryase and modulates cold stress- and osmotic stress- responsive gene expression. *Plant Cell* 13:2063-2083
- Xiong L, Lee H, Ishitani M and Zhu J-K.** (2002a) Regulation of osmotic stress responsive gene expression by *LOS6/ABA1* locus in *Arabidopsis*. *J Biol Chem* 277: 8588-8596
- Xiong L, Schumaker K and Zhu J-K.** (2002b) Cell signaling during cold, drought, and salt stress. *Plant Cell* 14: S165-S183
- Xu X, Chen C, Fan B and Chen Z.** (2006) Physical and functional interactions between pathogen-induced *Arabidopsis* WRKY18, WRKY40, and WRKY60 transcription factors. *Plant Cell* 18:1310-1326
- Xu YH, Wang JW, Wang S, Wang JY and Chen XY.** (2004) Characterization of GaWRKY1, a cotton transcription factor that regulates the sesquiterpene synthase gene (+)-delta-cadinene synthase-A. *Plant Physiol* 135, 507–515

- Xu Z, Escamilla-Trevino L, Zeng L, Lalgondar M, Bevan D, Winkel B, Mohamed A, Cheng CL, Shih MC, Poulton J and Esen A.** (2004) Functional genomic analysis of *Arabidopsis thaliana* glycoside hydrolase family 1. *Plant Mol Biol* 55: 343-367
- Yamaguchi-Shinozaki K and Shinozaki K.** (1993) Characterization of the expression of a desiccation-responsive *rd29* gene of *Arabidopsis thaliana* and analysis of its promoter in transgenic plants. *Mol Gen Genet* 236:331-340
- Yamaguchi-Shinozaki K and Shinozaki K.** (1994) A novel cis-acting element in an *Arabidopsis* gene is involved in responsiveness to drought, low-temperature, or high-salt stress. *Plant Cell* 6, 251–264
- Yang LX, Wang RY, Ren F, Liu J, Cheng J and Lu YT.** (2005) AtGLB1 enhances the tolerance of *Arabidopsis* to hydrogen peroxide stress. *Plant Cell Physiol* 46:1309-1316
- Yang P, Chen C, Wang Z, Fan B and Chen Z.** (1999) A pathogen- and salicylic acid-induced WRKY DNA-binding activity recognizes the elicitor response element to the tobacco class I chitinase gene promoter. *Plant J* 18, 141–149
- Yoda H, Ogawa M, Yamaguchi Y, Koizumi N, Kusano T and Sano H.** (2002) Identification of early-responsive genes associated with the hypersensitive response to tobacco mosaic virus and characterization of a WRKY-type transcription factor in tobacco plants. *Mol Genet Gen* 267, 154–161
- Zhang JZ.** (2003) Overexpression analysis of plant transcription factors. *Curr Opin Plant Biol* 6:430-440
- Zhang Y and Wang L.** (2005) The WRKY transcription factor superfamily: its origin in eukaryotes and expansion in plants. *BMC Evol Biol* 5:1
- Zhang ZL, Xie Z, Zou X, Casaretto J, Ho TD and Shen QJ.** (2004) A rice WRKY gene encodes a transcriptional repressor of the gibberellin signaling pathway in aleurone cells. *Plant Physiol* 134, 1500–1513
- Zheng Z, Qamar SA, Chen Z and Mengiste T.** (2006) *Arabidopsis* WRKY33 transcription factor is required for resistance to necrotrophic fungal pathogens. *Plant J* 48:592-605
- Zheng Z, Mosher SL, Fan B, Klessig DF and Chen Z.** (2007) Functional analysis of *Arabidopsis* WRKY25 transcription factor in plant defense against *Pseudomonas*

syringae. BMC Plant Biol 7: 2

Zhu J-K. (2001) Cell signaling under salt, water and cold stresses. Curr Opin Plant Biol 4: 401-406

Zhu J-K. (2002) Salt and drought stress signal transduction in plants. Annu Rev Plant Biol 53, 247-273

Zhu J-K. (2003) Regulation of ion homeostasis under salt stress. Curr Opin Plant Biol 6: 441-445

Zou X, Seemann JR, Neuman D and Shen QJ. (2004) A WRKY gene from *Creosote Bush* encodes an activator of the abscisic acid signaling pathway. J Biol Chem 279, 55770-55779

Chapter 6: General discussion and conclusions

Plant growth and productivity are adversely affected by a variety of abiotic stresses, such as low temperature, salt, drought, flooding, heat, oxidative stress and heavy metal toxicity. Plants are also challenged by pathogens including bacteria, fungi, and viruses as well as herbivores. All these stress factors are a menace for plants and prevent them from reaching their full genetic potential and limit the crop productivity worldwide. Abiotic stress is the major cause of crop failure worldwide, and it causes losses worth hundreds of millions of dollars each year. Because of increases in global population, food productivity also needs to be increased. Therefore, it is important to develop stress tolerant crops.

In response to abiotic stress conditions, expression of various genes in plants is up-regulated, which can mitigate the effect of stress and lead to adjustment of the cellular milieu and plant tolerance (Shinozaki and Yamaguchi-Shinozaki, 2000; Xiong and Zhu 2001). During the past two decades, great progress has been achieved in the study of abiotic stress, including identification of key components and the associated signaling pathways through a combination of forward and reverse genetic strategies (Xiong et al., 2002b; Yamaguchi-Shinozaki and Shinozaki K, 2006). In nature, stress does not generally come in isolation and many different stresses may challenge plants at the same time. Hence, there exists cross talk between abiotic stresses (Xiong *et al.*, 2002; Shinozaki and Yamaguchi-Shinozaki, 2000; Zhu 2002; Yamaguchi-Shinozaki and Shinozaki, 2006).

Among the abiotic stresses, soil salinity is one of the dominant factors limiting the harvestable yields of crop plants. Secondary salinization caused by poor irrigation water and subsequent drainage is damaging productive agricultural land worldwide. High salinity causes both hyperionic and hyperosmotic stress and can lead to plant demise. High salt depositions in the soil generate a low water potential zone in the soil making it increasingly difficult for the plant to acquire both water as well as nutrients. Therefore, salt stress essentially results in a water deficit condition in the plant and this is similar to drought. Hence, it is essential to understand the control of ion homeostasis and osmotic regulation, and to use this knowledge to engineer crop plants with

enhanced salt tolerance. *Arabidopsis* is a small model flowering plant that has many advantages. Research on salt toxicity and tolerance using *Arabidopsis* has produced significant results, and it is now widely held that NHX1 on the vacuolar membrane and SOS1 on the plasma membrane play an important role in decreasing the cytosolic sodium concentration (Apse and Blumwald, 2002; Zhu 2002), which is the basic mechanism for developing salt-tolerant plants. However, much is still unknown about the elusive salt-stress sensor and regulators of ion transporters.

Conventional methods of breeding salt-tolerant crops are rather slow and have had only limited success (Wang *et al.*, 2003), partly due to the fact that plant responses to salt stress are complex and the underlying mechanisms remain unclear. Developing salt-tolerant crops through genetic engineering, i.e. introducing genes from one plant to another or overexpressing them in plants has demonstrated exciting success (Bartels and Sunkar, 2005; Yamaguchi and Blumwald, 2005; Zhang *et al.*, 2005). To identify novel salt-responsive genes and proteins will not only contribute to understanding salt signaling mechanisms but also provide new target genes for engineering stress-tolerant plants.

Microarray is a widely used technology to identify stress-responsive genes in many plants (Kawasaki *et al.*, 2001; Seki *et al.*, 2002a, 2002b, Kreps *et al.*, 2002; Rabbani *et al.*, 2003; Wang *et al.*, 2003; Chao *et al.*, 2005; Ma *et al.*, 2006). With the completion of sequencing of genomes of *Arabidopsis* and rice, microarrays including almost the whole sets of genes encoded by *Arabidopsis* and rice genomes have been developed and widely used to identify stress-responsive genes (Chao *et al.*, 2005; Ma *et al.*, 2006). Root is the first organ to sense higher concentration of salt in the soil. Therefore, it is necessary to focus on root for salt stress research. However, there is no report using root as the material to study salt stress response. To identify a comprehensive set of salt-responsive genes, we used the QiagenOperon 26 k oligonucleotide microarray to monitor the transcriptome changes in *Arabidopsis* roots under the treatment of a moderate NaCl concentration (150 mM) by 6, 24 and 48 h (Jiang and Deyholos, 2006). Statistical analysis of our data enabled us to identify many novel salt-responsive genes and highlight potentially important roles for underappreciated gene families, including: several groups of transporters (e.g. MATE,

LeOPT1-like); signalling molecules (e.g. PERK kinases, MLO-like receptors), carbohydrate active enzymes (e.g. XTH18), transcription factors (e.g. members of ZIM, WRKY, NAC), and other proteins (e.g. 4CL-like, COMT-like, LOB-Class 1). The newly identified salt-responsive genes demonstrated the power of full-genome microarray and may be selected as target genes for developing stress-tolerant crops in the future.

Proteomics based on 2-D gel electrophoresis (2-DGE) is another powerful technology for identifying stress-responsive proteins. Coupled with mass spectrometry, 2-DGE has been applied to identify environmental stress responsive proteins in many plants (Agrawal *et al.*, 2005; Yan *et al.*, 2005 & 2006; Hajheidari *et al.*, 2005, 2007; Lee *et al.*, 2007; Li *et al.*, 2007; Nohzadeh *et al.*, 2007; Plomion *et al.*, 2006; Yang *et al.*, 2007). The type of 2-DGE-based comparative proteomics has yielded interesting new insights into stress responses in plants (Yan *et al.*, 2006; Hajheidari *et al.*, 2007; Nohzadeh *et al.*, 2007; Yang *et al.*, 2007). This is partly explained by the fact that mRNA levels of some genes do not show significant changes under stress condition and therefore fail to be captured by RNA-based techniques such as differential display (DD), suppressive subtractive hybridization (SSH), SAGE (serial analysis of gene expression) and microarray, whereas the changes at the protein levels under stress conditions are above a threshold and thus are detected by the 2-DGE based technology. Besides, mRNA levels are not always correlated with those of corresponding proteins, because of post-transcriptional regulation, such as post-translational modifications (PTMs) including glycosylation and phosphorylation. It has been reported that only moderate correlation exists between changes in the levels of specific mRNAs and their corresponding proteins in studies involving yeast (*Saccharomyces cerevisiae*), animals, or Arabidopsis (Gygi *et al.*, 1999; Tian *et al.*, 2004; Mooney *et al.*, 2006). DNA sequence and mRNA expression profiling provide little information about sub-cellular localization, beyond the identification of putative targeting signal sequences and structural domains. These biological realities motivated us to perform an analysis of NaCl stress responses at the proteome level, and to compare these results to our microarray-based studies of similarly treated Arabidopsis roots. We conducted a comparative proteomic analysis of Arabidopsis roots that had been exposed to 150 mM

NaCl for 6 or 48 h through utilizing two-dimensional electrophoresis coupled to tandem mass spectrometry (Jiang *et al.*, 2007). We successfully identified 86 of the differentially abundant proteins, which are implicated in ROS scavenging, signal transduction, translation, cell wall biosynthesis, protein translation, processing and degradation, and metabolism of energy, amino acids, and hormones. At the resolution of approximately 70 individual genes/proteins, we detected poor statistical correlation of these protein expression data with our previous microarray results, supporting the concept that post-transcriptional regulation plays an important role in stress-responsive gene expression (Jiang *et al.*, 2007). However, more proteins need to be identified to determine the generality of this conclusion under different abiotic stress treatments and measurement techniques.

2-DGE-based proteomic study has its own limitations, i.e. inability to resolve membrane proteins and detect low-abundant proteins (Rose *et al.*, 2004), therefore, complementary strategies at the protein and metabolite levels have been developed and are used to gain more insight into the intricate network of plant response to environmental stimuli. One of these is multidimensional protein identification technology (MudPIT), which was developed to allow automated analyses of peptide mixtures generated from complex protein samples (Link *et al.*, 1999). However, while MudPIT is able to provide a catalogue of all the proteins present in a particular sample, it does not generate useful quantitative information and even reliable qualitative comparisons between samples are still technically challenging and time consuming. A second technique, isotope-coded affinity tag (ICAT) in the LC-MS/MS system has also emerged as a tool for quantitative proteomics (Han *et al.*, 2001). However, the quantitative reproducibility and the number of replicate experiments required for statistical significance are yet to be completely resolved and application of ICAT has not been reported in plants (Rabilloud, 2002).

A common problem with current salt stress study is that many experiments are not designed to expose genes that confer salinity tolerance under natural conditions. Many experiments with NaCl do not include supplemental Ca^{2+} , and even NaCl concentrations as high as 250 or 300 mM have been given in one treatment, with no additional Ca^{2+} . It is widely reported that supplemental Ca^{2+} ameliorates Na^+ toxicity

symptoms in different plant species (Rengel, 1992). Supplemental Ca^{2+} may provide protection against salinity stress through stabilizing cell wall components, plasma membrane lipids, and proteins (Rengel, 1992; Kinraide, 1998). Supplemental Ca^{2+} in a physiological concentration range (1–10 mM) reduced, or even completely prevented, Na^+ -induced K^+ efflux from salinized root and leaf tissues in broad bean (*Vicia faba*) and barley (*Hordeum vulgare*) plants (Shabala, 2000; Shabala *et al.*, 2003). Extracellular Ca^{2+} ameliorates NaCl -induced K^+ loss from Arabidopsis root and leaf cells by controlling plasma membrane K^+ -permeable channels (Shabala *et al.*, 2006). In our microarray and 2-DGE experiments, Arabidopsis plants were treated by $\frac{1}{2}$ x MS with or without 150 mM NaCl . The concentration of CaCl_2 in $\frac{1}{2}$ x MS solution is 1.5 mM, which is within the range of physiological concentration. However, the issue with our NaCl treatment is that 150 mM NaCl was applied in one hit. Instead, the concentration of NaCl could start with a low concentration of NaCl , i.e. 50 mM, and then gradually increased to be 150 mM within a few days. In this way, more natural salt-responsive genes may be identified.

The discrepancy or low correlation between our microarray and 2-DGE data for some gene/protein was also observed. For instance, the manganese superoxide dismutase (MSD1, At3g10920) was identified to be down-regulated at the transcript level by salt treatment, but it was identified to up-regulated at the protein level (Table 3-1). The nucleoside-diphosphate kinase (NDPK1, At4g09320) was identified to be down-regulated at the transcript level by salt treatment, but it was not found to be repressed at the protein level (Table 3-1). To resolve the discrepancy between microarray and 2-DGE data, Western blotting using specific antibody could be used to validate the 2-DGE data at the protein level. However, this is not very feasible as it is not easy to obtain specific antibodies for these proteins, although some antibodies are commercially available from Agrisera (<http://www.agrisera.com/index.php>). Alternatively, for those proteins implicated in ROS scavenging such as SOD or APX, enzymatic assays could be performed to examine their activities in salt-treated roots relative to control.

Three transcription factor genes, *bHLH92*, *WRKY25* and *WRKY33*, whose transcript levels were found to be highly induced by 150 mM NaCl in Arabidopsis

roots, were chosen for further characterization. The bHLH transcription factor gene family is the second largest after MYB in Arabidopsis, and previous studies have shown that members of the bHLH family are involved in diverse physiological processes including cold (Chinnusamy *et al.*, 2003), ABA (Abe *et al.*, 2003), JA (Lorenzo *et al.*, 2004) and more recently, salt stress (Kim and Kim, 2006) signalling. Our study of *bHLH92* demonstrate its abundance was increased by salt, drought, mannitol, JA and cold treatments (Fig. 4-4B), suggesting that *bHLH92* is responsive to multiple abiotic stresses. We also found that the induction of *bHLH92* by NaCl is partly ABA- and SOS2-dependent but independent of SOS1 and SOS3 (Fig. 4-5). Promoter-GUS fusion assay demonstrate that promoter of *bHLH92* is salt, drought and JA inducible (Fig. 4-6). Knock-out mutant of *bHLH92* was sensitive to osmotic stress and slightly to NaCl stress (Fig. 4-8), while constitutive overexpression of *bHLH92* in Arabidopsis increased plants' tolerance to salt, osmotic and oxidative stresses (Fig. 4-9), which indicate that *bHLH92* plays an important role in stress signaling and tolerance. Microarray profiling of *bhlh92* mutant under NaCl treatment identified 19 genes that were differentially expressed between mutant and wild-type, and some genes were known to be involved in abiotic stress or associated with ROS scavenging (Table 4-2). Future study on *bHLH92* can be directed to identify upstream gene(s) regulating bHLH92 activity and its interacting partners by using yeast two-hybrid system or tandem affinity purification (TAP) coupled to LC/MS identification as demonstrated by previous studies (Rohila *et al.*, 2006; Miao and Zentgraf, 2007). We observe that the difference between mutant and WT under stress condition is not very large, and this may suggest a functional redundancy exist between some members of bHLH gene family. Therefore, double, triple or even quadruple knockout mutants for between *bHLH92* and other evolutionally close members of bHLH gene family (Fig. 4-2) could be generated to perform comparative analysis of these composite knockout mutants so that both distinct and overlapping functions of these *bHLH* genes could be revealed. Further work could also involve investigating the transcript levels of abiotic and ABA stress responsive marker genes, such as *RD29A*, *RD29B*, *ERD10*, *KIN1* and *CBF2*, in the *bHLH92*-overexpression lines under NaCl, mannitol or ABA treatment.

WRKY transcription factors have been shown to play an important role in

pathogen responses and leaf senescence (Eulgem *et al.*, 2000; Eulgem and Somssich, 2007). WRKYs form a large gene family with 72 members in Arabidopsis. In a previous microarray profiling assay, 18 WRKY TFs, including *WRKY25* (At2g30250) and *WRKY33* (At2g38470) that were highly induced by 150 mM NaCl treatment in Arabidopsis roots, were identified (Jiang and Deyholos, 2006). Previous cDNA microarray also indentified a few Arabidopsis WRKY genes which are responsive to multiple abiotic stresses (Seki *et al.*, 2002a, 2002b). Recent researches have shown that *WRKY25* and *WRKY33* play an important role in mediating disease resistance (Zheng *et al.*, 2006, 2007; Lippok *et al.*, 2007), which is proposed to be regulated through a MAPK cascade (Andreasson *et al.*, 2005). We hypothesized that *WRKY25* and *WRKY33* might also play an important role in abiotic stress response such salt stress tolerance. To test this, we used a combination of reverse genetic methods. We first demonstrate that both *WRKY25* and *WRKY33* were induced by salt, mannitol and cold treatments as well as by other stress treatments (Fig. 5-6). We observed that the inducibility of *WRKY25* by salt stress was ABA- and SOS-signaling pathways independent, whereas the induction of *WRKY33* by salt stress was partially dependent on ABA biosynthesis and signaling but was independent of the SOS pathway (Fig. 5-7). Stable transformed Arabidopsis plants harboring PWRKY25- and PWRKY33-GUS fusion constructs revealed that both promoters were inducible by salt stress as well as by other specific abiotic stresses (Fig. 5-8). Knock-out mutant of *WRKY33* only showed slight sensitivity to salt stress while *WRKY25* knock-out mutants did not exhibit any significant difference from WT under tested stress conditions, suggesting a functional redundancy between evolutionally close WRKY TFs (Fig. 5-2). We further produced *wrky25wrky33* double mutant and found that this double mutant was more sensitive to NaCl treatment than wild-type (Fig. 5-9E & H). Overexpression of *WRKY25* and *WRKY33* separately in Arabidopsis increased plant tolerance to salt but were more sensitive to oxidative and ABA treatments at the seedling stage (Fig. 5-10). These data suggest that *WRKY25* and *WRKY33* are positive regulators of salt stress response while a negative regular of ABA and oxidative stress responses. Finally, full-genome microarray was employed to identify the target genes of *WRKY25* and *WRKY33* under salt treatment, and 31 and 208 differentially expressed genes between *wrky25* or

wrky33 single mutant and wild-type were identified, respectively (Tables 5-2 to 5-5). Further examination of the 1 kb upstream regions of these differential genes demonstrated that W-box (C/TTGACC/T), which is the *cis*-element recognized and bound by WRKY TFs, is highly enriched in the target gene promoter regions (Table 5-6). Moreover, the different numbers and identity of differential expressed genes identified suggest that WRKY25 and WRKY33 might play a role in salt stress by targeting divergent downstream genes. On the basis of our results as well as those reported previously (Ichimura et al., 2000; Teige *et al.*, 2004; Andreasson *et al.*, 2005), we present a model describing the interrelationships among MAPK signaling components, MEKK1, MKK2, MPK4 as well as WRKY25 and WRKY33 in the salt signal transduction pathway (Fig. 5-12).

Future study can be directed to identify the direct targets of WRKY25 and WRKY33 as well as their interacting partners by using yeast two-hybrid system or tandem affinity purification (TAP) coupled to LC/MS identification as demonstrated by previous studies (Andreasson *et al.*, 2005; Rohila *et al.*, 2006; Miao and Zentgraf, 2007). Further work could also involve investigating the transcript levels of abiotic and ABA stress responsive marker genes, such as *RD29A*, *RD29B*, *ERD10*, *KINI* and *CBF2*, in the overexpression lines under NaCl treatment. It can also be directed to generate double, triple and quadruple knockout mutants of *WRKY25* or *WRKY33* and other related WRKY genes. Further comparative analysis of these composite knockout mutants should reveal both distinct and overlapping functions of the WRKY genes.

In conclusions, elevated concentration of salt has pleiotropic effects on plant development, morphology and physiology. The changes at both transcript and protein levels in *Arabidopsis* roots under a moderate salt treatment were monitored using microarray and 2-DGE, respectively. Many differentially expressed genes/proteins were identified, among which many are novel. These help us to better understand plant response to salt stress in the context of regulatory network and data presented in the thesis provide alternative choices for engineering salt-tolerant crops in the future. Characterization of three novel salt-inducible transcription factor genes, *bHLH92*, *WRKY25* and *WRKY33*, using both knock-out and overexpression strategies demonstrate that they play important roles in plant protection against different abiotic

stresses. These findings described in the thesis thus allow a better understanding of the roles of two classes of previously underappreciated TFs, bHLH and WRKY in abiotic stress. The results on WRKY25 and WRKY33 characterization also help to understand the complex interaction and cross-talk between components implicated in biotic and abiotic stresses as they appear to be key regulators connecting biotic stress and abiotic stress signalings.

References

- Abe H, Urao T, Ito T, Seki M, Shinozaki K, and Yamaguchi-Shinozaki K. (2003)** Arabidopsis AtMYC2 (bHLH) and AtMYB2 (MYB) function as transcriptional activators in abscisic acid signaling. *Plant Cell* 15, 63–78
- Agrawal GK, Yonekura M, Iwahashi Y, Iwahashi H and Rakwal R. (2005)** System, trends and perspectives of proteomics in dicot plants. Part III: Unraveling the proteomes influenced by the environment, and at the levels of function and genetic relationships. *J Chromatogr B Analyt Technol Biomed Life Sci* 815:137-145
- Andreasson E, Jenkins T, Brodersen P, Thorgrimsen S, Petersen NH, Zhu S, Qiu JL, Micheelsen P, Rocher A, Petersen M, Newman MA, Bjorn Nielsen H, Hirt H, Somssich I, Mattsson O and Mundy J. (2005)** The MAP kinase substrate MKS1 is a regulator of plant defense responses. *EMBO J* 24:2579-2589
- Apse MP and Blumwald E. (2002)** Engineering salt tolerance in plants. *Curr Opin Biotechnol* 13:146-150
- Bartels D and Sunkar R. (2005)** Drought and Salt Tolerance in Plants. *Crit Rev Plant Sci* 24:23–58
- Chao DY, Luo YH, Shi M, Luo D and Lin HX. (2005)** salt-responsive genes in rice revealed by cDNA microarray analysis. *Cell Res* 15: 796-810
- Chinnusamy V, Ohta M, Kanrar S, Lee BH, Hong X, Agarwal M and Zhu JK. (2003)** ICE1: a regulator of cold-induced transcriptome and freezing tolerance in Arabidopsis. *Genes Dev* 17:1043-1054
- Eulgem T, Rushton PJ, Robatzek S, and Somssich IE. (2000)** The WRKY superfamily of plant transcription factors. *Trends Plant Sci* 5, 199–206
- Eulgem T and Somssich IE. (2007)** Networks of WRKY transcription factors in

defense signaling. *Curr Opin Plant Biol* 10:366–371

Gygi SP, Rochon Y, Franz BR and Aebersold R. (1999) Correlation between protein and mRNA abundance in yeast. *Mol Cell Biol* 19:1720-1730

Hajheidari M, Abdollahian-Noghabi M, Askari H, Heidari M, Sadeghian SY, Ober ES, Salekdeh GH. (2005) Proteome analysis of sugar beet leaves under drought stress. *Proteomics*. 5:950-960

Hajheidari M, Eivazi A, Buchanan BB, Wong JH, Majidi I, Salekdeh GH. (2007) Proteomics uncovers a role for redox in drought tolerance in wheat. *J Proteome Res*. 6:1451-1460

Han DK, Eng J, Zhou H, Aebersold R. (2001) Quantitative profiling of differentiation-induced microsomal proteins using isotope-coded affinity tags and mass spectrometry. *Nat Biotech* 19, 946-951

Ichimura K, Mizoguchi T, Yoshida R, Yuasa T and Shinozaki K. (2000) Various abiotic stresses rapidly activate Arabidopsis MAP kinases ATMPK4 and ATMPK6. *Plant J* 24, 655-665

Jiang Y-Q and Deyholos MK. (2006) Comprehensive transcriptional profiling of NaCl-stressed Arabidopsis roots reveals novel classes of responsive genes. *BMC Plant Biol* 6, 25

Jiang Y-Q, Yang B, Harris NS and Deyholos MK. (2007) Comparative proteomic analysis of NaCl stress-responsive proteins in Arabidopsis roots. *J Exp Bot* 58: 3591-3607

Kawasaki S, Borchert C, Deyholos M, Wang H, Brazille S, Kawai K, Galbraith D and Bohnert HJ. (2001) Gene expression profiles during the initial phase of salt stress in rice (*Oryza sativa* L.). *Plant Cell* 13: 889-905

Kim J and Kim H-Y. (2006) Functional analysis of a calcium-binding transcription factor involved in plant salt stress signaling. *FEBS Lett* 580:5251-5256

Kinraide TB. (1998) Three mechanisms for the calcium alleviation of mineral toxicities. *Plant Physiol* 118: 513–520

Kreps JA, Wu Y, Chang H, Zhu T, Wang X and Harper JF. (2002) Transcriptome changes for *Arabidopsis* in response to salt, osmotic, and cold stress. *Plant Physiol* 130, 2129-2141

- Lee DG, Ahsan N, Lee SH, Kang KY, Bahk JD, Lee IJ and Lee BH.** (2007) A proteomic approach in analyzing heat-responsive proteins in rice leaves. *Proteomics* 7: 3369-3383
- Lorenzo O, Chico JM, Sanchez-Serrano JJ and Solano R.** (2004) *JASMONATE-INSENSITIVE1* encodes a MYC transcription factor essential to discriminate between different jasmonate regulated defense responses in *Arabidopsis*. *Plant Cell* 16:1938–1950
- Li K, Xu C, Zhang K, Yang A and Zhang J.** (2007) Proteomic analysis of roots growth and metabolic changes under phosphorus deficit in maize (*Zea mays* L.) plants. *Proteomics*. 7:1501-1512
- Link AJ, Eng J, Schieltz DM, Carmack E, Mize GJ, Morris DR, Garvik BM, Yates JR III.** (1999) Direct analysis of protein complexes using mass spectrometry. *Nat Biotechnol* 17:676-682
- Lippok B, Birkenbihl RP, Rivory G, Brümmer J, Schmelzer E, Logemann E and Somssich IE.** (2007) Expression of AtWRKY33 encoding a pathogen- or PAMP-responsive WRKY transcription factor is regulated by a composite DNA motif containing W box elements. *Mol Plant Microbe Interact* 20:420-429
- Ma SS, Gong QQ and Bohnert HJ.** (2006) Dissecting salt stress pathways. *J Exp Bot* 57:1097-1107
- Miao Y and Zentgraf U.** (2007) The antagonist function of *Arabidopsis* WRKY53 and ESR/ESP in leaf senescence is modulated by the jasmonic and salicylic acid equilibrium. *Plant Cell* 19:819-830
- Mooney BP, Miernyk JA, Greenlief CM and Thelen JJ.** (2006) Using quantitative proteomics of *Arabidopsis* roots and leaves to predict metabolic activity. *Physiol Plant* 128: 237-250
- Nohzadeh Malakshah S, Habibi Rezaei M, Heidari M and Hosseini Salekdeh G.** (2007) Proteomics reveals new salt responsive proteins associated with rice plasma membrane. *Biosci Biotechnol Biochem.* 71:2144-2154
- Plomion C, Lalanne C, Claverol S, et al.** (2006) Mapping the proteome of poplar and application to the discovery of drought-stress responsive proteins. *Proteomics* 6, 6509-6527

- Rabbani MA, Maruyama K, Abe H, Khan MA, Katsura K, Ito Y, Yoshiwara K, Seki M, Shinozaki K and Yamaguchi-Shinozaki K. (2003)** Monitoring expression profiles of rice genes under cold, drought, and high-salinity stresses and abscisic acid application using cDNA microarray and RNA gel-blot analyses. *Plant Physiol* 133:1755-1767
- Rabilloud T. (2002)** Two-dimensional gel electrophoresis in proteomics: old, old fashioned, but it still climbs up the mountains. *Proteomics* 2: 3-10
- Rengel Z. (1992)** The role of calcium in salt toxicity. *Plant Cell Environ* 15: 625–632
- Rohila JS, Chen M, Chen S, Chen J, Cerny R, Dardick C, Canlas P, Xu X, Gribskov M, Kanrar S, Zhu JK, Ronald P and Fromm ME. (2006)** Protein-protein interactions of tandem affinity purification-tagged protein kinases in rice. *Plant J* 46:1-13
- Rose JKC, Bashir S, Giovannoni JJ, Jahn MM and Saravanan RS. (2004)** Tackling the plant proteome: practical approaches, hurdles and experimental tools. *Plant J* 39, 715–733
- Seki M, Ishida J, Narusaka M, Fujita M, Nanjo T, Umezawa T, Kamiya A, Nakajima M, Enju A, Sakurai T, Satou M, Akiyama K, Yamaguchi-Shinozaki K, Carninci P, Kawai J, Hayashizaki Y and Shinozaki K. (2002a)** Monitoring the expression pattern of around 7,000 *Arabidopsis* genes under ABA treatments using a full-length cDNA microarray. *Funct Integr Genomics*. 2: 282-291
- Seki M, Narusaka M, Ishida J, , Nanjo T, Fujita M, Oono Y, Kamiya A, Nakajima M, Enju A, Sakurai T, Satou M, Akiyama K, Taji T, Yamaguchi-Shinozaki K, Carninci P, Kawai J, Hayashizaki Y and Shinozaki K. (2002b)** Monitoring the expression profiles of 7000 *Arabidopsis* genes under drought, cold and high-salinity stresses using a full-length cDNA microarray. *Plant J* 31: 279-292
- Shabala S, Demidchik V, Shabala L, Cuin TA, Smith SJ, Miller AJ, Davies JM and Newman IA. (2006)** Extracellular Ca^{2+} ameliorates NaCl-induced K^{+} loss from *Arabidopsis* root and leaf cells by controlling plasma membrane K^{+} -permeable channels. *Plant Physiol*. 141:1653-1665
- Shabala S and Newman I. (2000)** Salinity effects on the activity of plasma membrane H^{+} and Ca^{2+} transporters in bean leaf mesophyll: masking role of the cell wall. *Ann Bot*

(Lond) 85: 681–686

Shabala S, Shabala L and van Volkenburgh E. (2003) Effect of calcium on root development and root ion fluxes in salinised barley seedlings. *Funct Plant Biol* 30: 507–514

Shinozaki K and Yamaguchi-Shinozaki K. (2000) Molecular responses to dehydration and low temperature: differences and cross-talk between two stress signaling pathways. *Curr Opin Plant Biol* 3:217-223

Teige M, Scheikl E, Eulgem T, Dóczi R, Ichimura K, Shinozaki K, Dangl JL and Hirt H. (2004) The MKK2 pathway mediates cold and salt stress signaling in *Arabidopsis*. *Mol Cell* 15:141-152

Tian Q, Stepaniants SB, Mao M, Weng L, Feetham MC, Doyle MJ, Yi EC, Dai HY, Thorsson V, Eng J, Goodlett D, Berger JP, Gunter B, Linseley PS, Stoughton RB, Aebersold R, Collins SJ, Hanlon WA and Hood LE. (2004) Integrated genomic and proteomic analyses of gene expression in mammalian cells. *Mol Cell Proteomics* 3:960-969

Wang H, Miyazaki S, Kawai K, Deyholos M, Galbraith DW and Bohnert HJ. (2003) Temporal progression of gene expression responses to salt shock in maize roots. *Plant Mol Biol* 52: 873-891

Wang W, Vinocur B and Altman A. (2003) Plant responses to drought, salinity and extreme temperatures: towards genetic engineering for stress tolerance. *Planta* 218: 1–14

Xiong L and Zhu JK. (2001) Abiotic stress signal transduction in plants: Molecular and genetic perspectives. *Physiol Plant* 112:152-166

Xiong L, Schumaker KS and Zhu J-K. (2002b). Cell signaling during cold, drought, and salt stresses. *Plant Cell* 14: S165-183

Yamaguchi T and Blumwald E. (2005) Developing salt-tolerant crop plants: challenges and opportunities. *Trends Plant Sci* 10:615-620

Yamaguchi-Shinozaki K and Shinozaki K. (2006) Transcriptional regulatory networks in cellular responses and tolerance to dehydration and cold stresses. *Annu Rev Plant Biol* 57:781-803

Yang Q, Wang Y, Zhang J, Shi W, Qian C, Peng X. (2007) Identification of

aluminum-responsive proteins in rice roots by a proteomic approach: cysteine synthase as a key player in Al response. *Proteomics* 7:737-749

Yan S, Tang Z, Su W and Sun W. (2005) Proteomic analysis of salt stress-responsive proteins in rice root. *Proteomics* 5: 235-244

Yan SP, Zhang QY, Tang ZC, Su WA and Sun WN. (2006) Comparative proteomic analysis provides new insights into chilling stress responses in rice. *Mol Cell Proteomics* 5: 484-496

Zhang JZ, Creelman RA, and Zhu JK. (2004) From laboratory to field. Using information from *Arabidopsis* to engineer salt, cold, and drought tolerance in crops. *Plant Physiol* 135: 615-621

Zheng Z, Qamar SA, Chen Z and Mengiste T. (2006) *Arabidopsis* WRKY33 transcription factor is required for resistance to necrotrophic fungal pathogens . *Plant J* 48:592-605

Zheng Z, Mosher SL, Fan B, Klessig DF and Chen Z. (2007) Functional analysis of *Arabidopsis* WRKY25 transcription factor in plant defense against *Pseudomonas syringae*. *BMC Plant Biol* 7: 2

Zhu J-K. (2002) Salt and drought stress signal transduction in plants. *Annu Rev Plant Biol* 53, 247-273