

University of Alberta

**PLANKTONIC RESPONSES TO ELEVATED NITROGEN AND PHOSPHORUS
DEPOSITION - A REPLICATED NATURAL ALPINE POND EXPERIMENT**

by

James Paul Zettel

A thesis submitted to the Faculty of Graduate Studies and Research
in partial fulfillment of the requirements for the degree of

Masters of Science
in
Ecology

Biological Sciences

©James Paul Zettel

Fall 2010

Edmonton, Alberta

Permission is hereby granted to the University of Alberta Libraries to reproduce single copies of this thesis and to lend or sell such copies for private, scholarly or scientific research purposes only. Where the thesis is converted to, or otherwise made available in digital form, the University of Alberta will advise potential users of the thesis of these terms.

The author reserves all other publication and other rights in association with the copyright in the thesis and, except as herein before provided, neither the thesis nor any substantial portion thereof may be printed or otherwise reproduced in any material form whatsoever without the author's prior written permission.

Examining Committee

Dr. Rolf Vinebrooke

Dr. David Schindler

Dr. Vincent St. Louis

Dr. Marianne Douglas

Abstract

Several lines of evidence suggest small alpine lakes and ponds are sensitive to nitrogen deposition. Paleolimnological studies, nutrient bioassays, and mesocosm experiments show the positive effects of nitrogen on aquatic alpine primary producers. In particular, alpine pond ecosystems have been inferred to be nitrogen-limited based on low availability of dissolved inorganic nitrogen relative to total phosphorus. However, nitrogen-limitation of alpine ponds has never been tested at the whole-ecosystem level. I performed a replicated *in situ* whole-pond experiment, consisting of two crossed treatments (2 nitrogen x 2 phosphorus levels) applied across 16 natural alpine ponds (n = 4) immediately following ice-out in 2008. Surprisingly, neither nutrient amendment stimulated phytoplankton or zooplankton abundance although subtle shifts in community composition were detected over a two-month period. Intensive grazing pressure exerted by high densities (> 100 individuals/ m²) of herbivorous *Branchinecta paludosa* (fairy shrimp) may have suppressed planktonic responses to nutrient additions. Another ecological explanation for the lack of a positive effect of nutrients on phytoplankton abundance was competition from periphyton, which are comparatively more abundant in most shallow ponds on an areal basis. Therefore, density-dependent ecological interactions (competition and predation) may mediate the responses of phytoplankton to nitrogen deposition over ponds situated in extreme environments.

Table of Contents

Chapter 1 Introduction to alpine and arctic pond ecosystems 1

Introduction 1

Physical Characteristics..... 1

 Distribution 1

 Formation 2

 Temperature..... 3

 Light 4

Chemical composition..... 4

 Dissolved organic carbon 4

 Alkalinity, pH, conductivity and major ions 5

 Nitrogen..... 6

 Phosphorus 7

 Chemical Index of Nutrient Limitation (DIN:TP)..... 7

Biological Communities 9

 Pelagic 9

Primary producers (phytoplankton)	9
Bacteria	10
Zooplankton.....	11
Anostraca (Fairy Shrimp)	13
Benthic Communities.....	14
Periphyton	14
Zoobenthos	15
Study Rational	15
Literature Cited	17
Chapter 2 Planktonic responses to elevated nitrogen and phosphorus deposition-a replicated natural alpine pond experiment.....	26
Introduction.....	26
Materials and Methods	29
Study Site.....	29
Experimental Design	31
Sample Collection	32

Statistical Analysis.....	34
Results.....	35
Water chemistry	35
Phytoplankton.....	36
Zooplankton.....	37
Discussion	38
Literature Cited	43
Chapter 3 Conclusions	60
Synthesis	60
Future Research Directions	61
Literature Cited	65

List of Tables and Figures

Table 1 Average Total Phosphorus (TP) and Total Nitrogen (TN) concentrations and standard error (SE) of alpine and arctic ponds reported by various studies. Hyphenated sections represent data unavailable. 49

Table 2 Physical and chemical parameters of studied alpine ponds in the Cascade Valley, Banff National Park, Canada. 50

Figure 1 Study site location of surveyed alpine waterbodies in Banff National Park, Canada 51

Figure 2 (A) Total Nitrogen (TN) and (B) Total Phosphorus (TP) concentrations in the water column. C (Control), N (Nitrogen), P (Phosphorus), and NP (Nitrogen + Phosphorus) treatment concentrations were measured immediately following nutrient amendment. Error bars represent one standard-error unit based on $n = 3$ 52

Figure 3 (A) Total nitrogen (TN) and (B) total phosphorus (TP) concentrations in the water column 15 days following nutrient amendments: N, P, and NP treatments. Dashed line depicts the average concentration of each nutrient immediately following amendment (see Fig. 1). 53

Figure 4(A) Particulate nitrogen (PN) and (B) particulate phosphorus (PP) prior to each nutrient amendment during the 45-d experiment: C (Control), N (Nitrogen), P (Phosphorus), and NP (Nitrogen + Phosphorus). Error bars represent one standard-error unit based on $n = 3$ 54

Figure 5 Phytoplankton chlorophyll *a* concentration prior to each nutrient amendment during the 45-d experiment: C (Control), N (Nitrogen), P (Phosphorus), and NP (Nitrogen + Phosphorus). 55

Figure 6 Pigment-inferred abundance of (A) Chromophytes (Fucoxanthin) (B) Chlorophytes/Cyanobacteria (Lutein/Zeaxanthin) (C) Chlorophytes and Euglenophytes (Chlorophyll *b*) 56

Figure 7 (A) Ordination biplot of phytoplankton taxa and explanatory environmental variables during the 45-d experiment based on canonical correspondence analysis. Total phosphorus (TP), total nitrogen (TN) and dissolved organic carbon (DOC) are 57

Figure 8 Total zooplankton biomass prior to each nutrient amendment during the 45-d experiment: (C) Control, N (Nitrogen), P (Phosphorus), NP (Nitrogen+Phosphorus). Error bars represent one standard-error unit based on n = 3. 58

Figure 9 Ordination biplot of zooplankton taxa with species codes and explanatory environmental variables during the 45-d experiment. 59

Acknowledgements

Most importantly I would like to thank my supervisor Dr. Rolf Vinebrooke who has played a vital role in every component of my thesis. His helpful insight was much appreciated during the early stages of experimental design, planning, and assisting in field work during the “thunderflurries” and challenging mountain weather. Rolf always provided sound advice, and was full of positive encouragement even in the later stages of data analysis and report writing. I would also like to thank my committee members Dr. David Schindler, Dr. Vincent St Louis and Dr. Marianne Douglas for their constructive input and valuable comments.

Field support was provided by Lisa Porter, Conrad Murphy, Diana Tirlea, and Mirea Bartrons. In particular, I greatly appreciated Megan MacLennan’s effort in the field and in the lab preparing experiments in addition to our sanity saving Swiss Chalet escapes. Mark Graham and Devin Lyons were always willing to lend their time to solve any problems in the lab and provided great advice while writing.

I would like to thank my parents and siblings for their encouragement from the beginning of my academic career. Many thanks to friends who have helped me survive the dark winters of Edmonton by enjoying the outdoors to hike, hunt, fish, and climb that include: Christina Macdonald, Ben Dorsey, Simon Ham, Vij Patil, Rene Kohut, Lindsey Wilson, Fred Noddin, Kieren Odonovan, Jamie Hockenhill, Andrea Kortello, Brett Scheffers, Hannah Mckenzie, Dustin Raab, Joe Northrup, Jesse Tigner, Mark Wong, Jolene Swain, Jordan Messner, Heidi Ando, and Aaron Schafer. I would also like to

thank Sarah Trefry, Kim Dawe, Linnea Mowat, Isla Myers Smith, Jamie Gorrell and Michelle Bacon for their friendship and positive encouragement. A special thanks to Oliver for the good times fishing and hunting the Berland that has provided many delicious meals of elk.

This work was supported by a Alexander Graham Bell Canada Graduate Scholarship from the Natural Sciences and Engineering Research Council (NSERC), a Queen Elisabeth II Scholarship, and Graduate Teaching Assistant Scholarships awarded by the University of Alberta. Funding support was also provided by Alberta Conservation Authority, Alberta Water Research Institute and NSERC Discovery grants awarded to Rolf Vinebrooke.

Chapter 1 Introduction to alpine and arctic pond ecosystems

Introduction

A pond can be defined as a shallow waterbody of permanent or seasonal nature between 1 m² and 2 ha in area that may be of man-made or natural form (Biggs et al. 2005). A deep waterbody small in area could also be considered a pond. Alpine and arctic pond systems show many similar physical, chemical, and biological characteristics (Orbaek et al. 2007; Vincent and Laybourn-Parry 2008). These extreme environments are often remote and poorly characterized (Orbaek et al. 2007). Cloud-cover, wind turbulence and extreme daily fluctuations in temperature are common at high elevation sites (Linacre 1982). Polar regions receive pronounced seasonal oscillations in solar radiation and the light regime changes dramatically due to their position on the globe (Orbaek et al. 2007).

Physical Characteristics

Distribution

Global surveys have shown that inland waters are predominately composed of small (<10 ha surface area), shallow (<10 m mean depth) lakes (Downing et al. 2006). Ponds located in alpine and polar regions are more abundant than lakes that receive greater scientific attention (Orbaek et al. 2007; Vincent and Laybourn-Parry 2008). In comparison to lakes, ponds are considered biological “hotspots” that often contain greater microbial and invertebrate biodiversity (Williams et al., 2004; Biggs et al., 2005; Smol and Douglas 2007). Overall, the

globe is covered with a much greater proportion of small waterbodies and shallow lakes compared to large deep lakes, which have received greater scientific attention (Oertli et al. 2005; Downing et al. 2006; Moss 2010). For example, over 3000 small waterbodies can be found in Banff National Park relative to 480 lakes over the same region (McMaster and Schindler 2005). Alpine ponds are potentially sensitive to environmental inputs and changes in part due to factors such as high water renewal rates, thin soils often with poor buffering capacity, and small catchments (Hinden et al. 2005). Despite their high abundance and ecological importance, research on pond ecosystems is limited, especially in remote alpine and polar environments.

Formation

Thermokarst or thrust ponds are common in the alpine and polar regions across much of the northern hemisphere, including Alaska, Canada, and Siberia (Cote and Burn 2002; Hinkel et al. 2005; Walter et al. 2006). These small waterbodies are formed by physical depressions from melting ice and permafrost. They are typically found in alpine meadows lacking inflows/outflows, surrounded by steep banks, less than 1 m in depth and characterized by fine flocculent organic silt sediments (Vincent and Laybourn-Parry 2008).

Ponds of alpine and polar regions share many similar physical features (Neldner and Pennak 1951; Hobbie 1980; Douglas and Smol 1994). Water budgets of many ponds depend mainly on precipitation, evaporation and seepage through the sediment (Neldner and Pennak 1951). Surface drainage supplies much of the water in the early growing season from snowmelt in the surrounding catchment, while the ground is usually frozen (McMaster 2003). By July, showers and

seepage are the main sources of water. During warm-dry weather periods, pond volumes decrease most rapidly, but may rebound after major precipitation events (McMaster 2003). Therefore, many ponds may disappear because of climate change effects and decreases in precipitation/evaporation ratios (Douglas and Smol, 1994; Schindler and Smol, 2006). In early fall, alpine ponds freeze to the bottom and thaw in late spring from the top downwards (Daborn and Clifford 1974; McMaster 2003). Pools of water form overlying thick masses of ice bound to sand and gravel particles. Ponds are often subject to a short hydroperiod of less than 2 months (see Chapter 2) and short growing seasons with less than 100 ice-free days (Neldner and Pennak 1955; Schmitz 1959).

Temperature

Alpine ponds have highly variable thermal environments because of a limited capacity for heat storage during the ice-free period, often closely tracking daily changes in air temperatures ranging from below freezing to $>20^{\circ}\text{C}$ (Strecker et al. 2004; McMaster 2003). Most polar aquatic habitats experience temperatures close to 0°C most of the year, however, some shallow waterbodies high in light absorbing dissolved inorganic carbon (DOC) may reach temperatures above 15°C (Vincent and Laybourn-Parry 2008). The water column is often well-mixed, driven by high wind velocities. During the winter when air temperatures remain below zero for several months, the water surface of ponds freeze to the sediments in high elevation (Neldner and Pennak 1955) and high latitude regions (Daborn and Clifford 1974). Pond particulate matter and dissolved solids become increasingly more concentrated in the water column as overlying ice thickens (Daborn and Clifford 1974). The remaining solutes are forced into the sediment

when the pond freezes completely (Prentki 1980). Aquatic organisms must have the ability to migrate to a more suitable habitat, or mechanisms to avoid freezing (Danks 2004).

Light

Alpine and arctic ponds are exposed to relatively high levels of ultraviolet radiation (UV: 290 – 400 nm wavelength band) following ice-out during early summer (Laurion et al. 2000). For example, under clear sky conditions, solar UV- B (290-320 nm) radiation increased by almost 20% while UVA increased by approximately 10% per 1000 m gain in elevation (Blumthaler et al. 1997). Native populations of amphibians (Palen et al. 2000) heterotrophic protists, bacteria, and viruses are particularly sensitive to UVR (Sommaraga 2001) . In comparison, several primary producers and zooplankton appear tolerant of high solar irradiance (Sommaraga 2001), exhibiting high concentrations of photoprotective pigments (Quesada et al. 1999; Vincent and Roy 1993). Photodegradation changes the composition and structure of dissolved organic carbon producing less aromatic products (Bertilsson and Tranvik 2000), liberating phosphorus (Franko and Heath 1982; Cotner and Heath 1990) and nitrogen (Bushaw-Newton and Moran 1999; Kieber 2000).

Chemical composition

Dissolved organic carbon

DOC often regulates the underwater light environment of alpine and polar ponds by attenuating shorter wavelengths of the visible spectrum (400 – 700 nm) and ultraviolet (300 – 400 nm) band (Vincent and Roy 1993; Laurion et

al. 2000). In alpine and polar regions, ponds surrounded by terrestrial vegetation in their catchment generally appear more brown than lakes owing to high levels of DOC (Graneli 2004; Vinebrooke and Leavitt 1999). DOC concentrations in these regions can range from less than 1 mg C/L to higher than 20 mg C/L, owing to potentially pronounced terrestrial (allochthonous) inputs of humic substances and internal (autochthonous) production of organic exudates by benthic biofilms (McKnight et al. 1994; McMaster 2003; Lim et al 2005).

Alkalinity, pH, conductivity and major ions

The chemistry of alpine and arctic ponds may vary depending on the water budget characteristics, surrounding geology, and atmospheric inputs. Average conductivity measurements have ranged from as low as $9.7 \mu\text{S cm}^{-1}$ in alpine ponds of Switzerland (Oertli et al. 2008; Hinden et al. 2005) to $61 \mu\text{S cm}^{-1}$ in the Canadian Rocky Mountains (McMaster and Schindler 2005). Ponds are often weakly buffered (mean alkalinity 0.55 meq L^{-1}) and close to neutral in pH (7.2; McMaster and Schindler 2005; Hinden et al. 2005). The most common ions in solution consist of bicarbonate, magnesium and calcium (McMaster 2003). Seepage losses may be more important than evaporation in some alpine ponds, shown by relatively stable concentrations in alkalinity, pH, conductivity, bicarbonate, magnesium, and calcium (McMaster 2003). In contrast, arctic ponds have shown higher pH, ion concentrations, mean alkalinity, and conductivity presumably due differences in geology (Prentki et al. 1980; Douglas and Smol 1994; Lim et al. 2005).

Dissolved reactive silica concentrations are frequently very low in arctic (Prentki et al. 1980) and alpine ponds (McMaster 2003). Primary producers such as diatoms and other siliceous algae are often limited by this element at low concentrations ($<0.5 \text{ mg L}^{-1}$; Conley et al. 1993). Phycological analyses of alpine ponds in the Canadian Rocky Mountains showed diatoms were scarce in the alpine pond phytoplankton (McMaster 2003; Zettel and Vinebrooke, unpublished data). Epipellic diatoms were more abundant than pelagic diatoms but were low in concentration owing to their sensitivity to desiccation (Mosisch 2001), grazing (McMaster 2003) and low concentrations of silica (Douglas and Smol 1993; Lim et al. 2005).

Nitrogen

Ponds of alpine and polar regions contain total nitrogen (TN) concentrations that vary regionally (Table 1; Lim et al. 2005; Oertli et al. 2008; Murphy et al 2010; Zettel and Vinebrooke, unpublished data). TN consists mainly of total dissolved nitrogen (TDN). Particulate nitrogen (PN) accounts for only ~10% of the TN pool (Lim et al. 2005; Murphy et al 2010; Zettel and Vinebrooke, unpublished data). In certain small mountain lakes, TDN is primarily composed of dissolved organic nitrogen (DON). Low concentrations of bioavailable dissolved inorganic nitrogen (DIN), which is comprised of ammonia, nitrites, and nitrates (Bunting et al. 2010), suggests the potential for nitrogen-limitation in these ecosystems.

Phosphorus

Average total phosphorus (TP) concentrations reported in ponds of the Canadian Arctic and Rocky Mountains are reported in Table 1 and many ponds could be classified as meso- to eutrophic ecosystems (Lim et al. 2005; Murphy et al 2010; McMaster and Schindler 2005; Zettel and Vinebrooke, unpublished). Here, TP can contain approximately equal fractions of total dissolved phosphorus (TDP) and particulate organic phosphorus (PP), while soluble reactive phosphorus (SRP) is low, averaging approximately 1-2 $\mu\text{g/L}$ (Lim et al. 2005; McMaster and Schindler 2005; Van Geest 2007; Murphy et al. 2010). Organically bound phosphorus may however be made bioavailable as orthophosphate through the hydrolytic activity of microbial enzymes, such as phosphatases (Chrost and Overbeck 1987).

Chemical Index of Nutrient Limitation (DIN:TP)

Nutrient limitation status is often inferred using a dissolved inorganic nitrogen (DIN) to total phosphorous (TP) ratio (Morris & Lewis 1988). The DIN:TP index is favored over others because it most accurately represents N and P fractions truly available to phytoplankton communities from cellular reserves and surrounding water column (Morris and Lewis 1988). Short-term bioassays of mountain lakes in Colorado showed that nitrogen either alone or in combination with P limited phytoplankton in almost 80% of all cases (Morris and Lewis 1988). Similarly, positive bioassay responses to N amendment were matched by low DIN:TP values (Murphy et al. 2010). Historically, many lakes in the Sierra Nevada showed phosphorus (P)

limitation but as a result of increased P loading, N limitation is progressively more common (Sickman et al. 2003).

Phytoplankton may readily assimilate DIN while DON composed of high molecular weight compounds (peptides, humic acids, fulvic acids) is too large for cellular uptake (Morris and Lewis 1988). Although low molecular weight organic compounds such as urea and amino acids are also accessible to phytoplankton, they are usually found in low concentrations in inland waters (Bonin and Maestrini 1981; Styrett 1981). Therefore, DIN most closely approximates the amount of nitrogen available to the algal community.

TP is also a good predictor of nutrient limitation because numerous forms (SRP, TDP, PP) are available both extra- and intracellularly to primary producers. For example, SRP may be directly assimilated whereas dissolved organic phosphorus may be chemically broken down by extracellular phosphatase excreted by bacteria and algae (Chrost and Overbeck 1987). Algae also store excess P as particulate phosphorus (PP) and therefore it is an important component to measure when investigating nutrient status (Morris and Lewis 1988).

Empirical evidence of nutrient-limitation in alpine and polar pond ecosystems varies by region, type of primary producer (periphyton/phytoplankton), and scale of the experiment. A field-based study (McMaster and Schindler 2005) examining the correlation between environmental parameters and sediment-dwelling algae (epipelton) abundance found that nitrate and nitrite were the only significant predictor of epipelton abundance. However, phosphorus was a significant predictor of phytoplankton abundance. Benthic biofilms may

rapidly assimilate DIN in the water column and outcompete phytoplankton (Axler and Reuter 1996; Mcknight et al. 2004; Nydick et al. 2004). Laboratory bioassays suggested alpine ponds were most often co-limited (Vinebrooke, unpublished) and very few showed nitrogen (N) limitation (Murphy et al. 2010). Overall, increased N deposition in the Rocky Mountains was not suspected to result in widespread eutrophication of alpine waterbodies because the surrounding lakes of the region are primarily phosphorus-limited (Gardner et al. 2008; Murphy et al. 2010). In contrast, a 24 hour bioassay experiment conducted by Levine and Whalen (2001) proposed that N more commonly stimulated phytoplankton production in arctic ponds rather than phosphorus. However, bottle and mesocosm nutrient bioassays may not be ecologically realistic therefore there is a need for experiments at the whole ecosystem scale (Carpenter 1996; Schindler 1998).

Biological Communities

Pelagic

Primary producers (phytoplankton)

Ponds in arctic and alpine regions contain chlorophyll concentrations ranging from 1-27 $\mu\text{g/L}$ (Lim et al. 2005; McMaster and Schindler 2005; Murphy et al. 2010). Seasonal fluctuations in primary producer biomass in these systems have been attributed to variation in resources and predation (i.e., grazing pressure; Van Geest et al. 2007). Phytoplankton primary production was measured in arctic ponds between 2.1 – 3.8 $\mu\text{mol C/L/day}$ (Whalen et al. 2008). Low temperatures and supply of nutrients in the alpine limit autotrophic production (Ward 1985; Hinden et al. 2005).

Alpine and arctic ponds support a wide range of phytoplankton species that belong to the Bacillariophyta (diatoms), Chrysophyta (chrysophytes), Cryptophyta (cryptophytes), Chlorophyta (green algae), and Cyanophyta (cyanobacteria or “blue-green algae”) (Vincent and Laybourn-Parry 2008). Common genera in alpine ponds include *Gloeocapsa*, *Oscillatoria*, *Scenedesmus*, *Crucigenia*, *Synedra*, *Anabaena*, and *Tabellaria* (Strecker et al. 2004; Zettel and Vinebrooke, unpublished data). Phytoplankton from the genera such as *Chlamydomonas* and *Ochromonas* are found in many polar and alpine ponds (Laybourn et al. 1991). Shifts in alpine phytoplankton community composition have been documented in response to various environmental stressors, including environmental warming (Strecker et al. 2004; Hozapfel and Vinebrooke 2005; Flanagan et al. 2009; Parker et al. 2008), and nitrogen deposition (Baron et al. 2000; Lafrancois et al. 2003; Nydick et al. 2004; Thompson et al. 2008).

Bacteria

Bacterial production and biomass are more important than algae as food sources to higher trophic levels in some arctic ecosystems. For example, Rautio and Vincent (2006) showed that lakes and ponds had high zooplankton concentrations (up to 17 000 crustaceans per m³) despite clear waters low in phytoplankton and primary productivity. The planktonic resource pool was primarily composed of detritus (89–98%) instead of and bacteria biomass (up to 29 mg / m³) - the most important food item to zooplankton in this food web (Rautio and Vincent 2006). Graneli (2004) found that bacterial production and abundance were over 10 times higher in

the ponds than in lakes despite equally low P levels across both types of waterbodies. Bacterioplanktonic growth is well correlated with temperature, especially at low temperatures (<10°C; Scavia and Laird 1987; Morris and Lewis 1992; Coveney and Wetzel 1995). Therefore, in northern shallow lakes and ponds the classical foodweb where phytoplankton provides the sole food base for higher trophic levels does not occur. Instead, microbial mats and bacterioplankton are the primary food source to zooplankton.

There are many similarities in the bacterioplankton communities of polar regions. Pearce et al (2007) showed that freshwater ecosystems of the Arctic and Antarctic share many of the same bacterioplankton by employing RNA-sequencing techniques. Differences in bacterial community composition were evident between some lakes (Lindstrom & Leskinen 2002). Many bacterial species and protists found in polar regions resembled those found in temperate waters (Vincent and Laybourn-Parry 2008).

Zooplankton

Many alpine and arctic ponds have food webs that differ from nearby lakes because to they lack fish and have an abundance of basal resources. Dense populations of pigmented zooplankton (>17 000 crustaceans / m³) including cladocerans, anostracans (fairy shrimp) and copepods inhabit ponds but are less abundant in lakes of these regions (Bertilsson et al. 2003; Rautio and Vincent 2006). Large zooplankton are often present in these ponds because they lack predation pressure by fish (Parker et al. 2001).

Food is the most important environmental factor explaining zooplankton size distribution in polar waterbodies (Vincent and Laybourn-Parry 2008). Benthic microbial mats sustain large zooplankton in many highly productive ponds (Hansson and Tranvik 2003; Rautio and Vincent 2006). Therefore, grazing pressure exerted by zooplankton in ponds may be much stronger than in lakes (Granelli 2004). Zooplankton species richness typically decreases with elevation (Rautio 1998; Hessen et al. 2006).

Cladocerans such as *Daphnia pulex* are common to many alpine ponds and widely distributed (Schmitz 1959). During favourable environmental conditions cladocerans reproduce through parthenogenesis. Changes in food availability, water quality, photoperiod, population density can produce ephippia well in advance of the early winter freeze, which usually occurs in middle to late October on the alpine tundra (Schmitz 1959).

Several species of predatory copepods and anostracans that reproduce sexually may be distributed across many arctic and alpine ponds (Dodson 1974).

Planktonic rotifers such as *Keratella*, *Notholca*, and *Filinia* are generally uncommon in polar and alpine ponds, owing possibly to strong competitive interference by larger cladocerans and predation by copepods (Strecker et al. 2004; Vincent and Laybourn-Parry 2008).

Anostraca (Fairy Shrimp)

Fairy shrimp are very abundant in many ponds and may even be regarded as a dominant life form (i.e. keystone species) linking pelagic and benthic habitats. In an Antarctic study, more than 40000 juvenile *Branchinecta gaini* were reported in a cubic meter, feeding on 8 μg of chlorophyll *a* per hour per individual (Hawes 1985). Fairy shrimp are ubiquitous and one in particular, *Branchinecta paludosa* covers many regions of the globe, inhabiting alpine waterbodies and latitudes above 60°N in Europe, Russia and North America (Saunders et al. 1993). They frequently inhabit fishless ponds that have a short hydroperiod, and freeze to the bottom during winter months. On occasion, fairy shrimp do not hatch in a particular pond each year, possibly due to extirpation and recolonization, or specific physiological requirements that are not always met (Neldner and Pennak 1955, Donald 1983).

Although limited research has been conducted on their feeding habits and life history, most fairy shrimp are described as non-selective filter feeders that collect algal and detrital material by particle filtration using thoracic appendages (Brendonck 1993). They target a variety of food items that include phytoplankton (Mitchell 1991), benthic microbial mats (Rautio and Vincent 2006), rotifers and cladocerans (Sarma 2002; Hawes 2008). Fairy shrimp are described as generalist feeders that are also capable of feeding directly on dead conspecifics (Hawes 2008).

Benthic Communities

Periphyton

Benthic primary production (BPP; 8.4 g C / m²/year) was seven times greater than that of phytoplankton in some polar ponds, suggesting its importance to the food webs of shallow systems (Hobbie 1980). BPP (53 g C / m²/year) comprised more than 66% of the total primary production in shallow systems (Whalen et al. 2008) while abundances may range from 18 – 90 mg chlorophyll *a*/m² (Vincent and Laybourn-Parry 2008). Therefore, dense microbial mats may be more ecologically significant than the overlying dilute concentrations of phytoplankton.

The periphyton community of ponds located in the alpine and polar regions are often dominated by cyanobacterial mats (McMaster 2003, Vincent and Castenholz 1993; Quesada et al. 1999), which may be several centimeters thick and completely cover the bottom substrate. Genera that are common to the periphytic community of ephemeral ponds include: *Anabaena*, *Oscillatoria*, *Phormidium*, *Lyngbya*, *Microcoleus*, *Nodularia*, *Nostoc*, *Schizothrix*, *Synechococcus*, *Scytonema*, *Calothrix*, *Tolypothrix*, and *Scytonema* (Vincent and Castenholz 1993) while diatoms and chlorophytes are also present in small proportions (Quesada et al. 1999). Mats are bound by a cohesive mucilaginous layer, often overlying silt and sand particles. Tolerant of extreme environmental conditions (i.e., pH, temperature, salinity), these microbial mats are most abundant where macroinvertebrate grazing is limited (Palmisano et al. 1989). Benthic cyanobacterial mats quickly assimilate SRP and DIN, ultimately affecting the nutrient limitation status of smaller waterbodies (McKnight et al. 2004).

Zoobenthos

Benthic invertebrates of polar and alpine ponds often form the highest trophic level in the aquatic food web. Their diversity typically declines with elevation (Oertli et al. 2008) and latitude (Gibson and Bayly 2007) and is comprised of taxa belonging to Chironimidae, Coleoptera, Oligochaeta, Tricoptera, Heteroptera, Gastropoda, Odonata, and Ephemeroptera (Oertli et al. 2008). Top-down control on macroinvertebrates is usually weak as the majority of ponds are fishless therefore, invertebrate density is dependent on cold tolerance and food supply (Vincent and Laybourn-Parry 2008). Oertli et al. (2008) documented a weak connection of macroinvertebrate assemblage (composition, richness) to environmental parameters. Instead, regional processes were more important such as physical connection to other ponds and tributaries were better predictors of invertebrate communities.

Study Rational

Dramatic increases in wet deposition of nitrogen (N) over North America and Europe due to anthropogenic emissions are well known (Galloway and Cowling 2002; Fenn et al 2003). Decade-long impacts of increased N deposition on mountain lakes have been inferred paleolimnologically based on fossilized diatom assemblages (Wolfe et al. 2001; Saros et al. 2003). Experimental evidence has also shown the positive effects of nitrogen on alpine waters (Nydick et al 2004; Thompson et al 2008; Murphy et al. 2010). In particular, alpine ponds are more likely N limited compared to lakes based on lower DIN:TP ratios (Strecker et al. 2004; McMaster and Schindler 2005; Murphy et al. 2010). Increasingly, freshwater ecosystems are reported as

being co-limited by N and P (Vinebrooke, unpublished; Elser et al 1990, 2007; Ogbebo et al 2009). Consequently, the distinct effect of either nutrient N or P on primary producers is amplified by addition of the other, resulting in a synergistic interaction that denotes co-limitation.

I designed a replicated whole-pond experiment to determine the effects of N and P on phytoplankton and zooplankton communities using 16 alpine ponds located along the Eastern Front ranges of the Canadian Rocky Mountains. I proposed that the degree of the positive ecological effects of N would depend on P availability (i.e. nutrient co-limitation driven by phosphorus). The rationale for this hypothesis came from findings from earlier in vitro nutrient bioassays, which indicated that many of these ponds were co-limited by N and P (Vinebrooke, University of Alberta, unpublished data). Also, I expected the nutrient additions to stimulate the proliferation of large inedible phytoplankton, suppressing higher trophic levels by impairing energy transfer to herbivores (Beisner et al. 1997).

Literature Cited

Axler, R.P. and Reuter, J.E. (1996) Nitrate uptake by phytoplankton and periphyton: Whole-lake enrichments and mesocosm-¹⁵N experiments in an oligotrophic lake. *Limnology and Oceanography*. 41: 659-671.

Baron, J.S., Rueth, H.M., Wolfe, A.M., et al. (2000) Ecosystem responses to nitrogen deposition in the Colorado Front Range. *Ecosystems*. 3:352-368.

Beisner, B.E., McCauley, E., and Wrona, F.J. (1997) The influence of temperature and food chain length on plankton predator-prey dynamics. *Canadian Journal of Fisheries and Aquatic Sciences*. 54:586-595.

Bertilsson, S. and Tranvik, L.J. (2000) Photochemical transformation of dissolved organic matter in lakes. *Limnology and Oceanography*. 45:753-62.

Biggs, J., Williams, P., Whitfield, et al. (2005) 15 years of pond assessment in Britain: results and lessons learned from the work of pond conservation. *Aquatic Conservation: Marine and Freshwater Ecosystems*. 15: 693-714.

Blumthaler, M., Ambach, W. and Ellinger, R. (1997) Increase in solar UV radiation with altitude. *Journal of Photochemistry and Photobiology*. 39: 130-134.

Bonin, D.J. and Maestrini, S.Y. (1981) Importance of organic nutrients for phytoplankton growth: Implications for algal species succession. *Canadian Bulletin of Fisheries and Aquatic Science*. 210: 279-291.

Brendonck, L. (1993) Feeding in the fairy shrimp *Streptocephalus proboscideus* (Frauenfeld) (Branchiopoda: Anostraca). Aspects of the feeding biology. *Journal of Crustacean Biology*. 13: 235-244.

Bunting, L., Levitt, P.R., Weidman, P., et al. 2010. Regulation of the nitrogen biogeochemistry of mountain lakes by subsidies of terrestrial dissolved organic matter and the implications for climate studies. *Limnology and Oceanography*. 55: 333-345.

Bushaw-Newton, K.L. and Moran, M.A. (1999) Photochemical formation of biologically available nitrogen from dissolved humic substances in coastal marine systems. *Aquatic Microbial Ecology*. 18: 285-92.

Carpenter, S.R. (1996) Microcosm experiments have limited relevance for community and ecosystem ecology. *Ecology*. 77: 677-680.

Chrost, R.J. and Overbeck, J. (1987) Kinetics of alkaline phosphatase activity and phosphorus availability for phytoplankton and bacterioplankton in Lake Plusee (north German eutrophic lake). *Microbial Ecology*: 13: 229–248.

Conley, D.J., Schelske, C.L. and Stoermer, E.F. (1993) Modification of the biogeochemical cycle of silica with eutrophication. *Marine ecology progress series*. 101: 179-192

Coveney, M.F. and Wetzel, R.G. (1995) Biomass, production, and specific growth rate of bacterioplankton and coupling to phytoplankton in an oligotrophic lake. *Limnology and Oceanography*. 40: 1187-1200.

Cote, M.M. and Burn, C.R. (2002) The oriented lakes of Tuktoyaktuk Peninsula, Western Arctic Coast, Canada: a GIS based analysis. *Permafrost and Periglacial Processes*. 13: 61-70.

Cotner, J.B. and Heath, R.T. (1990) Iron redox effects on photosensitive phosphorus release from dissolved humic materials. *Limnology and Oceanography*. 35:1175–81.

Daborn, G.R. and Clifford H.F. (1974) Physical and chemical features of an aestival pond in Western Canada. *Hydrobiologia*. 44: 43-59.

Danks, H.V. (2004) Seasonal adaptations in arctic insects. *Integrative and Comparative Biology*. 44: 85–94.

Dodson, S.I. (1974) Zooplankton competition and predation: an experimental test of the size-efficiency hypothesis. *Ecology*. 55: 605-613.

Donald, D.B. (1983) Erratic occurrence of anostracans in a temporary pond: colonization and extinction or adaptation to variations in annual weather? *Canadian Journal of Zoology*. 61: 1492–1498.

Douglas, M.V and Smol J.P. (1993) Freshwater diatoms from high arctic ponds (Cape Herschel, Ellesmere-island, NWT) *Nova Hedwigia*. 57:511-552.

Douglas, M.V. and Smol, J.P. (1994) Limnology of high arctic ponds (Cape Herschel, Ellesmere Island, NWT). *Archiv Fur Hydrobiologie*. 131: 401-434.

Downing, J.A., Prairie, Y.T., Cole, J.J., et al. (2006) The global abundance and size distribution of lakes, ponds, and impoundments. *Limnology & Oceanography*. 51: 2388–2397.

Downing, J.A., Cole, J.J., Middelburg, J.J., et al. (2008) Sediment organic carbon burial in agriculturally eutrophic impoundments over the last century. *Global Biogeochemical Cycles*. 22: 1-10.

Elser, J.J., Marzolf, E.R., and Goldman, C.R. (1990) Phosphorus and nitrogen limitation of phytoplankton growth in the freshwaters of North America: a review and critique of experimental enrichments. *Canadian Journal of Fisheries and Aquatic Science*. 47: 1468–1477.

Elser, J.J., Bracken, M.E., Cleland, E.E., et al. (2007) Global analysis of nitrogen and phosphorus limitation of primary producers in freshwater, marine and terrestrial ecosystems. *Ecology Letters*. 10: 1135–1142.

Fenn, M., Haeuber, R., Tonnesen, G., et al. (2003) Nitrogen emissions, deposition, and monitoring in the western United States. *Bioscience*. 53: 391-403.

Flanagan, C.M., McKnight, D.M., Liptzin, D., et al. (2009) Response of the phytoplankton community in an alpine lake to drought conditions: Colorado Rocky Mountain Front Range, U.S.A. *Arctic, Antarctic and Alpine Research*. 41:191-203

Franko, D.A. and Heath, R.T. (1982) UV-sensitive complex phosphorus: association with dissolved humic material and iron in a bog lake. *Limnology and Oceanography*. 27:564-569.

Galloway, J.N. and Cowling E.B. (2002) Reactive nitrogen and the world: 200 years of change. *Ambio*. 31:64-71.

Gardner, E.M., Mcknight, D.M., Lewis, W.M., et al. 2008. Effects of nutrient enrichment on phytoplankton in an alpine lake, Colorado, U.S.A. *Arctic, Antarctic, and Alpine Research*. 40: 55–64.

Gibson, J.A. and Bayly, I.A. (2007) New insights into the origins of crustaceans of Antarctic lakes. *Antarctic Science*. 19:157-164

Graneli, W., Bertilsson, S. and Philibert, A. (2004) Phosphorus limitation of bacterial growth in high Arctic lakes and ponds. *Aquatic Sciences*. 66: 430-439.

Hansson, L.A. and Tranvik, L.J. (2003) Food webs in sub-Antarctic lakes: a stable isotope approach. *Polar Biology*. 26: 783–788.

- Hawes, I. (1985) Factors controlling phytoplankton populations in maritime Antarctic lakes. Antarctic Nutrient Cycles and Food Webs 4th scientific committee on Antarctic research symposium on Antarctic biology
- Hawes, T.C. (2008) Feeding behaviour in the Antarctic fairy shrimp, *Branchinecta gaini*. *Polar Biology*. 31:1287-1289.
- Hessen, D.O., Faafeng, B.A., Smith, V.H. et al. (2006) Extrinsic and intrinsic controls of zooplankton diversity in lakes. *Ecology*. 87:433–443.
- Hinden, H., Oertli, B., Menetrey, N., et al. (2005) Alpine pond biodiversity: what are the related environmental variables? *Aquatic Conservation: Marine and Freshwater Ecosystems*. 15: 613-624.
- Hinkel, K.M., Frohn, R.C., Nelson, F.E., et al. (2005) Morphometric and spatial analysis of thaw lakes and drained thaw lake basins in the Western Arctic Coastal Plain, Alaska. *Permafrost and Periglacial Processes*. 16:327-341.
- Hobbie, J. E., 1980: *Limnology of Tundra Ponds: Barrow Alaska*. Stroudsburg, Pennsylvania: Dowden, Hutchinson and Ross, Inc.
- Holzappel, A. M. and Vinebrooke, R. D. (2005) Environmental warming increases invasion potential of alpine lake communities by imported species. *Global Change Biology*. 11: 2009–2015.
- Kieber, D.J. (2000) Photochemical production of biological substrates. In: de Mora, S., Demers, S. and Vernet, M. (eds) *The effects of UV radiation in the marine environment*. Cambridge, England, Cambridge University Press. p 130–48.
- Lafrancois, B.M., Nydick, K.R. and Caruso, B. (2003) Influence of nitrogen on phytoplankton biomass and community composition in fifteen snowy range lakes (Wyoming, U.S.A.) *Arctic, Antarctic, and Alpine Research*. 35:499–508.
- Laurion, I., Ventura, M., Catalan, J., et al. (2000) Attenuation of ultraviolet radiation in mountain lakes: Factors controlling the among- and within-lake variability. *Limnology and Oceanography*. 45:1274-1288.
- Laybourn-Parry, J., Marchant, H.J. and Brown, P. (1991) The plankton of a large oligotrophic freshwater Antarctic lake. *Journal of Plankton Research*. 13:1137-1149.

Levine, M.A. and Whalen, S.C. (2001) Nutrient limitation of phytoplankton production in Alaskan Arctic foothill lakes. *Hydrobiologia*. 455: 189–201.

Lim, D.S., Douglas, M.S. and Smol, J.P. (2005) Limnology of 46 lakes and ponds on Banks Island, N.W.T., Canadian Arctic Archipelago. *Hydrobiologia*. 545: 11–32.

Linacre, E. (1982) The effect of altitude on the daily range of temperature. *Journal of Climatology*. 2: 375-382.

Lindstrom, E.S. and Leskinen, E. (2002) Do neighboring lakes share common taxa of bacterioplankton? Comparison of 16S rDNA fingerprints and sequences from three geographic regions. *Microbial Ecology*. 44: 1-9.

McKnight, D.M., Andrews, E.D., Spalding, S.A. et al. (1994) Aquatic Fulvic Acids in Algal-Rich Antarctic Ponds. *Limnology and Oceanography*. 39:1972-1979.

McKnight, D.M., Runkel, R.L., Tate C.M. et al. (2004) Inorganic nitrogen and phosphorus dynamics of Antarctic glacial meltwater streams as controlled by hyporheic exchange and benthic autotrophic communities. *Journal of North American Benthological Society*. 23: 171–188.

McMaster, N.L. (2003) Alpine ponds: Climate change and N deposition. MSc thesis, University of Alberta, Edmonton, Alberta.

McMaster, N.L., and Schindler, D.W. (2005) Planktonic and epipelagic algal communities and their relationship to physical and chemical variables in alpine ponds in Banff National Park, Canada. *Arctic, Antarctic, and Alpine Research*. 37: 337–347.

Mitchell, S. A. (1991) The growth rate and growth efficiency of *Streptocephalus macrourus* (Crustacea, Anostraca) cultured on microalgae. *Hydrobiologia*. 212: 1–10.

Morris, D. P. and Lewis, W. M. (1988) Phytoplankton nutrient limitation in Colorado mountain lakes. *Freshwater Biology*. 20: 315–327.

Morris, D.P. and Lewis, W.M. (1992) Nutrient limitation of bacterioplankton growth in Lake Dillon, Colorado. *Limnology and Oceanography*. 37: 1179-1192.

Mosisch, T.D. (2001) Effects of desiccation on stream epilithic algae. *New Zealand Journal of Marine and Freshwater Research*. 35: 173-179.
Moss, B. (2010) Climate change, nutrient pollution and the bargain of Dr Faustus. *Freshwater Biology*. 55: 175-187.

Murphy, C.A., Thompson, P.L., and Vinebrooke, R.D., 2010. Assessing the sensitivity of alpine lakes and ponds to nitrogen deposition in the Canadian Rocky Mountains. *Hydrobiologia*. 648: 83-90.

Neldner, K.H. and Pennak, R.W. (1955) Seasonal faunal variations in a Colorado alpine pond. *American Midland Naturalist*. 53: 419-430.

Nydick, K.R., Lafrancois B.M. and Baron, J.S. (2004) NO₃ uptake in shallow, oligotrophic, mountain lakes: the influence of elevated NO₃ concentrations. *Journal of the North American Benthological Society*. 23: 397-415.

Nydick, K.R., LaFrancois, B.M., Baron, J.S., et al. (2004) Nitrogen regulation of algal biomass, productivity, and composition in shallow mountain lakes, Snowy Range, Wyoming, USA. *Canadian Journal of Fisheries and Aquatic Sciences*. 61: 1256-1268.

Ogbebo, F.E.; Evans, M.S.; Waiser, M.J., et al. (2009) Nutrient limitation of phytoplankton growth in Arctic lakes of the lower Mackenzie River Basin, northern Canada. *Canadian Journal of Fisheries and Aquatic Science*. 66:247-260.

Oertli, B., Biggs, J., Cereghino, R., et al. (2005) Conservation and monitoring of pond biodiversity: introduction. *Aquatic Conservation: Marine and Freshwater Ecosystems*. 15: 535-540.

Oertli, B., Indermuehle, N., Angelibert, S., et al. (2008) Macroinvertebrate assemblages in 25 high alpine ponds of the Swiss National Park (Cirque of Macun) and relation to environmental variables. *Hydrobiologia*. 597:29-41.

Orbaek, R., Kallenborn, Tombre, I. et al. (2007) Arctic alpine ecosystems and people in a changing environment. Springer, Berlin.

Palmisano, A.C., Wharton, R.A., Cronin, S.E., et al. (1989). Lipophilic pigments from the benthos of a perennially ice-covered Antarctic lake. *Hydrobiologia*. 178: 73-80.

- Parker B.R., Schindler, D.W., Donald, D.B. et al. (2001) The effects of stocking and removal of a nonnative salmonid on the plankton of an alpine lake Ecosystems. 4: 334-345.
- Parker, B.R., Vinebrooke, R.D., Schindler, D.W. (2008) Recent climate extremes alter alpine lake ecosystems. Proceedings of the National Academy of Sciences of the United States of America. 105: 12927-12931.
- Pearce, D.A., Cockell, C.S., Lindstrom E.S., et al. (2007) First evidence for a bipolar distribution of dominant freshwater lake bacterioplankton. Antarctic Science. 19: 245-252.
- Palen, W.J., Schindler, D.E., Adams, M.J., et al. (2002) Optical characteristics of natural waters protect amphibians from UV-B in the U.S. pacific northwest. Ecology. 83: 2951-2957.
- Prentki, R.T., Miller, M.C., Barsdale, R.J., et al. (1980) Chemistry. In Hobbie, J.E. (editor) Limnology of Tundra Ponds. Dowden and Hutchinson and Ross, Pennsylvania, United States.
- Quesada, A., Vincent W.F., and Lean D.R. (1999) Community and pigment structure of Arctic cyanobacterial assemblages: the occurrence and distribution of UV-absorbing compounds. Fems microbiology ecology. 28: 315-323.
- Rautio, M. (1998) Community structure of crustacean zooplankton in subarctic ponds - effects of altitude and physical heterogeneity. Ecography. 21: 327-135.
- Rautio, M. and Vincent, W.F. (2006) Benthic and pelagic food resources for zooplankton in shallow high-latitude lakes and ponds. Freshwater Biology. 51: 1038-1052.
- Sarma, S.S and Nandini, S. (2002) Studies on functional response and prey selection using zooplankton in the anostracan Chirocephalus diaphanus Prevost, 1803. Hydrobiologia. 486: 169-174.
- Saros, J. E., Interlandi, S. J., Wolfe, A. P., et al. (2003) Recent changes in the diatom community structure of lakes in the Beartooth Mountain Range, U.S.A. Arctic, Antarctic, and Alpine Research. 35: 18-23.
- Scavia, D. and Laird, G.A. (1987) Bacterioplankton in Lake Michigan: dynamics, controls, and significance to carbon flux. Limnology and Oceanography. 32: 1017-1033.

- Schmitz, E.H. (1959) Seasonal biotic events in two Colorado alpine tundra ponds. *American Midland Naturalist*. 61: 424-446.
- Schindler, D.W. (1998) Replication versus realism: the need for ecosystem scale experiments. *Ecosystems*. 1: 323-334.
- Schindler, D.W. and Smol, J.P. (2006) Cumulative effects of climate warming and other human activities on freshwaters of arctic and subarctic North America. *Ambio*. 35: 160-168
- Sickman, J.O., Melack, J.M. and Clow, D.W. (2003) Evidence for nutrient enrichment of high-elevation lakes in the Sierra Nevada, California. *Limnology and Oceanography*. 48: 1885-1892.
- Smol, J.P. and Douglas, M.S. (2007) Crossing the final ecological threshold in high Arctic ponds. *Proceedings of the National Academy of Sciences of the United States of America*. 104: 12395-12397.
- Sommaruga, R. (2001) The role of solar UV radiation in the ecology of alpine lakes. *Journal of photochemistry and photobiology*. 62: 35-42.
- Strecker, A. L., Cobb, T. P. and Vinebrooke, R. D. (2004) Effects of experimental greenhouse warming on phytoplankton and zooplankton communities in fishless alpine ponds. *Limnology and Oceanography*. 49: 1182-1190.
- Styrett, P.J. (1981) Nitrogen metabolism of microalgae. *Canadian Bulletin of Fisheries and Aquatic Sciences*. 210: 182-210.
- Thompson, P.L., St-Jacques, M.C., and Vinebrooke, R.D. (2008) Impacts of climate warming and nitrogen deposition on alpine plankton in lake and pond habitats: an in vitro experiment. *Arctic, Antarctic, and Alpine Research*. 40: 192-198.
- Van Geest G. J., Hessen D.O., Spierenburg P., et al. (2007) Goose-mediated nutrient enrichment and planktonic grazer control in arctic freshwater ponds. *Oecologia*. 153: 653-662.
- Vincent, W.F., Castenholz, R.W., Downes, M.T., et al. (1993) Antarctic cyanobacteria – light, nutrients, and photosynthesis in the microbial mat environment. *Journal of phycology*. 29: 745-755.

Vincent, W.F. and Laybourn-Parry, J. (2008) Polar lakes and rivers: limnology of Arctic and Antarctic aquatic ecosystems. (eds.) Vincent, W.F. and Laybourn-Parry J.

Vincent, W.F. and Roy, S. (1993) Solar ultraviolet-B radiation and aquatic primary production: damage, protection and recovery. *Environmental Reviews* 1: 1-12.

Vinebrooke, R.D. and Leavitt, P.R. (1999) Direct and interactive effects of allochthonous dissolved organic matter, inorganic nutrients, and ultraviolet radiation on an alpine littoral food web. *Limnology and Oceanography*. 43: 1065-1081

Walter, K.M., Zimov, S.A., Chanton, J.P. et al. (2006) Methane bubbling from Siberian thaw lakes as a positive feedback to climate warming. *Nature*. 443:71-75.

Ward J.V. (1985) Thermal characteristics in running waters. *Hydrobiologia*. 125: 31-46.

Whalen, S.C., Chalfant, B.A. and Fischer, E.N. (2008) Epipelagic and pelagic primary production in Alaskan Arctic lakes of varying depth. *Hydrobiologia*. 614: 243-257.

Williams, P., Whitfield, M., Biggs, J., et al. (2004) Comparative biodiversity of rivers, streams, ditches and ponds in an agricultural landscape in Southern England. *Biological Conservation* 115: 329-341.

Wolfe A.P., Baron, J.S., and Cornett, J. (2001) Anthropogenic nitrogen deposition induces rapid ecological changes in alpine lakes of the Colorado Front Range (USA). *Journal of Paleolimnology*. 25: 1-7.

Chapter 2 Planktonic responses to elevated nitrogen and phosphorus deposition-a replicated natural alpine pond experiment

Introduction

Anthropogenic emissions have led to dramatic increases in wet deposition of nitrogen (N) globally (Galloway and Cowling 2002; Galloway et al. 2008). The present global rate of anthropogenic N deposition is over 10 times higher than a century ago (Steffan et al. 2004). Nitrogen deposition in western North America is highest downwind of urban areas or agricultural sources, ranging in concentrations from 30 – 90 kg per hectare per year (Fenn et al. 2003). In western Canada, deposition rates currently range from 5 to 10 kg/ha/yr (Murphy et al. 2010), having more than doubled over the past 20 years (Schindler et al. 2006). The severe, long-term ecological consequences of N deposition on ecosystem functioning have been documented mainly for terrestrial, but not aquatic, environments at a global scale (Vitousek 1997).

Several lines of evidence suggest that thousands of small lake and pond ecosystems of the North American Rockies are especially sensitive to increases in N deposition because this element limits their productivity (Baron et al. 2000; McMaster and Schindler 2005; Murphy et al. 2010). Decade-long impacts of increased N deposition on mountain lakes have been inferred paleolimnologically based on fossilized diatom assemblages (Wolfe et al. 2001; Saros et al. 2003). Nydick et al. (2004) reported that experimental addition of nitrogen to small enclosures (500 L) in an alpine lake altered phytoplankton community

composition and stimulated primary production. However, zooplankton did not respond significantly to the N amendments (Nydick et al. 2004), possibly because of the relatively small spatial and temporal scale of their experiment. Nutrient bioassays showed the positive effects of N on primary producers of alpine ponds and lakes in the Canadian Rockies (Murphy et al. 2010). Similarly, experimental N amendment did stimulate alpine primary producers and herbivorous zooplankton when cultured in warmed growth chambers (Thompson et al. 2008). In comparison, solid empirical evidence of the consequences of elevated N deposition on trophic energy transfer (e.g., grazing and predation rates) and higher trophic levels (e.g., fish) at the whole-ecosystem level is not yet available. Nevertheless, small alpine waterbodies are considered susceptible to increased N deposition because they are typically found in steep watersheds with sparse vegetation that retain only small amounts of deposited nutrients (Baron et al. 2000).

The widespread ecosystem effects of airborne (aeolian) phosphorus (P) deposition have also been recently reported for high-elevation lakes. Common sources of aeolian P include plowing, unpaved roads, biomass burning, fossil fuel combustion and deserts. Psenner (1999) found that Saharan dust inputs were important sources of nutrients for remote mountain lakes of the European Alps and Pyrenees. Similarly, anthropogenic sources of P deposition of the high Sierra Nevada (e.g., Emerald Lake) have resulted in the switch from phosphorus to nitrogen limitation in several remote mountain lakes (Sickman et al. 2003).

There is increasing evidence of freshwater ecosystems being co-limited by N and P (Elser et al. 1990, 2007; Ogbebo et al. 2009). Here, the single effect of either nutrient on primary production is amplified by addition of the other, resulting in a

synergistic interaction that denotes co-limitation. Further, nutrient co-limitation is defined by the combined net impact of both nutrients exceeding the sum of their individual effects (i.e. a non-additive effect), which can be detected statistically as a significant interaction using a two-way analysis of variance (ANOVA; Morris and Lewis 1988). *In vitro* fertilization bioassays also suggest that nutrient co-limitation is common among several small lakes and ponds of the Canadian Rocky Mountains (R.D. Vinebrooke, University of Alberta, unpublished data) and the Californian Sierra Nevada (Sickman 2001). Further, low dissolved inorganic nitrogen: total phosphorus (DIN:TP) mass ratios ranging between 2 - 8, of these and other high-elevation aquatic ecosystems highlight their sensitivity to the combined effect of elevated P and N deposition (Michel et al. 2006; Murphy et al. 2010).

Traditionally, ecological theory has predicted that abiotic or bottom-up forces (e.g., nutrient availability) are more important than biotic interactions in determining ecosystem functioning in extreme environments (Menge and Sutherland 1987). However, unexpectedly high abundances of consumers in extreme arctic environments suggest that herbivory may be an important ecological factor structuring primary producers. A study of arctic ponds across a nutrient gradient of N and P showed that zooplankton grazing suppressed primary producers (Van Geest et al. 2007) negating any observable nutrient effect. In addition, mountain lakes have shown strong top down cascading trophic interactions (Parker and Schindler 2006).

Numerous studies have addressed the impacts of nutrients on primary producers through small-scale experiments (Elser et al 2009, Murphy et al. 2010). Unlike such experiments (e.g., bottle bioassays and mesocosms),

ecosystem-level experiments do not suffer from a lack of ecological realism (Carpenter 1996; Schindler 1998). For example, heavy zooplankton grazing, predation, nutrient recycling, benthic microbial biofilms, realistic scale of space and time, temperature fluctuations, wind, and biogeochemical reactions may all play important roles in alpine pond ecosystems (Wilbur 1995) but short term bottle experiments would not capture these processes.

Therefore, I performed a whole-ecosystem experiment to test for the effects of N and P on phytoplankton and zooplankton communities in 16 alpine ponds located along the Eastern Front ranges of the Canadian Rocky Mountains. The driving hypothesis was that phosphorus amendment would amplify the positive effect of nitrogen on planktonic communities (i.e. a synergistic nutrient interaction). The basis for this hypothesis came from an earlier demonstration of nutrient co-limitation of phytoplankton from chemically inferred nitrogen-limited alpine pond ecosystems being driven by phosphorus (R.D. Vinebrooke, University of Alberta, unpublished data).

Materials and Methods

Study Site

The ponds are located near treeline (~ 2200 m above sea level) along Snow Pass at the top of the Cascade Valley catchment (Fig. 1), in the northeastern sector of Banff National Park, Canada (51° 36'N 115° 49'W). They ranged from 0.3 – 1.0 m in depth and were less than 2000 m² in area during the summer of 2008. Physical and chemical parameters of each of the ponds are shown in Table 2. Local geology consists of orthic eutric brunisol soil covered by sparse vegetation and exposed sedimentary rock such as

limestone, shale, and sandstone (Holland and Coen 1982). The surrounding vegetation includes alpine fir (*Abies lasiocarpa*), arctic willow (*Salix arctica*), bladder locoweed (*Oxytropis podocarpa*), Engelmann spruce (*Picea engelmannii*), lichens, sedges (*Carex* spp.), white heather (*Cassiope tetragona*), yellow heather (*Phyllodoce glanduliflora*) and white mountain avens (*Dryas hookeriana*). The nearest available historical climate records are from the townsite of Banff (1383 m a.s.l.; 51° 11' N, 115° 34' W), located approximately 50 km south of the study site, showing a mean summer (June – September) temperature of 12.1°C, and a daily temperature range of -1.4°C to 26.6°C. Small alpine ponds in the region typically freeze in September/October. Average annual precipitation of Banff was 467 mm with 267 mm as rain and 200 mm as snow (McMaster and Schindler 2005).

Experimental Design

A two-factor experiment with random blocking design was replicated four times for a total of 16 ponds. Blocking was based on the proximity of ponds to each other and shared landscape features (ridge vs gully setting; Figure 1). Three blocks of ponds were located atop corresponding ridges along the North Ridge, East Ridge and South Ridge of the study site. A Gully block was selected that contained ponds with a lower elevation and exhibited more willow along the bank than other blocks. The North Ridge block (N=4) was not included in the analysis owing to a brief hydroperiod.

Total volume for each pond was first estimated by measuring the maximum width, length, and depth, and then applying the volumetric formula for a best-fitting geometric shape. The required amounts of added nutrients were then calculated for each pond, ensuring volume-corrected equal dosages were applied across replicate ponds. Nutrients were dissolved in a 4-L fertilizer sprayer containing water, and broadcast over each pond on day 0, 15, and 30 of the experiment. Ammonium nitrate was added to achieve a target concentration of 1000 $\mu\text{g N/L}$ in N-amended ponds while phosphoric acid was added to achieve a final concentration of 30 $\mu\text{g/L}$ in P-amended ponds. These amendments were added simultaneously to all NP-amended ponds. No nutrients were added to a set of control ponds.

These nutrient amendments represented reported worst-case scenarios of anthropogenically elevated deposition over high-elevation sites (Fenn et al. 2003). For example, a target concentration of 1000 $\mu\text{g N/L}$ in the N-amended ponds was based on a deposition rate of 20 kg/ha/yr with the assumption of

no runoff and complete retention of N deposited on the surrounding catchment. In comparison, Fenn et al. (2003) reported annual nitrogen deposition rates in the western United States as high as 90 kg/ha/yr downwind of major urban and agricultural regions. Furthermore, numerous studies have shown that nutrient-rich dust high in phosphorus contribute to atmospheric P deposition in remote alpine environments (Ahl 1988, Psenner 1999, Sickman et al 2003).

Sample Collection

All ponds were sampled on day 0, 15, 30, and 45 of the experiment before and following nutrient amendments. Water samples for chemical and phytoplankton analyses were collected using a two-litre Van Dorn water bottle and acid-washed Nalgene sample containers. Samples for chemical analysis were stored in ice and returned to the laboratory within five days. Chemical analyses were performed by the University of Alberta Biogeochemical Analytical Facility (URL link:<http://www.biology.ualberta.ca/facilities/limnology>). Unfiltered water samples were used for determination of total nutrient concentrations while filtrate using a 0.45- μm filter (pre-ashed Whatman GF/F filters) was used for analysis of dissolved fractions. Total nitrogen (TN), total dissolved nitrogen (TDN), ammonium (NH_4), and nitrate/nitrite ($\text{NO}_3 + \text{NO}_2$) were measured colorimetrically using a Lachat Quikchem 8500 FIA automated ion analyzer with detection limits of 7, 7, 2 and 1 $\mu\text{g/L}$, respectively. Total phosphorus (TP), total dissolved phosphorus (TDP) were quantified by automated Ascorbic Acid Reduction Method using a Lachat QuickChem 8500 multi-channel flow analyzer with a detection limit of 1 $\mu\text{g/L}$. Dissolved organic carbon (DOC) was measured

using a Shimadzu 5000A TOC Analyzer with a detection limit of 0.1 mg/L.

Phytoplankton were concentrated on a Whatman GF/F filter by filtering one litre of pond water. Filters were wrapped in foil, placed in petri dishes, and stored on ice. Algal community structure was quantified using high performance liquid chromatographic analysis of taxonomically diagnostic chlorophylls and carotenoids (Vinebrooke et al. 2002). Total phytoplankton biomass was inferred based on concentrations of chlorophyll *a* (all algae), while community composition was estimated using chlorophyll *b* (chlorophytes), fucoxanthin (chrysophytes, diatoms, some dinoflagellates), lutein (chlorophytes and euglenoids), and zeaxanthin (cyanobacteria). Qualitative analysis of the preserved phytoplankton samples using light microscopy was conducted to determine taxonomic composition.

Zooplankton were collected using a 64-micron mesh-sized plankton net, which was hauled at a rate of 1 m/s horizontally across each pond basin. Samples were immediately preserved in 70% ethanol. A minimum of 300 individuals per sample were measured and identified to species using a stereomicroscope. Standard length-weight regression equations were applied to calculate invertebrate biomass (McCauley 1971). During the first sample period, juvenile fairy shrimp (*Branchinecta paludosa*) were effectively captured with the plankton net. However, capture efficiency appeared to decrease during the experiment as adult fairy shrimp were observed effectively evading net hauls.

Statistical Analysis

Fertilization effects on total chlorophyll-inferred algal abundance and total zooplankton biomass were tested statistically using a one factor Repeated Measure Analysis of Variance (RM-ANOVA) while Repeated Measure Multivariate Analysis of Variance (RM-MANOVA) of phytoplankton pigments (fucoxanthin, chlorophyll *b*, and lutein/zeaxanthin) was used to test for treatment effects on community composition. East Ridge Block P Treatment (ERP), South Ridge Block Nitrogen Treatment (SRN) and all ponds in the North Ridge Block with the exception of the NP treatment were dry by day 30 of the experiment (July). Gully Block phosphorus was not included in the analysis as it had abnormally high concentrations of total chlorophyll at the beginning of the experiment. Therefore a total of six ponds were not included in the analysis, resulting in an uneven loss of replicates across treatment combinations. Consequently, it was not possible to conduct a two-factor ANOVA of the zooplankton/phytoplankton data for the entire experiment

Canonical Correspondence Analysis (CCA) was performed using CANOCO version 4.5 (ter Braak 1998) to determine the environmental variables that best explained changes in zooplankton community composition and phytoplankton presence/absence data. CCA is a multivariate technique that tests differences in community structure between samples and compares the maximum amount of variation in the taxonomic data set to environmental variables. Each environmental variable (log-transformed) is represented by a CCA axis and its eigenvalue (λ), which is a measure of the explained species variance. The length and orientation of each environmental vector symbolizes

its importance and direction of maximum change in ordination space. The proximity and alignment of a species with a particular vector shows how closely the variation in its abundance is related to that environmental variable. Similarly, site score differences in ordination space suggest taxonomic dissimilarity of invertebrate assemblages between different ponds, or across time. Time, DOC, TP, TN, PN, PP, and chlorophyll *a* were included in the model.

Results

Water chemistry

Chemical conditions in all of the experimental ponds were similar prior to the start of the experiment (Table 2). Most importantly, TN, TDN, TP, TDP and DOC concentrations did not differ among the ponds on day 0. All ponds initially contained 14 – 33 $\mu\text{g TP/L}$, making them naturally mesotrophic. Further, ponds were noticeably stained in appearance, containing high concentrations of brownish DOC.

Immediately following fertilization, TP and TN concentrations exceeded target levels of 1000 $\mu\text{g N/L}$ and 30 $\mu\text{g P/L}$ (Figs. 2a, b) on day 0 through day 30 of the experiment. Nutrient target concentrations were exceeded during a few occasions probably due to over estimates of pond volumes. Upon returning to the ponds following a two-week period, TN and TP in the water column significantly declined in concentration (Figs. 3a, b). The majority of the added nutrients did not persist in the water column in a dissolved or particulate form two weeks following fertilization.

Low concentrations of particulate fractions of TN and TP did not differ significantly among the nutrient-amended ponds (Figs. 4a, b). TN consisted mainly of TDN relative to PN, which accounted for only ~10% of the total nitrogen pool. Similarly, TP was comprised primarily of TDP while PP represented only ~30% of the total phosphorus pool.

Phytoplankton

N and P did not significantly (fertilization effect, $F_{3,40} = 0.17$, $P = 0.91$) increase phytoplankton chlorophyll *a* concentrations despite three additions of nutrients over a 45-day period (Fig. 5). Instead, chlorophyll *a* concentrations remained low ($< 2 \mu\text{g/L}$), showing high variance among the ponds during the experiment. Chlorophyll *a* did not vary significantly over time (time effect, $F_{3,40} = 8.70$, $P = 0.10$), but steadily increased from day 0 to 30 before declining thereafter by day 45.

Fertilization also did not significantly ($F_{9,40} = 0.20$, $P = 0.49$) alter carotenoid-inferred phytoplankton community composition (Fig. 6). However, RM – MANOVA using three pigments common to all ponds (fucoxanthin, lutein/zeaxanthin, chlorophyll *b*) showed that community composition changed significantly over time (RM MANOVA; $F_{9,40} = 2.261$, $P = 0.038$). Specifically, three carotenoid concentrations increased during the first half of the experiment, before declining by day 45 (Fucoxanthin, $F_{3,40} = 3.09$, $P = 0.053$; Chlorophyll *b*, $F_{3,40} = 5.17$, $P = 0.009$; Lutein/zeaxanthin, $F_{3,40} = 5.52$, $P = 0.007$).

CCA using automatic backward selection showed that fertilization variables of (TN; $F = 0.97$, $P = 0.44$, TP; $F = 0.78$, $P = 0.71$), and DOC ($F = 0.84$, $P = 0.6$) were not significant predictors of phytoplankton community composition after 45 days (Fig. 7). The first axis was best explained by TN accounting for only 13% of the variance in the taxonomic presence/absence data. The second axis was best explained by TP accounting for only 8% of the variance in the taxonomic data.

Zooplankton

Fertilization also did not significantly affect total zooplankton biomass during the experiment (Fig. 8; time-fertilization effect, $F_{3,40} = 0.36$, $P = 0.78$). Total zooplankton biomass did not change significantly over time (time effect; $F_{3,40} = 0.45$, $P = 0.55$). However, two phosphorus-amended ponds did contain substantially greater, but also highly variable zooplankton biomass by the end of the experiment, consisting mainly of fairy shrimp.

CCA using automatic backward selection showed that time ($F = 10.49$, $P = 0.002$), TP ($F = 3.70$, $P = 0.008$) and TN ($F = 3.96$, $P = 0.03$) were significant predictors of zooplankton community composition across the experimental ponds (Fig. 9; App 1). The first axis was best explained by time accounting for 21% of the variance in the taxonomic abundance data. Ponds contained juvenile zooplankton (nauplii and copepodites, small fairy shrimp) early in the summer and contained more adults later on in the season. The second axis captured 12% of the variance in the taxonomic data, being best represented by opposing gradients of

N and P. Here, diverse assemblages of cladocerans and copepods were associated with higher concentrations of TN while adult fairy shrimp were more closely related to TP. Adult fairy shrimp were detected in all ponds at one point in time during the experiment but fertilization did affect fairy shrimp growth rate.

Discussion

My experimental findings did not support the co-limitation hypothesis, namely that phosphorus additions would amplify the positive effects of nitrogen on these chemically inferred nitrogen-limited alpine pond ecosystems. Despite low DIN:TP ratios across all ponds, neither nitrogen nor phosphorus amendments significantly stimulated phytoplankton or zooplankton abundance. Although aggregate community properties did not respond significantly to fertilization, elevated N and P levels were significant explanatory variables of species-level shifts in the zooplankton communities. Several potential ecological and methodological explanations could account for this surprising lack of statistically detectable effects of N and P.

One possible ecological explanation for the lack of a phytoplankton response to nutrient fertilization involves suppression by intensive grazing pressure (Schindler and Comita 1972; Elser et al. 2000). Two observations lend support to this explanation: very low chlorophyll concentrations relative to dissolved nutrient concentrations, and observed high densities of filter-feeding fairy shrimp. For instance, Van Geest et al. (2007) found strong top-down regulation of phytoplankton by cladocerans precluded any significant relationships between chlorophyll and measured nutrients in a series of arctic

ponds. Parker and Schindler (2006) also showed pronounced grazing pressure and a strong grazer-algal interaction in an alpine lake located near my study area.

Fairy shrimp were very abundant in the study ponds and may even be regarded as a keystone species. In an Antarctic study, more than 40000 juvenile *Branchinecta gaini* were reported in a cubic meter, feeding on 8 μg of chlorophyll *a* per hour per individual (Hawes 1985). However, limited research has been conducted on their feeding habits and life history. Hawes (1985) suggested that immature *Branchinecta gaini* feed in the water column but adopt a benthic habitat when mature. Although many fast-swimming fairy shrimp likely evaded horizontal net hauls with a plankton net during the sampling of my experimental ponds, observation of high densities of this species on surface sediments attest to its functional importance in these systems. Also, ponds drying down may have concentrated fairy shrimp and over-inflated biomass estimates.

Competition with periphyton may have also contributed to suppression of a phytoplankton response to added nutrients. Importantly, periphyton can be an order of magnitude more abundant than phytoplankton in shallow (~1-m depth) habitats, providing them with a competitive advantage over phytoplankton for resources (Hansson 1988; Vinebrooke & Leavitt 1999; Vadeboncoeur et al. 2003). A mesocosm experiment by Confer (1972) showed periphyton immediately sequestered large amounts of phosphorus and decreased soluble phosphorus as well as phytoplankton standing stock. Hansson (1990) also found that mesocosms without periphyton contained 32-44% more phosphorus in the water column compared to controls including

periphyton. Furthermore, a bioassay of shallow oligotrophic mountain lakes in the Colorado Front Range showed that N uptake by the benthic processes may be more important than in the pelagic (Nydick et al. 2004). Periphyton responded significantly to the addition of N in the ponds of this study (L Porter, University of Alberta, unpublished data).

Bacteria may also have taken up the added nutrients before an algal response was possible. Bacteria may represent a substantial fraction of the TPP given high concentrations of DOC likely fueling heterotrophy in these ponds (Siuda and Chrost 2002). Heterotrophic bacteria may be responsible for low chlorophyll concentrations relative to ambient nutrient levels in the study ponds. Similarly, laboratory and field based studies have shown that bacterial production in humic lakes is more important than algal primary production (Tranvik and Hofle 1987, Arvola and Kankaala 1989, Tranvik 1989, Jones 1990). However, significant bacterial biomass responses to the added nutrients would have been reflected in elevated particulate N and P fractions, which was not the case in my experiment.

Another possible explanation involves high concentrations of DOC in these ponds resulting in organic binding of amended nutrients, thereby reducing their bioavailability. DOC may inhibit microbial growth by binding with inorganic nutrients such as iron (Jackson and Hecky 1980; Guildford et al 1987) and phosphorus (Stewart and Wetzel 1981) as well as deactivating phosphatases (Wetzel 1992; Boavida and Wetzel 1998). Experimentally added orthophosphate may bind with DOC but may be less of a consequence than nitrogen-binding because of the likely presence of phosphatases (Jones et al. 1988). Therefore, chemical complexation of the added inorganic

nutrients may have precluded algal uptake, thereby dampening the expected positive effect of fertilization as measured by chlorophyll *a*-inferred phytoplankton biomass.

The observed general lack of detectable statistically significant responses of planktonic communities to nutrient amendments may also have been a consequence of poor study power owing to insufficient replication. Power analysis suggests more than sixty ponds would have been required to detect a significant difference between treatment groups based on an effect size of 0.26. There was a high degree of variation between study ponds with similar treatment. For instance, in the control treatment the coefficient of variation was 0.65 on day 30 of the experiment. Pond permanence is another important factor in determining zooplankton community structure (Wellborn et al. 1996; Brooks 2000; Rundle et al. 2002; Urban, 2004; Jocque et al., 2007). Ponds that dry out quickly cannot support long living species that require time to reproduce before desiccation.

Although the spatial scale of the experiment captured whole-ecosystem responses to nutrient amendments, its temporal scale was limited to a part of single growing season. I believe that if the experiment had run a month or two longer treatment effects may have been more evident. Large filamentous blooms of green algae were observed in two of the NP treatments in support of our hypothesis and another had turned dark brown. Therefore, had the experiment lasted a month longer we may have observed more pronounced effects between treatment groups, most notably in the NP treatment. The moment of ice out may have been unique to each pond studied corresponding to a phytoplankton bloom that we may have missed. Enrichment of ponds

during ice out may be critical and the seasonality influences of ponds may make them incomparable.

Though I have little empirical evidence, field observations suggest that unexpected swarms of fairy shrimp were in all likelihood the primary reason why added nutrients did not stimulate either phytoplankton or zooplankton abundance in these alpine ponds. Further research including the grazing effects on periphyton and phytoplankton communities may provide a greater understanding of the nutrient status of alpine ponds. Overall despite the high degree of variance found within treatment groups and the lack of replication, whole ecosystem projects are an invaluable component to understanding ecosystem processes and community interactions (Schindler 1998).

Biogeochemical interactions within sediment, presence of periphyton and multiple trophic levels, and longer experimental time periods are all factors left unconsidered in previous bottle experiments (Murphy et al. 2010).

Literature Cited

Ahl, T. (1988) Background yield of phosphorus from drainage area and atmosphere: An empirical approach. *Hydrobiologia*. 170: 35-44.

Arvola, L., and Kankaala, P. (1989) Winter and Spring Variability in Phytoplankton and Bacterioplankton in Lakes with Different Water Colour. *Aqua Fennica*. 19: 29-39.

Baron, J.S., Rueth, H.M., Wolfe, A.M., et al. (2000) Ecosystem responses to nitrogen deposition in the Colorado Front Range. *Ecosystems*, 3:352-368.

Boavida, M.J. and Wetzel, R.G. (1998) Inhibition of phosphatase activity by dissolved humic substances and hydrolytic reactivation by natural ultraviolet light. *Freshwater Biology*. 40: 285-293.

Brooks, R. T. (2000) Annual and seasonal variation and the effects of hydroperiod on benthic macroinvertebrates of seasonal forest (“vernal”) ponds in central Massachusetts, USA. *Wetlands*. 20: 707–715.

Carpenter, S.R. (1996) Microcosm experiments have limited relevance for community and ecosystem ecology. *Ecology*. 77: 677-680.

Carpenter, S.R., Kitchell, J.F., Cottingham, K.L., et al. (1996) Chlorophyll variability, nutrient input, and grazing: Evidence from whole-lake experiments. *Ecology*. 77: 725-735.

Confer, J.L. (1972) Interrations among plankton, attached algae, and phosphorus cycle in artificial open systems. *Ecological Monographs*. 42: 1-23.

Elser, J.J., Bracken, M.E., Cleland, E.E., et al. (2007) Global analysis of nitrogen and phosphorus limitation of primary producers in freshwater, marine and terrestrial ecosystems. *Ecology Letters*. 10: 1135–1142.

Elser, J.J., Kyle, M., Steger, L., et al. (2009) Nutrient availability and phytoplankton nutrient limitation across a gradient of atmospheric nitrogen deposition. *Ecology*. 90: 3062-3073.

Elser, J.J., Marzolf, E.R., and Goldman, C.R. (1990) Phosphorus and nitrogen limitation of phytoplankton growth in the freshwaters of North America: a review and critique of experimental enrichments. *Canadian Journal of Fisheries and Aquatic Science*. 47: 1468–1477.

- Elser, J.J., Sterner, R.W., Galford, A.E., et al. (2000) Pelagic C:N:P stoichiometry in a eutrophied lake: responses to a whole-lake food-web manipulation. *Ecosystems*. 3: 293-307.
- Fenn, M., Haeuber, R., Tonnesen, G., et al. (2003) Nitrogen emissions, deposition, and monitoring in the western United States. *Bioscience*. 53: 391-403.
- Galloway, J.N., and Cowling E.B. (2002) Reactive nitrogen and the world: 200 years of change. *Ambio*. 31: 64-71.
- Galloway, J.N., Townsend, A.R., Erisman, J.W., et al. (2008) Transformation of the nitrogen cycle: recent trends, questions, and potential solutions. *Science*. 320: 889-892.
- Guildford, S.J., Healey, F.P., and Hecky, R.E. (1987) Depression of primary production by humic matter and suspended sediment in limnocorral experiments at southern Indian lake, northern Manitoba. *Canadian Journal of Fisheries and Aquatic Sciences*. 44: 1408-1417.
- Hansson, L.A. (1988) Effects of competitive interactions on the biomass development of planktonic and periphytic algae in lakes. *Limnology and Oceanography*. 33: 121-128.
- Hansson, L.A. (1990) Quantifying the impact of periphytic algae on nutrient availability for phytoplankton. *Freshwater Biology*. 24: 265-273.
- Hawes, I. (1985) Factors controlling phytoplankton populations in maritime Antarctic lakes. *Antarctic Nutrient Cycles and Food Webs 4th scientific committee on Antarctic research symposium on Antarctic biology*.
- Holland, W.D. and Coen, G.M. (1982) Ecological (biophysical) land classification of Banff and Jasper National Parks. Volume II: soil and vegetation resources and map supplement. Alberta Institute of Pedology.
- Jackson, T.A. and Hecky, R.E. (1980) Depression of primary productivity by humic matter in lake and reservoir waters of the boreal forest zone. *Canadian Journal of Fisheries and Aquatic Sciences*. 37: 2300-2317.
- Jocque', M., Riddoch, B.J., and Brendonck, L. (2007) Successional phases and species replacements in freshwater rock pools: towards a biological definition of ephemeral systems. *Freshwater Biology*. 52: 1734-1744.

Jones, R.I., Salonen, K., and De Haan, H. (1988) Phosphorus transformations in the epilimnion of humic lakes: abiotic interactions between dissolved humic materials and phosphate. *Freshwater Biology*. 19: 357-369.

Jones, R.I. (1990) Phosphorus transformations in the epilimnion of humic lakes: biological uptake of phosphate. *Freshwater Biology*. 23: 323-337.

McCauley, E. (1971) The estimation of the abundance and biomass of zooplankton in samples. *In* Downing, J.A. and Rigler, F.H. (eds.) *A manual on methods for the assessment of secondary productivity in fresh waters*. Oxford. Blackwell Scientific Publications, 228-261.

McMaster, N.L. and Schindler, D.W. (2005) Planktonic and epipelagic algal communities and their relationship to physical and chemical variables in alpine ponds in Banff National Park, Canada. *Arctic, Antarctic, and Alpine Research*. 37: 337-347.

Menge, B.A. and Sutherland, J.P. (1987) Community regulation variation in disturbance, competition, and predation in relation to environmental stress and recruitment. *American Naturalist* 130: 730-757.

Michel, T.J., Saros, J.E., Interlandi, S.J., and et al. (2006) Resource requirements of four freshwater diatom taxa determined by in situ growth bioassays using natural populations from alpine lakes. *Hydrobiologia*. 568: 235-243.

Morris, D. P. and Lewis, W. M. (1988) Phytoplankton nutrient limitation in Colorado mountain lakes. *Freshwater Biology*. 20: 315-327.

Murphy, C.A., Thompson, P.L., and Vinebrooke, R.D. (2010) Assessing the sensitivity of alpine lakes and ponds to nitrogen deposition in the Canadian Rocky Mountains. *Hydrobiologia*. 648: 83-90.

Nydick, K. R., Lafrancois, B. M., Baron, J. S., et al. (2004) Nitrogen regulation of algal biomass, productivity, and composition in shallow mountain lakes, Snowy Range, Wyoming, USA. *Canadian Journal of Fisheries and Aquatic Sciences*. 61: 1256-1268.

Ogbebo, F.E.; Evans, M.S.; Waiser, M.J., et al. (2009) Nutrient limitation of phytoplankton growth in Arctic lakes of the lower Mackenzie River Basin, northern Canada. *Canadian Journal of Fisheries and Aquatic Sciences*. 66: 247-260.

- Parker, B.R. and Schindler, D.W. (2006) Cascading trophic interactions in an oligotrophic species-poor alpine lake. *Ecosystems*. 9: 157-166.
- Psenner, R. (1999) Living in a dusty world: airborne dust as a key factor for alpine lakes. *Water and Soil Pollution*. 112: 217-227.
- Rundle, S. D., Foggo, A., Choiseul, V., et al. (2002). Are distribution patterns linked to dispersal mechanism? An investigation using pond invertebrate assemblages. *Freshwater Biology*. 47: 1571–1581.
- Saros, J.E., Interlandi, S.J., Wolfe, A.P., et al. (2003) Recent changes in the diatom community structure of lakes in the Beartooth Mountain Range, USA. *Arctic, Antarctic, and Alpine Research*. 35: 18-23.
- Schindler, D.W. (1998) Replication versus realism: the need for ecosystem scale experiments. *Ecosystems*. 1: 323-334.
- Schindler, D.W. and Comita, G.W. (1972) The dependence on primary production upon physical and chemical factors in a small, senescing lake, including the effects of complete winter oxygen depletion. *Archiv fur hydrobiology*. 69: 413-451.
- Schindler, D.W., Dillon, P.J., Schreier, H. (2006) A review of anthropogenic sources of nitrogen and their effects on Canadian aquatic ecosystems. *Biogeochemistry*. 79: 25-44.
- Sickman, J. O. (2001) Comparative analyses of nitrogen biogeochemistry in high-elevation ecosystems. Doctoral dissertation, Univ. of California, Santa Barbara.
- Sickman, J.O., Melack, J.M. and Clow, D.W. (2003) Evidence for nutrient enrichment of high-elevation lakes in the Sierra Nevada, California. *Limnology and Oceanography*. 48: 1885-1892.
- Siuda, W. and Chrost, R.J. (2002) Decomposition and utilization of particulate organic matter by bacteria in lakes of different trophic status. *Polish Journal of Environmental Studies*. 11:53-65.
- Steffan, W., Sanderson, A., Tyson, P.D., et al. (2004) Global change and the earth system: a planet under pressure. IGBP global change series. Springer-Verlag, Berlin.

Stewart, A.J. and Wetzel R.G. (1981) Dissolved humic materials – photodegradation, sediment effects, and reactivity with phosphate and calcium carbonate precipitation. *Archiv fur hydrobiologie*. 92: 265-286.

ter Braak, C. and Milauer, P. (1998) CANOCO reference manual and user's guide to CANOCO for Windows: Software for Canonical community ordination (version 4). New York: Microcomputer Power.

Thompson, P.L., St-Jacques, M.C., and Vinebrooke, R.D. (2008) Impacts of climate warming and nitrogen deposition on alpine plankton in lake and pond habitats: an in vitro experiment. *Arctic, Antarctic, and Alpine Research*. 40: 192–198.

Tranvik L.J. (1989) Bacterioplankton growth, grazing mortality and quantitative relationship to primary production in a humic and a clearwater lake. *Journal of Plankton Research*. 11: 985-1000.

Tranvik, L.J. and Hofle M. (1987) Bacterial growth in mixed cultures on dissolved organic carbon from humic and clear waters. *Applied Environmental Microbiology*. 53: 482-488.

Urban, M. C. (2004) Disturbance heterogeneity determines freshwater metacommunity structure. *Ecology*. 85: 2971-2978.

Vadeboncoeur, Y., Jeppesen, E., Vander Zanden, M.J., et al. (2003) From Greenland to green lakes: cultural eutrophication and the loss of benthic pathways in lakes. *Limnology and Oceanography*. 48: 1408-1418.

Van Geest, G. J., Hessen, D.O., Spierenburg P., et al. (2007) Goose-mediated nutrient enrichment and planktonic grazer control in arctic freshwater ponds. *Oecologia*. 153: 653–662.

Vinebrooke, R.D., Dixit, S.S., Graham, M.D., et al. (2002) Whole-lake algal responses to a century of acidic deposition on the Canadian Shield. *Canadian Journal of Fisheries and Aquatic Sciences*. 59: 483-493.

Vinebrooke, R.D. and Leavitt, P.R. (1999) Direct and interactive effects of allochthonous dissolved organic matter, inorganic nutrients, and ultraviolet radiation on an alpine littoral food web. *Limnology and Oceanography*. 43: 1065-1081.

Vitousek, P., Aber, J., Howarth, R., et al. (1997) Human alteration of the global nitrogen cycle: Sources and consequences. *Ecological Applications*. 7: 737-750.

Wellborn, G.A., Skelly, D.K., Werner, E.E. (1996) Mechanisms creating community structure across a freshwater habitat gradient. *Annual Review of Ecology and Systematics*. 27: 337-363.

Wetzel, R.G. (1992) Gradient-dominated ecosystems: sources and regulatory functions of dissolved organic matter in freshwater ecosystems. *Hydrobiologia*. 229: 181-198.

Wilbur, H.M. (1995) Experimental ecology of food webs: complex systems in temporary ponds. *Ecology*. 78: 2279-2302.

Wolfe A.P., Baron, J.S., and Cornett, J. (2001) Anthropogenic nitrogen deposition induces rapid ecological changes in alpine lakes of the Colorado Front Range (USA). *Journal of Paleolimnology*. 25: 1-7.

Table 1 Average Total Phosphorus (TP) and Total Nitrogen (TN) concentrations and standard error (SE) of alpine and arctic ponds reported by various studies. Hyphenated sections represent data unavailable.

	TP ($\mu\text{g/L}$)	SE	TN ($\mu\text{g/L}$)	SE	n
Lim et al. 2005	21.8	6.3	631.3	80.6	29
McMaster and Schindler 2005	51.3	11.8	-	-	28
Oertli et al. 2008	-	-	240	22.5	16
Murphy et al. 2010	19.8	4.0	382.2	61.4	23
Zettel and Vinebrooke, unpubl.	23.5	1.8	350.0	28.7	12

Table 2 Physical and chemical parameters of studied alpine ponds in the Cascade Valley, Banff National Park, Canada.

Block	Treatment	Latitude (dd mm ss)	Longitude (dd mm ss)	Area (m ²)	Depth (m)	TN ($\mu\text{g L}^{-1}$)	TDN ($\mu\text{g L}^{-1}$)	PN ($\mu\text{g L}^{-1}$)	TP ($\mu\text{g L}^{-1}$)	TDP ($\mu\text{g L}^{-1}$)	PP ($\mu\text{g L}^{-1}$)	DOC (mg L ⁻¹)	Chl a ($\mu\text{g L}^{-1}$)
1	C	51° 36' 16 N	115° 49' 04 W	390	0.8	226	209	17	14	7	7	5.8	1.02
1	N	51° 36' 12 N	115° 49' 04 W	240	0.3	427	389	38	18	10	8	9.6	0.05
1	P	51° 36' 03 N	115° 49' 10 W	180	0.4	406	356	50	30	18	12	8.7	0.87
1	NP	51° 36' 15 N	115° 49' 00 W	120	0.3	353	349	4	27	12	15	6.9	0.13
2	C	51° 36' 01 N	115° 49' 11 W	176	0.6	299	266	33	14	9	4	8.6	0.52
2	N	51° 36' 45 N	115° 49' 46 W	308	0.3	385	316	31	22	21	1	8.9	0.19
2	P	51° 36' 44 N	115° 49' 48 W	1000	0.5	310	264	46	25	17	8	6.6	0.72
2	NP	51° 36' 02 N	115° 49' 09 W	286	0.4	549	525	24	30	21	9	11.5	0.57
3	C	51° 36' 23 N	115° 49' 24 W	862	1.0	487	472	15	*	*	0	8.2	1.07
3	N	51° 36' 45 N	115° 49' 36 W	690	0.6	290	266	24	28	20	8	7.2	0.30
3	P	51° 36' 17 N	115° 49' 08 W	324	0.3	673	638	217	129	16	9	10.2	8.85
3	NP	51° 36' 46 N	115° 49' 46 W	495	0.6	236	216	20	33	21	12	5.4	1.15
4	C	51° 36' 50 N	115° 49' 51 W	450	0.3	364	318	20	40	30	10	9.9	0.29
4	N	51° 36' 51 N	115° 49' 51 W	90	0.3	545	503	32	35	28	12	13.8	0.37
4	P	51° 36' 43 N	115° 49' 47 W	1000	0.5	325	284	32	28	20	27	8.7	0.82
4	NP	51° 36' 51 N	115° 49' 47 W	1512	0.5	233	203	30	19	12	7	6.2	0.44




Figure 1 Study site location of surveyed alpine waterbodies in Banff National Park, Canada




Figure 2 (A) Total Nitrogen (TN) and (B) Total Phosphorus (TP) concentrations in the water column. C (Control), N (Nitrogen), P (Phosphorus), and NP (Nitrogen +Phosphorus) treatment concentrations were measured immediately following nutrient amendment. Error bars represent one standard-error unit based on n = 3.




Figure 3 (A) Total nitrogen (TN) and (B) total phosphorus (TP) concentrations in the water column 15 days following nutrient amendments: N, P, and NP treatments. Dashed line depicts the average concentration of each nutrient immediately following amendment (see Fig. 1). Error bars represent one standard-error unit based on $n = 3$.




Figure 4(A) Particulate nitrogen (PN) and (B) particulate phosphorus (PP) prior to each nutrient amendment during the 45-d experiment: C (Control), N (Nitrogen), P (Phosphorus), and NP (Nitrogen + Phosphorus). Error bars represent one standard-error unit based on n = 3.




Figure 5 Phytoplankton chlorophyll a concentration prior to each nutrient amendment during the 45-d experiment: C (Control), N (Nitrogen), P (Phosphorus), and NP (Nitrogen +Phosphorus). Error bars represent one standard-error unit based on n = 3.




Figure 6 Pigment-inferred abundance of (A) Chromophytes (Fucoxanthin) (B) Chlorophytes/Cyanobacteria (Lutein/Zeaxanthin) (C) Chlorophytes and Euglenophytes (Chlorophyll *b*) prior to each nutrient amendment during the 45-d experiment: C (Control), N (Nitrogen), P (Phosphorus), and NP (Nitrogen + Phosphorus). Error bars represent one standard-error unit based on n = 3.




Figure 7 (A) Ordination biplot of phytoplankton taxa and explanatory environmental variables during the 45-d experiment based on canonical correspondence analysis. Total phosphorus (TP), total nitrogen (TN) and dissolved organic carbon (DOC) are shown as passive, statistically non-significant predictors of taxonomic composition.





Figure 8 Total zooplankton biomass prior to each nutrient amendment during the 45-d experiment: (C) Control, N (Nitrogen), P (Phosphorus), NP (Nitrogen+Phosphorus). Error bars represent one standard-error unit based on $n = 3$.



Zooplankton Species Codes

Juvenile <i>Branchinecta paludosa</i>	JuvBraPal
Adult <i>Branchinecta paludosa</i>	BraPal
Cyclopoid Nauplii	CycNau
Cyclopoid instar (1-3)	Cyc1_3
Cyclopoid instar (4-6)	Cyc4_6
Cyclops vernalis	CycVer
Calanoid nauplii	CalNau
Calanoid instar (1-3)	Cal1_3
Calanoid instar (4-6)	Cal4_6
<i>Diaptomus nudus</i>	DiaNud
<i>Diaptomus shoshone</i>	DiaSho
<i>Chydorus sphaericus</i>	ChySph
<i>Daphnia pulex</i>	DapPul
<i>Ceriodaphnia laticaudata</i>	CerLat

Figure 9 Ordination biplot of zooplankton taxa with species codes and explanatory environmental variables during the 45-d experiment. Total phosphorus (TP), total nitrogen (TN) and time are shown as statistically significant predictors of community composition.

Chapter 3 Conclusions

Synthesis

Higher trophic levels in alpine pond ecosystems such as zooplankton and fairy shrimp may be ecologically important and control the abundance and species composition of primary producers. These findings contradict experimental evidence and surveys that suggest shallow-water communities along the littoral area of alpine lakes are principally regulated by resources and not density-dependent species interactions (Vinebrooke and Leavitt 1998; Vinebrooke and Leavitt 1999). Numerous (> 100 individuals/ m^2) *Branchinecta paludosa* were observed in the ponds and intensive grazing pressure suppressed the response of primary producers to nutrient additions. Nitrogen and phosphorus amendments did not increase phytoplankton or zooplankton abundance suggesting strong biotic interactions (Dodson 1974). The importance of grazing overshadowing nutrient effects has also been reported in similar pond ecosystems (Van Geest et al. 2007).

Benthic processes that were not reported on in this study likely play very important functional roles (e.g., nutrient cycling, trophic energy transfer) in shallow-water ecosystems. Microbial mats of benthic algae may quickly outcompete phytoplankton and assimilate available DIN in the water column (Axler and Reuter 1996; Mcknight et al. 2004; Nydick et al. 2004). In combination with extensive grazing by zooplankton and fairy shrimp low concentrations of phytoplankton are therefore at a competitive disadvantage for nutrients in these systems.

Future Research Directions

Environmental monitoring in the remote regions of the mountain parks of Banff, Jasper and Yoho as well as the surrounding area including Kananaskis Country should be improved to better demonstrate changes in nitrogen and phosphorus deposition. High-elevation meteorological field stations are sparse throughout the North American Rockies, especially in Canada (Bradley et al. 2004). Therefore, automated or regular manual monitoring of oxidized forms of N (NO_x) and reduced forms of N (NH_3 and NH_4) in rainfall, snow and surface waters would be important to detect expected increases in nitrogen deposition in any of these regions (Fenn et al. 2003).

Limited research has been conducted on the physical, chemical, and biological patterns of alpine ponds in North America (McMaster and Schindler 2005). Additional research should include monthly surveys over a number of years throughout the ice free season to document physical, chemical and biological patterns to gain a greater understanding of these ecologically important and abundant shallow systems. A strong sampling emphasis on the periphyton and higher trophic levels could aid in our understanding of trophic dynamics and species interactions.

Mesocosm and split-basin experiments may be a more effective way to empirically demonstrate nutrient limitation in alpine ponds given the observed high degree of heterogeneity or variance that exists among these ecosystems. In other words, one should not assume that the similar physical appearance of these systems reflects habitat homogeneity among ponds.

Instead, a split-basin experiment would enable the investigator to control for site-specific differences in abiotic and biotic factors, which may otherwise regulate the effects of nutrient amendments. Further, factorial experiments could be conducted in each pond to examine the hypothesized mediating influences of fairy shrimp and periphyton on planktonic communities. For example, nutrients could be added directly to surface waters contained in mesocosms where either ambient densities of fairy shrimp are excluded or enhanced relative to controls. Similarly, open- versus closed-bottom mesocosms could be used to examine the interactive effect of benthos and nutrient fertilization on planktonic communities.

Grazing experiments examining the effects of different zooplankton taxonomic groups on primary producers may account for some of the unexplained variation in this study. Ponds were dominated by unequal abundances of a variety of taxonomic groups including copepods, cladocerans, fairy shrimp, as well as higher trophic levels such as *Chaoborus*. The dissimilarity in filtering rates, size selection strategies and natural history of these grazers may ultimately affect primary producer abundance in the benthic and planktonic regimes. Analysis incorporating both planktonic and benthic communities may also aid in a greater understanding of these unique systems.

The ecology of fairy shrimp should be directly examined in these systems as this ubiquitous species appears to play a keystone role in alpine and polar pond ecosystems. The sampling effort of fairy shrimp would have to be improved to generate more accurate and precise estimates of this species' abundance in these ponds. Sampling with a net haul at night may improve

capture efficiency because fairy shrimp may be less likely to see the moving net. Fairy shrimp were quickly able to evade an approaching net and were often observed burrowing under the fine pond sediment. Light traps could be set at night to capture fairy shrimp. Similar to fish stock assessment fairy shrimp abundance could be compared based on a catch per unit effort basis that may be more effective than net hauls.

Alpine ponds are extreme environments that show strong daily temperature fluctuations, lack thermal stratification, and exhibit rapid changes in the physical and chemical parameters over a short ice free season (McMaster 2003). Climate change will likely amplify the effects of these extreme conditions on the biological communities of ponds in polar and alpine regions through increased frequency of drought, pond desiccation and stronger temperature fluctuations (Douglas and Smol 1999). In addition to these stressors alpine pond communities may be particularly sensitive to increased warming, species invasions, nitrogen/phosphorus deposition, persistent organic pollutants and other atmospheric inputs.

Further research is required to understand the potential effects of increased atmospheric deposition on shallow waterbodies in the Canadian Rocky Mountains. Anthropogenic emissions estimates of N in Canada are over 1.4 million tonnes annually and are expected to increase further (Chambers et al. 2001). My limited research does not suggest that widespread eutrophication is likely under current deposition estimates but a multi year study with increased replication should be conducted to compare my findings. Studies have demonstrated that climate change and atmospheric nitrogen deposition alter alpine (Saros et al. 2003) and arctic (Wolfe et al. 2006) phytoplankton communities as

well as higher trophic levels (Thompson et al. 2008). These studies highlight the importance of understanding the effects of multiple stressors on alpine communities. Moreover, the combination of multiple stressors such as atmospheric deposition of pollutants, UV light, and climate change may be more realistic in a dynamic world, ultimately impacting aquatic biodiversity and ecosystem function in these extreme environments.

Literature Cited

Axler, R.P. and Reuter, J.E. (1996) Nitrate uptake by phytoplankton and periphyton: whole-lake. enrichments and mesocosm-¹⁵N experiments in an oligotrophic. *Limnology and Oceanography*. 41: 659-671.

Bradley, R.S., Keimig, F.T. and Diaz, H.F. (2004) Projected temperature changes along the American cordillera and the planned GCOS Network. *Geophysical Research Letters*. 31: L16210.

Chambers, P.A., Guy, M., Roberts, E.S., et al. (2001) Nutrients and their impact on the Canadian environment. Agriculture and Agri-Food Canada, Environment Canada, Fisheries and Oceans Canada, Health Canada and Natural Resources Canada, Ottawa.

Dodson, S.I. (1974) Zooplankton competition and predation: an experimental test of the size-efficiency hypothesis. *Ecology*. 55: 605-613.

Douglas, M.S.V. and Smol, J.P. (1999) Freshwater diatoms as indicators of environmental change in the high arctic. In *The Diatoms: Applications for the environmental and earth sciences*. Stoermer, E.F. and Smol, J.P. (eds.) Cambridge University Press, Cambridge.

Fenn, M., Haeuber, R., Tonnesen, G., et al. (2003) Nitrogen emissions, deposition, and monitoring in the western United States. *Bioscience*. 53: 391-403

McKnight, D.M., Runkel, R.L., Tate, C.M., et al. (2004) Inorganic nitrogen and phosphorus dynamics of Antarctic glacial meltwater streams as controlled by hyporheic exchange and benthic autotrophic communities. *Journal of North American Benthological Society*. 23: 171–188.

McMaster, N.L. (2003) Alpine ponds: climate change and N deposition. MSc thesis, University of Alberta, Edmonton, Alberta.

McMaster, N.L. and Schindler, D.W. (2005) Planktonic and epipelagic algal communities and their relationship to physical and chemical variables in alpine ponds in Banff National Park, Canada. *Arctic, Antarctic, and Alpine Research*. 37: 337–347.

Nydick, K. R., Lafrancois, B. M., Baron, J. S., et al. (2004) Nitrogen regulation of algal biomass, productivity, and composition in shallow mountain lakes, Snowy Range, Wyoming, USA. *Canadian Journal of Fisheries and Aquatic Sciences*. 61: 1256–1268.

Saros, J.E.; Interlandi, S.J.; Wolfe, A.P., et al. (2003) Recent changes in the diatom community structure of lakes in the Beartooth Mountain Range, USA. *Arctic Antarctic Alpine Research*. 35: 18-23

Thompson, P.L., St-Jacques, M.C., and Vinebrooke, R.D. (2008) Impacts of climate warming and nitrogen deposition on alpine plankton in lake and pond habitats: an in vitro experiment. *Arctic, Antarctic, and Alpine Research*. 40: 192–198.

Van Geest, G. J., Hessen, D.O., Spierenburg P., et al. (2007) Goose-mediated nutrient enrichment and planktonic grazer control in arctic freshwater ponds. *Oecologia*. 153: 653–662.

Vinebrooke, R.D. and Leavitt, P.R. (1998) Direct and interactive effects of allochthonous dissolved organic matter, inorganic nutrients, and ultraviolet radiation on an alpine littoral food web. *Limnology and Oceanography*. 43: 1065-1081

Vinebrooke, R.D. and Leavitt, P.R. (1999) Differential responses of littoral communities to ultraviolet radiation in an alpine lake. *Ecology*: 80: 223-237.

Wolfe, A.P., Cooke, C.A. and Hobbs, W.O. (2006) Are current rates of atmospheric nitrogen deposition influencing lakes in the Eastern Canadian Arctic? *Arctic, Antarctic, and Alpine Research*. 38: 465-476.